First report of marine sponge *Chelonaplysilla delicata* (Demospongiae: Darwinellidae) from

the Andaman Sea/Indian Ocean with a baseline information of epifauna on a mesophotic

shipwreck

Rocktim Ramen Das[1,2*], Titus Immanuel[1,3], Raj Kiran Lakra[1], Karan baath[4] and Ganesh

Thiruchitrambalam[1]

[1]Department of Ocean Studies and Marine Biology, Pondicherry University, brookshabad,

Andaman Islands, India

[2]Graduate School of Engineering and Science, University of the Ryukyus, Nishihara, Okinawa,

Japan

[3]Marine Biology Regional Centre (MBRC), Zoological Survey of India (ZSI), Chennai, Tamil Nadu,

India

[4]Infinity Scuba Andaman's, Chidiyatapu, Andaman Islands, India

Correspondence e-mail: k188604@cs.u-ryukyu.ac.jp; asomorlora@gmail.com

Abstract

During a biodiversity assessment on a wreck located in the Andaman Sea (Andaman Islands), a

single specimen of sponge *Chelonaplysilla delicata* was recorded. Our finding confirms the

species taxonomy and highlights the current observation as a first report from the Andaman

Sea/ Indian Ocean. The baseline information of epifauna is further stated in this study.

Keywords: Biodiversity, Porifera Taxonomy, Invasive, *Tubastraea*

INTRODUCTION

The Andaman Sea, an eastern subdivision of the Indian Ocean is bordered by countries like

Thailand, Myanmar on the east and the Andaman archipelago (Andaman & Nicobar

Islands/ANI) on the west (Brown 2007). (Figure 1). A large portion although falls within the

boundary of the Coral Triangle Initiative (CTI) (Rudi et al. 2012) studies related to its marine

biodiversity or the coral reef ecosystem has been comparatively understudied or scattered
(Aungtonya et al. 2000; Brown 2007). Additionally, the Andaman Sea possesses several
shipwrecks (Kheawwongjan & Kim 2012) acting as artificial reef ecosystems knowledge
pertaining to which is mostly limited in the region. These sunken structures provide space for
growth and establishment of various sessile marine communities like poriferans (Walker et al.
2007; Lira et al. 2010) and other non-native species (Patro et al. 2015; Soares et al. 2020).
Within the Indian EEZ, recent studies targeting shallow-water wrecks has filled important
knowledge gaps (example Mohan 2013; Das 2014; Yogesh-Kumar et al. 2015) (Table 1). This
article further adds essential information in these rarely studied ecosystems at a mesophotic
depth and reports a marine sponge from the Andaman Sea/Indian Ocean.

MATERIAL AND METHODS

The sponge *C.delicata* (Image 1), collected from the shipwreck HMIS SM* during a
survey conducted for documenting epifaunal diversity from February to March 2014. The wreck
is a 70m long Royal Indian navy minesweeper that sank in the year of 1942. At a depth of ≈33
meters, it lies at the edge of the Macpherson strait near Chidiyatapu (11^{o}28'38.02" N
92^{o}42'12.20" E) (Figure 1). Water transparency and temperature were recorded with Secchi
disc and dive calculator. Within 2 hours after collection, the specimen was preserved in 100%
ethanol. A surface peel of the easily separable cortex of the specimen was removed and placed
in xylene for 24 hours after which a permanent slide of the peel was mounted with DPX. A
single fibre with its base and branches intact was removed from the sponge for species-level
identification under a stereo microscope. (Image 1B-D). The specimen was identified following
Pulitzer-Finali & Pronzato (1999). The preserved specimen is deposited in the National
Zoological Collections (NZC) of the Andaman and Nicobar Regional Centre (ANRC), Zoological
Survey of India (ZSI), Port Blair.

Benthic cover was assessed by randomly placing 20 (0.25x0.25cm) quadrats (Image 2).
The photographs were analysed using open-sourced Coral-Net software (Beijbom et al. 2012)
and the epifauna was categorized into Unknown, Sponge, Hard Coral, *Ircinia* sp., Algae,
Iotrochota sp., Sediment, *Tubastraea* aff. *coccinea*, *Tubastraea micranthus*, Hard Substrate,

Ascidian, Bleached Coral (modified from Zintzen et al. 2006). Data from the annotated quadrats

was later transferred and processed in Microsoft Excel® (Microsoft 365 MSO,

16.0.13001.20266/32bit). Study maps were created using open-sourced Quantum Geographic

Information System (QGIS ver. 3.6).

RESULTS AND DISCUSSION

Systematics

Phylum: Porifera

Class: Demospongiae

Order: Dendroceratida

Family: Darwinellidae

Genus: *Chelonaplysilla*

Species: *C. delicata* Pulitzer-Finali & Pronzato, 1999

Paratype: 1 ex (Paratype)., ZSI/ANRC – 14321, India: Andaman Island: South Andaman:

Chidiyatapu (11°28'38.02" N; 92°42'12.20" E). Coll. Rocktim Ramen Das, 2014

Diagnosis

C.delicata predominantly thickly encrusting (< 10 mm) but has erect lobes that are about 4 - 5

cm high. The sponge surface is conulose, and the acute conules separated from each other by 2

- 5 mm. Oscules 1 - 3 mm in diameter, flush with the surface and unevenly distributed all over

on sponge surface. The texture is soft collapsible and feeble. The fresh specimen was dark

violet or purple in colour and retained its colour even in the preserved condition. Sponge

surface covered by structured regular reticulation of sand and spicule detritus, which forms

regular roundish or oval meshes of 90 - 155 μm. This reticulation is typical of the genus. Regular

rounded fibrous pores, inhalant in nature, is enclosed within these rounded meshes (Image.

1D). The skeleton is dendritic, made up of pigmented fibres fragile in nature with repeated

branching that originate from a basal spongin plate (Image 1B-C) and extends towards the

boundary. The primary fibre measured to be around 0.4 mm at its thickest. Spicules are absent.

Distribution: India: Andaman Sea (ANI, South Andaman, Present study). Elsewhere: Bismark Sea

(Papua New Guinea) (Pulitzer-Finali and Pronzato, 1999), Indonesia (Sulawesi) (GBIF, 2000),

Palau (Micronesia) (Ridley et al. 2005), French Polynesia (Alencar et al. 2017) (Figure 2).

Remarks: *C.delicata* is very similar to *C.erecta* (Tsurnamal, 1967); however, the latter has fibres

anastomosing in nature whereas the thickness of fibres in *C.delicata* fades in diameter. The

specimen mentioned in Pulitzer-Finali and Pronzato, 1999 is Gray whereas our specimen in dark

maroon in live condition (Image 1A). The specimen was initially misidentified as *C.erecta* (Das

2014; Das et al. 2016), thus there was a need for the update and fill the knowledge gap of this

species distribution range.

Comments: The family Darwinellidae possesses sponging fibres with proper skeleton and

fibrous spicules (Van Soest 1978; Bergquist & Cook 2002). It consists of four recognized genus

and forty-seven accepted species (one under "nomen nudum" status). *Chelonaplysilla* is the

only genus, which is devoid of spicules but consists of a fibrous dendritic skeleton that

possesses distinct laminated bark surrounding a central pith region. A structured and separable

cortex that is reinforced by a delicate reticulation of sand grains (Van Soest 1978) distinguishes

this genus.

Wreck Biodiversity: Benthic cover assessment (Image 2) reveal that sponges are the second

most abundant group on the surface of the wreck mostly encrusting in nature. In most

instances the encrusting sponge genus *Iotrochota* was readily visible. Ahermatypic and invasive

sun corals were abundant in selected localities and may have found a successful substrate for

further expansion (Image 2A, 3). Few polyps of *Tubastraea micranthus* had bleached, a stark

contrast to their ahermatypic nature. Updated and revised identification following Das et al.

(2016) on the wreck surface includes sclectenian genus *Favia*, *Symphyllia*, *Podabacia crustacea*,

and *Leptoseris*. A single individual of the Gastropod genus (*Chicoreus*), and few Crinoids. The

identified Poriferan families include Irciniidae (*Ircinia*), Chalinidae (*Haliclona* (*Reniera*));

Thorectidae (*Hyrtios*), Iotrochotidae (*Iotrochota baculifera*), Thorectidae (*Dactylospongia*).

Tunicates comprised of Didemnidae (*Didemnum*), Perophoridae (*Perophora*) and other
unidentified sp.

The Faunal organisms that thrive in Artificial Reefs (Shipwrecks) is an important part of
the marine community (Amaral et al. 2010; Zintzen et al. 2006). With increasing anthropogenic
impacts on natural coral reef habitats, artificial reefs are regarded as a successful alternative
(Perkol-Finkel & Benayahu 2005). As a result, it becomes important to understand the biological
communities growing on these habitats (Thanner et al. 2006). Sponges which naturally occupy
shipwrecks are one of the dominant organisms in such habitat, as evidenced in the present
study, however its diversity will be strictly limited to the environmental settings. For example,
some species of genus *Iotrochota* is found in sheltered environments (Cleary & Voogd 2007) as
seen in our observation (Image 2C). Similarly, shipwrecks are also known to act as a successful
substrate for many non-native species as reported from the Atlantic and the Red Sea (Perkol-
Finkel et al. 2006; Soares et al. 2020). Repeated encounter of *Tubastrea* aff. *coccinea* and T.
micranthus in the study site is a strong evidence from the Andaman Sea (Figure 1). The sponge
species reported in herein is in a much-extended depth compared to its initial described type
locality (see. Pulitzer-Finali & Pronzato 1999).

Technical difficulties are a foremost reason for these habitats to have limited studies at
mesophotic depths (Massin et al. 2002; Zintzen et al. 2006), but with the rapid scale
development of ROV's and submersibles, detailed exploration of these ecosystems can be well
predicted. Further these areas might be a hub for various underexplored flora and fauna and
might be effective in reviving threatened marine life due to the loss of natural ecosystems.

Author Contribution Statement

Rocktim Ramen Das: Conceptualization, Study Design, Writing – Original Draft, Reviewing &
Editing, Field Assessment, Sample Collection, Data Analysis, Laboratory Analysis.

Titus Immanuel: Writing Original Draft, Reviewing & Editing, Sample Identification, laboratory
Analysis.

Raj Kiran Lakra: Study Design; Writing, Reviewing & Editing, Field Assessment.

Karan Baath: Field Assessment

Ganesh Thiruchitrambalam: Study Design; Writing, Reviewing & Editing; Supervision.

Acknowledgements

The authors thank Dr. P.M. Mohan, Department of Ocean Studies and Marine Biology (DOSMB),
Pondicherry University, India for the necessary facilities. The authors acknowledge Dr. S.Venu
(DOSMB, Pondicherry University, India) for initial comments on the project. Drs. E. Hirose and F.
Sinniger (University of the Ryukyus, Japan) for valuable comments regarding this draft. Drs. K.
Wangkulangkul (PSU, Thailand), S.Y. Tenjing (NCSCM, India) for English editing.

Conflict of Interest

A preprint of this manuscript was uploaded on Bioarxiv preprint server on May 15th, 2019. The
preprint can be accessed at https://www.biorxiv.org/content/10.1101/636043v1.

References

Amaral, F. M. D., C. M. R. Farrapeira, S. M. A. Lira & C. A. C. Ramos (2010). Benthic macrofauna
inventory of two shipwrecks from Pernambuco Coast, Northeastern of Brazil. *Revista*
Nordestina de Zoologia-Recife 4(1): 24-41.
Alencar, Aline & Bourgeois, Bertrand & Butscher, John & Debitus, Cécile & Ekins, Merrick &
Fleurisson, Dominique & Folcher, Eric & Hall, Kathryn & Hertrich, Lionel & Hooper, John &
Lerouvreur, Franck & Levy, Peva & Maihota, Nicolas & Orempuller, Joel & Petek, Sylvain &
Pisera, Andrzej & Renaud, Armelle & Sutcliffe, Patricia & Vacelet, Jean. 2017. Sponges of
Polynesia. 827 pp.
Aungtonya, C., S. Thaipal & O. Tendal (2000). A preliminary report on the Thai-Danish bioshelf
surveys (1996-2000) of the west coast of Thailand, Andaman Sea. *Res. Bull. (Sun*
Chiwawitthaya thang Thale Phuket), 63: 53-76.

155 Beijbom, O., P. J. Edmunds, D. I. Kline, G. B. Mitchell & D. Kriegman. *"Automated Annotation of*
156   *Coral Reef Survey Images"*. IEEE Conference on Computer Vision and Pattern Recognition
157   (CVPR), Providence, Rhode Island, June 2012.

158 Bergquist, P. R. & Cook, S. C (2002). Family Darwinellidae Merejkowsky, 1879. In: Hooper J. N.
159   A., R. W. M Van-Soest (eds.). *Systema Porifera: A guide to the classification of*
160   *Sponges*, New York: Springer, 1068-1071 pp.

161 Brown B. E. 2007. Coral reefs of the Andaman Sea – An Integrated perspective In: Gibson R. N.,
162   R. J. A Atkinson, J. D. M Gordon (eds). *Oceanography and Marine Biology: An Annual*
163   *Review*, 547pp.

164 Cleary, D. F. R. & N. J. de Voogd. 2007. Environmental associations of sponges in the Spermonde
165   Archipelago, Indonesia. *Journal of the Marine Biological Association of the United*
166   *Kingdom*, 87: 1669-1676.

167 Das, R. R. (2014). An Inventory of Benthic Macrofauna from Two Shipwrecks along the Coast of
168   South Andaman. MSc Thesis, Department of Ocean Studies and Marine Biology,
169   Pondicherry University.

170 Das, R. R., R. K. Lakra, T. Immanuel, T. Ganesh (2016). A preliminary macrobenthic study of
171   World War two era wrecks along the coast of South Andaman. In: Bijukumar A, N. S.
172   Pradeep, K. G. Ajit Kumar, P. G. Rajendran (eds.) *Perspectives on Biodiversity of India*,
173   Centre for Innovation in Science & Social Action (CISSA), 83-92pp.

174 GBIF. 2000. https://www.gbif.org/species/2248714. Accessed: 12[th] September 2020.

175 Kheawwongjan, A. & D. S. Kim (2012). Present status and prospects of artificial reefs in
176   Thailand. *Ocean and Coastal Management* 57: 21-33.

177 Lira de S. M. A., C. M. R. Farrapeira, F. M. D. Amaral & C. A. C Ramos (2010). Sessile and
178   Sedentary Macrofauna from the Pirapama Shipwreck, Pernambuco, Brazil. *Biota*
179   *Neotropica* 10(9): 155-165*.*

180 Massin, C., A. Norro. and J. Mallefet (2002). Biodiversity of a shipwreck from the Belgian
181   Continental Shelf: monitoring using scientific diving Preliminary results. *Bulletin van het*
182   *Koninklijk Belgisch Instituut voor Natuurwetenschappen, Biologie*, 72: 67–72.

Mohan, H. (2013). Fish Assemblage on Artificial Reef and Natural Reef (A Comparative Study).
MSc Thesis, Department of Ocean Studies and Marine Biology, Pondicherry University.

Mondal, T. & C. Raghunathan (2017). Shipwrecks in Andaman and Nicobar Islands: An artificial
habitat for corals, *Journal of the Marine Biological Association of India* 59(2): 92-101.

Perkol-Finkel, S. & Y. Benayahu (2005). Recruitment of Benthic Organisms onto a Planned
Artificial Reef: Shifts in Community Structure One Decade Post Deployment. *Marine*
Environmental Research 59(2): 79–99.

Pulitzer-Finali, G. & R. Pronzato (1999). Horny Sponges from the North-Eastern Coast of Papua
New Guinea, Bismark Sea. *Journal of the Marine Biological Association of the United*
Kingdom 79(4): 593-607.

Patro, S., P. Krishnan, M. Gopi, S. Raja, P. Ramachandran & R. Ramesh (2015). Snowflake coral,
Carijoa riisei from Grand Island, Goa: a case of invasion of an alien species or re-
establishment of a native species.? *Current Science* 109(6): 1028-1030.

Rudi, E., S. J. Campbell, A. S. Hoey, N. Fadli, M. Linkie & A. H. Baird (2012). The coral triangle
initiative: what are we missing? A case study from Aceh. *Oryx* 46(4): 482-485.

Ridley, C. P., P. R. Bergquist, M. K. Harper, D. J. Faulkner, J. N. A. Hooper & M. G. Haygood
(2005). Speciation and Biosynthetic variation in four dictyoceratid sponges and their
cyanobacterial symbiont, *Oscillatoria spongeliae*. *Chemistry and Biology* 12: 397-406.

Soares, M. D. O., S. Salani, S. V. Paiva & M. D. A. Braga (2020). Shipwrecks help invasive coral to
expand range in the Atlantic Ocean, *Marine Pollution Bulletin* 158: 111394.

Thanner, S. E., T. L. Mcintosh & M. Stephen. 2006. Development of Benthic and Fish
Assemblages on Artificial Reef Materials Compared to Adjacent Natural Reef Assemblages
in Miami-Dade County, Florida. *Bulletin of Marine Science* 78(1): 57-70.

Tsurnamal, M. (1967). *Chelonaplysilla erecta* n. sp. (Demospongiae, Keratosa) from
Mediterranean Coast of Israel. *Israel Journal of Zoology* 16(2): 96-100.

Van Soest, R. W. M. (1978). Marine Sponges from Curaçao and Other Caribbean localities Part I.
Keratosa. Studies on the Fauna of Curaçao and other Caribbean Islands 56(1): 1-94.

Walker, S. J., T. A. Schlacher. & M. A. Schlacher-Hoenlinger (2007). Spatial Heterogeneity of

Epibenthos on Artificial Reefs: Fouling Communities in the Early Stages of Colonization on

an East Australian Shipwreck. *Marine Ecology* 28: 435-445.

Yogesh-Kumar J. S., S. Geetha, C. Raghunathan & K. Venkataraman (2015). An Assessment of

Faunal Diversity and its conservation of shipwrecks in Indian seas. In: Venkataraman K, C.

Sivaperuman C. (eds.) *Marine Faunal Diversity in India: Taxonomy, Ecology and*

Conservation, Elseiver USA, 441-450pp.

Zintzen, V., C. Massin, A. Norro & J. Mallelfet. 2006. Epifaunal inventory of two shipwrecks from

the Belgian Continental Shelf. *Hydrobiologia* 555(1): 207-219.

Figure 1. Location of the study area

Figure 2. Global distribution of *C. delicata* Pulitzer-Finali & Pronzato, 1999

Figure 3. Mean percent cover of epifauna obtained from twenty 0.25x0.25 quadrats

Image 1. *Chelonaplysilla delicata* [ZSI/ANRC-14321]: (A) Freshly collected specimen, (B) Branching fibres and basal sponging plate, (C) Closer view of pigmented, branching, dendritic spongin fibre, (D) Inhalant pores surrounded by rounded meshes reinforced by sand grains. Scale (A) 5mm (B) 2 mm, (C) 2 mm, (D) 155 μm.

Image 2 :A) *Tubastraea* aff. *coccinea* B) *Ircinia* sp. C) Mixed assemblage of communities. Encrusting sponge *Iotrochota* sp. (Yellow) D) Mixed assemblage, Sponge (Yellow), coral (green).

236

237   Image 3. A part of the wreck HMIS SM. (Arrow: high abundance of invasive *Tubastraea*

238   *micranthus*)

240   Table 1. Information on biodiversity studies in a wreck ecosystem in Andaman and Nicobar Islands

Wreck Name	Co-ordinates	Location	Date of Sinking	Depth (m)	Current Activities	Reference
SS Inchkeith	12°00'23.69"N 92°46'08.34"E	Kyd Island (South Andaman)	1955	14	Scuba	Mohan 2013
HMIS Sophie Marie	**11⁰28'38.02"N 92⁰42'12.20" E**	**Chidiyatapu (South Andaman)**	**1942**	**30 - 33**	**Scuba**	**Present Study**
MV Mars	11°55'54.98"N 92°57'24.12"E	Havelock (Ritchie's Archipelago)	2006	10 - 16	Scuba	Das R. R. (Pers. Obs.)
North Bay Wreck	11°43'00.56"N 92°45'60.60" E	Port Blair (South Andaman)	30 – 40 (yrs)	10	Scuba and Fishing	Yogesh-Kumar et al. 2015
Peel Wreck	12°03'84.20"N 92°57'81.10" E	Havelock (Ritchie's Archipelago)	8 - 10	9 - 12.	Scuba	Yogesh-Kumar et al. 2015
Japan Wreck	09°10'88.30"N 92°50'12.30" E	Car Nicobar (Nicobar Islands)	40 - 50	28	Fishing	Yogesh-Kumar et al. 2015
Sinclair Bay Shipwreck	11°39'873"N 92°45'488"E	Near Ross Island (South Andaman)	-	8		Mondal & Raghunathan 2017

241