Title

The effect of NMDA-R antagonist, MK-801, on neuronal

mismatch along the auditory thalamocortical pathway.

Authors

Gloria G Parras[1,2], PhD; Catalina Valdés-Baizabal[1,2], PhD, Lauren Harms[3], PhD,
Patricia Michie[3], PhD & Manuel S Malmierca, PhD [1,2,4]*

Affiliations

[1] Cognitive and Auditory Neuroscience Laboratory, Institute of Neuroscience of Castilla
y León (INCYL), Salamanca, Spain.

[2] The Salamanca Institute for Biomedical Research (IBSAL), Salamanca, Spain.

[3] School of Psychology, University of Newcastle, Callaghan, NSW, Australia; Priority
Centre for Brain and Mental Health Research, Callaghan, NSW, Australia; Hunter
Medical Research Institute, Newcastle, NSW, Australia.

[4] Department of Cell Biology and Pathology, Faculty of Medicine, University of
Salamanca, Salamanca, Spain.

Running title: MK-801 affects auditory neuronal mismatch

Contact Information

* msm@usal.es

Number of pages: 38   **Number of figures/tables:** 6/2

Number of word abstract: 242   **Introduction:** 677   **Discussion:** 1582

Competing interest.

The authors report no competing interests.

ABSTRACT

Efficient sensory processing requires that the brain is able to maximize its response to unexpected stimuli, while suppressing responsivity to expected events. Mismatch negativity (MMN) is an auditory event-related potential that occurs when a regular pattern is interrupted by an event that violates the expected properties of the pattern. MMN has been found to be reduced in individuals with schizophrenia in over 100 separate studies, an effect believed to be underpinned by glutamate N-methyl-D-aspartate receptor (NMDA-R) dysfunction, as it is observed that NMDA-R antagonists also reduce MMN in healthy volunteers. The aim of the current study is to examine this effect in rodents. Using single unit recording in specific auditory areas using methods not readily utilized in humans, we have previously demonstrated that neuronal indices of rodent mismatch responses recorded from thalamic and cortical areas of the brain can be decomposed into a relatively simple repetition suppression and a more sophisticated prediction error process. In the current study, we aimed to test how the NMDA-R antagonist, MK-801, affected both of these processes along the rat auditory thalamocortical pathway. We found that MK-801 had the opposite effect than expected, and enhanced thalamic repetition suppression and cortical prediction error. These single unit data correlate with the recordings of local field responses. Together with previous data, this study suggests that our understanding of the contribution of NMDA-R system to MMN generation is far from complete, and also has potential implications for future research in schizophrenia.

Keywords: Mismatch negativity, predictive coding, prediction error, repetition suppression, schizophrenia, auditory.

Significance Statement: (120 words – currently 119 words) In this study, we demonstrate that an NMDA-R antagonist, MK-801, differentially affects single neuron responses to auditory stimuli along the thalamocortical axis by increasing the response magnitude of unexpected events in the auditory cortex and intensifying the adaptation of responses to expected events in the thalamus. Thus, we provide evidence that NMDA-R antagonists alter the balance between prediction-error and repetition suppression processes that underlie the generation of mismatch responses in the brain, and these effects are differentially expressed at different levels of auditory processing. As effects of MK-801 were in the opposite direction to our expectations, it demonstrates that our understanding of role of NMDA-R in synaptic plasticity and the neural processes underpinning MMN generation are far from complete.

INTRODUCTION

Mismatch negativity (MMN) is an auditory event-related potential (ERP) that occurs when an unexpected stimulus (deviant, DEV) interrupts a train of regular stimuli (standards, STD) in an *oddball* sequence. MMN is commonly quantified as the difference between the size of the DEV and STD ERPs (Näätänen *et al.*, 1978).

The predictive coding framework has emerged as an appealing model of MMN (Randeniya *et al.*, 2018) and of how sensory information is processed. According to this model, the brain constantly generates top-down predictions from any regular ascending input that is compared with sensory bottom-up signals. Neural responses to stimuli that match predictions are suppressed, whereas unexpected stimuli discrepant with the prediction generate an enhanced error signal (Friston, 2005; Michie *et al.*, 2016; Carbajal & Malmierca, 2018). NMDA-R dependent plasticity is believed to underpin the capacity of the brain to adjust internal predictions and use memory of recent past inputs to anticipate future stimuli (Wacongne, 2016).

There are two mechanisms underlying the MMN difference signal according to the predictive coding model. First, MMN could reflect *repetition suppression*. When the same stimulus is repeatedly presented, neuronal populations sensitive to that stimulus undergo adaptation and neural responses decrease (Bendixen *et al.*, 2007). MMN could also reflect a process of *prediction error*, where the sensory memory of regular stimuli establishes a predictive model. Violation of this prediction upon presentation of an unexpected DEV stimuli results an enhanced neural response. Prediction error has been observed in human and rodent surface recordings when suitable control conditions have been included in the design of sound sequences (Nakamura *et al.*, 2011; Harms *et al.*,

2014; Parras *et al.*, 2017; Kurkela *et al.*, 2018). Essentially, there is evidence in both

humans and rodents that MMN receives contributions from both prediction error and

repetition suppression at various levels of the auditory system (Parras *et al.*, 2017;

Ishishita *et al.*, 2018).

MMN reduction occurs in healthy volunteers administered an NMDA-R

antagonist (Umbricht *et al.*, 2000) and has been reported in over 100 separate reports in

individuals diagnosed with schizophrenia (Todd et al., 2013), leading to assertions that

MMN indexes the functional state of NMDA-R neurotransmission (Umbricht *et al.*,

2002). The schizophrenia findings fit with current views that NMDA-R hypofunction

contributes to the neuropathology of the disorder (Krystal *et al.*, 2005; Javitt et al.,

1996). Mismatch like responses (MMRs) in animal models have shown a similar

sensitivity to NMDA-R antagonism (Javitt *et al.*, 1996; Siegel *et al.*, 2013; Harms,

2016; Featherstone *et al.*, 2018).

Our primary interest here is whether NMDA-R antagonists differentially affect

repetition suppression and prediction error at the single unit level at both the thalamic

level and cortex. While previous studies have demonstrated that MMN-like responses in

rodents are altered by NMDA-R antagonists (Ehrlichman *et al.*, 2008; Tikhonravov *et*

al., 2008, 2010; Sivarao *et al.*, 2014), only one report has examined the impact on

prediction error component of the MMN in surface recordings (Harms *et al.*, 2018).

There are no published data on the effects of NMDA-R antagonists on repetition

suppression. More importantly, there are no reports that have examined the effects of

NMDA-R antagonists on single-unit activity and local field potential recordings from

the thalamus and auditory cortex.

Thus, it is unknown (i) whether there are differential effects of NMDA-R

antagonism on prediction error as opposed to repetition suppression at the single unit or

local field potential level, and (ii) the regional specificity of where effects of NMDA-R

antagonists occur: in the lemniscal *vs.* non-lemniscal auditory areas, or the thalamus *vs.*

cortex. Therefore, we used an acute exposure to a low dose of MK-801 to examine the

impact of NMDA-R antagonism on individual responses of lemniscal and non-

lemniscal thalamocortical neurons while auditory oddball, many standards and cascade

control sequences were presented (figure 1a-b). This design allowed us to delineate

effects on repetition suppression *vs.* prediction error (figure 1c)(Opitz *et al.*, 2005;

Ruhnau *et al.*, 2012; Harms, 2016; Parras *et al.*, 2017). Unexpectedly, we found that

MK-801 increased repetition suppression in the thalamic regions, enhanced prediction

error in the cortex and increased neuronal mismatch at both locations.

MATERIAL AND METHODS

Experiments were performed on 48 (control=25; MK-801=23) adult, female Long-

Evans rats with body weights between 200-250g (aged 9 to 15 weeks). All experimental

procedures were performed at the University of Salamanca, and all procedures and

experimental protocols were in accordance with the guidelines of the European

Communities Directive (86/609/EEC, 2003/65/EC and 2010/63/EU) and the RD

53/2013 Spanish legislation for the use and care of animals. All the details of the study

were approved by the Bioethics Committee of the University of Salamanca (ref#

USAL-ID-195).

Surgical procedures: Anesthesia was induced and maintained with urethane (1.5g/kg,

i.p), with supplementary doses (0.5g/kg, i.p) given as needed. Dexamethasone

(0.25mg/kg) and atropine (0.1mg/kg) were administered at the beginning of the surgery

to reduce brain edema and bronchial secretions, respectively. Isotonic glucosaline

solution was administered periodically (5-10ml every 6-8h, s.c) to avoid dehydration.

During all experimental procedures, animals were artificially ventilated, and CO_2 and

temperature monitored (Pérez-González *et al.*, 2012; Duque & Malmierca, 2015).

The initial procedure was the same in each case, and the subsequent procedures differed

only in the craniotomy location, and the placement/orientation for the recording

electrode (animals per group/location: control MGB=16, AC=9; MK-801 MGB=15,

AC=8). For MGB recordings, a craniotomy (~2x2mm, from -5 to -6.5mm bregma and -

3.5mm lateral) was performed in the left parietal bone, dura was removed, and the

electrode advanced in a vertical direction (Antunes *et al.*, 2010). For AC recordings, the

skin and muscle over the left temporal bone was retracted and a 6x5mm craniotomy was

performed (between -2 and -6 from Bregma) over the temporal bone (Nieto-Diego &

Malmierca, 2016) dura was removed and the area was covered with a thin, transparent

layer of agar to prevent desiccation and stabilize recordings. Electrodes for AC

recording were inserted using a triple axis micromanipulator (Sensapex), forming a 30º

angle with the horizontal plane, to penetrate through all cortical layers of the same

cortical column.

For this study, animals in MK-801-treated group receive a systemic intraperitoneal

injection (0.1mg/kg) of a noncompetitive NMDA-R antagonist (MK-801 hydrogen

maleate, M107 Sigma-Aldrich). Control animals did not receive any injection.

Electrophysiological recording procedures. During all procedures, animals were placed in a stereotaxic frame fixed with hollow specula ear bars that housed the sound delivery system. One single neuron and local field potential (LFP) was recorded at a time, using the same tungsten electrode (1-4MΩ) inserted into a single auditory station (MGB or AC) in each individual animal. The signal recorded was pre-amplified (1000x) and band-pass filtered (1-3kHz) with a medusa preamplifier (TDT). This analog signal was digitalized 12k sampling rate and further band-pass filtered (TDT-RX6) separately for spikes (500Hz-3kHz) and LFP (3-50Hz). We used short trains of white noise bursts (30 ms, 5 ms rise-fall ramps) to search for neuronal activity. To prevent neuronal adaptation during the search, some parameters (frequency and intensity) and stimulus type (white noise, pure tone) were manually varied. Once a single neuron was isolated a frequency-response area (FRA) of the response magnitude for each frequency/intensity combination was first computed (Figure 1a). A randomized sequence of pure tones (from 1 to 44 KHz) was presented at a rate of 4Hz, with varying frequency and intensity, and with 3 repetitions of all tones.

For each animal treated with MK-801 the first single neuron was recorded ~15 min after the drug injection (Vezzani *et al.*, 1989). Ten evenly-spaced pure tones (0.5 octaves separation) at a fixed sound intensity (usually 20-30dB above the threshold) were selected to each neuron recorded to create the control sequences, cascades and many-standard (Ruhnau *et al.*, 2012; Parras *et al.*, 2017), and additionally, adjacent pairs of them were used to present various oddball sequences (Figure 1b). All sequences were 400 tones in length (75ms duration, 5ms rise-fall ramp and 250ms interstimulus interval), each tone in the control sequences was played 40 times, with the same overall presentation rate as deviants in the oddball sequence.

Oddball sequences were used to test the specific contribution of deviant tones in an

adaptation context. An oddball sequence consisted of a repetitive tone (standard 90%

probability), occasionally replaced by a tone of a different frequency (deviant 10%

probability), in a pseudorandom manner. We used two types of control sequences: the

many-standard and cascade sequences. Both containing the same 10 frequencies but

differing in the order of presentation. The many-standard control was randomly

presented, mimicking the presentation rate and the unpredictability of the deviant tones.

While cascades were played always in the same presentation order, ascending or

descending in frequency. Hence the cascade contains a regularity, mimic the

presentation rate of deviant sounds but in a predictable context and consequently do not

violate a regularity. These four conditions, and by extension responses to them, will be

denoted as deviant (DEV), standard (STD), cascade (CAS) and many-standard. Finally,

if the neuron could be held for long enough, the same protocol was repeated for

different frequencies and/or intensity.

Anatomical location. For MGB recording localization, at the end of each tract and

experiment, two electrolytic lesions were made to mark the end and the beginning of the

auditory signal (figure 6a). Then, animals were given a lethal dose of sodium

pentobarbital and perfused transcardially with phosphate buffered saline (0.5% $NaNO_3$

in Phosphate Buffered Saline) followed by a fixative mix of 1% paraformaldehyde and

1% glutaraldehyde). After fixation and dissection, the brain was cryoprotected in 30%

sucrose and sectioned into 40μm slices. Sections were Nissl stained with 0.1% cresyl

violet. Recording sites were marked on images from an adult rat brain atlas (Paxinos &

Watson, 2013) and neurons that were recorded from were assigned to one of the main

divisions of the MGB (dorsal, medial or ventral). This information was complemented

and confirmed by the stereotaxic coordinates as well as the depth of the neuron within a

tract.

For the AC experiments, a magnified picture (25x) of the exposed cortex and the

Bregma references was taken at the end of the surgery with a digital single lens reflex

camera (D5100, Nikon) coupled to the surgical microscope (Zeiss). The picture was

overlapped to guide and mark each electrode placement into a micrometric grid (250-

500 ~μm spacing; figure 6b). Then we performed several tracts recording multi-unit

activity frequency response area (FRA), the characteristic frequency arise from each

FRA was placed over the picture, resulting in a characteristic frequency map of each

animal. Boundaries were identify following the changes in the tonotopic gradient: high-

frequency reversal between the ventral and anterior auditory fields (rostrally), low-

frequency reversal between primary and posterior auditory field (dorsocaudally) and

high-frequency reversal between ventral and suprarhinal auditory field (ventrally)

(Nieto-Diego & Malmierca, 2016). Then, each recording was located in one of these

five fields. Nevertheless, the map was complemented during all electrophysiological

recording session with the characteristic frequency of each new tract.

Statistical analysis. All the data analyses were performed with MatlabTM software,

using the built-in functions, the Statistics and Machine Learning toolbox, or custom

scripts and functions developed in our laboratory. To test for significant excitatory

responses to tones we used a Monte Carlo approach, simulating 1000 peri-stimulus time

histogram (PSTH) using a Poison model with a constant firing rate equal to the

spontaneous firing rate. A null distribution of baseline-corrected spike counts was

generated from this collection of PSTH. Lastly, the *p* value of the baseline-corrected

spike count was empirically computed as $p = (g + 1)/(N + 1)$, where g is the count of null

measures greater than or equal to baseline-corrected spike count, and N=1000 is the size

of the null sample. Finally, we only included in the analysis neuron/frequency

combinations with significant excitatory response ($p > 0.05$) after the baseline-corrected

spike count to at least one of the conditions (DEV, STD, CAS). PSTH were used to

estimate the spike-density function (SDF) over the time, showing action potential

density over time (in action potentials per second) from −75 to 250ms around stimulus

onset, for the 40 trials available for each tone and condition (DEV, STD, CAS),

smoothed with a 6ms gaussian kernel ("ksdensity" function in Matlab) in 1ms steps.

The baseline spontaneous firing rate was determined as the average firing rate during

the 75ms preceding stimulus onset.

The excitatory response was measured as the area below the SDF and above the

baseline spontaneous firing rate, between 0 and 180ms after stimulus onset (positive

area patches only, to avoid negative response values). This measure will be referred to

as "baseline-corrected spike count".

Baseline-corrected spike count responses of a neuron to the same tone in the three

conditions (DEV, STD, CAS) were normalized using the formulas:

$DEV_{Normalized} = DEV/N;$

$STD_{Normalized} = STD/N;$

$CAS_{Normalized} = CAS/N;$

Where $N = \sqrt{DEV^2 + STD^2 + CAS^2}$, is the Euclidean norm of the vector (DEV, STD,

CAS) defined by the three responses. Normalized values were the coordinates of a 3D

unit vector (DEV$_{Normalized}$, STD$_{Normalized}$, CAS$_{Normalized}$) with the same direction of the

original vector (DEV, STD, CAS), and thus the same proportions between the three

response measures. This normalization procedure always results in a value ranging 0–1,

and has a straightforward geometrical interpretation.

From these normalized responses, indices of neuronal mismatch (iMM), repetition

suppression (iRS), and prediction error (iPE) were computed as:

iMM = DEV$_{Normalized}$ − STD$_{Normalized}$;

iPE = DEV$_{Normalized}$ - CAS$_{Normalized}$;

iRS = CAS$_{Normalize}$ - STD$_{Normalized}$;

These indices, consequently, always range between −1 and 1, and provide the

following quantitative decomposition of neuronal mismatch into repetition suppression

and prediction error: iMM = iRS + iPE. To test these indices over time, we divided the

whole response into 12 time windows, 20ms width, from -50 to 190ms with respect to

the stimulus onset. Then, we compared each time window against zero using a sign-rank

test, false discovery rate (FDR=0.1) corrected for the 12 windows.

For the analysis of the LFP signal, we aligned the recorded wave to the onset of the

stimulus for every trial, and computed the mean LFP for every recording site and

stimulus condition (DEV-LFP, STD-LFP and CTR-LFP), as well as the differences

between them, resulting in the three LFP-indices: "neuronal mismatch" (MM-LFP =

DEV-LFP – STD-LFP), "prediction error" (PE-LFP = DEV-LFP – CAS-LFP) and

"repetition suppression" (RS-LFP = CAS-LFP – STD-LFP). Then, grand-averages were

computed for all conditions and auditory station separately. The p value of the grand-

averaged for the three LFP-indices (MM-LFP, PE-LFP and RS-LFP) was determined

for every time point with a two-tailed t test (FDR corrected).

Our data set was not normally distributed, so we used distribution-free (non-

parametric) tests. These included the Wilcoxon signed-rank test and Friedman test (for

baseline-corrected spike counts, normalized responses, indices of neuronal mismatch,

repetition suppression and prediction error). Only the difference wave for the LFPs was

tested using a t-test, since each LFP trace is itself an average of 40 waves. For multiple

comparison tests, p values were FDR corrected using the Benjamini-Hochberg method.

Linear models were used to test for significant average iMM, iPE and iRS within each

auditory station. Significant effects of station, pathway, and interactions between them

were fitted using the 'fitlm' function in Matlab, with robust options. To estimate final

sample sizes required for the observed effects after the initial exploratory experiments,

we used the 'sampsizepwr' function in Matlab adjusted for the iPE for each region, to

obtain a statistical power of 0.8 for this index. Sample sizes were enlarged with

additional experiments until they were just greater than the minimum required (number

of points recorded, and the minimum required for each station; see Table 1).

To analyze the time course of adaptation we computed an averaged time course for all

the standard stimuli presented. Then, we fitted a power law function with a three

parameters model, $y(t)=a \cdot t^b + c$, where a indicates the responses beginning or the first

spike strength; b the sensitivity to repetitive stimuli, or the adaptation velocity, and c the

steady-state response. R^2 values indicated that the model fits very well for standard

responses in both groups, explaining between 60% and 78% of the response variability

within all regions.

To analyze spikes differences between MK-801 and control group we computed the

median values for each condition tested (DEV, STD and CAS) and their differences

(iMM, iRS and iPE) and calculated a ranksum test. To compare each time window

between groups a two-sample *t*-test (from 0 to 200ms, Bonferroni corrected for 200

comparisons with family-wise error rate FWER< 0.05) was used for the SDF and LFPs

to each stimulus condition and indices, using the 'ttest2' function in Matlab, for every

time point.

RESULTS

We recorded a total of 290 well isolated neurons, 143 from the control group and 147

from the MK-801-treated group. One single neuron and local field potential (LFP) was

simultaneously recorded at a time, using the same tungsten electrode. Recordings were

performed in the medial geniculate body (MGB) and in the auditory cortex (AC) while

playing oddball and control sequences (many-standards and cascade: CAS) in

anesthetized rats. Since we found no statistically significant differences between the use

of the cascade and many-standards sequences for the control group and MK-801 group,

except for the MGB_{NL} from the MK-801 group (table 1), the CAS sequence was chosen

to control for repetition effects. This is because the CAS paradigm not only controlled

for the presentation rate of the deviant stimuli, but also the frequency difference

(ascending or descending) between standards and deviants in the oddball sequences.

Effects of MK-801 on the neuronal firing rate. MK-801 injection significantly

reduced the responses to STD tones within all regions. By contrast, for responses to the

DEV tones, we observed a significant increment in responses in AC but not for the

MGB. When the firing rate of the cascade sequence was considered, MK-801

differentially affected the AC and MGB such that CAS responses were significantly

increased in the MGB_{NL} but decreased in AC. These results reveal a differential effect

of MK-801 on the refractoriness and salience of infrequent events at the single neuron

level (Figure 2a, table 2).

Effects of MK-801 on neuronal mismatch and its components. Next we analyzed the

differences between these normalized responses and computed three indexes (ranging

between -1 and +1): 1) the index of neuronal mismatch (iMM=DEV-STD), similar to

the typical SSA index used in previous single neurons studies; 2) the index of prediction

error (iPE= DEV-CAS), that shows the relative enhancement of DEV tones compared

with CAS tones and 3) the index of repetition suppression (iRS=CAS-STD) that

reflects the level of response suppression due to the repetition effect, and is obtained by

comparing the normalized responses to CAS and STD. It should be noted that the iMM

is the sum of iRS and iPE (iMM=iRS+iPE).

The analysis of the iMM after the injection of MK-801 demonstrated that iMM values

are significantly different from zero for all recording sites (figure 2b, table 1: Friedman

test). But when comparisons between groups were considered, the analysis revealed that

MK-801 increased the neuronal iMM (figure 2b-iMM; table 2). As described above,

these changes are largely due to a reduced response to STD tones in all recording

locations and an enhanced response to DEV in the AC.

Since iMM=iRS+iPE, an important advantage of these metrics is that we can determine

how much of the mismatch index is due to the regularity of the context (RS) and/or to

the occurrence of an infrequent event (PE). Thus, to determine which of these two

components of the iMM is affected by MK-801, we computed the indices of iPE and

iRS separately.

Interestingly, MGB neurons in the MK-801 group did not show any sign of genuine

deviance detection, as iPE values were almost zero and negative. While both AC

showed a significant positive iPE (figure 2c; iPE values in table 1). When comparison

between groups were analyzed an increased iPE for the MK-801 group in the AC were

found, and even a further decreased iPE for the MGB_{NL} in the MK-801 group (figure 2c

and e light and bright oranges; iPE in table 2). These data suggest that the MK-801

produces an augmentation of saliency for novel stimuli processed in the AC.

Yet, the detection of rare or novel stimuli requires the establishment of a regular context

or pattern. Therefore, we were also interested to find out if the refractoriness due to

regularity was altered by MK-801. We calculated the iRS by assessing the response of

the same tone when it was presented as CAS, with a 10% probability in a regular pattern

and presented as STD with a probability of 90%, within an oddball paradigm, so it is in

a much more regular context (Harms et al., 2014; Ruhnau et al., 2012). In both cases,

we assume some level of regularity adaptation, but only a genuine repetition

suppression can be determined if the responses to STD tones are lower than responses

to CAS. Our results demonstrate that there is a significant repetition suppression effect

in the MK-801 group along the thalamocortical pathway (figure 2d bright blue; iRS in

table 1). The analysis also revealed that MK-801 produced a significant increase in

repetition suppression at thalamic level but did not affect repetition suppression in the

AC when compared with controls (figure 2d-e light and bright blues; results in table 2).

These results show that the auditory thalamus and cortex differ in the way repetition

effects and prediction errors are processed. To confirm this hypothesis and considering

that we have previously found an increase in the level of iPE along the thalamocortical

hierarchy in awake and anesthetized animals (Parras et al., 2017), we fitted a linear

model to determine the effect of MK-801 on the degree of increase in iPE along the

auditory hierarchy relative to the control (urethane) condition. Using station (S: MGB

vs. AC), drug (D: Control vs. MK-801) and pathway (P: Lemniscal *vs.* Non-lemniscal)

and their interactions as factors, with control MGB_L as reference level for these factors,

the fitted model is as follows:

$iPE = -0.099 - 0.052D + 0.16P + 0.216S - 0.032DP + 0.297DS + 0.003SP + 0.1490DPS$.

Next, we applied an ANOVA to this model and found a significant effect of station

(F=209.24, p=1.21x10^{-44}), drug (F=10.82, p=0.001), pathway (F=42.93, p=7.66x10^{-11})

and for the interaction drug *x* station (F=37.9, p=9.41x10^{-10}) but not for the interactions

of drug *x* pathway (F=0.45, p=0.5011) nor station *x* pathway (F=0.004, p=0.9488), nor

the three way interaction (F=2.37, p=0.1236). That is, there is a significant station effect

(iPE is larger overall at AC than MGB), pathway (iPE is larger overall in the non-

lemniscal than the lemniscal pathway), and drug (iPE is increased overall with MK801

vs control) and a drug x station interaction indicating that while iPE at AC is increased

overall relative to MGB, the degree of iPE increase is moderated by MK801. Inspection

of Figure 2e indicates that the significant station difference is in the direction of being

larger under MK801 than urethane. These results indicate that indeed, the sensitivity to

detect novel stimuli increases significantly more along the thalamocortical axis in the

MK-801 than the control group (figure 2e, iPE in orange).

We also fitted a similar linear model to iRS. The resulting model was:

iRS = 0.467 + 0.167D - 0.118P - 0.125S +0.01DP - 0.159DS + 0.0405SP - 0.0424DPS.

ANOVA demonstrated a significant effect for drug (F=26.42, p=3.08x10^{-7}), station

(F=80.52, p=7.89x10^{-19}), pathway (F=24.65, p=7.6x10^{-7}) and the interaction of drug x

station (F=17.61, p=2.84x10^{-5}), but not for the interactions drug x pathway (F=0.078,

p=0.7790), station x pathway (F=1.45, p=0.2281) and the three way interaction (F=0.31,

p=0.5759). Once again, these results indicate that while iRS was more marked at MGB

than AC in general, this station difference was exacerbated with MK-801.

In summary, the changes described above demonstrate that NMDA-R antagonism has

distinct effects on auditory scene analysis, as measured by the iPE and iRS, at different

levels of the thalamocortical hierarchy.

Effect of MK-801 on Spike-Density Function and indexes. Next, we sought to

identify how MK-801 affected the temporal responses to auditory stimuli (DEV, STD

and CAS) by comparing spike-density functions (SDF) to each condition between

groups. Analysis revealed the latency of the main peak for the SDF to DEV tones was

mostly unaffected by MK-801 in the MGB, but it was clearly delayed by 40 and 60 ms

in the AC_L and AC_{NL}, respectively. Furthermore, the magnitude of the SDF was altered

at the AC and MGB_{NL}, with the early component being reduced and the later sustained

component being enhanced (figure 3a, horizontal white line for significant differences

at $p<0.05$). When the STD tones were considered, we observed a distinct and significant

decrease of the SDF mostly at the AC and only marginally at the subcortical levels

(figure 3b). Finally, MK-801 affected mostly the initial responses to cascade tones at all

regions, being reduced in the auditory cortex but was earlier and increased in MGB_L

(figure 3c). The sustained portion of the SDF was only significantly increased in the

MGB_{NL}. Results show that MK-801 has a profound effect on the spike-density

functions to DEV, STD and cascade stimuli.

Next, we studied where and when the MK-801 effect on the neuronal indices of iMM,

iPE and iRS was significantly different from control. Thus, we examined whether in

each group independently (MK-810 and control) these indices are different from zero,

i.e., is there a significant iMM, iPE or iRS at each time point. Figure 3d-f highlights the

significant time windows ($p<0.01$) with white and black asterisks for control and MK-

801, respectively. The analysis revealed that under MK-801, there was a significant

iMM along the thalamocortical axis (between 20-40ms for MGB_L, 20-80ms in MGB_{NL}

and from 20-190ms in both AC; Figure 3d, bright purple lines) and a significant iPE

between 20 and 180ms in both AC, and a late iPE in the lemniscal thalamus between

60-80ms and 140-190ms (Figure 3e, bright orange lines). We also found significant

thalamocortical iRS (figure 3f, bright cyan lines; between 20-40ms for MGB_L, 0-100ms

in MGB_{NL}, from 20-120ms in AC_L and between 40-100ms in AC_{NL}).

When we compared the two groups, the analysis revealed that MK-801 produced a

significant enhancement of iMM and iPE at both AC subdivisions ($p<0.000$ for iMM

between 60-190ms in both AC; and $p<0.05$ for iPE ranging between 100 and 190ms in

AC_L and between 60-190ms in AC_{NL}; white horizontal lines in Figures 3 d-f). By

contrast, iRS was affected more in the MGB ($p<0.000$ between 5-35ms in MGB_L;

$p<0.000$ between 40-110ms in MGB_{NL}; $p<0.05$ between 60-130ms in AC_L; and $p<0.05$

at 80ms in AC_{NL}; white horizontal lines in Figure 3f). Thus, MK-801 produces an

increase of iMM and iPE mostly in the late time window in AC, while iRS is much

affected in the MGB.

MK-801 affects the dynamics of adaptation. Since MK-801 lowered and flattened

responses to STD tones across the response window, we sought to assess the dynamics

and the time course of adaptation (figure 4a). Results show that the control group (light

gray arrows) exhibit a hierarchical timing for adaptation responses, becoming faster in

higher order areas (from top to down, responses reach the half of the initial values at the

fourth, ninth, twelfth and fourteenth standard tone, respectively). By contrast, results

from the MK-801 group exhibited much faster adaptation dynamics (figure 4b; 50% of

the initial response occurred at the third and second standard tones in MGB and AC,

respectively; b values for control group: $MGB_L=-0.1769$, $MGB_{NL}=-0.4174$, $AC_L=-$

0.6824 and $AC_{NL}=-1.175$; and for MK-801 group: $MGB_L=-0.8499$, $MGB_{NL}=-0.8853$,

$AC_L=-1.712$ and $AC_{NL}=-1.418$).

These data reveal that MK-801 alters the timing across the hierarchical organization of

the auditory system, resulting in the lemniscal thalamus having almost the same

adaptation velocity as the non-lemniscal cortex (arrows in Figure 4b). Furthermore,

MK-801 reduces (almost by half) the steady-state plateau in the AC (dotted lines in

Figure 4b; c values for control group: $MGB_L=0.0776$, $MGB_{NL}=0.2908$, $AC_L=0.6084$

and AC$_{NL}$=0.7740; and for MK-801 group: MGB$_L$=0.1428, MGB$_{NL}$=0.2884,

AC$_L$=0.3523 and AC$_{NL}$=0.3834).

All these results together support the idea that MK-801 produces a differential effect on

adaptation and deviance detection along the thalamocortical axis, providing new

evidence of a change in the firing pattern and temporal responses at single neuron level.

Delayed and broader larger-scaled LFP responses. Next, we wanted to check if the

single unit responses correlated with larger-scale measurements of neuronal activity.

The analysis of local field potentials (LFP) revealed that MK-801 produced significant

changes in MGB$_{NL}$ and AC (both in the lemniscal and non-lemniscal portions) for the

deviant, standard and cascade LFPs (Figure 5a-c), such that they exhibited broader and

longer responses for DEV-LFP and CAS-LFP in the auditory cortex, while the

waveforms of these LFPs were shifted in latency for the MGB$_{NL}$ due to a progressive

delay of N1, P1 and N2 (note that this terminology refers to the first negative peak, first

positive peak and second negative peak), showing delayed peaks of 8, 14 and 57ms for

DEV-LFP and 6, 26 and 45ms delay for CAS-LFP in N1, P1 and N2, respectively

(DEV-LFP: N1 peak for MK-801= -6.6μV at 20ms and control= -1.5μV at 12ms; P1

peak for MK-801=6.9μV at 41ms and control=6.8μV at 28ms; finally, N2 peak for MK-

801= -5.4μV at 102ms and control= -10.2μV at 45ms. CAS-LFP: N1 peak for MK-

801= -6.5μV at 18ms and control= -1.2μV at 12ms; P1 peak for MK-801=5.1μV at

53ms and control= 10.6μV at 27ms; finally, N2 peak for MK-801= -5.0μV at 91ms and

control= -10.3μV at 45ms).

Similarly, we also sought significant LFP signals for each computed index (Figure 5d-

f). The horizontal colored lines highlight the time at which significant deflections occur

to each index-LFP for control and MK-801 groups independently (light and bright

horizontal lines, respectively). Additionally, we compared these LFP indices between

groups. The analysis of the MM-LFP shows that MK-801 elicited stronger and broader

deflections within all regions (horizontal bright purple lines; Figure 5d) and abolished

the late negative component (N2) in the AC (MGB_L: N2 = 114-157ms; MGB_{NL}: N1 =

12-21ms, P1 = 32-63ms and N2 = 75-135ms; AC_L: N1 = 10-57ms and P1 = 60-147ms;

AC_{NL} N1 = 20-53 and P1 = 60-144ms). Our data also demonstrate that MK-801

produced a higher MM-LFP for virtually the whole LFP response within MGB_{NL} and

both AC, while no differences occurred in MGB_L.

Similar to the spike population analysis, and considering that the PE-LFP and RS-LFP

both contribute to the MM-LFP, we also wanted to understand how MK-801 shapes the

LFP for prediction error and repetition suppression. In response to MK-801, the PE-LFP

waveform was reduced at the early component of the MGB_{NL}, while it was increased

and delayed for the AC (orange horizontal lines in Figure 5e). Moreover, MK-801 also

abolished the N2 deflection (MGB_{NL}: N1 = 99-146ms; AC_L: N1 = 30-65ms and P1 =

87-180ms; AC_{NL} N1 = 30-67 and P1 = 106-180ms). When PE-LFP was compared

between groups, we only found differences in AC, mainly at the early (50-70ms) and

late components (120-180ms). In other words, the lemniscal thalamus does not exhibit

deviance detection, neither at the single neuron level nor at large-scale responses. Hence

PE-LFP confirm single unit population data, where MK-801 produced greater levels of

deviance detection in the auditory cortex (figure 2e).

Finally, MK-801 had similar effects on RS-LFP to those described above for MM-LFP

and PE-LFP, eliciting broader and larger waveforms for MGB_{NL} and AC (Figure 5f;

MGB_{NL}: N1 = 10-28ms, P1 = 34-63ms and N2 = 73-108ms; AC_L: N1 = 10-55ms, P1 =

67-140ms and N2 = 148-180ms; AC_{NL}: N1 = 10-51, P1 = 55-132ms and N2 = 141-

180ms). When differences between groups are considered, the non-lemniscal thalamus

exhibited a shift in the waveform between 15-100ms, while for the cortex, responses

over virtually the whole temporal window were increased by MK-801.

DISCUSSION

In this study, we demonstrate that the neuronal index derived from single cell

recordings of mismatch is profoundly affected along the auditory thalamocortical

system in rats treated acutely with a low dose of the NMDA-R antagonist, MK-801.

Importantly, we also reveal that the two elements that make up the index of mismatch

negativity, i.e., repetition suppression and prediction error, are differentially affected by

MK-801 in single neurons at auditory thalamus and cortex. MK-801 increases repetition

suppression in thalamus and prediction error in cortex. The increase in repetition

suppression is more prominent in lemniscal areas of the thalamus, while the increase in

prediction error is more evident in the non-lemniscal areas of cortex. Furthermore, our

results demonstrate that MK-801 alters the dynamics of neuronal adaptation along the

thalamocortical axis, becoming faster and stronger especially at thalamic level. The

pattern of results from single unit data were confirmed by recordings of large-scale

responses, LFPs, as the latter exhibit delayed and broader deflections. In summary, our

work demonstrates that the MK-801 increase of the neuronal mismatch in the auditory

cortex 60ms after stimulus onset is due to the combined effect of an increment in the

sustained responses to deviant tones and a decrement to standard tones. It should be

noted that, in contrast to most previous studies using recording procedures to study

neuronal population activity in rodents such as LFPs or EEG via skull screws, we have

recorded single-unit activity, a technique that has a much higher resolution level at the

single cell level for revealing patterns of activity underpinning mismatch responses.

It is well established that NMDA-R plays a fundamental role in neuronal

plasticity, controlling long-term potentiation and depression (Blanke & VanDongen,

2009). Further, it is generally accepted that human MMN is reduced after NMDA-R

antagonist treatments because NMDA-R antagonist blocks synaptic plasticity,

precluding the formation of a memory trace for the standard tones (Todd *et al.*, 2013).

As we have seen in our results, MK-801 reduces responses to standard tones thus

increasing repetition suppression.

Although this finding supports the hypothesis that NMDA-R antagonists alter

sensory-memory formation (Auksztulewicz & Friston, 2016), the findings that low dose

(0.1 mg/kg) MK-801 treatment produces a significant increment in the response to the

deviant tones, in prediction error and hence, an increment in the neuronal mismatch, are

in the opposite direction to expected. It is clear that the role of NMDA-R in the

generation of MMN is considerably more complex than thought (Harms *et al.*, 2018).

There have been suggestions in the literature of precedents for our observations. Even

considering that MK-801 has 160 times the affinity of ketamine to NMDA-R,

necessitating higher ketamine doses for similar drug effect (Schuelert *et al.*, 2018), our

results conform with those that report an increment in amplitude and latencies to deviant

responses after the acute ketamine treatment in rats (Ahnaou *et al.*, 2017) and with a

sub-anaesthetic dose of ketamine in healthy humans producing larger N100 to deviant

tones but not MMN (Oranje *et al.*, 2000). Interestingly, a dose response study of the

MK-801 effects on MMN-like responses in male rats showed that while a high dose

(0.5mg/kg) reduced late deviance detection (around 55ms), a medium dose (0.3mg/kg)

significantly enhanced early deviance detection effects (at about 13 ms) and some

evidence of enhanced late effects although not significantly (Harms *et al.*, 2018). We

used a single dose of 0.1mg/kg in female rats, as it has been demonstrated that females

are more sensitive to MK-801 than males (Andine *et al.*, 1999) and that this dose is

enough to induce behavioral effects (Meehan *et al.*, 2017). Importantly, memantine, a

low affinity uncompetitive agonist of NMDA-R, has been shown to (i) increase the

duration of rodent MMN-like responses (Tikhonravov *et al.*, 2010), (ii) increase MMN

amplitude in healthy individuals (Korostenskaja *et al.*, 2007), and (iii) in persons with

schizophrenia (Swerdlow *et al.*, 2016).

The memantine results suggest an interpretation of our findings in terms of the

mechanisms underpinning synaptic plasticity (Slutsky *et al.*, 2004). Partial blockade of

NMDA-R channels (such as mediated by memantine, or low dose MK-801) is also

likely to reduce background calcium flux resulting in homeostatic upregulation of

NR2B-containing NMDA-Rs leading in turn to the conversion of synapses to a plastic

state. That is, while these drugs reduce calcium influx during uncorrelated activity,

there is increased calcium influx during correlated activity (produced by physiological

stimuli), increased signal to noise, facilitated transmission and increased plasticity

(Slutsky *et al.*, 2010).

Other characteristics of the neuronal mechanisms and microcircuitry involving

the glutamate NMDA-R system are relevant to the effects we have observed on the

neuronal mismatch after the MK-801 treatment. NMDA-R are located, not only at

postsynaptic and presynaptic sites in excitatory neurons, but they are also found at

GABAergic inhibitory interneurons in neocortex (DeBiasi *et al.*, 1996). MK-801 has

demonstrated a preferential regulation of the firing rate of cortical GABA interneurons,

increasing the firing rate of the majority of pyramidal neurons (Homayoun &

Moghaddam, 2007) and therefore producing an imbalance in the excitatory/inhibitory

networks in the cortices (Bygrave *et al.*, 2016; Okada *et al.*, 2019). It is well known that

cortical GABAergic interneurons differentially amplify stimulus-specific adaptation (a

similar phenomenon to iMM) in excitatory pyramidal neurons in auditory cortex (Chen

et al., 2015). Moreover, a model of a mutually coupled excitatory/inhibitory network

can explain distinct mechanisms that allow cortical inhibitory neurons to enhance the

brain's sensitivity to deviant or unexpected sounds (Natan *et al.*, 2015). Further MK-

801 would alter the tonic inhibitory control of NMDA-R in cortical areas leading to the

activation of pyramidal neurons by subsequent deviant tones.

The increased repetition suppression we observed in the medial geniculate body

could also be by altered excitatory/inhibitory balance. Although the rat MGB lacks

GABAergic neurons, it receives GABAergic input from the thalamic reticular nucleus

(TRN) and the inferior colliculus (Malmierca, 2003). The latter is a source of bottom-up

inhibitory influences while the TRN provides the MGB with an indirect and inhibitory

feedback activation from AC (Bartlett, 2013). Cortical stimulation hyperpolarizes TRN

neurons and increases their inhibitory output to the MGB (Crabtree *et al.*, 2013) and

furthermore, TRN has been demonstrated to profoundly influence SSA in the MGB (Yu

et al., 2009). Changes in the thalamocortical neuronal firing pattern of thalamic neurons

into bursts have been suggested to provide an alerting signal to the cortex to enhance

stimulus detection (Hu & Agmon, 2016). Overall our results match the general concept

that when the system is adapted, it is more sensitive to detect changes in the

environment (Musall *et al.*, 2014), where a stronger thalamic repetition suppression (or

inhibition) support the increase in the prediction error signals (excitatory) at cortical

level, or vice versa. It would be interesting to test whether thalamic repetition

suppression is correlated with cortical prediction error signals, but this question awaits

future experiments.

Our study is important because it has revealed the involvement of two basic

mechanisms, i.e., repetition suppression and prediction error; and two different

pathways, lemniscal and non-lemniscal, underlying the neuronal mismatch in the

thalamocortical hierarchy. Predictive coding theory proposes that the brain constantly

tries to minimize the discrepancy between actual sensory input and internal

representations of the environment (Friston, 2005). What is new in our data is the

critical importance of the hierarchical organization of the auditory system in sharing the

'responsibility' for generating the representation and detecting the discrepancy, largely

attributable to thalamic and cortical processes. However, our data provide evidence that

the NMDA-synaptic plasticity and MMN relationship is not as simple as previously

surmised from human studies. Admittedly, we have only tackled the functional role of

the NMDA-R under a particular experimental manipulation and we cannot exclude the

possibility that larger doses of MK-801 would have generated different results. It is

also well known that other neuromodulatory systems such as the dopaminergic,

cholinergic and/or cannabinoid systems may be altered and interact with the NMDA-Rs

in normal brain function (Ayala *et al.*, 2016; Valdés-Baizabal *et al.*, 2017) as well as in

schizophrenia (Howes & Kaar, 2018; Lucatch *et al.*, 2018; Parr & Friston, 2018; Okada

et al., 2019). What are the implications of our findings for schizophrenia? Reduced

MMN is associated with poor global functioning (Light & Braff, 2005) and cognitive

deficits (Baldeweg *et al.*, 2004) in schizophrenia. Hence, if a safe drug were available

that targeted the relevant NMDA-R subunit, and facilitated neuroplasticity as indexed

by increased MMN, it offers opportunities for interventions to remediate cognitive

deficits that are a core feature of schizophrenia (Green *et al.*, 2000). Memantine which

has been shown to increase MMN amplitude in healthy individuals and in schizophrenia

has been used as an adjunctive therapy in schizophrenia for some time to improve

cognition in particular. While effects of adjunctive therapy are small, recent meta-

analysis suggests that there are improvements in global measures of cognition, but

improvements in more sensitive composite cognitive test scores have not been observed

(Kishi *et al.*, 2018). To date, there have been no attempts to utilize MMN enhancement

to memantine as an index of increased neuroplasticity that could be exploited in

remediation studies. Interestingly, both the moderate affinity antagonist, memantine,

and high affinity antagonist, MK-801, bind to the NR2B subunit of the NMDA-R at

very similar binding locations (Song *et al.*, 2018) but only memantine has been

approved for use in humans given evidence of neurotoxic effects of MK-801 in humans

(Olney *et al.*, 1989). One avenue of future research is the development of safe

compounds for human use that target similar binding locations to memantine and MK-

801.

Data availability. The data that support the findings of this study are available from the

corresponding author on reasonable request.

Acknowledgements

We thank Drs. Javier Nieto-Diego and David Perez-Gonzalez for their technical support

and constructive criticisms; and Drs. Nashat Abumaria and Juanita Todd for advice and

constructive comments on the discussion. We also thank Ms. Marianny Janiree Pernía

Rosales for helping with the histological photomicrography and Mr. Antonio Rivas

Cornejo for the technical support.

Funding

Financial support was kindly provided by the Spanish MINECO (SAF2016-75803-P)

and JCYL (SA023P17) to MSM. GGP held a fellowship from the Spanish MINECO

(BES-2014-069113) and CVB held a fellowship from Mexican CONACyT (216652).

LH was supported by an NHMRC project grant: APP1109283.

Competing interest.

The authors report no competing interests.

REFERENCES

Ahnaou, A., Huysmans, H., Biermans, R., Manyakov, N. V., & Drinkenburg, W.H.I.M.

(2017) Ketamine: differential neurophysiological dynamics in functional networks

in the rat brain. *Transl. Psychiatry*, **7**, e1237.

Andine, P., Widermark, N., Axelsson, R., Nyberg, G., Olofsson, U., Martensson, E., &

Sandberg, M. (1999) Characterization of MK-801-Induced Behavior as a Putative

Rat Model of Psychosis. *J. Pharmacol. Exp. Ther.*, **290**, 1393–1408.

Antunes, F.M., Nelken, I., Covey, E., & Malmierca, M.S. (2010) Stimulus-Specific

Adaptation in the Auditory Thalamus of the Anesthetized Rat. *PLoS One*, **5**,

e14071.

Auksztulewicz, R. & Friston, K. (2016) Repetition suppression and its contextual

determinants in predictive coding. *Cortex.*, **80**, 125–140.

Ayala, Y.A., Pérez-González, D., & Malmierca, M.S. (2016) Stimulus-specific

adaptation in the inferior colliculus: The role of excitatory, inhibitory and

modulatory inputs. *Biol. Psychol.*, **116**, 10–22.

Baldeweg, T., Klugman, A., Gruzelier, J., & Hirsch, S.R. (2004) Mismatch negativity

potentials and cognitive impairment in schizophrenia. *Schizophr. Res.*, **69**, 203–

217.

Bartlett, E.L. (2013) The organization and physiology of the auditory thalamus and its

role in processing acoustic features important for speech perception. *Brain Lang.*,

126, 29–48.

Bendixen, A., Roeber, U., & Schröger, E. (2007) Regularity extraction and application

in dynamic auditory stimulus sequences. *J. Cogn. Neurosci.*, **19**, 1664–1677.

Blanke, M.L. & VanDongen, A.M.J. (2009) *Activation Mechanisms of the NMDA*

Receptor (Chapter 13), Biology of the NMDA Receptor. CRC Press/Taylor &

Francis.

Bygrave, A.M., Masiulis, S., Nicholson, E., Berkemann, M., Barkus, C., Sprengel, R.,

Harrison, P.J., Kullmann, D.M., Bannerman, D.M., & Kätzel, D. (2016) Knockout

of NMDA-receptors from parvalbumin interneurons sensitizes to schizophrenia-

related deficits induced by MK-801. *Transl. Psychiatry*, **6**, e778.

Carbajal, G. V & Malmierca, M.S. (2018) The Neuronal Basis of Predictive Coding

Along the Auditory Pathway: From the Subcortical Roots to Cortical Deviance

Detection. *Trends Hear.*,.

Chen, I.-W.I.-W., Helmchen, F., & Lutcke, H. (2015) Specific Early and Late Oddball-

Evoked Responses in Excitatory and Inhibitory Neurons of Mouse Auditory

Cortex. *J. Neurosci.*, **35**, 12560–12573.

Crabtree, J.W., Lodge, D., Bashir, Z.I., & Isaac, J.T.R. (2013) GABAA, NMDA and

mGlu2 receptors tonically regulate inhibition and excitation in the thalamic

reticular nucleus. *Eur. J. Neurosci.*, **37**, 850–859.

DeBiasi, S., Minelli, A., Melone, M., Conti, F., & DeBiasi, S., Minelli, A., Melone, M.,

& Conti, F. (1996) Presynaptic NMDA receptors in the neocortex are both auto-

and heteroreceptors. *Neuroreport*, **7**, 2773–2776.

Duque, D. & Malmierca, M.S. (2015) Stimulus-specific adaptation in the inferior

colliculus of the mouse: anesthesia and spontaneous activity effects. *Brain Struct.*

706 *Funct.*, **220**, 3385–3398.

707 Ehrlichman, R.S., Maxwell, C.R., Majumdar, S., & Siegel, S.J. (2008) Deviance-

708 elicited changes in event-related potentials are attenuated by ketamine in mice. *J.*

709 *Cogn. Neurosci.*, **20**, 1403–1414.

710 Featherstone, R.E., Melnychenko, O., & Siegel, S.J. (2018) Mismatch negativity in

711 preclinical models of schizophrenia. *Schizophr. Res.*, **191**, 35–42.

712 Friston, K. (2005) A theory of cortical responses. *Philos. Trans. R. Soc. B Biol. Sci.*,

713 **360**, 815–836.

714 Green, M.F., Kern, R.S., Braff, D.L., & Mintz, J. (2000) Neurocognitive deficits and

715 functional outcome in schizophrenia: are we measuring the "right

716 stuff"? *Schizophr. Bull.*, **26**, 119–136.

717 Harms, L. (2016) Mismatch responses and deviance detection in N-methyl-D-aspartate

718 (NMDA) receptor hypofunction and developmental models of schizophrenia. *Biol.*

719 *Psychol.*, **116**, 75–81.

720 Harms, L., Fulham, W.R., Todd, J., Budd, T.W., Hunter, M., Meehan, C., Penttonen,

721 M., Schall, U., Zavitsanou, K., Hodgson, D.M., & Michie, P.T. (2014) Mismatch

722 negativity (MMN) in freely-moving rats with several experimental controls. *PLoS*

723 *One*, **9**, e110892.

724 Harms, L., Fulham, W.R., Todd, J., Meehan, C., Schall, U., Hodgson, D.M., & Michie,

725 P.T. (2018) Late deviance detection in rats is reduced, while early deviance

726 detection is augmented by the NMDA receptor antagonist MK-801. *Schizophr.*

727 *Res.*, **191**, 43–50.

Homayoun, H. & Moghaddam, B. (2007) NMDA Receptor Hypofunction Produces

Opposite Effects on Prefrontal Cortex Interneurons and Pyramidal Neurons. *J.*

Neurosci., **27**, 11496–11500.

Howes, O.H. & Kaar, S.J. (2018) Antipsychotic drugs: challenges and future directions.

World Psychiatry, **17**, 170–171.

Hu, H. & Agmon, A. (2016) Differential Excitation of Distally versus Proximally

Targeting Cortical Interneurons by Unitary Thalamocortical Bursts. *J. Neurosci.*,

36, 6906–6916.

Ishishita, Y., Kunii, N., Shimada, S., Ibayashi, K., Tada, M., Kirihara, K., Kawai, K.,

Uka, T., Kasai, K., & Saito, N. (2018) Deviance detection is the dominant

component of auditory contextual processing in the lateral superior temporal gyrus:

A human ECoG study. *Hum. Brain Mapp.*, **40**, 1184–1194.

Javitt, D.C., Steinschneider, M., Schroeder, C.E., & Arezzo, J.C. (1996) Role of cortical

N-methyl-D-aspartate receptors in auditory sensory memory and mismatch

negativity generation: implications for schizophrenia. *Proc. Natl. Acad. Sci. U. S.*

A., **93**, 11962–11967.

Kishi, T., Ikuta, T., Oya, K., Matsunaga, S., Matsuda, Y., & Iwata, N. (2018) Anti-

dementia Drugs for Psychopathology and Cognitive Impairment in Schizophrenia:

A Systematic Review and Meta-analysis. *Int. J. Neuropsychopharmacol.*, **21**, 748.

Korostenskaja, M., Nikulin, V. V., Kičić, D., Nikulina, A. V., & Kähkönen, S. (2007)

Effects of NMDA receptor antagonist memantine on mismatch negativity. *Brain*

Res. Bull., **72**, 275–283.

Krystal, J.H., Perry, E.B., Gueorguieva, R., Belger, A., Madonick, S.H., Abi-Dargham,

751    A., Cooper, T.B., MacDougall, L., Abi-Saab, W., & Cyril D'Souza, D. (2005)

752    Comparative and interactive human psychopharmacologic effects of ketamine and

753    amphetamine: Implications for glutamatergic and dopaminergic model psychoses

754    and cognitive function. *Arch. Gen. Psychiatry*, **62**, 985–995.

755 Kurkela, J.L.O., Lipponen, A., Kyläheiko, I., & Astikainen, P. (2018)

756    Electrophysiological evidence of memory-based detection of auditory regularity

757    violations in anesthetized mice. *Sci. Rep.*, **8**, 3027.

758 Light, G.A. & Braff, D.L. (2005) Mismatch negativity deficits are associated with poor

759    functioning in schizophrenia patients. *Arch. Gen. Psychiatry*, **62**, 127–136.

760 Lucatch, A.M., Coles, A.S., Hill, K.P., & George, T.P. (2018) Cannabis and Mood

761    Disorders. *Curr. Addict. Reports*, **5**, 336–345.

762 Malmierca, M.S. (2003) The Structure and Physiology of the Rat Auditory System: An

763    Overview. pp. 147–211.

764 Meehan, C., Harms, L., Frost, J.D., Barreto, R., Todd, J., Schall, U., Shannon Weickert,

765    C., Zavitsanou, K., Michie, P.T., & Hodgson, D.M. (2017) Effects of immune

766    activation during early or late gestation on schizophrenia-related behaviour in adult

767    rat offspring. *Brain. Behav. Immun.*, **63**, 8–20.

768 Michie, P.T., Malmierca, M.S., Harms, L., & Todd, J. (2016) Understanding the

769    neurobiology of MMN and its reduction in schizophrenia. *Biol. Psychol.*, **116**, 1–3.

770 Musall, S., von der Behrens, W., Mayrhofer, J.M., Weber, B., Helmchen, F., & Haiss,

771    F. (2014) Tactile frequency discrimination is enhanced by circumventing

neocortical adaptation. *Nat. Neurosci.*, **17**, 1567–1573.

Näätänen, R., Gaillard, A.W.K.W.K., & Mäntysalo, S. (1978) Early selective-attention

effect on evoked potential reinterpreted. *Acta Psychol. (Amst).*, **42**, 313–329.

Nakamura, T., Michie, P.T., Fulham, W.R., Todd, J., Budd, T.W., Schall, U., Hunter,

M., & Hodgson, D.M. (2011) Epidural Auditory Event-Related Potentials in the

Rat to Frequency and duration Deviants: Evidence of Mismatch Negativity? *Front.*

Psychol., **2**, 367.

Natan, R.G., Briguglio, J.J., Mwilambwe-Tshilobo, L., Jones, S.I., Aizenberg, M.,

Goldberg, E.M., & Geffen, M.N. (2015) Complementary control of sensory

adaptation by two types of cortical interneurons. *Elife*, **4**.

Nieto-Diego, J. & Malmierca, M.S. (2016) Topographic Distribution of Stimulus-

Specific Adaptation across Auditory Cortical Fields in the Anesthetized Rat. *PLOS*

Biol., **14**, e1002397.

Okada, M., Fukuyama, K., Kawano, Y., Shiroyama, T., & Ueda, Y. (2019) Memantine

protects thalamocortical hyper-glutamatergic transmission induced by NMDA

receptor antagonism via activation of system xc. *Pharmacol. Res. Perspect.*, **7**,

e00457.

Olney, J., Labruyere, J., & Price, M. (1989) Pathological cahnges induced in

cerebrocortical neurons by phencyclidine and related drugs. *Science (80-.).*, **244**,

1360–1362.

Opitz, B., Schröger, E., & Von Cramon, D.Y. (2005) Sensory and cognitive

mechanisms for preattentive change detection in auditory cortex. *Eur. J. Neurosci.*,

794 **21**, 531–535.

795 Oranje, B., Berckel, B. van, Kemner, C., Ree, J. van, Kahn, R., & Verbaten, M. (2000)

796  The Effects of a Sub-Anaesthetic Dose of Ketamine on Human Selective

797  Attention. *Neuropsychopharmacology*, **22**, 293–302.

798 Parr, T. & Friston, K.J. (2018) The Anatomy of Inference: Generative Models and Brain

799  Structure. *Front. Comput. Neurosci.*, **12**, 90.

800 Parras, G.G., Nieto-Diego, J., Carbajal, G. V., Valdés-Baizabal, C., Escera, C., &

801  Malmierca, M.S. (2017) Neurons along the auditory pathway exhibit a hierarchical

802  organization of prediction error. *Nat. Commun.*, **8**, 2148.

803 Paxinos, G. & Watson, C. (2013) *The Rat Brain in Stereotaxic Coordinates : Hard*

804  *Cover Edition.* Elsevier Science.

805 Pérez-González, D., Hernández, O., Covey, E., & Malmierca, M.S. (2012) GABA A-

806  mediated inhibition modulates stimulus-specific adaptation in the inferior

807  colliculus. *PLoS One*, **7**.

808 Randeniya, R., Oestreich, L.K.L., & Garrido, M.I. (2018) Sensory prediction errors in

809  the continuum of psychosis. *Schizophr. Res.*, **191**, 109–122.

810 Ruhnau, P., Herrmann, B., & Schröger, E. (2012) Clinical Neurophysiology Finding the

811  right control : The mismatch negativity under investigation. *Clin. Neurophysiol.*,

812  **123**, 507–512.

813 Schuelert, N., Dorner-Ciossek, C., Brendel, M., & Rosenbrock, H. (2018) A

814  comprehensive analysis of auditory event-related potentials and network

815  oscillations in an NMDA receptor antagonist mouse model using a novel wireless

816  recording technology. *Physiol. Rep.*, **6**, e13782.

817 Siegel, S.J., Talpos, J.C., & Geyer, M.A. (2013) Animal models and measures of

818  perceptual processing in schizophrenia. *Neurosci. Biobehav. Rev.*, **37**, 2092–2098.

819 Sivarao, D. V, Chen, P., Yang, Y., Li, Y.-W., Pieschl, R., & Ahlijanian, M.K. (2014)

820  NR2B Antagonist CP-101,606 Abolishes Pitch-Mediated Deviance Detection in

821  Awake Rats. *Front. psychiatry*, **5**, 96.

822 Slutsky, I., Abumaria, N., Wu, L.-J., Huang, C., Zhang, L., Li, B., Zhao, X.,

823  Govindarajan, A., Zhao, M.-G., Zhuo, M., Tonegawa, S., & Liu, G. (2010)

824  Enhancement of Learning and Memory by Elevating Brain Magnesium. *Neuron*,

825  **65**, 165–177.

826 Slutsky, I., Sadeghpour, S., Li, B., & Liu, G. (2004) Enhancement of Synaptic Plasticity

827  through Chronically Reduced Ca2+ Flux during Uncorrelated Activity. *Neuron*,

828  **44**, 835–849.

829 Song, X., Jensen, M.Ø., Jogini, V., Stein, R.A., Lee, C.-H., Mchaourab, H.S., Shaw,

830  D.E., & Gouaux, E. (2018) Mechanism of NMDA receptor channel block by MK-

831  801 and memantine. *Nature*, **556**, 515–519.

832 Swerdlow, N.R., Bhakta, S., Chou, H.-H., Talledo, J.A., Balvaneda, B., & Light, G.A.

833  (2016) Memantine Effects On Sensorimotor Gating and Mismatch Negativity in

834  Patients with Chronic Psychosis. *Neuropsychopharmacology*, **41**, 419–430.

835 Tikhonravov, D., Neuvonen, T., Pertovaara, A., Savioja, K., Ruusuvirta, T., Näätänen,

836  R., & Carlson, S. (2008) Effects of an NMDA-receptor antagonist MK-801 on an

837  MMN-like response recorded in anesthetized rats. *Brain Res.*, **1203**, 97–102.

Tikhonravov, D., Neuvonen, T., Pertovaara, A., Savioja, K., Ruusuvirta, T., Näätänen,

R., & Carlson, S. (2010) Dose-related effects of memantine on a mismatch

negativity-like response in anesthetized rats. *Neuroscience*, **167**, 1175–1182.

Todd, J., Harms, L., Schal l, U., & Michie, P.T. (2013) Mismatch negativity:

Translating the potential. *Front. Psychiatry*,.

Umbricht, D., Koller, R., Vollenweider, F.X., & Schmid, L. (2002) Mismatch negativity

predicts psychotic experiences induced by NMDA receptor antagonist in healthy

volunteers. *Biol. Psychiatry*, **51**, 400–406.

Umbricht, D., Schmid, L., Koller, R., Vollenweider, F.X., Hell, D., & Javitt, D.C.

(2000) Ketamine-induced deficits in auditory and visual context-dependent

processing in healthy volunteers: Implications for models of cognitive deficits in

schizophrenia. *Arch. Gen. Psychiatry*, **57**, 1139–1147.

Valdés-Baizabal, C., Parras, G.G., Ayala, Y.A., & Malmierca, M.S. (2017)

Endocannabinoid Modulation of Stimulus-Specific Adaptation in Inferior

Colliculus Neurons of the Rat. *Sci. Rep.*, **7**, 1–14.

Vezzani, A., Serafini, R., Stasi, M.A., Caccia, S., Conti, I., Tridico, R. V, & Samanin,

R. (1989) Kinetics of MK-801 and its effect on quinolinic acid-induced seizures

and neurotoxicity in rats. *J. Pharmacol. Exp. Ther.*, **249**, 278–283.

Wacongne, C. (2016) A predictive coding account of MMN reduction in schizophrenia.

Biol. Psychol., **116**, 68–74.

Yu, X., Xu, X., He, S., & He, J. (2009) Change detection by thalamic reticular neurons.

Nat. Neurosci., **12**, 1165–1170.

a) Frequency Response Area Example

b) Paradigms

Figure 1. Experimental design. a) Frequency response area example, with a representation of the ten selected tones to build the experimental paradigms. **b)** Stimulation sequences, the same tone could be presented in different experimental paradigms, thus we can compare same tone in different contexts to control adaptation and deviance detection; and conform the indices of neuronal mismatch (iMM), prediction error (iPE) and repetition suppression (iRS). Note that ascending and descending tones will be compared to the control ascending or descending, respectively. **c)** Sketch of summary results of mismatch responses for healthy and schizophrenia subjects under the classical analysis of mismatch. Second row decomposition of neuronal mismatch, under the assumption of predictive coding framework in healthy subjects, and the hypothetical decomposition of neuronal mismatch into prediction error and repetition suppression in schizophrenia.

Figure 2. **Single neuron spikes population analysis**. Results for firing rate analysis and their computed differences along the thalamocortical axis. **a)** Boxplot of median normalized responses for deviants (red), cascade (green) and standard (blue) for each group, control (light colors) and MK801 (bright colors), within each station and the statistical significance between groups (Wilcoxon signed-rank test, * $p<0.05$, ** $p<0.001$, *** $p<0.000$). **b-d)** Indices histograms displayed in a mirror-like manner for the two groups (controls upper and in light colors; MK801 under and in bright colors), showing the distribution of the three indexes for each neuronal response (ranging between -1 and +1, dotted lines indicate index=0). Vertical solid lines indicate their medians and the significant difference between groups is noted at the right of each histogram block. **e)** Median indices of Prediction Error (orange) and **d)** Repetition Suppression (blue), represented with respect to the baseline set by the cascade control (green line). Thereby, iPE upwards-positive while iRS is downwards-positive. Each median index corresponds to differences between normalized responses in a). Asterisks inside bars denote statistically significance of these indices against zero (Friedman test), while asterisks outside bars denote statistically significance between groups (Wilcoxon signed-rank test, * $p<0.05$, ** $p<0.001$, *** $p<0.000$).

Figure 3. Spike Density Function. Peristimulus time histogram along the thalamocortical axis. **a-c)** Averaged firing rate profiles for each condition as normalized spike-density function (light colors for control and bright color for MK801 group), and their respective differences (white dotted lines). Solid horizontal white lines represent the time in which the difference between groups is significant (two-sample t test $p<0.05$, Bonferroni corrected). **d-f)** Indices over time computed for 12 intervals (from -50 to 190ms) compared against zero (signed-rank test and FDR corrected for 12 comparisons; * $p<0.01$) for each group (light colors for control and bright color for MK801 group). Solid white lines denote differences between groups across time intervalss (two-sample t test for each of the 12-time windows, $p<0.05$).

a)

AC$_{NL}$

AC$_L$

MGB$_{NL}$

MGB$_L$

STD   CSC   DEV

MK-801

Control

Spikes per trial (average)

0   Time (s)   100

b)

STD

MK-801

Control

0   Time (s)   4

Figure 4. Time course for dynamical thalamocortical adaptation. **a)** Averaged time course for the stimulus played in relation to the time elapsed from the beginning of the sequence. **b)** The first fifteen standard stimuli showing the three parameters of the power low fitted: *a* initial average response; *b* adaptation velocity; and *c* the steady-state value (dotted lines) for each group. Arrows represent the 50% of the initial responses demonstrating faster adaptation in the MK801 group and the break down in the dynamical hierarchy of adaptation.

Figure 5. Local Field Potentials for each condition and their differences. a-c) Population grand-averaged LFP for each condition recorded (CAS, DEV and STD) within each group (controls and MK801). Grey panels under the main LFP representations shows the instantaneous p value (white trace) of corresponding stimulus condition LFP (critical threshold set at 0.05 represented as a horizontal dotted yellow line). The thick black horizontal bars in figure 5a-c highlights the time interval for which the LFP comparison between the control and MK801 groups is significant. **d-f)** Population grand-averaged LFP for and neuronal Mismatch (MM-LFP=LFP$_{DEV}$-LFP$_{STD}$), Prediction Error (PE-LFP=LFP$_{DEV}$-LFP$_{CAS}$), and Repetition Suppression (RS-LFP=LFP$_{STD}$-LFP$_{CAS}$) respectively Colored horizontal lines denote significative deflections (*t*-test, FDR corrected). Grey panels show the instantaneous p value (white trace) of corresponding stimulus condition LFP (critical threshold set at 0.05 represented as a horizontal dotted yellow line) and black horizontal lines the time interval in which MK801 and control are statistically different.

a) Medial Geniculate Body Histology

Dorsal division

Medial division

Ventral division

500 μm

Bregma -5.7

b) Auditory Cortex Tonotopic mapping

Ventral from bregma (mm)

Caudal from bregma (mm)

PAF

A1

AAF

VAF

SRAF

32 kHz
16 kHz
8 kHz
4 kHz
2 kHz
1 kHz

Figure 6. Anatomical recordings location. **a)** Photomicrography sample of a MGB Nissl-stained slice (10x), red arrows point the two electrolytic lesions. **b)** Example of localization all recordings made in the AC of one rat, each colored dot represent the characteristic frequency of each performed tract. A1: Primary Auditory Field; AAF: Anterior Auditory Field; VAF: Ventral Auditory Field; PAF: Posterior Auditory Field and SRAF: Suprarhinal Auditory Field.

	CONTROL				MK801			
	MGB		AC		MGB		AC	
	L	NL	L	NL	L	NL	L	NL
# Neurons	28	38	40	37	44	42	37	24
# Points/ required	111/72	240/224	295/87	309/29	156/81	228/544	163/21	94/7
DEV (spk)	0.7249	0.6805	0.9594	0.9694	0.5970	0.6723	1.0978	1.6450
STD (spk)	0.1500	0.1504	0.2363	0.2103	0.0338	0.0837	0.0906	0.1114
CSD (spk)	0.8292	0.5601	0.7719	0.5910	0.7329	0.7889	0.5913	0.5978
	Cascade vs. Many-Standard Controls analysis (median and Wilcoxon signed-rank values)							
Many-standard	0.8982	0.6499	0.8034	0.5237	0.7855	0.9966	0.5109	0.6861
Cascade	0.7625	0.5601	0.7720	0.5911	0.7326	0.7889	0.5913	0.5979
P value	*0.2231*	*0.1567*	*0.8266*	*0.3288*	*0.4103*	*0.0286*	*0.2543*	*0.0830*
	Friedman test analysis							
iMM	0.4240	0.5169	0.5283	0.6073	0.6075	0.5578	0.7734	0.8699
P value	1.01^{-09}	2.96^{-29}	7.52^{-51}	7.90^{-56}	6.96^{-24}	5.98^{-38}	1.63^{-41}	2.34^{-30}
iPE	-0.0950	0.0492	0.1221	0.2734	-0.0633	-0.0445	0.3723	0.5788
P value	*0.0130*	*0.0199*	*0.0012*	3.29^{-12}	*0.0092*	*0.5120*	1.88^{-07}	8.84^{-13}
iRS	0.5192	0.4678	0.4062	0.3339	0.6707	0.6023	0.4011	0.2910
P value	8.63^{-18}	5.55^{-19}	5.99^{-32}	1.68^{-18}	7.48^{-37}	9.89^{-42}	1.16^{-16}	1.68^{-05}

Table 1. **Spike population analysis** for each experimental group and auditory station independently: First row, number of recorded neurons; second row number of tested neuron/frequency combinations (points), along with estimated minimum sample size (of points) required for a statistical power (See Methods). Followed by median values for base-line corrected spike count (spikes) to the different conditions. Comparative analysis for control paradigms, median values and Wilcoxon signed-rank test values for each station and group. Median indices of neuronal mismatch (iMM), prediction error (iPE) and repetition suppression (iRS), and their corresponding *p* value.

Table 2: Firing Rate Comparisons

	MGB		AC	
	L	NL	L	NL
STD_control	0.1912	0.1717	0.1990	0.1856
STD_MK-801	0.0537	0.1047	0.0689	0.0547
P value	**<0.000**	**<0.000**	**<0.000**	**<0.000**
DEV_ control	0.6153	0.6887	0.7274	0.7930
DEV _ MK-801	0.6611	0.6620	0.8424	0.9247
P value	0.2309	0.2641	**<0.000**	**<0.000**
CAS_ control	0.7104	0.6395	0.6052	0.5196
CAS_ MK-801	0.7244	0.7063	0.4701	0.3458
P value	0.1169	**<0.000**	**<0.000**	**<0.000**
iMM_ control	0.3675	0.4897	0.5124	0.5781
iMM_ MK-801	0.5197	0.5221	0.7269	0.8088
P value	**<0.000**	**0.0409**	**<0.000**	**<0.000**
iPE_ control	-0.0670	0.0513	0.1060	0.2697
iPE_ MK-801	-0.0502	-0.0444	0.3592	0.5652
P value	0.9622	**0.0035**	**0.000**	**<0.000**
iRS_ control	0.5300	0.3672	0.3350	0.2935
iRSdrug	0.6812	0.5632	0.3514	0.2765
P value	**<0.000**	**<0.000**	0.5800	0.8934

16 **Table 2: Comparative analysis between control and MK-801 group.** Median spikes

17 to the former measures responses to standard (STD), deviant (DEV) and cascade (CAS)

18 tones, and their corresponding *p* value (ranksum test). Similarly, medians and their

19 associates *p* value for the index of mismatch (iMM), index of prediction error (iPE) and

20 index of repetition suppression (iRS).

21