Asymmetry in kinematic generalization between visual and passive lead-in movements are consistent with a forward model in the sensorimotor system

Abbreviated title: Spatiotemporal generalization of lead-in movements

Ian S. Howard[1], Sae Franklin[2] and David W. Franklin[3]

[1]Centre for Robotics and Neural Systems, School of Computing, Electronics and Mathematics, University of Plymouth, Plymouth, United Kingdom
[2]Institute for Cognitive Systems, Department of Electrical and Computer Engineering, Technical University of Munich, Germany
[3]Neuromuscular Diagnostics, Department of Sport and Health Sciences, Technical University of Munich, Germany

Correspondence should be sent to:
Ian S. Howard,
Centre for Robotics and Neural Systems, School of Computing, Electronics and Mathematics, University of Plymouth, Plymouth, United Kingdom

Email: ian.howard@plymouth.ac.uk
Telephone: 0044 1752 586324

Keywords: motor learning, context, transfer, dynamic learning, motor memories, human

Abstract

In our daily life we often make complex actions comprised of linked movements, such as reaching for a cup of coffee and bringing it to our mouth to drink. Recent work has highlighted the role of such linked movements in the formation of independent motor memories, affecting the learning rate and ability to learn opposing force fields. In these studies, distinct prior movements (lead-in movements) allow adaptation of opposing dynamics on the following movement. Purely visual or purely passive lead-in movements exhibit different angular generalization functions of this motor memory as the lead-in movements are modified, suggesting different neural representations. However, we currently have no understanding of how different movement kinematics (distance, speed or duration) affect this recall process and the formation of independent motor memories. Here we investigate such kinematic generalization for both passive and visual lead-in movements to probe their individual characteristics. After participants adapted to opposing force fields using training lead-in movements, the lead-in kinematics were modified on random trials to test generalization. For both visual and passive modalities, recalled compensation was sensitive to lead-in duration and peak speed, falling off away from the training condition. However, little reduction in force was found with increasing lead-in distance. Interestingly, asymmetric transfer between lead-in movement modalities was also observed, with partial transfer from passive to visual, but very little vice versa. Overall these tuning effects were stronger for passive compared to visual lead-ins demonstrating the difference in these sensory inputs in regulating motor memories. Our results suggest these effects are a consequence of state estimation, with differences across modalities reflecting their different levels of sensory uncertainty arising as a consequence of dissimilar feedback delays.

Introduction

Recent studies have highlighted key aspects for neural rehabilitation using robotic systems [1]. However, continual progress in this area depends on understanding the mechanisms of human sensorimotor learning in order to determine the optimal presentation of sensory information to improve the rate, retention and generalization of adaptation. Although adaptation is often studied on single movements in the laboratory, we rarely produce movements in isolation in everyday life. Rather, one movement often directly leads into another. For example, to catch a ball, we make use of visual motion information to estimate its state in order to plan and execute an interception movement. Thus, natural movements often follow directly from previous movements, or from visual motion.

This scenario can be investigated in the laboratory by using two-part
movements. Here, the first part consists of a lead-in movement from a start
location to an intermediate via point, which is followed closely in time by a
second movement to a final target location. Recent work has shown that closely
linking multiple movements together in time in this fashion reduces interference
in learning opposing tasks [2-5]. In particular, distinct past movements act like
a contextual cue, enabling adaptation to opposing viscous curl fields when the
adaptation movements are preceded by unique lead-in motions, each
associated with one of the dynamics [2]. This shows that motor learning and
recall depends not only on the current state of the arm during a movement, but
also on its preceding states. Interestingly, active, passive or visual lead-in
movements were all equally effective at reducing interference. This indicates
that sensory feedback relating to motion is sufficient to affect adaptation, even
when no active movement is involved. The contextual effect of this prior
movement disappears as the time between lead-in and adaptation movements
exceeds about half a second, indicating that the representation of past state
decays quickly over time. This suggests a strong link between the
representation of state and the theory of neural population dynamics [6,7].

Dynamic adaptation to a single force field occurs locally; after training in a
specific movement, the recall of predictive compensation decreases as the
movement angle [8-11] or distance [12-14] deviates from the training condition.
The Gaussian-like angular generalization observed in these studies has also
be found for lead-in movements, with different lead-in modalities exhibiting
different characteristics, both in terms of their absolute level of influence, but
also in their sharpness of tuning. In particular both active [15] and passive lead-
in movements [10] show narrower and deeper tuning than visual lead-in
movements [11].

Interference studies have been widely adopted to investigate contextual effects
on motor learning, and to examine if contextual cues can assist in the learning
of opposing dynamics [14,16-21]. Such interference paradigms are more
sensitive to generalization effects of contextual cues than single field
paradigms, and have been used effectively to examine the angular
generalization characteristics of lead-in movements [10,11]. Using these
paradigms it has been possible to extract features of the neural basis functions
underlying dynamical adaptation, allowing for the development of simple
computational models [4]. However, we still lack basic information on the
generalization features of lead-in movements for different kinematics such as
duration or distance.

Here, we first characterize the generalization of passive and visual lead-in
movements across different kinematics using an interference paradigm. In two

separate experiments, we examine generalization across distance and duration (and the dependent variable of speed) of passive and visual lead-in movements. Second, in order to gain insight into any commonality between the neural resources employed in passive and visual lead-in movements, we also investigate how adaptation transfers between these two different lead-in modalities.

Methods

Experimental Design

Subjects. Sixteen human participants were randomly allocated to two experimental groups that each performed one experiment. Eight participants (7 female, aged 24.8 ± 5.0 years, mean ± sd) performed the passive lead-in experiment. Eight further participants (6 female; aged 27.4 ± 6.7 years) participated in the visual lead-in experiment. All participants were right handed according to the Edinburgh handedness questionnaire [22], and naïve to the aims of the study. All participants provided written informed consent to the protocol before participating in the experiment, which had been approved by the University of Cambridge Ethics Committee. The methods were carried out in accordance with the approved guidelines. Some of the data (visual lead-in experiment) was previously presented in a conference paper [23].

Apparatus. Experiments were performed using a vBOT planar robotic manipulandum and its associated virtual reality system [24]. Handle position is measured using optical encoders sampled at 1000 Hz, and motors operating under torque control allow the application of end-point forces. A force transducer (Nano 25; ATI), mounted under the handle, measures the applied forces, and its output signals were low-pass filtered at 500 Hz using analogue 4th pole Bessel filters prior to digitization. To reduce body movement, participants were seated in a sturdy chair in front of the apparatus and firmly strapped against the backrest with a four-point seatbelt. During an experiment, participants grasped the robot handle in their right hand while their right forearm was supported by an air sled, constraining arm movement to the horizontal plane. Participants could not view their hand directly. Instead veridical visual feedback was used to overlay images of the starting location, via point, final target, (all 1.25 cm radius disks) and a hand cursor (0.5 cm radius red disk) using the virtual reality system. This ensured that the visual cursor appeared to the participant in the same plane and at the same location as their hand. Data was collected at 1000 Hz and logged to disk for offline analysis using Matlab (Matlab, The MathWorks Inc., Natick, MA, USA).

157   *Force Fields.* In the adaptation movement, participants performed reaching
158   movements either in a null field condition, a velocity-dependent curl force field
159   [14], or a mechanical channel [25]. The curl force field was implemented as:
160

161
$$\begin{bmatrix} F_x \\ F_y \end{bmatrix} = k \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} \dot{x} \\ \dot{y} \end{bmatrix} \tag{1}$$

where the field constant k was set to a value of ±16 Nm^{-1}s, and the sign
determines the direction (CW or CCW) of the force-field. Each participant
experienced both force field directions. The direction of the force field was
always associated with a specific direction of a prior contextual movement. The
relationship between the contextual movement direction and curl field direction
(CW/CCW) was counterbalanced across participants.

Mechanical channel trials were implemented using a spring constant of 6,000
Nm^{-1} and a damping constant of 30 Nm^{-1}s perpendicular to the direction of
motion throughout the movement between the central location and the final
target. Channel trials were only produced on the movements to the 0° target
with corresponding lead-in movements starting at 135° or 225°, and never
presented on consecutive trials.

Protocol

Two separate experiments were performed to examine the generalization of
the learning associated with one contextual movement to other contextual
movements with different kinematic profiles (within the same modality), as well
as the transfer of learning between passive and visual lead-in conditions
(across the modalities).

After an initial pre-learning session in a null field, participants were exposed to
the curl force fields (learning phase). Channel trials were used to examine
adaptation to the novel dynamics, in which the lead-in movement duration,
speed and distance were varied. In addition, the modality of the lead-in
movement was occasionally changed to examine transfer. The trial parameters
for both experiments are shown in Table 1 and the kinematics of the lead-in
movements can be seen in Fig 1. On these trials, the lead-in movement was
chosen from one of 15 different movements with distances ranging from 3 cm
to 20 cm and durations ranging between 210 ms to 1400 ms.

Trial Organization. All trials consisted of a two-part movement: An initial lead-
in movement followed directly by an adaptation movement. The first part was a
contextual lead-in movement from a starting location to a central via point. This
contextual lead-in movement was 10 cm in length during all null and force field

training conditions. The second part was an 18 cm adaptation movement to the final target. The participants only experienced a force field or channel trial during this adaptation part of the movement. Defining angular locations relative to the via point, in each experiment there were two target locations, at 0° and at 270° respectively. The 0° target location was associated with lead-in start locations at +135° and 235°, and the 270° target location was associated with lead-in start locations at +45° and 135°. For each target location, the associated start locations were indicative of the direction of the curl force field on the adaptation movement (clockwise or counter-clockwise). All in all, this resulted in four possible distinct two-part movements (combinations of lead-in and adaptation movement).

Fig. 2A-C shows start and target locations only for the 0° target location case. When the lead-in starts from the 235° location, the adaptation movement is associated with a CW field (Fig. 2A). When the lead-in starts from the 135° location, the adaptation movement is associated with a CCW field (Fig. 2B). Fig. 2C shows one of the two possible channel trials, in this case with the lead-in starting from the 235° location (the other case is with a lead-in from the 135° start location but is not shown here). This relationship between starting location and the CW and CCW field directions shown here was switched for half of the participants, to counterbalance the relationship of lead-in direction and field direction across the experiments.

Session Design. As training an interference paradigm can take a significant number of trials and a large number of channel trials were required to collect the generalization data and enable low variance estimation of compensation to be made, each experiment was performed in two separate sessions on different days. This procedure was adopted to limit fatigue effects that might have been experienced from one single long experimental session. As some forgetting may result between sessions, the second session commenced with another training phase before proceeding with the generalization phase. In total, there were 1546 and 1580 trials on days 1 and 2 respectively, leading to a total of 3126 trials overall. In particular there were 2248 training field exposure trials and 544 channel trials use to probe generalization characteristics.

Probe Trials. In order to probe the generalization characteristics there were 17 different lead-in conditions for the channel trials (Table 1). Each lead-in was performed from the two possible starting locations to probe recall of both the CW and CCW field contexts. This gave rise to a total number of 34 distinct channel trial lead-ins (17 x 2 starting locations). Each of these 34 distinct channel trials was repeated 16 times during the generalization phases of the experiments. These generalization lead-in conditions were always followed by

a channel trial on the adaptation movement. Fifteen of these seventeen lead-in
movements were chosen to sample lead-in distances between 3 - 20 cm, peak
speeds between 8.04 - 53.57 cms^{-1} and durations between 210 – 1400 ms (Fig
1). One of these 15, had the exact motion as the training movements (Fig 1,
black line). The sixteenth lead-in movement was designed to test transfer
between passive and visual lead-in movements. That is, in the passive lead-in
experiment, this lead-in was a visual lead-in with the same kinematics as the
training movement. In the visual lead-in experiment this was a passive lead-in
movement. Finally, the seventeenth condition was a reversed visual cursor
condition. This "lead-away" condition was similar to a lead-in movement
between a start and via-position, except the cursor started at the central via-
point and moved to what was previously the start position. This condition
examined if the reversal of visual motion would still lead to a recall of dynamics
during the second probe phase, or whether the form of visual movement
needed to be consistent with the lead-in movement used during training to elicit
an appropriate contextual effect or transfer. Both the trained lead-in probe trial
and the opposite modality lead-in probe trials were also repeated throughout
the pre-exposure and training phase of experiment, such that each was
repeated a total of 38 times. The trained lead-in probe trials provided a means
to assess learned compensation as the experiment progressed.

Trial block organization. The experiment was organized in blocks. These blocks
contained different numbers and types of trials in the pre-exposure, training and
test phases of the experiment. Within a block, trials were sorted pseudo-
randomly, with the constraint that channel trials were not allowed to be adjacent
or the first trial in a block. Participants were required to take short rest breaks
approximately every 200 trials (195-205 trials) but could rest at any time
between trials. The blocks and trials were organized as follows:

Day 1

Pre-exposure: The pre-exposure phase started with 2 blocks of 40 trials. A
block consisted of 36 Null trials and 4 channel trials. The four channel trials
were 2 channel trials in the training condition (one for each lead-in direction)
and 2 channel trials in the transfer condition (that is, in the other modality, with
one trial for each lead-in direction). The 36 null trials were evenly split between
movements to the 0° direction target and the 270° direction target, with equal
numbers of each lead-in direction for each target (9 repetitions of each of the
four trial types). Next, participants were provided with three repetitions of each
of the 17 generalization channel trials for each of the two possible lead-in
directions (102 generalization condition channel trials) to ensure prior
experience of all the generalization trial conditions. Finally, 2 blocks of 40 trials
were again performed as described above (36 null and 4 channel).

Field-exposure Training: During the exposure phase, participants were
exposed to the curl force fields during the adaptation movement. This phase
consisted of 12 blocks of 40 trials (36 field trials and 4 channel trials in a block)
arranged as in the pre-exposure phase. This was a total of 480 trials (432 force
field trials, 24 training condition channel trials and 24 transfer condition channel
trials).
289
Generalization Testing: This phase examined generalization of the learned
predictive compensation by pseudo-randomly interspersing curl force field trials
with trials in which channel trials were preceded by the full range of contextual
34 movements (17 different generalization trial types x 2 lead-in directions).
The generalization phase consisted of 6 blocks of 134 trials. Each block
consisted of one of each of the 34 channel trials and 100 curl force field trials
(25 of each of the four types). This provided 6 repetitions of each of the 34
generalization channel trials. In this phase there was a total of 804 trials (600
curl field trials and 204 generalization channel trials).
299
300　Day 2
301
302　*Exposure Training:* At the beginning of the second session, training was briefly
303　resumed. The phase consisted of 6 blocks of 40 trial blocks (36 curl field trials
304　+ 4 channel trials) for a total of 240 trials.
305
306　Generalization Testing: Similar to the session of Day 1, participants performed
307　10 blocks of 134 trials (Total 1340 trials: 1000 curl field trials and 340
308　generalization condition channel trials), such that each of the 34 different probe
309　trials (17 x 2 lead-in directions) was repeated 10 times.
310
311
312　*Experiment 1. Passive lead-in movements.*
313
In experiment 1, the contextual lead-in movement was comprised of passive
movement of the participant's hand. This passive movement was produced by
the robotic manipulandum passively moving the participant's hand while no
cursor was presented. Each trial began by displaying the start location for the
lead-in movement, the central location and final target. The vBOT then moved
the participant's hand to the lead-in start location. Once the handle was
stationary within the start location for 300 ms, a beep was generated indicating
the start of the trial. At this time, the handle of the robotic system moved to the
central via-point following a minimum jerk trajectory. The training contextual
movement was a 10 cm movement of duration 700 ms. Once the hand reached
the central location, participants were required to produce an active adaptation
movement from the central location to the final target location. The dwell time
of the hand within the central via point was required to be between 0-250 ms,

otherwise a warning was provided. If dwell time exceeded 500 ms then the trial
was aborted and repeated. If the second movement (adaptation movement)
duration was between 450 ms and 600 ms a "Great" message was displayed;
otherwise an appropriate "Too Fast" or "Too Slow" warning was shown. Force
fields and channel trials were only ever presented during this second
movement.
333
334 *Experiment 2. Visual lead-in movements.*
335
Experiment 2 had a similar design to Experiment 1 and used the same block
structure. The only difference was that the contextual lead-in movements for
both training and generalization testing consisted of a visual movement of the
cursor. This is illustrated in Fig. 2D-F for the 0° target location condition. The
training contextual lead-in movement again followed a minimum jerk trajectory
of duration 700 ms from the start to the central location. During this time, the
participant's hand remained stationary at the central location. Immediately after
the cursor reached the central location, the participant made an active reaching
adaptation movement from the central location to the final target. The same
variations of generalization movement trials were performed (but with visual
instead of passive motion). In addition, a transfer condition was used in which
a passive movement lead-in was performed. Again, a reversed visual cursor
condition was also employed.
349
350 *Participant instructions*
351
352 Verbal instructions were used to instruct participants how to take part in the
353 experiments. Participants were first asked to read the ethics form and sign if
354 they wished to proceed with the experiment, which would last a few hours over
355 two days.
356
Participants in the passive condition were informed that at the start of a trial that
the robot would pull them to the via point and they were required to immediately
try to move to the target. They were told that this robot pulled movement could
come from different directions or with different lengths of movements, but that
their task was the same – to make a movement from this via-point to the target.
They were asked to move briskly and that feedback on movement speed and
any start delay would be reported after each trial. Specifically, they were
informed that the movement should be a point-to-point goal directed reaching
movement completed in a single movement without overshooting or
undershooting the target. They were informed that they should go roughly
straight to the target using a natural relaxed movement. They were also told
that at some point during the experiment some kind of disturbance would occur
in which the machine might push them to the left or the right of their movement
and disturb their trajectory. They were also informed that to complete the

movement they need to get the cursor to the target regardless of this
disturbance. Finally, they were also informed that approximately every 200
trials there would be a short break. They were also made aware that they could
take a break at any time if they released the handle switch on the robot.
Instructions for the visual condition were identical, except participants were
informed that on commencement of a trial the cursor would move to the center
position while their hand was stationary at a via point, and then they were
required to try to move quickly to the target, as in the passive experiment.
379
380 **Data Analysis**
381
The experimental data was analyzed offline using Matlab R14. Statistics to
examine differences between the generalization from visual lead-in and passive
lead-in movements were performed in JASP 0.11.1 (JASP Team, 2019) using
a repeated measure ANOVA. T-tests were performed within Matlab. To
examine learning, kinematic error on the adaptation movements and force
compensation on the channel trials were used.
388
Kinematic error. The kinematic error was calculated on the adaptation portion
of the movement, but only during the null and curl field trials. To be consistent
with our previous work, this was quantified as the maximum perpendicular error
(MPE) for each trial, which is the maximum deviation of the hand path to the
straight line joining the movement starting location to the target. It was
necessary to have different experimental block sizes in various phases of the
experiment to accommodate the training and generalization trials. Therefore, to
calculate a consistent averaged-across-trials of MPE, an analysis block size of
8 was chosen, because this number of trials could be used consistently
throughout the experiment. During calculation of the analysis block MPE
average, the sign of each trial MPE was flipped appropriately so that results
from CW and CCW field trials could be averaged together. The mean and
standard error (SE) of the averaged-across-trials MPE values was then
computed across all participants.
403
404
Force compensation. On each channel trial, during the adaptation movement
between the via point to the final target, velocity of the movement towards the
target and the force exerted by participants perpendicularly into the wall of the
simulated channel were simultaneously recorded at 1000Hz, to enable the
estimation of predictive feedforward adaptation [25], which is an established
technique in the analysis of dynamic learning in viscous curl fields [26]. Using
this paradigm, the channel clamps movement perpendicular to the direction of
movement to small values by resisting it using a high spring constant (6000
Nm^{-1}) as well as viscous damping (30 $Nm^{-1}s$), thereby minimizing lateral
movement error. As there is no lateral error on these trials, there is no error-

induced feedback component. Since channel trials occur sparsely within blocks
of curl fields trials, this means that any measured perpendicular force exerted
into the channel will be dominated by the predictive feedforward force that has
been learned to compensate the curl field.
419
To estimate the level of force field compensation of the participant, the
measured perpendicular channel force samples during movement along the
channel were regressed against their corresponding forward velocity samples
scaled by curl field strength. This calculation was performed for each movement
over the period from leaving the via-point until entering the target. This yielded
an estimate of the level of force compensation present in each channel trial
[26]. For each participant, the force compensation values were averaged
across 2 sign-corrected sequential channel trials (since they corresponded to
opposite curl field directions). The mean and standard error (SE) of
compensation was then computed across participants. This method to assess
adaptation to the novel dynamics is preferable to relying on a reduction in
kinematic error during force field learning, since the latter can also arise from
muscle co-contraction [27-29].
433
434
Results
436
In the passive lead-in experiment, participants performed active reaching
movements to a target after being passively moved from a start position to a
central target. After initial movements in a null field, participants were presented
with a curl force field during the active movement. The direction of the curl field
depended on the angle between the passive movement and active movement
(Fig. 2A,B). When presented with the curl force field, participants' adaptation
movements were disturbed, producing large errors that were gradually reduced
over the exposure phase (Fig. 3A). Throughout the experiment channel trials
were introduced on random trials in order to measure the predictive force
compensation throughout adaptation (Fig. 2C). Over a similar timescale as the
reduction in kinematic error, force compensation increased, reaching just over
62% compensation averaged over both force fields (Fig. 3B). A small but
significant increase in the kinematic error can be seen between day 1 and the
start of day 2 (paired t-test between final block on day 1 and first block on day
2: $t_7=3.49$; p=0.01), but the associated decrease in force compensation was not
significant ($t_7=2.09$; p=0.075). Compensation to each field direction separately
is shown in Fig. 3F. The final levels of force compensation (mean of final 4
blocks) were not significantly different between the two lead-in directions
(paired t-test: $t_7=0.743$; p=0.48).
456
457 Participants in the visual lead-in experiment performed a similar protocol but

where the lead-in movements were purely visual in nature (Fig. 2 D-F). Again, when presented with the curl force field, participants' adaptation movements were disturbed, producing large errors that were gradually reduced over the exposure phase (Fig. 3D). Again, a small but not significant reduction in the force compensation (t_7=1.67; p=0.14) and a small but significant increase in the kinematic error (t_7=4.19; p=0.004) can be seen between day 1 and the start of day 2 (final block day one compared to first block day 2 with paired t-test). Over a similar timescale, force compensation increased, reaching approximately 70% compensation averaged over both force fields. Mean and SE of percentage force compensation showing compensation for each field direction separately is shown in Fig. 3F. The final levels of force compensation (mean of final 4 blocks) were again not significantly different between the two lead-in directions (paired t-test: t_7=0.402; p=0.70).

On random trials late in the adaptation phase, channel trials were applied with a range of different lead-in movement kinematics (Fig. 1) in order to examine generalization. After learning the force fields with the passive lead-in movement, variations in the kinematics of this lead-in movement produced a range of generalization levels (Fig. 4A). As the testing lead-in movements varied further away from the training kinematics the predictive force level decreased. A repeated measures ANOVA with a within subject effect of lead-in kinematics (15 levels) demonstrated a significant main effect ($F_{14,98}$=33.057; p<0.001). A similar finding is shown for the generalization after learning a visual lead-in movement (Fig. 4B). Although in this condition only small decreases in the predictive force are seen over a wide range of changes in the lead-in kinematics, the repeated measures ANOVA again showed a significant main effect of lead-in kinematics ($F_{14,98}$=23.397; p<0.001) demonstrating that the lead-in kinematics affected the predictive force on these channel trials.

Across the different lead-in movement kinematics, several conditions had the same duration, peak velocity or distance as the learned training condition (dotted lines in Fig. 4). We examined the predictive force compensation values over these conditions in more detail along iso-contours for lead-in distance, duration and speed (Fig. 5). Significant differences for select comparisons are reported from post-hoc tests following the significant main within subject effect of the repeated measures ANOVA reported above (Holm corrected for multiple comparisons across all 15 levels). The results for the passive lead-in condition show strong variations over changes in lead-in duration, distance and speed (Fig. 5A-C).

In panel A, lead-in distance is constant and the movement duration (shown on the x-axis) and speed vary. In panel B, lead-in duration is constant and the movement distance (shown on the x-axis) and speed vary. In panel C, lead-in

speed is constant and the movement duration (shown on the x-axis) and the
distance vary. The dotted lines indicate the training values of lead-in distance
and lead-in duration. It can be seen that the recall of predictive compensation
was strongly affected by changes in the duration of the movement (shown for
constant distance conditions in Fig. 5A and constant speed conditions on Fig.
5C). There was a strong tuning effect centered around the movement duration
used for training (Fig 5A). Specifically, compared to the training value, the force
compensation significantly decreased as movement duration either increased
to 1.4s (p=0.016) or decreased to 0.35s (p=0.033). Similar results were seen
for a constant peak speed (Fig 5C), but while compensation significantly
decreased as movement duration decreased to 0.21s (p=0.01) this did not
reach significance for the longer duration (p=0.062). However, changing lead-
in distance produced different effects. While reducing lead-in distance to 3cm
reduced compensation (p=0.004), increasing movement distance from the
training value to 20cm had essentially no effect (p=1.0) as shown in Fig. 5B.

In the visual lead-in condition, we found a slightly less pronounced reduction in
recalled compensation as the some of the kinematics were varied while others
remained constant (Fig. 5 D-F). Again, it can be seen that compensation was
affected by changes in the duration of the movement, with compensation
significantly falling off as movement duration decreased to 0.35s (p<0.001).
Although there was a tuning effect centered around the movement duration
used for training, as duration deviated from the training value the fall off was
appeared less pronounced than in the passive lead-in condition (Fig. 5D).
Similar results were seen for a constant peak speed (Fig 5F), with
compensation significantly falling off as movement duration decreased to 0.21s
(p<0.001). However, as the training distance varied but duration was fixed (Fig
5E, we found no significant differences (all p>0.1).

If we compare the generalization of the predictive compensation across the two
experiments, we can see one major finding; namely that the overall tuning
effects were more pronounced for the passive lead-in condition compared to
the visual lead-in condition. Visual lead-in generalization showed less
sensitivity to variations in lead-in kinematics. Indeed, whereas passive lead-ins
resulted in a 2D monotonic curved surface in the dimensions of duration and
distance (Fig. 4A), the corresponding surface for visual lead-ins (Fig. 4B)
exhibits a large region consisting of a flat planar surface. To test if this
difference in the generalization across kinematics between passive and visual
lead-in movements is statistically significant, we performed ANOVAs on the
force compensation results (with main effects of kinematic condition (14 levels)
and lead-in modality (2 levels: visual or passive). To do so, the force
compensation for each testing condition was normalized with respect to the
value at the trained condition for each participant. We then compared the
generalization across kinematic conditions using a repeated measures ANOVA

with a repeated measure factor of lead-in condition (15 levels) and between
subject factors of experimental condition (2 levels: passive and visual lead-ins).
We report the Greenhouse-Geisser sphericity corrected values. There was both
a significant main-effect of lead-in condition ($F_{5.127,71.782}$=56.627; p<0.001),
experimental condition ($F_{1,14}$=8.073; p=0.013), and interaction between these
two ($F_{5.127,71.782}$=3.061; p=0.014). As the predicted force compensation level
was normalized to 100% for the training condition, the presence of a significant
main effect of experimental condition demonstrates that there were differences
between the shape of this generalization between the passive and visual lead-
in movements. This highlights a clear difference between visual inputs and
passive inputs as a contextual signal for motor adaptation, extending our
previous findings [10,11].
558
We also investigated how learning opposing force fields with contextual cues
in one sensory modality would transfer to the other sensory modality. To
investigate this, occasional channel trials were used with a lead-in in the other
modality. It can be seen that there is asymmetric transfer between passive lead-
in and visual lead-in movements (Fig. 6). Although there was partial transfer
from passive to visual lead-in movements (Fig. 6A) with values reaching just
above 20%, there was much less transfer from visual to passive lead-in
movements (Fig. 6B) with values just under 10%. To compare the level of
transfer between the two modalities, the transferred adaptation was scaled
according to the final level of adaptation in each experiment and compared
using a t-test. The transfer from passive to visual (35.2% ± 12.1 %; mean ± std)
was significantly larger (t_{14}=3.966; p=0.0014) than the transfer from visual to
passive (13.0% ± 10.2 %). Thus, there is a clear asymmetry between the
transfer of adaptation between these two sensory modalities.
573
Finally, we examined how learning the visual lead-in movement would transfer
to a completely reversed visual cursor (with the same duration and distance).
To balance conditions and the number of trials across experiments, this was
also tested for the passive lead-ins. To compare between the two experiments,
the predictive compensation of the reversed visual cursor was scaled by the
predictive compensation on the training condition. There was no significant
difference in transfer to reverse visual movement between participants trained
with passive lead-in movements (22.7% ± 17.3; mean ± std) versus those
trained with visual lead-in movements (29.3% ± 23.0) using a t-test (t_{14}=0.641;
p=0.53). This shows that the predictive compensation is sensitive to the
direction of the visual motion.
585

586   **Discussion**
587
We investigated the kinematic generalization characteristics of passive and

visual lead-in movements using a force field interference paradigm. Participants
first experienced a lead-in movement and then immediately made an active
movement in a curl force field where the field direction was associated with the
lead-in movement. Channel trials within the active movement examined how
predictive compensation varied as lead-in kinematics were varied. In the first
experiment lead-in movements were passive, whereas in the second
experiment they were visual. For both modalities, recall of predictive
compensation decreased as the duration of the lead-in movements deviated
from the training condition. Reducing lead-in distance also reduced
compensation but increasing lead-in distance had little effect on the force
generalization. Our results show that although passive and visual lead-in
movements influence memory formation and recall in subsequent movement,
passive motion exhibits narrower generalization characteristics, whereas visual
motion is less sensitive to kinematic change.

These generalization results further characterize the neural tuning exhibited by
lead-in movements, extending beyond the directional tuning seen previously.
The observation that passive lead-ins were more sensitive to changes in
kinematics than visual lead-ins is consistent with the prior observations
examining angular generalization [10,11,15]. Namely that active and passive
tuning was narrower than the wider tuning seen in the visual condition. Here
we describe this gradual change in the predictive force output on the adaptation
movement as the lead-in motion is changed as generalization. Nevertheless, it
is important to note that while the direction of the force field depended on the
lead-in direction, variations in the lead-in motion were never used to produce
changes in the forces arising from the force field. This means that participants
could in principle have simply learned a binary left-right mapping to the force
fields, in which case we would be a constant predictive compensation for all
variations of the lead-in kinematics. However, it can be seen that variations in
the kinematics of the passive (and to a lesser extent visual) lead-in movements
produced clear variations in the predictive forces. This shows that the learning
of this mapping was at least partially local and associated with the specific
kinematics of the lead-in movement. The continuous nature of this effect, which
exhibits a smooth transition in compensation between training conditions,
agrees with what was seen previously for angular generalization for both
passive [10] and visual lead-in movements [11].

More recently it was shown that the tuning characteristics of different lead-in
modalities could explain why angular variability of active lead-in movements
affects the learning rate in two-part movement tasks, whereas no such effect
exists for visual lead-in movements [4]. Our current results suggest that
variations in the speed or duration of lead-in movements could provide similar
decrements in learning rate, whereas an increase in movement distance would
not have much effect. This might have important implications for rehabilitation,

suggesting learning and recovery would be faster for training routines with
consistent lead-in kinematics. One caveat is that such routines might also
produce less generalization across tasks, as the adaptation is more likely to be
learned specifically for the trained lead-in movement.

In both experiments, to examine transfer of adaptation across modality, a visual
lead-in cursor motion occasionally replaced the passive lead-in, and vice versa.
Interestingly, there was an asymmetric transfer between passive and visual
lead-in movements, with partial transfer from passive lead-in movements to
visual lead-in movements, but almost no transfer from visual to passive lead-in
movements. Transfer could arise because passive lead-in movement partially
engages neural mechanisms shared by the visual observation of movement,
but not the converse. This result may be due to asymmetry in the connections
between the neural substrates. Alternatively, it could arise because the visual
feedback pathway has a lower gain due to the uncertainly introduced by the
longer time delay associated with visual information [30]. The current
observation that passive lead-in are more strongly tuned in duration than visual
lead-ins, as well as the former results that the absolute level of influence of
passive lead-ins [10] is higher than for visual lead-in [11] supports the latter
hypothesis.

The wide ranging results from studies examining contextual cues for learning
opposing dynamics have demonstrated that not all sensory signals are able to
influence motor learning [15,20,31-37]. For example, color has essentially no
effect [20]. In addition to the strong effects of prior movements, it has been
shown that particularly effective contextual cues relate to state; for example
limb state and physical locations [38,39], or different visual locations of the
cursor and targets [33]. Indeed, a location cue could constitute a complete
physical shift of the movement task, or just a shift of one of its two essential
components; namely a change in the location of the visual feedback, or a
change in the physical location of the task with identical visual feedback. Other
experiments have shown that future state also effects motor learning in an
analogous way [3], with this effect depending on movement planning rather
than execution [5].

On the face of it, it appears that there are multiple types of contextual cues that
strongly influence motor memory formation. Here we propose that a factor they
all have in common is that they are related to either past, current, or future state
of the limb; or are signals used in the estimation of such limb states. That is,
setting up the sensorimotor system in a different state before (or at the end of)
a movement allows the formation and recall of different motor memories. This
suggests that some contextual cues (such as visual lead-in movement or
location in the visual workspace) are simply effective because the motor system
makes use of these signals within a state estimation framework to determine

the state of the arm. Such state estimation can only be made on the basis of
sensory feedback and efference copy. This hypothesis would be consistent with
the observation that visual or proprioceptive movements are as effective an
active movement. It would also explain why a visual change of state can be as
effective as a complete change in the physical state of the limb. Moreover, it
can explain why vestibular inputs could also be used to learn opposing
dynamics [15] but why color cues have much less effect on the adaptation
system [20].
685
In order to reach with our arm to a specific location, our sensorimotor control
system needs to know the initial limb state, and then activate the appropriate
muscles in a specific pattern to generate forces that bring the arm into the final
state to meet the task requirements. To make this movement robust in the face
of noise and disturbances, this process does not simply rely on feedforward
control, but makes use of sensory feedback of our arm's state, enabling online
correction in any task-relevant deviation from the goal of our movement. Arm
state can be estimated through the combination of appropriate sensory
feedback signals such as proprioception from the skin muscles and joints,
visual information, and vestibular inputs. However due to neural signal
transmission and processing delays, motor responses to proprioceptive and
visual feedback only start producing force after delays of 50 ms and 140 ms
respectively. Such delays represent a challenge in the design of a feedback
control systems, since using direct feedback from delayed signals can lead to
instability.
701
To deal with delays in sensory feedback, Smith proposed an architecture which
involves using immediate feedback from the output of a forward model of the
plant, rather than the actual output of the plant itself [40]. In this architecture,
the forward model estimates plant output without delay, thereby avoiding the
instabilities that delay can introduce, Miall and Wolpert suggested that the
Smith predictor architecture could account for delays in the human motor
system [41,42]. In control engineering, forward models form part of observers,
which are systems used to estimate the full state of a plant, which often cannot
be observed directly. Such full state estimation can be used as the basis of full
state feedback control of a plant, which is more effective than only basing
control on the observable plant output. As with the Smith's predictor, such
observers can also be constructed to estimate plant state without the delay,
providing an elegant way to control plants that have inherent delay in their
sensory feedback paths.
716
To make an observer robust to inaccuracies in its forward model and to deal
with unpredictable disturbances, there is normally a state correction pathway
term based on actual output error calculated as the difference between the
actual output of the plant, and an appropriately delayed prediction of plant

output. This results in the state prediction based on the efference copy of the
motor command being combined with a state prediction error correction term
based on the delayed sensory feedback, which is something that has been
shown to occur during the control of human movement [43].
725
Within such an observer-based controller framework, the observer performs
state estimation for an active movement using efference copy, while improving
the estimate using the delayed feedback signals. In the case of a purely visual
observation or passive movement of the arm, the observer can still make a
state estimate, but only based on the state corrections from feedback. From the
premise that state is the key issue in formation of separate motor memory, such
a framework would account for the observation that either active, passive or
visual lead-in movements would influence state estimation.
734
To conclude, we have shown the current and previous observations of lead-in
phenomena are consistent with the hypothesis that the human motor system
operates as an observer-based controller mechanism, that makes use of a
forward model to estimate state. In particular, our results support the proposal
[30] that even though the variances of visual positional information is known to
be lower than that obtained from proprioception, its longer temporal delay
reduces its weighting in state estimation. As a consequence of this, visual
information has less effect on the motor system than proprioceptive information,
an effect that we have extended to the learning and generalization of opposing
dynamics.
745
746 **Author contributions**
747
748 ISH and DWF designed and implemented the study. SF performed data
749 collection. ISH performed data analysis. ISH, SF and DWF wrote the
750 manuscript.
751
752 **Competing interest statement**
753
754 The authors declare that they have no financial, personal, or professional
755 interests that could be construed to have influenced the paper.

Condition	Duration [ms]	Distance [cm]	Peak velocity
P/V Null/Training	700 ms	10 cm	26.8 cms-1
P Channel	700 ms	10 cm	26.8 cms-1
V Channel	700 ms	10 cm	26.8 cms-1
P/V Channel	1400 ms	20 cm	26.8 cms-1
P/V Channel	1050 ms	15 cm	26.8 cms-1
P/V Channel	420 ms	6 cm	26.8 cms-1
P/V Channel	210 ms	3 cm	26.8 cms-1
P/V Channel	1400 ms	10 cm	13.4 cms-1
P/V Channel	1050 ms	10 cm	17.9 cms-1
P/V Channel	420 ms	10 cm	44.6 cms-1
P/V Channel	350 ms	10 cm	53.6 cms-1
P/V Channel	700 ms	20 cm	53.6 cms-1
P/V Channel	700 ms	15 cm	40.2 cms-1
P/V Channel	700 ms	6 cm	16.1 cms-1
P/V Channel	700 ms	3 cm	8.0 cms-1
P/V Channel	1050 ms	20 cm	35.7 cms-1
P/V Channel	350 ms	3 cm	16.1 cms-1
Reverse V Channel	700 ms	10 cm	-26.8 cms-1

Table 1. Durations, distances and peak velocities of the lead-in movements for
the training and generalization conditions. Passive and visual lead-in
movements are represented by P and V respectively. The P/V Training
condition used a passive lead-in for experiment 1 and with a visual lead-in for
experiment 2. In both experiments, test trials with a passive lead-in and a visual
lead-in were performed with the same kinematics as the training conditions.
Similarly, a P/V channels represents channel conditions used with a passive
lead-in for experiment 1 and with a visual lead-in for experiment 2. The Reverse
V Channel was used with a visual lead-in in both experiments.
774

Figure 1. Kinematics of lead-in movements used for testing generalization. **A** Profiles of movement distance versus duration of lead-in probe conditions across all conditions. Thick black line indicates the training lead-in motion. Colors indicate specific conditions that are matched across duration (red), peak speed (green), or duration (blue). **B** Lead-in movement kinematics of peak speed as a function of duration.

784
785
Figure 2. Experimental Design. **A-C**: Passive lead-in generalization experiment.
A Participants first experienced an initial passive lead-in motion from a starting
position (grey circle) to the central target (green circle) and then immediately
made a second active movement to the target (yellow circle) on which a curl
force field (blue arrows) could be applied. **B** An initial movement from a different
starting target was associated with the opposite force field on the second
movement. The direction of curl force field and lead-in movements were
counterbalanced across participants. **C** In order to examine learning rate and
generalization, random trials in which the contextual movement was followed
by a mechanical channel on the second movement to the target were applied.
D-F: Visual lead-in generalization experiment. **D** Participants initially observed
an initial visual cursor movement (red circle) from the grey starting circle to the
central target (green circle). Once the cursor entered the central target,
participants immediately performed a second active movement to the target
(yellow circle) on which a curl force field (blue arrows) could be applied. **E** An
initial cursor movement from a different starting target was associated with the
opposite force field on the second movement. **F** On random trials, after the
visual lead-in motion, a mechanical channel was applied on the active
movement to the target to measure predictive compensation.
805

806
Figure 3. Adaptation to two opposing force fields. **A** Mean and SE of MPE
across over 8 participants for the passive lead-in experiment as a function of
blocks of 8 trials. **B** Mean and SE of percentage force compensation for pairs
of channel trials (one for each force field direction) throughout the passive lead-
in experiment where the lead-in movement was the same as the training trials.
C Mean and SE of percentage force compensation as B but showing
compensation for each field direction separately. **D** Mean and SE of MPE for
the visual lead-in experiment. **E** Mean and SE of percentage force
compensation for the visual lead-in experiment. **F** Mean and SE of percentage
force compensation as E but showing compensation for each field direction
separately.

Figure 4. Generalization surface plots for passive and visual lead-in movements. **A** Surface plot of generalization for passive lead-in movements. The percentage force compensation is represented by color and plotted against lead-in duration and lead-in distance. The black circle with a white center indicates the result at the training condition. The solid black dots indicate points for which measurements were made on probe trials. The black dotted lines correspond to conditions with the same training lead-in distance of 10 cm, same training lead-in duration of 0.7s or same training lead-in speed of 26.8 cm/s. The legend shows the correspondence between color and percentage perfect force compensation. **B** Surface plot of generalization for the visual lead-in condition.

Figure 5. Generalizations results for both passive and visual lead-in conditions plotted for fixed values of lead-in distance, lead-in duration and lead-in speed. **A-C** Results of passive lead-in experiments. The dotted lines indicate the training values of lead-in distance and lead-in duration. Error bars indicate standard error of the mean. Significant differences are indicated for post-hoc comparisons (Holm corrected for multiple comparisons). **A** Effect of changing lead-in duration for fixed 10 cm lead-in distance. In this panel, lead-in distance is held constant and the movement duration (shown on the x-axis) and speed vary. **B** Effect of changing lead-in distance (and peak speed) across conditions with a fixed 700 ms lead-in duration. **C** Effect of changing lead-in duration (and distance) across conditions with a fixed 26.7 cm/s lead-in speed. **D-F** Corresponding results for visual lead-in condition.

Figure 6. Comparison of transfer across sensory modality or to reversed visual cursor. **A** Transfer from passive lead-in to visual lead-in or reversed visual lead-in. For comparison the learned force compensation on passive movements is shown. Black circles indicate the results of individual participants. Error bars indicate standard error of the mean. **B** Transfer from visual lead-in to passive lead-in or reversed visual lead-in.

References

1. Reinkensmeyer DJ, Burdet E, Casadio M, Krakauer JW, Kwakkel G, Lang CE, et al. Computational neurorehabilitation: modeling plasticity and learning to predict recovery. J NeuroEngineering Rehabil. BioMed Central; 2016;13: 42. doi:10.1186/s12984-016-0148-3

2. Howard IS, Ingram JN, Franklin DW, Wolpert DM. Gone in 0.6 seconds: the encoding of motor memories depends on recent sensorimotor States. Journal of Neuroscience. 2012;32: 12756–12768. doi:10.1523/JNEUROSCI.5909-11.2012

3.   Howard IS, Wolpert DM, Franklin DW. The value of the follow-through
derives from motor learning depending on future actions. Curr Biol.
Elsevier; 2015;25: 397–401. doi:10.1016/j.cub.2014.12.037

4.   Howard IS, Ford C, Cangelosi A, Franklin DW. Active lead-in variability
affects motor memory formation and slows motor learning. Scientific
Reports. Nature Publishing Group; 2017;7: 7806. doi:10.1038/s41598-
017-05697-z

5.   Sheahan HR, Franklin DW, Wolpert DM. Motor Planning, Not
Execution, Separates Motor Memories. Neuron. 2016;92: 773–779.
doi:10.1016/j.neuron.2016.10.017

6.   Churchland MM, Cunningham JP, Kaufman MT, Foster JD, Nuyujukian
P, Ryu SI, et al. Neural population dynamics during reaching. Nature.
Nature Publishing Group; 2012;: 1–8. doi:10.1038/nature11129

7.   Pandarinath C, Gilja V, Blabe CH, Nuyujukian P, Sarma AA, Sorice BL,
et al. Neural population dynamics in human motor cortex during
movements in people with ALS. Elife. 2015;4: e07436.
doi:10.7554/eLife.07436

8.   Donchin O, Francis JT, Shadmehr R. Quantifying generalization from
trial-by-trial behavior of adaptive systems that learn with basis functions:
theory and experiments in human motor control. Journal of
Neuroscience. 2003;23: 9032–9045.

9.   Thoroughman KA, Shadmehr R. Learning of action through adaptive
combination of motor primitives. Nature. 2000;407: 742–747.
doi:10.1038/35037588

10.   Howard IS, Franklin DW. Neural Tuning Functions Underlie Both
Generalization and Interference. Gribble PL, editor. PLoS ONE. Public
Library of Science; 2015;10: e0131268.
doi:10.1371/journal.pone.0131268

11.   Howard IS, Franklin DW. Adaptive tuning functions arise from visual
observation of past movement. Scientific Reports. Nature Publishing
Group; 2016;6: 28416. doi:10.1038/srep28416

12.   Mattar AAG, Ostry DJ. Modifiability of Generalization in Dynamics
Learning. Journal of Neurophysiology. 2007;98: 3321–3329.
doi:10.1152/jn.00576.2007

13. Goodbody SJ, Wolpert DM. Temporal and amplitude generalization in
motor learning. Journal of Neurophysiology. 1998;79: 1825–1838.
doi:10.1152/jn.1998.79.4.1825

14. Gandolfo F, Mussa-Ivaldi FA, Bizzi E. Motor learning by field
approximation. Proc Natl Acad Sci USA. 1996;93: 3843–3846.

15. Sarwary AME, Stegeman DF, Selen LPJ, Medendorp WP.
Generalization and transfer of contextual cues in motor learning.
Journal of Neurophysiology. 2015;: jn.00217.2015.
doi:10.1152/jn.00217.2015

16. Caithness G, Osu R, Bays P, Chase H, Klassen J, Kawato M, et al.
Failure to consolidate the consolidation theory of learning for
sensorimotor adaptation tasks. Journal of Neuroscience. 2004;24:
8662–8671. doi:10.1523/JNEUROSCI.2214-04.2004

17. Brashers-Krug T, Shadmehr R, Bizzi E. Consolidation in human motor
memory. Nature. 1996;382: 252–255. doi:10.1038/382252a0

18. Krakauer JW, Ghilardi MF, Ghez C. Independent learning of internal
models for kinematic and dynamic control of reaching. Nat Neurosci.
1999;2: 1026–1031. doi:10.1038/14826

19. Karniel A, Mussa-Ivaldi FA. Does the motor control system use multiple
models and context switching to cope with a variable environment? Exp
Brain Res. 2002;143: 520–524. doi:10.1007/s00221-002-1054-4

20. Howard IS, Wolpert DM, Franklin DW. The effect of contextual cues on
the encoding of motor memories. Journal of Neurophysiology.
2013;109: 2632–2644. doi:10.1152/jn.00773.2012

21. Nozaki D, Kurtzer I, Scott SH. Limited transfer of learning between
unimanual and bimanual skills within the same limb. Nat Neurosci.
2006;9: 1364–1366. doi:10.1038/nn1785

22. Oldfield RC. The assessment and analysis of handedness: the
Edinburgh inventory. Neuropsychologia. 1971;9: 97–113.

23. Howard IS, Franklin S, Franklin DW. Characterization of Neural Tuning:
Visual Lead-in Movements Generalize in Speed and Distance.
Converging Clinical and Engineering Research on Neurorehabilitation
III. Cham: Springer, Cham; 2018. pp. 1030–1033. doi:10.1007/978-3-
030-01845-0_206

24.   Howard IS, Ingram JN, Wolpert DM. A modular planar robotic
manipulandum with end-point torque control. Journal of Neuroscience
Methods. 2009;181: 199–211. doi:10.1016/j.jneumeth.2009.05.005

25.   Scheidt RA, Reinkensmeyer DJ, Conditt MA, Rymer WZ, Mussa-Ivaldi
FA. Persistence of motor adaptation during constrained, multi-joint, arm
movements. Journal of Neurophysiology. 2000;84: 853–862.

26.   Smith MA, Ghazizadeh A, Shadmehr R. Interacting adaptive processes
with different timescales underlie short-term motor learning. Plos Biol.
2006;4: e179. doi:10.1371/journal.pbio.0040179

27.   Milner TE, Franklin DW. Impedance control and internal model use
during the initial stage of adaptation to novel dynamics in humans. The
Journal of Physiology. 2005;567: 651–664.
doi:10.1113/jphysiol.2005.090449

28.   Franklin DW, Osu R, Burdet E, Kawato M, Milner TE. Adaptation to
stable and unstable dynamics achieved by combined impedance control
and inverse dynamics model. Journal of Neurophysiology. 2003;90:
3270–3282. doi:10.1152/jn.01112.2002

29.   Burdet E, Osu R, Franklin DW, Milner TE, Kawato M. The central
nervous system stabilizes unstable dynamics by learning optimal
impedance. Nature. Nature Publishing Group; 2001;414: 446–449.
doi:10.1038/35106566

30.   Crevecoeur F, Munoz DP, Scott SH. Dynamic Multisensory Integration:
Somatosensory Speed Trumps Visual Accuracy during Feedback
Control. J Neurosci. Society for Neuroscience; 2016;36: 8598–8611.
doi:10.1523/JNEUROSCI.0184-16.2016

31.   Addou T, Krouchev N, Kalaska JF. Colored context cues can facilitate
the ability to learn and to switch between multiple dynamical force fields.
Journal of Neurophysiology. 2011;106: 163–183.
doi:10.1152/jn.00869.2010

32.   Cothros N, Wong J, Gribble PL. Visual cues signaling object grasp
reduce interference in motor learning. Journal of Neurophysiology.
2009;102: 2112–2120. doi:10.1152/jn.00493.2009

33.   Hirashima M, Nozaki D. Distinct motor plans form and retrieve distinct
motor memories for physically identical movements. Curr Biol. 2012;22:
432–436. doi:10.1016/j.cub.2012.01.042

34.   Hwang EJ, Donchin O, Smith MA, Shadmehr R. A Gain-Field Encoding
of Limb Position and Velocity in the Internal Model of Arm Dynamics.
Plos Biol. Public Library of Science; 2003;1: e25.
doi:10.1371/journal.pbio.0000025

35.   Yokoi A, Hirashima M, Nozaki D. Gain field encoding of the kinematics
of both arms in the internal model enables flexible bimanual action.
Journal of Neuroscience. 2011;31: 17058–17068.
doi:10.1523/JNEUROSCI.2982-11.2011

36.   Wainscott SK. Internal Models and Contextual Cues: Encoding Serial
Order and Direction of Movement. Journal of Neurophysiology. 2004;93:
786–800. doi:10.1152/jn.00240.2004

37.   Nozaki D, Yokoi A, Kimura T, Hirashima M, Orban de Xivry J-J. Tagging
motor memories with transcranial direct current stimulation allows later
artificially-controlled retrieval. Elife. 2016;5. doi:10.7554/eLife.15378

38.   Hwang EJ, Shadmehr R. Internal models of limb dynamics and the
encoding of limb state. J Neural Eng. 2005;2: S266–S278.
doi:10.1088/1741-2560/2/3/S09

39.   Yeo S-H, Wolpert DM, Franklin DW. Coordinate Representations for
Interference Reduction in Motor Learning. Balasubramaniam R, editor.
PLoS ONE. Public Library of Science; 2015;10: e0129388.
doi:10.1371/journal.pone.0129388

40.   Smith OJM. A controller to overcome dead time. ISA J. 1959;6: 28–33.

41.   Miall RC, Weir DJ, Wolpert DM, Stein JF. Is the cerebellum a smith
predictor? J Mot Behav. 1993;25: 203–216.
doi:10.1080/00222895.1993.9942050

42.   Miall RC, Wolpert DM. Forward Models for Physiological Motor Control.
Neural Netw. Pergamon; 1996;9: 1265–1279.

43.   Wolpert DM, Ghahramani Z, Jordan MI. An internal model for
sensorimotor integration. Science. 1995;269: 1880–1882.