Social foraging extends associative odor-food memory expression in fruit flies

(*Drosophila melanogaster*)

Aarti Sehdev[1], Yunusa Mohammed[1], Cansu Tafrali[1], Paul Szyszka[1,2,*]

[1]University of Konstanz, Department of Biology, Neurobiology, Konstanz 78457, Germany

[2]Present address: University of Otago, Department of Zoology, Dunedin 9054, New Zealand

* Correspondence: paul@szyszkalab.com

Keywords: social interactions, social information transfer, odor-food learning, associative

memory, collective foraging, *Drosophila melanogaster*, automated classical conditioning

ABSTRACT

Animals socially interact during foraging to share information about the quality and location

of food sources. The mechanisms of social information transfer during foraging have been

mostly studied at the behavioral level, and its underlying neural mechanisms are largely

unknown. The fruit fly *Drosophila melanogaster* has become a model for studying the neural

bases of social information transfer, as fruit flies show a rich repertoire of social behaviors

and provide a well-developed genetic toolbox to monitor and manipulate neuronal activity.

Social information transfer has already been characterized for fruit flies' egg laying, mate

choice, foraging and aversive associative learning, however the role of social information

transfer on associative odor-food learning during foraging are unknown. Here we present an

automated foraging assay for groups of walking flies that allows studying the effect of group

size on the formation and expression of associative odor-food memories. We found that

inter-fly attraction increases with group size and that groups of flies exhibit extended odor-

food memory expression, as compared to single or pairs of flies. We discuss possible

behavioral and neural mechanisms of this social effect on odor-food memory expression.

This study opens up opportunities to investigate how social interactions are relayed in the

neural circuitry of learning and memory expression.

INTRODUCTION

Vertebrates often forage in groups to get a more accurate estimate of the location and quality of resources (Giraldeau and Caraco, 2000; Templeton and Giraldeau, 1996; Valone, 1989; Ward and Zahavi, 1973). Insects also convey information about the location and quality about a food source through social interactions. For example, honey bees signal the direction and distance of food locations to other bees (Frisch, 1965), ants complement their individual memory of a route to food using trail pheromones left by scouts (Czaczkes et al., 2011), and stimulus enhancement and local enhancement at the food source improves foraging efficiency in bumble bees (Alem et al., 2016; Avarguès-Weber and Chittka, 2014; Leadbeater and Dawson, 2017; Worden and Papaj, 2005). These social effects on foraging have been mostly studied at the level of behavioral outcome, and the neural mechanisms of how social information transfer improves foraging are still unknown.

The fruit fly, *Drosophila melanogaster*, is a suitable model organism for studying the effects of social interactions on foraging at both the behavioral and the neuronal level. Fruit flies are gregarious (Lefranc et al., 2001; Navarro and del Solar, 1975) and demonstrate a rich repertoire of social behaviors that encompass communication about internal states (Suh et al., 2004), social information spread during odor avoidance (Ramdya et al., 2014), foraging (Abu et al., 2018; Lihoreau et al., 2016; Tinette et al., 2004) and predator-induced egg-retention (Kacsoh et al., 2015). Moreover, fruit flies socially learn (Battesti et al., 2012a; Danchin et al., 2018; Durisko and Dukas, 2013; Golden and Dukas, 2014; Lin et al., 2015; Mery et al., 2009; Sarin and Dukas, 2009) and they show improved aversive odor memory retrieval when in groups (Chabaud et al., 2009). However, it is still unknown whether social information transfer affects flies' associative odor-food learning during foraging.

The mechanistic understanding of foraging in fruit flies is unparalleled, both in regard to the neural mechanisms of odor-guided search (Galizia, 2014; Haverkamp et al., 2018; Wilson, 2013) and feeding (Itskov and Ribeiro, 2013), and of associative odor-food learning (Burke et al., 2012; Huetteroth et al., 2015; Liu et al., 2012; Owald and Waddell, 2015; Schwaerzel et al., 2003; Tempel et al., 1983; Thum et al., 2007), making the fruit fly a good model for studying the neural mechanisms of social interactions during foraging.

Here we investigated whether fruit flies socially interact during foraging and whether group size affects associative odor-food memory expression. We developed an automated foraging assay to study classical odor-reward conditioning in single flies and in groups of flies. We

conditioned flies to associate an odorant with a sucrose patch, and then tested their associative odor-food memory expression. We found that small groups (3 - 4 flies) and large groups (7 - 8 flies) showed extended odor-food memory expression compared to pairs of flies or single flies, and pairs of flies showed shorter memory expression than single flies. Moreover, flies in small or large groups, but not in pairs, were attracted to each other. These data suggest that flies socially interact during foraging and that these social interactions increase the efficiency of odor-guided food search.

MATERIALS AND METHODS

Animals

Drosophila melanogaster wild type Canton S were raised on a standard food medium (100 mL contain 6.7 g fructose, 2.4 g dry yeast, 0.7 g agar, 2.1 g sugar beet syrup, 0.282 g ethyl paraben and 0.61 ml propionic acid). Flies were raised in a room with normal day light cycle, with an average temperature of 23.5 °C and 32% relative humidity. One to four days old flies were anesthetized with CO_2 and female flies were collected. Flies were starved for 2-3 days to motivate them to search for food. Flies were starved in a fly vial with filter paper soaked in water.

Conditioning apparatus

To condition groups of flies, we used an automated rotating platform with four concentric arenas (Figure 1A, B). The arenas were covered with a watch glass (7 cm diameter, 8 mm height in the center) which was coated on the inner side with Sigmacote (Sigma-Aldrich) to prevent flies from walking on the inside of the glass. The floor was made of a Teflon-coated fiberglass fabric (441.33 P, FIBERFLON, Konstanz). Pure odorants (ethyl acetate and 2,3-butanedione, Sigma-Aldrich) were stored in 20 ml vials (Schmidlin Labor and Service). The vials were mounted under the platform and the lid was pierced with a needle (Hypodermic-needle; 0.45 x25 mm, Sterican), allowing the odorant to diffuse through a hole (diameter: 5 mm) in the platform through the Teflon fabric and into the arena. Each arena had two odorant sources. One odorant was used as sucrose-paired conditioned stimulus (CS+) and the other odorant was used as unpaired conditioned stimulus (CS-). At the location of the CS+, 20 µl of oversaturated sucrose-ethanol solution was pipetted onto the fiberglass fabric

and blow-dried for 20 minutes, producing a thin layer of pure sucrose on a round patch with

a diameter of 10 mm. The position of the CS+ and CS- were always switched between the

conditioning and the test (e.g. if CS+ was at the inside position during conditioning, it was at

the outside position during the test, and in half of the experimental runs the CS+ was at the

inside position during conditioning, and in the other half at the outside position); each

odorant was used equally often as CS+ and CS-. To change the floor between experimental

phases, the platform was rotated underneath the arenas; the arenas themselves did not

move. The angular rotation speed of the platform was 360°/25 s which corresponded to a

speed of 2.6 cm/s in the center of the arena (the distance between center of the table and

the center of the arena is 10.25 cm). The conditioning apparatus was placed in an air suction

hood in order to remove odorants. All experiments were performed in the dark to eliminate

visual stimuli. The arena was back-illuminated with infrared light (850 nm, SOLAROX LED

Strip), which is not visible to flies, and experiments were video recorded with an infrared

sensitive camera (infrared Camera Module v2, Pi NoiR, connected to a Raspberry Pi 3, Model

B V1.2) at 5 frames/s. The rotating motor was controlled via TTL pulses through the

Raspberry Pi. The rotation of the table and the video recordings were controlled with

custom-written software in Python (Stefanie Neupert).

Odor-food conditioning

All experiments were done between 10:00 and 12:00 or after 15:00 during periods when

flies show higher foraging activity (Breugel et al., 2017). Each experimental run contained 4

differently sized groups ("single", "pair", "small group", "large group"), and the positions of

the 4 arenas used for the 4 differently sized groups were balanced across experimental runs.

One experimental run consisted of four phases:

Acclimatization (Figure 1B, solid arcs): Flies were sucked out from the starvation vials using a

tube aspirator and placed into an arena that had no odorant source and were allowed to

acclimatize for 10 minutes.

Conditioning (Figure 1B, dotted arcs): The floor was rotated counterclockwise by 22.5° and

the CS+ paired with sucrose and the CS- without sucrose were presented for 7 minutes.

Pause (Figure 1B, dashed arcs): The floor was rotated by 22.5° and replaced by a new floor

without odorants or sucrose. The pause lasted for 5 seconds.

<u>Test:</u> The floor was rotated by 22.5° and replaced by a floor which had the CS+ and CS- but

without sucrose. CS+ and CS- positions were switched from conditioning. The test phase

lasted for 7 minutes.

Videos were recorded during the conditioning and test. After each experimental run, all flies

were discarded and the Teflon fabric floor was rinsed with hot water and soap to remove the

odorants and the sucrose patch.

Fly tracking

Video recordings were analyzed using the software Fiji (ImageJ 1.51s Wayne Rasband NIH,

USA). We removed the first 30 frames due to compression artifacts and converted the video

to grayscale. Then we did a Z projection to get the maximum intensity projection over the

whole video, and calculated the difference per frame between the maximum intensity

projection and the original video. This gave us a clear image of flies moving around the arena

for tracking. We used this output to track flies using the plugin TrackMate (Version: 3.5.3,

Tinevez et al., (2017)). We used a Downsample LoG detector to identify flies (blob diameter

= 13 pixels, downsampling = 3, threshold = 6-8). To generate the tracks, we used the Simple

LAP Tracker, with the following parameters: linking distance = 150 pixels, max gap closing =

150 pixels and maximal frame gap = 3 frames. For the conditioning data, we only extracted

the x and y coordinates of each fly per frame. For the test data, we extracted the x and y

coordinates per frame as well as the identity of the fly throughout the recording. We

inspected all tracking results visually and corrected the tracks manually to connect the

missing links and afterwards we extracted the x and y coordinates for the analysis.

Data analysis

Normalizing arenas for comparison

For both the conditioning and the test datasets, we centralized each arena so that the center

point of the circular arena was at (0, 0). The center point was determined by taking the

midpoint between the CS+ and CS- locations; the x and y coordinates of the CS+ and CS-

were recorded manually. We then converted each Cartesian coordinate to polar

coordinates, in order to rotate each arena so that the CS+ location was at the top of the

arena and the CS- was at the bottom. We took the distance of the CS+ to the center as a

reference radius of 1, and normalized all coordinates to this radius. We then filtered out any

points that had a radius equal to or greater than 1.3 to remove tracking errors. Note that for

the conditioning dataset, only the x and y coordinates of a fly per frame were recorded. For

the test dataset, the x and y coordinates per frame were recorded, but also the identity of

the fly across frames. Thus the analysis for the two datasets differs slightly.

Visit probability maps

Visit probability maps were generated only for the test dataset. For every individual fly, we

divided the arena into 20 x 20 pixel bins. For each frame, we gave the pixel bin that

contained the coordinate of the fly a score of 1, and gave all of the other bins a score of 0.

We summed the scores of each pixel bin over all frames and then normalized by the track

length for each individual. For each group size, we then took the mean of each pixel bin over

all individual fly tracks. We used the same analysis for the time-binned visit probability maps

by looking only at the frames that occurred during the time bin.

Calculating the distance of individual flies to the CS+ and CS-

164 We calculated the distance of each fly to the CS+ and to the CS- in every frame using:

165 Euclidean distance = $\sqrt{(x - x_0)^2 + (y - y_0)^2}$

166 Where x and y are the Cartesian coordinates of the fly, and x_0 and y_0 are the Cartesian

167 coordinates of either the CS+ or the CS-.

Approach probability

For the conditioning data, the coordinates of the flies were tracked for each frame. For every

frame of the experiment, we scored each point: if the fly was closer to the CS+ than the CS-,

it was given a score of 1, otherwise it was given a score of 0. We then divided the data into

one-minute time bins. For every experiment we calculated the number of ones and zeros in

each time bin; if there were more ones than zeros, the time bin was given a score of 1 for

that experiment, otherwise it was given a score of 0. We then took the mean for each time

bin over all experimental runs.

For the test data, we used a similar method, however since we had the fly identities over the

whole video, we calculated the approach probability using the individual fly tracks. This

allowed us to calculate the approach score for each individual fly. A fly received a score of 1

when she spent more time closer to the CS+ than the CS- and a 0 when she spent more time

closer to the CS- than to the CS+ during the experimental run. If more flies in a particular

experimental run scored a 1 than scored a 0, then the experimental run was scored a 1,

otherwise it was scored a 0. Moreover, we divided each track into one-minute time bins and

for each experimental run each time was scored 1 (flies were closer to the CS+) or 0 (flies

were closer to the CS-).

Latency to the CS+ and the CS-

For the test dataset, we identified for each individual the first frame (latency) that the

individual was 0.5 cm or closer to the center of the CS+ or CS-. We firstly took the mean

latency to the CS+ or CS- across all flies in one experimental run, and then took the mean

across the 30 experimental runs for each group size.

Calculating the distance between flies

For the test dataset, we used the rotated Cartesian coordinates to calculate the Euclidean

distance between every fly in each frame of the experiment. We divided the distances into

bins of 5 mm and counted the occurrences of each distance per experimental run.

Simulating distances between flies due to chance

For the test dataset, we selected flies according to their group size, and randomly sampled

entire fly tracks from different experimental runs of the same experiment. We simulated as

many experimental runs as there were real experimental runs, and we also simulated as

many flies as were in each experimental run. We then overlaid these tracks and calculated

the Euclidean distance between flies for the simulated experiments as we did for the real

experiments.

Calculating encounters between flies

We defined an encounter as the center of one fly being maximum two fly lengths (5 mm)

away from the center of another fly. We calculated the Euclidean distance between all flies

as before. We selected the distances that were less than or equal to 5 mm (the encounter

distances). Since we had the identity of every fly per experimental run, we could calculate

the number of encounters for every fly and the length of these encounters.

For the mean encounter number per fly per experimental run, we calculated the total

number of encounters for one experimental run, multiplied it by two as there were two flies

involved in each encounter, and then divided it by the number of flies in the arena. For the

mean encounter length per experimental run, we summed the length of all encounters and

divided by the total number of encounters per experimental run. We repeated this analysis

for the simulated data.

Statistical analysis

For all data analysis, R version 3.5.0 was used (R Core Team, 2018). All statistics were

performed using Bayesian data analysis, based on Korner-Nievergelt et al., (2015).

To investigate the effect of group size on conditioned approach probability for the CS+, we

used a binomial generalized linear model (GLM), with conditioned approach probability as

the binary response variable (1 = approach, 0 = no approach). We used the logistic

regression (logit) link function. The group size ("single", "pair", "small group" and "large

group") was used as the explanatory variable. We used an improper prior distribution (flat

prior) and simulated 100 000 values from the posterior distribution of the model parameters

using the function "sim" from the package "arm". The means of the simulated values from

the posterior distributions of the model parameters were used as estimates, and the 2.5 %

and 97.5 % quantiles as the lower and upper limits of the 95 % credible intervals. We used

this GLM to compare the conditioned approach probability of each group size against

chance. For each group size, we calculated the proportion of simulated values from the

posterior distribution that were larger than 0.5. A value of 0.5 indicates that flies spend

equal time in the two halves of the arena, whereas a value of 1 indicates that flies spent all

of the time in the half of the arena containing the CS+, and a value of 0 indicates that flies

spent all of the time in the half of the arena containing the CS-. If the proportion of

simulated values was significantly greater than 0.5, we determined that flies spent more

time in the half of the arena containing the CS+ than predicted by chance (represented by

filled circles in plots). To test for differences between conditioned approach probabilities for

different group sizes, we calculated the proportion of simulated values from the posterior

distribution that were larger for one group compared to another group. We declared an

effect to be significant if the proportion was greater than 0.95 (*). Proportions greater than

0.99 are marked "**" and greater than 0.999 marked "***". We performed this analysis for

the whole recording: for the different time bins, we compared the conditioned approach

probability of the different group sizes within a single time bin, not between them.

To test whether the latency to reach the CS+ or CS- depends on group size, we used a linear model (LM). The latency to the CS+ or to the CS- was the response variable, and the group size was used as the explanatory variables. We used the same methodology as previously used in the GLM to simulate values from the posterior distribution and generate the means and the 95 % credible intervals. To test for differences, we calculated the proportion of draws from the posterior distribution for which the mean of each draw was smaller in the experimental dataset than the mean of each draw of the simulated dataset. We declared an effect to be significant if the proportion was greater than 0.95.

To investigate whether grouped flies differ in the number of their inter-fly encounters from random (simulated data), we used an LM for each distance bin. The number of occurrences of that distance was the response variable, and the type of data (experimental or simulated data) was used as the explanatory variable. We used the same method as specified above to test for differences.

To investigate whether the encounter number and lengths were different to random (simulated data), we used an LM with either encounter number or encounter length as the response variable, and the type of data (experimental or simulated data) as the explanatory variables. We used the same method as specified above to test for differences.

To determine whether the mean encounter number per fly differed between group sizes, we randomly assigned pairs of experimental and simulated encounter numbers from different experimental runs for each group size (Figure 2F). For each pair, we then subtracted the simulated encounter number value from the real encounter number value (difference between encounters). This allowed us to compare between group sizes as by removing the simulated value, we remove the number of encounters that could be due to chance, which is correlated with group size. To test for differences between group sizes, we used an LM. The response variable was the "difference between encounters". The explanatory variable was the different group size ("pair", "small group" and "large group"). The large group was used as the reference level. We used the same method as specified above to draw inferences about the differences between the large group and the other two group sizes.

RESULTS

To investigate whether group size affects associative odor-food memory in flies, we developed an automated conditioning assay to condition four groups of flies simultaneously (Figure 1A, B). The learning assay consisted of a computer-controlled rotating table platform with four concentric arenas. This design allowed us to transfer flies from conditioning to the test without anesthesia and with minimal mechanical disturbance, which can alter fruit flies' behavior (Barron, 2000; Bartholomew et al., 2015; Trannoy et al., 2015). We compared four different sized groups of flies: one fly (single), two flies (pair), three or four flies (small group) and seven or eight flies (large group) (Figure 1C). Both the training phase and test phase lasted for 7 minutes. During the 7-minutes long conditioning, two odorants (2,3-butanedione and ethyl acetate) were presented at opposite sides of the arena; one odorant (conditioned stimulus, CS+) was paired with dried sucrose and the other odorant was not (CS-). During conditioning, flies were allowed to forage around the arena and find the food and odorant source.

Walking flies aggregate on the odorant-sucrose patch during conditioning and learn to associate the odorant with sucrose

Flies of all group sizes showed a high probability to approach the odorant (CS+)-sucrose patch, implying that they were feeding on the sucrose (mean approach probability over the last minute- single, small group and large group: >0.999, pair: 0.97) (Figure 1D). For a given experimental run the approach probability was 1 if the flies spent more time closer to the CS+ than to the CS- and 0 if the fly spent more time closer to the CS- than to the CS+. We did this for one-minute time bins, and took the mean for each bin across experimental runs to get a mean approach probability per group size.

Figure 1: Automated odor-food conditioning assay.

(**A**) Conditioning assay for conditioning and testing four groups of flies in parallel.

(**B**) Conditioning assay viewed from above through the tracking camera. The locations of the CS+ and CS- in each arena are indicated. The arcs indicate the position of the arena in different experimental phases. Full lines outline the acclimatization phase, dotted lines outline the conditioning phase, and dashed lines outline the pause phase before the test.

(**C**) Same as (B) but with examples of individual fly trajectories during the test overlaid. Trajectories are shown for a single fly (top left), a pair of flies (top right), a group of 4 flies (small group, bottom right) and a group of 8 flies (large group, bottom left).

(**D**) Approach probability for the odorant (CS+)-sucrose patch for the single flies (red), the pairs (blue), the small group (yellow) and the large group (pink) during conditioning. Each minute bin shows the mean of the binary approach scores across all experimental runs (N= 30 experimental runs). The dashed line

represents the approach probability of 0.5 (chance).
(**E**) Visit probability maps for each group size (columns) for the entire duration of the recording, the first minute
of recording and the second minute of recording (rows). Each bin shows the mean binary value across all
experimental runs. N = 30 experimental runs for all groups; single flies: N = 30 individuals; pairs: N = 60
individuals; small group: N = 118 individuals; large group: N = 229 individuals.

Flies were transferred from the conditioning to the test by rotating the platform (Figure 1B).

In between conditioning and test there was a pause, where the platform was rotated to a

neutral segment where there were no odorants or sucrose present. The pause lasted 5 s,

and then platform was rotated to the test segment. During the test, the positions of the CS+ and the CS- were switched and there was no sucrose, thus flies could not rely on remembering the location of the sucrose patch (Kim and Dickinson, 2017) and had to follow the olfactory CS+ to search for the expected food. To visualize conditioned approach to the CS+, we projected the flies' trajectories on a plane and calculated the probability across flies to visit a particular pixel bin (Figure 1E). Throughout conditioning, flies of all group sizes showed highest visit probabilities to areas around the CS+. The increased visit probabilities around the CS+ persisted over the entire 7 minutes of the test, confirming that flies learned to associate the CS+ with food.

Flies conditioned and tested in groups exhibit extended associative odor-food memory compared to flies conditioned alone or in pairs

We next asked whether group size affects the expression of associative odor-food memory during the test and determined the conditioned approach probability for the CS+ (Figure 2A). We calculated the conditioned approach probability for the CS+ by the following steps: 1) If a fly spent more time near the CS+ than the CS-, it scored a one, otherwise it scored a zero. 2) If an experimental run had more flies scored with one than zero, the experimental run itself was scored as 1, otherwise zero. 3) We took the mean across experimental runs to get a conditioned approach probability per group size. During the test, all group sizes showed a conditioned approach probability for the CS+ that was significantly higher than chance (*p(large group > 0.5)* > 0.999, *p(small group > 0.5)* > 0.999, *p(pair > 0.5)* = 0.983, *p(single > 0.5)* > 0.999), showing that they had formed an associative odor-food memory (Figure 2A). Both the large group and the small group showed significantly higher conditioned approach probabilities than the pair group (*p(large group > pair)* = 0.983, *p(small group > pair)* = 0.968), however there were no differences between any of the other group sizes (see Table S1). There was no differences between group sizes for how quickly the flies approached the CS+ (Figure S1A), however the pair approached the CS- significantly faster than the single fly or the small group (*p(single > pair)* = 0.982, *p(small group > pair)* = 0.981) (Figure S1B).

Figure 2: Group size affects associative odor-food memory expression and inter-fly encounters during the test.
(**A**) Conditioned approach probability for the CS+ for the single fly group (red), the pair group (blue) the small group (yellow) and the large group (pink) during the test. Bars represent the mean binary score across all experimental runs. Whiskers represent the 95 % credible intervals. The dashed line indicates the conditioned approach probability due to chance (0.5). Stars represent significant differences in conditioned approach probability between groups.
(**B**) Conditioned approach probability for the CS+ over one-minute time bins during the test. Colors, dashed line and whiskers are the same as in (A). The points represent the mean binary score across all experimental runs within a time bin. Filled points are conditioned approach probabilities that are significantly different to chance, empty points are not significantly different to chance. For statistical comparisons between groups, see Table S1.
(**C**) Mean inter-fly encounter number per fly per experimental run for the pair (blue) and the simulated pair

(grey). Bars represent the mean across experimental runs. Vertical lines represent the 95 % credible intervals.
(**D**) Same as (C) for the small group (yellow) and the simulated small group (grey).
(**E**) Same as (C) for the large group (pink) and the simulated large group (grey). Stars represent significant
differences in mean encounter length between the real and simulated group (**: *p(experimental > simulated) >*
0.99).
(**F**) Paired differences between mean encounter numbers across experimental runs. Stars represent significant
differences between groups (*: *p(large group > small group) > 0.95*, **: *p(large group > pair) > 0.999*). Black
circles represent the mean difference in encounter number between experimental runs. Colors and whiskers
are the same as in A.
424

To investigate the time course of memory expression, we calculated the conditioned

approach probability over one-minute time bins (Figure 2B). During the first minute of the

test, the single fly showed a higher conditioned approach probability than the pair (*p(pair >*

single) = 0.031). Between the 2nd to 7th minutes, the large group showed higher conditioned

approach probabilities than both the single fly group and the pair group throughout most

other bins tested (Table S1). The large group also showed higher conditioned approach

probability than the small group during the 4th minute and the 6th minute. The pair group

showed the lowest conditioned approach probability which was only significantly different

to chance during the second minute (2nd minute: *p(pair > 0.5)* = 0.997), and during the 3rd to

7th minutes, the paired group showed a significant lower approach probability than the large,

small and single fly groups (Table S1, S2). These results suggest that flies conditioned and

tested in a large group express a longer associative odor-food memory than flies conditioned

and tested in smaller groups or single flies, and flies conditioned and tested in pairs express

shorter associative odor-food memory than single flies or flies in groups.

Flies tested in groups – but not in pairs – exhibit more inter-fly encounters than random

We next asked whether flies socially interact, which could potentially allow them to share

information with each other about the location of the predicted food source and thus could

explain the extended associative memory of the large group. To assess whether group size

affects the frequency of social interactions we measured the number of inter-fly distances

and compared it with the number of expected random distances. We calculated the

distances between all flies conditioned in groups in each video frame during conditioning to

see if they approached each other. To determine whether these distance distributions could

be explained merely due to flies randomly encountering each other in the arena, we

simulated 30 new experimental runs by randomly sampling fly locations from all

experimental runs for each video frame. We then calculated the distances between these

simulated groups of flies in each video frame (Figure S2A, Table S3). For the large group

there were more short inter-fly distances (0-5 and 5-10 mm) than for the simulated group.

We chose a distance of 5 mm between fly centers as a threshold for inter-fly encounters

where flies could potentially socially interact. Flies in the small and large group made

significantly more encounters (approached each other by 5 mm or less) than the simulated

groups of flies, but not the pair (*p(large group > simulated large group)* > 0.999, *p(small*

group > simulated small group) = 0.996, *p(pair > simulated pair)* = 0.917) (Figure 2C-E).

To compare encounter number across group sizes, we needed to correct for trivial

differences in encounters that are just due to differences in the group sizes (in larger groups

there is a higher chance for random inter-fly encounters). We corrected for these trivial

differences in encounter number by the following procedure: We randomly took an experimental run from the experimental and simulated datasets and subtracted the number of encounters between the two experimental runs (Figure 2F). By subtracting the number of encounters in the simulated runs, we removed the number of encounters per experimental run that could be due to random encounters. The encounter number was significantly higher for the large group compared to the small group and the pair (*p(large group > pair)* > 0.999, *p(large group > small group)* = 0.996). There were no differences in encounter length between any group size and their simulated groups (Figure S2B).

The increased number of encounters in the larger group indicates that flies are more attracted to each other when they are in large groups than when they are in small groups or pairs. More encounters allow more opportunities for social interactions between flies, which in turn could underlie the longer associative memory expression of the large group as compared to smaller groups or single flies.

DISCUSSION

We developed an automated conditioning assay for walking fruit flies that allows analyzing the behavior of individual flies during foraging. We used this assay to investigate how social interactions affect odor-food learning in single flies and in groups of flies. Flies in groups showed extended expression of the associative odor-food memory as compared to single flies or pairs of flies. Moreover, flies in larger groups were more attracted to each other than flies in smaller groups. These data suggest that foraging in groups of more than two flies facilitates social information transfer about the quality of a food source (during odor-food learning) or about the location of the predicted food source (during odor-guided food search).

Social information transfer during foraging

Fruit flies accumulate on fermenting fruit which they find by following the odorants released by fermenting fruit (Becher et al., 2012; Kellogg et al., 1962; Semmelhack and Wang, 2009) and by following aggregation pheromones released by males (Bartelt et al., 1985; Lin et al., 2015; Mercier et al., 2018) and females (Lebreton et al., 2017). In the group, fruit flies transfer information to other flies about their internal state, such as stress (Suh et al., 2004)

and about the location and quality of resources during mate choice (Danchin et al., 2018; Mery et al., 2009), egg laying (Battesti et al., 2012b; Durisko and Dukas, 2013; Lin et al., 2015; Sarin and Dukas, 2009), and foraging (Lihoreau et al., 2016; Tinette et al., 2004). During foraging, primer flies search for the most favorable food patch and other flies aggregate at those food patches after socially interacting with the primer flies (Tinette et al., 2004), thus through local enhancement of food patches, the entire group of flies appears to benefit from the transmitted information from primer flies.

Since fruit flies forage, lay eggs and mate during aggregation, it is plausible that they have adapted to transmit information to each other about the environment through social interactions. Our finding of extended expression of associative odor-food memories in groups together with the positive correlation between group size and inter-fly attraction suggests that associative odor-food learning or memory expression also benefits from social information transfer during aggregation. The positive correlation between group size and inter-fly attraction that we found is in line with a previous study where the number of inter-fly attraction was higher in larger than in smaller groups (20-40 versus 10 flies) (Simon et al., 2012). To our knowledge, such an increase in inter-animal attraction with increasing group size has not yet been reported in vertebrates (Miller and Stephen, 1966).

Social effects on odor-food learning

While we found extended memory expression in larger groups, we found shorter memory expression in pairs of flies as compared to single flies or groups of 4 or 8 flies. This finding was surprising, and we have no good explanation for this phenomenon. We will therefore constrain our discussion the social effects on odor-food learning and memory in groups larger than two.

The extended expression of associative odor-food memory in groups of flies could be a result of social information transfer during the formation of the odor-food memory (during conditioning) or during the retrieval of the odor-food memory (during the memory test). Since our experiments were performed in the dark, flies could have transferred information socially via olfactory stimuli (Jallon, 1984; Keesey et al., 2016; Lebreton et al., 2017; Lin et al., 2015), gustatory stimuli (Schneider et al., 2012) sound (Tauber and Eberl, 2003), substrate-borne vibration (Fabre et al., 2012) and touch (Ramdya et al., 2014), but not via visual cues

(Danchin et al., 2018; Ferreira and Moita, 2019; Golden and Dukas, 2014; Mery et al., 2009; Sarin and Dukas, 2009).

During conditioning, the presence of other flies at the sucrose patch could increase the reinforcing strength of the sucrose since the presence of other flies indicates that the food patch is good. Indeed, flies prefer food sources with other flies present over food sources without any flies (Lihoreau et al., 2016; Tinette et al., 2004). Alternatively, the presence of other flies at the sucrose patch may be an additional appetitive reinforcing stimulus. The role of conspecifics as a positive reinforcer has been previously demonstrated in honey bees, where antennal touching of a nestmate acts as positive reinforcer during odor conditioning (Cholé et al., 2019). The fact that fruit flies are attracted to each other (Lefranc et al., 2001; Simon et al., 2012; Tinette et al., 2004) makes it plausible to assume that flies could also act as an additional positive reinforcer at the sucrose patch.

The reinforcing function of other flies could be mediated by dopaminergic neurons, because dopaminergic neurons mediate the reinforcing function of sucrose (Liu et al., 2012) and because dopamine itself has an effect on the sociality of flies: inter-fly attraction decreases in flies that have a deficiency in dopamine released from neurons and hypodermal cells (Fernandez et al., 2017).

Social effects on odor-food memory expression

Besides there being a learning effect, the extended odor-food memory expression could be a memory retrieval effect. Memory retrieval could be affected by the social interactions during the memory test, as flies that had learned the association between CS+ and sucrose could transfer information about the location of the predicted sucrose patch to flies that have failed to learn. Information transfer from experienced to naïve flies can affect group level behavior during odor avoidance (Ramdya et al., 2014), aversive memory retrieval (Chabaud et al., 2009), mate choice (Danchin et al., 2018; Mery et al., 2009), oviposition site choice (Battesti et al., 2012b; Sarin and Dukas, 2009) and predator-induced egg-retention (Kacsoh et al., 2015). Moreover, a theoretical study predicted that social interactions can increase performance during odor-guided foraging (Torney et al., 2009).

Alternatively, flies that located the CS+ first during the test could serve as an attractive reinforcing stimulus (see discussion above and (Lihoreau et al., 2016; Tinette et al., 2004)),

thus appetitive learning of the CS+ could continue throughout the test. This ongoing appetitive learning of the CS+ during the test would appear as extended associative memory expression in our study.

Another possible explanation for the extended memory expression could be reduced memory extinction due to social interaction at the CS+. Memories can be extinguished when the CS+ is presented without reinforcement (Lagasse et al., 2009; Schwaerzel et al., 2002), which is effectively what occurs throughout the test in our study. In flies, extinction of odor-sucrose memories is mediated by dopaminergic neurons that encode reinforcement during aversive odor learning (Felsenberg et al., 2017): lack of reward during CS+ induced memory retrieval activates those dopaminergic neurons, and this activation counteracts the associative odor-food memory. The presence of other flies at the CS+ could provide an appetitive stimulus and thereby prevent the activation of these extinction mediating dopaminergic neurons.

Technical considerations and outlook

We analyzed the walking behavior of individual flies during the memory test, but not during the conditioning because we could not separate flies from each other when they clustered at the sucrose patch due to a lack of spatial resolution. By using cameras with higher spatial resolution, this assay can be extended to a high-throughput assay for multiple parallel fly groups that would allow automatic tracking of all individuals, classifying of patterns of pairwise and higher-order interactions between individuals, as well as stereotyped behaviors in individuals (Berman et al., 2016; Branson et al., 2009). This assay could help reveal external factors (e.g., fly density, the ratio of informed to uninformed flies) and internal factors (e.g., metabolic, genetic, or circadian states) that influence learning in social contexts, and it would also allow studying the neural basis of social effects on foraging, by disentangling sensory processing and memory formation. To identify the sensory bases of information transmission between flies, one could test the effect of temporarily perturbing their ability to smell, see and mechanosense by expressing a temperature-sensitive switch for synaptic transmission in defined neuron populations (Kitamoto, 2001). Likewise, neuronal perturbation experiments would help identifying the neurons that encode the valence of social information and reveal how these neurons integrate with the neurons known to encode the hedonic and caloric value of food (Huetteroth et al., 2015). Moreover,

to investigate whether information transmission during foraging is affected by the fly's predisposition to forage, one could use the two naturally occurring *foraging* gene *Drosophila* mutants. "Rovers" move more during foraging and demonstrate improved short term memory, whereas "sitters" move less and show an improved long term memory (Mery et al., 2007; Osborne et al., 1997). Since both foraging and aversive memory expression are affected by social context (Kohn et al., 2013), experiments using these morphs would help to assess whether improved memory expression or learning during foraging in groups is genetically predetermined.

CONFLICT OF INTEREST

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

AUTHOR CONTRIBUTIONS

PS conceptualized and designed the study. CT performed the data collection. CT and YM performed the pilot experiments. AS, CT and YM prepared the video data for analysis. AS performed the statistical analysis. AS, PS and YM wrote the manuscript. PS supervised the study.

ACKNOWLEDGEMENTS

We thank C. Giovanni Galizia for discussions initiating this research, Stefanie Neupert for programming the acquisition software and comments on the manuscript, Jana Hörsch for contributing to pilot experiments and comments on the manuscript, Charles Ellen for comments on the manuscript, FIBERFLON (Konstanz) for donating Teflon-coated fiberglass fabric.

REFERENCES

Abu, F., Wang, J. G., Oh, Y., Deng, J., Neubert, T. A. and Suh, G. S. B. (2018). Communicating the nutritional value of sugar in *Drosophila*. *Proc. Natl. Acad. Sci.* **115**, 201719827.

Alem, S., Perry, C. J., Zhu, X., Loukola, O. J., Ingraham, T., Søvik, E. and Chittka, L. (2016). Associative Mechanisms Allow for Social Learning and Cultural Transmission of String Pulling in an Insect. *PLOS Biol.* **14**, e1002564.

Avarguès-Weber, A. and Chittka, L. (2014). Local enhancement or stimulus enhancement? Bumblebee social learning results in a specific pattern of flower preference. *Anim. Behav.* **97**, 185–191.

Barron, A. B. (2000). Anaesthetising Drosophila for behavioural studies. *J. Insect Physiol.* **46**, 439–442.

Bartelt, R. J., Schaner, A. M. and Jackson, L. L. (1985). cis-Vaccenyl acetate as an aggregation pheromone in Drosophila melanogaster. *J. Chem. Ecol.* **11**, 1747–1756.

Bartholomew, N. R., Burdett, J. M., Vandenbrooks, J. M., Quinlan, M. C. and Call, G. B. (2015). Impaired climbing and flight behaviour in Drosophila melanogaster following carbon dioxide anaesthesia. *Sci. Rep.* **5**, 1–10.

Battesti, M., Moreno, C., Joly, D. and Mery, F. (2012a). Spread of Social Information and Dynamics of Social Transmission within Drosophila Groups. *Curr. Biol.* **22**, 309–313.

Battesti, M., Moreno, C., Joly, D. and Mery, F. (2012b). Spread of social information and dynamics of social transmission within Drosophila groups. *Curr. Biol.* **22**, 309–313.

Becher, P. G., Flick, G., Rozpędowska, E., Schmidt, A., Hagman, A., Lebreton, S., Larsson, M. C., Hansson, B. S., Piškur, J., Witzgall, P., et al. (2012). Yeast, not fruit volatiles mediate Drosophila melanogaster attraction, oviposition and development. *Funct. Ecol.* **26**, 822–828.

Berman, G. J., Bialek, W. and Shaevitz, J. W. (2016). Predictability and hierarchy in Drosophila behavior. *Proc. Natl. Acad. Sci. U. S. A.* **113**, 11943–11948.

Branson, K., Robie, A. A., Bender, J., Perona, P. and Dickinson, M. H. (2009). High-throughput ethomics in large groups of Drosophila. *Nat. Methods* **6**, 451–7.

Breugel, F. van, Huda, A. and Dickinson, M. H. (2017). Drosophila have distinct activity-gated pathways that mediate attraction and aversion to CO2. *bioRxiv* 227991.

Burke, C. J., Huetteroth, W., Owald, D., Perisse, E., Krashes, M. J., Das, G., Gohl, D., Silies, M., Certel, S. and Waddell, S. (2012). Layered reward signalling through octopamine and dopamine in Drosophila. *Nature* **492**, 433–437.

Chabaud, M. A., Isabel, G., Kaiser, L. and Preat, T. (2009). Social Facilitation of Long-Lasting Memory Retrieval in Drosophila. *Curr. Biol.* **19**, 1654–1659.

Cholé, H., Carcaud, J., Mazeau, H., Famié, S., Arnold, G. and Sandoz, J.-C. (2019). Social Contact Acts as Appetitive Reinforcement and Supports Associative Learning in Honeybees. *Curr. Biol.* **29**, 1–7.

Czaczkes, T. J., Grüter, C., Jones, S. M. and Ratnieks, F. L. W. (2011). Synergy between social and private information increases foraging efficiency in ants. *Biol. Lett.* **7**, 521–524.

Danchin, E., Nöbel, S., Pocheville, A., Dagaeff, A.-C., Demay, L., Alphand, M., Ranty-Roby, S., van Renssen, L., Monier, M., Gazagne, E., et al. (2018). Cultural flies: Conformist social learning in

fruitflies predicts long-lasting mate-choice traditions. *Science (80-.).* **362**, 1025–1030.

Durisko, Z. and Dukas, R. (2013). Attraction to and learning from social cues in fruitfly larvae. *Proc R*
Soc B **280**, 20131398.

Fabre, C. C. G., Hedwig, B., Conduit, G., Lawrence, P. A., Goodwin, S. F. and Casal, J. (2012).
Substrate-Borne Vibratory Communication during Courtship in Drosophila melanogaster. *Curr.*
Biol. **22**, 2180–2185.

Felsenberg, J., Barnstedt, O., Cognigni, P., Lin, S. and Waddell, S. (2017). Re-evaluation of learned
information in Drosophila. *Nature* **544**, 240–244.

Fernandez, R. W., Akinleye, A. A., Nurilov, M., Feliciano, O., Lollar, M., Aijuri, R. R., O'Donnell, J. M.
and Simon, A. F. (2017). Modulation of social space by dopamine in Drosophila melanogaster,
but no effect on the avoidance of the Drosophila stress odorant. *Biol. Lett.* **13**,.

Ferreira, C. H. and Moita, M. A. (2019). Safety in numbers is mediated by motion cues and depends
on lobula columnar neurons in Drosophila melanogaster. *bioRxiv* 629311.

Frisch, K. Von (1965). *Die tänze der Bienen*.

Galizia, C. G. (2014). Olfactory coding in the insect brain: Data and conjectures. *Eur. J. Neurosci.* **39**,
1784–1795.

Giraldeau, L.-A. and Caraco, T. (2000). *Social foraging theory*. Princeton University Press.

Golden, S. and Dukas, R. (2014). The value of patch-choice copying in fruit flies. *PLoS One* **9**,.

Haverkamp, A., Hansson, B. S. and Knaden, M. (2018). Combinatorial Codes and Labeled Lines: How
Insects Use Olfactory Cues to Find and Judge Food, Mates, and Oviposition Sites in Complex
Environments. *Front. Physiol.* **9**, 49.

Huetteroth, W., Perisse, E., Lin, S., Klappenbach, M., Burke, C. and Waddell, S. (2015). Sweet taste
and nutrient value subdivide rewarding dopaminergic neurons in drosophila. *Curr. Biol.* **25**, 751–
758.

Itskov, P. M. and Ribeiro, C. (2013). The dilemmas of the gourmet fly: the molecular and neuronal
mechanisms of feeding and nutrient decision making in Drosophila. *Front. Neurosci.* **7**, 12.

Jallon, J.-M. (1984). A few chemical words exchanged by Drosophila. *Behav. Genet.* **14**, 441–478.

Kacsoh, B. Z., Bozler, J., Ramaswami, M. and Bosco, G. (2015). Social communication of predator-
induced changes in Drosophila behavior and germ line physiology. *Elife* **4**,.

Keesey, I. W., Koerte, S., Retzke, T., Haverkamp, A., Hansson, B. S. and Knaden, M. (2016). Adult
Frass Provides a Pheromone Signature for Drosophila Feeding and Aggregation. *J. Chem. Ecol.*
42, 739–747.

Kellogg, F., Frizel, D. and Wright, R. (1962). The Olfactory Guidance of Flying Insects. IV. Drosophila.
Can. Entomol. **94**, 884–888.

Kim, I. S. and Dickinson, M. H. (2017). Idiothetic Path Integration in the Fruit Fly Drosophila
melanogaster. *Curr. Biol.* **27**, 2227-2238.e3.

Kitamoto, T. (2001). Conditional modification of behavior in Drosophila by targeted expression of a
temperature-sensitive shibire allele in defined neurons. *J. Neurobiol.* **47**, 81–92.

Kohn, N. R., Reaume, C. J., Moreno, C., Burns, J. G., Sokolowski, M. B. and Mery, F. (2013). Social
Environment Influences Performance in a Cognitive Task in Natural Variants of the Foraging
Gene. *PLoS One* **8**, e81272.

Korner-Nievergelt, F., Roth, T., von Felten, S., Guelat, J., Almasi, B., Korner-Nievergelt, P., Guélat, J.,
Almasi, B. and Korner-Nievergelt, P. (2015). *Bayesian data analysis in ecology using linear*
models with R, BUGS, and Stan. Academic Press.

Lagasse, F., Devaud, J.-M. and Mery, F. (2009). A Switch from Cycloheximide-Resistant Consolidated
Memory to Cycloheximide-Sensitive Reconsolidation and Extinction in Drosophila. *J. Neurosci.*
29, 2225–2230.

Leadbeater, E. and Dawson, E. H. (2017). A social insect perspective on the evolution of social
learning mechanisms. *Proc. Natl. Acad. Sci.* **114**, 7838–7845.

Lebreton, S., Borrero-Echeverry, F., Gonzalez, F., Solum, M., Wallin, E. A., Hedenström, E., Hansson,
B. S., Gustavsson, A. L., Bengtsson, M., Birgersson, G., et al. (2017). A Drosophila female
pheromone elicits species-specific long-range attraction via an olfactory channel with dual
specificity for sex and food. *BMC Biol.* **15**, 88.

Lefranc, A., Jeune, B., Thomas-Orillard, M. and Danchin, E. (2001). Non-independence of individuals
in a population of Drosophila melanogaster: effects on spatial distribution and dispersal. *C. R.*
Acad. Sci. III. **324**, 219–27.

Lihoreau, M., Clarke, I. M., Buhl, J., Sumpter, D. J. T. and Simpson, S. J. (2016). Collective selection
of food patches in Drosophila. *J. Exp. Biol.* **219**, 668–675.

Lin, C.-C., Prokop-Prigge, K. A., Preti, G. and Potter, C. J. (2015). Food odors trigger Drosophila males
to deposit a pheromone that guides aggregation and female oviposition decisions. *Elife* **4**, 1–26.

Liu, C., Plaçais, P.-Y., Yamagata, N., Pfeiffer, B. D., Aso, Y., Friedrich, A. B., Siwanowicz, I., Rubin, G.
M., Preat, T. and Tanimoto, H. (2012). A subset of dopamine neurons signals reward for odour
memory in Drosophila. *Nature* **488**, 512–516.

Mercier, D., Tsuchimoto, Y., Ohta, K. and Kazama, H. (2018). Olfactory Landmark-Based
Communication in Interacting Drosophila. *Curr. Biol.* **28**, 2624-2631.e5.

Mery, F., Belay, A. T., So, A. K.-C., Sokolowski, M. B. and Kawecki, T. J. (2007). Natural
polymorphism affecting learning and memory in Drosophila. *Proc. Natl. Acad. Sci.* **104**, 13051–
13055.

Mery, F., Varela, S. A. M., Danchin, É., Blanchet, S., Parejo, D., Coolen, I. and Wagner, R. H. (2009).
Public Versus Personal Information for Mate Copying in an Invertebrate. *Curr. Biol.* **19**, 730–734.

Miller, R. and Stephen, W. (1966). Spatial Relationships in Flocks of Sandhill Cranes (Grus
Canadensis). *Ecology* **47**, 323–327.

Navarro, J. and del Solar, E. (1975). Pattern of spatial distribution in Drosophila melanogaster.
Behav. Genet. **5**, 9–16.

Osborne, K. A., Robichon, A., Burgess, E., Butland, S., Shaw, R. A., Coulthard, A., Pereira, H. S.,
Greenspan, R. J. and Sokolowski, M. B. (1997). Natural Behavior Polymorphism Due to a cGMP-
Dependent Protein Kinase of Drosophila. *Science (80-.).* **277**, 834–836.

Owald, D. and Waddell, S. (2015). Olfactory learning skews mushroom body output pathways to
steer behavioral choice in Drosophila. *Curr. Opin. Neurobiol.* **35**, 178–184.

R Core Team (2018). R: A Language and Environment for Statistical Computing.

Ramdya, P., Lichocki, P., Cruchet, S., Frisch, L., Tse, W., Floreano, D. and Benton, R. (2014).
Mechanosensory interactions drive collective behaviour in Drosophila. *Nature* **519**, 233–236.

Sarin, S. and Dukas, R. (2009). Social learning about egg-laying substrates in fruitflies. *Proceedings.*

Biol. Sci. **276**, 4323–8.

Schneider, J., Dickinson, M. H. and Levine, J. D. (2012). Social structures depend on innate
determinants and chemosensory processing in Drosophila. *Proc. Natl. Acad. Sci.* **109**, 17174–
17179.

Schwaerzel, M., Heisenberg, M. and Zars, T. (2002). Extinction antagonizes olfactory memory at the
subcellular level. *Neuron* **35**, 951–960.

Schwaerzel, M., Monastirioti, M., Scholz, H., Friggi-Grelin, F., Birman, S. and Heisenberg, M. (2003).
Dopamine and octopamine differentiate between aversive and appetitive olfactory memories in
Drosophila. *J. Neurosci.* **23**, 10495–502.

Semmelhack, J. L. and Wang, J. W. (2009). Select Drosophila glomeruli mediate innate olfactory
attraction and aversion. *Nature* **459**, 218–223.

Simon, A. F., Chou, M. T., Salazar, E. D., Nicholson, T., Saini, N., Metchev, S. and Krantz, D. E. (2012).
A simple assay to study social behavior in Drosophila: Measurement of social space within a
group. *Genes, Brain Behav.* **11**, 243–252.

**Suh, G. S. B., Wong, A. M., Hergarden, A. C., Wang, J. W., Simon, A. F., Benzer, S., Axel, R. and
Anderson, D. J.** (2004). A single population of olfactory sensory neurons mediates an innate
avoidance behaviour in Drosophila. *Nature* **431**, 854–859.

Tauber, E. and Eberl, D. F. (2003). Acoustic communication in Drosophila. *Behav. Processes* **64**, 197–
210.

Tempel, B. L., Bonini, N., Dawson, D. R. and Quinn, W. G. (1983). Reward learning in normal and
mutant Drosophila. *Proc. Natl. Acad. Sci. U. S. A.* **80**, 1482–6.

Templeton, J. J. and Giraldeau, L.-A. (1996). Vicarious sampling; the use of personal and public
information by starlings.pdf. *Behav Ecol Sociobiol.* **38**, 105–114.

Thum, A. S., Jenett, A., Ito, K., Heisenberg, M. and Tanimoto, H. (2007). Multiple memory traces for
olfactory reward learning in Drosophila. *J. Neurosci.* **27**, 11132–8.

Tinette, S., Zhang, L. and Robichon, A. (2004). Cooperation between Drosophila flies in searching
behavior. *Genes, Brain Behav.* **3**, 39–50.

**Tinevez, J. Y., Perry, N., Schindelin, J., Hoopes, G. M., Reynolds, G. D., Laplantine, E., Bednarek, S.
Y., Shorte, S. L. and Eliceiri, K. W.** (2017). TrackMate: An open and extensible platform for
single-particle tracking. *Methods* **115**, 80–90.

Torney, C., Neufeld, Z. and Couzin, I. D. (2009). Context-dependent interaction leads to emergent
search behavior in social aggregates. *Proc. Natl. Acad. Sci.* **106**, 22055–22060.

Trannoy, S., Chowdhury, B. and Kravitz, E. A. (2015). Handling alters aggression and loser effect
formation in Drosophila melanogaster. *Learn. Mem.* **22**, 64–68.

Valone, T. J. (1989). Group Foraging, Public Information, and Patch Estimation. *Oikos* **56**, 357–363.

Ward, P. and Zahavi, A. (1973). Importance of certain assemblages of birds as information centers
for food-finding. *Ibis (Lond. 1859).* **115**, 517–534.

Wilson, R. I. (2013). Early olfactory processing in Drosophila: mechanisms and principles. *Annu. Rev.
Neurosci.* **36**, 217–41.

Worden, B. D. and Papaj, D. R. (2005). Flower choice copying in bumblebees. *Biol. Lett.* **1**, 504–507.

SUPPLEMENTARY MATERIALS

Table S1: Bayesian probabilities comparing conditioned approach probability between groups.
Red values indicate significant differences.

Time (minute)	Group	Pair >	Small >	Large >
All test (1st-7th)	Single	0.187	0.856	0.926
	Pair		0.968	0.983
	Small			0.677
1st	Single	0.031	0.277	0.195
	Pair		0.906	0.856
	Small			0.393
2nd	Single	0.502	0.840	0.976
	Pair		0.839	0.976
	Small			0.904
3rd	Single	0.009	0.740	0.976
	Pair		0.998	> 0.999
	Small			0.941
4th	Single	0.214	0.713	0.991
	Pair		0.912	0.998
	Small			0.974
5th	Single	0.033	0.727	0.945
	Pair		0.991	> 0.999
	Small			0.857
6th	Single	0.301	0.792	0.993
	Pair		0.906	0.998
	Small			0.968
7th	Single	0.221	0.906	0.992
	Pair		0.980	> 0.999
	Small			0.887

776 **Table S2: Bayesian probabilities comparing conditioned approach probability against chance (0.5).**
777 Red values indicate significant differences.
778

Time (minute)	Group	p(Conditioned approach probability > 0.5)
All test (1st – 7th)	Single	> 0.999
	Pair	> 0.999
	Small	0.983
	Large	> 0.999
1st	Single	0.997
	Pair	0.640
	Small	0.983
	Large	0.963
2nd	Single	0.998
	Pair	0.997
	Small	> 0.999
	Large	> 0.999
3rd	Single	0.997
	Pair	0.359
	Small	> 0.999
	Large	> 0.999
4th	Single	0.964
	Pair	0.767
	Small	0.993
	Large	> 0.999
5th	Single	0.993
	Pair	0.499
	Small	> 0.999
	Large	> 0.999
6th	Single	0.861
	Pair	0.64253
	Small	0.983
	Large	> 0.999
7th	Single	0.642
	Pair	0.232
	Small	0.983
	Large	> 0.999

781 **Table S3: Bayesian probabilities comparing distance distributions between experimental (Exp.) groups and**
782 **simulated (Sim.) groups.**
783 Red values indicate significant differences.
784

Group	Distance bin (mm)	*p(Sim. group > Exp. group)*
Pair	0-5	0.220
	5-10	0.260
	10-15	0.231
	15-20	0.141
	20-25	0.367
	25-30	0.978
	30-35	0.977
	35-40	0.742
	40-45	0.625
	45-50	0.244
	50-55	0.220
	55-60	0.555
	60-65	0.844
Small	0-5	0.069
	5-10	0.054
	10-15	0.483
	15-20	0.556
	20-25	0.759
	25-30	0.810
	30-35	0.782
	35-40	0.679
	40-45	0.813
	45-50	0.797
	50-55	0.554
	55-60	0.429
	60-65	0.992
Large	0-5	0.003
	5-10	0.040
	10-15	0.243
	15-20	0.373
	20-25	0.492
	25-30	0.772
	30-35	0.894
	35-40	0.971
	40-45	0.959
	45-50	0.956
	50-55	0.778
	55-60	0.627
	60-65	0.887

788
Figure S1: Latency to approach the CS+ and CS- in the test.
(**A**) Latency to the CS+: the first time point an individual fly reaches the CS+ for the single fly (red), pair (blue),
small group (yellow) and the large group (pink) during the test. Points indicate the mean latency per group size;
whiskers indicate the 95 % credible intervals. Stars represent significance between groups (*: *p(small group >*
pair) > 0.95).
(**B**) Same as (A) but for the CS-.
795

796
797
Figure S2: Quantifying inter-fly encounters during the test.
(**A**) The number of inter-fly distances during the test for flies conditioned in pairs (left), small groups (middle)
and large groups (right. The first bin ranges from 0 mm to 2.5 mm, and the following bins represent a range of
5 mm i.e. the 10 mm bin ranges from 7.5 mm to 12.5 mm. Bars represent the mean number of distances per
bin between real groups of flies (pair: blue, small group: yellow, large group: pink) and between simulated
groups of flies (grey). Vertical lines represent the 95 % credible intervals. Videos were recorded at 15 frames /
s. Bars of the experimental and simulated number of distances within the same bin were compared to each
other. Stars indicate significantly different numbers of distances between the experimental and simulated
distances within a bin (N = 30 real or simulated experimental runs).
(**B**) Mean encounter length per experimental run for the pair (blue, left), small group (yellow, middle) and large
group (pink, right). The corresponding simulated groups of flies are shown in grey. Bars represent the mean
across experimental runs. Whiskers represent the 95 % credible intervals. Stars represent significant
differences between the real and simulated groups.