Survival status and predictors of mortality among Breast Cancer patients in Adult Oncology Unit at Black Lion Specialized Hospital, Addis Ababa, Ethiopia,2018.

Habtamu Abera Areri [2]*, Wondimeneh Shibabaw[1]**,** Tefera Mulugeta[2],Yared Asmare[1],Tadesse Yirga[3]

[1]Institute of medicine and health science, Debre Berhan University, Debre Berhan, Ethiopia

[2]College of Health Science, Addis Ababa University, Addis Ababa, Ethiopia.

[3]College of Health Science, Debre Markos University, Debre Markos, Ethiopia.

Address

* Corresponding author: *habtamu.abera64@gmail.com*

 P. O. Box 4412, Addis Ababa University, Ethiopia

Co-authors emails.

WS: *wshibabaw2015@gmail.com*

TM: tmulugeta79@yahoo.com

TY:tadesseyirga680@gmail.com

YA:yaredasmare123@gmail.com

Abstract

Introduction: Breast cancer is a foremost cause of death worldwide, ranks fifth among causes of death from all types of cancers; this is the most common cause of cancer death in women among both developing and developed countries. Breast cancer ranks first among most frequent cancers in women of Ethiopia. In spite of the high incidence, mortality rate, and survival status among breast cancer patients was not determined in Ethiopia.

Objective: The main aim of the study is to assess the survival status and predictor the mortality among Breast Cancer patients in Adult Oncology Unit at Black Lion Specialized Hospital in 2018.

Methods: An institution based retrospective follow up study was conducted in Adult Oncology Unit at Black Lion Specialized Hospital. All cases of breast cancer registered from 1st January 2012 to 31th December,2014 were followed for the six-year survival (until 31th December, 2017). Kaplan-Meier survival curve together with log rank test was deployed to test for variations in the survival among predictor variables. Cox regression was used at 5% level of significance to determine the net effect of each independent variable on time to death of breast cancer clients.

Results: The results indicate that the incidence rate of mortality was 9.8 per 100 person/ years (95% CI: 8.49- 11.47).The overall median survival time was 56.5(95% CI (53.46 - 60.83)) months. The overall estimated survival rate was recorded 27% (95% CI, 17.09 to 36.67 %) at 72 months of follow up, whereas at odd years (1, 3, and 5 years) were, 97.2%, 80.8%, and 46.2% respectively. Predictors of mortality were assessed at clinical stage (III&IV),(AHR =1.86), poorly differentiated histology (AHR: 3.1) & positive lymph node status (AHR:3.13),Whereas adjuvant hormone therapy (AHR: 0.67) and chemotherapy (AHR:0.72) were protective.

Conclusion: The overall probability of survival in Ethiopia was inferior when compared with other high and middle-income countries. Predictors of mortality were at advanced clinical stage, poorly differentiated histology grade, surgical margin involvement and positive lymph node status. In contrary, adjuvant hormone therapy, modified radical mastectomy and chemotherapy were protective factors. Hence, special emphasis could be given to early screening, stage diagnosis and initiation of treatment.

Keywords:-Breast cancer, mortality, predictors**,** survival, and Ethiopia.

Introduction

Breast cancer is an emerging public health danger in the worldwide, responsible for the fifth cause of death among all forms of cancers. Globally breast cancer is the second most common cancer next to lung cancer (1). Similarly, which accounts for 25% of cancer cases and 15% of cancer deaths among women (2). According to global burden of cancer 2018 estimation, globally there will be about 2.1 million new female breast cancer cases (3).

Based on 2019 estimation in U.S, an estimated 268,600 new cases of invasive breast cancer are anticipate to be diagnosed in women and about 41,760 women are likely to die(4). Globally, each year above 1.1 million women were newly diagnosed and responsible for 1.6% of female deaths from cancer causes (5). In low-income countries, breast cancer incidence was comparably low, but the most common leading cause of death among women, whereas now a day in high income countries, mortality became reduced (6). The mortality rates of breast cancer developed regions was less than the incidence because of early stage diagnosis and advanced treatment (7). Even though breast cancer has higher incidence in developed countries. But more than half of new breast cancer diagnosis and about 60% of breast cancer mortality happen in the developing countries(8).

Currently breast cancer is a rising public health problem for Sub-Saharan Africa (9). Likewise, in Ethiopia breast cancer became the public health disaster and one study indicate that breast cancer responsible for about 20.8 % of all cancers (10). In addition, Addis Ababa city cancer registry reports, shows that breast cancer accounts for about 34% of all female cancer cases (11). Worldwide, the cumulative 5-year survival rates of breast cancer were varying greatly, which is more than 85% in the high-income countries, while it is 60% or lower in many LMICs (12). Overall, in high-income countries, because of screening program, and early stage diagnosis, the prognosis is good. However, in low and middle-income countries, women diagnosis at advanced stage with more aggressive histological features, and related with a poorer survival (13,14).

In Sub-Sahara Africa, the 5-year survival rate of women who had breast cancer revealed that below 50%, in contrasting with 73% black women in US (9). According to GLOBOCAN data in 2012, the mortality/incidence ratios was 0.55 in Central Africa and 0.16 in the U.S (15). As a result, now a day mortality rates became escalating in certain LMICs, while it decline in most high-income countries (16). Furthermore, clients with late-stage diagnosis,

limited access to standard treatment and those who present with triple-negative subtype tumors had lower survival rates (17).

In Ethiopia according to World Health Organization estimation, an age standardize incidence case of breast cancer was 12,956 and mortality rate was 25 per 100,000 women (18). As a result, to avert this burden currently the Ethiopian Federal Ministry of Health (EFMOH) prepared a task force to address the issue of non-communicable diseases with special emphasis on cancer. Overall, the fundamentals parts of the strategic framework is to reduce the incidence, mortality and improve the quality of life of cancer patients (19). Regardless of the adequate data on the incidence and survival rates of breast cancer in the western countries, survival data were scanty in Africa, and Asia countries (20-22).

In Ethiopia, most of previous studies were focused on metastasis free survival (23), trends of breast cancer (24), treatment outcome (25) and distribution of molecular subtypes of breast cancer clients (26). Despite, the government concern, to decrease the incidence and mortality rate of breast cancer patients, the survival status were still not known in Ethiopia particularly at BLSH. In line with this strategy, the current study aimed to assess the survival status and predictor of mortality among breast cancer patients at Black Lion specialized hospital, Addis Ababa, Ethiopia. Findings of this study could help to provide evidence based information to health care authorities, other stakeholders in order to design interventions that can decrease the incidence and mortality of breast cancer clients.

Methods

Study design, setting and population

A six-year institution based, retrospective follow up study was conducted at BLSH. Patients who have been newly diagnosed and enrolled in breast cancer treatment, from 1st January, 2012 to 31st December 2014 were followed until the end of the study (31st December 2017). Black Lion Specialized Hospital, is located in Addis Ababa, the capital of Ethiopia. It is a teaching; central tertiary comprehensive referral hospital and has 800 beds, renders diagnostic and treatment service for about 370,000 - 400,000 patients per year. The BLSH is the only specialized hospital in the radiotherapy treatment of cancer in Ethiopia.

It has a number of services specialized in the treatment of cancers such as radiotherapy, medical oncology, anatomic pathology, nuclear medicine, gynaecology and surgery. In BLSH oncology unit, there are three senior oncologists, one palliative care specialist, nine residents, five radiotherapist, four medical Physicist, and twenty-three nurses. The most common cancer cases seen in this hospital were breast, cervical, colon and sarcomas. This study was conducted at the adult oncology unit, one of the specialty units of the hospital (27).

Inclusion and exclusion criteria

All adult breast cancer patients who were newly diagnosed and enrolled at BLSH during the study time (*i.e.* 1st January 2012 to 31st December 2014) were included. On the other hand, women whose medical charts were incomplete, not found, those had previous diagnosis of breast cancer, a patient who had diagnosis at other hospitals and subsequently referred to BLSH for further treatment were excluded.

Sample size and sampling technique

All breast cancer patients who attended the oncology unit of BLSH between 1st January 2012 to 31th December,2014 and fulfilled the inclusion criteria of the study were included in the study. This period was selected in order to have the nearest six year follow up period. A total of 1,083 breast cancer patients were registered in this period. Out of them, 627 participants who fulfilled the inclusion criteria were included in the study (**see figure1**).

Figure 1:-Flow diagram showing the final sample size included in the current study at black lion specialized hospital, Addis Ababa, Ethiopia, 2018.

Data collection tool and procedure

Data collection tool were developed from different literature (5,11,16). All available information on patient records were checked and formats from different literatures were reviewed with modification. Then, appropriate data extraction format were developed in English, to extract all the relevant variables from patient charts to meet the study objectives. Training on record review was given to data collectors and supervisors for 01 day before actual data collection task on the already existing records half-day theoretical and half-day practical training. The starting point for follow-up was the time between 1st January 2012 to 31th December 2014 with first confirmed diagnosis of breast cancer and the endpoint was date of death, date of lost to follow up, date of last contact until 31th December 2017, whichever comes first.

All sociodemographic, clinical, pathological and treatment related data were obtained from medical charts of all breast cancer cases diagnosed from 1st January 2012 to 31th December 2014 at black lion specialized hospital were reviewed from cancer registries. The records of all study participants were selected according to the eligibility criteria. The survival status of patients was obtained from the medical record. The time of survival was calculated as the time between the date of diagnosis of breast cancer to the date of death, or the end of study.

Before collecting the data, the records were reviewed (both baseline and follow up records), death certificate complemented by registration was identified from their medical record number. The data were collected by three oncology nurses who were working at the cancer treatment center. Furthermore, the data were extracted and reviewed from the charts.

Definition and measurement of variables

Age at diagnosis was classified into four groups (<40 years, 40-49 years, 50-59 years, >60+ years) based on the American Cancer Society facts and figures for breast cancer. Comorbidity conditions was taken from the Charlson index (28, 29). The presence of any of these conditions at diagnosis was designated as "yes", while the absence of these condition at diagnosis was denoted as "no". Menopausal status (Premenopausal/Postmenopausal), Marriage status (Single, Married, Widowed, Divorced), and place of residence (Urban/Rural) were also recorded.

Clinical stage at diagnosis was assigned to each patient by using the TNM classification scheme (30). In this research, the coding for those diagnosed at stages I and IV remained. Stages IIA and IIB were merged as stage II and similarly stages IIIA, IIIB and IIIC were combined as stage III. The Nottingham Grading System as grade 1, grade 2 and grade 3 measured histological grade. Tumor size was classified based on AJCC guidelines (≤2 cm, >2–5 cm, and >5cm), margins (clear/involved), axillary node status (positive/negative), Hormonal therapy (yes/no), histology type (ductal in situ, ductal invasive, lobular in situ, and lobular Invasive) were also recorded. The type of treatment was classified as chemotherapy with surgery, chemotherapy with radiotherapy, surgery alone and surgery with radiotherapy. The time (measured in months) till the death was used for the survival analysis.

Outcome variable

The primary outcome variable was time to death of breast cancer clients in months. Breast cancer patient's data were followed until the date of death, failure to follow-up, transferring out, or at the end of the study. Individuals who were failure to follow-up, alive and had transferred out at the end of the study period were considered as censored. The time of survival was calculated in months using the time between the dates of diagnosis to the date of the death or date of censoring.

Data entry and analysis

Data were coded, cleaned, edited and then entered using Epi-data VS 3.1 and exported to Stata VS 14 statistical software for further analysis. Data exploration were carry out to see if there are odd codes or items that were not logical and then subsequent editing was made. Descriptive statistics were carried out in terms of central tendency for continuous data and frequency distribution for categorical data. In the current study incidence, density rate of mortality among breast cancer clients was calculated. Subsequently, the number of mortality within the follow up was divided by the total person time at risk on follow up and reported per 100 P/Y. Time to death of breast cancer clients were estimated using Kaplan-Meier survival curve and the presence of variations in survival among categories of co-variates were deployed using log rank test. Before running the cox proportional hazard regression analysis, assumption of proportional-hazard and multi-co linearity were checked.

Those independent variables, at bivariate cox regression having p-value of ≤ 0.25 was used as a cut of point to transfer in the multivariate cox regression analysis. Cox regression at 5% level of significance was used to determine the association between each independent variable to the outcome variable. The assumption of cox-proportional hazard model was checked using schoenfeld residual test and variables having P-value > 0.1 were considered as fulfilling the assumption. Hence, all the variables were fulfilling the assumption. Furthermore, residuals were checked using goodness-of-fit test with cox snell residuals, and which satisfied the model test.

Results

Socio-demographic characteristics

Between 1st January 2012 to 31st December 2014, 1,083 breast cancer patients were enrolled to Black Lion Specialized Hospital, of them, 627 were eligible for this study. Among the study participants, 458 were censored and the rest 169 develop the event. The mean age of study participants at the time of breast cancer diagnosis was 42.61 years with SD $\pm$ 12.28 years. Almost nearly half, 279 (44.5%) of the age group was less than 40 years old. About two-thirds, 403 (64.3%) of patients were married; most, 433 (69.1%) of the women was premenopausal (age less than 50 years old); slightly more than one-third, 224 (35.7%) were having preexisting medical problem during diagnosis. More than half, 366 (58.4%) of the participants were urbanites are shown below (**Table 1**).

Table 1:-Socio-demographic characteristics of breast cancer patients at Black Lion Specialized Hospital, Addis Ababa, Ethiopia, from 1st January 2012 to 31st December 2017 (n=627).

Covariate	category	status at last contact		Total No. (%)
		Censored No. (%)	Death No. (%)	
Age in years at time of diagnosis	<40	212(75.9)	67(24.1)	279(44.49)
	40-49	111(68.09)	52(31.91)	163(26.0)
	50-59	93(78.81)	25(21.19)	118(18.82)
	>60+	42(62.28)	25(37.31)	67(10.68)
Place of residence	Urban	285(77.86)	81(22.14)	366(58.37)
	Rural	173(66.28)	88(33.71)	261(41.63)
Marital status	Single	71(76.34)	22(23.66)	93(4.83)
	Married	301(74.68)	102(25.32)	403(64.27)
	Divorced	63(63.64)	36(36.36)	99(15.78)
	Widowed	23(71.87)	9(28.13)	32(5.1)
Menopause status	Premenopausal	316(72.97)	117(27.03)	433(69.05)
	Postmenopausal	142(73.19)	52(26.81)	194(30.95)
Preexisting medical diagnosis	No	323(80.14)	80(19.86)	403(64.27)
	Yes	135(60.26)	89(39.74)	224(35.73)

Clinical, Histopathological and treatment characteristics

More than half, 357 (57%) of the women were clinical stage III at the time of diagnosis. Invasive Ductal Carcinoma (IDC) was predominant, 427 (68.1%) histology type of cancer. Nearly half, 304 (48.5%) of the histology grade were moderately differentiated; Surgery associated with radiotherapy was the common mode of treatment which accounts for 256 (40.8%). More than half, 369 (58.85%) of tumor size was less than 2.5 cm on presentation. About two third, 407 (64.91 %) of the study participants were reported receiving hormone therapy. Nearly half, 229 (48.11%) of patients had modified radical mastectomy. The clinical, histopathological and treatment characteristics of the study participants are shown below (**Table 2**).

Table 2:-Baseline clinical, histological and treatment information of breast cancer patients at Black Lion Hospital, Addis Ababa, Ethiopia, from 1st January 2012 to 31st December 2017 (n=627)

Variable	Category	Vital status at last contact		Total
		Censored No. (%)	Death No. (%)	No. (%)
Stage of breast cancer	I	21(95.45)	1	22(3.5)
	II	145(86.82)	22(13.18)	167(26.63)
	III	245(68.62)	112(31.38)	357(56.93)
	IV	47(58.02)	34(41.98)	81(12.91)
Histology grade	Grade I	84(91.3)	8	92(14.67)
	Grade II	237(77.96)	67(22.04)	304(48.48)
	Grade III	137(59.3)	94(40.7)	231(36.84)
Histology type	Ductal insitu	136(83.44)	27(16.56)	163(26.3)
	Invasive ductal	294(68.85)	133(31.15)	427(68.1)
	Lobular insitu	15(75)	5	20(3.2)
	Invasive lobular	11(73.4)	4	15(2.4)
Deep surgical margin	Free	393(80.2)	97(19.8)	490(79.28)
	involved	58(45.3)	70(54.7)	128(20.72)
Node status	Negative	364(80)	91(20)	455(72.68)
	Positive	93(54.38)	78(45.62)	171(27.32)
Metastasis at time diagnosis	No	348(87.21)	51(12.79)	399(63.64)
	Yes	110(48.24)	118(51.76)	228(36.36)
Tumor size	≤2.5cm	289 (81.17)	67(18.82)	356(56.78)
	2.5-5cm	152(63.34)	88(36.66)	240(38.27)
	>5cm	17(54.8)	14(45.2)	31 (4.95)
Number of Positive node	<2	284(75.53)	92(24.47)	376(59.97)
	≥2	174(69.32)	77(30.68)	251(40.03)
Treatment mode	Surgery associated with radiotherapy [a]	165(64.45)	66(25.65)	256(40.82)
	Surgery[b]	101(72.66)	38(27.34)	139(22.18)
	others[c]	191(82.68)	65(28.13)	231 (36.8)

Type of surgery	Breast-conserving surgery	65 (58.56)	46(41.44)	111(23.32)
	total mastectomy	86 (76.1)	27 (23.89)	113(23.74)
	Modified radical mast	180 (78.6)	49 (21.4)	229(48.11)
	Axillary node dissection	14(60.86)	9(39.13)	23(4.83)
Chemotherapy	No	67(72.8)	25(27.2)	92 (14.7)
	Yes	391 (73.1)	144(26.9)	535(85.3)
Endocrine therapy	No	137(62.27)	83(37.72)	220(35.08)
	Yes	321(78.86)	86(21.13)	407(64.92)

[a]Surgery associated with radiotherapy (or also combined with chemotherapy)

[b]Surgery only (or also combined with chemotherapy) ;[c] radiotherapy with chemotherapy(surgery therapy).

Survival status of breast cancer patients

The overall mortality rate in the cohort during the 1,712 person-years of observation (PYO) was 9.87 per 100 (95% CI: 8.49-11.47) person-years follow up. The cumulative incidence of this study was 169 (27%) with the confidence interval (95% CI, 23.6-30.3%) of patients were died over six years. However, 458 (73.04%) were censored till the end of the study. Of these, 194 (42.27%) were failure to follow up, 132 (28.76%) were alive, 119 (25.9%) was against medical advice and the rest were transferred to other institution during the overall study follow up period.

Survival probability of breast cancer patients

In the current study, 627 breast cancer participants were followed for a total of 72 months. The estimated median survival time was 56.5 (95% CI, 53.46 - 60.83) months. Kaplan-Meier survival estimation showed that overall estimated survival rate after diagnosis of breast cancer was 26.42% (95% CI, 17.09 to 36.67 %) at 72 months of follow up. Likewise, the estimated cumulative survival rate was 97.2%, 89.8%, 80.8%, 66.33%, 46.2% and 26.4% at 12, 24, 36, 48, 60 and 72 months respectively **(see figure 2)**.

Figure 2. Kaplan-Meir survival estimate on the overall survival time of breast cancer clients at Black Lion Specialized Hospital, Addis Ababa, Ethiopia, 2018 (n = 627).

Survival function among different predictors

Log-rank test was carried out to check the presence of any significant differences in survival time among various levels of the categorical variables. According to the present study it was found that the median survival time for those who had clinical stage I, II or III at baseline had a longer survival time than those who had advanced clinical stage (IV) (45.6 months, 95% :41.04- 50.57) at baseline with a p-value < 0.000. The median survival time for those who have negative lymph node status had a longer survival time 63.6 months than those who had positive lymph node at baseline (47.34 months, 95% CI: 41.2-49.7),which is supported by the p-value of 0.000.

Among cases diagnosed in early stage (I & II), cumulative survival was 56.65 % (95 % CI: 35.94-72.94 %), while those cases detected at late stage (III&IV), had a survival rate of 18.49% (95 % CI: 10.03-28.98%) with the testing equality among the groups with p value of 0.000. A 6-year survival rate for those diagnosed with IDC (invasive ductal carcinoma) was 20.2% (95%CI,10.82-30%) which is lower as compared to other histological types 43.26%(22.39-62.56) shown below (**table 3 and figure 3**).

Table 3:-Survival time, cumulative survival probability, significance and log rank test for the study population according to different characteristics of patients during 6-year of follow-up (Kaplan-Meier Method) of breast cancer clients at Black Lion Specialized Hospital, Ethiopia from 1st January 2012 to 31st December 2017 (n=627).

Covariate	Survival time, mo, (95% CI)	Overall 6-year survival probability (%)	Log rank test (p–value)
	Median(95%CI)		
Residence			
Urban	59.7 (54.47-..)	36.57	(8.83)*
Rural	50.73 (49.3 -60.67)	17.26	
Menopause status			
Premenopausal	60.7 (53.47- 63.63)	28.56	
Postmenopausal	54.5 (48.64-58.84)	19.67	0.63
comorbidity			
No	63.64 (60.84- .)	35.62	(23.18)*
Yes	50.04 (45.6-53.37)	12.89	
Stage			
I		92.86	
II	61.54 (59.7-..)	43.56	(41.46)*
III	54.46 (50.24-60.67)	20.18	
IV	45.6 (41.04- 50.57)	16.8	
histology grade			
Grade I		66.4	
Grade II	60.74 (55.76- 67.54)	34.6	(62.38)*
Grade III	48.63 (42.96 -50.73)	0	
Deep surgical margin			
free			
involved	62.3 (59.7- .)	41.31	(56.32)*
	42.97 (39.74-48.34)	2.93	
Number of involved lymph nodes			
<2	60.74 (54.47-66.5)	30.21	
>=2	53.96 (49.6-59.7)	18.93	3.65
Tumor size			
≤2.5	61.53 (60.67- …)	45.65	(26.42)*
2-5	50.57 (47.96-54.33)	11.35	
>5	45.36 (41.84-…)	0	
Metastasis at diagnosis			
No	66.5 (60.84- .)	48.5	(99.9)*
Yes	42.5 (40.04- 47.34)	5.4	
Endocrine therapy			
No	47.97 (41.84-51.64)	6.8	(40.65)*
Yes	65.3 (58.84- .)	40.58	
Lymph node status			
Negative	63.64 (60.67- .)	38.27	
Positive	47.34 (41.2-49.7)	5.1	(48.95)*

Neo adjuvant Chemotherapy			
No	53.47 (42.5-	24.9	2.35
Yes	58.84 (53.97-61.54	27.52	
Type of surgery Breast conserving surgery	49.6 (43.03-54.34)	12.6	(19.3)*
Total mastectomy	60.74 (51.64- .)	39.2	
MRM	62.3 (58.84- .)	39.4	
Axillary dissection	50.03 (14.64- .)	0	

mo; months, CI; confidence interval, SE: Standard Error, MRM; Modified Radical Mastectomy,

∗ indicates that the variables have significantly difference in survival among groups at 95% level of significant (*P*< 0.05).

A   B

Figure 3. Kaplan-Meier Survival function among different groups of breast cancer clients (A), stage at diagnosis (B), histologic grade at diagnosis at Black Lion Specialized Hospital, Addis Ababa, Ethiopia, 2018 (n=627).

Testing overall model fit

In order to check the overall model fit we compare the jagged line with the reference line, and we observe that, the cox model fit these data to reasonable extent. In addition, if the cox regression model fits the data, these residuals should have a standard censored exponential distribution with hazard ratio. Overall, the hazard function follows the 45° line very closely.

Hence, the output shows cox–snell residuals were satisfied the overall model fitness test (**see figure 4**).

Figure 4 .This figure shows cox-snell residual Nelson -Aalen cumulative hazard graph on breast cancer clients at Black Lion Specialized Hospital, Addis Ababa, Ethiopia, 2018.

Predictors of mortality

In multivariable cox regression analysis explanatory variables with p-value < 0.25 at the bivariate analysis and non-collinear independent variables were included. The result of multivariable analysis revealed that women with late stage of breast cancer (III and IV) were 1.86 times higher hazard of death than those who had early stage breast cancer at the time of diagnosis (I and II) (AHR : 1.86, 95%CI: 1.13- 3.08). Those women whose surgical margin were involved at baseline 3.13 times more likely to die than those women whose surgical margin was not involved (AHR: 3.13, 95% CI: 2.140- 4.573).

Patients having positive lymph node status were 1.83 times higher hazard of death as compared to those having negative lymph node at time of diagnosis (AHR: 1.83, 95% CI:1.217- 2.736). Women those who are being histologic grade III at the beginning of breast cancer diagnosis were 3.12 times higher risk of death as compared to those who have grade I histology(AHR: 3.12, 95% CI:1.16-8.39). Furthermore, women who have taken endocrine therapy during the six year follow up time have reduced mortality by 33% compared to those who are not taken endocrine therapy (AHR: 0.67, 95% CI: 0.451- 0.989).

Table 4:-Results shows cox proportional hazard regression analysis to identify predictors of mortality among breast cancer patients at Black Lion Specialized Hospital, Addis Ababa, Ethiopia, 1st January 2012 to 31st December 2017 (n=627).

Characteristics	Bivariable CHR (95% CI)	Multivariable AHR (95% CI)
Age		
<40	1	1
40-49	1.45(1.010-2.089)*	1.21(.766- 1.898)
50-59	1.21(0.761-1.919)	0.88(0.468- 1.660)
>60+	1.84(1.080-3.141)*	1.62(0.766- 3.447)
Place of residence		
Urban	1	1
Rural	1.57(1.164-2.130)*	1.48(0.999-2.195)
Marital status		
Married	1	1
Single	1.07(.708- 1.629)	0.95(0.551- 1.638)
Widowed	1.77(1.224- 2.568)*	0.69(0.395- 1.224)
Divorced	1.34(0.826- 2.363)	0.67(0.318 -1.422)
comorbidity		
no	1	1
yes	2.07(1.530- 2.808)*	1.49(0.980- 2.290)
Stage of breast cancer		
Early (I&II)	1	1
Advanced (III&IV)	3.11(2.01-4.84)*	1.86(1.127-3.080)**
Histology grade		
Grade I	1	1
Grade II	2.92(1.40-6.09)*	1.73(0.67-4.53)
Grade III	6.79(3.28-14.06)*	3.12(1.16-8.36)**
Histology type		
Invasive Ductal	1.87(1.298-2.714)*	1.04(0.644- 1.697)
Others	1	1
Surgical margin involved		
No	1	1
Yes	3.06(2.252-4.168)*	3.13(2.140- 4.573)**
No of involved lymph node		
<2	1	1
>=2	1.34(0.991-1.822)	0 .79(0.517- 1.205)
Lymph node status		
Negative	1	1
Positive	1.61(1.396-1.848)*	1.83(1.217- 2.736)**
Tumor size		
≤2.5	1	1
2-5	2.19(1.593-3.021)*	1.15(0.736- 1.832)
>5	2.08(1.172-3.717)*	2.31(0.891-4.123)

Type of surgery		
Breast conserving surgery	1	1
Simple (total) mastectomy	0.61(0.378- 0 .986)	0 .68(0.413- 1.134)
MRM	0.48(0.318-0.714)*	0.45(0.280- 0 .690)**
Axillary node dissection	1.51(0.735-3.084)	1.25(0.593- 2.660)
Radiotherapy		
No	1	1
Yes	0.79(0.586-1.074)	0.95(0.654- 1.381)
Neo adjuvant Chemotherapy		
No	1	1
Yes	0.72(0.467-1.099)	0.27(0.130-0.565)**
Hormone therapy		
No	1	1
Yes	0.38(0.281-0.519)*	0.67(0.451-0.989)**

CI; Confidence Interval, aHR; Adjusted Hazard ratio, cHR: Crude Hazard Ratio, MRM; Modified Radical Mastectomy,*shows that significantly associated variables with the outcome at bivariable analysis ($P< 0.05$). **shows that significantly associated variables with the outcome at multivariable analysis ($P< 0.05$).

Discussion

In the current study, the overall mortality rate of breast cancer patients were 9.8/100 women-years. Our finding was much higher than the study done in Brazil (31), where age-standardized mortality rate within 6-year follow up period was 431.8/100 000 women-years. The observed variation in mortality rates might be because of the fact that the more advanced stage (III & IV) breast cancer clients were presented in our finding. Other possible explanation could be methodology difference; they have been used ecological analysis.

In this 6-year retrospective follow up study, the cumulative survival probability at 1, 3, and 5 years were 97.2%, 80.8% and 46.2% respectively. This finding is in line with studies conducted in Malaysia (70.8%, 56.9% and 49.4%) (32) and Ghana, (47.9%) (21). However, this finding is higher than in Tanzania (21.8%) (33), Cameroon (30%)(34), Uganda (51.8 %) (35) and Vietnam, (94%, 83% and 74%)(5). On the contrary, survival probability is still lower when compared with studies done in Northwest Iran (96%, 86%, and 81%), Germany (83%) and the Qidong (83.61%, 67.53% and 58.75%) respectively (36-38). Similarly, the overall cumulative 6 years estimated survival rate of the current study was 26.42 % (95 % CI: 17.09-36.67 %) at 72 months of follow up. This result is inferior than the study done in Qidong (56.04%) (38).

This gap might be due to several reasons. This perhaps could be due to advanced stage at diagnosis (69.5%). However, 52.3% in Ghana, 27.6% in Vietnam, 67.8% in Cameroon is due

to lack of early screening programs, limited treatment facilities, and financial problems. Other explanation could be the facilities to treat cancer are located in the capital city of the country. This could result in delay of diagnosis and treatment. Additionally, possible explanation on discrepancies in survival might be due to different methodologies applied in each studies, as well as variances in local cancer care. Furthermore, the molecular and genetic differences among breast cancer clients may also contribute to variations in survival probability.

In the current study, the overall median survival for histologically confirmed breast cancer was 56.5 (95% CI, 53.46 - 60.83) months. The results were higher than the previous studies in which the median survival was 24 months in Cameroon (34), 40 months in Sudan (39), but lower than the results reported from Malaysia the median survival was 68.1 months (32). Although our study showed higher overall median survival than that of the Cameroon and Sudan studies, the lower value obtained in comparison with Malaysia is most probably due to the shorter period of study. In addition, it could be difference in health seeking behavior and treatment adherence among those studies.

In the log rank test, it is observed that the median survival time for those who had clinical stage I, II or III at baseline had a longer survival time than those in advance clinical stage (IV) (45.6 months, 95% :41.04- 50.57). Similarly, the median survival time for those who have negative lymph node status had a longer survival time (63.6month, 95%CI: 60.67-) than those who had positive lymph node at baseline (47.34 months, 95% CI: 41.2-49.7). These values are in line with the studies conducted in different African countries (39,40).

In the present study, women who had positive lymph node at diagnosis were nearly 2 times prone for death than women who had negative lymph node at baseline. These results were in agreement with different studies conducted in African countries (21,41) and the study done in Iran (42) in which lymph node status was an important determinant of survival. In addition, as the number of lymph nodes increases, also does the relapse rate, while the survival rate decreases. Women who had histologic grade III at diagnosis were nearly 3 times more likely to die as compared to women who had well differentiated histologic (grade I) at baseline. These results were in agreement with the studies conducted in different Asian countries (32, 42, 43).

According to the results obtained, women who had positive surgical margin were 3.13 times more prone to death than women who were free from surgical margin at time of diagnosis.

Most of the previous studies go hand in hand with the present findings (5,35). These findings might be explained by the fact that the residual disease at the surgical margins could increase the risk of local recurrence and possibly death through the years.

As shown by others studies, women who had advanced stage of breast cancer (III and IV) at diagnosis were nearly 2 times more vulnerable to death than women who had early stage breast cancer (I&II) at time of diagnosis. The same line of results were presented by several studies conducted in different African and Asian countries (5,35, 37,43). This reflect that stage was a major predictor of mortality among breast cancer clients and had the most significant influence on patient outcome in the present study.

The results indicate that patients who had undergone modified radical mastectomy (MRM) experienced 55% reduced chances of mortality than those who underwent breast-conserving surgery. The observed results are in contrary with the studies done in different African and Asian countries (34,36,42). The difference in survival between two groups could be because of late stages at presentation and poor infrastructure for treatment of breast cancer has made this approach of surgical treatment more common in many developing countries including Ethiopia. Scientifically, breast-conserving surgery is usually recommended for early stage breast cancer in contrast to modified radical mastectomy. Other possible explanation could be the options for treating breast cancer patient, which depends on the stage of disease.

Unlike most of previous findings (40,44), those women who were receiving neo-adjuvant chemotherapy was found to be reduce the chances of death by 73% than those patients without neo-adjuvant chemotherapy. However, the current result is in agreement with study done in France (45), which revealed a 25% reduction in the relative risk of death. Overall, the present study found a protective effect of neo-adjuvant chemotherapy on survival of breast cancer patients. This might be due to patients with advanced stage breast cancer, neo-adjuvant chemotherapy would be applied in order to shrink the tumor size and facilitate surgery. In our study, the better survival rate was related to patients receiving neo-adjuvant chemotherapy. Additional studies should be conducted to confirm a protective effect of neo-adjuvant chemotherapy on survival rate of breast cancer clients.

The final predictor of survival identified in the study was hormone therapy, which reduced the chances of death by 33% than those who could not undergo hormone therapy. The findings are in consistent with previous findings that were conducted in different continents (5, 40, 44) which reflect that the effect of hormone therapy on survival rate was often stated along with the impact of hormone receptors.

This study has some limitations too. First, cause specific survival was not determined due to lack of data on specific cause of death. This may over estimate breast cancer related mortality rate. Second, limited data on hormone receptors; estrogen receptor, progesterone receptor and Human Epidermal Growth Factor Receptor 2 prevented us from analyzing the role of them on survival time. Likewise, data were not recorded on socio economic status, such as occupation and educational level. Selection bias was possibly introduced during secondary data collection because patient's with incomplete charts were excluded. Moreover, the data were collected over the period 1st January 2012 to 31st December 2014 and do not reflect current utilization of advanced treatment methods and new medications for breast cancer treatment.

Conclusion

The overall probability of survival in patient's diagnosis of breast cancer was 27% at 72 months of follow up in Ethiopia that is inferior when compared with those of high and middle-income countries. Significant predictors of mortality after diagnosis of breast cancer were advanced clinical stage, grade III histology, involvement of surgical margin and positive lymph node status. In contrast, hormone therapy, modified radical mastectomy and chemotherapy reduced mortality. Hence, a special emphasis should be given to early screening, diagnosis and initiation of treatment before advanced stage. leading to high mortality.

Abbreviations

AAU=Addis Ababa University, AJCC=American Joint Committee on Cancer, BLSH=Black Lion Specialized Hospital, ER=Estrogen Receptor,EFMOH= Ethiopian Federal Ministry of Health, GLOBOCAN=Global Burden of Cancer, HER2= Human Epidermal Growth Factor Receptor 2,HIC=High Income Country, IDC=Invasive Ductal Carcinoma, LMICs=Low and Middle Income Countries,PR= Progesterone Receptor,SSA=Sub-Saharan Africa.

Ethics approval and consent to participate

Ethical clearance was obtained from Addis Ababa University, ethical review board. Permission for conducting the study was gained from Adult Oncology Unit, Black Lion Specialized Hospital. The data were collected from the client charts, as a result the individual clients were not exposed to any harm as far as the privacy was kept. To keep the confidentiality, all collected data was coded and locked in a separate room before being entered into the computer.

Consent for publication

Not applicable.

Availability of data and materials

Data will be shared upon request and will be obtained by emailing to the corresponding author using "wshibabaw2015@gmail.com".

Competing interest

The authors confirmed that, we have no any competing interest

Funding

The study was funded by Addis Ababa University, Ethiopia. We would like to acknowledge the university. However, the funder has no role on study design, data collection, analysis, interpretation of data, and preparation of manuscript or decision to publication.

Authors' contributions

WS, TM, and HA were involved in conceptualize, design, methodology, statistical analysis, developing the initial drafts of the manuscript and revising. Whereas YA and TY involved in software, supervision and subsequent drafts. In addition, WS, TM, and HA prepared the final draft of the manuscript, reviewed and edited. All authors read and agreed on the final draft of the manuscript.

Acknowledgment

We would like to acknowledge Addis Ababa University for funding and Black Lion Specialized Hospital for permission to undertake this study. Our heartfelt thanks to Black Lion Specialized Hospital Managers, all oncology unit staffs, card room officer and data collectors for their cooperation during data collection.

Reference

1.World Health Organization (2017). Cancer Facts & Figures 2016-2017. Available at: http:// http://www.who.int/mediacentre/factsheets/fs297/en/. Accessed on: 12/31/17.

2.Ferlay J, Soerjomataram I, Ervik M, Dikshit R, Eser S, Mathers C, et al. GLOBOCAN 2012 v1. 0, Cancer Incidence and Mortality *Worldwide*: IARC CancerBase No. 11 [Internet].

2013; Lyon, France: International Agency for Research on Cancer. globocaniarcfr/Default aspx. 2014.

3. Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. CA: a cancer journal for clinicians. 2018 Nov;68(6):394-424.

4.Breast Cancer Statistics.https://www.breastcancer.org/symptoms/understand_bc/statistics [Accessed4 June 2019].

5. Lan N, Laohasiriwong W, Stewart J. Survival probability and prognostic factors for breast cancer patients in Vietnam. Global health action. 2013;6(1):18860.

6. Ferlay J, Soerjomataram I, Dikshit R, Eser S, Mathers C, Rebelo M, et al. Cancer incidence and mortality worldwide: sources, methods and major patterns in GLOBOCAN 2012. International journal of cancer. 2015;136(5).

7. Cancer IAfRo. GLOBOCAN 2012: estimated cancer incidence, mortality and prevalence worldwide in 2012. 2012.

8.Jemal A, Bray F, Center MM, Ferlay J, Ward E, Forman D. Global cancer statistics. CA Cancer J Clin. 2011;61(2):69–90**.

9. McKenzie F, Zietsman A, Galukande M, Anele A, Adisa C, Cubasch H, et al. African Breast Cancer—Disparities in Outcomes (ABC-DO): protocol of a multicountry mobile health prospective study of breast cancer survival in sub-Saharan Africa. BMJ open. 2016;6(8):e011390.

10. Abate S, Yilma Z, Assefa M, Tigeneh W. Trends of Breast Cancer in Ethiopia. *Int J Cancer Res Mol Mech*. 2016;2(1).

11.Top 10 Female Cancer in Addis Ababa [online]. Available: [http://afcrn.org/membership/membership-list/100-addisababa] [Accessed 21 Apr 2017].

12. Allemani C, Weir HK, Carreira H, Harewood R, Spika D, Wang X-S, et al. Global surveillance of cancer survival 1995–2009: analysis of individual data for 25 676 887 patients from 279 population-based registries in 67 countries (CONCORD-2). The Lancet. 2015;385(9972):977-1010.

13. Claudia A, Weir HK, Carreira H, Harewood R, Spika D, Wang X-S, et al. da Silva et al.". Wan-Qing Chen, Olufemi J Ogunbiyi, Bernard Rachet, Matthew J Soeberg, Hui You, Tomohiro Matsuda, Magdalena Bielska-Lasota, Hans Storm, Thomas C Tucker, Michel P Coleman, and the CONCORD Working Group*--Global surveillance of cancer survival. 1995;2009.

14. Danforth Jr DN. Disparities in breast cancer outcomes between Caucasian and African American women: a model for describing the relationship of biological and nonbiological factors. Breast cancer research. 2013;15(3):208.

15. Torre LA, Bray F, Siegel RL, Ferlay J, Lortet-Tieulent J, Jemal A. Global cancer statistics, 2012. CA: a cancer journal for clinicians. 2015 Mar;65(2):87-108.

16. DeSantis CE, Bray F, Ferlay J, Lortet-Tieulent J, Anderson BO, Jemal A. International variation in female breast cancer incidence and mortality rates. Cancer Epidemiology and Prevention Biomarkers. 2015;24(10):1495-506.

17. Xia C, Ma F, Zeng H, Li H, Yang L, Liu L, et al. The influence of stage at diagnosis and molecular subtype on breast cancer patient survival: a hospital-based multi-center study. Chinese Journal of Cancer. 2017;36(1):84.

18. Organization WH. cancer country profiles 2014. 2014.

19. Nuño R, Coleman K, Bengoa R, Sauto R. Integrated care for chronic conditions: the contribution of the ICCC Framework. Health Policy. 2012;105(1):55-64.

20. Kene TS, Odigie VI, Yusufu LM, Yusuf BO, Shehu SM, Kase JT. Pattern of presentation and survival of breast cancer in a teaching hospital in north Western Nigeria. Oman medical journal. 2010;25(2):104.

21. Mensah AC, Yarney J, Nokoe SK, Opoku S, Clegg-Lamptey J. Survival outcomes of breast cancer in Ghana: an analysis of clinicopathological features. Open Access Library J. 2016;3:1-11.

22. Parkin DM, Bray F, Ferlay J, Jemal A. Cancer in africa 2012. Cancer Epidemiology and Prevention Biomarkers. 2014;23(6):953-66.

23. Kantelhardt EJ, Zerche P, Mathewos A, Trocchi P, Addissie A, Aynalem A, Wondemagegnehu T, Ersumo T, Reeler A, Yonas B, Tinsae M. Breast cancer survival in Ethiopia: a cohort study of 1,070 women. International journal of cancer. 2014 Aug 1;135(3):702-9.

24. Abate S, Yilma Z, Assefa M, Tigeneh W. Trends of Breast Cancer in Ethiopia. Int J Cancer Res Mol Mech. 2016;2(1).

25. SelamawitD.etal, treatment outcome and associated factors of breast cancer at black lion specialized hospital, Ethiopia, master thesis, accessed from Addis ababa university library catalog march 2018.

26. Hadgu E, Seifu D, Tigneh W, Bokretsion Y, Bekele A, Abebe M, Sollie T, Merajver SD, Karlsson C, Karlsson MG. Breast cancer in Ethiopia: evidence for geographic difference in the distribution of molecular subtypes in Africa. BMC women's health. 2018 Dec;18(1):40.

27. Black lion specialized hospital oncology unit manager office D, 2017.

28. Maskarinec G, Pagano I, Lurie G, Bantum E, Gotay CC, Issell BF. Factors affecting survival among women with breast cancer in Hawaii. Journal of Women's Health. 2011;20(2):231-7.

29. Kartal M, Tezcan S, Canda T. Diagnosis, treatment characteristics, and survival of women with breast cancer aged 65 and above: a hospital-based retrospective study. BMC women's health. 2013;13(1):34.

30.American Cancer Society. Breast Cancer Facts & Figures 2017-2018. Atlanta: American Cancer Society in 2017.

31. Diniz CSG, Pellini ACG, Ribeiro AG, Tedardi MV, de Miranda MJ, Touso MM, et al. Breast cancer mortality and associated factors in São Paulo State, Brazil: an ecological analysis. BMJ open. 2017;7(8):e016395.

32. Abdullah NA, Wan Mahiyuddin W, Muhammad NA, Ali ZM, Ibrahim L, Ibrahim Tamim N, et al. Survival rate of breast cancer patients in Malaysia: a population-based study. Asian Pac J Cancer Prev. 2013;14(8):4591-4.

33. Mabula JB, Mchembe MD, Chalya PL, Giiti G, Chandika AB, Rambau PF, et al. Stage at diagnosis, clinicopathological and treatment patterns of breast cancer at Bugando Medical Centre in north-western Tanzania. Tanzania journal of health research. 2012;14(4).

34. Ngowa J, Kasia J, Yomi J, Nana A, Ngassam A, Domkam I, et al. Breast Cancer Survival in Cameroon: Analysis of a Cohort of 404 Patients at the Yaoundé General Hospital. Advances in Breast Cancer Research. 2015;4(02):44.

35. Galukande M, Wabinga H, Mirembe F. Breast cancer survival experiences at a tertiary hospital in sub-Saharan Africa: a cohort study. World journal of surgical oncology. 2015;13(1):220.

36. Ziaei JE, Sanaat Z, Asvadi I, Dastgiri S, Pourzand A, Vaez J. Survival analysis of breast cancer patients in northwest Iran. Asian Pacific Journal of Cancer Prevention. 2013;14(1):39-42.

37. Holleczek B, Jansen L, Brenner H. Breast cancer survival in Germany: a population-based high resolution study from Saarland. PloS one. 2013;8(7):e70680.

38. Zhu J, Chen J-G, Chen Y-S, Zhang Y-H, Ding L-L, Chen T-Y. Female breast cancer survival in Qidong, China, 1972–2011: a population-based study. BMC cancer.2014;14(1):318.

39. Elhaj AM, Abdalsalam A, Abuidris A, Eltayeb A. Overall survival of females with breast cancer in the National Cancer Institute, University of Gezira, Sudan. Sudan Medical Monitor. 2015;10(1):1.

40. Aziz KKA, Tawfik EA, Shaltout EA, Moneum RAMA. Clinical outcome and survival of breast cancer patients treated at the Clinical Oncology Department, Menoufia University. Menoufia Medical Journal. 2015;28(2):333.

41. Seedhom AE, Kamal NN. Factors affecting survival of women diagnosed with breast cancer in El-Minia Governorate, Egypt. International journal of preventive medicine. 2011;2(3):131.

42. Baghestani A, Moghaddam S, Majd H, Akbari M, Nafissi N, Gohari K. Survival Analysis of Patients with Breast Cancer using Weibull Parametric Model. Asian Pac J Cancer Prev. 2015;16(18):8567-71.

43. Abadi A, Yavari P, Dehghani-Arani M, Alavi-Majd H, Ghasemi E, Amanpour F, et al. Cox models survival analysis based on breast cancer treatments. Iranian journal of cancer prevention. 2014;7(3):124.

44. Balabram D, Turra CM, Gobbi H. Association between age and survival in a cohort of Brazilian patients with operable breast cancer. Cadernos de saudepublica. 2015;31(8):1732-42.

45. Rossi L SD, Pierga J-Y, Lerebours F, Reyal F, Robain M, et al. (2015) Impact of Adjuvant, e0132853.doi:10.1371/journal.pone.0132853 CoBCSAR-WPPO.