Detection and quantification of house mouse *Eimeria* at the species level –

challenges and solutions for the assessment of Coccidia in wildlife

Jarquín-Díaz, Víctor Hugo[1,2]; Balard, Alice[1,2]; Jost, Jenny[1]; Kraft, Julia[1]; Dikmen, Mert Naci[1];

Kvičerová, Jana[3] and Heitlinger, Emanuel[1,2*]

[1]Department of Molecular Parasitology, Institute for Biology, Humboldt University Berlin (HU).

Philippstr. 13, Haus 14, 10115, Berlin, Germany.

[2]Research Group Ecology and Evolution of molecular Parasite-Host Interactions, Leibniz-Institute

for Zoo and Wildlife Research (IZW) im Forschungsverbund Berlin e.V.. Alfred-Kowalke-Straße 17,

10315, Berlin, Germany.

[3]Department of Parasitology, Faculty of Science, University of South Bohemia, Branišovská 1760,

370 05 České Budějovice,

24 *Corresponding author: emanuelheitlinger@gmail.com

Abstract

Detection and quantification of coccidia in studies of wildlife can be challenging. Therefore, prevalence of coccidia is often not assessed at the parasite species level in non-livestock animals. Parasite species – specific prevalences are especially important when studying evolutionary questions in wild populations. We might expect, for example, highly prevalent parasite species to have a lower virulence than lowly prevalent species.

We studied free-living commensal populations of the house mouse (*Mus musculus*) in Germany, and established a strategy to detect and quantify *Eimeria* infections. We show that a novel diagnostic primer targeting the apicoplast genome (Ap5) and coprological assessment after flotation provide complementary detection results increasing sensitivity. Genotyping PCRs confirm detection in a subset of samples and cross-validation of different PCR markers does not indicate bias towards a particular parasite species in genotyping. We were able to detect double infections and to determine the preferred parasite occurrence along the distal-proximal axis of the intestine. Parasite genotyping from tissue samples provides additional indication for the absence of species bias in genotyping amplifications. Three *Eimeria* species were found infecting house mice at different prevalences: *Eimeria ferrisi* (16.1%; 95% CI 12.7 – 20.2), *E. falciformis* (4.2%; 95% CI 2.6 – 6.8) and *E. vermiformis* (1.1%; 95% CI 0.4 – 2.7)*.*

We provide methods for the assessment of prevalences of coccidia at the species level in rodent systems. We discuss the need for broader species level assessment in Coccidia. Prevalence negatively correlates with virulence for *Eimeria* species of house mice, as the more prevalent *E. ferrisi* has been shown to be less virulent than *E. falciformis*. It is an open question whether house mouse *Eimeria* are host specialist and whether prevalence in our system correlates with host range.

Key words

Eimeria, Coccidia, house mice, diagnostic PCR, species specific prevalence, qPCR

Highlights

- Flotation and PCR provide complementary results for *Eimeria* detection in house mice.

- Genotyping PCRs confirm detections.

- *E. ferrisi*, *E. falciformis,* and *E. vermiformis* infect natural populations of *M. musculus*.

- Double infections and preferentially infected tissues could be identified using qPCR.

- Potential virulence prevalence trade-off for *Eimeria* of house mice.

1. Introduction

House mice (*Mus musculus*) are the most commonly used mammalian model for biomedical research worldwide (Houdebine, 2004; Vandenbergh, 2013). Laboratory mouse strains are derived mainly from the subspecies *M. m. domesticus* with genetic contributions from other subspecies (*M.m musculus* and *M. m. castaneus*) (Nishioka, 2011; Yang et al., 2007). Establishment of suitable mouse models to better understand infections with coccidia is an ongoing process. Wild rodents and especially wild house mice are an attractive system for first steps in this direction (Ehret et al., 2017).

Eimeria (Schneider, 1875) is with around 1700 species the most specious genus in the phylum Apicomplexa (Duszynski, 2011; Perkins et al., 2000). For economical reasons, the most studied parasites in this group are those infecting livestock (Shirley et al., 2005; Su et al., 2003). However, at least one third of the total number of described species infects rodents (Levine and Ivens, 1965; Zhao and Duszynski, 2001).

The most used detection and identification method for coccidia is the flotation and microscopical observation of oocysts shed in faeces during the patent period of infection (Ryley et al., 1976). However, unsporulated oocyst are difficult to differentiate into different species (Levine and Ivens, 1988, 1965) and a sporulation process under specific conditions and high degree of expertise in the observation are required (Duszynski and Wilber, 1997; Long and Joyner 1984). For that reason, tools based on DNA amplification and sequencing have been included as routine strategy not only for detection, but also for taxonomic assessment (Fernandez et al., 2003; Hnida and Duszynski, 1999; Kawahara et al., 2010; Morris et al., 2007; Schnitzler et al., 1999; Su et al., 2003; Vrba et al., 2010).

Up to 16 species of *Eimeria* have been described from house mice (Levine and Ivens, 1965) and some of them use different niches in the intestine. The reasons for this diversity are still elusive (Zhao and Duszynski, 2001) and artificial splitting of morphologically plastic forms of the same species (in the same of different hosts) might contribute to this.

Eimeria species described from house mice include *E. falciformis,* the first coccidia described in

house mice (Eimer, 1870), which has sometimes been regarded as the most prevalent species in mice (Becker, 1934; Owen, 1976). This species (and especially the BayerHaberkorn1970 isolate) are the most commonly studied coccidia model in laboratory mice. Life cycle progression (Haberkorn, 1970) and host response (Mesfin et al., 1978; Schelzke et al., 2009; Schmid et al., 2012) are relatively well studied and the whole genome of this species has been sequenced and annotated in detail (Heitlinger et al., 2014).

E. vermiformis was first described in 1971 (Ernst et al., 1971) but since then, to our knowledge, not reported in wild house mice. Similar to *E. falciformis*, most of the information on this species comes from laboratory infection experiments (Figueiredo-Campos et al., 2018; Todd and Lepp, 1971; Rose et al., 1990; Rose and Millard, 1985), making the timing of life cycle progression and its effect on the host relatively well studied.

E. ferrisi was originally described from *M. m. domesticus* from North America (Ankrom et al., 1975a; Levine and Ivens, 1965) but since then to our knowledge investigated only in a single recent laboratory infection showing its virulence to be lower and life cycle to be shorter than that of *E. falciformis* (Al-khlifeh et al., 2019).

To the best of our knowledge, no investigation of prevalences and intensities of coccidia has been conducted in free-living populations of *M. musculus* and its sub-species. In the present work we studied the prevalence of *Eimeria* in house mice from a transect of the well-studied European house mice hybrid zone (HMHZ) (Boursot et al., 2003; Ďureje et al., 2012; Phifer-Rixey and Nachman, 2015). We established methods for detection, species identification and quantification of *Eimeria* in these wild commensal populations of house mice.

2. Material and methods

2.1. Collection of samples

Between 2015 and 2017, 378 house mice (*Mus musculus*) were captured in 96 farms and private properties in a transect 152.27 km long and 114.48 km wide, within the German federal state of Brandenburg (capture permit No. 2347/35/2014) (Figure 1A, Supplementary data 1). On average 20 traps were set overnight per locality. Faeces for microscopical diagnosis of *Eimeria* spp. were preserved in potassium dichromate ($K_2Cr_2O_7$) 2.5% (w/v) and stored at 4°C until further processing,

colon content was preserved in liquid nitrogen and stored at -80°C. For a subset of 163 mice (from

Brandenburg in 2016) tissue samples from cecum and ileum were collected for DNA extraction and

molecular identification of *Eimeria* spp. All samples were kept in liquid nitrogen during

transportation and maintained at -80°C until processing.

2.2. Flotation and microscopical analysis of oocyst

Fecal samples were washed with tap water to eliminate potassium dichromate and homogenized.

After oocyst were flotated using a saturated salt solution (specific gravity = 1.18—1.20), they were

collected by centrifugation (3,234 × g/room temperature/ 10 minutes)*,* washed with distilled water

(1,800 × g/room temperature/10 minutes). The flotations were screened for the presence of oocyst

using a Leica® DM750 M light microscope under the 10X objective. To estimate the intensity of

infection, flotated oocysts were counted using a Neubauer chamber and the results were

expressed as oocyst per gram (OPG) of faeces. Samples were then preserved in a fresh solution

of potassium dichromate 2.5% (w/v) and sporulated in a water bath at 30°C for 10-12 days for

further characterisation.

Eimeria isolates, corresponding to different phylogenetic groups (see below), were selected to take

photomicrographs of sporulated oocysts using a Carl-Zeiss microscope at 100x magnification.

Measurements were made on ~30 oocysts and ~30 sporocysts, using Adobe Photoshop CC

v14.2.1 (3778 pixels = 100, 0000 µm). Length and width were measured and reported in

micrometers. The Length/Width (L/W) ratio was calculated for both oocysts and sporocysts

including means, standard deviation and variation coefficients. Additionally, main morphological

traits (oocyst wall, oocyst residuum, micropyle, polar granule, sporocyst residuum, refractile bodies

and Stieda body) were described, according to the protocol of Duszynski and Wilber (1997).

2.3. DNA extraction

DNA from colon content was extracted using the NucleoSpin® Soil kit (MACHEREY-NAGEL

GmbH & Co. KG, Düren, Germany) according to the manufacturer's recommendations, adding a

mechanical lysis process in a Mill Benchtop Mixer MM 2000 (Retsch GmbH, Haan, Germany).

DNA from cecum and ileum tissues was isolated using the innuPREP DNA Mini Kit (Analytik Jena AG, Jena, Germany) following the instructions of the manufacturer after disruption of the tissue with liquid nitrogen in a mortar. Quality and quantity of isolated DNA was measured spectrophotometrically in a NanoDrop 2000c (Thermo Scientific, Waltham, USA).

2.4. PCR amplification for detection (ap tRNA) and identification (nu 18S rDNA and mt COI)

For detection of *Eimeria*, amplification of a conserved tRNA region of the apicoplast genome (Ap5) was used. Primers Ap5_Fwd (YAAAGGAATTTGAATCCTCGTTT) and Ap5_Rev (YAGAATTGATGCCTGAGYGGTC) were designed based on the complete apicoplast genomes sequences available in the GenBank from *Eimeria tenella* (NC_004823.1), *E. falciformis* (CM008276.1) and *E. nieschulzi* (JRZD00000000.1).

For all samples with oocysts detected during flotation or successful amplification of Ap5, genotyping PCRs were performed to confirmation of detection and further identification of parasite species. A fragment of nuclear small subunit ribosomal RNA (~1,500 bp) and mitochondrial cytochrome C oxidase subunit I (~ 800 bp) were amplified using primers 18S_EF and 18S_ER (Kvičerová et al., 2008) and Cocci_COI_For/Rev (Ogedengbe et al., 2011), respectively. An alternative pair of primers was used in case of failure to amplify COI: Eim_COI_M_F (ATGTCACTNTCTCCAACCTCAGT) and Eim_COI_M_R (GAGCAACATCAANAGCAGTGT). These primers were designed based on the mitochondrial genome of *E. falciformis* (CM008276.1) (Heitlinger et al., 2014) and amplify a ~700 bp fragment of COI gene.

PCR reactions were carried out in a Labcycler (SensoQuest GmbH, Göttingen, Germany) using 0.025 U/µL of DreamTaq™ DNA Polymerase (Thermo Scientific, Waltham, USA), 1X DreamTaq Buffer, 0.5 mM dNTP Mix, 0.25 µM from each primer and 1—20 ng/µL of DNA template in 25 µL reaction. A concentration of 0.25 mM dNTP Mix and a supplementation with 2 mM $MgCl_2$ was used for the amplification of Ap5. The thermocycling protocol consist of 95 °C initial denaturation (4 min) followed by 35 cycles of 92 °C denaturation (45 s), annealing at 52°C (30 s/Eim_COI); 53 °C (45 s/18S_E); 55 °C (30 s/Cocci_COI); 56 °C (30 s/Ap5); 72 °C extension 90 s (18S_E), 20 s (Cocci_COI/Eim_COI) or 45s (Ap5) and a final extension at 72 °C (10 min). DNA from oocyst of *E. falciformis* BayerHaberkorn1970 strain and DNA from colon content of a non-infected NMRI mouse

were used as positive and negative controls, respectively.

All PCR products with the expected size were purified using the SAP-Exo Kit (Jena Bioscience

GmbH, Jena, Germany) and Sanger sequenced from both directions by LGC Genomics (Berlin,

Germany). Quality assessment and assembly of forward and reverse sequence was performed in

Geneious v6.1.8. All sequences were submitted to NCBI GenBank (Accession numbers: nu 18S

rDNA [MH751925—MH752036] and mt COI [MH777467—MH777593 and MH755302—

MH755324] (Supplementary data S2).

2.5. Eimeria detection in tissue by qPCR

For mice collected in 2016 (*n*= 163) cecum and ileum tissue was screened using qPCR. Primers

targeting a short fragment of mt COI were used to amplify DNA from from intracellular stages of

Eimeria (Eim_COI_qX-F, TGTCTATTCACTTGGGCTATTGT; Eim_COI_qX-R

GGATCACCGTTAAATGAGGCA). Amplification reactions with a final volume of 20 µL contained 1X

iTaq™ Universal SYBR® Green Supermix (Bio-Rad Laboratories GmbH, München, Germany),

400 nM of each primer and 50 ng of DNA template. Cycling in a Mastercycler® RealPlex 2

(Eppendorf, Hamburg, Germany) was performed with the following program: 95 °C initial

denaturation for 2 min, followed by 40 cycles of denaturation at 95 °C for 15 s, annealing at 55 °C

for 15 s and extension 68 °C for 20 s. Melting curves were analysed to detect eventual primer

dimer formation or non-specific amplification. As internal reference for relative quantification the

CDC42 gene from the nuclear genome of the house mouse was amplified (Ms_gDNA_CDC42_F

CTCTCCTCCCCTCTGTCTTG; Ms_gDNA_CDC42_R TCCTTTTGGGTTGAGTTTCC). Infection

intensity was estimated as the ΔCt between mouse and *Eimeria* amplification (CtMouse-

Ct*Eimeria*). To correct for background noise a detection threshold was estimated at ΔCt = -5 and

only results above this value were considered infected. ΔCtIleum and ΔCtCecum were compared

for samples above the threshold in both tissues to assess primary tissue occurrence (Ahmed et al.,

2019). In samples positive for qPCR, *Eimeria* genotyping was performed based on DNA extracted

from tissue, as described above (see *2.4*).

2.6. Molecular identification of Eimeria spp. isolates: 18S and COI phylogenetic analysis.

As strategy for molecular identification, datasets of nu 18S and mt COI sequences were compiled.

Sequences generated for the present work were compared to databases sequences using NCBI

BLAST and most similar sequences were selected. Based on this, sequences for *E. falciformis, E.*

vermiformis and *E. ferrisi* were downloaded from GenBank as a reference. COI sequences were

aligned by translation using the Multiple Align algorithm and translation frame 1 with the genetic

code for "mold protozoan mitochondrial", 18S sequences were aligned using MUSCLE (Edgar,

2004), both through Geneious v6.1.8.

Phylogenetic trees for all datasets were constructed using Maximum Likelihood (ML) and Bayesian

inference (BI) methods, implemented in PhyML v3.0 (Guindon et al., 2010) and MrBayes v3.2.6

(Huelsenbeck and Ronquist, 2001; Ronquist et al., 2012), respectively. The most appropriate

evolutive models for each datasets were determined in JModelTest v2.1.10 (Posada, 2008). For

ML trees, a bootstrap analysis with 1000 replicates was performed, whereas MCMC for BI was run

with two cold and two hot chains for 1,000,000 generations or until the average split frequency was

below 0.05. The concatenated dataset was analysed using partitions and locus-specific models.

Visualization of the trees was done with FigTree v1.4.2 (Rambaut, 2012).

2.7. Statistical analysis

All statistical analyses were performed in R (R Development Core Team, 2008). Prevalence of

Eimeria was calculated as the proportion of positive samples in the total number of analysed

samples. The 95% confidence interval [CI 95%] was calculated using the Sterne's exact method

(Sterne, 1954) implemented in the package "epiR" v0.9-99 (Nunes et al., 2018).

To assess the significance in primer bias, logistic regression models were used to estimate the

probability to successfully amplify and sequence a specific genetic marker for each *Eimeria*

species. The response variable in these models was the amplification and sequencing success

with a particular primer pair (COCCI_COI_F/R, Eim_COI_M_F/R or 18S_E F/R), and the predictors

were the species identity (as determined with the other markers only, to make response and

predictors independent) and additionally the detection of an infection with Ap5 and Flotation. These

models were fitted first for COCCI_COI_F/R as response, then for the combined probability of

successful COI genotyping and finally for 18S genotyping as response using the package "stats"

(R Development Core Team). Tables were produced for the summary of models using the package

"jtools" v2.0.0 (Long, 2019).

Differences on oocyst and sporocyst L/W ratios between *Eimeria* species were tested for

significance with an analysis of variance fitting a linear model using the species assignment as

predictor with a Tukey HSD post hoc test adjusting for multiple comparisons, with the package

"stats" (R Development Core Team).

3. Results

3.1. Sampling and Eimeria spp*. detection*

We used flotation of oocyst from faeces and PCR amplification of a novel diagnostic maker (Ap5)

from colon content DNA to detect *Eimeria* parasites in a total of 378 house. Overall prevalence was

25.9% [95% CI = 21.7 – 30.7] (98/378) for PCR and 27.0% [95% CI = 22.7 – 31.7%] (102/378) for

flotation. These estimates are not significantly different. Nevertheless, both techniques considered

together estimate a higher prevalence of 37.6% [95% CI = 32.8 – 42.6] (142/378), meaning that

some positive results were detected by only one of the techniques (Fig. 1B). We further aimed to

provide species specific specific identification and to consolidate results from the two different

detection methods.

3.2. Molecular identification of Eimeria isolates - (phylogenetic analysis nu 18S and mt COI)

Eimeria species were identified by phylogenetic analysis of nu 18S and mt COI sequences, the

most commonly used molecular markers of apicomplexan parasites. To identify our isolates,

sequences were compared with references from the NCBI database. Sequences from three

previously described *Eimeria* species infecting *M. musculus* showed highest BLAST similarities

and phylogenetic clustering. This approach ignores the problem whether isolates from different

hosts would be assigned to the same phylogenetic clusters while they are regarded different

species by taxonomists.

The nu 18S phylogenetic tree was inferred based on 80 sequences (540—1,795 bp), 73 of them

from wild house mice generated in our study (3 from ileum tissue, 16 from cecum tissue and 54

from colon content, see below). *Eimeria* species previously described in house mouse were

represented by *E. falciformis* (AF080614), *E. vermiformis* (KT184355) and *E. ferrisi* (KT360995).

In addition, one newly generated sequence from *E. falciformis* strain BayerHaberkorn1970

(MH751998) was also included. Sequence identity of our isolates to this references sequences

was above 98% and even 100% in most of the cases for this marker. *Isospora* sp. sequences

identified in *Talpa europaea* moles were used as outgroup. Both ML and BI rooted trees shared a

topology placing our sequences at the same positions in relation to reference sequences with high

support (bootstrap values and posterior probabilities are shown in Fig. 2A). The sequences

clustered in three well supported monophyletic groups (Fig. 2A).

The phylogenetic tree for mt COI was based on 103 sequences (519—804 bp), 97 of which were

obtained from *Eimeria* infecting wild house mice (3 from ileum, 16 from cecum tissue and 78 from

colon content). Reference sequences from house mouse *Eimeria* (*E. ferrisi*, KT361028; *E.*

falciformis, KX495129 and MH777539; *E. vermiformis* JN205071) identified by BLAST searches

showed an identity of above 98% to respective groups of our isolates. We defined *Isospora* sp.

from *Talpa europaea* as outgroup for rooting. ML and BI rooted trees based on alignments of these

COI sequences shared a general topology with respect to the placement of our isolates in relation

to reference sequences. Bootstrap values and posterior probabilities for support of these

placements are shown in Fig. 2B. The sequences derived from house mice cluster in three

monophyletic groups including reference sequences for *E. falciformis* (*n*= 17, sequences from our

study), *E. ferrisi* (*n*= 72) and *E. vermiformis* (*n*= 8) (Fig 2B). Phylogenies based on concatenated

supermatrices for the two markers show the same topology concerning placement of isolates from

the present study (Supplementary data S3 and S4).

3.3. Morphometrical and morphological comparison of oocysts

For further support assignment of the three phylogenetic groups of *Eimeria* from house mouse, we

characterized sporulated oocysts morphologically (Table 1). *E. falciformis, E. ferrisi* and *E.*

vermiformis oocyst shared most of the traits we evaluated and showed overlapping morphometry

(Fig. 3A). The length/width ratio of *E. vermiformis* oocysts, however, was significantly higher (1.29;

95%CI = 1.26—1.33; *n*= 35) than that of *E. falciformis* (1.17; 95%CI = 1.14—1.20; *n*= 31) and *E.*

ferrisi oocysts (1.23; 95%CI = 1.21—1.25; *n*= 127) (Tukey HSD, *p* <0.05) (Fig. 3B; Supplementary

data S5). This means that *E. vermiformis* has more ellipsoidal oocysts than the other two species.

Other morphological characteristics of oocysts (smooth wall, absence of micropyle, presence of

polar granule and absence of oocyst residuum) are very similar or identical between the three

species (Table 1).

Morphological measurements of sporocysts are not significantly different between the three

species found in house mice (Fig. 3C). We also observed the presence of a sporocyst residuum

and refractile bodies uniformly in all species. Nevertheless, Stieda bodies were absent in *E.*

falciformis only, in disagreement with previous descriptions (Eimer, 1870; Haberkorn, 1970).

3.5. Proximal-distal occurrence of infection and double infections

We detected DNA from endogenous stages by qPCR in 27 of 163 samples analysed

(Supplementary data S6). We differentiate detection between small and large intestine, analysing

ileum as the most distal tissue of the small intestine and cecum as the most proximal tissue of the

large intestine. Detection was either limited just to cecum (n = 19), to ileum (n = 2) or possible in

both tissues (n = 6). Infections in cecum were identified as *E. falciformis* (n = 4), *E. ferrisi* (n = 17)

and *E. vermiformis* (n = 1) Detections in ileum (n = 2) were identified as *E. vermiformis*. In two mice

positive in both tissues, it was possible to identify *E. ferrisi* in cecum and *E. vermiformis* in ileum,

providing evidence that these animals presented a double infection (that is, simultaneous infections

with different isolates; Fig. 4).

3.6. Amplification efficiency of different markers

As *Eimeria* detection by PCR and the determination of species identity could be biased especially

in cases of double infections, we analysed differences in amplification efficiency of the three primer

pairs used for the molecular identification of *Eimeria* species (Table 2). In a cross-validation

approach we compare the likelihood to amplify a maker given the species identification with the

other marker. The amplification and sequencing efficiency of Cocci_COI primer pair was

significantly higher for *E. ferrisi* isolates (logistic regression, $p < 0.001$) than for *E. falciformis*

isolates (the letter determined by the 18S marker). Using the novel primer pair Eim_COI_M_F/R

the sequencing results were complemented and we detected no significant differences in the

combined amplification efficiency for both primers (logistic regression; $p = 0.62$).

Similarly, we did not detect significant differences in the probability to obtain an 18S sequence

(using the 18S_EF/R primer pair) between *Eimeria* species as determined by (combined) COI

assessment (logistic regression; $p = 0.25$). Differences in PCR efficiency for Cocci_COI make it

likely that markers amplify different species in case of double infections in a single isolate. Both

detection by flotation and diagnostic (Ap5) PCR significantly increase the likelihood amplification

for all primers (COI or 18S). The statistical models for biased amplification hence also confirm that

both detection methods provide complementary results, even while controlling for detection with

the other method (Table 2).

3.7. Prevalence of the different Eimeria species

Genotyping amplification and sequencing with either 18S or COI primers (or both) was possible for

samples in which infections had been detected (n=81) (Fig. 1B). Both COI and 18S genotyping

PCRs hence support detection by flotation and diagnostic PCRs in this subset of samples.

Furthermore amplification of both or either one maker was fully sufficient to assign the isolate to an

Eimeria species (Fig. 2), allowing us to resolve prevalence on the species level.

These corrections and controls allow us to determine prevalence at the species level: *E. ferrisi* was

identified at a higher prevalence of 16.1% (61/378, 95%CI = 12.7 — 20.2) in comparison to *E.*

falciformis (16/378, 4.2% [95%CI = 2.6—6.8]) and *E. vermiformis* (4/378, 1.1% [0.4—2.7]).

4. Discussion

In this study we identify *Eimeria* species in wild commensal populations of house mice (*Mus*

musculus). We show that detection and identification of this group of rodent coccidia can be

challenging and propose to complement classical coprological assessment with molecular tools: a

highly sensitive detection PCR, genotyping PCRs for species identification and qPCR for

localization and detection of double infections. Based on this we identified three different *Eimeria*

species in the house mouse: *E. ferrisi*, *E. falciformis* and *E. vermiformis.* Morphological

characteristics and preferential occurrence were congruent with the assignment of isolates to the

above species. We give examples for the need for *Eimeria* species identification in predictions for

two evolutionary relevant hypotheses: virulence-prevalence and host range-prevalence trade-off.

Only few studies report prevalences of *Eimeria* in wild populations of rodents. Prevalences ranging

between 20 and 40% for isolates unclassified as either as *E. falciformis*, *E. ferrisi* and/or *E.*

vermiformis were estimated before for wild populations of *Mus musculus* in UK, USA and Australia

(Ball and Lewis, 1984; Moro et al., 2003; Tattersall et al., 1994). A recent study in rodents (other

then *Mus musculus*) reported an *Eimeria* spp. prevalence of 32.7% based on coprological

observations (Mácová et al., 2018). In comparison, overall prevalence for all *Eimeria* species in our

study is only slightly higher (37.6%) in the house mouse. While flotation is the most commonly

used for detection and quantification of coccidia (Hobbs et al., 1999; Hu et al., 2016; Rinaldi et al.,

2011), we here used a complementary approach of flotations and diagnostic PCRs.

Traditional identification of *Eimeria*, depends on the expertise to recognise the morphology of

sporulated oocyst (Levine and Ivens, 1965). We show that interpretation of morphometrical data is

complex due overlap between species while measurement means agree with literature (Table 1).

Considering the challenges of identification and characterisation of *Eimeria* isolates from field

samples, we conclude that characterisation of *Eimeria* species requires molecular markers and

phylogenetic analysis.

Sequence identity of our isolates to reference sequences from *Eimeria* species previously

described in *M. musculus* was above 98% for COI, which is sometimes assumed sufficient

correspond differences within species of *Eimeria* (Yang et al., 2015). We confirm taxonomic

assignment based on highly supported maximum likelihood and Bayesian phylogenetic clustering

of 18S and COI sequences. Moreover, the three identity of the three *Eimeria* species was

supported by phenotypic characteristics: morphometry of oocysts and tissue occurrence of the

infection.

For some *Eimeria* species precise tissue localization is described based on histology or electron microscopy (Ankrom et al., 1975; Šlapeta et al., 2001). Both methods provide detailed information on developmental stages, but are also time consuming and require a high level of expertise. As an alternative to determine (only) the rough occurrence of the infection along the proximal-distal axis of the intestine, a DNA based qPCR method allowed us not only to detect the presence of *Eimeria*, but also to estimate tissue specific intensity of infection. While infection with rodent *Eimeria* in general can be limited to the duodenum and jejunum (Kvičerová et al., 2007), house mouse *Eimeria* have been described to be mostly found in either the small or the large intestine (Ankrom et al., 1975; Ernst et al., 1971; Haberkorn, 1970; Levine and Ivens, 1965). The ileum as the most distal part of the small intestine and the cecum as the most proximal part of the large intestine thus provide an easy differentiation in these large small intestinal categories of infection. Genotyping DNA derived from tissue allowed to detect double infections with *E. vermiformis* in the small intestine and *E. ferrisi* in the large intestine. Localization generally agrees with previous descriptions for the isolates we identified as *E. ferrisi*, *E. falciformis* and *E. vermiformis* by phylogenetic clustering. To our knowledge we provide the first report of double infections in wild rodents.

Double infections can be problematic for identification of species by genotyping. Simultaneous infection of the caecum with *E. ferrisi* and *E. falciformis* would not be recognized with our qPCR method. We shown that amplification of COI with the commonly used primer pair Cocci_COI (Ogedengbe et al., 2011) is differentially efficient for different *Eimeria* isolates. This primer preference can lead to a misidentification in double infections due to the generation of "chimeric" isolates that present different and contradictory information for different markers. Attention to such discrepancies is needed when collating database sequences and when developing multi-marker approaches in general. For rodent *Eimeria* systems, we develop an alternative COI primer pair and find no evidence for differential amplification bias in our cross-validation of the different primer pairs.

E. ferrisi is by far the most prevalent species in our study area infecting *M. musculus*. Concerning the house mouse hybrid zone we find infections in *M. m. domesticus, M. m. musculus* and hybrids, suggesting that there are no strict geographical or host subspecies constraints for this species.

Population structure for *E. ferrisi* (which could in turn correspond to host subspecies (Goüy de

Bellocq et al., 2018; Kváč et al., 2013) can not be found at the resolution the analysed markers

provide. In addition to the currently used *E. falciformis* and *E. vermiformis* (Al-khlifeh et al., 2019)

Differences in prevalence between parasites found in wildlife systems can be integrated in an

evolutionary perspective, allowing predictions on virulence and host range:

A lower virulence (reduction of host fitness caused by a parasite (Frank, 1996) should be

associated with higher prevalences, because of virulence-transmission trade-offs (Anderson and

May, 1982). Especially in relatively virulent parasites (such as *Eimeria* in general) infected

individuals might suffer from moribundity and mortality reducing the rate of interaction between

infected and uninfected hosts and the length of transmission periods (Leggett et al., 2013). While

both the highly prevalent *E. ferrisi* and the lowly prevalent *E. falciformis* and *E. vermiformis* has

been described pathogenic and potentially lethal in mice in laboratory infections (Blagburn and

Todd, 1984; Klesius and Hinds, 1979; Mesfin et al., 1978), a direct comparison found virulence of

E. ferrisi to be lower than that of *E. falciformis* (Al-khlifeh et al., 2019). In this regard the species

specific prevalence in our system matches a prediction from one of the most central hypotheses in

evolutionary epidemiology of parasites.

Two different hypotheses predict different prevalence of parasites with different host ranges. Under

the trade-off hypothesis (Poulin and Combes, 1999) specialism is associated with higher

prevalence as generalist parasites are less well adapted to infect many individuals in single

species. Under the resource breadth hypothesis (Krasnov et al., 2005), however, species with a

broad host range would be generally better at exploiting all host species and have higher

prevalence. Predictions from both hypotheses might be integrated when taking into account the

structure of differences between hosts (Pinheiro et al., 2016). We emphasize that we did not test

host range of rodent *Eimeria* in the present work but used the assumption of biologically

meaningful taxonomy and traditional "morphospecies" within house mice. This means that open

questions on host specificity of rodent *Eimeria* (Mácová et al., 2018) could in future be investigated

on the background of prevalences resolved on the level of "morphospecies" in the house mouse

host.

In conclusion we argue that *Eimeria* in wildlife populations should be identified more frequently at

the level of species previously described by taxonomist. We propose to integrate a set of simple

methods into a reproducible procedure to achieve this aim. For Coccidians, as important parasites

of vertebrates, only species specific assessment will allow to test hypotheses under the framework

of evolutionary epidemiology.

5. References

Ahmed, N., Heitlinger, E., Affinass, N., Kühl, A.A., Xenophontos, N., Jarquin, V.H., Jost, J.,
Steinfelder, S., Hartmann, S., 2019. A Novel Non-invasive Method to Detect RELM Beta
Transcript in Gut Barrier Related Changes During a Gastrointestinal Nematode Infection.
Front. Immunol. 10, 1–11. https://doi.org/10.3389/fimmu.2019.00445

Al-khlifeh, E., Balard, A., Jarquín-Díaz, V.H., Weyrich, A., Wibbelt, G., Heitlinger, E., 2019. Eimeria
falciformis BayerHaberkorn1970 and novel wild derived isolates from house mice : differences
in parasite lifecycle , pathogenicity and host immune reactions. bioRxiv 1–33.
https://doi.org/http://dx.doi.org/10.1101/611277

Anderson, R.M., May, R.M., 1982. Coevolution of Hosts and Parasites. Parasitology 85, 411–426.
https://doi.org/10.1017/S0031182000055360

Ankrom, S.L., Chobotar, B., Ernest, J. V., 1975. Life Cycle of Eimeria ferrisi Levine & Ivens, 1965 in
the Mouse, Mus musculus. J. Protozool. 22, 317–323.
https://doi.org/https://doi.org/10.1111/j.1550-7408.1975.tb05177.x

Ball, S.J., Lewis, D.C., 1984. Eimeria (Protozoa: Coccidia) in wild populations of some British
rodents S. J. Zool. 202, 373–381. https://doi.org/https://doi.org/10.1111/j.1469-
7998.1984.tb05089.x

Becker, E.R., 1934. Coccidia and Coccidiosis of Domesticated, Game and Laboratory Animals and
of Man, JAMA: The Journal of the American Medical Association. Akademiai Kiado., Berlin;
Verlag Paul Parey: Budapest, Hungary. https://doi.org/10.1001/jama.1934.02750260071034

Blagburn, B.L., Todd, K.S., 1984. Pathological Changes and Immunity Associated with
Experimental Eimeria vermiformis Infections in Mus musculus. J. Protozool. 31, 556–561.
https://doi.org/10.1111/j.1550-7408.1984.tb05502.x

Boursot, P., Auffray, J.-C., Britton-Davidian, J., Bonhomme, F., 2003. The Evolution of House Mice.
Annu. Rev. Ecol. Syst. 24, 119–152. https://doi.org/10.1146/annurev.es.24.110193.001003

Ďureje, L., Macholán, M., Baird, S.J.E., Piálek, J., 2012. The mouse hybrid zone in Central Europe:
From morphology to molecules. Folia Zool. 61, 308–318.
https://doi.org/https://doi.org/10.25225/fozo.v61.i3.a13.2012

Duszynski, D.W., 2011. Eimeria, in: ELS. John Wiley & Sons, Ltd, Chichester, UK, pp. 1192–1196.
https://doi.org/10.1002/9780470015902.a0001962.pub2

Duszynski, D.W., Wilber, P.G., 1997. A guidelines for the preparation of species description in the
Eimeriidae. J. Parasitol. 83, 333–336. https://doi.org/10.2307/3284470

Edgar, R.C., 2004. MUSCLE: Multiple sequence alignment with high accuracy and high
throughput. Nucleic Acids Res. 32, 1792–1797. https://doi.org/10.1093/nar/gkh340

Ehret, T., Seeber, F., Torelli, F., Klotz, C., Pedersen, A.B., 2017. Translational Rodent Models for
Research on Parasitic Protozoa—A Review of Confounders and Possibilities. Front. Cell.
Infect. Microbiol. 7. https://doi.org/10.3389/fcimb.2017.00238

Eimer, T., 1870. Ueber die ei-oder kugelförmigen sogenannten Psorospermien der Wirbelthiere:
Ein Beitrag Zur Entwicklungsgeschichte Der Gregarinen Und Zur Kenntniss Dieser Parasiten
Als Krankheitsursache. A. Stuber.

Ernst, J. V., Chobotar, B., Hammond, D.M., 1971. The Oocysts of Eimeria vermiformis sp. n. and E.
papillata sp. n. (Protozoa: Eimeriidae) from the Mouse Mus musculus. J. Protozool. 18, 221–
223.

Fernandez, S., Pagotto, A.H., Furtado, M.M., Katsuyama, Â.M., Madeira, A.M.B.N., Gruber, A.,
2003. A multiplex PCR assay for the simultaneous detection and discrimination of the seven
Eimeria species that infect domestic fowl. Parasitology 127, 317–325.
https://doi.org/10.1017/S0031182003003883

Figueiredo-Campos, P., Ferreira, C., Blankenhaus, B., Veldhoen, M., 2018. Eimeria vermiformis
Infection Model of Murine Small Intestine. BIO-PROTOCOL 8.
https://doi.org/10.21769/BioProtoc.3122

Frank, S.A., 1996. Models of Parasite Virulence. Q. Rev. Biol. 71, 37–78.

Goüy de Bellocq, J., Wasimuddin, Ribas, A., Bryja, J., Piálek, J., Baird, S.J.E., 2018. Holobiont
suture zones: Parasite evidence across the European house mouse hybrid zone. Mol. Ecol.
5214–5227. https://doi.org/10.1111/mec.14938

Guindon, S., Dufayard, J.-F., Lefort, V., Anisimova, M., Hordijk, W., Gascuel, O., 2010. New
Algorithms and Mehtods to Estimate Maximum-Likelihood Phylogenies: Asessing the
Performance of PhyML 2.0. Syst. Biol. 59, 307–321. https://doi.org/10.1093/sysbio/syq010

Haberkorn, A., 1970. Die Entwicklung von Eimeria falciformis (Eimer 1870) in der weissen Maus
(Mus musculus). Zeitschrift für Parasitenkunder 34, 49–67.
https://doi.org/10.1007/BF00629179

Heitlinger, E., Spork, S., Lucius, R., Dieterich, C., 2014. The genome of Eimeria falciformis -
reduction and specialization in a single host apicomplexan parasite. BMC Genomics 15, 1–17.
https://doi.org/10.1186/1471-2164-15-696

Hnida, J. a, Duszynski, D.W., 1999. Taxonomy and phylogeny of some Eimeria
(Apicomplexa:Eimeriidae) species of rodents as determined by polymerase chain
reaction/restriction-fragment-length polymorphism analysis of 18S rDNA. Parasitol. Res. 85,
887–94. https://doi.org/10.1007/s004360050653

Hobbs, R.P., Twigg, L.E., Elliot, A.D., Wheeler, A.G., 1999. Factors Influencing the Fecal Egg and
Oocyst Counts of Parasites of Wild European Rabbits Oryctolagus cuniculus (L .) in Southern
Western Australia Author (s): Russell P . Hobbs , Laurie E . Twigg , Aileen D . Elliot and
Amanda G . Wheeler Published by. J. Parasitol. 85, 796–802.

Houdebine, L.-M., 2004. Chapter 6 - The Mouse as an Animal Model for Human Diseases BT - The
laboratory mouse, The laboratory mouse. Elsevier Ltd. https://doi.org/10.1016/B978-0-12-
336425-8.50059-5

Hu, X.L., Liu, G., Wang, W.X., Zhou, R., Liu, S.Q., Li, L.H., Hu, D.F., 2016. Methods of preservation
and flotation for the detection of nematode eggs and coccidian oocysts in faeces of the forest

musk deer. J. Helminthol. 90, 680–684. https://doi.org/10.1017/S0022149X15000942

Huelsenbeck, J.P., Ronquist, F., 2001. MrBayes: Bayesian inference of phylogenetic trees.
Bioinformatics 17, 754–755. https://doi.org/10.1093/bioinformatics/17.8.754

Kawahara, F., Zhang, G., Mingala, C.N., Tamura, Y., Koiwa, M., Onuma, M., Nunoya, T., 2010.
Genetic analysis and development of species-specific PCR assays based on ITS-1 region of
rRNA in bovine Eimeria parasites. Vet. Parasitol. 174, 49–57.
https://doi.org/10.1016/j.vetpar.2010.08.001

Kenneth S. Todd, J., Lepp, D.L., 1971. The Life Cycle of Eimeria vermiformis Ernst, Chobotar and
Hammond, 1971 in the Mouse Mus musculus. J. Protozool. 18, 332–337.

Klesius, P.H., Hinds, S.E., 1979. Strain-dependent differences in murine susceptibility to coccidia.
Infect. Immun. 26, 1111–1115.

Krasnov, B.R., Poulin, R., Shenbrot, G.I., Mouillot, D., Khokhlova, I.S., 2005. Host specificity and
geographic range in haematophagous ectoparasites. Oikos 108, 449–456.
https://doi.org/10.1111/j.0030-1299.2005.13551.x

Kváč, M., McEvoy, J., Loudová, M., Stenger, B., Sak, B., Květoňová, D., Ditrich, O., Rašková, V.,
Moriarty, E., Rost, M., Macholán, M., Piálek, J., 2013. Coevolution of Cryptosporidium tyzzeri
and the house mouse (Mus musculus). Int. J. Parasitol. 43, 805–817.
https://doi.org/10.1016/j.ijpara.2013.04.007

Kvičerová, J., Pakandl, M., Hypša, V., 2008. Phylogenetic relationships among Eimeria spp.
(Apicomplexa, Eimeriidae) infecting rabbits: Evolutionary significance of biological and
morphological features. Parasitology 135, 443–452.
https://doi.org/10.1017/S0031182007004106

Kvičerová, J., Ptáčková, P., Modrý, D., 2007. Endogenous development, pathogenicity and host
specificity of Eimeria cahirinensis Couch, Blaustein, Duszynski, Shenbrot and Nevo, 1997
(Apicomplexa: Eimeriidae) from Acomys dimidiatus (Cretzschmar 1826) (Rodentia: Muridae)
from the Near East. Parasitol. Res. 100, 219–226. https://doi.org/10.1007/s00436-006-0251-7

Leggett, H.C., Buckling, A., Long, G.H., Boots, M., 2013. Generalism and the evolution of parasite
virulence. Trends Ecol. Evol. 28, 592–596. https://doi.org/10.1016/j.tree.2013.07.002

Levine, N.D., Ivens, V., 1988. Cross-Transmission of Eimeria spp. (Protozoa, Apicomplexa) of
Rodents -A Review. J. Protozool. 35, 434–437.

Levine, N.D., Ivens, V., 1965. The Coccidian Parasites (Protozoa, Sporozoa) of Rodents. The
University of Ilinois Press, Urbana, Ill.

Long, J., 2019. jtools: Analysis and Presentation of Social Scientific Data.

Long, P.L., Joyner, L.P., 1984. Problems in the identification of species of *Eimeria*. J. Protozool. 31,
535–541.

Mácová, A., Hoblíková, A., Hypša, V., Stanko, M., Martinů, J., Kvičerová, J., 2018. Mysteries of
host switching: Diversification and host specificity in rodent-coccidia associations. Mol.
Phylogenet. Evol. 127, 179–189. https://doi.org/10.1016/j.ympev.2018.05.009

Mesfin, G.M., Bellamy, J.E., Stockdale, P.H., 1978. The pathological changes caused by Eimeria
falciformis var pragensis in mice. Can. J. Comp. Med. 42, 496–510.

Moro, D., Lawson, M.A., Hobbs, R.P., Thompson, R.C.A., 2003. Pathogens of House Mice on Arid
Boullanger Island and Subantarctic Macquarie Island, Australia. J. Wildl. Dis. 39, 762–771.
https://doi.org/10.7589/0090-3558-39.4.762

Morris, G.M., Woods, W.G., Grant Richards, D., Gasser, R.B., 2007. The application of a
polymerase chain reaction (PCR)-based capillary electrophoretic technique provides detailed
insights into Eimeria populations in intensive poultry establishments. Mol. Cell. Probes 21,
288–294. https://doi.org/10.1016/j.mcp.2007.03.001

Nishioka, Y., 2011. The origin of common laboratory mice. Genome 38, 1–7.
https://doi.org/10.1139/g95-001

Nunes, M.S., Heuer, T.C., Marshall, J., Sanchez, J., Thornton, R., Reiczigel, J., Robison-Cox, J.,
Sebastiani, P., Solymos, P., Yoshida, K., Jones, G., Pirikahu, S., Firestone, S., Kyle, R., Popp,
J., Jay, M., 2018. epiR: Tools for the analysis of epidemiological data.

Ogedengbe, J.D., Hanner, R.H., Barta, J.R., 2011. DNA barcoding identifies Eimeria species and
contributes to the phylogenetics of coccidian parasites (Eimeriorina, Apicomplexa, Alveolata).
Int. J. Parasitol. 41, 843–850. https://doi.org/10.1016/j.ijpara.2011.03.007

Owen, D., 1976. Some parasites and other organisms of wild rodents in the vicinity of an SPF unit.
Lab. Anim. 10, 271–278. https://doi.org/10.1258/002367776781035314

Perkins, F.O., Barta, J.R., Clopton, R.E., Peirce, M.A., Upton, S.J., 2000. Apicomplexa, in: Lee, J.,
Leedale, G., Bradbury, P. (Eds.), The Illustrated Guide to the Protozoa. Allen Press, pp. 190–
304.

Phifer-Rixey, M., Nachman, M.W., 2015. Insights into mammalian biology from the wild house
mouse Mus musculus. Elife 2015, 1–13. https://doi.org/10.7554/eLife.05959

Pinheiro, R.B.P., Félix, G.M.F., Chaves, A. V., Lacorte, G.A., Santos, F.R., Braga, É.M., Mello,
M.A.R., 2016. Trade-offs and resource breadth processes as drivers of performance and
specificity in a host-parasite system: A new integrative hypothesis. Int. J. Parasitol. 46, 115–
121. https://doi.org/10.1016/j.ijpara.2015.10.002

Posada, D., 2008. jModelTest: Phylogenetic model averaging. Mol. Biol. Evol. 25, 1253–1256.
https://doi.org/10.1093/molbev/msn083

Poulin, R., Combes, C., 1999. The concept of virulence: Interpretations and implications. Parasitol.
Today 15, 474–475. https://doi.org/10.1016/S0169-4758(99)01554-9

Rambaut, A., 2012. FigTree.

Rinaldi, L., Coles, G.C., Maurelli, M.P., Musella, V., Cringoli, G., 2011. Calibration and diagnostic
accuracy of simple flotation, McMaster and FLOTAC for parasite egg counts in sheep. Vet.
Parasitol. 177, 345–352. https://doi.org/10.1016/j.vetpar.2010.12.010

Ronquist, F., Teslenko, M., Van Der Mark, P., Ayres, D.L., Darling, A., Höhna, S., Larget, B., Liu, L.,
Suchard, M. a., Huelsenbeck, J.P., 2012. Mrbayes 3.2: Efficient bayesian phylogenetic
inference and model choice across a large model space. Syst. Biol. 61, 539–542.
https://doi.org/10.1093/sysbio/sys029

Rose, M.E., Millard, B.J., 1985. Host specificity in eimerian coccidia: development of *Eimeria*
vermiformis of the mouse, *Mus musculus*, in *Rattus norvegicus*. Parasitology.

Rose, M.E., Wakelin, D., Hesketh, P., 1990. Eimeria vermiformis: Differences in the course of

primary infection can be correlated with lymphocyte responsiveness in the BALB/c and
C57BL/6 mouse, Mus musculus. Exp. Parasitol. 71, 276–283. https://doi.org/10.1016/0014-
4894(90)90032-8

Ryley, J.F., Meade, R., Hazelhurst, J., Robinson, T.E., 1976. Methods in coccidiosis research:
Separation of oocysts from faeces. Parasitology 73, 311–326.
https://doi.org/10.1017/S0031182000046990

Schelzke, K., Lucius, R., Liesenfeld, O., Steinfelder, S., Pogonka, T., Stange, J., Papadakis, K.,
2009. CD8+ cells protect mice against reinfection with the intestinal parasite Eimeria
falciformis. Microbes Infect. 12, 218–226. https://doi.org/10.1016/j.micinf.2009.12.005

Schmid, M., Lehmann, M.J., Lucius, R., Gupta, N., 2012. Apicomplexan parasite, Eimeria
falciformis, co-opts host tryptophan catabolism for life cycle progression in mouse. J. Biol.
Chem. 287, 20197–20207. https://doi.org/10.1074/jbc.M112.351999

Schneider, A., 1875. Note sur la psorospermie oviforme du poulpe. Arch. Zool. exp. gen, 4, 4045.

Schnitzler, B.E., Thebo, P.L., Tomley, F.M., Uggla, A., Shirley, M.W., 1999. PCR identification of
chicken Eimeria: A simplified read-out. Avian Pathol. 28, 89–93.
https://doi.org/10.1080/03079459995091

Shirley, M.W., Smith, A.L., Tomley, F.M., 2005. The biology of avian Eimeria with an emphasis on
their control by vaccination. Adv. Parasitol. 60, 285–330. https://doi.org/10.1016/S0065-
308X(05)60005-X

Šlapeta, J.R., Modrý, D., Votýpka, J., Jirku, M., Oborník, M., Lukeš, J., Koudela, B., 2001. Eimeria
telekii n.sp. (Apicomplexa: Coccidia) from Lemniscomys striatus (Rodentia: Muridae):
Morphology, pathology and phylogeny. Parasitology 122, 133–143.
https://doi.org/10.1017/S0031182001007259

Sterne, T.E., 1954. Some remarks on confidence or fiducial limits. Biometrika 41, 275–278.
https://doi.org/10.1093/biomet/41.1-2.275

Su, Y.C., Fei, A.C.Y., Tsai, F.M., 2003. Differential diagnosis of five avian Eimeria species by
polymerase chain reaction using primers derived from the internal transcribed spacer 1 (ITS-
1) sequence. Vet. Parasitol. 117, 221–227. https://doi.org/10.1016/j.vetpar.2003.07.028

Tattersall, F.H., Nowell, F., Smith, R.H., 1994. A review of the endoparasites of wild House Mice
Mus domesticus. Mamm. Rev. 24, 61–71. https://doi.org/10.1111/j.1365-2907.1994.tb00135.x

Vandenbergh, J.G., 2013. Use of House Mice in Biomedical Research. ILAR J. 41, 133–135.
https://doi.org/10.1093/ilar.41.3.133

Vrba, V., Blake, D.P., Poplstein, M., 2010. Quantitative real-time PCR assays for detection and
quantification of all seven Eimeria species that infect the chicken. Vet. Parasitol. 174, 183–
190. https://doi.org/10.1016/j.vetpar.2010.09.006

Yang, H., Bell, T.A., Churchill, G.A., Pardo-Manuel De Villena, F., 2007. On the subspecific origin of
the laboratory mouse. Nat. Genet. 39, 1100–1107. https://doi.org/10.1038/ng2087

Yang, R., Brice, B., Ryan, U., 2015. Morphological and molecular characterization of Eimeria
haematodi, coccidian parasite (Apicomplexa: Eimeriidae) in a rainbow lorikeet (Trichoglossus
haematodus). Exp. Parasitol. 153, 123–128. https://doi.org/10.1016/j.exppara.2015.03.005

Zhao, X., Duszynski, D.W., 2001. Phylogenetic relationships among rodent Eimeria species

determined by plastid ORF470 and nuclear 18S rDNA sequences. Int. J. Parasitol. 31, 715–
719. https://doi.org/10.1016/S0020-7519(01)00136-9

662 **Tables**

Table 1. Morphological and morphometrical characteristics from *Eimeria* wild-derived isolates and reference morphotypes[§].

Species	Oocysts							
	Shape	Wall	Micropyle	Polar granule	Residuum	Length (µm)	Width (µm)	L/W ratio
E. falciformis	Spherical/Ellipsoidal	Smooth	Absent	Present	Absent	18.62 (15.29 – 20.76)	15.92 (13.89 – 17.6)	1.17 (1.00 – 1.36)
E. falciformis	Ovoid/Spherical	Smooth	Absent	Present	Absent/Present	21 (15 – 26)	18 (13 – 24)	1.17 (1.1 – 1.2)
E. ferrisi	Spherical/Ellipsoidal	Smooth	Absent	Present*	Absent	17.32 (13.59 – 21.47)	14 (10.97 – 17.44)	1.23 (1.02 – 1.56)
E. ferrisi	Ellipsoidal/Subspherical	Smooth	Absent	Present	Absent	17 (12 – 22)	14 (11 – 18)	1.22 (1.0 – 1.6)
E. vermiformis	Spherical/Ellipsoidal	Smooth	Absent	Present*	Absent	20.02 (16.22 – 22.8)	15.57 (12.37 – 18.44)	1.29 (0.99 – 1.45)
E. vermiformis	Spherical/Ellipsoidal	NA	Absent	Present	Absent	23.1 (18 – 26)	18.4 (15 – 21)	1.25 (1.1 – 1.4)

Species	Sporocysts					Tissue localization	Reference
	Residuum	Refractile body	Stieda body	Length (µm)	Width (µm)		
E. falciformis	Present	Present	Absent	8.88 (7.03 – 10.30)	5.52 (4.16 – 6.93)	Cecum	This work
E. falciformis	Present	Present	Present	11 (10 – 12)	7 (6 – 8)	Ileum,**cecum**and colon	Eimer, 1870; Haberkorn, 1970
E. ferrisi	Present	Present	Present	7.75 (5.17 – 11.47)	5.03 (2.88 – 7.73)	Cecum	This work
E. ferrisi	Absent/Present	Present	Present	10.5 (10 – 11)	5.5 (5 – 6)	**Cecum**and colon	Levine & Ivens, 1965; Ankrom et al., 1975
E. vermiformis	Present	Present	Present	8.35 (6.18 – 11.29)	5.43 (4.19 – 7.00)	Cecum and ileum	This work
E. vermiformis	Present	Present	Present	12.8 (11 – 14)	7.9 (6 – 10)	Jejunum,**ileum**and cecum	Ernst et al., 1971

§ Measurements are means in micrometers with ranges in parenthesis
*Observed in more than 80% of the oocysts

Table 2. Statistical models used to analyse primer preference to different *Eimeria* species.

Predictors	Model 1 (Cocci_COI)	Model 2 (Cocci_COI + Eim_COI)	Model 3 (18S_EF/R)
Intercept	-5.80 ***	-1.55	-1.40
	(1.11)	(1.27)	(0.87)
Ap5	3.50 ***	3.99 ***	0.78
	(0.59)	(0.60)	(0.62)
Flotation	1.64 **	2.15 ***	1.59 **
	(0.52)	(0.54)	(0.51)
Identification *E. ferrisi* (other marker)	5.10 ***	0.28	0.32
	(1.14)	(1.35)	(0.63)
Identification *E. vermiformis* (other marker)	17.23	11.98	1.71
	(1385.38)	(1385.38)	(1.33)
No identification (other marker)	0.97	-3.57 **	-3.44 ***
	(0.92)	(1.23)	(0.87)
Identification *Eimeria* spp. (COI)			-14.80
			(1455.40)
N	378	378	378
AIC	128.09	119.18	143.39
BIC	151.70	142.79	170.93
Pseudo R2	0.77	0.82	0.70

The upper number represents the estimate and numbers in brackets represent standard error for each predictor.

Intercept is *Eimeria falciformis* identification with other marker

Other marker refers to 18S based identification for COI models or vice versa

N (Total number of samples), AIC (Akaike information criterion), BIC (Bayesian information criterion).

*** p < 0.001; ** p < 0.01; * p < 0.05.

Figure captions

Figure 1. Geographical localization of house mice (*Mus musculus*) collected for this study and comparison of diagnostic methods for *Eimeria*. A) Localization from the 378 mice included in the present study, colors indicate the *Eimeria* species identified for each. B) Venn diagram showing the overlap between detection methods and successful genotyping identification of the isolates.

Figure 2. Phylogenetic trees based on 18S rRNA and COI sequences. Sequences of 18S A) and COI B) were used to infer the molecular identification of wild-derived isolates of *Eimeria*. In both phylogenies, our isolates clustered in three groups one close to *E. falciformis* (red), other close to *E. ferrisi* (green) and finally one to *E. vermiformis* (yellow). Numbers in the branches represent the Bayesian posterior probability and the non-parametric bootstrap value. In bold are indicate the reference sequences for each species. CE and IL make reference to sequence derived from cecum or ileum tissue DNA, respectively.

Figure 3. Morphological and morphometrical characteristics of *Eimeria* oocyst isolated from *Mus musculus*. a) Photomicrographs at 1000x amplification of *Eimeria* oocyst from the three species isolated from *Mus musculus* (red = *E. falciformis*; green = *E. ferrisi* and yellow = *E. vermiformis*). Length/Width ratio from b) oocyst and c) sporocysts corresponding to each species (*E. falciformis* n= 31; *E. ferrisi* n= 127 and *E. vermiformis* n= 35). Mean +/- 95% Confidence Interval is plotted. * Represent significant difference (Tukey HSD, $p<0.05$).

Figure 4. qPCR detection of intracellular stages of *Eimeria* in cecum and ileum from *Mus musculus*. -Delta Ct value ($Ct_{Eimeria}$ - Ct_{Mouse}) from each tissue for 164 mice are plotted on the graph. The dotted line indicate the threshold of -6, values above the line are considered positive for the corresponding tissue. Circles represent negative samples, triangles indicate samples with *Eimeria* species identification and colors correspond to the *Eimeria* species identified in those samples.

Supplementary data S1. Geographical coordinates for collected mice in Brandenburg,

Germany.

Supplementary data S2. Accession numbers for 18S and COI sequences from *Eimeria* spp.

wild isolates.

Supplementary data S3. Maximum-Likelihood phylogenetic tree inferred for 18S/COI

sequences supermatrix.

Supplementary data S4. Bayesian phylogenetic tree inferred for 18S/COI sequences

supermatrix.

Supplementary data S5. Oocysts measurements from wild-derive isolates of *Eimeria*

falciformis*, *E. ferrisi* and *E. vermiformis

Supplementary data S6. Ct values from the qPCR detection of *Eimeria* in tissue.

A)

B)

ΔCt Ileum

ΔCt Cecum

Eimeria **identification**

△ Double △ *E. falciformis* △ *E. ferrisi*
△ *E. vermiformis* ⬤ Negative