Diversification in evolutionary arenas – assessment and synthesis

Nicolai M. Nürk*[1], H. Peter Linder*[2], Renske E. Onstein[3], Matthew J. Larcombe[4], Colin E. Hughes[2], Laura Piñeiro Fernández[2,5], Philipp M. Schlüter[5], Luis Valente[6,7], Carl Beierkuhnlein[8], Vanessa Cutts[9], Michael J. Donoghue[10], Erika J. Edwards[10], Richard Field[9], Suzette G.A. Flantua[11], Steven I. Higgins[12], Anke Jentsch[13], Sigrid Liede-Schumann[1], Michael D. Pirie[14,15]

[1] Department of Plant Systematics, Bayreuth Center of Ecology and Environmental Research (BayCEER), University of Bayreuth, Universitätsstraße 30, 95447 Bayreuth, Germany. nuerk@uni-bayreuth.de, orcid.org/0000-0002-0471-644X; sigrid.liede@uni-bayreuth.de, orcid.org/0000-0003-2707-0335.

[2] Department of Systematic & Evolutionary Botany, University of Zurich, Zollikerstrasse 107, 8008 Zurich, Switzerland. peter.linder@systbot.uzh.ch, orcid.org/0000-0002-1373-2708; colin.hughes@systbot.uzh.ch, orcid.org/0000-0002-9701-0699; laura.pineiro@systbot.uzh.ch, orcid.org/0000-0002-8617-6490.

[3] German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Deutscher Platz 5e, 04103 Leipzig, Germany. onsteinre@gmail.com, orcid.org/0000-0002-2295-3510.

[4] Department of Botany, University of Otago, PO Box 56, Dunedin New Zealand 9054. matt.larcombe@otago.ac.nz, orcid.org/0000-0003-1632-9591.

[5]† Department of Botany, University of Hohenheim, Garbenstraße 30, 70599 Stuttgart, Germany. philipp.schlueter@uni-hohenheim.de, orcid.org/0000-0002-6057-0908.

[6]† Naturalis Biodiversity Center, Understanding Evolution Group, 2332 AA Leiden, the Netherlands. luis.valente@naturalis.nl, orcid.org/0000-0003-4247-8785.

[7] Groningen Institute for Evolutionary Life Sciences, University of Groningen, P.O. Box 11103, Groningen 9700 CC, the Netherlands

[8] Department of Biogeography, Bayreuth Center of Ecology and Environmental Research (BayCEER), University of Bayreuth, Universitätsstraße. 30, 95440 Bayreuth, Germany. carl.beierkuhnlein@uni-bayreuth.de.

[9] School of Geography, University of Nottingham, Nottingham, NG7 2RD, United Kingdom. vanessa.cutts@nottingham.ac.uk, orcid.org/0000-0002-4986-2934. richard.field@nottingham.ac.uk, orcid.org/0000-0003-2613-2688

[10] Department of Ecology and Evolutionary Biology, Yale University, PO Box 208105, New Haven CT 06511. michael.donoghue@yale.edu; erika.edwards@yale.edu.

[11] Department of Biological Sciences, University of Bergen, PO Box 7803, 5020, Bergen, Norway. s.g.a.flantua@gmail.com, orcid.org/0000-0001-6526-3037.

[12] Plant Ecology, University of Bayreuth, Universitätsstraße 30, 95447 Bayreuth, Germany. steven.higgins@uni-bayreuth.de, orcid.org/ 0000-0001-5695-9665.

[13] Department of Disturbance Ecology, Bayreuth Center of Ecology and Environmental Research (BayCEER), University of Bayreuth, Universitätsstraße. 30, 95440 Bayreuth, Germany. anke.jentsch@uni-bayreuth.de, orcid.org/0000-0002-2345-8300.

[14] Johannes Gutenberg-Universität, Saarstraße 21, 55099 Mainz, Germany. pirie@uni-mainz.de.

[15]† University Museum, University of Bergen, Postboks 7800, N-5020 Bergen, Norway. michael.pirie@uib.no, orcid.org/0000-0003-0403-4470

* contributed equally; nuerk@uni-bayreuth.de (+49 921 55 2466), peter.linder@systbot.uzh.ch (+41 44 634 84 10).

Abstract

Understanding how and why rates of evolutionary diversification vary is a key issue in evolutionary biology, ecology, and biogeography, and the metaphorical concepts of adaptive radiation and evolutionary stasis describe two opposing aspects causing variation in diversification rates. Here we review the central concepts in the evolutionary diversification literature and synthesize these into a simple, general framework for studying rates of diversification and quantifying their underlying dynamics, which can be applied across clades and regions and across spatial and temporal scales. Our framework describes the diversification rate (d) as a function of the abiotic environment (a), the biotic environment (b) and clade-specific phenotypes or traits (c); thus $d \sim a, b, c$. We refer to the four components (a–d) and their interactions collectively as the 'Evolutionary Arena'. We outline analytical approaches to this framework and present a case study on conifers, for which we parameterise the general model. We also discuss three conceptual examples: the *Lupinus* radiation in the Andes in the context of emerging ecological opportunity and fluctuating connectivity due to climatic oscillations; oceanic island radiations in the context of island formation and erosion; and biotically driven radiations of the Mediterranean orchid genus *Ophrys*. The results of the conifer case study are consistent with the long-standing scenario that low competition and high rates of niche evolution promote diversification. The conceptual examples illustrate how using the synthetic Evolutionary Arena framework helps to identify and structure future directions for research on evolutionary radiations. In this way, the Evolutionary Arena framework promotes a more general understanding of variation in evolutionary rates by making quantitative results comparable between case studies, thereby allowing new syntheses of evolutionary and ecological processes to emerge.

Key words

Introduction

Biologists have long been fascinated by the circumstances under which species diversification and trait

disparification rates – evolutionary radiations – are accelerated. Studies in recent decades on

evolutionary **radiations** (words in bold are in the Glossary) have produced a proliferation of terminology

and new statistical approaches. These developments in (macro-) evolution are largely based on the

adaptive radiation paradigm (Osborn, 1902; Simpson, 1944, 1953; Schluter, 2000), a metaphorical

concept describing the evolution of a multitude of ecological forms from a single common ancestor. The

paradigm, however, complicates quantitative comparisons of the trajectories and correlates of

diversification between evolutionary lineages (species, clades) and among geographical regions, and

does not address the circumstances under which evolutionary stasis or decline may occur. Here, we

build on current theoretical foundations and propose a conceptual framework for the integrative study

of shifts and stasis in diversification rates. It is not our aim to thoroughly review the literature on

evolutionary radiations; rather, we provide an overview of recent developments, and integrate these

into a framework that can in principle be quantified in all systems, from cellular to global spatial scales

and spanning ecological to evolutionary time frames.

A short history of diversification theory

Darwin, in sharp contrast to early-nineteenth-century dogma, envisioned evolution to be gradual, with

small changes accumulating from generation to generation, eventually leading to species divergence

(Orr, 2005). This gradualist view was soon challenged and seemingly contradicted by the fossil record,

leading to the appreciation that rates of divergent evolution are uneven through time and among

clades, sometimes generating species and ecomorphological diversity in evolutionary radiations (Mayr,

1954; Stanley, 1979), while at other times demonstrating long-term stasis (Eldredge & Gould, 1972;

Gould & Eldredge, 1977; Flegr, 2010) or decline in diversity (Benton, 1995; Rohde & Muller, 2005).

The development of phylogenetic theory (Hennig, 1950, 1965) followed by the generation of

massive DNA sequence datasets, increased computing power, and the proliferation of analytical

methods (i.a., maximum likelihood, Felsenstein, 1973, 1981; Bayesian inference, Huelsenbeck, Ronquist,

Nielsen, & Bollback, 2001; Bayesian molecular dating, Drummond, Ho, Phillips, & Rambaut, 2006;

multispecies coalescence, Degnan & Rosenberg, 2006; Edwards, 2009) have resulted in a vast

accumulation of progressively higher quality phylogenies Maddison, 1997; e.g., Brassac & Blattner,

2015), leading to recognition of monophyletic groups and estimates of the temporal dynamics of

evolutionary radiations (Nee, May, & Harvey, 1994; Magallón & Sanderson, 2001; Alfaro et al., 2009;

Stadler, 2011; Morlon, 2014; Rabosky, 2014). These have revealed orders-of-magnitude differences in

clade diversification rates, exemplified by *Amborella trichopoda*, an understory shrub endemic to New

Caledonia, which is the only species of an angiosperm clade that is sister to, and therefore just as old as,

the clade that contains all remaining ca. 400,000 species of flowering plants (Albert et al., 2013). Placing

such salient diversification rate differences into a striking temporal context, Salzburger (2018, p. 705)

recently noted that within "the time span that it took for 14 species of Darwin's finches to evolve on the

Galapagos archipelago […], about 1,000 cichlid species evolved in Lake Malawi alone".

Drivers of evolutionary radiations

The importance of **traits** and environments for understanding the mechanisms underlying evolutionary

radiations was emphasized by Simpson (1944), and he postulated that most radiations are underpinned

by **adaptation**. His adaptive radiation model envisioned diversification to take place in **adaptive zones**. A

species may enter new adaptive zones by the evolution of new traits, climate change, or the formation

of novel landscapes and disturbance regimes, such as newly emerged volcanic islands or human-induced

night-light environments around the globe (Simpson, 1953; Erwin, 1992). The adaptive zone is a

metaphor for the ways in which evolutionary innovations interact with environmental factors to

modulate species **diversification** and trait **disparification** rates (de Vladar, Santos, & Szathmary, 2017;

Olson, Arroyo-Santos, & Vergara-Silva, 2019), and Simpson's concept of adaptive radiation underpinned

and inspired a highly productive period of research on evolutionary radiations (e.g., Fryer, 1969; Losos,

1994; Sanderson & Donoghue, 1994; Baldwin, 1997; Hughes & Eastwood, 2006; Losos & Ricklefs, 2009;

Wagner, Harmon, & Seehausen, 2012; Cooney et al., 2017; Marques, Meier, & Seehausen, 2019).

The effects of traits on species diversification (the interplay of speciation and extinction rates),

combined with the temporal sequence of geographic movement and environmental change, estimated

across a phylogenetic tree, has led to the recognition of **key innovations** (Miller, 1949; Van Valen, 1971;

Liem, 1973; Sanderson & Donoghue, 1994; Heard & Hauser, 1995; Hunter, 1998). Key innovations are

exemplified by freezing tolerance in Antarctic fishes (Portner, 2002) and herbaceous life-history

strategies for occupying seasonally freezing environments by flowering plants (Zanne et al., 2014). In

addition, phylogenetic comparative studies (Felsenstein, 1985; Harmon, 2018) have revealed the

importance of **key events**, such as mass extinctions (e.g., of dinosaurs), climate change (e.g., late

Miocene aridification), and orogeny (e.g., of the Andes and the New Zealand alps). Including

evolutionary changes in genomic structure has led to the recognition that the connections between key

innovations, key events, and diversification rate shifts can be complex (Erwin, 2001, 2017), for example,

in context of hybridization and whole-genome duplications in flowering plants (Tank et al., 2015; Landis

et al., 2018; Naciri & Linder, 2020) or African Rift Lake cichlids (Meier et al., 2017; Irisarri et al., 2018).

Generally, however, it is the interaction between variable **intrinsic** (e.g., genome duplication) and

extrinsic (e.g., climate change) factors that is thought to modulate diversification rates (for a review on

context-dependent diversification see Donoghue & Sanderson, 2015; on the interplay of dispersal and

biome shifts see Donoghue & Edwards, 2014).

The interaction between intrinsic (lineage-specific) and extrinsic (environmental) factors provides

the **ecological opportunity** for adaptive radiations to occur (Erwin, 2015). Simpson (1953) summarised

such opportunities in terms of three factors: (i) physical access to an environment, resulting from

dispersal, or from the change of geo-ecological conditions in the region where a lineage already occurs;

(ii) lack of effective competition in the environment, because no suitably adapted lineages already occur

there; (iii) genetic capacity and adaptability of a lineage, which can be manifested in the evolution of key

innovations, or more generally, in the ability to more readily explore the character space of certain trait

innovations (Nürk, Atchison, & Hughes, 2019). All three conditions have to be met for successful

adaptive radiation to start (Donoghue & Edwards, 2014; Stroud & Losos, 2016).

The evolution of diversity requires the evolution of reproductive isolation, which can be promoted

by geographic fragmentation. Geographic isolation can result in stochastic divergence, i.e., by intensified

genetic drift in smaller populations (Kimura, 1968; Duret, 2002), eventually leading to allopatric

speciation. Repeated allopatric speciation can result in "non-adaptive" radiation (Gittenberger, 1991;

Comes, Tribsch, & Bittkau, 2008; Verboom, Bergh, Haiden, Hoffmann, & Britton, 2015). Such (mainly)

isolation-driven processes are contrasted to ecological speciation, which is driven by divergent selection

pressures from the environment, implying that there will be (some) adaptation. Ecological speciation

can result in repeated evolution of phenotypes and trait–environment interactions in adaptive

radiations (trait utility; Schluter, 2000). However, in most radiations both ecological adaptation (natural

selection) and geographic isolation (intensified genetic drift) are involved (Gittenberger, 2004; Brawand

et al., 2014), although their relative contributions to diversification may vary between study systems

(Rundell & Price, 2009; Czekanski-Moir & Rundell, 2019; Naciri & Linder, 2020).

Variation in the relative contributions of non-adaptive and adaptive processes to diversification

was encapsulated by Simpson (1953) in his distinction between 'access to an environment' and 'lack of

competition in that environment'. In ecological niche theory, it is well appreciated that species live in

environments that can be described by abiotic and biotic factors (for a review on the various aspects of

niche concepts see McInerny & Etienne, 2012; McInerny, Etienne, & Higgins, 2012). Both abiotic and

biotic factors can influence diversification rates (Holt, 2009) and may have varying or even opposite

effects (Bailey, Dettman, Rainey, & Kassen, 2013) on speciation probability and extinction risk (Ezard,

Aze, Pearson, & Purvis, 2011). Species interactions can drive diversification (Brodersen, Post, &

Seehausen, 2018; Gavini, Ezcurra, & Aizen, 2019), or can have negative effects on species diversity, for

example under competition for limited resources (Rosenblum et al., 2012; Harpole et al., 2016). On the

other hand, heterogeneous abiotic conditions appear to be generally associated with higher species

diversity (Rainey & Travisano, 1998; see also Erwin, 2001) potentially due to higher carrying capacities of

larger and more heterogeneous areas (Field et al., 2009; Wagner, Harmon, & Seehausen, 2014; Storch &

Okie, 2019), greater topographic complexity (Badgley et al., 2017; Sundaram et al., 2019), more climatic

variability (Weigelt, Steinbauer, Cabral, & Kreft, 2016; Flantua, O'Dea, Onstein, Giraldo, & Hooghiemstra,

2019), or more complementary disturbance dynamics concordant or discordant with life histories

(Jentsch & White, 2019). Indeed, abiotic and biotic factors and their effects on diversification are firmly

established in research on evolutionary radiations (Ezard et al., 2011; Aguilée, Gascuel, Lambert, &

Ferriere, 2018; Condamine, Romieu, & Guinot, 2019) and both factors are part of the extrinsic

environment in which evolution of a lineage takes place.

In general, it is not the particular sequence of trait evolution or access to a novel environment

that **triggers** a radiation, but the establishment/evolution of a complementary state – either the

establishment of an environment that fits a pre-evolved trait or an **exaptation**, or the evolution of the

trait that is an adaptation to a pre-existing environment (Kozak & Wiens, 2010; Wagner et al., 2012;

Bouchenak-Khelladi, Onstein, Xing, Schwery, & Linder, 2015; Nürk, Michling, & Linder, 2018). Exploring

the theoretical arguments that underpin such context-dependent radiations, Donoghue and Sanderson

(2015) coined the terms **synnovation** for interacting combinations of (several) innovative traits, and

confluence to describe sequential combinations of a set of traits and events along the stem lineages of

radiating clades. The idea that evolutionary radiations are the product of synnovations and confluences

of multiple intrinsic and extrinsic factors has gained momentum (Arakaki et al., 2011; Wagner et al.,

2012; Guerrero, Rosas, Arroyo, & Wiens, 2013; Seehausen, 2015; Linder & Bouchenak-Khelladi, 2017;

Harmon et al., 2019; Nürk, Atchison, et al., 2019).

The Evolutionary Arena framework

Description of the framework

Here we synthesize these insights into the drivers of evolutionary radiations – the context-dependent

interplay between clade-specific intrinsic and extrinsic biotic and abiotic factors – into a simple

framework, which we call the 'Evolutionary Arena' (EvA). In EvA the diversification (or disparification)

rate of a focal lineage is a function of three components into which all macroevolution-relevant

processes can be grouped and parameterised:

$d \sim a, b, c$

where d = **d**iversification or **d**isparification rate, a = **a**biotic environment, b = **b**iotic environment, and c =

clade-specific phenotypes or traits (Fig. 1).

Figure 1. The Evolutionary Arena framework. The four components of the Evolutionary Arena (boxes) are
illustrated with interactions (arrows) among the components, with those influencing diversification rates in

black (larger arrows). We refer to the environment (abiotic [*a*] and the biotic [*b*] components; blue boxes) in combination with clade traits (*c*) and diversification / disparification rate (*d*) of an evolutionary lineage (green boxes) as the Evolutionary Arena. This framework is potentially dynamic, because interactions among the components allow for feedback, together shaping evolution. Note that, although all components can affect each other (positively or negatively; indicated by small and/or grey arrows), we focus here on the dependence of diversification on environmental (extrinsic biotic and abiotic) and clade-specific (intrinsic trait) factors.

In more detail, these four components are:

d: diversification or *disparification* is the rate of evolution in the broadest sense and depends on the values of the *a*, *b*, and *c* components (note, *d* might influence the other components as well; Fig. 1). Here, *d* can be expressed by the rate of change in taxonomic diversity (number of species), interspecific morphological/phenotypic disparity, DNA nucleotide diversity and genetic differentiation (e.g., functional variation of expressed genes or metabolites), physiological diversity (e.g., photosynthetic modes), or niche diversity (e.g., diversity of ecological niches occupied) and differentiation (e.g., rate of ecological expansion). This list is not exhaustive, and which expression of *d* is used depends on the question being investigated. The instantaneous rate of diversification, defined as speciation minus extinction per unit time (Nee et al., 1994), is the simplest expression of *d* and can be directly inferred from a dated species-level phylogeny (Magallón & Sanderson, 2001). Disparification can be measured, for example, by (relative) evolutionary rate estimates (Butler & King, 2004), the change in trait variance in a clade through time (Rolshausen, Davies, & Hendry, 2018), or by transition rates between discrete character states (Huelsenbeck, Nielsen, & Bollback, 2003). The diversification rate may be positive, resulting in an increase in diversity, or negative, resulting in a decrease in diversity.

a: abiotic environment incorporates abiotic factors, such as climate, soil, habitat variables, or fragmentation of the species range. It can also include biotic elements, particularly with respect to their physical characteristics, such as vegetation types (which, as classes, cannot evolve). Component *a* can be measured as absolute values, e.g., area or niche space, or as the variance in these across space and time, e.g., variance in mean annual precipitation, in number of vegetation types or soil types, physiographic heterogeneity, and the functional and structural connectivity. Ideally, the abiotic environment is described by processes that generate this environment, such as erosion or orogeny, or patterns of change, such as climate or vegetation change.

b: biotic environment captures the interactions of the focal lineage with all other species (including

species both within and outside the clade and also including trophic interactions). The interaction(s) can

be, for example, mutualistic or commensalistic (e.g. pollinators or dispersers, interspecific facilitation, or

mycorrhiza), antagonistic (e.g., herbivores, diseases, interspecific competition, or hosts for parasites), or

genetic (e.g., hybridization/introgression and horizontal gene transfer). Note, however, that the *capacity*

to hybridize may be treated as a trait, and so categorised under *c*. These biotic interactions can also be

indirect, if seen as part of the extended phenotype of the focal lineage (e.g., habitat modification such as

niche construction, or grasslands increasing fire frequency, or stoichiometric needs of organisms

modifying available resources; Jentsch & White, 2019). There is a rich theory surrounding biotic

interactions (e.g., the macro-evolutionary Red Queen hypothesis [Benton, 2009, see also Van Valen,

1973], niche construction [Laland, Matthews, & Feldman, 2016]) suggesting they can have a powerful

influence on diversification rates. However, the biotic environment is complex; quantifying and including

it in comparative analyses is a challenging task (Harmon et al., 2019). Indeed, recent studies indicate

that inter-lineage competition (e.g., Silvestro, Antonelli, Salamin, & Quental, 2015; Pires, Silvestro, &

Quental, 2017) or interactions with possible dispersal agents (Onstein et al., 2018) may modulate

diversification rates.

c: clade-specific traits include the phenotypic characteristics of the focal species or lineage. Included

here are, among others, physiological characteristics (e.g., variation in photosynthetic mode),

anatomical or morphological traits, life history strategies, pollination strategies and dispersal modes.

Clade-specific traits are part of the phenotype and can be labile or phylogenetically conserved. This is

illustrated by the remarkable floral variation in the orchid genus *Disa* (Johnson, Linder, & Steiner, 1998),

and the impact of a lack of floral morphology variation on diversification in the oil-bee pollinated

Malpighiaceae (Davis et al., 2014). The phenotype (or the extended phenotype) is a manifestation, or

function, of the genome (via the genotype–phenotype map, inasmuch as this is independent of the

environment), or genome diversity (e.g., structural variation such as ploidy level variation, variation in

genome size, or DNA nucleotide sequence variation) within the focal group. Therefore, this parameter

should ultimately be genetically measurable.

The formulation of the EvA framework is general because it is simple and all-encompassing. The key

challenge faced in studies addressing evolutionary radiations is to disentangle the effects of the

different components at different moments in time. The ultimate in understanding radiations and

evolutionary stasis would be the joint estimation of all components at all times, but we cannot analyse

an infinite number of variables. On the other hand, studies sometimes assign an increase or decrease of

diversification to particular factors that happen to have been investigated and quantified, disregarding

the possible effects of other factors that may be driving or constraining diversification. We argue that

the complete EvA, as expressed by the abiotic and biotic environment, as well as the traits of organisms,

should be considered when attempting to explain variation in diversification rates. Not only does this

ensure that all relevant factors are taken into account, but the same set of components are considered

in all analyses. In this way, the EvA framework can provide new insights by comparing diversification

between clades directly.

To make EvA operational requires parameterising it appropriately, which means making it more

specific and detailed. Here, we describe four simple extensions to illustrate how the EvA framework can

be enriched by more properties to provide insights into different hypotheses in evolutionary

diversification. We end this section by outlining general analytical approaches and possibilities for null

hypothesis formulation.

1) *Direction of effects*: The three components *a*, *b*, and *c* can have a significant positive (+) or

negative (-) effect on *d*, thus causing the diversification rate to increase or decrease, or even show a

false absence of change as the summed end result of the three. This process of 'nullification', or less

increase or decrease than expected, is sometimes ignored and interpreted as a lack of power by the

factors influencing diversification. Statistical approaches comparing different systems could provide

insights in cases in which diversification rates are higher or lower than expected.

2) *Complex conditions*: The framework can be expanded to include any set of variables per

component. For example, the abiotic environment could be described by climatic factors such as mean

annual precipitation, temperature, and by disturbance factors such as fire frequency. The set of

variables selected depends on the hypothesis being tested. If, for example, the hypothesis is that *d* is

related to the total variance in the abiotic environment *a*, then $a=\sum_{i=1}^{n} a_i$, where a_i is the i^{th} variable of *a*

(up to the last variable a_n) given as a measure of variance. Thus, the abiotic environment *a* (as well as

the other components in EvA) can be decomposed into diverse measures.

3) *Rate shifts*: Explanations for shifts in diversification rates can be sought by testing for changes

in the EvA component values. This can be done by including initial values at time *t* (ancestral) and the

values after a time interval Δ*t* (derived), for example, at time *t*+Δ*t* after an event: $\Delta d = (a, b, c)_{t+\Delta t} -$

$(a, b, c)_t$ or simply $\Delta d \sim \Delta a, \Delta b, \Delta c$. If we, for example, hypothesise that geographic movement to a new

region is a key event, the condition of *a* varies between *t* and *t*+Δ*t*.

4) *Interaction of components*. We can incorporate interactions between components, such as *ac*,

bc, or *abc*: $d \sim a, b, c, ab, ac, bc, abc$. This allows us to analyse context-dependence: whether the single

components, or the interactions among them modulate the diversification rate, i.e., driving or

constraining the evolution of diversity. Such interactions can be exemplified by ecosystem engineers

modifying disturbance regimes (Jentsch & White, 2019), such as impacts of grass invasion on forests

increasing fire frequency and ultimately transforming the environment (Bond, Woodward, & Midgley,

2005; Beerling & Osborne, 2006). Similar interactions (or transformations) have been proposed for

changes in fruit size (component *c*) as consequences of mega-herbivore extinctions (component *b*) and

climate change (component *a*; Onstein et al., 2018). A further dynamic factor is provided by interactions

between the diversification rate (*d*) and the predictor variables (*a, b, c*). Such feedback mechanisms can

broadly be summarised by the concept of niche construction (Laland et al., 2016), and are exemplified

by the 'Viking Syndrome' described in reference to grasses (Linder, Lehmann, Archibald, Osborne, &

Richardson, 2018). The latter proposes that global grass success (in species richness, environmental

range, ecological dominance and geographical distribution) is due to the high invasiveness of grasses,

which results from their high rate of dispersal, effective establishment, ecological flexibility and

disturbance tolerance (all component *c*), and ability to transform environments by increasing the

frequency of fire (component *a*) and the density of grazers (component *b*). These scenarios describe

feedback systems where through increased diversity and dominance, the diversifying clade increasingly

modifies the abiotic and biotic environment.

In this way, the EvA framework can facilitate the direct testing of competing hypotheses about the

diversification of a group, using standard model selection approaches (e.g., likelihood ratio tests for

nested models, AIC or Bayes factors; Burnham & Anderson, 2002). Phylogenetic pseudoreplication

(Maddison & FitzJohn, 2015), which describes the non-independence of, or the autocorrelation among,

species' traits due to shared ancestry, is a basic property of comparative analyses (Felsenstein, 1985).

Phylogenetically independent contrasts (PIC; Felsenstein, 1985) and phylogenetic generalised least

squares (PGLS; Grafen, 1989; Martins & Hansen, 1997) are methods for analysing comparative

phylogenetic data by accounting for the covariances between traits resulting from shared phylogenetic

history. These methods may be generally useful for exploring the relationships between components in

the EvA framework. Although both methods can be thought of as "analogous to data transformations

made to better approximate the assumptions of standard statistical tests" (Huey, Garland, & Turelli,

2019, p 762), they, as well as most other phylogenetic comparative methods (but see Rolshausen et al.,

2018), implicitly assume a specific process underlying character state change along the evolutionary

lineage (i.e., a model of character evolution such as Brownian motion; Boucher, Demery, Conti, Harmon,

& Uyeda, 2018). The appropriate data transformation model in relation to the hypothesis being tested is

a non-trivial question in comparative analyses (Uyeda, Zenil-Ferguson, & Pennell, 2018); graphical

models depicting hypothesized causal links (Höhna et al., 2014) can help here. On the other hand, the

case of a singular, unreplicated event in the evolutionary history of a lineage challenges the statistical

power of comparative phylogenetic methods (Maddison & FitzJohn, 2015). An approach to overcome the

this limitation might be the combined application of hypothesis testing and exploratory methods

('phylogenetic natural history') as outlined by Uyeda et al. (2018).

Associations between diversification rates and the other components in the EvA framework can

be tested using state-dependent speciation and extinction (SSE) models (such as BiSSE): a birth–death

process where the diversification rates depend on the state of an evolving (binary) character (Maddison,

Midford, & Otto, 2007), given a phylogeny and trait data. For complex components, the multistate SSE

(MuSSE) model can be applied to accommodate several qualitative character states, or multiple (binary

state) characters following Pagel (1994) for state recoding (FitzJohn, 2012). Although SSE model

extensions for quantitative characters (QuaSSE; (FitzJohn, 2010) and geographic range evolution

(GeoSSE; Goldberg, Lancaster, & Ree, 2011) (to name just a few; for a critical introduction to SSE models

see O'Meara & Beaulieu, 2016) have been developed, in the SSE model-family none is currently available

for the likelihood calculation of a model considering both quantitative and qualitative variables (but see

Felsenstein, 2012; Revell, 2014). Nor do current SSE models allow the analysis of interactions, and so

context-dependent radiations. This situation could be common under the EvA framework and

represents a priority for method development.

Comparing a biologically meaningful and appropriately complex null hypothesis to the goodness-

of-fit of alternative (H1) models is essential for detecting whether a character state-dependent model

can explain more of the observed variation than could be expected under random diversification rates

(Caetano, O'Meara, & Beaulieu, 2018). It has, for example, been shown for SSE models that the variation

in the diversification rate observed in a phylogenetic tree is not necessarily explained by the focal factor

(character) under study (Rabosky & Goldberg, 2015). False-positives can potentially result because the

null hypotheses did not account for the possibility that diversification rates can be "independent of the

character but not constant through time" (Harmon, 2018, p 215). Hidden state model (HSM) approaches

(Marazzi et al., 2012; Beaulieu, O'Meara, & Donoghue, 2013; Beaulieu & O'Meara, 2016; Caetano et al.,

2018), which incorporate unobserved ('hidden') factors as model parameters equivalent to observed

ones, offer a solution to this problem. Comparing goodness-of-fit between 'hidden state' null models

and those representing the focal factor(s) provides appropriately complex null hypotheses that can be

used for testing differently parameterised EvA models, and thereby allows identification of "the

meaningful impact of [the] suspected 'driver[s]' of diversification" (Caetano et al., 2018, p 2308).

Advantages of the Evolutionary Arena framework

The Evolutionary Arena framework does not contribute any new concepts or terms but is built on the

concepts developed over the past few decades, reviewed above (section "Drivers of evolutionary

radiations"). However, what is still lacking is, as noted by Donoghue and Sanderson (2015), a single,

simple formula with which to develop convincing hypotheses of the drivers of evolutionary rate

changes. This we attempt to provide with EvA. The basic four components – diversification /

disparification, clade-specific intrinsic traits, extrinsic abiotic and biotic factors – and their interactions

can be compared between systems to gain more general insights into the factors that underpin

evolutionary diversification. Considering all four components together in a single framework fosters a

holistic approach. EvA consequently incorporates the full complexity of triggers, synnovations, and

confluences associated with evolutionary radiations (Bouchenak-Khelladi et al., 2015; Donoghue &

Sanderson, 2015). This facilitates comparative analyses of evolutionary radiations, or evolutionary stasis

and decline, using phylogenetic comparative methods. This is possible because d can be positive or

negative/smaller, so the correlates (e.g., a x c in EvA) of diversification increase or decrease (e.g.,

density-dependent slowdowns) can be sought. EvA does not present any new analytical methods,

analyses within this framework can be done using existing packages and software (it may also indicate

priorities for method development). Particularly important is the central notion that no single factor is a

sufficient explanation for an evolutionary rate change, but that the interaction between external and

internal factors results in shifts in diversification and/or disparification rates (Givnish, 2015). Overall,

there are three heuristic advantages to couching evolutionary radiation studies in the EvA framework:

Firstly, this framework, similar to a model, predicts which factors may be drivers of evolutionary

radiations. This reduces the risk of missing important drivers, and so stimulates the development of

comprehensive models, rather the simple exploration of the effect of a factor on diversification rates. In

addition, it encourages taking recent advances in understanding context-dependence into account.

Secondly, this framework is readily quantified, for example as a regression model. Quantification

both facilitates and encourages data transparency (i.e. what data sets are used, and how these data are

transformed). This transparency becomes more important as the model is expanded to reflect the

complexity of the predictor factors.

Thirdly, it provides a single, general framework within which to analyse all or any evolutionary

radiations. The framework can be applied to any biological organisms, geographical regions, or

ecosystems. This facilitates the comparison among taxa and regions as to the processes underlying

diversification, even if the studies were by different people. This will ease the progression from case

studies to general syntheses.

401 Case study: Conifers

We use a case study of conifer radiation to illustrate EvA implementation and component quantification.

In contrast to the conceptual simplicity of the framework, obtaining data for all components across

multiple lineages can be challenging, although well-developed phylogenies over a wide range of taxa are

increasingly available. Here, we use published data on 455 conifer species (Larcombe, Jordan, Bryant, &

Higgins, 2018) that enable parameterisation of the $d \sim a, b, c$ framework. The conifers provide an

excellent study clade for comparative analysis: the lineage is rich in species grouped into well-defined

clades, geographically widespread, and well-studied with excellent distribution data (Farjon, 2018).

Although conifers originated ca. 300 million years (Ma) ago, with the main clades thought to have

diverged between the early Triassic (ca. 240 Ma) and mid-Cretaceous (ca. 100 Ma), most modern species

arose in the Neogene or Quaternary (23 Ma–present; Leslie et al., 2012). We used the dated phylogeny

of Leslie et al. (2012; inferred from a Bayesian analysis assuming an uncorrelated lognormal clock model,

based on two nuclear and two plastid genes, and calibrated with 16 fossils) to define 70 reciprocally

monophyletic or single-species groups; using a stem age cut-off at 33.9 Ma (the Eocene/Oligocene

boundary) in order to focus on the variables which could explain Neogene–Quaternary diversification

rate variation (Larcombe et al., 2018). Forty-one of these groups have more than one species, the most

species-rich 52 species, and range in age from 34 to 146 Ma.

The factors that contribute to *a*, *b* and *c* in the conifers model were derived from the output of a

process-based niche model (Larcombe et al., 2018). This niche modelling method is described in detail in

Higgins et al. (2012), and is based on a mechanistic model of plant growth, the 'Thornley Transport

Resistance' model, which models resource acquisition, transport and allocation between roots and

shoots, based on environmental information extracted from species distribution data. This produces two

types of information for each species: 1) estimates of the geographic distribution of the potential niche

of each species (i.e., a species distribution model [SDM]); and 2) estimates of the physiological

parameters that describe the niche of each species (Higgins et al., 2012). We used these metrics, and

occurrence data and species richness per clade to parametrise $d \sim a, b, c$ as follows. Note that all four

factors are values for the 41 multispecies clades, not for the constituent species individually (Larcombe

et al., 2018):

d = the diversification rates for each clade: calculated using the method-of-moments estimator of

Magallón and Sanderson (2001) as $r_\varepsilon = \frac{1}{t} ln[n(1 - \varepsilon) + \varepsilon]$, where r_ε is the net diversification rate

assuming a relative extinction fraction ε = 0.9, n the number of extant species, and t the stem age of the

clade.

a = abiotic environment, quantified by the clade's potentially suitable area size: the projected

geographical range reflecting the potential niche of all species within the clades. This is calculated per

clade as the number of ¼° grid cells across the globe that at least one species of the clade can occupy,

based on the physiological SDM and corrected for clade species numbers (i.e., rarefied to the clade with

the lowest diversity to remove sampling effects; Larcombe et al., 2018). This means that if the score is

small, the niche size of a clade is expected to be narrow, and if the score is large, the niche size of the

clade is large so that the clade comprises ecologically more generalist species or the species in the clade

might be specialised but different from one another. Despite its simplifying assumptions about the

spatial distribution of environmental variation (some types of environment are more common than

others), this clade-wise suitable area size is an appealing measure of a because it approximates the

potential niche, consequently biotic interactions and effects of traits can be estimated separately. Figure

3 shows the combined potential niche for all 455 species in the dataset, i.e., the abiotic arena of the

conifers.

Potential SR

☐ 1 – 2
☐ 2 – 10
☐ 10 – 20
☐ 20 – 40
☐ 40 – 60
☐ 60 – 80
☐ 80 – 100
☐ 100 – 175

Figure 2. Projected potential species richness (SR) of conifers representing their abiotic arena. The abiotic
arena of conifers is defined by geographic locations (quarter degree grid cells) that can support one or more of
the 455 conifer species, based on projections from the process-based physiological niche model
(approximating the fundamental conifer niche). The projected potential SR (yellow-green shading) highlights
areas that are suitable for conifer species. The abiotic arena alone does not always predict patterns of conifer
diversity. For example, the eastern Congo is predicted to be climatically suitable for many species but has
relatively low conifer diversity (Farjon, 2018). It is likely that clade-specific traits (*c*) and biotic interactions (*b*)
limit the diversity in certain regions (see main text).

b = competitive interactions estimated at the species level: we determine the expected
competition for each species with all members within its clade (one of the 41 clades defined above) as
the product of niche and geographic overlap between species. The metric underestimates competition
because it excludes competitive processes with other clades and other indirect competitive processes
(see discussion in Larcombe et al., 2018). Geographic overlap is estimated based on the occurrence data.
Niche overlap between each species pair was calculated using Schoener's niche overlap metric *D*
(Schoener, 1968) based on the potential distributions from the SDM analysis. We then scaled these two
numbers to range from 0 to 1 for each species pair and multiplied them to provide a competition index
(Larcombe et al., 2018). This means that if either score is zero, the competition score is zero, and if they
have the same (potential) niche and the same (realized) range then the competition score is 1. The
species-level estimates were averaged to provide a clade-level competition score.

c = clade-specific rate of niche evolution: we used eleven physiological traits (see Fig. 6 in

Larcombe et al., 2018) that were identified as being most important for defining the overall niche space

of conifers (Larcombe et al., 2018). Although an effectively limitless number of physiological traits could

be defined, our method provides an objective selection criterion of ecologically appropriate measures.

These eleven traits were fitted together in a multivariate Brownian motion model of evolution (Butler &

King, 2004) on the conifer phylogeny of Leslie et al. (2012), and the diagonal elements of the resulting

variance–covariance matrix for the species traits represent the phylogenetic rate of evolution (O'Meara,

Ané, Sanderson, & Wainwright, 2006). These were summed and scaled to provide a multidimensional

clade-level niche evolution rate.

Our expectation is that conifer diversification rates (*d*) are positively affected by the available

abiotic environment (*a*) and the rate of niche evolution (*c*), and negatively by inter-specific competition

(*b*). We fitted the conifers EvA model $\sqrt{d} \sim \ln(a) + b + \ln(c)$ by means of phylogenetic generalised

least squares (PGLS), controlling for the non-independence between cases resulting from phylogenetic

structure in the data using the R v3.5.3 (R Core Team, 2013) library 'phylolm' v2.6 (Ho & Ané, 2014).

Note that the variables *a, c* and *d* were transformed to reduce skewness in the data. PGLS estimates the

regression parameters of all variables (the scaled variables used to parameterise the components of

EvA), adjusted for the phylogenetic signal in the model residuals. We accessed the standardised

coefficients and calculated the variance explained by the full model using the coefficient of

determination (R^2) to measure goodness-of-fit, and also assessing partial r^2 (variance explained per

predictor variable *a*, *b*, and *c*) using the R library 'rr2' v1.0.1 (Ives & Li, 2018; Ives, 2019). R scripts are

available in Dryad (Nürk, Linder, et al., 2019).

The full model accounted for 64% of variation in diversification rates among the conifer clades

(adjusted R^2 = 0.638). The predictor variables *a, b,* and *c* in the conifer EvA model differentially

contributed to explaining *d* (table 1). Against our expectations, the abiotic environment of clades (*a*)

showed no relationship to diversification rate (slope -0.003, P = 0.89), neither did the rate of niche

evolution (*c*; slope 0.166, P = 0.36). Contrarily, competitive interactions (*b*) between the species in a

clade indicated a significant negative relationship to diversification (slope -0.581, P < 0.001), supporting

our expectation.

Figure 3. **Quantified EvA model for the conifers**. Coloured trees illustrate the distribution of abiotic environment (*a*), competitive interactions (*b*), rate of niche evolution (*c*), and diversification rates (*d*) across the conifer phylogeny with bars at tips detailing the values per clade and variable. The estimated effects on net diversification rates r_ε are indicated by arrows scaled to the standardised coefficients (slopes) also showing significant levels (ns, non-significant; ***, P <0.001). Colours on branches are rate estimates (obtained using the R package phytools' fastAnc function; Revell, 2012; see R scripts in Nürk, Linder, et al., 2019) and illustrate parameter distribution on the tree. When competition among species is low and the rate of niche evolution in a clade is pronounced, the diversification rate of that clade accelerates.

The significantly negative effect of competition (*b*) on diversification (*d*) indicates higher diversification rates in clades where competition among species is low. This result is consistent with the concept of diversity-dependent diversification (Foote, 2000; Rabosky, 2013) and suggests that diversity-dependent relationships are more important among the conifers in regulating diversification rate than potential area size (*a*) or rates of physiological trait evolution (niche evolution; *c*). However, the rates of niche evolution (*c*) among the conifer clades show a very similar, although inverse, pattern to that of competition (*b*) (Fig. 4). Estimates of a model accounting for interactions among predictor variables (results not shown due to lack of statistical power using n = 41; see R scripts available in Nürk, Linder, et al., 2019) indicated that the two-way interaction *b:c* (competition interacting with rate of niche evolution) influenced diversification in the conifers, in line with findings by Larcombe et al. (2018), who showed that conifer evolution is jointly shaped by bounded and unbounded evolutionary processes (e.g., Harmon & Harrison, 2015). The two-

way interaction *b:c* may enhance or relax diversity-dependent processes so as to promote or constrain

diversification (Larcombe et al., 2018). This is also consistent with the concept that spatial (or temporal)

variation in trait disparity can result in variation in competitive pressure (McPeek, 2008; Marshall &

Quental, 2016). The interaction of competition and rate of niche evolution suggests that the fastest

diversification in conifers is found when competition is low (increased ecological opportunity), which

could be the result of fast trait/niche evolution (high adaptability of the lineage).

The case presented here shows the potential to infer general patterns using the EvA framework.

However, it is in no way a full exploration of the approach, and more sophisticated analyses are likely to

prove more informative. For example, our analysis assumes that rates for *d, b,* and *c* are fixed, which is

an over-simplification. It could be interesting to repeat the analysis using species instead of clades, as

this allows us to account for phylogenetic structure within the clades, ecologically highly variable

species, and diversification stasis. However, there are issues interpreting tip-diversification rates (Title &

Rabosky, 2019). Methods are available to reconstruct ancestral sympatry and infer the effect of

competition on trait divergence and lineage diversification (Aristide & Morlon, 2019; Harmon et al.,

2019). Methods that reconstruct and evolve ancestral states along phylogenies for *a, b,* and *c*, are also

appealing (Uyeda et al., 2018), and with increasing complexity of the EvA model, approaches such as

hidden states will be important for rigorous testing against equivalently complex null hypotheses

(Caetano et al., 2018). This illustrates the value of EvA in making data assumptions explicit.

Conceptual examples

Lupinus continental radiation

Among the most intensively investigated radiations are several in the tropical alpine environments of

the high-elevation Andean grasslands. These environments emerged as a result of the most recent

Pliocene uplift of the Northern Andes, and consequently the radiations themselves are largely confined

to the Pleistocene (Luebert & Weigend, 2014; Hughes & Atchison, 2015). These are exemplified by the

diversification of *c.* 85 species of *Lupinus* L. (lupines, atmospheric nitrogen-fixing Leguminosae) within

the last 1.2–3.5 Ma. The Andean *Lupinus* radiation has been attributed to a combination of intrinsic

evolutionary (trait) innovation and extrinsic ecological opportunity (Hughes & Atchison, 2015). The shift

from an annual to a perennial life history (i.e., evolution of a clade-specific phenotype = *c* in EvA; Fig. 4)

is hypothesised to have acted as a key innovation facilitating occupation of mesic montane habitats

(Drummond, Eastwood, Miotto, & Hughes, 2012), also enabling accelerated disparification of plant

growth forms in the Andes (Nürk, Atchison, et al., 2019). This is because perennials have different cold

tolerance strategies than annuals, underpinning their adaptation to high-elevation ecosystems, and a

fundamentally greater potential growth-form disparity than annuals (Ogburn & Edwards, 2015; Nürk,

Atchison, et al., 2019). At the same time, extrinsic ecological opportunities for diversification were

available in the island-like high-elevation habitats that emerged during the last few million years due to

Andean uplift and cooling of global temperatures (i.e., abiotic factors = a in EvA), prompting Hughes and

Eastwood (2006) to refer to the Andean *Lupinus* clade as an example of "island-like radiation on a

continental scale". In this example, the evolution of secondary perenniality, the shift from lowland to

montane habitats, and the primary shift to higher rates of species diversification all coincide on the

same branch of the phylogeny, presenting an example of a "key confluence" *sensu* Donoghue and

Sanderson (2015). In the Andean *Lupinus* clade, d has been estimated as the rate of species

diversification (Drummond et al., 2012), phenotypic trait evolution or disparification of plant growth

forms (Nürk, Atchison, et al., 2019), and coding DNA sequence evolution (Nevado, Atchison, Hughes, &

Filatov, 2016). All of these estimates of d show accelerated rates across the western New World

montane radiation when compared to the earlier diverging lineages of more slowly diversifying lowland

western New World *Lupinus* annuals.

Lupinus thus provides an apparently straightforward example of a confluence of intrinsic

innovation and extrinsic opportunity as the trigger for accelerated diversification of species and

disparification of plant growth forms. This explanation assumes that ecological opportunity presented

by empty sky island habitats and the means to take advantage of those opportunities (secondary

perenniality) are driving diversification. However, treating the biotic environment (b in EvA) as zero, or

empty of competition is clearly an over-simplification, given that there were apparently many plant

radiations playing out during the Pleistocene across the high-elevation Andean grasslands, presumably

in parallel with each other (Luebert & Weigend, 2014). The detailed order of timing of these radiations

and their interactions remain unknown, just as interactions among sympatric and more or less

contemporaneous radiations have been difficult to tease out more generally (Tanentzap et al., 2015).

The EvA framework draws attention to the fact that biotic factors have not been critically investigated

beyond the simple idea of lack of competition in the newly emerged tropical alpine sky island habitat

(Fig. 5).

Figure 4. **The EvA model for *Lupinus***: $(d_1, d_2, d_3) \sim (a_1, a_2, a_3)$, b_1, c_1. Note that rates of growth form disparification and accelerated gene evolution can influence diversification rates; consequently, the grouping of variables as the components of the EvA framework depends on the questions being investigated.

A more detailed analysis might indicate what factors of these high-elevation habitats are important for the observed high rates of diversification. Indeed, it is aspects of the abiotic environment (*a* in EvA) that are most often put forward as the central explanation for the numerous rapid recent radiations in the high-elevation Andean grassland. Foremost amongst these aspects are: (i) the large continental-scale extent of the high-elevation Andes; (ii) the extreme physiographic heterogeneity; and (iii) the rapid fluctuation in the extent and connectivity between the north Andean alpine sky islands (páramo) during the Pleistocene glacial-interglacial climate cycles. Physiographic heterogeneity of the Andes, spanning steep and extended environmental gradients (e.g., temperature and rainfall), has long been considered as a key factor driving Andean radiations (Hughes & Eastwood, 2006) and indeed of diversification more generally (e.g., Rangel et al., 2018). It has also long been recognised that the area and connectivity of the high-elevation Andean grasslands have varied dramatically through the Pleistocene due to elevational shifts in vegetation zones and species distributions imposed by glacial–interglacial periods. However, it is only recently that area and connectivity have been modelled and quantified in sufficient detail through the Pleistocene (Flantua et al., 2019) to assess the potential of such an alpine 'flickering connectivity system' (Flantua & Hooghiemstra, 2018) to further enhance diversification (e.g., Nevado et al., 2016). Such models demonstrate the need to quantify attributes of the abiotic environment (*a* in

EvA) through time as well as the potential of such time-dependent models to make more realistic

estimates of the impact on diversification rates (d in EvA).

Considering the Andean *Lupinus* radiation in the light of the EvA framework highlights the lack of

knowledge of the biotic interactions involved (*b*). This illuminates that explanations using only the

abiotic environment (*a*) and intrinsic traits (*c*) may be incomplete. Using the EvA framework suggests

new research questions.

Island radiations

Volcanic islands that arise *de novo* on the oceanic crust show a typical lifecycle. In contrast to

islands on the continental shelf, oceanic islands emerge following a volcanic eruption, grow rapidly in

area and elevation, and then erode down to the sea level over 5 to 30 million years, depending on the

substrate and climatic conditions. Even though oceanic islands may show a flickering connectivity effect

as a result of Pleistocene climate fluctuations (e.g., lower sea levels would have resulted in larger islands

areas leaving an imprint on current biodiversity; Weigelt et al., 2016), we here focus on the entire

oceanic island lifecycle (Whittaker, Triantis, & Ladle, 2008; Borregaard et al., 2017). At this scale, island

ontogeny can be considered unimodal in its key properties, and so differs from a flickering model as

described for terrestrial high mountains (Flantua & Hooghiemstra, 2018; Flantua et al., 2019), which is

multimodal. The island lifecycle is an ontogenetic geomorphological trajectory of area, elevation and

habitat diversity – from island birth, through maturity, until island submergence. Consequently, this can

be analysed as a continuous time series. The potential effects of this ontogeny on evolutionary

processes have been described in the general dynamic model of island biogeography (Whittaker et al.,

2008), which provides a temporal framework for variations in island features (Lim & Marshall, 2017)

such as area, topographical complexity, isolation, and habitat diversity (*a* in EvA; Fig. 5). The model has

already been implemented using quantitative methods (Valente, Etienne, & Phillimore, 2014;

Borregaard, Matthews, & Whittaker, 2016) and therefore can define the temporal dimension of the

abiotic arena in an insular context (Fig. 5).

Figure 5. **The EvA model for island radiations**: $(d_1, d_2) \sim (a_1, a_2, a_3), b_1, c_1$. Island ontogeny describes the
typical life-cycle of oceanic islands; isolation the degree of isolation of the insular system (e.g. distance from
the continent); geographical fragmentation could be the number of islands (if dealing with an archipelago).
Incorporating the lineage-specific traits is complex, as several independent lineages may be involved. Trait
disparification is particularly interesting in islands, for example, Hawaiian honeycreepers and silverswords.

In the EvA framework, oceanic islands offer a convenient conceptual aspect, in that the abiotic,

geographical component of the Evolutionary Arena – the island or archipelago – has discrete

boundaries. While many classic studies have analysed insular diversification and disparification (*d* in

EvA) in single monophyletic radiations (e.g., Hawaiian silverswords, Baldwin & Sanderson, 1998;

Madagascan vangas, Jønsson et al., 2012), a potential of islands is that entire communities resulting

from multiple colonisations of the same island can be studied simultaneously, because the abiotic

environment is the same for all included taxa with similar dispersal capacity. For instance, by including

all terrestrial birds of the Galápagos islands in the same model, Valente, Phillimore, and Etienne (2015)

showed that Darwin's finches have statistically exceptional rates of species diversification (i.e.

significantly different from the 'background' rates of all terrestrial Galápagos birds). The effect of the

phenotype (*c* in EvA; Fig. 5) can be considered on a lineage-specific basis (e.g., Givnish et al., 2009) or,

potentially, using a community-level multi-lineage approach, where the effects of given traits are

assessed across multiple insular radiations within the same EvA model. Regarding the effect of the biotic

interactions on islands (*b* in EvA) it needs to be considered that in most cases, there are precursors to

current-day islands that have been eroded to the Pleistocene sea level and are now submerged as guyot

seamounts. These previous islands may explain the fact that the evolutionary age of lineages can be

older than the respective island where they are endemic today (Pillon & Buerki, 2017). Also, while

methods for assessing diversity-dependent effects within single lineages already exist (Rabosky & Glor,

2010; Valente et al., 2015), we currently lack an approach for testing how the interaction of habitat

heterogeneity, island size and present diversity can affect all lineages on an island-wide basis.

EvA provides a heuristic framework for the integration of time-dependent model and multi-clade

analyses. Once several analyses of clades or archipelagos are available in this framework, it should be

possible to combine them to develop a single model for the evolution of diversity within island systems.

Furthermore, island disparification and diversification can be compared using the same analytical

framework, allowing us to test the hypothesis that they respond to the same factors. The simple EvA

framework makes explicit these research question.

Ophrys biotically driven radiation

It is thought that biotic interactions (*b* in EvA) have been a dominant driver of the radiation of the

Mediterranean orchid genus *Ophrys* L., which has produced two parallel adaptive radiations within the

last ~1 Ma. Both of these radiations are characterised by a shift to (mostly) *Andrena* Fabricius solitary

bees as highly specific pollinator species, and by rapid disparification of flowers (Paulus & Gack, 1990;

Breitkopf, Onstein, Cafasso, Schlüter, & Cozzolino, 2015). In this system, pollinators mediate strong

reproductive isolation in the absence of any measurable post-pollination barriers to gene flow among

closely related species (Xu et al., 2011; Sedeek et al., 2014). Consequently, pollinators may drive

speciation in these two parallel radiations.

The high specificity of the pollinators in the *Ophrys* system is due to the plants' chemical mimicry

of the pollinator females' sex pheromones (i.e., phenotypic traits, *c* in EvA), which is predominantly

mediated by alkenes (Schiestl et al., 1999; Xu, Schlüter, Grossniklaus, & Schiestl, 2012). A simple genetic

basis underlies alkene biosynthesis, with only two loci being sufficient to completely change the

pollinator-important alkene double-bond profile sensed by insects (Schlüter et al., 2011; Sedeek et al.,

2016). Selection on alkene composition, and on loci putatively involved in their biosynthesis (*c* in EvA), is

in stark contrast to the rest of the loci in the genome, where abundant polymorphisms are shared across

closely related species (Sedeek et al., 2014). Simulations suggest that this simple trait architecture could

lead to rapid pollinator-driven divergence (Xu & Schlüter, 2015). Overall, the available data suggest that,

given the trait architecture of pollinator attraction, pollinators may be a key factor driving the *Ophrys*

radiations (Fig. 6). However, the relative importance of other factors remains unknown. For example,

what is the potential contribution to phenotypic variation (i.e. disparification; d in EvA) attributable to

the highly heterogeneous (dynamic) abiotic environment (a in EvA) providing habitats for plants and

pollinators (b in EvA) during the last million years?

Figure 6. **The EvA model for *Ophrys***: $d_1 \sim a_1, b_1, (c_1, c_2)$. Floral traits ($c_{1.1}$ geometry, $c_{1.2}$ epidermis
micromorphology, $c_{1.3}$ colour and patterns), also including scent, are probably the most important regulators
of the highly specific pollination system, but the role of mycorrhiza and geographical isolation on the
Mediterranean islands is poorly understood.

Potentially, proxies for all components are measurable here. Due to the extensive amount of allele

sharing, and gene coalescence times frequently predating 'speciation' times (establishment of

reproductive barriers), diversification (d in EvA) among very recent groups of *Ophrys* may best be

assessed not by phylogenetic means, but by within-group pairwise estimates among closely related

species, either in terms of genetic differentiation (e.g., F_{ST}) or phenotypic measurements. Among the

abiotic measurables (a in EvA) would be estimates of habitat fragmentation (also, for example,

estimates from biogeographic and niche modelling approaches) over the estimated age of target

clades/species groups. Biotic interactions (b in EvA) can be represented as matrices of interactions at

varying levels; in its simplest form, orchid species vs. pollinator species. Due to the occurrence of (i)

parallel use of the same pollinators in different lineages and (ii) repeated use of the same pollinators by

allopatric orchid species, such an interaction matrix could initially take the shape of a variance-

covariance matrix as in comparative analyses (O'Meara, 2012). Since the importance of different floral

traits (including morphology and chemistry) for successful interaction with pollinators can be

experimentally measured by quantifying insect responses (e.g., Xu et al., 2012), such an interaction

matrix may eventually contain explicit likelihoods of plant/pollinator interactions (cf. pollination

probabilities in Xu & Schlüter, 2015). Clade-specific intrinsic effects (c in EvA) essentially refer to the

complexity of genetic/genomic change needed to effect a change in a relevant trait. This trait change

would ideally be based upon mechanistic molecular knowledge that could then be quantified

numerically. As a first step in this direction, it may be possible to construct a statistic that summarises

trait–gene associations derived from genomic/transcriptomic data on gene polymorphisms and/or

expression from phenotyped individuals. Overall, although the data to formally test the components of

the EvA framework in the *Ophrys* example are not yet collected, doing so seems at least theoretically

straightforward.

EvA adds a comparative structure to the questions developed above (Table 2). It may also provide

a semantic to bridge micro- and macroevolution in systems where investigations span different levels,

from populations to species and clades.

Conclusions

In this review we synthesize the central concepts of the evolutionary diversification literature and

present the Evolutionary Arena framework (EvA), a heuristic for exploring the modulators of

diversification rates, in terms of the (extrinsic) biotic and abiotic environment and the (intrinsic) traits of

the focal lineage (Table 2), i.e. we integrate three comprehensive classes of diversification rate

modulators. The framework encourages us to organise knowledge about the factors regulating

evolutionary radiations, evolutionary stasis and evolutionary decline of lineages. It does so in a

generalized framework and helps recognise missing information. Particularly in the exploratory phases

of research, EvA may support the search for more complete explanations of diversification rate

variation.

The framework is very flexible, facilitating the incorporation of detailed variables, interactions

among the components, changes in the direction of effect of these components, and interpretation of

phylogenetic conservatism and trait lability. EvA advocates a multivariate perspective on radiations and

can be readily expanded to accommodate increasing levels of complexity, to test for the interactions

among variables, or to rank variables according to their relative influence on diversification rates.

Whether the specific results are tallied, or whether the predictors are collapsed, will most likely depend

on the type of question being asked, and on the power available in the study system. EvA can be

formulated as a hypothesis-testing framework to test whether the likelihood of observing the data

under a favourite particular model provides better fit than an appropriate null model, or to compare

models of varying complexity. The framework may be particularly useful in parameterising data-rich,

broad-scale analyses comparing different systems, such as evolutionary radiations of clades across

different regions, or between different clades within the same region, for example, the plant radiations

in the north Andean páramos. Applying these analyses within the framework allows us to identify the

important components that account for differences in diversification rates between clades and regions.

The Evolutionary Arena framework thus encourages a more comparative approach to exploring

phylogenetic and geographical variation in the correlates and drivers of speciation and extinction.

741   Glossary

Adaptation: a trait is an adaptation to a selective regime if it evolved in response to selection by that

regime (Gould & Vrba, 1982)

Adaptive zone: a fitness peak in a set of related niches (the adaptive grid or macroevolutionary

landscape) that a lineage occupies by virtue of a novel trait(s) that confer fitness in these niches.

Confluence: the sequential coming together of a set of traits (innovations and synnovations),

environmental changes, and geographic movements along the branches of a phylogenetic tree

(Donoghue & Sanderson, 2015).

Disparification: increase in trait variance in a clade through time. That is, increase in measurable

phenotypic differences among taxa, whereas the traits in question may be morphological,

anatomical, physiological, genetic, behavioural, etc. Disparification is a characteristic of adaptive

radiation.

Diversification: increase in the taxonomic diversity in a clade through time. The diversification rate is

defined as speciation *minus* extinction and can thus be negative.

Ecological opportunity: lineage-specific environmental conditions that contain both niche availability
and niche discordance, favouring adaptation and promoting diversifying selection within the
lineage (adapted from Wellborn & Langerhans, 2015).

Exaptation: a trait that has evolved for one use, and that is later useful for another usage (sometimes
deceptively termed 'pre-adaptation'; adapted from de Vladar et al., 2017). The original
definition of Gould and Vrba (1982) is: "features that now enhance fitness but were not built by
natural selection for their current role".

Extrinsic factor: environmental factors such as abiotic and biotic niche parameters, not inherited
genetically by the focal lineage.

Intrinsic factor: phenotypic (morphological, physiological) or genetic trait(s), inherited by the focal
lineage.

Key event: events that trigger a shift in diversification rates.

Key innovation: new trait which facilitates the occupation of a new adaptive zone, or which breaks an
evolutionary constraint; i.e. "phenotypes that allowed a species to interact with the
environment in a novel way" (Stroud & Losos, 2016, p 508).

Phenotype: a set of features of an individual that stems from the interactions between genotype and
environment.

Radiation: accelerated proliferation of species and/or phenotypes, in the sense of significant increase in
the diversification and/or disparification rate compared to background rates (without a shift/
significant rate increase, it is not a radiation but [background] diversification / disparification).
Radiation (diversification / disparification) can be combined with an epithet, such as adaptive,
geographical, ecological, or genetical (gene flow) to further describe the nature of the
evolutionary forces and situations (e.g., 'sexual radiation' may refer to radiation driven by sexual
selection, or 'montane radiation' to radiation in mountains, 'insular radiation' to radiations on
islands, or simply 'cichlid radiation' to refer to a certain lineage, etc.). We refer to biological
radiation most generally as '**evolutionary radiation**' as the change in diversification /
disparification rates has macro-evolutionary consequences. In addition, there are two
prominent concepts that refer to the process underlying the evolutionary radiations:

Adaptive radiation: proliferation of species driven by the evolution of phenotypic (ecological
and/or morphological) diversity that can be linked to adaptation to an environment. The

environment may act as a modulator, driving (potentially sympatric) speciation and/or slowing

extinction.

Geographic radiation: proliferation of species driven by enhanced opportunities for allopatric

speciation (reproductive isolation resulting from spatial barriers) in a particular region (modified

from Simões et al., 2016). Also referred to as 'non-adaptive radiation' (= geographic), or 'climatic

radiation' when differing climates are thought important. Note that the adaptive and geographic

categories are simplified: both adaptive and neutral processes likely play a role in modulating

diversification rates in most radiations, but their relative contributions differ. For example,

ecological factors may enhance the opportunity for reproductive isolation, and species

divergence in adaptive radiations may additionally be promoted by spatial isolation (see main

text, Drivers of evolutionary radiations).

Synnovation: interacting combination of traits with a particular consequence (Donoghue & Sanderson,

2015).

Trait: a heritable attribute of evolutionary lineages (genes, individuals, populations, species, clades) that

can be observed.

Trigger: Event or situation starting a radiation.

Data Accessibility Statement

803   The phylogenetic tree, the data matrix and R scripts used in the analysis of conifers are available

804 at Dryad: doi: 10.5061/dryad.2bvq83bkx (Nürk, Linder, et al., 2019).

Competing Interests Statement

806   None declared.

Author Contributions section

808   HPL and NMN developed research concept, all authors contributed to developing the Evolutionary

809 Arena framework, MJL, NMN and REO analysed the data, CEH, LPF, LV, and PMS contributed the

810   conceptual examples. HPL and NMN wrote the manuscript and all authors contributed critically to the

811   drafts and gave final approval for publication.

812   Acknowledgements

We especially thank Seraina Klopfstein, Susanne Renner and Bob Ricklefs for discussions and

important comments, Andreas Franzke for commenting on an earlier version of this manuscript, and

Isabell Niclas, Stefan Pelka and Bernhard Maier for their help during an ambulatory workshop meeting in

August 2018 where the ideas presented here were discussed and developed, and which was supported

by a WinUBT (Bayreuth) conference grant to NMN. REO acknowledges the support of the German

Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig funded by the Deutsche

Forschungsgemeinschaft (DFG, German Research Foundation)—FZT 118. SGAF acknowledges the ERC

Advanced Grant 741413 Humans on Planet Earth (HOPE).

822   References

Aguilée, R., Gascuel, F., Lambert, A., & Ferriere, R. (2018). Clade diversification dynamics and the biotic and abiotic controls of
speciation and extinction rates. *Nature Communications, 9*(1), 3013. doi:10.1038/s41467-018-05419-7

Albert, V. A., Barbazuk, W. B., dePamphilis, C. W., Der, J. P., Leebens-Mack, J., Ma, H., . . . Tomsho, L. (2013). The *Amborella*
genome and the evolution of flowering plants. *Science, 342*(6165), 1241089–1241089. doi:10.1126/science.1241089

Alfaro, M. E., Santini, F., Brock, C., Alamillo, H., Dornburg, A., Rabosky, D. L., . . . Harmon, L. J. (2009). Nine exceptional
radiations plus high turnover explain species diversity in jawed vertebrates. *Proceedings of the National Academy of*
Sciences of the United States of America, 106(32), 13410–13414. doi:10.1073/pnas.0811087106

Arakaki, M., Christin, P.-A., Nyffeler, R., Lendel, A., Eggli, U., Ogburn, R. M., . . . Edwards, E. J. (2011). Contemporaneous and
recent radiations of the world's major succulent plant lineages. *Proceedings of the National Academy of Sciences of*
the United States of America, 108(20), 8379–8384. doi:10.1073/pnas.1100628108

Aristide, L., & Morlon, H. (2019). Understanding the effect of competition during evolutionary radiations: An integrated model
of phenotypic and species diversification. *Ecology Letters, 22*(12), 2006–2017. doi:10.1111/ele.13385

Badgley, C., Smiley, T. M., Terry, R., Davis, E. B., DeSantis, L. R. G., Fox, D. L., . . . Yanites, B. J. (2017). Biodiversity and
topographic complexity: Modern and geohistorical perspectives. *Trends in Ecology & Evolution, 32*(3), 211–226.
doi:10.1016/j.tree.2016.12.010

Bailey, S. F., Dettman, J. R., Rainey, P. B., & Kassen, R. (2013). Competition both drives and impedes diversification in a model
adaptive radiation. *Proceedings of the Royal Society B: Biological Sciences, 280*(1766), 20131253.
doi:10.1098/rspb.2013.1253

Baldwin, B. G. (1997). Adaptive radiation of the Hawaiian silversword alliance: Congruence and conflict of phylogenetic
evidence from molecular and non-molecular investigations. In T. J. Givnish & K. J. Sytsma (Eds.), *Molecular evolution*
and adaptive radiation (pp. 103—128). London: Cambridge University Press, UK.

Baldwin, B. G., & Sanderson, M. J. (1998). Age and rate of diversification of the Hawaiian silversword alliance (Compositae).
Proceedings of the National Academy of Sciences of the United States of America, 95(16), 9402–9406.
doi:10.1073/pnas.95.16.9402

Beaulieu, J. M., & O'Meara, B. C. (2016). Detecting hidden diversification shifts in models of trait-dependent speciation and
extinction. *Systematic Biology, 65*(4), 583–601. doi:10.1093/sysbio/syw022

Beaulieu, J. M., O'Meara, B. C., & Donoghue, M. J. (2013). Identifying hidden rate changes in the evolution of a binary
morphological character: The evolution of plant habit in campanulid angiosperms. *Systematic Biology, 62*(5), 725–
737. doi:10.1093/sysbio/syt034

Beerling, D. J., & Osborne, C. P. (2006). The origin of the savanna biome. *Global Change Biology, 12*(11), 2023–2031.
doi:10.1111/j.1365-2486.2006.01239.x

Benton, M. J. (1995). Diversification and extinction in the history of life. *Science, 268*(5207), 52–58.
doi:10.1126/science.7701342

Benton, M. J. (2009). The Red Queen and the Court Jester: Species diversity and the role of biotic and abiotic factors through
time. *Science, 323*(5915), 728–723. doi:10.1126/science.1157719

Bond, W. J., Woodward, F. I., & Midgley, G. F. (2005). The global distribution of ecosystems in a world without fire. *New
Phytologist, 165*(2), 525–537. doi:10.1111/j.1469-8137.2004.01252.x

Borregaard, M. K., Amorim, I. R., Borges, P. A. V., Cabral, J. S., Fernandez-Palacios, J. M., Field, R., . . . Whittaker, R. J. (2017).
Oceanic island biogeography through the lens of the general dynamic model: Assessment and prospect. *Biological
Reviews, 92*(2), 830–853. doi:10.1111/brv.12256

Borregaard, M. K., Matthews, T. J., & Whittaker, R. J. (2016). The general dynamic model: Towards a unified theory of island
biogeography? *Global Ecology and Biogeography, 25*(7), 805–816. doi:10.1111/geb.12348

Bouchenak-Khelladi, Y., Onstein, R. E., Xing, Y., Schwery, O., & Linder, H. P. (2015). On the complexity of triggering evolutionary
radiations. *New Phytologist, 207*(2), 313–326. doi:10.1111/nph.13331

Boucher, F. C., Demery, V., Conti, E., Harmon, L. J., & Uyeda, J. (2018). A general model for estimating macroevolutionary
landscapes. *Systematic Biology, 67*(2), 304–319. doi:10.1093/sysbio/syx075

Brassac, J., & Blattner, F. R. (2015). Species-level phylogeny and polyploid relationships in *Hordeum* (Poaceae) inferred by next-
generation sequencing and *in silico* cloning of multiple nuclear loci. *Systematic Biology, 64*(5), 792–808.
doi:10.1093/sysbio/syv035

Brawand, D., Wagner, C. E., Li, Y. I., Malinsky, M., Keller, I., Fan, S. H., . . . Di Palma, F. (2014). The genomic substrate for
adaptive radiation in African cichlid fish. *Nature, 513*(7518), 375–381. doi:10.1038/nature13726

Breitkopf, H., Onstein, R. E., Cafasso, D., Schlüter, P. M., & Cozzolino, S. (2015). Multiple shifts to different pollinators fuelled
rapid diversification in sexually deceptive *Ophrys* orchids. *New Phytologist, 207*(2), 377–389. doi:10.1111/nph.13219

Brodersen, J., Post, D. M., & Seehausen, O. (2018). Upward adaptive radiation Cascades: Predator diversification induced by
prey diversification. *Trends in Ecology & Evolution, 33*(1), 59–70. doi:10.1016/j.tree.2017.09.016

Burnham, K. P., & Anderson, D. R. (2002). *Model selection and multimodel inference: A practical information-theoretic approach*
(2nd ed.). New York, Berlin, Heidelberg: Springer Verlag.

Butler, M. A., & King, A. A. (2004). Phylogenetic comparative analysis: A modeling approach for adaptive evolution. *American
Naturalist, 164*(6), 683–695. doi:10.1086/426002

Caetano, D. S., O'Meara, B. C., & Beaulieu, J. M. (2018). Hidden state models improve state-dependent diversification
approaches, including biogeographical models. *Evolution, 27*(11), 2308–2324. doi:10.1111/evo.13602

Comes, H. P., Tribsch, A., & Bittkau, C. (2008). Plant speciation in continental island floras as exemplified by *Nigella* in the
Aegean Archipelago. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences,
363*(1506), 3083–3096. doi:10.1098/rstb.2008.0063

Condamine, F. L., Romieu, J., & Guinot, G. (2019). Climate cooling and clade competition likely drove the decline of lamniform
sharks. *Proceedings of the National Academy of Sciences, 116*(41), 20584–20590. doi:10.1073/pnas.1902693116

Cooney, C. R., Bright, J. A., Capp, E. J. R., Chira, A. M., Hughes, E. C., Moody, C. J. A., . . . Thomas, G. H. (2017). Mega-
evolutionary dynamics of the adaptive radiation of birds. *Nature, 542*(7641), 344–347. doi:10.1038/nature21074

Czekanski-Moir, J. E., & Rundell, R. J. (2019). The Ecology of Nonecological Speciation and Nonadaptive Radiations. *Trends in
Ecology & Evolution, 34*(5), 400–415. doi:10.1016/j.tree.2019.01.012

Davis, C. C., Schaefer, H., Xi, Z., Baum, D. A., Donoghue, M. J., & Harmon, L. J. (2014). Long-term morphological stasis
maintained by a plant-pollinator mutualism. *Proceedings of the National Academy of Sciences of the United States of*
America, 111(16), 5914–5919. doi:10.1073/pnas.1403157111

de Vladar, H. P., Santos, M., & Szathmary, E. (2017). Grand views of evolution. *Trends in Ecology & Evolution, 32*(5), 324–334.
doi:10.1016/j.tree.2017.01.008

Degnan, J. H., & Rosenberg, N. A. (2006). Discordance of species trees with their most likely gene trees. *PLoS Genetics, 2*(5),
762–768. doi:10.1371/journal.pgen.0020068

Donoghue, M. J., & Edwards, E. J. (2014). Biome shifts and niche evolution in plants. *Annual Review of Ecology, Evolution, and*
Systematics, 45(1), 547–572. doi:10.1146/annurev-ecolsys-120213-091905

Donoghue, M. J., & Sanderson, M. J. (2015). Confluence, synnovation, and depauperons in plant diversification. *New*
Phytologist, 207(2), 260–274. doi:10.1111/nph.13367

Drummond, A. J., Ho, S. Y. W., Phillips, M. J., & Rambaut, A. (2006). Relaxed phylogenetics and dating with confidence. *PLoS*
Biology, 4(5), 699–710. doi:10.1371/journal.pbio.0040088

Drummond, C. S., Eastwood, R. J., Miotto, S. T. S., & Hughes, C. E. (2012). Multiple continental radiations and correlates of
diversification in *Lupinus* (Leguminosae): Testing for key innovation with incomplete taxon sampling. *Systematic*
Biology, 61(3), 443–460. doi:10.1093/sysbio/syr126

Duret, L. (2002). Evolution of synonymous codon usage in metazoans. *Current Opinion in Genetics & Development, 12*(6), 640–
649. doi:10.1016/S0959-437x(02)00353-2

Edwards, S. V. (2009). Is a new and general theory of molecular systematics emerging? *Evolution, 63*(1), 1–19. doi:Doi
10.1111/J.1558-5646.2008.00549.X

Eldredge, N., & Gould, S. J. (1972). Speciation and punctuated equilibria: An alternative to phyletic gradualism. In T. J. M. Schopf
(Ed.), *Models in Paleobiology* (pp. 82–115). San Francisco: Freeman, Cooper and Company.

Erwin, D. H. (1992). A preliminary classification of evolutionary radiations. *Historical Biology, 6*(2), 133–147.
doi:10.1080/10292389209380423

Erwin, D. H. (2001). Lessons from the past: Biotic recoveries from mass extinctions. *Proceedings of the National Academy of*
Sciences, 98(10), 5399–5403. doi:10.1073/pnas.091092698

Erwin, D. H. (2015). Novelty and Innovation in the history of life. *Current Biology, 25*(19), R930—R940.
doi:10.1016/j.cub.2015.08.019

Erwin, D. H. (2017). The topology of evolutionary novelty and innovation in macroevolution. *Philosophical transactions of the*
Royal Society of London. Series B, Biological sciences, 372(1735), 20160422. doi:10.1098/rstb.2016.0422

Ezard, T. H. G., Aze, T., Pearson, P. N., & Purvis, A. (2011). Interplay between changing climate and species' ecology drives
macroevolutionary dynamics. *Science, 332*(6027), 349–351. doi:10.1126/science.1203060

Farjon, A. (2018). The Kew review: Conifers of the world. *Kew Bulletin, 73*(8), 1–16. doi:10.1007/s12225-018-9738-5

Felsenstein, J. (1973). Maximum-likelihood estimation of evolutionary trees from continuous characters. *American Journal of*
Human Genetics, 25(5), 471–492. doi:10.2307/2412304

Felsenstein, J. (1981). Evolutionary trees from DNA sequences: A maximum likelihood approach. *Journal of Molecular Evolution,*
17(6), 368–376. doi:10.1007/BF01734359

Felsenstein, J. (1985). Phylogenies and the comparative method. *American Naturalist, 125*(1), 1–15. doi:10.1086/284325

Felsenstein, J. (2012). A comparative method for both discrete and continuous characters using the threshold model. *American*
Naturalist, 179(2), 145–156. doi:10.1086/663681

Field, R., Hawkins, B. A., Cornell, H. V., Currie, D. J., Diniz-Filho, J. A. F., Guegan, J. F., . . . Turner, J. R. G. (2009). Spatial species-
richness gradients across scales: A meta-analysis. *Journal of Biogeography, 36*(1), 132–147. doi:10.1111/j.1365-
2699.2008.01963.x

FitzJohn, R. G. (2010). Quantitative traits and diversification. *Systematic Biology, 59*(6), 619–633. doi:10.1093/sysbio/syq053

FitzJohn, R. G. (2012). Diversitree: Comparative phylogenetic analyses of diversification in R. *Methods in Ecology and Evolution,*
3(6), 1084–1092. doi:10.1111/j.2041-210X.2012.00234.x

Flantua, S. G. A., & Hooghiemstra, H. (2018). Historical connectivity and mountain biodiversity. In C. Hoorn, A. Perrigo, & A.
Antonelli (Eds.), *Mountains, climate and biodiversity* (pp. 171—186): Wiley-Blackwell.

Flantua, S. G. A., O'Dea, A., Onstein, R. E., Giraldo, C., & Hooghiemstra, H. (2019). The flickering connectivity system of the north
Andean páramos. *Journal of Biogeography, 46*(1), 1808–1825. doi:10.1111/jbi.13607;
https://www.youtube.com/watch?v=-Wcp18vBDK4

Flegr, J. (2010). Elastic, not plastic species: Frozen plasticity theory and the origin of adaptive evolution in sexually reproducing
organisms. *Biology Direct, 5*, 2. doi:10.1186/1745-6150-5-2

Foote, M. (2000). Origination and extinction components of taxonomic diversity: Paleozoic and post-Paleozoic dynamics.
Paleobiology, 26(4), 578–605. doi:10.1666/0094-8373(2000)026<0578:Oaecot>2.0.Co;2

Fryer, G. (1969). Speciation and adaptive radiation in African lakes. *SIL Proceedings, 1922-2010, 17*(1), 303–322.
doi:10.1080/03680770.1968.11895855

Gavini, S. S., Ezcurra, C., & Aizen, M. A. (2019). Plant–plant interactions promote alpine diversification. *Evolutionary ecology,*
33(2), 195–209. doi:10.1007/s10682-019-09972-5

Gittenberger, E. (1991). What about non-adaptive radiation. *Biological Journal of the Linnean Society, 43*(4), 263–272.
doi:10.1111/j.1095-8312.1991.tb00598.x

Gittenberger, E. (2004). Radiation and adaptation, evolutionary biology and semantics. *Organisms Diversity & Evolution, 4*(3),
135–136. doi:10.1016/j.ode.2004.04.002

Givnish, T. J. (2015). Adaptive radiation versus "radiation' and "explosive diversification': Why conceptual distinctions are
fundamental to understanding evolution. *New Phytologist, 207*(2), 297–303. doi:10.1111/nph.13482

Givnish, T. J., Millam, K. C., Mast, A. R., Paterson, T. B., Theim, T. J., Hipp, A. L., . . . Sytsma, K. J. (2009). Origin, adaptive radiation
and diversification of the Hawaiian lobeliads (Asterales: Campanulaceae). *Proceedings of the Royal Society of London.*
Series B, Biological sciences, 276(1656), 407–416. doi:10.1098/rspb.2008.1204

Goldberg, E. E., Lancaster, L. T., & Ree, R. H. (2011). Phylogenetic inference of reciprocal effects between geographic range
evolution and diversification. *Systematic Biology, 60*(4), 451–465. doi:10.1093/sysbio/syr046

Gould, S. J., & Eldredge, N. (1977). Punctuated equilibria: The tempo and mode of evolution reconsidered. *Paleobiology, 3*(2),
115–151. doi:10.1017/S0094837300005224

Gould, S. J., & Vrba, E. S. (1982). Exaptation – A missing term in the science of form. *Paleobiology, 8*(1), 4–15.
doi:10.1017/S0094837300004310

Grafen, A. (1989). The phylogenetic regression. *Philosophical Transactions of the Royal Society of London Series B-Biological*
Sciences, 326(1233), 119–157. doi:10.1098/rstb.1989.0106

Guerrero, P. C., Rosas, M., Arroyo, M. T. K., & Wiens, J. J. (2013). Evolutionary lag times and recent origin of the biota of an
ancient desert (Atacama-Sechura). *Proceedings of the National Academy of Sciences of the United States of America,*
110(28), 11469–11474. doi:10.1073/pnas.1308721110

Harmon, L. J. (2018). *Phylogenetic comparative methods, version 1.4.* https://lukejharmon.github.io/pcm/.

Harmon, L. J., Andreazzi, C. S., Débarre, F., Drury, J., Goldberg, E. E., Martins, A. B., . . . Matthews, B. (2019). Detecting the
macroevolutionary signal of species interactions. *Journal of Evolutionary Biology, 32*(8), 769–782.
doi:10.1111/jeb.13477

Harmon, L. J., & Harrison, S. (2015). Species diversity is dynamic and unbounded at local and continental scales. *American*
Naturalist, 185(5), 584–593. doi:10.1086/680859

Harpole, W. S., Sullivan, L. L., Lind, E. M., Firn, J., Adler, P. B., Borer, E. T., . . . Wragg, P. D. (2016). Addition of multiple limiting
resources reduces grassland diversity. *Nature, 537*(7618), 93–96. doi:10.1038/nature19324

Heard, S. B., & Hauser, D. L. (1995). Key evolutionary innovations and their ecological mechanisms. *Historical Biology, 10*(2),
151–173. doi:10.1080/10292389509380518

Hennig, W. (1950). *Grundzüge einer Theorie der phylogenetischen Systematik*. Berlin: Deutscher Zentralverlag.

Hennig, W. (1965). Phylogenetic systematics. *Annual Review of Entomology, 10*, 97–116. doi:papers://45568F70-9A94-472F-
BFDF-D607B4F5E0A4/Paper/p965

Higgins, S. I., O'Hara, R. B., Bykova, O., Cramer, M. D., Chuine, I., Gerstner, E. M., . . . Midgley, G. F. (2012). A physiological
analogy of the niche for projecting the potential distribution of plants. *Journal of Biogeography, 39*(12), 2132–2145.
doi:10.1111/j.1365-2699.2012.02752.x

Ho, L. S. T., & Ané, C. (2014). A linear-time algorithm for Gaussian and non-Gaussian trait evolution models. *Systematic Biology,*
63(3), 397–408. doi:10.1093/sysbio/syu005

Höhna, S., Heath, T. A., Boussau, B., Landis, M. J., Ronquist, F., & Huelsenbeck, J. P. (2014). Probabilistic graphical model
representation in phylogenetics. *Systematic Biology, 63*(5), 753–771. doi:10.1093/sysbio/syu039

Holt, R. D. (2009). Bringing the Hutchinsonian niche into the 21st century: Ecological and evolutionary perspectives. *Proceedings*
of the National Academy of Sciences of the United States of America, 106, 19659–19665.
doi:10.1073/pnas.0905137106

Huelsenbeck, J. P., Nielsen, R., & Bollback, J. P. (2003). Stochastic mapping of morphological characters. *Systematic Biology,*
52(2), 131–158. doi:10.1080/10635150390192780

Huelsenbeck, J. P., Ronquist, F., Nielsen, R., & Bollback, J. P. (2001). Bayesian inference of phylogeny and its impact on
evolutionary biology. *Science, 294*(5550), 2310–2314.

Huey, R. B., Garland, T., & Turelli, M. (2019). Revisiting a key innovation in evolutionary biology: Felsenstein's "phylogenies and
the comparative method". *The American Naturalist, 193*(6), 755–772. doi:10.1086/703055

Hughes, C. E., & Atchison, G. W. (2015). The ubiquity of alpine plant radiations: From the Andes to the Hengduan Mountains.
New Phytologist, 207(2), 275–282. doi:10.1111/nph.13230

Hughes, C. E., & Eastwood, R. (2006). Island radiation on a continental scale: Exceptional rates of plant diversification after
uplift of the Andes. *Proceedings of the National Academy of Sciences of the United States of America, 103*(27), 10334–
10339. doi:10.1073/pnas.0601928103

Hunter, J. P. (1998). Key innovations and the ecology of macroevolution. *Trends in Ecology & Evolution, 13*(1), 31–36.
doi:10.1016/S0169-5347(97)01273-1

Irisarri, I., Singh, P., Koblmüller, S., Torres-Dowdall, J., Henning, F., Franchini, P., . . . Meyer, A. (2018). Phylogenomics uncovers
early hybridization and adaptive loci shaping the radiation of Lake Tanganyika cichlid fishes. *Nature Communications,*
9(1), 3159. doi:10.1038/s41467-018-05479-9

Ives, A. R. (2019). R2s for correlated data: Phylogenetic models, LMMs, and GLMMs. *Systematic Biology, 68*(2), 234–251.
doi:10.1093/sysbio/syy060

Ives, A. R., & Li, D. (2018). rr2: An R package to calculate R2s for regression models. *Journal of Open Source Software, 3*(30),
1028. doi:10.21105/joss.01028

Jentsch, A., & White, P. S. (2019). A theory of pulse dynamics and disturbance in ecology. *Ecology, 100*(7), e02734.
doi:10.1002/ecy.2734

Johnson, S. D., Linder, H. P., & Steiner, K. E. (1998). Phylogeny and radiation of pollination systems in *Disa* (Orchidaceae).
American Journal of Botany, 85(3), 402–411. doi:10.2307/2446333

Jønsson, K. A., Fabre, P. H., Fritz, S. A., Etienne, R. S., Ricklefs, R. E., Jørgensen, T. B., . . . Irestedt, M. (2012). Ecological and
evolutionary determinants for the adaptive radiation of the Madagascan vangas. *Proceedings of the National*
Academy of Sciences of the United States of America, 109(17), 6620–6625. doi:10.1073/pnas.1115835109

Kimura, M. (1968). Evolutionary rate at the molecular level. *Nature, 217*(5129), 624–626. doi:10.1038/217624a0

Kozak, K. H., & Wiens, J. J. (2010). Accelerated rates of climatic-niche evolution underlie rapid species diversification. *Ecology*
Letters, 13(11), 1378–1389. doi:10.1111/j.1461-0248.2010.01530.x

Laland, K. N., Matthews, B., & Feldman, M. W. (2016). An introduction to niche construction theory. *Evolutionary ecology,*
30(1), 191–202. doi:10.1007/s10682-016-9821-z

Landis, J. B., Soltis, D. E., Li, Z., Marx, H. E., Barker, M. S., Tank, D. C., & Soltis, P. S. (2018). Impact of whole-genome duplication
events on diversification rates in angiosperms. *American Journal of Botany, 105*(3), 348–363. doi:10.1002/ajb2.1060

Larcombe, M. J., Jordan, G. J., Bryant, D., & Higgins, S. I. (2018). The dimensionality of niche space allows bounded and
unbounded processes to jointly influence diversification. *Nature Communications, 9*(1), 4258. doi:10.1038/s41467-
018-06732-x

Leslie, A. B., Beaulieu, J. M., Rai, H. S., Crane, P. R., Donoghue, M. J., & Mathews, S. (2012). Hemisphere-scale differences in
conifer evolutionary dynamics. *Proceedings of the National Academy of Sciences of the United States of America,*
109(40), 16217–16221. doi:10.1073/pnas.1213621109

Liem, K. F. (1973). Evolutionary strategies and morphological innovations: Cichlid pharyngeal jaws. *Systematic Zoology, 22*(4),
425—441. doi:10.2307/2412950

Lim, J. Y., & Marshall, C. R. (2017). The true tempo of evolutionary radiation and decline revealed on the Hawaiian archipelago.
Nature, 543(7647), 710–713. doi:10.1038/nature21675

Linder, H. P., & Bouchenak-Khelladi, Y. (2017). Adaptive radiations should not be simplified: The case of the danthonioid
grasses. *Molecular Phylogenetics and Evolution, 117*, 179–190. doi:10.1016/j.ympev.2017.10.003

Linder, H. P., Lehmann, C. E. R., Archibald, S., Osborne, C. P., & Richardson, D. M. (2018). Global grass (Poaceae) success
underpinned by traits facilitating colonization, persistence and habitat transformation. *Biological Reviews, 93*(2),
1125–1144. doi:10.1111/brv.12388

Losos, J. B. (1994). Integrative approaches to evolutionary ecology - Anolis Lizards as model systems. *Annual Review of Ecology*
and Systematics, 25(1), 467–493. doi:10.1146/annurev.es.25.110194.002343

Losos, J. B., & Ricklefs, R. E. (2009). Adaptation and diversification on islands. *Nature, 457*(7231), 830–836.
doi:10.1038/nature07893

Luebert, F., & Weigend, M. (2014). Phylogenetic insights into Andean plant diversification. *Frontiers in Ecology and Evolution, 2*,
27. doi:10.3389/fevo.2014.00027

Maddison, W. P. (1997). Gene trees in species trees. *Systematic Biology, 46*(3), 523–536. doi:10.1093/sysbio/46.3.523

Maddison, W. P., & FitzJohn, R. G. (2015). The unsolved challenge to phylogenetic correlation tests for categorical characters.
Systematic Biology, 64(1), 127–136. doi:10.1093/sysbio/syu070

Maddison, W. P., Midford, P. E., & Otto, S. P. (2007). Estimating a binary character's effect on speciation and extinction.
Systematic Biology, 56(5), 701–710. doi:10.1080/10635150701607033

Magallón, S. S., & Sanderson, M. J. (2001). Absolute diversification rates in Angiosperm clades. *Evolution, 55*(9), 1762–1780.
doi:10.2307/2680421

Marazzi, B., Ané, C., Simon, M. F., Delgado-Salinas, A., Luckow, M., & Sanderson, M. J. (2012). Locating evolutionary precursors
on a phylogenetic tree. *Evolution, 66*(12), 3918–3930. doi:10.1111/j.1558-5646.2012.01720.x

Marques, D. A., Meier, J. I., & Seehausen, O. (2019). A combinatorial view on speciation and adaptive radiation. *Trends in*
Ecology & Evolution, 34(6), 531–544. doi:10.1016/j.tree.2019.02.008

Marshall, C. R., & Quental, T. B. (2016). The uncertain role of diversity dependence in species diversification and the need to
incorporate time-varying carrying capacities. *Philosophical transactions of the Royal Society of London. Series B,*
Biological sciences, 371(1691), 20150217. doi:10.1098/rstb.2015.0217

Martins, E. P., & Hansen, T. F. (1997). Phylogenies and the comparative method: A general approach to incorporating
phylogenetic information into the analysis of interspecific data. *American Naturalist, 149*(4), 646–667.
doi:10.1086/286013

Mayr, E. (1954). Geographic speciation in tropical Echinoids. *Evolution, 8*(1), 1–18. doi:10.1111/j.1558-5646.1954.tb00104.x

McInerny, G. J., & Etienne, R. S. (2012). Stitch the niche – a practical philosophy and visual schematic for the niche concept.
Journal of Biogeography, 39(12), 2103–2111. doi:10.1111/jbi.12032

McInerny, G. J., Etienne, R. S., & Higgins, S. (2012). Ditch the niche - is the niche a useful concept in ecology or species
distribution modelling? *Journal of Biogeography, 39*(12), 2096–2102. doi:10.1111/jbi.12033

McPeek, M. A. (2008). The ecological dynamics of clade diversification and community assembly. *The American Naturalist,*
172(6), E270—E284. doi:10.1086/593137

Meier, J. I., Marques, D. A., Mwaiko, S., Wagner, C. E., Excoffier, L., & Seehausen, O. (2017). Ancient hybridization fuels rapid
cichlid fish adaptive radiations. *Nature Communications, 8*, 14363. doi:10.1038/ncomms14363

Miller, A. H. (1949). Some ecologic and morphologic considerations in the evolution of higher taxonomic categories. In E. Mayr
& E. Schüz (Eds.), *Ornithologie als Biologische Wissenschaft: 28 Beiträge als Festschrift zum 60. Geburtstag von Erwin*
Stresemann (22. November 1949) (pp. 84–88). Heidelberg: Carl Winter.

Morlon, H. l. n. (2014). Phylogenetic approaches for studying diversification. *Ecology Letters, 17*(4), 508–525.
doi:10.1111/ele.12251

Naciri, Y., & Linder, H. P. (2020). The genetics of evolutionary radiations. *Biological Reviews*, 1–71.

Nee, S., May, R. M., & Harvey, P. H. (1994). The reconstructed evolutionary process. *Philosophical transactions of the Royal*
Society of London. Series B, Biological sciences, 344(1309), 305–311. doi:10.1098/rstb.1994.0068

Nevado, B., Atchison, G. W., Hughes, C. E., & Filatov, D. A. (2016). Widespread adaptive evolution during repeated evolutionary
radiations in New World lupins. *Nature Communications, 7*(12384), 1–9. doi:10.1038/ncomms12384

Nürk, N. M., Atchison, G. W., & Hughes, C. E. (2019). Island woodiness underpins accelerated disparification in plant radiations.
New Phytologist, 224(1), 518—531. doi:10.1111/nph.15797

Nürk, N. M., Linder, H. P., Onstein, R. E., Larcombe, M. J., Hughes, C. E., Piñeiro Fernández, L., . . . Pirie, M. D. (2019). *Data from:*
Diversification in evolutionary arenas – assessment and synthesis. Retrieved from:
https://doi.org/10.5061/dryad.2bvq83bkx

Nürk, N. M., Michling, F., & Linder, H. P. (2018). Are the radiations of temperate lineages in tropical alpine ecosystems pre-
adapted? *Global Ecology and Biogeography, 27*(3), 334–345. doi:10.1111/geb.12699

O'Meara, B. C. (2012). Evolutionary inferences from phylogenies: A review of methods. *Annual Review of Ecology, Evolution,*
and Systematics, 43(1), 267–285. doi:10.1146/annurev-ecolsys-110411-160331

O'Meara, B. C., Ané, C., Sanderson, M. J., & Wainwright, P. C. (2006). Testing for different rates of continuous trait evolution
using likelihood. *Evolution, 60*(5), 922–933. doi:10.1111/j.0014-3820.2006.tb01171.x

O'Meara, B. C., & Beaulieu, J. M. (2016). Past, future, and present of state-dependent models of diversification. *American*
Journal of Botany, 103(5), 792–795. doi:10.3732/ajb.1600012

Ogburn, R. M., & Edwards, E. J. (2015). Life history lability underlies rapid climate niche evolution in the angiosperm clade
Montiaceae. *Molecular Phylogenetics and Evolution, 92*(1), 181–192. doi:10.1016/j.ympev.2015.06.006

Olson, M. E., Arroyo-Santos, A., & Vergara-Silva, F. (2019). A user's guide to metaphors in ecology and evolution. *Trends in*
Ecology & Evolution, 34(7), 605–615. doi:10.1016/j.tree.2019.03.001

Onstein, R. E., Baker, W. J., Couvreur, T. L. P., Faurby, S., Herrera-Alsina, L., Svenning, J. C., & Kissling, W. D. (2018). To adapt or
go extinct? The fate of megafaunal palm fruits under past global change. *Proceedings of the Royal Society of London.*
Series B, Biological sciences, 285(1880), 20180882. doi:10.1098/rspb.2018.0882

Orr, H. A. (2005). The genetic theory of adaptation: A brief history. *Nature Reviews Genetics, 6*(2), 119–127.
doi:10.1038/nrg1523

Osborn, H. F. (1902). The law of adaptive radiation. *The American Naturalist, 36*(425), 353—363.
doi:www.jstor.org/stable/2453444

Pagel, M. (1994). Detecting correlated evolution on phylogenies: A general method for the comparative analysis of discrete
characters. *Proceedings of the Royal Society of London. Series B, Biological sciences, 225*(1342), 37–45.
doi:10.1098/rspb.1994.0006

Paulus, H. F., & Gack, C. (1990). Pollinators as prepollinating isolation factors: Evolution and speciation in *Ophrys* (Orchidaceae).
Israel Journal of Botany, 39(1-2), 43–79. doi:10.1080/0021213X.1990.10677133

Pillon, Y., & Buerki, S. (2017). How old are island endemics? *Biological Journal of the Linnean Society, 121*(2), 469–474.
doi:10.1093/biolinnean/blx005

Pires, M. M., Silvestro, D., & Quental, T. B. (2017). Interactions within and between clades shaped the diversification of
terrestrial carnivores. *Evolution, 71*(7), 1855–1864. doi:10.1111/evo.13269

Portner, H. O. (2002). Climate variations and the physiological basis of temperature dependent biogeography: Systemic to
molecular hierarchy of thermal tolerance in animals. *Comparative Biochemistry and Physiology a-Molecular and*
Integrative Physiology, 132(4), 739–761. doi:10.1016/S1095-6433(02)00045-4

R Core Team. (2013). R: A language and environment for statistical computing. Vienna, Austria: R foundation for statistical
computing. Retrieved from http://www.R-project.org

Rabosky, D. L. (2013). Diversity-dependence, ecological speciation, and the role of competition in macroevolution. *Annual*
Review of Ecology, Evolution, and Systematics, Vol 44, 44(1), 481–502. doi:10.1146/annurev-ecolsys-110512-135800

Rabosky, D. L. (2014). Automatic detection of key innovations, rate shifts, and diversity-dependence on phylogenetic trees. *PloS*
One, 9(2), e89543. doi:10.1371/journal.pone.0089543.t001

Rabosky, D. L., & Glor, R. E. (2010). Equilibrium speciation dynamics in a model adaptive radiation of island lizards. *Proceedings*
of the National Academy of Sciences of the United States of America, 107(51), 22178–22183.
doi:10.1073/pnas.1007606107

Rabosky, D. L., & Goldberg, E. E. (2015). Model inadequacy and mistaken inferences of trait-dependent speciation. *Systematic*
Biology, 64(2), 340–355. doi:10.1093/sysbio/syu131

Rainey, P. B., & Travisano, M. (1998). Adaptive radiation in a heterogeneous environment. *Nature, 394*(6688), 69–72.
doi:10.1038/27900

Rangel, T. F., Edwards, N. R., Holden, P. B., Diniz, J. A. F., Gosling, W. D., Coelho, M. T. P., . . . Colwell, R. K. (2018). Modeling the
ecology and evolution of biodiversity: Biogeographical cradles, museums, and graves. *Science, 361*(6399), eaar5452.
doi:10.1126/science.aar5452

Revell, L. J. (2012). phytools: An R package for phylogenetic comparative biology (and other things). *Methods in Ecology and*
Evolution, 3(2), 217–223. doi:10.1111/j.2041-210X.2011.00169.x

Revell, L. J. (2014). Ancestral character estimation under the threshold model from quantitative genetics. *Evolution, 68*(3), 743–
759. doi:10.1111/evo.12300

Rohde, R. A., & Muller, R. A. (2005). Cycles in fossil diversity. *Nature, 434*(7030), 208–210. doi:10.1038/nature03339

Rolshausen, G., Davies, T. J., & Hendry, A. P. (2018). Evolutionary rates standardized for evolutionary space: Perspectives on
trait evolution. *Trends in Ecology & Evolution, 33*(6), 379–389. doi:10.1016/j.tree.2018.04.001

Rosenblum, E. B., Sarver, B. A. J., Brown, J. W., Des Roches, S., Hardwick, K. M., Hether, T. D., . . . Harmon, L. J. (2012). Goldilocks
meets Santa Rosalia: An ephemeral speciation model explains patterns of diversification across Ttime scales.
Evolutionary Biology, 39(2), 255–261. doi:10.1007/s11692-012-9171-x

Rundell, R. J., & Price, T. D. (2009). Adaptive radiation, nonadaptive radiation, ecological speciation and nonecological
speciation. *Trends in Ecology & Evolution, 24*(7), 394–399. doi:10.1016/j.tree.2009.02.007

Salzburger, W. (2018). Understanding explosive diversification through cichlid fish genomics. *Nature Reviews Genetics, 19*(11),
705–717. doi:10.1038/s41576-018-0043-9

Sanderson, M. J., & Donoghue, M. J. (1994). Shifts in diversification rate with the origin of Angiosperms. *Science, 264*(5165),
1590–1593. doi:10.1126/science.264.5165.1590

Schiestl, F. P., Ayasse, M., Paulus, H. F., Löfstedt, C., Hansson, B. S., Ibarra, F., & Francke, W. (1999). Orchid pollination by sexual
swindle. *Nature, 399*, 421–422. doi:10.1038/20829

Schluter, D. (2000). *The ecology of adaptive radiation*. Oxford, New York: Oxford University Press.

Schlüter, P. M., Xu, S., Gagliardini, V., Whittle, E. J., Shanklin, J., Grossniklaus, U., & Schiestl, F. P. (2011). Stearoyl-acyl carrier
protein desaturases are associated with floral isolation in sexually deceptive orchids. *Proceedings of the National*
Academy of Sciences of the United States of America, 108(14), 5696–5701. doi:10.1073/pnas.1013313108

Schoener, T. W. (1968). The Anolis lizards of Bimini: Resource partitioning in a complex fauna. *Ecology, 49*(4), 704–726.
doi:10.2307/1935534

Sedeek, K. E. M., Scopece, G., Staedler, Y. M., Schönenberger, J., Cozzolino, S., Schiestl, F. P., & Schlüter, P. M. (2014). Genic
rather than genome-wide differences between sexually deceptive *Ophrys* orchids with different pollinators. *Molecular*
Ecology, 23(24), 6192–6205. doi:10.1111/mec.12992

Sedeek, K. E. M., Whittle, E., Guthörl, D., Grossniklaus, U., Shanklin, J., & Schlüter, P. M. (2016). Amino acid change in an orchid
desaturase enables mimicry of the pollinator's sex pheromone *Current Biology, 26*(11), 1505–1511.
doi:10.1016/j.cub.2016.04.018

Seehausen, O. (2015). Process and pattern in cichlid radiations – inferences for understanding unusually high rates of
evolutionary diversification. *New Phytologist, 207*(2), 304–312. doi:10.1111/nph.13450

Silvestro, D., Antonelli, A., Salamin, N., & Quental, T. B. (2015). The role of clade competition in the diversification of North
American canids. *Proceedings of the National Academy of Sciences of the United States of America, 112*(28), 8684–
8689. doi:10.1073/pnas.1502803112

Simpson, G. G. (1944). *Tempo and mode in evolution*. New York: Columbia University Press.

Simpson, G. G. (1953). *The major features of evolution*. New York: Columbia University Press.

Stadler, T. (2011). Inferring speciation and extinction processes from extant species data. *Proceedings of the National Academy*
of Sciences of the United States of America, 108(39), 16145–16146. doi:10.1073/pnas.1113242108

Stanley, S. M. (1979). *Macroevolution – Pattern and process*. San Francisco, CA, USA: WH Freeman & Co.

Storch, D., & Okie, J. G. (2019). The carrying capacity for species richness. *Global Ecology and Biogeography, 28*(10), 1519–1532.
doi:10.1111/geb.12987

Stroud, J. T., & Losos, J. B. (2016). Ecological opportunity and adaptive radiation. *Annual Review of Ecology, Evolution, and*
Systematics, 47(1), 507–532. doi:10.1146/annurev-ecolsys-121415-032254

Sundaram, M., Donoghue, M. J., Farjon, A., Filer, D., Mathews, S., Jetz, W., & Leslie, A. B. (2019). Accumulation over
evolutionary time as a major cause of biodiversity hotspots in conifers. *Proceedings of the Royal Society B: Biological*
Sciences, 286(1912), 20191887. doi:10.1098/rspb.2019.1887

Tanentzap, A. J., Brandt, A. J., Smissen, R. D., Heenan, P. B., Fukami, T., & Lee, W. G. (2015). When do plant radiations influence
community assembly? The importance of historical contingency in the race for niche space. *New Phytologist, 207*(2),
468–479. doi:10.1111/nph.13362

Tank, D. C., Eastman, J. M., Pennell, M. W., Soltis, P. S., Soltis, D. E., Hinchliff, C. E., . . . Harmon, L. J. (2015). Nested radiations
and the pulse of angiosperm diversification: Increased diversification rates often follow whole genome duplications.
New Phytologist, 207(2), 454–467. doi:10.1111/nph.13491

Title, P. O., & Rabosky, D. L. (2019). Tip rates, phylogenies and diversification: What are we estimating, and how good are the
estimates? *Methods in Ecology and Evolution, 10*(6), 821–834. doi:10.1111/2041-210x.13153

Uyeda, J. C., Zenil-Ferguson, R., & Pennell, M. W. (2018). Rethinking phylogenetic comparative methods. *Systematic Biology,*
67(6), 1091–1109. doi:10.1093/sysbio/syy031

Valente, L. M., Etienne, R. S., & Phillimore, A. B. (2014). The effects of island ontogeny on species diversity and phylogeny.
Proceedings of the Royal Society of London. Series B, Biological sciences, 281(1784), 20133227.
doi:10.1098/rspb.2013.3227

Valente, L. M., Phillimore, A. B., & Etienne, R. S. (2015). Equilibrium and non-equilibrium dynamics simultaneously operate in
the Galápagos islands. *Ecology Letters, 18*(8), 844–852. doi:10.1111/ele.12461

Van Valen, L. (1971). Adaptive zones and the orders of Mammals. *Evolution, 25*(2), 420–428. doi:10.1111/j.1558-
5646.1971.tb01898.x

Van Valen, L. (1973). A new evolutionary law. *Evolutionary Theory, 1*(1), 1–30.

Verboom, G. A., Bergh, N. G., Haiden, S. A., Hoffmann, V., & Britton, M. N. (2015). Topography as a driver of diversification in
the Cape Floristic Region of South Africa. *New Phytologist, 207*(2), 368–376. doi:10.1111/nph.13342

Wagner, C. E., Harmon, L. J., & Seehausen, O. (2012). Ecological opportunity and sexual selection together predict adaptive
radiation. *Nature, 487*(7407), 366—370. doi:10.1038/nature11144

Wagner, C. E., Harmon, L. J., & Seehausen, O. (2014). Cichlid species-area relationships are shaped by adaptive radiations that
scale with area. *Ecology Letters, 17*(5), 583–592. doi:10.1111/ele.12260

Weigelt, P., Steinbauer, M. J., Cabral, J. S., & Kreft, H. (2016). Late Quaternary climate change shapes island biodiversity. *Nature,*
532(7597), 99–102. doi:10.1038/nature17443

Wellborn, G. A., & Langerhans, R. B. (2015). Ecological opportunity and the adaptive diversification of lineages. *Ecology and*
Evolution, 5(1), 176–195. doi:10.1002/ece3.1347

Whittaker, R. J., Triantis, K. A., & Ladle, R. J. (2008). A general dynamic theory of oceanic island biogeography. *Journal of*
Biogeography, 35(6), 977–994. doi:10.1111/j.1365-2699.2008.01892.x

Xu, S., & Schlüter, P. M. (2015). Modeling the two-locus architecture of divergent pollinator adaptation: How variation in *SAD*
paralogs affects fitness and evolutionary divergence in sexually deceptive orchids. *Ecology and Evolution, 5*, 493–502.
doi:10.1002/ece3.1378

Xu, S., Schlüter, P. M., Grossniklaus, U., & Schiestl, F. P. (2012). The genetic basis of pollinator adaptation in a sexually deceptive
orchid. *PLoS Genetics, 8*(8), e1002889. doi:10.1371/journal.pgen.1002889

Xu, S., Schlüter, P. M., Scopece, G., Breitkopf, H., Gross, K., Cozzolino, S., & Schiestl, F. P. (2011). Floral isolation is the main
reproductive barrier among closely related sexually deceptive orchids. *Evolution: International Journal of Organic*
Evolution, 65(1), 2606–2620. doi:10.1111/j.1558-5646.2011.01323.x

Zanne, A. E., Tank, D. C., Cornwell, W. K., Eastman, J. M., Smith, S. A., FitzJohn, R. G., . . . Beaulieu, J. M. (2014). Three keys to
the radiation of angiosperms into freezing environments. *Nature, 506*(7486), 89–92. doi:10.1038/nature12872

1225

1226 Tables

1227 **Table 1.** Estimated effects of abiotic environment (*a*), competitive interactions (*b*), and rate of niche

1228 evolution (*c*), on net diversification rate r_ε (*d*) across the conifer clades (samples size n = 41).

Source	Slope	Std. error	t value	P	partial r^2
Abiotic environment (*a*)	-0.003	0.148	-0.018	0.985	0.000
Competition (*b*)	-0.581	0.127	4.563	<0.001	0.360
Rate of niche evolution (*c*)	0.166	0.181	0.919	0.364	0.022
(Intercept)	0.601	0.160	3.761	<0.001	—

1229

1230 **Table 2.** Summary of EvA components for the four examples detailed above: worldwide conifers

1231 (quantified), Andean *Lupinus*, island radiations and Mediterranean *Ophrys*.

Case	*d*	*a*	*b*	*c*
Conifers	Net diversification rate	Potential area size (0.001 % r^2)	Competition (36 % r^2)	Physiological traits (2.2 % r^2)
Lupinus	Speciation Growth form disparification Genetic differentiation	Extent of páramo Physiographical heterogeneity Flickering connectivity	Low competition (?)	Secondary perenniality
Islands	Diversification Disparification	Island ontogeny Isolation Fragmentation	Competition for resources	Idiosyncratic (?)
Ophrys	Speciation	Fragmentation of range	Pollinators	Flower morphology Floral traits, esp. odours Genetics

1232