Rapid adaptation to elevated extracellular potassium in the

pyloric circuit of the crab, *Cancer borealis*

Abbreviated title: Loss and recovery of circuit activity in high [K+]

Lily S. He*, Mara C.P. Rue*, Ekaterina O. Morozova*, Daniel J. Powell, Eric J. James+,

Manaswini Kar o, and Eve Marder

Biology Department and Volen Center, Brandeis University, Waltham, MA 02454

+ Biology Department, Aldephi University, Garden City, NY 11530

o Department of Neurobiology, University of Pittsburgh, Pittsburgh, PA 15213

* these authors contributed equally to this work

Corresponding author: Eve Marder: marder@brandeis.edu

Number of pages: 42

Number of figures: 6

Abstract: 223 words

Introduction: 625 words

Discussion: 1484 words

The authors declare no conflicts of interest.

Acknowledgments: Supported by NSF Grant *R35 NS097343*. The authors would like to

thank Daniel Shin for assistance with dissections, and Stephen Van Hooser for statistical

advice, and Sonal Kedia and Jason Pipkin for careful reading of this manuscript.

Abstract

Elevated extracellular $[K^+]$ is associated with many disorders including epilepsy, traumatic brain injury, ischemia and kidney failure. Experimentally, elevated $[K^+]$ is used to increase excitability in neurons and networks, by shifting the potassium equilibrium potential (E_K) and consequently, the resting membrane potential. We studied the effects of increased extracellular $[K^+]$ on the well-described pyloric circuit of the crab, *Cancer borealis*, while recording pyloric network activity extracellularly and the activity of Pyloric Dilator neuron (PD) intracellularly. A 2.5-fold increase in extracellular $[K^+]$ ($2.5x[K^+]$) depolarized PD neurons and resulted in an unexpected short-term loss of their normal bursting activity. This period of silence was followed by the recovery of spiking and/or bursting activity during the continued superfusion of $2.5x[K^+]$ saline. In contrast, when PD neurons were pharmacologically isolated from pyloric presynaptic inputs, they exhibited no loss of spiking activity in $2.5x[K^+]$, suggesting the existence of an acute inhibitory effect mediated by circuit interactions. Action potential threshold in PD neurons decreased markedly over the course of exposure to $2.5x[K^+]$ concurrent with the recovery of spiking and/or bursting activity. This study illustrates a case of rapid adaptation to a global perturbation that is influenced by local synaptic connections. Moreover, the complex response of pyloric neurons to elevated $[K^+]$ demonstrates that electrophysiological recordings are necessary to determine how neuronal and circuit activity are affected by altered K^+ concentrations.

Significance Statement

To characterize the sensitivity of a neuronal circuit to global perturbation, we tested the response of the well-described pyloric circuit of the crab stomatogastric ganglion to saline with elevated [K$^+$]. Unexpectedly, a 2.5-fold increase in extracellular [K$^+$] led to a temporary loss of activity in pyloric neurons that is not due to depolarization block. This was followed by a rapid increase in excitability and concurrent recovery of spiking activity within minutes. In contrast, when presynaptic inputs to pyloric neurons were blocked, there was no temporary loss of spiking activity in elevated [K$^+$]. This is a case of rapid adaptation that restores neuronal activity disrupted by global depolarization.

Introduction

Neuronal circuits must be robust to various environmental challenges. This is especially true for central pattern generators (CPGs) that produce essential motor patterns such as breathing, walking, and chewing (Marder and Calabrese, 1996). Maintaining stability over a range of perturbations involves multiple intrinsic and synaptic mechanisms that operate across minutes to days (Von Euler, 1983; Marder and Bucher, 2001; Harris-Warrick, 2010). Complicating matters, both theoretical and experimental evidence suggest that robust CPGs with similar activity patterns can have widely variable underlying cell intrinsic and synaptic conductances (Prinz et al., 2004; Marder and Goaillard, 2006; Schulz et al., 2006; Schulz et al., 2007; Goaillard et al., 2009; Norris et al., 2011; Roffman et al., 2011). These individually variable circuits, nevertheless, must be reliable, and it remains an open question how such circuits respond and adapt to environmental challenges.

In the context of neuronal circuits, global perturbations involve changes to properties of the environment in which the neurons reside and thus may exert a wide influence over most circuit neurons. For instance, changes in temperature and pH will influence cellular function by altering how basic biochemical processes occur. Global perturbations also include changes to the ionic composition of extracellular fluid that then alter the electrochemical driving forces important for neuronal activity. In particular, elevated extracellular potassium concentration ($[K^+]$), is a physiologically relevant depolarizing stimulus associated with a wide array of conditions including thermal stress, epileptic seizures, kidney failure, traumatic brain injury, and stroke (Baylor and Nicholls, 1969; Katayama et al., 1990; Pérez-Pinzón et al., 1995; Jensen and Yaari, 1997; Rodgers et al., 2007; Krishnan and Kiernan, 2009; Morrison et al., 2011;

Arnold et al., 2014; Chauvette et al., 2016). Experimentally, increased extracellular $[K^+]$

is often used to induce changes in neuronal activity. For instance, in rodent

hippocampal cultures, increased $[K^+]$ were used to depolarize neurons and activate

calcium-dependent transcription pathways (Lin et al., 2008; Sharma et al., 2015).

Increased $[K^+]$ is often applied to pre-Bötzinger complex neurons in acute slice

preparations to reestablish rhythmic firing lost in the absence of excitatory inputs

(Ballerini et al., 1999; Panaitescu et al., 2009; Ruangkittisakul et al., 2011; Rybak et al.,

2014). Although our understanding of the basic depolarizing effect of increased $[K^+]$ on

neuronal membrane potential is well defined (Somjen, 1979), the circuit level and long-

term consequences of global changes in $[K^+]$ are less well understood. Here, we

characterized the effects of increased $[K^+]$ on a well described motor circuit of the crab,

Cancer borealis.

The crustacean stomatogastric nervous system (STNS) has been extensively

studied and is a highly advantageous system for the study of fundamental mechanisms

of circuit dynamics, pattern generation, and neuromodulation (Selverston, 1976;

Selverston and Moulins, 1987; Marder and Bucher, 2007; Marder et al., 2014). The

combination of intrinsic variability, well-established connectivity, and production of

behaviorally relevant fictive activity, make the STNS an attractive model to study

underlying network dynamics and robustness in response to a global perturbation

(Selverston and Miller, 1980; Eisen and Marder, 1982; Miller and Selverston, 1982;

Selverston et al., 1982). Previous studies have shown that the pyloric rhythm of the

STNS is extraordinarily robust to changes in both temperature and pH (Tang et al.,

2010; Tang et al., 2012; Soofi et al., 2014; Marder et al., 2015; Haddad and Marder,

2018; Haley et al., 2018; Kushinsky et al., 2019). Extracellular potassium concentrations

can act on all neurons in a circuit simultaneously and affect a multitude of cellular

processes, making it an attractive model with which to study the effects of a global

perturbation on neural circuits (Somjen, 2002; Misonou et al., 2004).

Here, we characterize the effects of elevated $[K^+]$ on the pyloric rhythm and

quantify the time course of these effects. We describe both some expected and

unanticipated effects of the treatment.

Materials and Methods

Animals and dissections

Adult male Jonah Crabs, *Cancer borealis*, (N = 73) were obtained from Commercial

Lobster (Boston, MA) between December 2016 and March 2019 and maintained in

artificial seawater at 10-12˚C in a 12-hour light/dark cycle. On average, animals were

acclimated at this temperature for one week before use. Prior to dissection, animals

were placed on ice for at least 30 min. Dissections were performed as previously

described (Gutierrez and Grashow, 2009). In short, the stomach was dissected from the

animal and the intact stomatogastric nervous system (STNS) was removed from the

stomach including the commissural ganglia, esophageal ganglion and stomatogastric

ganglion (STG) with connecting motor nerves. The STNS was pinned in a Sylgard-

coated (Dow Corning) dish and continuously superfused with 11˚C saline.

Solutions

Physiological *Cancer borealis* saline was composed of 440 mM NaCl, 11 mM KCl, 26

mM $MgCl_2$, 13 mM $CaCl_2$, 11 mM Trizma base, 5 mM maleic acid, pH 7.4-7.5 at 23˚C

(approximately 7.7-7.8 pH at 11˚C). High - 1.5x, 2x, 2.5x, and 3x[K^+] - salines (16.5, 22,

27.5 and 33mM KCl respectively) were prepared by adding more KCl salt to the normal

saline formula. Picrotoxin (PTX) used to block glutamatergic synapses was added to

normal or 2.5x[K^+] saline at a 10^{-5}M concentration for isolated pacemaker experiments

(Marder and Eisen, 1984).

Electrophysiology

Intracellular recordings from somata were performed in the desheathed STG with 10–

30 MΩ sharp glass microelectrodes filled with internal solution (10 mM $MgCl_2$, 400 mM

potassium gluconate, 10 mM HEPES buffer, 15 mM $NaSO_4$, 20 mM NaCl (Hooper et al.,

2015). Intracellular signals were amplified with an Axoclamp 900A amplifier (Molecular

Devices, San Jose). Extracellular nerve recordings were made by building wells around

nerves using a mixture of Vaseline and mineral oil and placing stainless-steel pin

electrodes within the wells to monitor spiking activity. Extracellular nerve recordings

were amplified using model 3500 extracellular amplifiers (A-M Systems). Data were

acquired using a Digidata 1440 digitizer (Molecular Devices, San Jose) and pClamp data

acquisition software (Molecular Devices, San Jose, version 10.5). For identification of

Pyloric Dilator (PD) and Lateral Pyloric (LP) neurons, somatic intracellular recordings

were matched to action potentials on the pyloric dilator nerve (*pdn*), lateral pyloric

nerve (*lpn*) and/or the lateral ventricular nerve (*lvn*).

Elevated [K$^+$] saline application

Prior to all applications of elevated [K$^+$] saline, baseline activity was recorded for 30

minutes in physiological saline. Following the baseline recording, the entire preparation

was superfused with elevated [K$^+$] saline in concentrations ranging from 1.5x to 3x [K$^+$]

for 90 minutes. The preparation was then washed with physiological saline for 30

minutes. Recordings from PD neurons in the isolated pacemaker kernel were made by

superfusing with 10^{-5}M Picrotoxin (PTX) saline until all inhibitory synaptic potentials in

the PD neurons disappeared (for at least 20 minutes). These preparations were then

exposed to 2.5x[K$^+$] PTX saline for 90 minutes, followed by a 30-minute wash in PTX

saline.

Threshold and excitability measurements

To measure the action potential threshold and excitability of PD neurons, two-electrode

current clamp was used to apply slow ramps of current from -4nA to +2nA over 60s.

Resting membrane potential and input resistance were measured during both baseline

conditions and after the application of elevated $[K^+]$ to ensure the integrity of the

preparation (neurons with input resistances <4MΩ were discarded). Three ramps were

performed during baseline at 10-minute intervals, and ramps were performed in

2.5x$[K^+]$ at 5, 10, 20, 30, 40, 50, 60, 70, 80 and 90 minutes after the start of elevated

$[K^+]$ superfusion. After the preparation was returned to physiological saline, three

ramps were performed again at 10-minute intervals. In recordings from the PD neurons

with glutamatergic synapses blocked by PTX, baseline ramps were performed as

described above, followed by three ramps in PTX saline. Preparations were then

superfused with 2.5x$[K^+]$ PTX saline and washed in PTX saline following the same ramp

procedure as above.

Data acquisition and analysis

Recordings were acquired using Clampex software (pClamp Suite by Molecular Devices,

San Jose, version 10.5) and were visualized and analyzed using custom MATLAB

waveform analysis scripts. These scripts were used to detect and measure voltage

response amplitudes and membrane potentials, plot raw recordings and processed data,

generate spectrograms, and perform some statistical analyses.

Spectral analysis

Spectrograms were calculated using the Burg (1967) method for estimation of the power

spectrum density in each time-window. The Burg method (1967) fits the autoregressive

(AR) model of a specified order p in the time series by minimizing the sum of squares of

the residuals. The fast-Fourier transform (FFT) spectrum is estimated using the

previously calculated AR coefficients. This method is characterized by higher resolution

in the frequency domain than traditional FFT spectral analysis, especially for a relative

short time window (Buttkus, 2000). We used the following parameters for the spectral

estimation: data window of 3.2 s, 50% overlap to calculate spectrogram, number of

estimated AR-coefficients p=window/4+1. Before the analysis, voltage traces were low

pass filtered to 5 Hz using a six-order Butterworth filter and down-sampled. PD neuron

burst frequency was calculated as the mean frequency at the peak spectral power in each

sliding window.

Analysis of interspike interval distributions

Intracellular voltage traces were thresholded to obtain spike times. Distributions of

inter-spike intervals (ISIs) were calculated within 2-minute bins. Hartigan's dip test of

unimodality (Hartigan and Hartigan, 1985) was used to obtain the dip statistic for each

of these distributions. This dip statistic was compared to Table 1 in Hartigan and

Hartigan (1985) to find the probability of multi-modality. The test creates a unimodal

distribution function that has the smallest value deviations from the experimental

distribution function. The largest of these deviations is the dip statistic. The dip statistic

shows the probability of the experimental distribution function being bimodal. Larger

value dips indicate that the empirical data are more likely to have multiple modes

(Hartigan and Hartigan, 1985).

Activity pattern plots

For all recordings, we determined the time of spikes over the course of the experiment.

For the recovery time plots, silence was defined as no more than 2 spikes in a 30-second

sliding window. Otherwise, all spike behaviors (even if irregular) were counted as active

spiking.

To more broadly determine the activity pattern of each PD neuron across the

experiment, we analyzed the distribution of inter-spike intervals (ISI) in 2-minute bins

using Hartigan's dip statistic, as described above. If the dip statistic was 0.05 or higher

the neuron was considered to be bursting. If the dip statistic was lower than 0.05 the

neuron was considered to be tonically firing. Neurons with some spikes, but not enough

ISIs to calculate the dip, were classified as sparsely firing. Neurons with no ISIs in the

observed window were classified as silent. We then plotted the activity pattern of the

neuron in these four categories – bursting, tonic, sparse firing and silent – for each PD

neuron across the entire experiment.

Identification of the spike threshold

The spike threshold was identified as the voltage point of the maximum curvature before

the first spike. Specifically, we calculated the first derivative of the voltage (dV/dt) and

defined the spike onset as the point when dV/dt crosses the threshold value of 10

mV/ms.

All electrophysiology analysis scripts are available at the Marder lab GitHub

(https://github.com/marderlab).

Results

Network activity in the pyloric circuit

The entire stomatogastric nervous system (STNS) of the crab *Cancer borealis* was isolated intact from the stomach and pinned in a dish, allowing us to change the composition of the continuously flowing superfused saline (Fig.1A). The stomatogastric ganglion (STG) contains identified neurons that drive the pyloric rhythm that filters food through the animal's foregut. Figure 1B illustrates the stereotypical triphasic pyloric pattern which is comprised of the activity from lateral pyloric (LP), pyloric (PY) and pyloric dilator (PD). The rhythm is recorded extracellularly from motor axons contained in the lateral ventricular nerve (*lvn*) and other nerves. Figure 1C illustrates the connectivity diagram of the pyloric network. The anterior burster (AB) neuron, the intrinsic oscillator that drives the circuit, is strongly electrically coupled to two PD neurons, which burst synchronously with the AB neuron. Together the AB and two PD neurons form the pacemaker kernel of the network, and their coordinated burst of spikes initiates each triphasic cycle (Maynard, 1972). Synaptic connections between neurons in the STG are all inhibitory and are both graded and spike mediated (Graubard et al., 1980; Manor et al., 1997). Rhythmic inhibition from the pacemaker drives bursting activity resulting from post-inhibitory rebound in the LP neuron and PY neurons which reciprocally inhibit each other (Hartline and Gassie, 1979; Selverston and Miller, 1980). As a result, LP bursting and PY bursting compose the second and third phases of the pyloric rhythm respectively.

The pyloric rhythm is disrupted by high extracellular potassium

To test the response of the pyloric rhythm to changes in extracellular $[K^+]$, we

switched from superfusion of normal physiological saline to superfusion of saline with

elevated $[K^+]$ over the STNS while continuously recording the activity of pyloric neurons

extracellularly from the *lvn*. We tested concentrations of $[K^+]$ that were 1.5, 2, 2.5 and 3-

times physiological concentrations to study the dose-dependent responses of pyloric

neurons to changes in extracellular $[K^+]$. When extracellular $[K^+]$ was changed slightly,

to 1.5x the physiological concentration (Fig. 1*D*, N = 4), the pyloric rhythm remained

triphasic and was negligibly affected. When exposed to 2x $[K^+]$ (Fig. 1*E*, N = 20), the

response of pyloric neurons was more variable; in some cases (N = 9/20) there was a

short disruption of the triphasic pyloric rhythm, which was followed by the recovery of

spiking activity.

Higher concentrations of extracellular $[K^+]$ produced more pronounced effects on

pyloric activity. Superfusion of 2.5x $[K^+]$ (Fig. 1*F,* N = 12 extracellular recordings)

reliably and profoundly altered the pyloric rhythm. During the application of 2.5x$[K^+]$

saline, all preparations exhibited a surprising disruption of action potentials from

pyloric neurons, followed by recovery of spiking activity during continued exposure to

2.5x$[K^+]$ saline. Similarly, application of 3x$[K^+]$ saline to the STNS resulted in consistent

cessation of the pyloric rhythm (Fig. 1*G,* N = 5). However, at 3x$[K^+]$, very few of the

preparations recovered consistent activity. Based on these responses, we settled on a

concentration of 2.5 times the control saline K^+ concentration (2.5x$[K^+]$) for further

study, which reliably disrupted the pyloric rhythm and was accompanied by a consistent

recovery of spiking or bursting activity during the continued application of 2.5x$[K^+]$.

The pattern of loss and recovery of pyloric activity in 2.5x$[K^+]$ saline was

consistent across all experiments; however, the precise nature of each neuron's response

cannot be determined from extracellular data alone. Therefore, to obtain more detailed

information on the effects of increased [K$^+$], we recorded intracellularly from pyloric

neurons while superfusing 2.5x[K$^+$] saline over the STNS.

Pyloric neurons PD and LP depolarize and temporarily lose spiking

activity in high extracellular potassium

With intracellular recordings, we saw a marked loss of spiking activity (crash) of

pyloric neurons in response to the application of 2.5x[K$^+$] saline that was consistent with

the previously described extracellular recordings. In the representative example shown

in Figure 2, the PD and LP neurons burst robustly in normal physiological saline (Fig.

2*Ai*). Within a few minutes of the start of 2.5x[K$^+$] saline application, the minimum

membrane potential of the PD and LP neurons depolarized by 15 and 22mV respectively

(Fig. 2*Aii*), which was coincident with a reduction in firing frequency. The activity of

both the LP and PD neurons became more burst-like over the course of the 90-minute

application of 2.5x[K$^+$] saline (Fig. 2*Aiii-iv*) and recovered to normal baseline behavior

when returned to physiological saline (Fig. 2*Av*). The pattern of depolarization and

recovery of spiking can be more clearly depicted by plotting time-condensed voltage

traces for the PD neuron (the response of the LP neuron closely resembles that of the PD

neuron) for the entire 150 minute experiment. In this trace, the membrane potential

depolarizes in 2.5x[K$^+$] saline followed by a loss of spiking activity (Fig. 2*B*). To visualize

spiking behavior over the course of the experiment, we plotted the instantaneous

interspike intervals (ISI) of the PD neuron for the whole experiment on a log scale

(log$_{10}$(ISIs), Fig. 2*C*). All healthy PD neurons in physiological saline have regular

bursting activity that yields a bimodal distribution of ISIs, reflecting the relatively longer

ISI period between bursts and the shorter ISIs of spikes within a burst. Although the

initial depolarization in 2.5x[K$^+$] saline caused a very brief increase in burst and spike

frequency, the increase was immediately followed by a loss of all spiking activity (Fig.

2*C*). Over the course of the 2.5x[K$^+$] saline application, both the PD and LP neuron

recovered rhythmic bursts of action potentials, which is clear from the re-emergence of

two ISI bands (Fig. *2C)*. Bursting activity is suggestive of the re-appearance of slow

membrane potential oscillations. These slow oscillations are best visualized by

spectrograms of the neuron's membrane potential; recovery of bursting activity in

elevated [K$^+$] saline can be seen by the re-appearance of a strong frequency band in the

voltage spectrogram (Fig. *2D*). We then calculated the most hyperpolarized point of the

membrane potential in each burst averaged over five-minute bins for all PD neurons to

determine the overall depolarizing effect of 2.5x[K$^+$] saline. Individual PD neurons

depolarized upon application of 2.5x[K$^+$] saline and remained depolarized throughout

the application with no repolarization of the membrane potential (Fig. 2*E*). Over all

preparations, PD neurons depolarized upon application of 2.5x[K$^+$] saline (Fig. 2*F,*

repeated measures ANOVA, Tukey post-hoc $p < 0.05$, average depolarization after 10

minutes 14.5 ± 3.3mV) After the initial change in the first 10 minutes in 2.5x[K$^+$], the

minimum membrane potentials did not change for the remainder of the elevated [K$^+$]

application ($p > 0.05$). The minimum membrane potential returned to baseline levels

when the preparations were returned to physiological saline (n.s., $p > 0.05$). The

behavior of LP neurons in 2.5x[K$^+$] was very similar to that of PD neurons; LP neurons

depolarized by 16.5 ± 2.9mV after 10 minutes in 2.5x[K$^+$] (n = 5) and remained

depolarized for the duration of the elevated [K$^+$] application.

Variability in the response of PD neurons to 2.5x[K⁺] saline

Although pyloric activity and circuit connectivity is highly conserved across

animals, responses of individual PD neurons to 2.5x[K⁺] saline varied substantially

across preparations. Superfusion of 2.5x[K⁺] saline led to a period of silence in 9 of the

13 PD neurons, with striking variability in the duration of silence and extent of recovery

across animals. Figure 3 shows the responses of four PD neurons from four different

animals to a 90-minute application of 2.5x[K⁺] saline. Across all preparations (n = 13),

the time of silence elicited by 2.5x[K⁺] saline application varied from 1 to 62 minutes

(time of silence was 10.9 ± 5.8 minutes SD). We characterized the recovery of PD

neuron activity by comparing the ISI distributions over time. In some cases, the PD

neurons exhibited only tonic firing activity in 2.5x[K⁺] saline which is reflected by the

presence of a single ISI band (Fig. 3*A, B*). In other cases, PD neurons regained burst-like

activity, which was reflected in the re-emergence of two ISI bands in 2.5x[K⁺] saline

(Fig. 3*C, D*).

In all PD neuron recordings, the variable period of silence upon application of

2.5x[K⁺] saline was followed by the recovery of spiking activity. For all PD neurons, we

calculated the "time to recovery," defined as the length of time between the silencing of

the neuron and when the neuron recovered at least two action potentials in a 30-second

sliding window during the application of 2.5x[K⁺] saline. We were interested in

determining whether aspects of baseline activity of each PD neuron influenced the

neuron's time to recovery. Therefore, for each PD neuron, we calculated the mean

minimum membrane potential during baseline recordings (Fig. 3*E*), the change in

membrane potential upon application of 2.5x[K⁺] saline (Fig. 3*F*) and the baseline

bursting frequency (Fig. 3*G*) and compared these values to the time to recovery for each

corresponding neuron. We found no correlation between any of these values and the

time elapsed until recovery of spiking activity (R^2=0.203, R^2=0.104, R^2=0.012

respectively).

PD neurons in the isolated pacemaker kernel continue spiking in 2.5x[K$^+$]

saline

From the previous experiments, it was unclear to what extent the crash and

recovery of activity in 2.5x[K$^+$] saline was due to presynaptic inputs to PD neurons.

Because the PD and AB neurons receive only glutamatergic input from other pyloric

network neurons, the pacemaker kernel can be studied in isolation from the pyloric

network neurons by superfusing saline with 10^{-5}M picrotoxin (PTX), which blocks

ionotropic glutamatergic synapses in the STG (Fig. 4*A*) (Bidaut, 1980).

The response of PD neurons in the presence of PTX (PTX(+)) to 2.5x[K$^+$] saline

was markedly different from the behavior of PD neurons in control conditions (in the

absence of PTX) (PTX(-)). Figure 4 illustrates a representative example of the responses

of PD neurons to 2.5x[K$^+$] PTX saline; the preparation initially switched from rhythmic

bursting to tonic spiking activity in 2.5x[K$^+$] PTX saline (Fig. 4*Biii*), followed by recovery

of bursting activity that became more pronounced with time (Fig. 4*B iv − vi*). There was

no interruption of (firing) activity upon the superfusion of 2.5x[K$^+$] PTX saline (Fig. 4C)

in PD neurons as compared to 25.x[K$^+$] alone. The recovery of bursting in 2.5x[K$^+$] PTX

saline can also be seen in the emergence of two distinct ISI bands (Fig. 4*D*) and the

emergence of a robust frequency band in the spectrogram of the PD intracellular voltage

trace (Fig. 4*E*).

We further quantified the response of PD neurons to 2.5x[K+] PTX saline by

calculating the mean minimum membrane potential in five-minute bins for each neuron

across the experiment (n = 8, Fig. *4F*). Similar to PD neurons in the intact circuitry, all

PTX(+) PD neurons depolarized in 2.5x[K+] saline (Fig. *4G,* repeated measures ANOVA,

Tukey post-hoc p < 0.05 , average depolarization after 10 minutes 12.6 ± 3.0mV). The

minimum membrane potential of PD neurons in PTX then remained stable during the

application of 2.5x[K+] and returned to baseline levels when returned to physiological

extracellular [K+] (Fig. *4G,* p > 0.05). Importantly, the fact that PTX(+) PD neurons

maintain tonic firing upon the application of 2.5x[K+] saline despite marked

depolarization indicates that the crash observed in PTX(-) PD neurons is unlikely to be

due to depolarization block.

Synaptic inputs alter response of PD neurons to 2.5x[K+] saline

The initial responses of PD neurons to 2.5x[K+] saline application differed in the

presence or absence of PTX. This difference indicates the existence of a circuit-driven

response to elevated [K+]. To quantify this effect, we used values from the Hartigan's dip

test on 2-minute bins of log(ISI) to determine the time that each PD neuron was either

bursting, tonically firing, or silent throughout the experiment and plotted the activity of

each PD neuron over time (see methods).

In the intact circuit, the majority (N = 9 of 13) of PD neurons exhibited a period

of silence following the application of 2.5x[K+] saline, then recovered spiking activity

over a variable amount of time (Fig. *5A*). In 2.5x[K+] PTX saline, PD neurons either

remained active or only briefly went silent, then demonstrated robust recovery of

spiking or bursting activity (Fig. *5B*). The average time of PD silence in 2.5x[K+] saline

was significantly different with the addition of PTX (Fig. 5*C*, Wilcoxon rank-sum test p =

0.013), as was the average time of PD sparse firing between PTX(+) and PTX(-) PD

neurons (Fig. 5*D*, Wilcoxon rank-sum test p = 0.034). In both the time of silence and

sparse firing comparisons, the difference between PTX(+) and PTX(-) PD neurons was

still significant when the largest point in the PTX(-) group was removed. Neither the

average time of tonic firing (Fig. 5*E*, n.s., p = 0.77), nor the average time of burst firing

significantly were significantly different in the presence or absence of PTX (Fig. 5*F*, n.s.,

p = 0.51).

Excitability of PD neurons changes rapidly during exposure to 2.5x[K$^+$]

saline

We wished to see whether periods of silence in high K$^+$ were due to

depolarization block. Therefore, we applied 60 second steady ramps of current from -

4nA to +2nA to PD neurons at several time points during the period of silence elicited

by application of 2.5x[K$^+$] saline. In silent PD neurons, action potentials could always be

induced by injecting positive current, indicating that this period of silence is not due to

depolarization block (Fig. 6A, ramps at 5-minutes and 10-minutes).

To determine the excitability of PD neurons during application of 2.5x[K$^+$], we

repeated the slow current ramps from -4nA to +2nA at 5, 10, 20, 30, 40, 50, 60, 70, 80,

and 90 minutes after the beginning of the 2.5x[K$^+$] saline application in the presence or

absence of PTX. In the representative example shown, there was a clear change in the

number and frequency of spikes elicited in the PD neuron by the current ramp as a

function of time in 2.5x[K$^+$] saline (Fig. 6*A*). In addition, as time in 2.5x[K$^+$] increased,

more spikes were elicited at the same membrane potentials during the current ramp

(Fig. 6*B*).

For each individual PTX(-) preparation (N = 5), spike threshold became more

hyperpolarized in the PD neuron as a function of time in 2.5x[K$^+$] saline (Fig. 6*C*). In

contrast, when the same current ramps were applied to PTX(+) PD neurons (N = 6),

there was no clear trend in spike threshold as a function of time in 2.5x[K$^+$] saline (Fig.

6*D*). Across all PD neurons, we calculated the percent change in spike threshold from

baseline for each current ramp. There was a significant drop in spike threshold over

time for PD neurons in 2.5x[K$^+$] saline between the 5-minute the 60 through 90-minute

time points and also between the 10-minute and the 60 through 90-minute time points

(Fig. 6*F* blue, one-way repeated measures ANOVA, Tukey Post-Hoc test, all p < 0.05).

For most of these neurons the biggest changes in firing rates and spike threshold

occurred during the first 10 minutes of 2.5x[K$^+$] saline application, which is similar to

the time in which most PD neurons recover spiking activity in elevated [K$^+$]. Across all

PTX(+) PD neurons there was no significant percent change in spike threshold over

time in 2.5x[K$^+$] (Fig. 6*G* red, one-way repeated measures ANOVA, Tukey Post-Hoc test,

n.s., all p > 0.05).

To visualize overall changes in PD neuron excitability, we then calculated average

F-I curves for all neurons at the beginning (5-minute ramp) and end (90-minute ramp)

of the 2.5x[K$^+$] saline application. Control PD neurons become more excitable as time in

2.5x[K$^+$] increases; there is a shift in the average F-I curve between the 5- and 90-

minute time points (Fig. 6*F* solid lines, two-way repeated measures ANOVA, p = 0.009).

In addition, at the 5-minute time point in 2.5x[K$^+$], PTX(+) PD neurons are more

excitable than PTX(-) PD neurons (Fig 6*F* purple lines, two-way repeated measures

ANOVA, p = 0.036). This initial difference in excitability is consistent with the fact that

PTX(+) neurons remain active upon application of $2.5x[K^+]$, while PTX(-) PD neurons

typically lose spiking activity. The excitability of PTX(+) PD neurons does not change

over time in $2.5x[K^+]$; there was no significant difference between the 5- and 90-minute

PTX(+) PD neuron F-I curves (Fig 6*F* dashed lines, two-way repeated measures

ANOVA, p = 0.097). In addition, by the end of the $2.5x[K^+]$ application (90-minutes),

there was no significant difference between the F-I curves of PTX(+) and PTX(-) PD

neurons (Fig 6*F* red lines, two-way repeated measures ANOVA, p = 0.375).

Discussion

Surprising inhibitory effect of a depolarizing stimulus

In the STG, the pyloric rhythm is robust to multiple perturbations including changes in temperature, pH and neuromodulatory state (Tang et al., 2010; Tang et al., 2012; Temporal et al., 2014; Hamood et al., 2015; Haddad and Marder, 2018; Haley et al., 2018). Here, we tested whether pyloric neurons are similarly robust to global depolarization through changes to extracellular [K^+]. It is generally assumed that positive current input or depolarization of a neuron's membrane potential will lead to an increase in neuronal activity, and that extreme depolarizations can lead to loss of activity through depolarization block. Instead, we found a type of transient neuronal silence in response to elevated [K^+] that is not due to a depolarization block.

In the pyloric circuit, all local synaptic connections are inhibitory (Eisen and Marder, 1982; Miller and Selverston, 1982). Synaptic transmission in the STG is both graded and spike-mediated, meaning that action potentials are not required for inhibitory synapses to function (Graubard et al., 1980; Manor et al., 1997; Nadim et al., 1997). Therefore, the observed decrease in both bursting and spiking activity in elevated extracellular [K^+] may be caused by global depolarization leading to an increase in graded inhibition that suppresses spiking and bursting activity. In support of this theory, elevated [K^+] does not have the same inhibitory effect on PD neurons with glutamatergic synapses blocked compared to PD neurons with intact synaptic connections. Similarly, in the proprioceptive neurons of the blue crab and the muscle receptor organ of the crayfish, it was recently reported that increased [K^+] also has an inhibitory effect at concentrations not thought to cause a depolarization block (Malloy et al., 2017).

Adaptation to global perturbation

The PD/AB pacemaker unit of the pyloric circuit exhibited rapid adaptation to the disruptive stimulus of increased extracellular $[K^+]$. Although the triphasic pyloric rhythm was not fully restored in $2.5x[K^+]$ saline, PD neurons exhibited rapid changes in excitability over several minutes, which corresponded to the recovery of spiking and, in many cases, bursting activity.

Classical homeostatic plasticity takes place over hours to days and requires changes in intracellular calcium concentrations which then subsequently drive changes in gene expression, leading to cell-intrinsic changes in excitability (Desai et al., 1999; Cudmore and Turrigiano, 2004; Turrigiano, 2012). There is evidence that this mechanism can be induced by changes in extracellular $[K^+]$. Rat myenteric neurons cultured in elevated $[K^+]$ serum for several days exhibit long-lasting changes in Ca^{2+} channel function (Franklin, 1992). Similarly, culturing rat hippocampal pyramidal cells in high $[K^+]$ medium for several days leads to activation of calcium-dependent changes in input resistance and the resting membrane potential that regulates the intrinsic excitability of the neurons (O'Leary et al., 2010). Changes in K^+ channel densities are also associated with homeostatic regulation of neuronal activity. Depolarization of crustacean motor neurons with current pulses for several hours alters K^+ channel densities in a cell-specific manner through a calcium-dependent mechanism (Golowasch et al., 1999). In addition, cerebellar granule cells in which inhibitory GABA receptors are blocked for several days maintain their response to excitatory input by strengthening voltage-independent K^+ conductances (Brickley et al., 2001). Adaptation to global perturbation over several hours to days is well described by models via calcium signals

that influence expression levels of ion channels (O'Leary et al., 2014; O'Leary, 2018).

Pyloric neurons in the STG also exhibit these long-term adaptations; preparations in

which neuromodulatory inputs are removed initially lose rhythmicity and gradually

recover over the course of several days (Thoby-Brisson and Simmers, 1998, 2002;

Luther et al., 2003; Gray and Golowasch, 2016; Gray et al., 2017).

Intriguingly, many cases of rapid adaptation in neuronal circuits are related to

changes in signaling between neurons. At the *Drosophila* neuromuscular junction,

blocking glutamatergic signaling results in long-lasting changes in synaptic strength

within minutes (Frank et al., 2006; Müller and Davis, 2012; Müller et al., 2012). In the

crustacean cardiac ganglion, blocking delayed rectifier K^+ currents led to adaptation of

the circuit and an increase in electrical coupling strength within one hour (Lane et al.,

2016). Although these rapid changes in synaptic strength are often associated with

Hebbian plasticity and learning, rapid alterations in synapses can also lead to a

homeostatic-like maintenance of circuit activity.

The rapid adaptation of PD neurons in elevated extracellular $[K^+]$ is most likely

due to a combination of cell-intrinsic conductance changes and synaptic modulation. In

crustacean motor neurons, cell-intrinsic changes in current densities have been shown

to be modulated by second-messenger kinase pathways activated by global

depolarization (Ransdell et al., 2012). Rapid changes in circuit state could also be

influenced by neuromodulation, as these experiments were conducted with the entire

STNS, including connections to the upstream modulatory ganglia which were also

exposed to the elevated $[K^+]$. In the STNS, the upstream commissural ganglia and the

esophageal ganglion release a wide range of neuromodulators onto the STG that affect

excitability of cells and the pyloric rhythm (Marder and Bucher, 2007; Marder, 2012). In

particular, proctolin released by the MPN and MCN1 neurons endogenously drives the

pyloric pacemaker neuron AB (Hooper and Marder, 1987; Nusbaum and Marder, 1989;

Wood et al., 2000; Stein et al., 2007). Dopamine is also released by modulatory neurons

in the STNS, and exogenous application of dopamine can cause rapid changes in

electrical coupling strength and has been shown to rapidly regulate potassium currents

in both mammalian and invertebrate neurons (Tornqvist et al., 1988; Harris-Warrick et

al., 1998; Gruhn et al., 2005; Rodgers et al., 2013).

Variable responses to similar perturbations

The effects of perturbations are often represented by averaging the experimental

results from a number of individuals, but there are several reasons why this approach is

not always appropriate. Circuits with apparently identical outputs under control

conditions can present distinctly different responses to perturbation due to underlying

differences in network parameters (Tang et al., 2012; Hamood and Marder, 2014;

Haddad and Marder, 2018; Haley et al., 2018; Alonso and Marder, 2019). Indeed, in this

study we observed a wide range of responses in identified PD neurons to the same

experimental conditions, 2.5x[K^+], which may uncover individual differences that are

not evident during control conditions. Within the STG, conductance densities and

strengths of synaptic connections can vary up to five-fold in magnitude between

individuals (Schulz et al., 2006; Schulz et al., 2007; Goaillard et al., 2009; Shruti et al.,

2014; Temporal et al., 2014). Variability in neuronal conductances underlying similar

activity patterns has also been demonstrated across phyla (Swensen and Bean, 2005;

Nelson and Turrigiano, 2008; Tran et al., 2019). In addition, computational modeling of

the pyloric network revealed that multiple combinations of independent circuit

parameters can give rise to functionally identical activity patterns (Goldman et al., 2001;

Prinz et al., 2004; Taylor et al., 2009; Marder et al., 2015). The application of elevated

[K^+] saline to identified STG neurons provides additional evidence that differences in

individual conductance parameters influence circuit responses to global perturbation

(Alonso and Marder, 2019).

Implications for disease states

Hyperkalemia is well documented in many nervous system disorders; however, it

remains unclear how the changes in [K^+] acutely affect neuronal activity. Chronic Kidney

Disease (CKD) leads to increases in serum [K^+] up to three times normal levels, directly

affecting neuronal excitability, and changes in neuronal properties (Krishnan and

Kiernan, 2009; Arnold et al., 2014). Similarly, increased activity of a group of neurons

can increase the extracellular [K^+] in the surrounding tissue (Baylor and Nicholls, 1969;

Kříž et al., 1974) and epileptic seizures and brain trauma can lead to increases in [K^+] in

surrounding brain regions (Moody et al., 1974; Katayama et al., 1990; Silver and

Erecinska, 1994; Fröhlich et al., 2008). These transient changes in extracellular [K^+]

have been tied to long lasting changes in the organization and phosphorylation pattern

of K^+ channels (Misonou et al., 2004), which could lead to long-lasting changes in circuit

function (Somjen, 2001, 2002; Rodgers et al., 2007).

In this study, pyloric neurons respond to elevated [K^+] with a rapid adaptation of

excitability; another example of how the such adaptations occur. The fast adaptation of

pyloric neurons to global changes in [K^+] also suggests that rapid swings in [K^+] may be

more damaging to neurons than gradual shifts or sustained changes in concentration.

Reassessing global perturbation

The results of this study highlight the unexpected complexity of seemingly simple

perturbations. In this classical manipulation of increased extracellular $[K^+]$ we observed

a paradoxical decrease in the activity of PD neurons upon bath application of high $[K^+]$,

followed by a recovery of activity in a short period time. Despite knowing the network

connectivity, circuit properties and behavior of identified neurons within the STG, we

were still unable to predict or fully explain the effects of increased $[K^+]$ on circuit

performance. The complex interaction between circuit level effects and cell intrinsic

responses to simple changes in ion concentrations underscores the importance of

recording and reporting neuronal activity during such manipulations in any experiment.

References

Alonso LM, Marder E (2019) Visualization of currents in neural models with similar behavior and

different conductance densities. eLife 8:e42722.

Arnold R, Pussell BA, Howells J, Grinius V, Kiernan MC, Lin CSY, Krishnan AV (2014) Evidence for

a causal relationship between hyperkalaemia and axonal dysfunction in end-stage

kidney disease. Clin Neurophysiol 125:179-185.

Ballerini L, Galante M, Grandolfo M, Nistri A (1999) Generation of rhythmic patterns of activity

by ventral interneurones in rat organotypic spinal slice culture. J Physiol 517:459-475.

Baylor DA, Nicholls JG (1969) Changes in extracellular potassium concentration produced by

neuronal activity in the central nervous system of the leech. J Physiol 203:555-569.

Bidaut M (1980) Pharmacological dissection of pyloric network of the lobster stomatogastric

ganglion using picrotoxin. J Neurophysiol 44:1089-1101.

Brickley SG, Revilla V, Cull-Candy SG, Wisden W, Farrant M (2001) Adaptive regulation of

neuronal excitability by a voltage- independent potassium conductance. Nature 409:88-

92.

Burg JP (1967) Maximum entropy spectral analysis. In: Proc. 37th Meeting Society of

Exploration Geophysicist. Oklahoma City, OK.

Buttkus B (2000) Spectral Analysis and Filter Theory. Berlin: Springer.

Chauvette S, Soltani S, Seigneur J, Timofeev I (2016) *In vivo* models of cortical acquired epilepsy.

Neurosci Methods 260:185-201.

Cudmore RH, Turrigiano GG (2004) Long-term potentiation of intrinsic excitability in LV visual

cortical neurons. J Neurophysiol 92:341-348.

Desai NS, Rutherford LC, Turrigiano GG (1999) Plasticity in the intrinsic excitability of cortical

pyramidal neurons. Nat Neurosci 2:515.

Eisen JS, Marder E (1982) Mechanisms underlying pattern generation in lobster stomatogastric

ganglion as determined by selective inactivation of identified neurons. III. Synaptic

connections of electrically coupled pyloric neurons. J Neurophysiol 48:1392-1415.

Frank CA, Kennedy MJ, Goold CP, Marek KW, Davis GW (2006) Mechanisms underlying the

rapid induction and sustained expression of synaptic homeostasis. Neuron 52:663-677.

Franklin JL (1992) Long-term Regulation of Neuronal Calcium Currents by Prolonged Changes of

Membrane Potential. J Neurosci 12:1726-1735.

Fröhlich F, Bazhenov M, Iragui-Madoz V, Sejnowski TJ (2008) Potassium dynamics in the

epileptic cortex: new insights on an old topic. The Neuroscientist 14:422-433.

Goaillard J-M, Taylor AL, Schulz DJ, Marder E (2009) Functional consequences of animal-to-

animal variation in circuit parameters. Nat Neurosci 12:1424-1430.

Goldman MS, Golowasch J, Marder E, Abbott L (2001) Global structure, robustness, and

modulation of neuronal models. J Neurosci 21:5229-5238.

Golowasch J, Abbott LF, Marder E (1999) Activity-dependent regulation of potassium currents

in an identified neuron of the stomatogastric ganglion of the crab *Cancer borealis*. J

Neurosci 19:RC33-RC33.

Graubard K, Raper JA, Hartline DK (1980) Graded synaptic transmission between spiking

neurons. Proc Natl Acad Sci USA 77:3733-3735.

Gray M, Golowasch J (2016) Voltage dependence of a neuromodulator-activated ionic current.

eneuro 3.

Gray M, Daudelin DH, Golowasch J (2017) Activation mechanism of a neuromodulator-gated

pacemaker ionic current. J Neurophysiol 118:595-609.

Gruhn M, Guckenheimer J, Land B, Harris-Warrick RM (2005) Dopamine modulation of two

delayed rectifier potassium currents in a small neural network. J Neurophysiol 94:2888-

2900.

Gutierrez GJ, Grashow RG (2009) *Cancer borealis* stomatogastric nervous system dissection.

JoVE (J Vis Exp):e1207.

Haddad SA, Marder E (2018) Circuit robustness to temperature perturbation is altered by

neuromodulators. Neuron 100:609-623. e603.

Haley JA, Hampton D, Marder E (2018) Two central pattern generators from the crab, *Cancer*

borealis, respond robustly and differentially to extreme extracellular pH. eLife 7:e41877.

Hamood AW, Marder E (2014) Animal-to-Animal Variability in Neuromodulation and Circuit

Function. Cold Spring Harbor Symp Quant Biol 79:21-28.

Hamood AW, Haddad SA, Otopalik AG, Rosenbaum P, Marder E (2015) Quantitative

reevaluation of the effects of short- and long-term removal of descending modulatory

inputs on the pyloric rhythm of the crab, *Cancer borealis*. eNeuro 2:0058-0014.

Harris-Warrick RM (2010) General principles of rhythmogenesis in central pattern generator

networks. Prog Brain Research 187:213-222.

Harris-Warrick RM, Johnson BR, Peck JH, Kloppenburg P, Ayali A, Skarbinski J (1998) Distributed

Effects of Dopamine Modulation in the Crustacean Pyloric Network a. Annals of the New

York Academy of Sciences 860:155-167.

Hartigan JA, Hartigan PM (1985) The dip test of unimodality. Ann Stat 13:70-84.

Hartline DK, Gassie DV, Jr. (1979) Pattern generation in the lobster (*Panulirus*) stomatogastric

ganglion. I. Pyloric neuron kinetics and synaptic interactions. Biol Cybern 33:209-222.

Hooper SL, Marder E (1987) Modulation of the lobster pyloric rhythm by the peptide proctolin. J

Neurosci 7:2097-2112.

Hooper SL, Thuma JB, Guschlbauer C, Schmidt J, Büschges A (2015) Cell dialysis by sharp

electrodes can cause nonphysiological changes in neuron properties. J Neurophysiol

114:1255-1271.

Jensen MS, Yaari Y (1997) Role of intrinsic burst firing, potassium accumulation, and electrical

coupling in the elevated potassium model of hippocampal epilepsy. J Neurophysiol

77:1224-1233.

Katayama Y, Becker DP, Tamura T, Hovda DA (1990) Massive increases in extracellular

potassium and the indiscriminate release of glutamate following concussive brain injury.

J Neurosurgery 73:889-900.

Krishnan AV, Kiernan MC (2009) Neurological complications of chronic kidney disease. Nat Rev

Neurol 5:542-551.

Kříž N, Syková E, Ujec E, Vyklický L (1974) Changes of of extracellular potassium concentration

induced by neuronal activity in the spinal cord of the cat. J Physiol 238:1-15.

Kushinsky D, Morozova E, Marder E (2019) *In vivo* effects of temperature on the heart and

pyloric rhythms in the Crab, *Cancer borealis*. J Exp Biol 222:jeb199190.

Lane BJ, Samarth P, Ransdell JL, Nair SS, Schulz DJ (2016) Synergistic plasticity of intrinsic

conductance and electrical coupling restores synchrony in an intact motor network.

eLife 5:e16879.

Lin Y, Bloodgood BL, Hauser JL, Lapan AD, Koon AC, Kim T-K, Hu LS, Malik AN, Greenberg ME

(2008) Activity-dependent regulation of inhibitory synapse development by Npas4.

Nature 455:1198-1204.

Luther JA, Robie AA, Yarotsky J, Reina C, Marder E, Golowasch J (2003) Episodic bouts of activity

accompany recovery of rhythmic output by a neuromodulator-and activity-deprived

adult neural network. J Neurophysiol 90:2720-2730.

Malloy C, Dayaram V, Martha S, Alvarez B, Chukwudolue I, Dabbain N, Mahmood DD, Goleva S,

Hickey T, Ho A (2017) The effects of potassium and muscle homogenate on

proprioceptive responses in crayfish and crab. J Exp Zool 327:366-379.

Manor Y, Nadim F, Abbott L, Marder E (1997) Temporal dynamics of graded synaptic

transmission in the lobster stomatogastric ganglion. J Neurosci 17:5610-5621.

Marder E (2012) Neuromodulation of neuronal circuits: back to the future. Neuron 76:1-11.

Marder E, Eisen JS (1984) Transmitter identification of pyloric neurons: electrically coupled

neurons use different transmitters. J Neurophysiol 51:1345-1361.

Marder E, Calabrese RL (1996) Principles of rhythmic motor pattern generation. Physiol Rev

76:687-717.

Marder E, Bucher D (2001) Central pattern generators and the control of rhythmic movements.

Curr Biol 11:R986-R996.

Marder E, Goaillard JM (2006) Variability, compensation and homeostasis in neuron and

network function. Nat Rev Neurosci 7:563-574.

Marder E, Bucher D (2007) Understanding circuit dynamics using the stomatogastric nervous

system of lobsters and crabs. Annu Rev Physiol 69:291-316.

Marder E, Goeritz ML, Otopalik AG (2015) Robust circuit rhythms in small circuits arise from

variable circuit components and mechanisms. Curr Opin Neurobiol 31:156-163.

Marder E, Goeritz ML, Gutierrez GJ, Hamood A, Brookings T, Caplan J, Haddad S, Kispersky T,

Shruti S (2014) The Crustacean Stomatogastric Nervous System. Nervous Systems and

Control of Behavior 3:337-361.

Maynard DM (1972) Simpler networks. Annals of the New York Academy of Sciences 193:59-72.

Miller JP, Selverston AI (1982) Mechanisms underlying pattern generation in lobster

stomatogastric ganglion as determined by selective inactivation of identified neurons.

IV. Network properties of pyloric system. J Neurophysiol 48:1416-1432.

Misonou H, Mohapatra DP, Park EW, Leung V, Zhen D, Misonou K, Anderson AE, Trimmer JS

(2004) Regulation of ion channel localization and phosphorylation by neuronal activity.

Nat Neurosci 7:711-718.

Moody WJ, Jr. , Futamachi KJ, Prince DA (1974) Extracellular potassium activity during

epileptogenesis. Exp Neurol 42:248-263.

Morrison B, III, Elkin BS, Dollé J-P, Yarmush ML (2011) In vitro models of traumatic brain injury.

Ann Rev Biomedical Engineering 13:91-126.

Müller M, Davis GW (2012) Transsynaptic control of presynaptic Ca2+ influx achieves

homeostatic potentiation of neurotransmitter release. Curr Biol 22:1102-1108.

Müller M, Liu KSY, Sigrist SJ, Davis GW (2012) RIM controls homeostatic plasticity through

modulation of the readily-releasable vesicle pool. J Neurosci 32:16574-16585.

Nadim F, Manor Y, Abbott L, Marder E (1997) Strength and timing of graded synaptic

transmission depend on frequency and shape of the presynaptic waveform. In:

Computational Neuroscience, pp 19-22: Springer.

Nelson SB, Turrigiano GG (2008) Strength through diversity. Neuron 60:477-482.

Norris BJ, Wenning A, Wright TM, Calabrese RL (2011) Constancy and variability in the output of

a central pattern generator. J Neurosci 31:4663-4674.

Nusbaum MP, Marder E (1989) A modulatory proctolin-containing neuron (MPN). I.

Identification and characterization. J Neurosci 9:1591-1599.

O'Leary T, Williams AH, Franci A, Marder E (2014) Cell types, network homeostasis, and

pathological compensation from a biologically plausible ion channel expression model.

Neuron 82:809-821.

O'Leary T (2018) Homeostasis, failure of homeostasis and degenerate ion channel regulation.

Curr Opin Physiol 2:129-138.

O'Leary T, van Rossum MCW, Wyllie DJA (2010) Homeostasis of intrinsic excitability in

hippocampal neurones: dynamics and mechanism of the response to chronic

depolarization: Homeostatic regulation of intrinsic excitability. J Physiol 588:157-170.

Panaitescu B, Ruangkittisakul A, Ballanyi K (2009) Silencing by raised extracellular Ca^{2+} of pre-

Bötzinger complex neurons in newborn rat brainstem slices without change of

membrane potential or input resistance. Neurosci Lett 456:25-29.

Pérez-Pinzón M, Tao L, Nicholson C (1995) Extracellular potassium, volume fraction, and

tortuosity in rat hippocampal CA1, CA3, and cortical slices during ischemia. J

Neurophysiol 74:565-573.

Prinz AA, Bucher D, Marder E (2004) Similar network activity from disparate circuit parameters.

Nat Neurosci 7:1345-1352.

Ransdell JL, Nair SS, Schulz DJ (2012) Rapid homeostatic plasticity of intrinsic excitability in a

central pattern generator network stabilizes functional neural network output. J

Neurosci 32:9649-9658.

Rodgers CI, Armstrong GA, Shoemaker KL, LaBrie JD, Moyes CD, Robertson RM (2007) Stress

preconditioning of spreading depression in the locust CNS. PLoS One 2:e1366.

Rodgers EW, Krenz W-D, Jiang X, Li L, Baro DJ (2013) Dopaminergic tone regulates transient

potassium current maximal conductance through a translational mechanism requiring

D1Rs, cAMP/PKA, Erk and mTOR. BMC Neurosci 14:143.

Roffman RC, Norris BJ, Calabrese RL (2011) Animal-to-animal variability of connection strength

in the leech heartbeat central pattern generator. J Neurophysiol 107:1681-1693.

Ruangkittisakul A, Panaitescu B, Ballanyi K (2011) K^+ and Ca^{2+} dependence of inspiratory-related

rhythm in novel "calibrated" mouse brainstem slices. Respir Physiol Neurobiol 175:37-

48.

Rybak IA, Molkov YI, Jasinski PE, Shevtsova NA, Smith JC (2014) Rhythmic Bursting in the Pre-

Bötzinger Complex. In: Progress in Brain Research, pp 1-23: Elsevier.

Schulz DJ, Goaillard J-M, Marder E (2006) Variable channel expression in identified single and

electrically coupled neurons in different animals. Nat Neurosci 9:356-362.

Schulz DJ, Goaillard JM, Marder EE (2007) Quantitative expression profiling of identified

neurons reveals cell-specific constraints on highly variable levels of gene expression.

Proc Natl Acad Sci USA 104:13187-13191.

Selverston A, Moulins M (1987) Comparison with other systems. In: The Crustacean

Stomatogastric System, pp 307-313. Berlin: Springer.

Selverston AI (1976) Neuronal mechanisms for rhythmic motor pattern generation in a simple

system. In: Neural control of locomotion, pp 377-399. Boston, MA: Springer.

Selverston AI, Miller JP (1980) Mechanisms underlying pattern generation in lobster

stomatogastric ganglion as determined by selective inactivation of identified neurons. I.

Pyloric system. J Neurophysiol 44:1102-1121.

Selverston AI, Miller JP, Wadepuhl M (1982) Local circuits for the generation of rhythmic motor

patterns. Journal de Physiologie 78:748-754.

Sharma N, Gabel Harrison W, Greenberg Michael E (2015) A Shortcut to Activity-Dependent

Transcription. Cell 161:1496-1498.

Shruti S, Schulz DJ, Lett KM, Marder E (2014) Electrical coupling and innexin expression in the

stomatogastric ganglion of the crab Cancer borealis. J Neurophysiol 112:2946-2958.

Silver IA, Erecinska M (1994) Extracellular glucose concentration in mammalian brain:

continuous monitoring of changes during increased neuronal activity and upon

limitation in oxygen supply in normo-, hypo-, and hyperglycemic animals. J Neurosci

14:5068-5076.

Somjen GG (1979) Extracellular potassium in the mammalian central nervous system. Ann Rev

Physiol 41:159-177.

Somjen GG (2001) Mechanisms of Spreading Depression and Hypoxic Spreading Depression-

Like Depolarization. Physiol Rev 81:1065-1096.

Somjen GG (2002) Ion Regulation in the Brain: Implications for Pathophysiology. The

Neuroscientist 8:14.

Soofi W, Goeritz ML, Kispersky TJ, Prinz AA, Marder E, Stein W (2014) Phase maintenance in a

rhythmic motor pattern during temperature changes *in vivo*. J Neurophysiol 111:2603-

2613.

Stein W, DeLong ND, Wood DE, Nusbaum MP (2007) Divergent co-transmitter actions underlie

motor pattern activation by a modulatory projection neuron. Eur J Neurosci 26:1148-

1165.

Swensen AM, Bean BP (2005) Robustness of burst firing in dissociated Purkinje neurons with

acute or long-term reductions in sodium conductance. J Neurosci 25:3509-3520.

Tang LS, Taylor AL, Rinberg A, Marder E (2012) Robustness of a rhythmic circuit to short- and

long-term temperature changes. J Neurosci 32:10075-10085.

Tang LS, Goeritz ML, Caplan JS, Taylor AL, Fisek M, Marder E (2010) Precise temperature

compensation of phase in a rhythmic motor pattern. PLoS Biol 8:e1000469.

Taylor AL, Goaillard J-M, Marder E (2009) How multiple conductances determine

electrophysiological properties in a multicompartment model. J Neurosci 29:5573-5586.

Temporal S, Lett KM, Schulz DJ (2014) Activity-dependent feedback regulates correlated ion

channel mRNA levels in single identified motor neurons. Curr Biol 24:1899-1904.

Thoby-Brisson M, Simmers J (1998) Neuromodulatory inputs maintain expression of a lobster

motor pattern-generating network in a modulation-dependent state: evidence from

long-term decentralization in vitro. J Neurosci 18:2212-2225.

Thoby-Brisson M, Simmers J (2002) Long-Term Neuromodulatory Regulation of a Motor

Pattern–Generating Network: Maintenance of Synaptic Efficacy and Oscillatory

Properties. J Neurophysiol 88:2942-2953.

Tornqvist K, Yang X, Dowling J (1988) Modulation of cone horizontal cell activity in the teleost

fish retina. III. Effects of prolonged darkness and dopamine on electrical coupling

between horizontal cells. J Neurosci 8:2279-2288.

Tran T, Unal CT, Severin D, Zaborszky L, Rotstein HG, Kirkwood A, Golowasch J (2019) Ionic

current correlations are ubiquitous across phyla. Sci Rep 9:1687.

Turrigiano G (2012) Homeostatic synaptic plasticity: local and global mechanisms for stabilizing

neuronal function. Cold Spring Harb Perspect Biol 4:a005736.

Von Euler C (1983) On the central pattern generator for the basic breathing rhythmicity. J Appl

Physiol 55:1647-1659.

Wood DE, Stein W, Nusbaum MP (2000) Projection neurons with shared cotransmitters elicit

different motor patterns from the same neural circuit. J Neurosci 20:8943-8953.

 FIGURE LEGENDS

Figure 1. **The pyloric rhythm is disrupted by large changes in extracellular [K$^+$].** The pyloric rhythm remains robust when exposed to 1.5x[K$^+$] saline and is disrupted at higher [K$^+$]. (A) Diagram of the dissected stomatogastric nervous system (STNS). The entire STNS was superfused with saline with altered [K$^+$]. (B) The triphasic pyloric rhythm is illustrated in an extracellular recording from *lvn,* which contains axons from LP, PY and PD neurons. (C) Connectivity diagram of the pyloric circuit of the crab *Cancer borealis*. (D - G) *lvn* recordings of spiking activity during application of elevated [K$^+$] saline (green boxes) in concentrations of 1.5x, 2.0x, 2,5x and 3.0x the physiological concentration of potassium.

Figure 2. Representative activity of pyloric circuit neurons, pyloric dilator (PD) and lateral pyloric (LP) neurons 2.5x[K$^+$] saline. Green shaded boxes indicate the period of 2.5x[K$^+$] saline superfusion. (A) 3-second segments of PD and LP activity (i-v) are shown in physiological saline, 10, 20 and 70 min into application of 2.5x[K$^+$] saline and upon return to physiological saline. (B) Voltage trace of a PD neuron's activity for the entire representative experiment. (C) Interspike intervals of the PD neuron's activity over the course of 2.5x[K$^+$] saline application. ISIs are plotted on a log scale, with the presence of only one band indicating a tonic firing regime and two bands indicating bursting activity. (D) Spectrogram of the PD neuron's voltage trace. The color code in each spectrogram represents the amplitude density. (E) Mean minimum membrane potential of all PD neurons in five-minute bins for each preparation. (F) Average change in PD neurons' minimum membrane potential

824 compared to baseline in five-minute bins for all preparations. Error bars represent

825 standard deviations. Minimum membrane potential increased significantly when

826 $2.5x[K^+]$ saline was applied (repeated measures ANOVA, Tukey post-hoc $p<0.05$), but

827 did not change significantly between 10 and 90 minutes in $2.5x[K^+]$ (n.s., all $p > 0.05$).

828 The PD minimum membrane potential returned to baseline levels in wash of

829 physiological saline (n.s., baseline compared to wash, all $p > 0.05$).

831 **Figure 3. Responses to $2.5x[K^+]$ saline are highly variable across**

832 **preparations.** Shaded green boxes indicate the period of $2.5x[K^+]$ saline superfusion**.**

833 Example traces of PD neurons from four different preparations (A-D) with a

834 characteristic crash and recovery of activity in $2.5x[K^+]$ saline are shown along with the

835 corresponding log(ISIs) for each preparation. (E) Time to recovery of activity in PD

836 neurons is not correlated to the baseline minimum membrane potential ($R^2 = 0.203$),

837 (F) the change in minimum membrane potential when $2.5x[K^+]$ is applied ($R^2 = 0.104$),

838 or (G) the burst frequency of the PD neuron at baseline conditions ($R^2 = 0.012$).

840 **Figure 4. The representative activity of a PD neuron with presynaptic**

841 **glutamatergic synapses blocked with picrotoxin (PTX) in $2.5x[K^+]$ saline.**

842 (A) Connectivity diagram of the pyloric network with picrotoxin (PTX) blocking

843 glutamatergic signaling. (B) 3-second segments of PD activity (i-vi) in physiological

844 saline, 20 min into application of 10^{-5}M PTX saline, and at 10, 20 and 70 min into

845 application of $2.5x[K^+]$ PTX saline and upon return to physiological saline (C). Voltage

846 trace of the PD neuron over the entire experiment. The blue shaded boxes indicate time

847 of 10^{-5}M PTX saline superfusion, and the green shaded box indicates the 90-minute

period of 2.5x[K^+] PTX saline superfusion. This color scheme is maintained D, F, and G.

(D) ISIs over the course of the experiment plotted on a log scale, with the presence of

one band indicating a tonic firing regime and two bands indicating bursting activity. (E)

Spectrogram of PD neuron's voltage trace. The color code reflects the amplitude density.

(F) Mean minimum membrane potential of all PTX(+) PD neurons in five-minute bins

for each preparation. (G) Average change in PD neurons' minimum membrane potential

in five-minute bins compared to baseline for all preparations. Error bars represent

standard deviations. Minimum membrane potential increases significantly when

2.5x[K^+] PTX saline is applied (repeated measures ANOVA, Tukey post-hoc comparing

baseline to 2.5x[K^+], all $p<0.05$) but does not change significantly after 10 minutes in

2.5x[K^+] PTX for the duration of the elevated [K^+] application (n.s., all $p > 0.05$). The

minimum membrane potential returned to baseline levels in wash of physiological

saline (n.s., baseline compared to wash, all $p > 0.05$).

Figure 5. PD neurons respond differently to 2.5x[K^+] saline in the presence

and absence of PTX. (A) Activity patterns of PD neurons in control conditions

exposed to 2.5x[K^+] saline. (B) Activity patterns of PTX(+)PD neurons exposed to

2.5x[K^+] PTX saline. (C) Control PD neurons exhibit a longer period of silence upon

2.5x[K^+] saline application compared to PD neurons in PTX (Wilcoxon rank-sum test p

= 0.0129). (D) Control PD neurons exhibit a longer period of sparse firing upon 2.5x[K^+]

saline application compared to PD neurons in PTX (Wilcoxon rank-sum test p = 0.034).

(E) Control PD neurons do not show significant differences in the amount of time in

tonic firing (n.s., p = 0.77) or in (F) burst firing upon 2.5x[K^+] saline application

compared to PD neurons in PTX (n.s., p = 0.51).

Figure 6. Action potential thresholds rapidly change in PD neurons during exposure to 2.5x[K$^+$]. (A) Two-electrode current clamp was used to inject current ramps from -4nA to +2nA over 60 seconds. Representative activity during ramps from a PD neuron in 2.5x[K$^+$] at 5, 10, 40, and 80 minutes after the onset of 2.5x[K$^+$] saline application. (B) From the representative neuron shown in A, for each ramp in 2.5x[K$^+$] saline, the average firing frequencies in 5-second bins are plotted against the average membrane potential of the corresponding bin. For C, D and E, green boxes indicate time of 2.5x[K$^+$] saline superfusion, and blue boxes indicate time of PTX superfusion. (C) PD neuron spike thresholds for each ramp are plotted for each preparation (different colors) in the absence of PTX. (D) PD neuron spike thresholds for each ramp plotted for each preparation in PTX. (E) Percent change in the spike threshold over the course of the experiment (as compared to the last baseline ramp) for PD neurons in the presence and absence of PTX. Error bars are SEM, one-way ANOVA with Tukey Post-Hoc test (* = $p < 0.05$, all comparisons between brackets). (F) Average F-I curves for PD neurons (solid lines) and PD neurons in PTX (dashed lines) at five minutes after the onset of 2.5x[K$^+$] saline application (purple lines) and at 90 minutes after the onset of 2.5x[K$^+$] saline application (red lines). There was a significant difference between the F-I curve for control PD neurons between the 5- and 90-minute time points (Two-way repeated measures ANOVA, $p = 0.009$). In addition, there was a significant difference in the F-I curves of PTX(+) and PTX(-) PD neurons after 5 minutes in 2.5x[K$^+$] saline (Two-way repeated measures ANOVA, $p = 0.036$).

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6