Seasonal influenza: Modelling approaches to capture immunity propagation.

Edward M. Hill[1,2*], Stavros Petrou[3], Simon de Lusignan[4,5], Ivelina Yonova[4,5], Matt J. Keeling[1,2,6]

1 Zeeman Institute: Systems Biology and Infectious Disease Epidemiology Research (SBIDER), University of Warwick, Coventry, CV4 7AL, United Kingdom.
2 Mathematics Institute, University of Warwick, Coventry, CV4 7AL, United Kingdom.
3 Warwick Clinical Trials Unit, Warwick Medical School, University of Warwick, Coventry, CV4 7AL, United Kingdom.
4 Department of Clinical and Experimental Medicine, University of Surrey, Guildford, GU2 7XH, United Kingdom.
5 Royal College of General Practitioners, London, NW1 2FB, United Kingdom.
6 School of Life Sciences, University of Warwick, Coventry, CV4 7AL, United Kingdom.

* Corresponding Author. Email: Edward.Hill@warwick.ac.uk

Abstract

Seasonal influenza poses serious problems for global public health, being a significant contributor to morbidity and mortality. In England, there has been a long-standing national vaccination programme, with vaccination of at-risk groups and children offering partial protection against infection. Transmission models have been a fundamental component of analysis, informing the efficient use of limited resources. However, these models generally treat each season and each strain circulating within that season in isolation. Here, we amalgamate multiple data sources to calibrate a susceptible-latent-infected-recovered type transmission model for seasonal influenza, incorporating the four main strains and mechanisms linking prior season epidemiological outcomes to immunity at the beginning of the following season. Data pertaining to nine influenza seasons, starting with the 2009/10 season, informed our estimates for epidemiological processes, virological sample positivity, vaccine uptake and efficacy attributes, and general practitioner influenza-like-illness consultations as reported by the Royal College of General Practitioners (RCGP) Research and Surveillance Centre (RSC). We performed parameter inference via approximate Bayesian computation to assess strain transmissibility, dependence of present season influenza immunity on prior protection, and variability in the influenza case ascertainment across seasons. This produced reasonable agreement between model and data on the annual strain composition. Parameter fits indicated that the propagation of immunity from one season to the next is weaker if vaccine derived, compared to natural immunity from infection. Projecting the dynamics forward in time suggests that while historic immunity plays an important role in determining annual strain composition, the variability in vaccine efficacy hampers our ability to make long-term predictions.

Introduction

As a significant contributor to global morbidity and mortality, seasonal influenza is an ongoing public health concern. Worldwide, these annual epidemics are estimated to result in about three

to five million cases of severe illness, and about 290,000 to 650,000 respiratory deaths [1]. In England, seasonal influenza inflicts a stark burden on the health system during winter periods, being linked with approximately 10% of all respiratory hospital admissions and deaths [2].

Influenza vaccination can offer some protection against seasonal influenza infection for the individual, while contributing to reduced risk of ongoing transmission via establishment of herd immunity [3, 4]. Influenza vaccines are designed to protect against three or four different influenza viruses; two influenza A viruses (an A(H1N1)pdm09 subtype and A(H3N2) subtype) and either one or two influenza B viruses (covering one or both of the B/Yamagata and B/Victoria lineages). In 2013, 40% of countries worldwide recommended influenza vaccination in their national immunisation programmes, although vaccine uptake varies [5–7].

For England (and elsewhere), the need to deploy updated vaccines on an annual basis means influenza vaccination programmes are costly. Yet, predictions of vaccine impact are difficult due to stochasticity in the annual strain composition, the potential misalignment between vaccine and the dominant co-circulating strains and the interaction between multiple seasons. Analysis informing the efficient use of limited resources is therefore vital, with the use of quality-assured analytical models advocated [8].

The previous ten years have seen the fruitful development of influenza transmission models tied to available real-word and experimental data sources [9–15]. Furthermore, combining parameterised transmission models with health economic evaluations permits assessments of changes to vaccination programmes, providing evidence to inform vaccine policy decisions [16, 17]. An eminent study by Baguelin *et al.* [12] connected virological data from England and Wales to a deterministic epidemiological model within a Bayesian inference framework. The transmission model was subsequently interfaced with a health economic analysis model, leading to the recommendation of introducing a paediatric seasonal influenza vaccination programme in the United Kingdom [17]. Subsequent studies have analysed the effect of mass paediatric influenza vaccination on existing influenza vaccination programmes in England and Wales [18], evaluated cost-effectiveness of quadrivalent vaccines versus trivalent vaccines [19], and cost effectiveness of high-dose and adjuvanted vaccine options in the elderly [20].

Prior modelling studies have typically treated each season and each strain circulating within that season independently. Though there are instances of model conceptualisations incorporating waning immunity for natural infection and vaccination, these assumed one generic influenza virus [13], or treated one or both of the influenza A and influenza B types as sole entities rather than explicitly distinguishing between the two common influenza A subtypes and two co-circulating influenza B lineages [9, 11, 16]. Here, we present a multi-strain, non-age structured, susceptible-latent-infected-recovered (SEIR) type transmission model for influenza incorporating a mechanism to link prior season epidemiological outcomes to immunity at the beginning of the following season. Incorporation of a mechanism for the building and propagation of immunity facilitates investigation of the impact of exposure in the previous influenza season, through natural infection or vaccination, on the disease transmission dynamics and overall disease burden in subsequent years. Accordingly, we sought insights to aid understanding of the longer-term dynamics of the influenza virus and its interaction with immunity at the population level.

In this study, we examine the contribution of the differing sources of immunity propagation between years on seasonal influenza transmission dynamics. To this end, we amalgamate multiple sources of epidemiological and vaccine data for England covering the last decade, and fit model outcomes to the available longitudinal data of seasonal rates of general practice (GP) consultations for influenza-like-illness (ILI), scaled by virological surveillance information. We demonstrate that natural infection plays a more prominent role in propagation of immunity to

the next influenza season compared to residual vaccine immunity. We conclude by inspecting forward projections under disparate vaccine efficacy assumptions to determine long-term patterns of seasonal influenza infection.

Methods

Historical data

Throughout, we define a complete epidemiological influenza season to run from week 36 to week 35 of the subsequent calendar year. We chose week 36 (which typically corresponds to the first full week in September) as the start week to match the start of the epidemic with the reopening of schools.

Consultations in General Practices

Since January 1967, the Royal College of General Practitioners (RCGP) Research and Surveillance Centre (RSC) has monitored the activity of acute respiratory infections in GPs. The Weekly Return reports the number of persons in the monitored RCGP RSC network practices registered population consulting for ILI (a medical diagnosis of possible influenza with a set of common symptoms, cough and measured or reported fever $\geq 38^{\circ}$C, with onset within the last 10 days [21]).

Importantly, the dataset is nationally representative both demographically and spatially; the demographics of the sample population closely resembling the country as a whole, with the RCGP RSC network of practices spread across England in order to reflect the distribution of the population. In May 2017, the RCGP RSC network included 1,835,211 patients (approximately 3.3% of the total population in England) from 174 practices [22].

For the influenza seasons 2009/2010 until 2017/2018 inclusive, RCGP RSC provided weekly, age-stratified records containing the size of the monitored RCGP RSC population, the number of individuals in the monitored population consulting for ILI, and ILI rates per 100,000. These data were aggregated across ages to produce weekly ILI rates per 100,000 for the entire sample population. We then summed the weekly ILI rates (weeks 36 through to week 35 in the following calendar year) to produce season-by-season ILI rates per 100,000. Expanded information pertaining to the data extraction is provided in Section 1.1 of the Supporting Information.

Respiratory Virus RCGP RSC Surveillance

To complement syndromic surveillance, a subset of GPs in the RCGP RSC submit respiratory samples from patients presenting in primary care with an ILI. These respiratory samples undergo virological testing to ascertain presence or absence of influenza viruses. Given not all patients reporting ILI are infected with an influenza virus, these virological sample positivity data yield the fraction of ILI reported primary care consultations that were attributable to influenza.

We collected relevant data from figures within Public Health England (PHE) annual influenza reports (for the 2009/10 to 2017/18 influenza seasons inclusive) [23, 24], and PHE Weekly National Influenza reports (for the 2013/14 influenza season onwards) [25]. (For full details, see Section 1.2 of the Supporting Information.) By scaling the longitudinal data of GP consultations

for ILI by the virological surveillance information we garnered an overall estimate of ILI GP consultations attributable to influenza.

Circulating strain composition

We derived seasonal rates of primary care consultations attributable to each of the two influenza A subtypes (A(H1N1)pdm09 and A(H3N2)) and the two influenza B lineages (B/Victoria and B/Yamagata) per 100,000 population. We computed the circulating strain distribution in each influenza season using publicly available data from FluNet [26], a global web-based tool for influenza virological surveillance, using data for the United Kingdom (additional details are provided in Section 1.3 of the Supporting Information).

ILI GP consultations attributable to influenza

We applied the circulating strain composition quantities to the overall estimate of ILI GP consultations attributable to influenza, disaggregating the estimate for all influenza strains of interest into strain-specific quantities:

$$\text{GP consultation rate for strain } m \text{ in season } y = C_{m,y} = \text{GP ILI consultation rate} \times \ldots$$
$$\text{Proportion of ILI samples influenza positive} \times \ldots \quad (1)$$
$$\text{Proportion of influenza viruses in circulation of strain type } m$$

These computations resulted in seasonal rates of GP consultations attributable for the two influenza A subtypes and two influenza B lineages per 100,000 population (Fig. 1(a)). Uncertainty distributions for these values were obtained via bootstrapping given the sample sizes for the three components.

Fig. 1: Empirical, strain-stratified data for ILI GP consultations attributable to influenza per 100,000 population. In both panels, stacked horizontal bars in the left-half depict the cumulative total of ILI GP consultations attributable to type A influenza per 100,000 population (red shading denoting the A(H1N1)pdm09 subtype, orange shading the A(H3N2) subtype). In an equivalent manner, on the right-half of each panel stacked horizontal bars present similar data for type B influenza (cyan shading denoting the B/Victoria lineage, dark blue shading the B/Yamagata lineage). **(a)** Influenza seasons 2009/10 to 2017/18 (inclusive). In the 2009/10 and 2011/12 influenza seasons, all samples identified as influenza B were untyped (represented by black filled bars). **(b)** Influenza seasons 2012/13 to 2017/18 (inclusive), the time span for which we performed parameter inference.

Vaccine uptake and efficacy

For the 2009/10 influenza season, we acquired vaccine uptake profiles from survey results on H1N1 vaccine uptake amongst patient groups in primary care [27]. For the 2010/11 influenza season onward we collated vaccine uptake information from PHE official statistics [24, 25].

We took age adjusted vaccine efficacy estimates for each historical influenza season (2009/2010-2017/2018 inclusive) from publications detailing end-of-season seasonal influenza vaccine effectiveness in the United Kingdom [28–34]. In seasons where equivalent publications were not available, we used mid-season or provisional end-of-season age adjusted vaccine efficacy estimates from PHE reports [35, 36].

There has been substantial variation in vaccine efficacy between seasons and also within seasons across strains, with estimates spanning 23-92% (Table 1). Further, there have been two instances of a mismatch between the vaccine and circulating viruses causing the vaccine to be ineffective (i.e. 0% efficacy) against a particular strain group (B/Yamagata in the 2010/11 influenza season, A(H3N2) in the 2017/18 influenza season). As a sole exception, for the 2009/10 influenza season we used the pandemic vaccine (rather than the seasonal influenza vaccine efficacy) to inform efficacy against A(H1N1)pdm09, with the effectiveness against all other strain types set to zero [28].

The empirical data did not provide individual vaccine efficacy estimates for each influenza A subtype and influenza B lineage (Table S1). We therefore used a set of simplifying assumptions to produce the strain-specific, vaccine efficacy point estimates used within our study (Table 1). The time-varying vaccine uptake data, together with the season-specific information on vaccine efficacy, act as forcing for the within-season epidemiological model (details are described in Sections 1.4-1.5 of the Supporting Information).

Modelling overview

Our motivation was to study the role of immunity propagation in seasonal influenza transmission dynamics by developing and applying a modelling framework incorporating multiple mechanisms linking prior season infection (or vaccination) to immunity in subsequent seasons.

Influenza transmission dynamics are highly complex, with many temporal and structural heterogeneities that will influence the precise pattern of recorded infection. Consequently, our aim was to capture the general trends in the data, such as the pattern of high and low infection levels, rather than the precise values for each influenza season. We, therefore, deliberately chose a parsimonious mechanistic modelling framework, without age-structure, to highlight the impact of immunity propagation.

Following the visualisation convention used in [12], we display a directed acyclic graph representing the model structure and incorporation of the data streams (Fig. 2). The model takes the form of a deterministic continuous-time set of ordinary differential equations (ODEs), which determines the within-season epidemiological dynamics, and a discrete-time map, informing the propagation of immunity from one season to the next.

Our model captures the four strains targeted by the quadrivalent seasonal influenza vaccine, namely, two influenza A subtypes, A(H1N1)pdm09 and A(H3N2), and two influenza B lineages, B/Victoria and B/Yamagata. The multi-strain model construction permits exploration of cross-reactive immunity mechanisms and their impact on the disease dynamics. Multiple infections per influenza season and/or co-infection events were not permitted. In other words, it was presumed

Table 1: Age adjusted, influenza vaccine efficacy point estimates (by season and strain type) All estimates are presented as percentages. The empirical adjusted influenza vaccine efficacy estimates by season and strain type (presented in Table S1) did not provide individual vaccine efficacy estimates for each influenza A subtype and influenza B lineage. We therefore implemented a series of assumptions to produce the strain-specific, vaccine efficacy point estimates used within our study (described in Section 1.5 of the Supporting Information)).

Season	Vaccine efficacy				Source
	A(H1N1)pdm09	A(H3N2)	B/Victoria	B/Yamagata	
2009/10	72.0	0.0	0.0	0.0	[28]*
2010/11	56.0	56.0	78.0	0.0	[29]
2011/12	23.0	23.0	92.0	92.0	[30]
2012/13	73.0	26.0	51.0	51.0	[31]
2013/14	61.0	61.0	61.0	61.0	[35]†
2014/15	29.9	29.3	46.3	46.3	[32]
2015/16	54.5	54.5	57.3	54.2	[33]
2016/17	36.7	31.6	54.5	58.5	[34]
2017/18	66.3	0.0	24.7	24.7	[36]‡

*: The adjusted seasonal influenza vaccine efficacy in the 2009/10 season was -30% (-89%, 11%) [28]. We therefore used the pandemic vaccine to inform efficacy against A(H1N1)pdm09, with the effectiveness against all other types set to zero.
†: Mid-season estimate of seasonal influenza vaccine effectiveness from [35]. Low incidence throughout the 2013/2014 influenza season meant reliable end-of-season estimates for the vaccine efficacy could not be attained (Personal communication, PHE).
‡: Provisional end-of-season influenza vaccine effectiveness results.

that individuals may only be infected by one strain of influenza virus per season, analogous to natural infection eliciting short-term cross immunity to all other strain types [37].

Next, we summarise, in turn, the vaccination, immunity propagation, epidemiological and observation model components.

Vaccination model

Epidemiological states were compartmentalised based on present season vaccine status (indexed by N for non-vaccinated or V for vaccinated). We obtained weekly time-varying vaccine uptake rates, ν, at the population level by computing a weighted average of the individual age group uptake values, thereby accounting for the population distribution in each given influenza season. We assumed the rate of vaccination ν to be constant over each weekly period. Vaccine efficacy depended upon the extent to which the strains in the vaccine matched the circulating strain in that influenza season (Table 1).

We assumed a 'leaky' vaccine, offering partial protection to every vaccinated individual, thus acting to reduce the overall susceptibility of the given group receiving vaccination. The 'leaky' vaccine action contrasts with an 'all-or-nothing' vaccine assumption, which assumes complete protection to a subset of the vaccinated individuals but no protection in the remainder of vaccinated individuals [38]. Explicitly, under a 'leaky' vaccine action with a vaccine efficacy against strain m of α_m, the relative susceptibility of the vaccinated group towards strain m is $1 - \alpha_m$. Additionally, we assumed those administered vaccines had unmodified transmission (i.e. those

Fig. 2: Schematic showing the links between the vaccination, immunity propagation, epidemiological and observation model components. We adopt the visualisation conventions of [12], with ellipses indicating variables, and rectangles indicating data. Dotted arrows indicate relationships between prior season epidemiological outcomes and immunity propagation factors. Solid arrows indicate within-season processes. Circled capitalised letters indicate the relationships connecting the variables or data involved. These relationships are: process A, propagation of immunity as a result of exposure to influenza virus in the previous influenza season (through natural infection or vaccination); process B, modulation of current influenza season virus susceptibility; process C, estimation of influenza case load via the SEIR model of transmission; process D, ascertainment of cases through ILI recording at GP.

receiving the 'leaky' vaccine were equally as infectious as those who did not). Vaccination is 165
modelled irrespective of infection status (susceptible, latent, infected and recovered individuals), 166
as vaccine history impacts the immunity levels for next season. 167

Immunity propagation model 168

A novel aspect of our model framework was the potential for exposure to influenza virus in 169
the previous influenza season, through natural infection or vaccination, to modulate current 170
influenza season susceptibility. Accordingly, we tracked immunity derived from natural infection 171
and vaccination separately (Fig. 2, process A), requiring ten distinct exposure history groupings. 172
Three susceptibility modifying factors of interest were: (i) modified susceptibility to strain m 173
given infection by a strain m type virus the previous season, denoted a; (ii) carry over cross- 174
reactivity protection between influenza B lineages, denoted b (to account for infection with one 175
influenza B virus lineage being potentially beneficial in protecting against subsequent infection 176
with either influenza B virus lineage [39]); (iii) residual strain-specific protection carried over 177
from the prior season influenza vaccine, denoted c_m. We mandated that $0 < a,b,c_m < 1$. We 178
let $f(h,m)$ denote, for those in exposure history group h, the susceptibility to strain m. The 179
collection of ten exposure history groupings and associated strain-specific susceptibilities were 180

consolidated into a single susceptibility array (Fig. 2, process B; Fig. 3). 181

We assumed the immunity propagation due to vaccination in the previous season (c_m) to be 182 associated to the strain-specific vaccine efficacy in the previous season (α_m^{y-1}). In particular, we 183 introduce a linear scaling factor $\xi \in (0, 1)$, which reduces the level of vaccine derived immunity 184 between seasons: $c_m^y = 1 - \xi \alpha_m^{y-1}$. 185

On a related note, we applied a further assumption in order to parameterise susceptibility values 186 amongst exposure groups capturing those both vaccinated and naturally infected in the prior 187 season (Fig. 3: rows 7-10). For this collection of exposure histories, susceptibility to a subset of 188 strains may conceivably be collectively modified via natural infection and vaccination immunity 189 propagation pathways. In these instances, we treated the immunity propagation mechanisms 190 independently, with the modified susceptibility set by the dominant immunity propagation entity 191 (i.e. $\min(a, c_m)$, $\min(b, c_m)$). In other words, we took a pessimistic stance by assuming no 192 boosting of the immunity propagation response as a result of dual influenza virus exposure 193 (from both natural infection and vaccination). 194

All three propagation parameters (a, b, ξ) were inferred from epidemiological data. A mathe- 195 matical description of the immunity mapping between seasons is given in Section 2.2 of the 196 Supporting Information. 197

In light of there being uncertainty around the precise time scale for which immunity to seasonal 198 influenza viruses may be retained, with individual- and population-level models having estimated 199 infection acquired immunity to wane over a timescale of two to ten years [40–42]), we also 200 considered an extended variant of the immunity propagation model component. 201

In the model extension, we fit an additional parameter (δ) representing the proportion of those 202 who began the influenza season in an exposure history group linked to natural infection (Fig. 3: 203 rows 2-5, 7-10), and who were also unexposed to influenza virus during the current season (who at 204 the end of of the current influenza season remained susceptible), that retained their pre-existing 205 natural infection acquired immunity. Those keeping immunity arising from natural infection 206 were mapped to the relevant prior infection exposure history group dependent upon current 207 season vaccination status. The remaining proportion $(1 - \delta)$ transitioned in the same manner 208 as in the original model, reverting to either the naive exposure history group or vaccinated only 209 exposure history group (Figure S20). For clarity, we did not introduce a mechanism to confer 210 vaccine-induced immunity beyond one influenza season (i.e. vaccine-induced immunity could be 211 retained for, at most, a single additional influenza season). 212

With the more complex immunity structure, we did not gain noticeable improvements in cor- 213 respondence of model outputs with the data (relative to fits with a model using the simpler 214 immunity propagation setup). Therefore, we do not discuss any further here the alternative 215 immunity propagation structure, instead giving a complete description of the extended model 216 and associated outcomes in Section 5 of the Supporting Information. 217

Epidemiological model 218

The within-season transmission dynamics were performed by an ODE epidemiological model 219 based on SEIR-type dynamics. At the start of each season we passed into the epidemiological 220 model the vaccination attributes (uptake and efficacy), immunity considerations, virus trans- 221 mission rates and the initial proportion of the population infectious per strain. These collection 222 of epidemiological inputs act as forcing parameters (Fig. 2, process C). 223

<div style="text-align:center">8</div>

Exposure history (h)	Strain susceptibility			
	A(H1N1)pdm09	A(H3N2)	B/Victoria	B/Yamagata
Naïve	1	1	1	1
A(H1N1)pdm09	a	1	1	1
A(H3N2)	1	a	1	1
B/Yamagata	1	1	a	b
B/Victoria	1	1	b	a
Vacc. (V)	$c_{A(H1N1)}$	$c_{A(H3N2)}$	$c_{B/Victoria}$	$c_{B/Yamagata}$
A(H1N1)pdm09 & V	$\min(a, c_{A(H1N1)})$	$c_{A(H3N2)}$	$c_{B/Victoria}$	$c_{B/Yamagata}$
A(H3N2) & V	$c_{A(H1N1)}$	$\min(a, c_{A(H3N2)})$	$c_{B/Victoria}$	$c_{B/Yamagata}$
B/Victoria & V	$c_{A(H1N1)}$	$c_{A(H3N2)}$	$\min(a, c_{B/Victoria})$	$\min(b, c_{B/Yamagata})$
B/Yamagata & V	$c_{A(H1N1)}$	$c_{A(H3N2)}$	$\min(b, c_{B/Victoria})$	$\min(a, c_{B/Yamagata})$

Fig. 3: Infographic presenting the interaction between exposure history and susceptibility.
The interaction between exposure history h and susceptibility to strain m in the current season, $f(h, m)$, was classified into ten distinct groups: One group for the naive (uninfected and not vaccinated, row 1); one group per strain, infected but not vaccinated (rows 2-5); one group for those vaccinated and experiencing no natural infection (row 6); one group per strain for being infected and vaccinated (rows 7-10). We let a denote modified susceptibility to strain m given infection by a strain m type virus the previous season (dark green shading), b modified susceptibility due to cross-reactivity between type B influenza lineages (dark blue shading), and c_m the change in susceptibility to strain m given vaccination in the previous season (gold shading). Unmodified susceptibilities retained a value of 1 (red shading). We enforced $0 < a, b, c_m < 1$.

The epidemiological model is a deterministic, non-age, multi-strain structured compartmental based model capturing demography, influenza infection status (with susceptible-latent-infected-recovered, SEIR, dynamics) and vaccine uptake. For influenza, it is commonplace to also consider a risk-stratified population, divided into individuals at low or high risk of complications associated with influenza. However, it is health outcomes, rather than epidemiological processes, that are contingent upon risk status [12] and therefore this additional structure can be ignored for the purposes of this model. We assumed disease transmission to be frequency-dependent, such that fundamental quantities (like the basic reproductive ratio, R_0) are unaffected by changes to population size.

We assumed exponentially-distributed latent and infectious periods (arising from rates of latency loss and recovery, respectively, being constant). Loss of latency rates $\gamma_{1,m}$ were strain-dependent. From Lessler *et al.* [43], we chose rates corresponding to average latent periods of 1.4 days and 0.6 days for influenza A and influenza B associated strains respectively. Following Cauchemez *et al.* [44], we set the rate of loss of infectiousness $\gamma_2 = 1/3.8$, corresponding to an average infectious period of 3.8 days, independent of strain.

We set the mortality rate based upon a gender-averaged life expectancy for England, approximated from male and female specific statistics from Office for National Statistics data [45] (see Table 2).

The ODE equations of the SEIR epidemiological model are given in Section 2.3 of the Supporting Information. The strain specific force of infection, λ_m, satisfies

$$\lambda_m = \beta_m \sum_X I_m^X, \qquad (2)$$

Table 2: Overview of parameters in the model.

Description	Notation	Value	Sources
Fixed parameters			
Mortality rate (day^{-1})	B,D	$\frac{1}{81\times365}$	[45]
Rate of latency loss, influenza A subtypes (day^{-1})	$\gamma_{1,A}$	$\frac{1}{1.4}$	[43]
Rate of latency loss, influenza B lineages (day^{-1})	$\gamma_{1,B}$	$\frac{1}{0.6}$	[43]
Recovery rate (day^{-1})	γ_2	$\frac{1}{3.8}$	[44]
Time-varying parameters			
Vaccination rate at time t	$\nu(t)$	—	[24, 25, 27]
Vaccine efficacy, season y strain m	α_m^y	—	[28–36]

Inferred parameter description	Notation	Prior	
Influenza virus transmissibility, strain m	β_m	$\mathcal{U}(0.2632,0.7896)$	
Modified susceptibility given natural infection in prior season	a	$\mathcal{U}(0,1)$	
Modified susceptibility due to type B influenza cross-reactivity	b	$\mathcal{U}(0,1)$	
Proportion of prior season vaccine efficacy carried over	ξ	$\mathcal{U}(0,1)$	
Ascertainment probability in season y	ϵ_y	$\mathcal{U}(0,0.05)$	

where β_m is the transmission rate for strain m, implicitly comprising the contact rate and the transmissibility of the virus (the probability that a contact between an infectious person and a susceptible person leads to transmission), and the superscript $X \in \{N,V\}$ denotes vaccination status (indexed by N for non-vaccinated, V for vaccinated).

For use in our observation model, we tracked the incidence $Z_m(y)$ of new strain m influenza infections in season y as a rate per 100,000 population:

$$Z_m(y) = \left(\int_{y-1}^{y} \gamma_{1,m}(E_m^N + E_m^V)\, \mathrm{d}t \right) \times 100,000. \tag{3}$$

Observation model

As discussed above, individuals consulting a GP for ILI do not necessarily directly correspond with individuals infected with one of the circulating influenza strains. To address this discrepancy, the final segment of the mathematical model linked the GP consultation data with the number of infections due to circulating influenza viruses in the population (Fig. 2, process D).

Our interest resided in the number of ascertainable cases, where we assumed that each individual infected by the strain of influenza under consideration had a probability ϵ of being ascertainable, i.e. going to the GP, being recorded as having ILI, and having a detectable influenza viral load. We assumed the ascertainment probability was season-specific (but strain agnostic) to account for a range of influences such as climate or co-circulating infections. We obtained model estimates of the proportion of individuals experiencing cases of ascertainable influenza (of strain type m)

in season y through scaling the incidence $Z_m(y)$ by the ascertainment probability in influenza season y, ϵ_y. Consequently, the ascertainable influenza cases in season y, $Z_m^+(y)$, obeys:

$$Z_m^+(y) = \epsilon_y Z_m(y). \qquad (4)$$

The parameter ϵ_y is inferred by comparing the total predicted incidence in a given year to the calculated GP consultation rate (Eq. (1)).

Parameter inference

To realise a model capable of generating influenza-attributed ILI GP consultations estimates that resemble the empirical data (Eq. (1)), we sought to fit the following collection of model parameters (Table 2): transmissibility of each influenza virus strain (β_m), modified susceptibility to strain m given infection by a strain m type virus the previous season (a), carry over cross-reactivity protection between influenza B lineages (b), residual protection carried over from the prior season influenza vaccine (ξ), and an ascertainment probability per influenza season (ϵ_y). One may notice the ascertainment probabilities were the sole group of parameters that could vary between influenza seasons, and we acknowledge that one may feasibly apply temporal (influenza season) dependencies to the transmission rates and immunity propagation parameters. Nevertheless, despite the the integration of additional parameters potentially improving the fit to the data, we would be running the risk of the model containing more parameters than can be justified by the data (overfitting). Instead, our intention was to extract a signal across the considered time frame, if present, identifying the most prominent transmission rates (comparing across the four influenza viruses) and immunity propagation mechanisms.

For inferring posterior parameter distributions, we employed an adaptive-population Monte Carlo approximate Bayesian computation (ABC) algorithm [46] combined with a local perturbation kernel (multivariate normal kernel with optimal local covariance matrix) [47]. Prior distributions were uniform for all parameters (Table 2).

Simulations began in the 2009/2010 influenza season and ran for a specified number of seasons with one parameter set. For the 2009/2010 influenza season, only the A(H1N1)pdm09 strain was present in our simulation (in accordance with the strain composition data where proportions of samples for the remaining three strains were very low, Fig. 1(a)), with the initial proportion of the population infected being 1×10^{-5} (one per 100,000). In all subsequent influenza seasons, with all four influenza virus strains being present, the initial proportion of the population infected by each strain was 2.5×10^{-6}. We fit to the 2012/13 season onward, which omitted the influenza seasons (2009/10 and 2011/12) that had no influenza B lineage typing data (Fig. 1). This selected time frame contained a regime in which the dominant circulating influenza A subtype generally switched annually between H1N1(2009) and H3N2, while for type B influenza the Yamagata lineage incidence typically exceeded Victoria lineage incidence.

The summary statistics for our ABC procedure were comprised of two parts. The first component was a within-season temporal profile check; the peak in influenza infection (combining those in latent and infectious states) could not occur after February in any season. The second component defined the metric to measure the correspondence of the model predicted influenza-attributed ILI GP consultations versus the observed data. We worked with a metric akin to Poisson deviance. Compared to a sum of squared residuals error metric, a measure akin to Poisson deviance penalises with greater severity poor fits to data points of small magnitude. Accounting

for season y and strain stratification m, we defined our deviance measure to be

$$\mathrm{DEV} = 2 \sum_y \sum_m \left(C_{m,y} \ln \left(\frac{C_{m,y}}{M_{m,y}} \right) - (C_{m,y} - M_{m,y}) \right), \tag{5}$$

with $C_{m,y}$ the observed value for strain m in season y, and $M_{m,y}$ the model estimate for strain m in season y.

We amassed 10,000 particles representing a sample from the posterior distribution (see Section 3 of the Supporting Information for expanded details on the parameter estimation methodology).

Forward simulations

We used the parameter set returning the lowest error (in the inference procedure) to explore potential strain dynamics up to the 2029/30 influenza season. Vaccination attributes (uptake and efficacy) for the 2009/10 to 2017/18 influenza seasons were taken from the observed data. For the forecasted seasons (2018/19 influenza season to the end of each simulation), we studied four scenarios arising from different hypotheses concerning future vaccine efficacy: (i) randomly sampled from all previous vaccine efficacy values (1,000 simulation replicates); (ii) 'expected' scenario (single simulation), strain-specific vaccine efficacies set at the median attained efficacy across 2010/11-2017/18 seasonal influenza vaccines (note, efficacy estimates from 2009/10 influenza season were not included as for 2009/10 we used pandemic influenza vaccine efficacy estimates); (iii) pessimistic scenario (single simulation), strain-specific vaccine efficacies set at the minimum attained efficacy across 2010/11-2017/18 seasonal influenza vaccines; and (iv) optimistic scenario (single simulation), strain-specific vaccine efficacies set at the maximum attained efficacy across 2010/11-2017/18 seasonal influenza vaccines. In addition, vaccine uptake in all projected influenza seasons matched that of the 2017/18 influenza season.

Simulation and software specifics

All model simulations began at week 36 of the epidemiological season. Inference and model simulations were performed in JULIA v0.6, with the system of ODEs solved numerically with the package DifferentialEquations v2.2.1. Data processing and production of figures were carried out in MATLAB R2019a.

Results

Parameter inference

We invoked the adaptive-population Monte Carlo ABC algorithm, accumulating 10,000 parameter sets (representing a sample from the posterior distribution) to determine credible values of the model parameters (Table 3). Samples were obtained after completion of 600 generations of the adaptive-population Monte Carlo ABC scheme, at which point the generation-by-generation tolerance level updates were minor (Fig. 4(a)). Accordingly, additional simulations would only marginally change the posterior distributions. The model parameters were well defined, with Gaussian-shaped histograms (Fig. 4(b)). All parameter sets adhered to the mandatory temporal criteria of peak infection in each influenza season occurring prior to March (Figure S5).

Table 3: Values (posterior weighted quantiles) for the transmission, exposure history and ascertainment probability parameters inferred fitting to the empirical data. Numbers inside brackets indicate 95% credible intervals. All values are given to 4 d.p.

Description	Notation	Median [95% credible interval]
Transmission parameters		
A(H1N1)pdm09	$\beta_{\text{A(H1N1)pdm09}}$	0.3913 [0.3900, 0.4041]
A(H3N2)	$\beta_{\text{A(H3N2)}}$	0.3917 [0.3917, 0.4074]
B/Victoria	$\beta_{\text{B/Victoria}}$	0.3510 [0.3510, 0.3606]
B/Yamagata	$\beta_{\text{B/Yamagata}}$	0.3653 [0.3653, 0.3759]
Exposure history parameters		
Natural infection in prior season	a	0.7883 [0.7847, 0.7964]
Type B influenza cross-reactivity	b	0.9703 [0.9558, 0.9988]
Prior season vaccine efficacy propagation	ξ	0.0051 [0.0004, 0.0143]
Ascertainment probabilities		
2012/13	$\epsilon_{2012/13}$	0.0017 [0.0015, 0.0018]
2013/14	$\epsilon_{2013/14}$	0.0009 [0.0009, 0.0012]
2014/15	$\epsilon_{2014/15}$	0.0024 [0.0022, 0.0026]
2015/16	$\epsilon_{2015/16}$	0.0037 [0.0032, 0.0039]
2016/17	$\epsilon_{2016/17}$	0.0014 [0.0012, 0.0015]
2017/18	$\epsilon_{2017/18}$	0.0055 [0.0047, 0.0055]

Inspecting the attained posterior distributions for virus transmissibility, estimates for the two influenza A subtypes are similar; both are larger than the corresponding estimates for the two type B lineages. However, the B/Yamagata transmissibility estimates exceeded those for B/Victoria, although the values are relatively close.

Amongst the three exposure history parameters (a, b, ξ), the parameter having a notable impact on the transmission dynamics was the modification to susceptibility due to natural infection in the previous influenza season (a). If infected by a given strain last season, the inferred weighted median estimate of 0.7883 corresponds to an approximate 21% reduction in susceptibility to the current season variant of that strain type. Strikingly, there was little carry over of prior season vaccine efficacy (ξ), with the majority of the posterior distribution massed near 0. Similarly, being previously infected by one of the type B influenza virus conferred little immunity against the remaining type B lineage virus in the next season.

Inferred ascertainment probabilities (ϵ), across all seasons, were typically within the range of 0.001-0.006. The highest ascertainment probabilities were found for the recent 2017/18 influenza season, attaining a median of 0.005; but this still suggests that only one in two hundred infections was reported to their GP and correctly identified as influenza.

We verified model robustness by performing parameter fits using two subsets of the historical influenza season data, spanning the time periods 2012/13-2015/16 and 2012/13-2016/17 respectively. Fitting to these differing influenza season ranges, outcomes generally exhibited qualitative consistency; in particular, amongst the three mechanisms for immunity propagation, current season susceptibility was modulated greatest by natural infection in the previous influenza season. The only notable quantitative differences when fitting to the two shortened

time frames (compared to the distributions inferred when fitting to the complete time period) were elevated transmissibility levels (β), counteracted by an enlarged effect of prior infection (a) and vaccination propagation (ξ); further details are given in Section 4.2 of the Supporting Information.

In addition, we assessed the parameter identifiability competency of our ABC fitting scheme via generation of synthetic data from known parameters. Employing our inference scheme to fit the model to the synthetic data, we ably recovered the parameter values from which the synthetic data had been generated, giving extra confidence to our results (see Section 4.3 of the Supporting Information).

(a)

(b)

Fig. 4: Results of the ABC scheme, fitting to the empirical data. (a) Summary metric threshold value upon completion of each generation of the inference scheme. The inset panel displays the latter quarter of generations. **(b)** Inferred parameter distributions estimated from 10,000 retained samples following completion of 600 generations of the inference scheme. Vertical red lines indicate the (non-weighted) median values for the model constants estimated from the inference procedure. Particularly noteworthy outcomes include: transmissibility of type A viruses exceeding type B viruses; prior season influenza B cross-reactivity and vaccine carry over had little impact on present season susceptibility; the highest ascertainment probability occurred in the 2017/18 influenza season.

Evaluating model fit

To assess the goodness-of-fit between our model and the available data, we performed 1,000
independent simulations using parameter sets drawn from the ABC inference procedure. There-
fore, although the model is deterministic, we generated variability in epidemic composition due
to the posterior distribution for the underlying parameters. The incidence of each of the four
influenza types over each season was converted to attributed GP consultations (per 100,000),
with both incidence and GP consultation outputs compared to the data (Fig. 5). We generally
obtained a reasonable fit to the data, both in terms of the severity of infection each year (as
controlled by the ascertainment ϵ_y) but also in terms of type and subtype composition (which
is largely driven by epidemiological dynamics and propagation of immunity).

Considering type A influenza, we clearly captured the alternating pattern of dominance by
A(H3N2) and A(H1N1)pdm09 subtypes for the years 2012/13 to 2016/17; but more importantly
the model was able to predict the unexpected result that A(H3N2) dominated in both 2016/17
and again in 2017/18. For influenza B, we obtained modest agreement for the overall magnitude
of GP consultations as a result of type B, but this was not always in complete agreement with
the subtype composition (Fig. 5(a)).

A more detailed comparison of predicted and observed subtypes in each season allowed us to
analyse discrepancies in more detail (Fig. 6). The model predictions are shown with a violin plot
to capture the variability in the distribution due to parameter uncertainty; while the observed
data has confidence intervals (calculated from bootstrapping) due to the finite sample sizes. In
general there was good agreement between the data and model, especially for years where par-
ticular subtypes dominated (e.g. A(H1N1)pdm09 in 2013/14 and 2015/16, A(H3N2) in 2014/15
and 2016/17, and B/Yamagata in 2012/13 and 2017/18). However, there were also discrepan-
cies, most notably in 2015/16 where the majority of the B-influenza cases were predicted to
the wrong lineage; a potential consequence of pursuing model parsimony, with preference for
ensuring the generic behaviour for B/Yamagata across all other influenza seasons in the study
period was correct, rather than seeking to capture every changing nuance when fitting to the
data. It was also clear that the model generally performed less well when there were very low
levels of a given subtype in a particular year.

(a)

(b)

Fig. 5: Posterior predictive distributions for strain composition and influenza positive GP consultations per 100,000 population. Stratified by influenza season, we present back-to-back stacked bars per simulation replicate, with 1,000 replicates performed, each using a distinct parameter set representing a sample from the posterior distribution. Each influenza season is topped out by a thicker stacked horizontal bar plot, corresponding to the strain-stratified point estimates for the empirical data. **(a)** Posterior predictive distributions for circulating influenza virus subtype/lineage composition. **(b)** Posterior predictive distributions for influenza positive GP consultations per 100,000 population. In both panels, the left side depicts data pertaining to type A influenza viruses (red shading denoting the A(H1N1)pdm09 subtype, orange shading the A(H3N2) subtype). In an equivalent manner, the right side stacked horizontal bars present similar data for type B influenza (cyan shading denoting the B/Victoria lineage, dark blue shading the B/Yamagata lineage). We see a reasonable qualitative model fit to the data, especially for the two influenza A subtypes.

Fig. 6: Posterior predictive influenza positive GP consultation distributions. We generated the estimated distributions from 1,000 model simulations, each using a distinct parameter set from the retained collection of particles. Crosses represent the observed data, with solid bars the range of the bootstrapped empirical data. Shaded violin plots denote outcomes from model simulations using the empirical data on vaccination, with filled circles the median value across the simulated replicates. **(a)** A(H1N1)pdm09; **(b)** A(H3N2); **(c)** B/Victoria; **(d)** B/Yamagata.

Forward simulations

Using the parameter set from the inference scheme returning the closest correspondence with the data (lowest DEV value, Eq. (5)), we explored potential strain dynamics up to the 2029/30 influenza season under contrasting vaccine efficacy settings (Fig. 7).

When maintaining 'expected' vaccine efficacy (A(H1N1)pdm09: 55.25%; A(H3N2): 30.45% B/Victoria: 55.90%; B/Yamagata: 52.60%) across all future seasons (Fig. 7(a), single bars), we witness minor variability in predicted influenza incidence for both influenza types. Total influenza B incidence ranged between 2,700-10,250 cases per 100,000 each season, and seasonal influenza A incidence consistently reached 25,000-40,000 cases per 100,000. Furthermore, we predict periodic behaviour for the portion of influenza A viruses in circulation ascribed to the subtypes A(H1N1)pdm09 and A(H3N2), mirroring the observed pattern from 2012/13 to 2016/17. Influenza seasons ending in even numbered years (e.g. 2029/30) had a relatively even split between the two subtypes; whereas for influenza seasons ending in odd numbered years (e.g. 2028/29) the A(H3N2) subtype dominated the A(H1N1)pdm09, with influenza A viruses apportioned to A(H3N2) and A(H1N1)pdm09 subtypes being 70-85% and 15-30%, respectively.

An alternative approach is to use the single (best-fit) parameter set, but to sample the vaccine efficacies from historical estimates. The most striking factor was the amount of variability that comes as a direct results of uncertainty in the vaccine efficacies. As such all between-season patterns were swamped by the within-season variability, leading to distributions that were remarkably consistent between seasons. The only notable exception was 2018/19, where the definitive knowledge of the 2017/18 season had a clear impact of the predictions for 2018/19 leading to broadly higher levels of influenza A and lower levels of influenza B (Fig. 7(a)).

Comparing subtype distributions when vaccine efficacies were either sampled from historical estimates or fixed at 'expected' values, we found that under randomly sampled vaccine efficacies incidence of B/Yamagata tended to exceed B/Victoria in each season (holding in more than 70% of simulations, Figure S18). However, we obtained greater levels of variation for the influenza A subtypes. In an 'expected' vaccine efficacy setting (single bars), less than half of total influenza A cases in any season were attributed to the A(H1N1)pdm09 subtype (rather than A(H3N2)), whereas under variable vaccine efficacies (distributed bars) A(H1N1)pdm09 incidence was greater than A(H3N2) incidence in 30-50% of all simulations (Figure S18). Furthermore, in each future influenza season the majority of simulations (∼60-75%) performed with seasonally variable vaccine efficacies resulted in overall incidence rates exceeding the equivalent seasonal incidence estimates generated under 'expected' vaccine efficacy conditions.

The impact of vaccine efficay on projected incidence is further exemplified by the stark contrast in predictions when comparing the pessimistic scenario (A(H1N1)pdm09: 23.00%; A(H3N2): 0.00%; B/Victoria: 24.70%; B/Yamagata: 0.00%) to the optimistic scenario (A(H1N1)pdm09: 73.00%; A(H3N2): 61.00% B/Victoria: 92.00%; B/Yamagata: 92.00%). Under pessimistic vaccine efficacy conditions, both influenza types underwent vast fluctuations in incidence from one influenza season to the next. Type A was more severe in odd numbered years (e.g. 2028/29). Similarly, in even numbered years (e.g. 2029/30) type B incidence exceeded type A incidence. For the pessimistic vaccine efficacy scenario (compared to the other fixed vaccine efficacy scenarios) we also observe the strongest re-emergence of the alternating behaviour between dominant influenza A subtypes (Fig. 7(b)).

Maintaining a high vaccine efficacy, both influenza types had consistent incidence estimates. Whilst combined influenza B incidence was low, ranging approximately between 600-1700 cases per 100,000 each season, seasonal influenza A incidence frequently reached 25,000-30,000 cases

per 100,000. Although we, once more, predict periodic behaviour for the portion of influenza A viruses in circulation ascribed to the subtypes A(H1N1)pdm09 and A(H3N2), under these optimistic vaccine efficacy conditions it was influenza seasons ending in odd numbered years having a relatively even split between the two subtypes; whereas for influenza seasons ending in even numbered years the A(H3N2) subtype dominated A(H1N1)pdm09 (Fig. 7(c)).

Fig. 7: Projected influenza seasonal incidence (per 100,000 population) up to 2029/30. For all scenarios, in forward simulated seasons vaccine uptake matched that of the 2017/18 influenza season. In all panels, the left side depicts the seasonal incidence of type A influenza per 100,000 population (red shading denoting the A(H1N1)pdm09 subtype, orange shading the A(H3N2) subtype). The right side stacked horizontal bars present projected seasonal incidence of type B influenza per 100,000 population (cyan shading denoting the B/Victoria lineage, dark blue shading the B/Yamagata lineage).**(a)** Each influenza season is topped out by a thicker stacked horizontal bar plot, corresponding to the simulated estimate under the 'expected' vaccine efficacy scenario; efficacies by strain were as follows - A(H1N1)pdm09: 55.25%; A(H3N2): 30.45%; B/Victoria: 55.90%; B/Yamagata: 52.60%. Thin bars correspond to simulation runs where vaccine efficacy against each strain were randomly sampled from the empirical distribution (totalling 1,000 replicates). Simulations are arranged such that the largest epidemics are in the centre of the bar. **(b)** Projected incidence under a pessimistic vaccine efficacy scenario; efficacies by strain were as follows - A(H1N1)pdm09: 23.00%; A(H3N2): 0.00%; B/Victoria: 24.70%; B/Yamagata: 0.00%. **(c)** Projected incidence under an optimistic vaccine efficacy scenario; efficacies by strain were as follows - A(H1N1)pdm09: 73.00%; A(H3N2): 61.00%; B/Victoria: 92.00%; B/Yamagata: 92.00%.

Discussion

Understanding the processes that drive seasonal influenza transmission patterns is an important 451 public-health question, given the economic cost and burden to human health afflicted by the 452 condition [1, 2]. During recent years there has been marked progress in synthesising real-world 453 data to inform mathematical model parameterisation. Yet, prior studies have tended to model 454 each season and each strain circulating within that season independently [12, 14, 15]. Thus, the 455 presence of strain and immunity interactions have been omitted. Using data subsequent to the 456 2009 pandemic for England, we have developed a dynamic multi-strain SEIR-type transmission 457 model for seasonal influenza, explicitly incorporating immunity propagation mechanisms between 458 seasons. 459

With a view to minimising the number of independent parameters, we fit a parsimonious mecha- 460 nistic model to seasonal-level data on strain competition. In spite of the multi-strain complexity 461 and time scale of the study period (six influenza seasons), predictions from the model attain a 462 strong qualitative resemblance (in terms of strain composition and overall quantity of GP con- 463 sultations) to the empirical subtype data from England. We attribute much of the discrepancies 464 to the homogeneous way in which we have treated immunity propagation, as in practise this is 465 driven by complex patterns of waning and cross immunity as well as somewhat irregular genetic 466 drift. 467

Inferred transmissibility of type A influenza strains exceeding those of type B, which reflects 468 type A being the predominant class of influenza virus in circulation over the studied time pe- 469 riod [26]. Moreover, concentrating on the parameters relevant to immunity propagation, we 470 uncover evidence against vaccination stimulating similar long-term immunity responses as for 471 natural infection. These conclusions corroborate previous immunological studies showing that 472 infection with influenza virus can induce broader and longer-lasting protection than vaccina- 473 tion [37, 48, 49], and authenticate prior work signalling that vaccine-mediated immunity rapidly 474 wanes [50]. There are also indications that prior natural infection boosts vaccine responses 475 against antigenically drifted strains, whereas prior vaccination does not [51]. The clinically ob- 476 served impact of prior infection for enhancing vaccine efficacy was long-lasting, which may be 477 used to instruct further model refinements. 478

In the case of finding minimal support for carry over cross-reactivity protection between in- 479 fluenza B lineages (parameter b), we have reported the best parameter fits under the modelling 480 assumptions made. Considered collectively, the inferred parameter fits signify that, at the pop- 481 ulation level, a combination of transmissibility rates, season-specific ascertainment probabilities 482 and modifications to susceptibility by lineage-specific immunity propagation (parameter a) and 483 within-season vaccination are sufficient to obtain the closest correspondence to the data under 484 the modelling assumptions, without requiring a prominent amount of influenza B cross-reactive 485 immunity propagation. In other words, if there were presence of propagation of protection be- 486 tween influenza B lineages, it is dwarfed by other processes within the biological system. The lack 487 of propagation of cross-reactivity protection between influenza B lineages may be unforeseen, 488 as it has been documented that there is potential for viral interference between the influenza 489 B virus lineages; infection with one influenza B virus lineage may be beneficial in protecting 490 against subsequent infection with either influenza B virus lineage [39]. However, these experi- 491 ments (using a ferret model of human influenza) had a short separation between primary and 492 challenge virus infections (at most 28 days), whereas our timescale of interest is of the order 493 of six months (spanning between peak infection in one influenza season to the initiation of the 494 seasonal epidemic in the next). 495

The extent of immunity propagation stemming from earlier influenza infection and vaccination occurrences may conceivably be quantified through serological surveys, potentially the most direct and informative technique available to infer the dynamics of a population's susceptibility and level of immunity [52]. There is scope to implement concepts monitoring changes in the immune response to infections in a rapid and cost-effective manner, by using existing primary care sentinel networks [53]. In this fashion, in England an age-stratified pilot serology study was recently initiated utilising the RCGP RSC to collect serological specimens and associated patient data to measure seropositivity and seroincidence due to seasonal influenza [54]. These progressive developments make usage of serological data to validate our study findings a realistic prospect.

Furthermore, though there have been studies carried out to quantify the strength of vaccine induced herd immunity for seasonal influenza [3, 4], the role of immunity originating from non-immunising exposure to influenza viruses in inducing herd immunity is unexplored. An upsurge in serology data would promote such assessments.

Ascertainment probabilities are key for linking population levels of influenza infection to GP-reported incidence. For greater parsimony we have assumed that ascertainment is independent of subtype, which could be an over-simplification in some years; but allowing ascertainment to depend on both season and subtype would over-fit the model. Across all considered influenza seasons, we found ascertainment probabilities were of a similar order (and concurred with the magnitudes of inferred ascertainment probabilities in [12]), with the greatest values obtained for the 2017/18 influenza season. The 2017/18 season saw moderate to high levels of influenza activity observed in the UK [24], exacerbated by a mismatch of the A(H3N2) component of the available vaccine towards the A(H3N2) strain in circulation [36]. The effects that consequential increased media coverage of an outbreak have on transmission dynamics can be complex [55], with the perception of the disease conceivably changing among the population [56]. Chiefly in this instance, for those infected and symptomatic the propensity to consult a GP may have been abnormally raised, which would result in an increased ascertainment probability relative to prior influenza seasons.

While historic immunity is predicted to play an important role in determining annual strain composition (leading to subtype patterns between seasons), our projections of the dynamics forward in time exemplify how variability in vaccine efficacy hampers our ability to make long-term predictions. Spanning our pessimistic to optimistic vaccine efficacy scenarios, we generally attained a model predicted incidence (per influenza season) of 30-50%. With volunteer challenge studies indicating that the majority of those infected by influenza are asymptomatic [57], plus only around 10% of those with ILI thought to consult a GP [58, 59], our model predicted influenza incidence pairs satisfactorily with previous influenza burden estimates covering England [60]. Additionally, influenza burden estimates in the USA (provided by the CDC) covering the 2017/18 and 2018/19 influenza seasons ascribe more than 48.8 million and 37.4-42.9 million influenza associated illnesses respectively, which is in the realms of 10-15% of the national population suffering symptomatic infection [61, 62].

Consistently high vaccine efficacies would aid forecasting efforts, providing additional benefits beyond the reduction in disease offered by new vaccines with improved clinical efficacy and effectiveness [63]. Most importantly, with annual emergence of drift variants of influenza viruses, cross-protective vaccines are needed to heighten protective effectiveness [64].

The development of the influenza transmission model presented here was built upon a collection of simplifying assumptions. A limitation of the model is fixing the influence of prior expo-sure history to a single season. Studies of repeated vaccination across multiple seasons suggest

that vaccine effectiveness may be influenced by more than one prior season [65]. In an United Kingdom-centric analysis, Pebody *et al.* [34] scrutinised the possible effect of prior season vaccination on 2016/17 influenza season vaccine effectiveness; while there was no evidence that prior season vaccination significantly reduced the effectiveness of influenza vaccine during the current season in adults, it was deemed to increase effectiveness in children. Extending exposure history and susceptibility interaction functionality to encompass immunity propagation dependencies beyond one prior influenza season would assist efforts evaluating the potential effect of repeat influenza vaccinations in the presence of vaccine interference [66].

Second, on the basis of model parsimony, within the immunity propagation model component we limited the number of mechanisms linking prior influenza season exposure history to susceptibility in the subsequent influenza season. Though we did include a parameter corresponding to immunity proliferating across influenza B lineages between seasons, we did not include influenza A heterosubtypic immunity amongst the immunity propagation mechanisms. Recent work has found H1 and H3 influenza infection in humans induces neuraminidase-reactive antibodies displaying broad binding activity spanning the entire history of influenza A virus circulation (in humans) [37]; these developments motivate continued work and data acquisition to elicit the timespan over which cross-reactive antibodies (arising from natural influenza infection) may significantly reduce susceptibility to unrelated influenza A subtypes and influenza B lineages.

Third, we assumed the mechanism underpinning propagation of immunity from influenza vaccination in the previous season behaved linearly. We recognise a linear dependency is a strong generalisation. Given the vaccine-induced protection to an influenza virus strongly depends on the level of mismatch of the strain contained in the vaccine and the circulating strain, a vaccine that is very protective in the current influenza season might be ineffective the following year due to the influenza virus undergoing antigenic mutation. Thus, alternative non-linear formulations for the vaccine-derived immunity propagation process warrant consideration, if there is sufficiently detailed data to underpin the models.

Relatedly, to parameterise the amended strain-specific susceptibility values within exposure history groups encapsulating those both experiencing a natural infection event and undergoing vaccination in the prior influenza season, we assumed no interaction between residual vaccine and natural infection immunity (we simply enforced the dominating process). Other formulations governing the interaction between residual vaccine and natural infection immunity may be scrutinised, conditional on the emergence of relevant data to parameterise the process.

Fourth, for simplicity we assumed each season was initialised with a low level of all subtypes; in practise the subtypes seeding each epidemic will be contingent on the dynamics in the rest of the world. Finally, we assumed the value of the ascertainment probability to be constant over the course of the influenza season. The underlying assumption is that the quantities contributing towards ascertainment remain stable over time. Yet, in reality there are contributory factors that are likely to vary during the course of each seasonal outbreak; for example, the propensity to consult with a GP if inflicted with symptomatic illness.

We view the dynamic influenza transmission model presented here as a foundation stage for influenza modelling constructs incorporating strain and immunity interactions. Heterogeneity in social contact patterns and the role of age in influenza transmission potential and severity of health outcomes are important considerations, with children identified as the main spreaders of influenza infection [67, 68] and most influenza associated deaths occurring among elderly adults and those with co-morbid conditions that place them at increased risk [2, 69]. As a consequence, next steps should include augmenting population mixing patterns and age-structure into the mathematical framework. In addition, coupling the transmission model together with economic

evaluation frameworks will permit cost-effectiveness appraisals of prospective vaccination programmes.

In summary, through the use of mathematical modelling and statistical inference, our analysis of seasonal influenza transmission dynamics in England quantifies the contribution of distinct immunity propagation mechanisms. We find that susceptibility in the next season to a given influenza strain type is modulated to the greatest extent through natural infection by that strain type in the current season, with residual vaccine immunity having a lesser role and inconsequential support for carry over type B cross-reactivity. As such, the approach utilised here offers a preliminary basis for long-term influenza modelling constructs incorporating strain and immunity interactions, although forecasts are strongly influenced by vaccine efficacy. We suggest that the adoption of influenza transmission modelling frameworks with immunity propagation provides a comprehensive manner to assess the impact of seasonal vaccination programmes.

Acknowledgements

We thank Rachel Byford, Ana Correa, Chris McGee, Julian Sherlock and Sameera Pathiran-nehelage for their collective contribution towards the production of the RCGP RSC data extract utilised in this study. We thank Tom Irving and colleagues at Department of Health and Social Care for helpful discussions. We acknowledge Trystan Leng, Ben Atkins, Glen Guyver-Fletcher and Susie Cant for their constructive feedback on the manuscript. This report is independent research funded by the National Institute for Health Research (NIHR) (Policy Research Programme, Infectious Disease Dynamic Modelling in Health Protection, 027/0089). The views expressed are those of the authors and not necessarily those of the NIHR or the Department of Health and Social Care.

Author contributions

Conceptualisation: Edward M. Hill, Matt J. Keeling, Stavros Petrou.

Data curation: Simon de Lusignan, Ivelina Yonova.

Formal analysis: Edward M. Hill.

Funding acquisition: Matt J. Keeling, Stavros Petrou.

Investigation: Edward M. Hill, Matt J. Keeling.

Methodology: Edward M. Hill, Matt J. Keeling.

Software: Edward M. Hill.

Supervision: Matt J. Keeling, Stavros Petrou.

Validation: Edward M. Hill, Matt J. Keeling.

Visualisation: Edward M. Hill, Matt J. Keeling.

Writing - original draft: Edward M. Hill.

Writing - review & editing: Edward M. Hill, Matt J. Keeling, Stavros Petrou, Simon de Lusignan, Ivelina Yonova.

Data availability

The GP consultation data contain confidential information, with public data deposition non-permissible for socioeconomic reasons. The GP consultation data resides with the RCGP Research and Surveillance Centre and are available via the RCGP RSC website (www.rcgp.org.uk/rsc). All other raw data utilised in this study are publicly available; relevant references and data repositories are stated within the main manuscript and Supporting Information. Code and processed data used for the study is available at https://github.com/EdMHill/SeasonalFluImmunityPropagation.

Competing interests

The authors declare that they have no competing interests.

References

[1] Iuliano AD, Roguski KM, Chang HH, Muscatello DJ, Palekar R, *et al.* Estimates of global seasonal influenza-associated respiratory mortality: a modelling study. *Lancet* (2017). doi: 10.1016/S0140-6736(17)33293-2.

[2] Cromer D, van Hoek AJ, Jit M, Edmunds WJ, Fleming D, *et al.* The burden of influenza in England by age and clinical risk group: A statistical analysis to inform vaccine policy. *J. Infect.* **68**(4):363–371 (2014). doi:10.1016/j.jinf.2013.11.013.

[3] Loeb M, Russell ML, Moss L, Fonseca K, Fox J, *et al.* Effect of Influenza Vaccination of Children on Infection Rates in Hutterite Communities. *JAMA* **303**(10):943–950 (2010). doi:10.1001/jama.2010.250.

[4] Pebody RG, Green HK, Andrews N, Zhao H, Boddington N, *et al.* Uptake and impact of a new live attenuated influenza vaccine programme in England: early results of a pilot in primary school-age children, 2013/14 influenza season. *Eurosurveillance* **19**(22):pii=20823 (2014).

[5] Members of the Western Pacific Region Global Influenza Surveillance and Response System. Seasonal influenza vaccine policies, recommendations and use in the World Health Organization's Western Pacific Region. *West. Pacific Surveill. Response J.* **4**(3):51–59 (2013). doi:10.5365/wpsar.2013.4.1.009.

[6] Palache A, Oriol-Mathieu V, Abelin A, Music T. Seasonal influenza vaccine dose distribution in 157 countries (2004–2011). *Vaccine* **32**(48):6369–6376 (2014). doi:10.1016/j.vaccine.2014.07.012.

[7] Palache A, Oriol-Mathieu V, Fino M, Xydia-Charmanta M. Seasonal influenza vaccine dose distribution in 195 countries (2004–2013): Little progress in estimated global vaccination coverage. *Vaccine* **33**(42):5598–5605 (2015). doi:10.1016/j.vaccine.2015.08.082.

[8] Macpherson, N. Review of quality assurance of Government analytical models. HM Treasury, London. (2013). URL https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/206946/review_of_qa_of_govt_analytical_models_final_report_040313.pdf. [Online] (Accessed: 09 May 2019).

[9] Vynnycky E, Pitman R, Siddiqui R, Gay N, Edmunds WJ. Estimating the impact of childhood influenza vaccination programmes in England and Wales. *Vaccine* **26**(41):5321–5330 (2008). doi:10.1016/j.vaccine.2008.06.101.

[10] Baguelin M, Hoschler K, Stanford E, Waight P, Hardelid P, *et al.* Age-Specific Incidence of A/H1N1 2009 Influenza Infection in England from Sequential Antibody Prevalence Data Using Likelihood-Based Estimation. *PLoS One* **6**(2):e17074 (2011). doi:10.1371/journal.pone.0017074.

[11] Pitman R, White L, Sculpher M. Estimating the clinical impact of introducing paediatric influenza vaccination in England and Wales. *Vaccine* **30**(6):1208–1224 (2012). doi:10.1016/j.vaccine.2011.11.106.

[12] Baguelin M, Flasche S, Camacho A, Demiris N, Miller E, *et al.* Assessing Optimal Target Populations for Influenza Vaccination Programmes: An Evidence Synthesis and Modelling Study. *PLoS Med.* **10**(10):e1001527 (2013). doi:10.1371/journal.pmed.1001527.

[13] Goeyvaerts N, Willem L, Van Kerckhove K, Vandendijck Y, Hanquet G, *et al.* Estimating dynamic transmission model parameters for seasonal influenza by fitting to age and season-specific influenza-like illness incidence. *Epidemics* **13**:1–9 (2015). doi:10.1016/j.epidem.2015.04.002.

[14] Rajaram S, Wiecek W, Lawson R, Blak BT, Zhao Y, *et al.* Impact of increased influenza vaccination in 2–3-year-old children on disease burden within the general population: A Bayesian model-based approach. *PLoS One* **12**(12):e0186739 (2017). doi:10.1371/journal.pone.0186739.

[15] Backer J, Wallinga J, Meijer A, Donker G, van der Hoek W, *et al.* The impact of influenza vaccination on infection, hospitalisation and mortality in the Netherlands between 2003 and 2015. *Epidemics* **26**:77–85 (2019). doi:10.1016/j.epidem.2018.10.001.

[16] Pitman R, Nagy L, Sculpher M. Cost-effectiveness of childhood influenza vaccination in England and Wales: Results from a dynamic transmission model. *Vaccine* **31**(6):927–942 (2013). doi:10.1016/j.vaccine.2012.12.010.

[17] Baguelin M, Camacho A, Flasche S, Edmunds WJ. Extending the elderly- and risk-group programme of vaccination against seasonal influenza in England and Wales: a cost-effectiveness study. *BMC Med.* **13**(1):236 (2015). doi:10.1186/s12916-015-0452-y.

[18] Hodgson D, Baguelin M, van Leeuwen E, Panovska-Griffiths J, Ramsay M, *et al.* Effect of mass paediatric influenza vaccination on existing influenza vaccination programmes in England and Wales: a modelling and cost-effectiveness analysis. *Lancet Public Heal.* **2**(2):e74–e81 (2017). doi:10.1016/S2468-2667(16)30044-5.

[19] Thorrington D, van Leeuwen E, Ramsay M, Pebody R, Baguelin M. Cost-effectiveness analysis of quadrivalent seasonal influenza vaccines in England. *BMC Med.* **15**:166 (2017). doi:10.1186/s12916-017-0932-3.

[20] Thorrington D, van Leeuwen E, Ramsay M, Pebody R, Baguelin M. Assessing optimal use of the standard dose adjuvanted trivalent seasonal influenza vaccine in the elderly. *Vaccine* **37**(15):2051–2056 (2019). doi:10.1016/j.vaccine.2019.03.002.

[21] World Health Organisation. Global Epidemiological Surveillance Standards for Influenza (2014). URL https://www.who.int/influenza/resources/documents/WHO_Epidemiological_Influenza_Surveillance_Standards_2014.pdf. [Online] (Accessed: 09 May 2019).

[22] Royal College of General Practitioners. RCGP RSC Weekly Returns Service Annual Report 2016-2017 (2017).

[23] Health Protection Agency. Surveillance of influenza and other respiratory viruses in the UK publications (2014). URL https://webarchive.nationalarchives.gov.uk/20140629102650/http://www.hpa.org.uk/Publications/InfectiousDiseases/Influenza/. [Online] (Accessed: 09 May 2019).

[24] Public Health England. Official Statistics: Annual flu reports (2018). URL https://www.gov.uk/government/statistics/annual-flu-reports. [Online] (Accessed: 09 May 2019).

[25] Public Health England. Collection: Weekly national flu reports (2018). URL https://www.gov.uk/government/collections/weekly-national-flu-reports. [Online] (Accessed: 09 May 2019).

[26] World Health Organisation. FluNet. (2019). URL http://www.who.int/influenza/gisrs_laboratory/flunet/en/. [Online] (Accessed: 09 May 2019).

[27] Department of Health. Pandemic H1N1 (Swine) Influenza Vaccine Uptake amongst Patient Groups in Primary Care in England (2010). URL https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/215977/dh_121014.pdf. [Online] (Accessed: 09 May 2019).

[28] Hardelid P, Fleming DM, McMenamin J, Andrews N, Robertson C, *et al.* Effectiveness of pandemic and seasonal influenza vaccine in preventing pandemic influenza A(H1N1)2009 infection in England and Scotland 2009-2010. *Eurosurveillance* **16**(2):pii=19763 (2011). doi:10.2807/ese.16.02.19763-en.

[29] Pebody RG, Andrews N, Fleming DM, McMenamin J, Cottrell S, *et al.* Age-specific vaccine effectiveness of seasonal 2010/2011 and pandemic influenza A(H1N1) 2009 vaccines in preventing influenza in the United Kingdom. *Epidemiol. Infect.* **141**(3):620–630 (2013). doi:10.1017/S0950268812001148.

[30] Pebody RG, Andrews N, McMenamin J, Durnall H, Ellis J, *et al.* Vaccine effectiveness of

2011/12 trivalent seasonal influenza vaccine in preventing laboratory-confirmed influenza in primary care in the United Kingdom: evidence of waning intra-seasonal protection. *Eurosurveillance* **18**(5):20389 (2013). doi:10.2807/ese.18.05.20389-en.

[31] Andrews N, McMenamin J, Durnall H, Ellis J, Lackenby A, *et al.* Effectiveness of trivalent seasonal influenza vaccine in preventing laboratory-confirmed influenza in primary care in the United Kingdom: 2012/13 end of season results. *Eurosurveillance* **19**(27):20851 (2014). doi:10.2807/1560-7917.ES2014.19.27.20851.

[32] Pebody R, Warburton F, Andrews N, Ellis J, von Wissmann B, *et al.* Effectiveness of seasonal influenza vaccine in preventing laboratory-confirmed influenza in primary care in the United Kingdom: 2014/15 end of season results. *Eurosurveillance* **20**(36):30013 (2015). doi:10.2807/1560-7917.ES.2015.20.36.30013.

[33] Pebody R, Warburton F, Ellis J, Andrews N, Potts A, *et al.* Effectiveness of seasonal influenza vaccine for adults and children in preventing laboratory-confirmed influenza in primary care in the United Kingdom: 2015/16 end-of-season results. *Eurosurveillance* **21**(38):30348 (2016). doi:10.2807/1560-7917.ES.2016.21.38.30348.

[34] Pebody R, Warburton F, Ellis J, Andrews N, Potts A, *et al.* End-of-season influenza vaccine effectiveness in adults and children, United Kingdom, 2016/17. *Eurosurveillance* **22**(44):pii=17–00306 (2017). doi:10.2807/1560-7917.ES.2017.22.44.17-00306.

[35] Public Health England. Surveillance of influenza and other respiratory viruses in the United Kingdom: Winter 2013/14 (2014). URL https://assets.publishing. service.gov.uk/government/uploads/system/uploads/attachment_data/file/325203/ Flu_annual_report_June_2014.pdf. [Online] (Accessed: 09 May 2019).

[36] Public Health England. Influenza vaccine effectiveness (VE) in adults and children in primary care in the United Kingdom (UK): provisional end-of-season results 2017-18 (2018). URL https://assets.publishing.service.gov.uk/government/uploads/system/uploads/ attachment_data/file/779474/Influenza_vaccine_effectiveness_in_primary_care_2017_2018. pdf. [Online] (Accessed: 09 May 2019).

[37] Chen YQ, Wohlbold TJ, Zheng NY, Huang M, Huang Y, *et al.* Influenza Infection in Humans Induces Broadly Cross-Reactive and Protective Neuraminidase-Reactive Antibodies. *Cell* **173**(2):417–429.e10 (2018). doi:10.1016/j.cell.2018.03.030.

[38] Halloran ME, Haber M, Longini IM. Interpretation and Estimation of Vaccine Efficacy under Heterogeneity. *Am. J. Epidemiol.* **136**(3):328–343 (1992). doi:10.1093/oxfordjournals. aje.a116498.

[39] Laurie KL, Horman W, Carolan LA, Chan KF, Layton D, *et al.* Evidence for Viral Interference and Cross-reactive Protective Immunity Between Influenza B Virus Lineages. *J. Infect. Dis.* **217**(4):548–559 (2018). doi:10.1093/infdis/jix509.

[40] Axelsen JB, Yaari R, Grenfell BT, Stone L. Multiannual forecasting of seasonal influenza dynamics reveals climatic and evolutionary drivers. *Proc. Natl. Acad. Sci.* **111**(26):9538–9542 (2014). doi:10.1073/pnas.1321656111.

[41] Du X, King AA, Woods RJ, Pascual M. Evolution-informed forecasting of seasonal influenza A (H3N2). *Sci. Transl. Med.* **9**(413):eaan5325 (2017). doi:10.1126/scitranslmed.aan5325.

[42] Ranjeva S, Subramanian R, Fang VJ, Leung GM, Ip DKM, *et al.* Age-specific differences in the dynamics of protective immunity to influenza. *Nat. Commun.* **10**:1660 (2019). doi: 10.1038/s41467-019-09652-6.

[43] Lessler J, Reich NG, Brookmeyer R, Perl TM, Nelson KE, *et al.* Incubation periods of acute respiratory viral infections: a systematic review. *Lancet Infect. Dis.* **9**(5):291–300 (2009). doi:10.1016/S1473-3099(09)70069-6.

[44] Cauchemez S, Carrat F, Viboud C, Valleron AJ, Boëlle PY. A Bayesian MCMC approach to study transmission of influenza: application to household longitudinal data. *Stat. Med.*

23(22):3469–3487 (2004). doi:10.1002/sim.1912.

[45] Office for National Statistics. Dataset: National life tables: England (2018). URL https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/ lifeexpectancies/datasets/nationallifetablesenglandreferencetables. [Online] (Accessed: 09 May 2019).

[46] Lenormand M, Jabot F, Deffuant G. Adaptive approximate Bayesian computation for complex models. *Comput. Stat.* **28**(6):2777–2796 (2013). doi:10.1007/s00180-013-0428-3.

[47] Filippi S, Barnes CP, Cornebise J, Stumpf MP. On optimality of kernels for approximate Bayesian computation using sequential Monte Carlo. *Stat. Appl. Genet. Mol. Biol.* **12**(1):87–107 (2013). doi:10.1515/sagmb-2012-0069.

[48] Margine I, Hai R, Albrecht RA, Obermoser G, Harrod AC, *et al.* H3N2 Influenza Virus Infection Induces Broadly Reactive Hemagglutinin Stalk Antibodies in Humans and Mice. *J. Virol.* **87**(8):4728–4737 (2013). doi:10.1128/JVI.03509-12.

[49] Nachbagauer R, Choi A, Hirsh A, Margine I, Iida S, *et al.* Defining the antibody cross-reactome directed against the influenza virus surface glycoproteins. *Nat. Immunol.* **18**(4):464–473 (2017). doi:10.1038/ni.3684.

[50] Kissling E, Rondy M, study team IMM. Early 2016/17 vaccine effectiveness estimates against influenza A(H3N2): I-MOVE multicentre case control studies at primary care and hospital levels in Europe. *Eurosurveillance* **22**(7):30464 (2017). doi:10.2807/1560-7917.ES. 2017.22.7.30464.

[51] Kim JH, Liepkalns J, Reber AJ, Lu X, Music N, *et al.* Prior infection with influenza virus but not vaccination leaves a long-term immunological imprint that intensifies the protective efficacy of antigenically drifted vaccine strains. *Vaccine* **34**(4):495–502 (2016). doi:10.1016/j.vaccine.2015.11.077.

[52] Metcalf CJE, Farrar J, Cutts FT, Basta NE, Graham AL, *et al.* Use of serological surveys to generate key insights into the changing global landscape of infectious disease. *Lancet* **388**(10045):728–730 (2016). doi:10.1016/S0140-6736(16)30164-7.

[53] de Lusignan S, Correa A. Opportunities and challenges of a World Serum Bank. *Lancet* **389**(10066):250–251 (2017). doi:10.1016/S0140-6736(17)30046-6.

[54] de Lusignan S, Borrow R, Tripathy M, Linley E, Zambon M, *et al.* Serological surveillance of influenza in an English sentinel network: pilot study protocol. *BMJ Open* **9**(3):e024285 (2019). doi:10.1136/bmjopen-2018-024285.

[55] Tchuenche JM, Dube N, Bhunu CP, Smith RJ, Bauch CT. The impact of media coverage on the transmission dynamics of human influenza. *BMC Public Health* **11 Suppl 1**:S5 (2011). doi:10.1186/1471-2458-11-S1-S5.

[56] Young ME, Norman GR, Humphreys KR. Medicine in the Popular Press: The Influence of the Media on Perceptions of Disease. *PLoS One* **3**(10):e3552 (2008). doi:10.1371/journal. pone.0003552.

[57] Carrat F, Vergu E, Ferguson NM, Lemaitre M, Cauchemez S, *et al.* Time Lines of Infection and Disease in Human Influenza: A Review of Volunteer Challenge Studies. *Am. J. Epidemiol.* **167**(7):775–785 (2008). doi:10.1093/aje/kwm375.

[58] Tilston NL, Eames KT, Paolotti D, Ealden T, Edmunds WJ. Internet-based surveillance of Influenza-like-illness in the UK during the 2009 H1N1 influenza pandemic. *BMC Public Health* **10**:650 (2010). doi:10.1186/1471-2458-10-650.

[59] Brooks-Pollock E, Tilston N, Edmunds WJ, Eames KT. Using an online survey of healthcare-seeking behaviour to estimate the magnitude and severity of the 2009 H1N1v influenza epidemic in England. *BMC Infect. Dis.* **11**:68 (2011). doi:10.1186/1471-2334-11-68.

[60] Pitman R, Melegaro A, Gelb D, Siddiqui M, Gay N, *et al.* Assessing the burden of influenza and other respiratory infections in England and Wales. *J. Infect.* **54**(6):530–538 (2007). doi:

10.1016/j.jinf.2006.09.017.

[61] Centers for Disease Control and Prevention. Estimated Influenza Illnesses, Medical visits, Hospitalizations, and Deaths in the United States — 2017–2018 influenza season (2018). URL https://www.cdc.gov/flu/about/burden/2017-2018.htm. [Online] (Accessed: 15 July 2019).

[62] Centers for Disease Control and Prevention. 2018-2019 U.S. Flu Season: Preliminary Burden Estimates (2019). URL https://www.cdc.gov/flu/about/burden/preliminary-in-season-estimates.htm. [Online] (Accessed: 15 July 2019).

[63] Osterholm MT, Kelley NS, Sommer A, Belongia EA. Efficacy and effectiveness of influenza vaccines: a systematic review and meta-analysis. *Lancet Infect. Dis.* **12**(1):36–44 (2012). doi:10.1016/S1473-3099(11)70295-X.

[64] Carrat F, Flahault A. Influenza vaccine: The challenge of antigenic drift. *Vaccine* **25**(39-40):6852–6862 (2007). doi:10.1016/j.vaccine.2007.07.027.

[65] Belongia EA, Skowronski DM, McLean HQ, Chambers C, Sundaram ME, *et al.* Repeated annual influenza vaccination and vaccine effectiveness: review of evidence. *Expert Rev. Vaccines* **16**(7):723–736 (2017). doi:10.1080/14760584.2017.1334554.

[66] Shim E, Smith KJ, Nowalk MP, Raviotta JM, Brown ST, *et al.* Impact of seasonal influenza vaccination in the presence of vaccine interference. *Vaccine* **36**(6):853–858 (2018). doi:10.1016/j.vaccine.2017.12.067.

[67] Viboud C, Boëlle PY, Cauchemez S, Lavenu A, Valleron AJ, *et al.* Risk factors of influenza transmission in households. *Br. J. Gen. Pract.* **54**(506):684–689 (2004).

[68] Cauchemez S, Valleron AJ, Boëlle PY, Flahault A, Ferguson NM. Estimating the impact of school closure on influenza transmission from Sentinel data. *Nature* **452**(7188):750–754 (2008). doi:10.1038/nature06732.

[69] Mauskopf J, Klesse M, Lee S, Herrera-Taracena G. The burden of influenza complications in different high-risk groups: a targeted literature review. *J. Med. Econ.* **16**(2):264–277 (2013). doi:10.3111/13696998.2012.752376.

Supporting information items

Supporting Information

Supporting information for 'Seasonal influenza: Modelling approaches to capture immunity propagation'. This supplement consists of the following parts: (1) Data descriptions; (2) Complementary details of the modelling approach; (3) Parameter inference; (4) Additional results; (5) Model extension: Immunity propagation across multiple influenza seasons.

Figure S1

Breakdown of subtype/lineage composition of sampled influenza viruses, per influenza type, from the United Kingdom in each influenza season (2009/10 onward). **(a)** Type A influenza, stratified by subtype; **(b)** Type B influenza, stratified by lineage.

Figure S2

Breakdown of subtype/lineage composition of sampled influenza viruses, as a proportion of all influenza samples, from the United Kingdom in each influenza season (2009/10 onward). Sequenced samples confirmed as influenza positive, stratified by type and subtype/lineage. We assumed the fraction of undetermined samples ascribed to each subtype/lineage matched the observed proportions (for influenza seasons where strain-specific information were available).

Figure S3

All age, overall population vaccination uptake for 2009/10-2017/18 influenza seasons. Vaccine uptake per influenza season were as follows (to 1 d.p.): 2009/10 - 21.0%; 2010/11 - 21.0%; 2011/12 - 21.7%; 2012/13 - 21.7%; 2013/14 - 23.0%; 2014/15 - 22.9%; 2015/16 - 22.9%; 2016/17 - 24.4%; 2017/18 - 26.0%.

Figure S4

Epidemiological model schematic. Diagram of the compartmentalisation of the population based on an SEIR model, in conjunction with non-vaccinated (N) and vaccinated (V) statuses under a 'leaky' vaccination assumption. Epidemiological processes are represented by dashed lines. Vaccination processes are represented by solid lines. Demographic process (births and deaths) have been omitted. Compartments situated within the shaded region signify vaccinated groups, with vaccination occurring at rate $/mu$. Transmission events lead to movement from a susceptible state (S) to latent (E, infected but not yet infectiousness). Latency is lost at rate $\gamma_{1,m}$. Infected (I) transition to recovered (R) at rate γ_2. Equivalent formulation for each strain m.

Figure S5

Example time series displaying proportion of population in infected states (latent or infectious). Grey lines represent the simulated infected prevalence time series for 10 retained parameter sets from the inference procedure. With each simulation replicate starting on 1st September, red dashed vertical lines designate the beginning of March in each year. As stipulated by our parameter selection criteria, peak infection occurs prior to the 1st March.

Figure S6

Posterior predictive influenza positive GP visit distributions: 2012/13 influenza season. Empirical data point estimate and bootstrap samples are depicted by the red diamond and green shaded regions respectively. Blue dots denote posterior predicted replicates. **(left)** Type A versus type B; **(top right)** influenza A subtypes, A(H1N1)pdm09 versus A(H3N2); **(bottom right)** influenza B lineages, B/Yamagata versus B/Victoria.

Figure S7

Posterior predictive influenza positive GP visit distributions: 2013/14 influenza season. Empirical data point estimate and bootstrap samples are depicted by the red diamond and green shaded regions respectively. Blue dots denote posterior predicted replicates. **(left)** Type A versus type B; **(top right)** influenza A subtypes, A(H1N1)pdm09 versus A(H3N2); **(bottom right)** influenza B lineages, B/Yamagata versus B/Victoria.

Figure S8

Posterior predictive influenza positive GP visit distributions: 2014/15 influenza season. Empirical data point estimate and bootstrap samples are depicted by the red diamond and green shaded regions respectively. Blue dots denote posterior predicted replicates. **(left)** Type A versus type B; **(top right)** influenza A subtypes, A(H1N1)pdm09 versus A(H3N2); **(bottom right)** influenza B lineages, B/Yamagata versus B/Victoria.

Figure S9

Posterior predictive influenza positive GP visit distributions: 2015/16 influenza season. Empirical data point estimate and bootstrap samples are depicted by the red diamond and green shaded regions respectively. Blue dots denote posterior predicted replicates. **(left)** Type A versus type B; **(top right)** influenza A subtypes, A(H1N1)pdm09 versus A(H3N2); **(bottom right)** influenza B lineages, B/Yamagata versus B/Victoria.

Figure S10

Posterior predictive influenza positive GP visit distributions: 2016/17 influenza season. Empirical data point estimate and bootstrap samples are depicted by the red diamond and green shaded regions respectively. Blue dots denote posterior predicted replicates. **(left)**

Type A versus type B; **(top right)** influenza A subtypes, A(H1N1)pdm09 versus A(H3N2); **(bottom right)** influenza B lineages, B/Yamagata versus B/Victoria.

Figure S11

Posterior predictive influenza positive GP visit distributions: 2017/18 influenza season. Empirical data point estimate and bootstrap samples are depicted by the red diamond and green shaded regions respectively. Blue dots denote posterior predicted replicates. **(left)** Type A versus type B; **(top right)** influenza A subtypes, A(H1N1)pdm09 versus A(H3N2); **(bottom right)** influenza B lineages, B/Yamagata versus B/Victoria.

Figure S12

Results of the ABC scheme, fitting to empirical data covering 2012/13-2015/16 influenza seasons (inclusive). (a) Summary metric threshold value upon completion of each generation of the inference scheme. The inset panel displays the latter quarter of generations. **(b)** Inferred parameter distributions estimated from 10,000 retained samples following completion of 1,100 generations of the inference scheme. Vertical red lines indicate the (non-weighted) median values for the model constants estimated from the inference procedure.

Figure S13

Results of the ABC scheme, fitting to empirical data covering 2012/13-2016/17 influenza seasons (inclusive). (a) Summary metric threshold value upon completion of each generation of the inference scheme. The inset panel displays the latter quarter of generations. **(b)** Inferred parameter distributions estimated from 10,000 retained samples following completion of 1,000 generations of the inference scheme. Vertical red lines indicate the (non-weighted) median values for the model constants estimated from the inference procedure.

Figure S14

Posterior predictive influenza positive GP consultation distributions, post-fitting to the empirical data covering the 2012/13-2015/16 influenza seasons (inclusive). We generated the estimated distributions from 1,000 model simulations, each using a distinct parameter set from the retained collection of particles. **(a)** Back-to-back stacked bars per simulation replicate. Each influenza season is topped out by a thicker stacked horizontal bar plot, corresponding to the strain-stratified point estimates for the empirical data. **(b)** Comparison of model simulated outcomes (shaded violin plots, with filled circles corresponding to the median value across the simulated replicates) versus the observed data (crosses denote the point estimate, with solid bars the range of the bootstrapped empirical data).

Figure S15

Posterior predictive influenza positive GP consultation distributions, post-fitting to the empirical data covering the 2012/13-2016/17 influenza seasons (inclusive). We generated the estimated distributions from 1,000 model simulations, each using a distinct

parameter set from the retained collection of particles. **(a)** Back-to-back stacked bars per simulation replicate. Each influenza season is topped out by a thicker stacked horizontal bar plot, corresponding to the strain-stratified point estimates for the empirical data. **(b)** Comparison of model simulated outcomes (shaded violin plots, with filled circles corresponding to the median value across the simulated replicates)versus the observed data (crosses denote the point estimate, with solid bars the range of the bootstrapped empirical data).

Figure S16

Results of the ABC scheme, fitting to synthetic data. (a) Summary metric threshold value upon completion of each generation of the inference scheme. The inset panel displays the latter quarter of generations. **(b)** Inferred parameter distributions estimated from 10,000 retained samples following completion of 1,000 generations of the inference scheme. Vertical red lines indicate the (non-weighted) median values for the model constants estimated from the inference procedure, and vertical black lines correspond to the true values of the parameters from which the data were generated. We ably recovered the parameter values from which the synthetic data had been generated.

Figure S17

Posterior predictive distributions for influenza positive GP consultations per 100,000 population, using the parameter sets generated when fitting to the synthetic data. Stratified by influenza season, we present back-to-back stacked bars for 1,000 simulation replicates, each using a distinct parameter set representing a sample from the posterior distribution when fitting our mathematical model to the synthetic data. Each influenza season is topped by a thicker stacked horizontal bar plot, corresponding to the synthetic data values. The left side depicts the cumulative total of ILI GP consultations attributable to type A influenza per 100,000 population (red shading denoting the A(H1N1)pdm09 subtype, orange shading the A(H3N2) subtype). In an equivalent manner, the right side stacked horizontal bars present similar data for type B influenza (cyan shading denoting the B/Victoria lineage, dark blue shading the B/Yamagata lineage). Overall, simulated outcomes from each parameter set had strong agreement with the synthetic data.

Figure S18

Quantification of the projected dominating influenza A subtype and influenza B lineage by influenza season, up to 2029/30. The left half depicts the proportion of simulations in which the major subtype contributor (>50% of cases) to the overall type A influenza incidence was A(H1N1)pdm09 (red bars) or A(H3N2) (orange bars). In an equivalent manner, the right half depicts the proportion of simulations in which the major lineage contributor (>50% of cases) to the overall type B influenza incidence was B/Victoria (cyan bars) or B/Yamagata (dark blue bars). Constructed from simulation runs where vaccine efficacy against each strain were randomly sampled from the empirical distribution (totalling 1,000 replicates). For all forward simulated seasons, vaccine uptake matched that of the 2017/18 influenza season.

Figure S19

Schematic showing the links between the vaccination, immunity propagation, epidemiological and observation model components for the extended model. We adopt the visualisation conventions of [12], with ellipses indicating variables, and rectangles indicating data. Dotted arrows indicate relationships between prior season epidemiological outcomes and immunity propagation factors. Solid arrows indicate within-season processes. Circled capitalised letters indicate the relationships connecting the variables or data involved. These relationships are: process A, propagation of immunity as a result of exposure to influenza virus in the previous influenza season (through natural infection or vaccination); process B, modulation of current influenza season virus susceptibility; process C, estimation of influenza case load via the SEIR model of transmission; process D, ascertainment of cases through ILI recording at GP.

Figure S20

Mechanism for conferring immunity to influenza across multiple influenza seasons. At the beginning of a generic influenza season y, those susceptible may be divided into two categories based on exposure history: (i) susceptibles in either the naive exposure history group , with unmodified susceptibility to infection, or in the vaccinated only exposure history group (labelled S); (ii) all those that are susceptible and in an exposure history group involving natural infection (labelled $\hat{S}$). During the influenza season, those that were initially susceptible may be infected (state I). Of those remaining in state $\hat{S}$ at the end of season y, a proportion δ are retained in state $\hat{S}$ at the beginning of season $y+1$. The remainder $(1 - \delta)$ transition to state S. Note, $\delta = 0$ corresponds to no residual immunity being retained due to natural infection beyond one subsequent influenza season, while $\delta = 1$ coincides with residual immunity being kept in all future seasons until the occurrence of a new infection event.

Figure S21

Results of the ABC scheme, fitting the extended model to the empirical data. (a) Summary metric threshold value upon completion of each generation of the inference scheme. The inset panel displays the latter quarter of generations. **(b)** Inferred parameter distributions estimated from 10,000 retained samples following completion of 750 generations of the inference scheme. Vertical red lines indicate the (non-weighted) median values for the model constants estimated from the inference procedure. Particularly noteworthy outcomes include: transmissibility of type A viruses exceeding type B viruses; vaccine carry over had little impact on present season susceptibility; support for immunity stemming from natural infection being retained long term; the highest ascertainment probability occurred in the 2017/18 influenza season.

Figure S22

Posterior predictive influenza positive GP consultation distributions, post-fitting the extended model to the empirical data covering the 2012/13-2017/18 influenza seasons (inclusive). We generated the estimated distributions from 1,000 model simulations, each using a distinct parameter set from the retained collection of particles. **(a)** Back-to-back stacked bars per simulation replicate. Each influenza season is topped out by a thicker stacked horizontal bar plot, corresponding to the strain-stratified point estimates for the empirical data.

(b) Comparison of model simulated outcomes (shaded violin plots, with filled circles corresponding to the median value across the simulated replicates) versus the observed data (crosses denote the point estimate, with solid bars the range of the bootstrapped empirical data).

Figure S23

Results of the ABC scheme, fitting the extended model to synthetic data. (a) Summary metric threshold value upon completion of each generation of the inference scheme. The inset panel displays the latter tenth of generations. **(b)** Inferred parameter distributions estimated from 10,000 retained samples following completion of 1,650 generations of the inference scheme. Vertical red lines indicate the (non-weighted) median values for the model constants estimated from the inference procedure, and vertical black lines correspond to the true values of the parameters from which the data were generated. We ably recovered the parameter values from which the synthetic data had been generated.

Figure S24

Posterior predictive distributions for influenza positive GP consultations per 100,000 population, using the parameter sets generated fitting the extended model to the synthetic data. Stratified by influenza season, we present back-to-back stacked bars for 1,000 simulation replicates, each using a distinct parameter set representing a sample from the posterior distribution when fitting our mathematical model to the synthetic data. Each influenza season is topped by a thicker stacked horizontal bar plot, corresponding to the synthetic data values. The left side depicts the cumulative total of ILI GP consultations attributable to type A influenza per 100,000 population (red shading denoting the A(H1N1)pdm09 subtype, orange shading the A(H3N2) subtype). In an equivalent manner, the right side stacked horizontal bars present similar data for type B influenza (cyan shading denoting the B/Victoria lineage, dark blue shading the B/Yamagata lineage). Overall, simulated outcomes from each parameter set had strong agreement with the synthetic data.

Table S1

Adjusted vaccine effectiveness estimates for influenza by season and strain type. 95% confidence intervals are stated within parentheses.

Table S2

Values (posterior weighted quantiles) for the transmission, exposure history and ascertainment probability parameters inferred fitting to 2012/13-2015/16, 2012/13-2016/17 and 2012/13-2017/18 influenza season data. Numbers inside brackets indicate 95% credible intervals. All values are given to 4 d.p.

Table S3

Values (posterior weighted quantiles) for the transmission, exposure history and ascertainment probability parameters inferred fitting to synthetic data. Numbers

inside brackets indicate 95% credible intervals. All values are given to 4 d.p.

Table S4

Overview of parameters in the extended model.

Table S5

Values (posterior weighted quantiles) for the transmission, exposure history and ascertainment probability parameters inferred fitting the extended model to the empirical data. Numbers inside brackets indicate 95% credible intervals. All values are given to 4 d.p.

Table S6

Values (posterior weighted quantiles) for the transmission, exposure history and ascertainment probability parameters inferred fitting the extended model to synthetic data. Numbers inside brackets indicate 95% credible intervals. All values are given to 4 d.p.