A light-dependent molecular link between competition cues and defense

responses in plants

G. L. Fernández-Milmanda[1], C. D. Crocco[1], M. Reichelt[2], C. A. Mazza[1], T. G. Köllner[2], T. Zhang[3,5],

M. D. Cargnel[1], M. Z. Lichy[1], A. J. Koo[3], A. T. Austin[1], J. Gershenzon[2], C. L. Ballaré[1,4]†

[1]IFEVA, Consejo Nacional de Investigaciones Científicas y Técnicas – Universidad de Buenos Aires, Ave.

San Martín 4453, C1417DSE, Buenos Aires, Argentina

[2]Max Planck Institute for Chemical Ecology, Hans-Knöll-Str. 8, 07745 Jena, Germany

[3]Department of Biochemistry, University of Missouri, Columbia 65211, USA

[4]IIBIO, Consejo Nacional de Investigaciones Científicas y Técnicas–Universidad Nacional de San Martín,

B1650HMP Buenos Aires, Argentina

[5]Current address: College of Agriculture, South China Agricultural University, Guangdong, China

† Corresponding author. Email: ballare@ifeva.edu.ar

ABSTRACT

One of the principal internal signals controlling plant growth and defense is jasmonate (JA), a

potent growth inhibitor that is simultaneously a central regulator of plant immunity to

herbivores and pathogens. When shade-intolerant plants perceive the proximity of

competitors using the photoreceptor phytochrome B (phyB), they accelerate growth and

down-regulate JA responses. However, the mechanisms by which photoreceptors relay light

cues to the JA signaling pathway are not understood. Here we identify a sulfotransferase

(ST2a) that is strongly up-regulated by plant proximity perceived by phyB via the phyB-

Phytochrome Interacting Factor (PIF) signaling module. By catalyzing the formation of a

sulfated JA derivative, ST2a acts to degrade bioactive forms of JA and represents a direct

molecular link between photoreceptors and hormone signaling in plants. The enzyme

provides a molecular mechanism for prioritizing shade avoidance over defense under close

plant competition.

RESULTS and DISCUSSION

Growth responses to competition with other plants (*1*) and defense responses to the attack of

consumer organisms (*2*) are two paradigmatic examples of adaptive phenotypic plasticity in

plants. However, the mechanistic and functional links between these responses are not well

understood. JAs are potent growth inhibitors (*3*) and regulators of cell division (*4, 5*), and their

role in balancing growth and defense is evolutionarily conserved in land plants, from

bryophytes (*6*) to angiosperms (*7*). When shade-intolerant plants perceive a high risk of

competition for light with neighboring individuals, they activate the shade-avoidance

syndrome (SAS), which allows them to position their leaves in well-illuminated areas of the

canopy. Under these competitive conditions, plants also often attenuate the expression of JA-

mediated defense responses against pathogens and herbivore (*8*). This attenuation of defense

presumably allows the plant to efficiently focus its resources and developmental decisions on

escaping shade, sacrificing plant parts that are unlikely to contribute to resource capture.

Plants perceive the proximity of competitors using photoreceptors. Low ratios of red (R) to far-

red (FR) radiation (R:FR ratio), which indicate a high risk of competition, result in partial

inactivation of the photoreceptor phyB, which in turn promotes growth-related hormonal

pathways (*9*), and attenuates signaling mechanisms involved in the activation of defense

responses, such as the JA and salicylic acid signaling pathways (*8*). The attenuation of defense

responses triggered by supplemental FR radiation or mutations in the *PHYB* gene is not merely

a consequence of redirecting resources to growth (i.e., a simple reflection of an energetic

tradeoff between growth and defense)(*10, 11*). Attenuation of JA responses under low R:FR

ratios has been associated with FR-induced changes in the balance between DELLA and JA-ZIM-

domain (JAZ) repressor proteins (*12, 13*), and decreased stability of key transcription factors,

such as MYC2 (*13*). However, the specific molecular links between the phyB and JA signaling

pathways have not been demonstrated.

Most JA-upregulated genes and metabolites are down-regulated by low R:FR ratios or phyB

inactivation (*8, 11*). In order to identify signaling elements that could be involved in the

suppression of defense in plants exposed to competition cues, we searched for genes with an

inverse pattern of regulation (i.e., JA-related genes showing increased expression under

conditions that inactivate phyB). In a microarray experiment, we found a small cluster of

approximately 100 genes that were positively regulated by JA whose expression was promoted

by phyB inactivation; this cluster included various *JAZ* genes, and a gene of the

sulfotransferase family annotated as *ST2a/SOT15* (Fig. S1A). Moerover, in an analysis of

expression patterns of JA-related genes using publicly-available microarray databases, we

discovered that this *ST2a* gene was consistently and strongly upregulated by low R:FR ratios

(Fig. S1B). Experimentally, we confirmed a strong upregulation of *ST2a* mRNA by FR radiation

treatments that mimicked the effects of plant proximity and mild suppression by UV-B

radiation (Fig. S1C), suggesting that this gene could indeed be involved in relaying information

on the canopy light conditions to the JA signaling pathway. Chromatin immunoprecipitation

sequencing (ChIP-seq) results indicate that *ST2a* is among the direct targets of Phytochrome

Interacting Factors (PIFs) (*14*). PIFs are growth-promoting transcription factors (*15*), and PIF4,

PIF5, and PIF7 have been shown to link phyB inactivation with growth responses to shade

signals (*16, 17*). We tested a *pif4 pif5 pif7* triple knock out mutant (*18*) and found that *ST2a*

mRNA was upregulated by tissue damage just like in Col-0, but the transcriptional response of

ST2a to supplemental FR radiation was completely lost (Fig. S1D). These results demonstrate

that low R:FR ratios upregulate the transcription of *ST2a* via the phyB-PIF transcription

module.

ST2a belongs to a family of 21 sulfotransferase-encoding genes in Arabidopsis (*19, 20*), and

shows sequence similarity to proteins in many dicotyledonous species (Fig. S2). In vitro, the

ST2a protein catalyzes the sulfation of 12-hydroxy JA (OH-JA) to form JA sulfate (HSO_4-JA) (*21*).

First described in the late 1800s (*22*), sulfation consists of the transfer of a sulfate residue to a

hydroxyl or amino group. In mammalian systems, sulfation represents an important pathway

for the biotransformation of hormones, neurotransmitters, and numerous xenobiotics (*23, 24*),

which in most cases results in increased water solubility and decreased biological activity.

Based largely on their activities and substrate specificities in vitro, sulfotransferases have also

been proposed to be important in the regulation of hormonal signaling in plants (*19, 20*). For

example, in Arabidopsis, a tyrosylprotein-sulfotransferase was reported to regulate the activity

of hormone peptides involved in the control of cell proliferation (*25*). However, there is very

little evidence from functional genetic studies that sulfotransferases are involved in the

adaptive modulation of phytohormonal pathways, and no information about their ecological

role in the regulation of plant responses to environmental cues.

We reasoned that a plausible physiological role for the activation of *ST2a* transcription via the

phyB-PIF module could be the attenuation of JA-mediated responses in plants exposed to

competition signals (low R:FR ratios). To test this hypothesis, we treated Arabidopsis plants

with methyl JA (MeJA) and evaluated the formation of JA-related metabolites and

accompanying changes in the expression of genes related to JA metabolism. Plants were kept

under either white light (Amb light treatment) or white light supplemented with FR radiation

(FR treatment). MeJA treatment induced rapid (≤30 min) increases in concentrations of JA and

the bioactive conjugate JA-Ile, which were followed by increases in further metabolites,

including OH-JA, OH-JA-Ile, and later-on by a marked increase in COOH-JA-Ile and HSO_4-JA (Fig.

1). Addition of FR to ambient light significantly decreased the abundance of JA and JA-Ile, and

their oxidized derivatives. In contrast, the pool of HSO_4-JA was significantly increased by

supplemental FR. The increase in HSO_4-JA concentration under FR correlated well with a

dramatic increase in *ST2a* mRNA level (Fig. S3). Genes encoding other enzymes involved in the

metabolism of JAs, such as *IAR3*, *ILL6*, and *CYP94B3* were also up-regulated by FR but to a

much lesser extent than *ST2a* (Fig. 1, Fig. S3). In summary, low R:FR ratios decreased the pools

of JA and JA-Ile, and this effect coincided with a massive up-regulation of *ST2a* gene

expression.

In the field, shade treatments, compared to full sunlight, have been reported to attenuate the

JA burst induced by insect herbivory, which correlates with attenuated production of defense

metabolites (*26*). Are the effects of shade on JA accumulation mediated by phyB inactivation

and functionally connected with *ST2a* gene expression? To address this question, we first

isolated and characterized two *ST2a* null alleles (*st2a-1* and *st2a-2*), and demonstrated that

both knock-out mutants produced only trace levels of HSO_4-JA (Fig. S4). The mutant carrying

the *st2a-1* allele was used for further functional characterization. In Col-0 plants, supplemental

FR reduced the JA burst induced by wounding, and this effect correlated with an increase in

the pool of HSO_4-JA (Fig. 2). Importantly, in *st2a-1* plants, the effect of FR attenuating the JA

response burst was completely missing (Fig. 2). OH-JA was reduced by FR in Col-0, and it was

more abundant in *st2a-1* than in wild type plants (Fig. S5). Within the resolution of our

sampling, the levels of JA-Ile were very low and variable, and most of the JA-Ile conjugates

were present in the carboxylated form (COOH-JA-Ile) 4 h after wounding. The concentrations

of the sum of JA-Ile conjugates was significantly lower in FR plants than in plants of the

ambient light treatment (Fig. S5), and significantly higher in *st2a-1* than in Col-0 plants at 4 h.

Overaccumulation of COOH-JA-Ile has also been reported in lines lacking JOX/JAOs (*27*), the

enzymes responsible for generating OH-JA (the putative substrate of ST2a). Thus, genetic

ablation of *JAO/JOX* or *ST2a* results in increased flux through JA-Ile metabolism and catabolite

accumulation. Because *ST2a* transcription in response to FR supplementation was minimal in

the *pif4 pif5 pif7* triple mutant (Fig. S1D), we used this mutant as a complementary genetic

tool to manipulate *ST2a* mRNA levels. Plants of *pif4 pif5 pif7* accumulated HSO_4-JA in response

to wounding, but failed to do so in response to supplemental FR (Fig. S6A). Furthermore, when

ST2a expression was physiologically manipulated in Col-0 plants using various combinations of

FR and wounding treatments, the variation in the HSO_4-JA content at 4 h after wounding was

largely explained by the variation in the levels of *ST2a* mRNA detected by qPCR (Fig. S6B).

Finally, when we overexpressed *ST2a* under a strong promoter, we obtained JA metabolite

profiles that were qualitatively similar to those generated in response to FR radiation (HSO_4-JA

was greatly increased, with a concomitant reduction in OH-JA, JA, and oxidized JA-Ile

conjugates; Fig S7). Neither wounding nor supplemental FR radiation affected the abundance

of transcripts of S*T2b*, a gene closely related to *ST2a* (Fig. S2), and a *st2b* null mutant showed

normal levels of HSO_4-JA (Fig. S8). These results provide compelling evidence that ST2a is the

sole sulfotransferase responsible for the generation of HSO_4-JA in Arabidopsis, and that the

increased sulfation of OH-JA under low R:FR ratios is functionally connected with the

transcriptional upregulation of the *ST2a* gene mediated by the phyB-PIF module.

Pharmacological experiments indicate that exogenous applications of OH-JA, the preferred

ST2a substrate in vitro, can enhance some JA-Ile triggered responses in Arabidopsis (*27*) and

induce the expression of certain genes, including *ST2a* (*21*). Therefore, sulfation of OH-JA by

ST2a may be critical for the generation of a genuinely inactive metabolite, channeling JA

molecules away from bioactive pools.

To define the functionality of ST2a, we measured JA-response markers in plants exposed to

mechanical wounding under contrasting light conditions. Genes involved in JA biosynthesis

(*LOX2*), JA signaling (*MYC2*), and JA response (*VSP2*) were regulated as expected in Col-0, with

FR repressing the response to wounding (Fig.3A). In the *st2a-1* null mutant, the basal

expression of these genes was higher than in Col-0, and the suppressing effect of FR radiation

completely disappeared (Fig. 3A). RNAseq analysis of samples from wounded rosettes revealed

a statistically significant overlap between the genes downregulated by FR in Col-0 plants and

those upregulated by the *st2a-1* mutation under FR radiation. The group of overlapping genes

was significantly enriched in the GO terms "Response to JA" and "JA biosynthetic process" (Fig.

3B and Data File S1). Consistent with the pattern of expression of JA biosynthetic genes in Col-

0 and *st2a-1*, we found that FR reduced the accumulation of *cis*-12-oxo-phytodienoic acid (*cis*-

OPDA) in Col-0, particularly at high rates of FR supplementation, but this effect of FR was less

marked in *st2a-1* plants (Fig. S9). Glucosinolates (GS) are important defense compounds in

Arabidopsis, which are often regulated by JA (*28*) (Fig. S10). In Col-0, the accumulation of these

JA-dependent compounds was attenuated when plants were exposed to supplemental FR

radiation (Fig. 3C), as expected (*29*). In contrast, in *st2a-1* plants, FR failed to inhibit GS

accumulation (Fig. 3C). Collectively, these data (Fig. 3) indicate that the sulfation reaction

catalyzed by ST2a plays a central role suppressing JA-dependent responses in plants

undergoing shade avoidance.

To investigate the functional role of changes in JA metabolism caused by ST2a activity, we

tested the *st2a-1* null mutant in bioassays with larvae of *Spodoptera littoralis* (a chewing

insect) and *Botrytis cinerea* (a necrotrophic pathogen). In Col-0, supplemental FR radiation

caused increased growth of *S. littoralis* caterpillars that fed on the plants, and increased the

size of necrotic lesions generated by *B. cinerea* (Fig. 4A). These FR effects were missing in

plants of *st2a-1*, which correlated strongly with the lack of effect of FR reducing the

concentration of JA (Fig. 2), JA marker gene transcripts, and defense compounds (Fig. 3).

Furthermore, the *pif4 pif5 pif7* triple mutant, which did not upregulate the transcription of

ST2a in response to supplemental FR, was significantly more resistant to *B. cinerea* than Col-0

under low R:FR ratios (Fig. 4A). These data provide compelling empirical support for a

functional connection between *ST2a* transcription, increased JA catabolism, and reduced

defense under low R:FR ratios.

Rosettes of the *st2a-1* null mutant appeared similar to those of Col-0 under ambient light, and

they showed normal morphological responses to supplemental FR radiation (leaf hyponasty

and petiole elongation) (Fig. 4B). However, when plants were treated with MeJA, the shade-

avoidance response to supplemental FR radiation was significantly attenuated in *st2a-1*; in

contrast, Col-0 plants were capable of reconfiguring their morphology, and showed a normal

response to FR even under MeJA elicitation (Fig. 4B and Fig. S11). Taken together, these results

suggest that the key function of the sulfotransferase ST2a under low R:FR ratios is to facilitate

the inactivation of JA, thereby allowing the plant to express its full repertoire of shade-

avoidance responses and maximize its competitive ability in crowded stands.

Failure to respond to competition signals with a rapid reconfiguration of shoot architecture

and leaf traits carries a disproportionate fitness penalty for plants competing for light in fast

growing stands. Under these conditions, suppression of the 'growth brake' (*3, 4*) imposed by JA

could be a key determinant of success, even if it comes at the cost of attenuating defense

responses. Our results demonstrate the molecular mechanism that links neighbor perception

via phyB with the attenuation of JA signaling (Fig. 4C), and provide a compelling example of the

role of sulfotransferases in the adaptive modulation of hormonal metabolism in plants. This

phyB-dependent sulfation mechanism generates a metabolic sink for bioactive JA, and allows

the plant to refocus its strategy on rapid growth when the perceived risk of competition for

light is strong.

1. J. Schmitt, J. R. Stinchcombe, M. S. Heschel, H. Huber, The adaptive evolution of

plasticity: phytochrome-mediated shade avoidance responses. *Integr. Comp. Biol.* **43**,

459 (2003).

2. E. E. Farmer, *Leaf Defence*. (Oxford University Press, Oxford, UK, 2014), pp. 224.

3. Y. Yan *et al.*, A downstream mediator in the growth repression limb of the jasmonate

pathway. *Plant Cell* **19**, 2470 (2007).

4. Y. Zhang, J. G. Turner, Wound-Induced endogenous jasmonates stunt plant growth by

inhibiting mitosis. *PLoS ONE* **3**, e3699 https://doi.org/10.1371/journal.pone.0003699

(2008).

5. W. Zhou *et al.*, A jasmonate signaling network activates root stem cells and promotes

regeneration. *Cell*, doi.org/10.1016/j.cell.2019.03.006 (2019).

6. I. Monte *et al.*, A single JAZ repressor controls the jasmonate pathway in *Marchantia

polymorpha*. *Mol. Plant* **12**, 185 (2019).

7. Q. Guo *et al.*, JAZ repressors of metabolic defense promote growth and reproductive

fitness in *Arabidopsis*. *P. Natl. Acad. Sci. USA* **115**, E10768 (2018).

8.   C. L. Ballaré, Light regulation of plant defense. *Annu. Rev. Plant Biol.* **65**, 335 (2014).

9.   M. de Wit, V. C. Galvão, C. Fankhauser, Light-mediated hormonal regulation of plant
growth and development. *Annu. Rev. Plant Biol.* **67**, 613 (2016).

10.   J. E. Moreno, Y. Tao, J. Chory, C. L. Ballaré, Ecological modulation of plant defense via
phytochrome control of jasmonate sensitivity. *P. Natl. Acad. Sci. USA* **106**, 4935 (2009).

11.   M. L. Campos *et al.*, Rewiring of jasmonate and phytochrome B signalling uncouples
plant growth-defense tradeoffs. *Nat. Comm.* **7**, 12570 | DOI: 10.1038/ncomms12570 |
(2016).

12.   M. Leone, M. M. Keller, I. Cerrudo, C. L. Ballaré, To grow or defend? Low red:far-red
ratios reduce jasmonate sensitivity in Arabidopsis seedlings by promoting DELLA
degradation and increasing JAZ10 stability. *New Phytol.* **204**, 355 (2014).

13.   J. M. Chico *et al.*, Repression of jasmonate-dependent defenses by shade involves
differential regulation of protein stability of MYC transcription factors and their JAZ
repressors in Arabidopsis. *Plant Cell* **26**, 1967 (2014).

14.   E. Oh, J.-Y. Zhu, Z.-Y. Wang, Interaction between BZR1 and PIF4 integrates
brassinosteroid and environmental responses. *Nat. Cell Biol.* **14**, 802 (2012).

15.   P. Leivar, P. H. Quail, PIFs: pivotal components in a cellular signaling hub. *Trends Plant*
Sci. **16**, 19 (2011).

16.   S. Lorrain, T. Allen, P. D. Duek, G. C. Whitelam, C. Fankhauser, Phytochrome-mediated
inhibition of shade avoidance involves degradation of growth-promoting bHLH
transcription factors. *Plant J.* **53**, 312 (2008).

17.   L. Li *et al.*, Linking photoreceptor excitation to changes in plant architecture. *Gene.*
Dev. **26**, 785 (2012).

18.   M. de Wit, K. Ljung, C. Fankhauser, Contrasting growth responses in lamina and petiole
during neighbor detection depend on differential auxin responsiveness rather than
different auxin levels. *New Phytol.* **208**, 198 (2015).

19.   F. Hirschmann, F. Krause, J. Papenbrock, The multi-protein family of sulfotransferases
in plants: composition, occurrence, substrate specificity, and functions. *Front. Plant*
Sci. **5**, (2014).

20.   A. Koprivova, S. Kopriva, Sulfation pathways in plants. *Chem. Biol. Int.* **259**, 23 (2016).

21.   S. K. Gidda *et al.*, Biochemical and molecular characterization of a hydroxyjasmonate
sulfotransferase from Arabidopsis thaliana. *J. Biol. Chem.* **278**, 17895 (2003).

22.   E. Baumann, Ueber Sulfosäuren im Harn. *Ber. Dtsch. Chem. Ges.* **9**, 54 (1876).

23.   K. Nagata, Y. Yamazoe, Pharmacogenetics of sulfotransferase. *Annu. Rev. Pharmacol.*
Toxicol. **40**, 159 (2000).

24.   N. Gamage *et al.*, Human sulfotransferases and their role in chemical metabolism.

Toxicol. Sci. **90**, 5 (2006).

25.   R. Komori, Y. Amano, M. Ogawa-Ohnishi, Y. Matsubayashi, Identification of

tyrosylprotein sulfotransferase in Arabidopsis. *P. Natl. Acad. Sci. USA* **106**, 15067

(2009).

26.   A. Agrawal, E. Kearney, A. Hastings, T. Ramsey, Attenuation of the jasmonate burst,

plant defensive traits, and resistance to specialist monarch caterpillars on shaded

common milkweed (*Asclepias syriaca*). *J. Chem. Ecol.* **38**, 893 (2012).

27.   E. Smirnova *et al.*, Jasmonic acid oxidase 2 hydroxylates jasmonic acid and represses

basal defense and resistance responses against *Botrytis cinerea* infection. *Mol. Plant*

10, 1159 (2017).

28.   B. A. Halkier, J. Gershenzon, Biology and biochemistry of glucosinolates. *Annu. Rev.*

Plant Biol. **57**, 303 (2006).

29.   M. D. Cargnel, P. V. Demkura, C. L. Ballaré, Linking phytochrome to plant immunity:

low red: far-red ratios increase Arabidopsis susceptibility to *Botrytis cinerea* by

reducing the biosynthesis of indolic glucosinolates and camalexin. *New Phytol.* **204**,

342 (2014).

30.   C. Wasternack, I. Feussner, The Oxylipin Pathways: Biochemistry and Function. *Annu.*

Rev. Plant Biol. **69**, 363 (2018).

ACKNOWLEDGMENTS

We thank I. Cerrudo, P. Karssemeijer and B. F. Alliani for help with the initial characterization

of *st2a* lines and P. V. Demkura and B. Rothe for technical assistance. **Funding:** Supported by

grants of the Agencia Nacional de Promoción Científica y Tecnológica, Universidad de Buenos

Aires and The New Phytologist Trust (C.L.B. and A.T.A.), the Max Planck Society (J.G.), National

Science Foundation (IOS-1557439; A.J.K.), a Georg Forster Research Award from the Alexander

von Humboldt Foundation (to C.L.B.) and a Deutscher Akademischer Austauschdienst

Fellowship (to G.L.F.-M.). **Author contributions:** G.L.F.-M. contributed to all aspects of this

research; C.D.C. contributed to experimental design, data collection, analysis and

interpretation; M.R. designed and executed protocols for metabolite and hormone analyses;

T.Z. and A.J.K. designed and performed hormone profiling; M.D.C. carried out bioassays and

M.Z.L. screened mutant lines and helped with gene expression analyses; C.A.M. and T.G.K.

performed analyses of transcriptomic data and helped with data interpretation; A.T.A. and J.G.

contributed to the general conception of the project and data interpretation; C.L.B. conceived

the project and contributed to data generation and analysis, and wrote manuscript with input

from all co-authors. **Competing interests:** The authors declare that they have no competing

interests. **Data and materials availability:** All data needed to evaluate the conclusions in the

paper are present in the main text or the supplementary materials.

SUPPLEMENTARY MATERIALS

The file includes:

Materials and Methods

Figs. S1 to S12

Tables S1 and S2

References

Data Files S1 and S2

Fig. 1. FR supplementation reduces the pool of bioactive JAs and increases *ST2a* transcription

and JA sulfation. Col-0 Arabidopsis plants were sprayed with 200 μM MeJA and harvested at

the indicated time points for measurements of JA pools and gene expression. The color of the

box outline indicates the direction of the FR effect: Blue = downregulation; Yellow = transient

upregulation; Red = upregulation; unboxed genes were not significantly regulated by FR.

Metabolic map adapted from Wasternack and Feussner (*30*). The bar charts show quantitative

data for metabolite concentrations (thin bars indicate 1 SE; n = 3 biological replicates). For

gene expression data, see Fig. S3.

Fig. 2. FR attenuates the JA burst triggered by mechanical wounding and increases the
concentration of HSO₄-JA in a *st2a*-dependent manner. Significant genotype x light (GxL)
interaction terms are indicted. For each time point, different letters indicate significant
differences between means (P < 0.05); thin bars indicate 1 SE (n = 6 biological replicates). DW =
dry weight. For additional jasmonate pools, see Fig. S5.

Fig. 3. FR downregulates gene and metabolite markers of jasmonate signaling in an *ST2a-dependent manner.* (A) qPCR results for selected markers of JA synthesis, signaling and response. (B) Summary of RNAseq results demonstrating a significant overlap between the genes downregulated by FR in Col-0 plants and those upregulated by the *st2a-1* mutation in wounded plants. The table shows the GO categories overrepresented in the set of overlapping genes (for details on analysis see Data File S1). (C) Suppression by FR of glucosinolate accumulation in plants treated with MeJA, mechanical wounding or insect herbivory (*Spodoptera littoralis*) was missing in a *st2a-1* null mutant. For specific data on 4MSOB and I3M in wounded plants, see Fig. S10 B). For induced plants, the significance of the genotype x light (GxL) interaction term is indicated in panels A and C. Different letters indicate significant (P < 0.05) differences between means; thin bars indicate 1 SE (n = 6 biological replicates for glucosinolate data o 3 for transcriptomic data).

Fig. 4. Sulfotransferase ST2a is key in regulating the growth/defense balance in response to changes in the R:FR ratio. (A) Under FR supplementation, plants that do not upregulate *ST2a* expression are better defended than their Col-0 counterparts. Bioassays were carried out comparing Col-0 and *st2a-1* rosettes using larvae of *Spodoptera littoralis* (upper panel) and inoculations with *Botrytis cinerea* spore suspensions (middle panel), and also comparing Col-0 and *pif4pif5pif7* triple mutants inoculated with *B. cinerea* spore suspensions (lower panel). The significance of the genotype x light interaction term (GxL) is indicated for each factorial experiment; different letters indicate significant (p< 0.05) differences between means. (B) *st2a-1* rosettes display normal phenotypes under control conditions but, compared with Col-0 rosettes, they display impaired shade-avoidance responses when exposed to low doses of MeJA (100 μM). **, p<0.01; for full dataset, see Fig. S10. (C) Conceptual model linking the perception of low R:FR ratios via phyB with the modulation of jasmonate metabolism and signaling through regulation of *ST2a* transcription via the phyB-PIF transcription module. Pr, inactive form of phytochrome; Pfr, active form of phytochrome.