Neuron-specific knockouts indicate the importance of network communication to

***Drosophila* rhythmicity**

M Schlichting[1], MM Diaz[1], J Xin[1] and M Rosbash[1*]

[1] Howard Hughes Medical Institute and Department of Biology

Brandeis University, Waltham, MA 02454, USA

* Correspondence should be addressed to rosbash@brandeis.edu

The authors declare no competing interests.

Abstract (150 words; limit 150)

Animal circadian rhythms persist in constant darkness and are driven by intracellular

transcription-translation feedback loops. Although these cellular oscillators communicate,

isolated mammalian cellular clocks continue to tick away in darkness without intercellular

communication. To investigate these issues in *Drosophila*, we assayed behavior as well as

molecular rhythms within individual brain clock neurons while blocking communication within

the ca. 150 neuron clock network. We also generated CRISPR-mediated neuron-specific

circadian clock knockouts. The results point to two key clock neuron groups: loss of the clock

within both regions but neither one alone has a strong behavioral phenotype in darkness;

communication between these regions also contributes to circadian period determination. Under

these dark conditions, the clock within one region persists without network communication. The

clock within the famous PDF-expressing s-LNv neurons however was strongly dependent on

network communication, likely because clock gene expression within these vulnerable sLNvs

depends on neuronal firing or light.

Introduction:

Neuronal networks make myriad contributions to behavior and physiology. By definition,

individual neurons within a network interact, and different networks also interact to coordinate

specialized functions. For example, the visual cortex and motor output centers must coordinate to

react properly to environmental changes. In a less immediate fashion, sleep centers and circadian

clocks are intertwined to properly orchestrate animal physiology. The circadian system is of

special interest: it not only times and coordinates physiology within neuronal tissues but also

sends signals to the body to keep the entire organism in sync with the cycling external

environment (Mohawk et al., 2012).

The small, circumscribed *Drosophila* clock network is ideal to address circadian

communication issues. The comparable region in mammals, the suprachiasmatic nucleus, is

composed of thousands of cells depending on the species. There are in contrast only 75 clock

neurons per hemisphere in *Drosophila*. These different clock neurons can be divided into several

subgroups according to their location within the fly brain. There are 4 lateral and 3 dorsal neuron

clusters, which have different functions in controlling fly physiology (Helfrich-Förster et al.,

2007).

The four small ventro-lateral neurons (sLNvs) are arguably the most important of the 75

clock neurons. This is because ablating or silencing these neurons abolishes rhythms in constant

darkness (DD). They reside in the accessory medulla region of the fly brain, an important

pacemaker center in many insects (Helfrich-Förster, 1997), and express the neuropeptide PDF. In

addition, they are essential for predicting dawn (Depetris-Chauvin et al., 2011; Grima et al.,

2004; Nitabach et al., 2002; Stoleru et al., 2004). A very recent study suggests that the sLNvs are

also able to modulate the timing of the evening (E) peak of behavior (Schlichting et al., 2019).

The other ventral-lateral group, the four large-ventro-lateral neurons (lLNvs), also express PDF

and send projections to the medulla, the visual center of the fly brain; they are important arousal

neurons (Shang et al., 2008; Sheeba et al., 2008). Consistent with the ablation experiments

mentioned above, the absence of *pdf* function or reducing PDF levels via RNAi causes

substantial arrhythmic behavior in DD (Renn et al., 1999; Shafer and Taghert, 2009).

Other important clock neurons include the dorso-lateral neurons (LNds), which are essential

for the timing of the E peak and adjustment to long photoperiods (Grima et al., 2004;

Kistenpfennig et al., 2018; Stoleru et al., 2004). Two other clock neuron groups, the lateral-

posterior neurons (LPN) and a subset of the dorsal neurons (DN1s), were recently shown to

connect the clock network to sleep centers in the fly central complex (Guo et al., 2018, 2016;

Lamaze et al., 2018; Ni et al., 2019). The DN2 neurons are essential for temperature preference

rhythms (Hamada et al., 2008), whereas no function has so far been assigned to the DN3s.

Despite these distinct functions, individual clock neuron groups are well-connected to each

other. At the anatomical level, all lateral neuron clusters and even DN1 dorsal neurons send

some of their projections into the accessory medulla, where they can interact. A second area of

common interaction is the dorsal brain; only the lLNvs do not project there (Helfrich-Förster et

al., 2007).

Several studies have investigated interactions between different clock neurons. Artificially

expressing kinases within specific clock neurons causes their clocks to run fast or slow and also

changes the overall free-running period of the fly, indicating that network signaling adjusts

behavior (Yao and Shafer, 2014). Similarly, speeding up or slowing down individual neurons is

able to differentially affect behavioral timing in standard light-dark (LD) cycles (Stoleru et al.,

2005; Yao et al., 2016). A high level of neuronal plasticity within the network also exists: axons

of individual cells undergo daily oscillations in their morphology (Fernández et al., 2008), and

neurons change their targets depending on the environmental condition (Gorostiza et al. 2014;

Chatterjee et al. 2018).

How neuronal communication influences the fly core feedback loop is not well understood.

The latter consists of several interlocked transcriptional-translational feedback loops, which

probably underlie rhythms in behavior and physiology (Hardin, 2011). A simplified version of

the core feedback loop consists of the transcriptional activators Clock (CLK) and Cycle (CYC)

and the transcriptional repressors Period (PER) and Timeless (TIM). CLK and CYC bind to E-

boxes within the *period* (*per*) and *timeless* (*tim*) genes (among other clock-controlled genes) and

activate their transcription. After PER and TIM synthesis in the cytoplasm, they form a

heterodimer and enter the nucleus towards the end of the night. There they interact with CLK

and CYC, release them from their E-box targets and thereby inhibit their own transcription. All

75 pairs of clock neurons contain this canonical circadian machinery, which undergoes daily

oscillations in level. Indeed, the immunohistochemical cycling of PER and TIM within these

neurons is a classic assay to visualize these molecular oscillations (Menegazzi et al., 2013).

Silencing PDF neurons stops their PER cycling, indicating an important role of neuronal firing

in maintaining circadian oscillations. However, only two time points were measured, and the

results were possibly confounded by developmental effects (Depetris-Chauvin et al., 2011;

Nitabach et al., 2002). PDF neuron silencing also phase advances PER cycling in downstream

neurons, suggesting that PDF normally serves to delay cycling in target neurons (Wu et al.,

2008). This is consistent with experiments showing that PDF signaling stabilizes PER (Li et al.,

2014). In addition, neuronal activation is able to mimic a light pulse and phase shift the clock

due to firing-mediated TIM degradation (Guo et al., 2014).

To investigate more general features of clock neuron interactions on the circadian machinery,

we silenced the majority of the fly brain clock neurons and investigated behavior and clock

protein cycling within the circadian network in a standard light-dark cycle (LD) as well as in

constant darkness (DD). Silencing abolished rhythmic behavior but had no effect on clock

protein cycling in LD, indicating that the silencing affects circadian output but not oscillator

function in a cycling light environment. Silencing similarly abolished rhythmic behavior in DD

but with very different effects on clock protein cycling. Although protein cycling in the LNds

was not affected by neuronal silencing in DD, the sLNvs dampened almost immediately.

Interestingly, this differential effect is under transcriptional control, suggesting that some

Drosophila clock neurons experience activity-regulated clock gene transcription. Cell-specific

CRISPR/Cas9 knockouts of the core clock protein PER further suggests that network properties

are critical to maintain wild-type activity-rest rhythms. Our data taken together show that clock

neuron communication and firing-mediated clock gene transcription are essential for high

amplitude and synchronized molecular rhythms as well as rhythmic physiology.

Results:

To investigate the effects of clock network communication on fly behavior, we silenced most

adult brain clock neurons using UAS-*Kir* (Johns et al., 1999). To this end, we used the *clk856*-

GAL4 driver, which is expressed in most clock neurons (Gummadova et al., 2009) and first

addressed locomotor activity behavior in a 12:12 LD cycle.

Both control strains show the expected morning and evening (M and E) anticipation

increases, which are normal behavioral manifestations of clock function (Fig. 1A and 1C). There

is however no discernable activity anticipation in the silenced flies. Only brief activity increases

are visible, precisely at the day/night and night/day transitions (Fig. 1B); these are startle

responses (Rieger et al., 2003). Flies lacking PER show similar behavior (*per[01]* Fig. 1D).

To address possible developmental defects, we added *tub-GAL80ts* as an additional

transgene to silence the clock network in an adult-specific manner. In this system, GAL80 is

active at low temperatures (18ºC) and inhibits GAL4 expression. By increasing the temperature

to 30ºC, GAL80 is inactivated, GAL4 is then functional and the *clk856* network silenced

(McGuire et al., 2003).

At the low temperature, the controls and experimental lines show a typical wild-type

bimodal activity pattern, which disappeared in experimental flies after switching to the high

temperature (Suppl. Fig. 1). This shows that the *clk856>Kir* phenotype is not caused by defects

during development.

We next compared the behavior to flies with silenced PDF neurons. Adult-specific

silencing of the PDF neurons using the gene-switch system reduced M anticipation and

significantly advanced the timing of the E peak (Fig. 1E and 1F), which reproduces previously

published results (Depetris-Chauvin et al., 2011; Nitabach et al., 2002). However, the

comparison with the *clk856* results shown above indicates that silencing the whole clock neuron

network causes a much more severe behavioral phenotype than only silencing the PDF cells, i.e.,

network silencing completely abolishes rhythmic LD behavior, similar to clock mutant flies.

How does network silencing affect the circadian molecular feedback loop? To address

this issue, we assayed PER as well as PDP1 protein levels in individual clock neuron clusters at

four different times during the LD cycle. Both proteins show robust cycling: PER peaks at the

end of the night (ZT0), and PDP1 peaks slightly earlier than PER as expected (Hardin, 2011)

(Fig. 1G-1L). This indicates that network silencing has no detectable effect on clock protein

timing or cycling amplitude in LD. These data further suggest that either the different neuron

clocks are self-sustained, comparable to the mammalian liver, or that light can drive rhythmic

gene expression even in absence of neuronal communication.

To distinguish between these possibilities, we assayed behavior and molecular cycling in

constant darkness (DD). Only 17% of the silenced flies were rhythmic, indicating that network

silencing causes high levels of DD arrhythmicity (Fig. 2A). To rule out developmental effects,

we applied the *tub-GAL80ts* system as described above: 80 percent of the experimental flies were

rhythmic at 18°C, but they were profoundly arrhythmic at 30°C with only two rhythmic flies

(Suppl. Fig. 2). In contrast, adult-specific silencing of only the PDF neurons more weakly

reduced rhythmicity (Fig. 2A) and also caused a short period (Fig. 2B), phenotypes that are

essentially indistinguishable from those of the classical pdf^{01} mutant (Renn et al., 1999).

To address why network silencing has such a profound effect, we assayed PER and PDP1

protein cycling after five days in constant darkness (DD5). As expected, all assayed clock

neurons from control strains maintain robust and coordinated cycling in DD (Fig. 2C-H); the

sLNvs, LNds and DN1s peak slightly sooner than in LD, consistent with the slightly less than 24

hr circadian period in DD (Fig. 2B).

In striking contrast, silencing the clock network causes clock protein cycling within the

individual neuronal subgroups to differ strongly from each other, in amplitude and in phase.

Clock protein cycling in the LNds is least affected by neuronal silencing and with little to no

change in phase or amplitude, suggesting a robust and possibly self-autonomous clock in these

neurons; see Discussion (Fig. 2D and 2G). The sLNvs in contrast dampen and rapidly become

arrhythmic, suggesting that these cells are rather weak oscillators and require network activity or

light for proper molecular rhythms (Fig. 2C and 2F). The DN1s also dampen but less strongly.

They manifest low amplitude cycling, which is phase-advanced; this intermediate situation

suggests a fast and somewhat network dependent clock in DN1s (Fig. 2E and 2H). The DN2s

were similar to the DN1s (data not shown). A comparable set of effects were observed in adult-

specific silencing experiments (Suppl. Fig. 3).

To further address the molecular basis of the silencing dependence, we applied a

fluorescent in-situ hybridization (fish) protocol to whole-mount *Drosophila* brains. Because *per*

mRNA was undetectable, likely due to low expression within the clock network (data not shown

and (Abruzzi et al., 2017)), we *assayed tim* mRNA cycling in LNds and in sLNvs as a proxy for

clock gene transcription/mRNA levels. In control flies under LD conditions, *tim*-RNA cycles

robustly in both sLNvs and LNds with a peak towards the beginning of the night as expected. In

addition, clock network silencing had no effect on *tim* mRNA cycling amplitude or phase in LD,

which parallels the protein cycling results (Fig. 3A and 3B). In constant darkness (DD5), the

controls show robust cycling in both sLNvs and LNds as expected, but silencing causes a

profound decrease in *tim* mRNA signal in the sLNvs; the LNds cycle normally (Fig. 3C and 3D).

These data indicate a direct correlation between neuronal activity and *tim* RNA levels at least in

the sLNvs and suggest that the silencing-mediated changes in clock protein cycling are in part

transcriptional in origin.

Network silencing therefore reveals different levels of autonomy and endogenous speeds

among clock neuron clusters. This leads to a drifting apart of the different subgroups from their

usual well-synchronized and robust clock protein expression pattern. Interestingly, it appears that

these phase differences are too big to re-establish coordinated rhythms after one week of

silencing; there is no indication of rhythmic behavior upon lowering the temperature in the

tubGAL80ts experiment (Suppl. Fig. 4).

The results to this point indicate that neuronal activity/communication is essential for

rhythmicity as well as synchronized, high amplitude clock protein cycling in DD conditions.

However, these results do not provide a hierarchy among the different groups, nor do they

address a need for the circadian clock within these neurons. To distinguish between these

possibilities and to develop a general knock-out strategy within the adult fly brain, we

established a cell-specific CRISPR/Cas9 strategy to eliminate the circadian clock in individual

clock neuron groups (Fig. 4A). We applied the guide protocol introduced by Park and Bullock

(2016) and cloned three guides targeting the coding sequence of *per* under UAS control and

generated UAS-*per-g* flies. For a first experiment, we expressed the *per*-guides and *Cas9* in most

of the clock neuron network under *clk856* control and performed behavioral (Fig. 4B-4D) and

immunocytochemical (Fig. 4E-4G) assays.

This PERKO strategy abolished M and E anticipation in LD behavior without affecting

the startle responses (Fig. 4C), and it also reduced the level of DD rhythmicity to below 10%;

this reproduced network silencing as well as the canonical per^{01} behavioral phenotypes (Fig. 1D

and 2A). Not surprisingly perhaps given these robust phenotypes, immunohistochemistry

indicates that the PERKO strategy works at more than 90% efficiency. For example, there was

no detectable nuclear PER signal in all PDF cells or in the DN2s (Fig. 4E and 4G). There is also

a marked reduction in the number of PER-positive DN1s in the dorsal brain; this is expected as

the *clk856*-GAL4 line does not express in all DN1 neurons (Gummadova et al., 2009) (Fig. 4G).

Similarly, most LNds are PER-negative. There are however two LNds that remain PER-positive

for some reason (Fig. 4F), i.e., there are a few cell escapers. We note that the PERKO strategy is

also effective with weaker and more narrowly expressed GAL4 lines (Suppl. Fig. 5), indicating

that it can be used to investigate the contribution of clocks in individual neuron subgroups to

circadian behavior.

We next addressed the contribution of clocks within individual neuron subgroups to DD

rhythmicity. Previous work assigned a central role to PDF neurons and specifically to the small

LNvs: ablating these cells eliminates DD rhythms, and expressing *per* in these same neurons

restores DD rhythms to per^{01} flies (Grima et al., 2004; Stoleru et al., 2004). We were therefore

surprised that the PERKO with *pdf*-GAL4 had no discernable effect on DD rhythmicity

compared to the controls (Fig. 4I). Similarly, a PERKO in the cells important for controlling E

activity (E cells: 3 LNds and the 5th sLNv) with *MB122*-split-GAL4 had no effect on

rhythmicity (Fig. 4J). However, a PERKO in both groups achieved with *Mai179*-GAL4, lowered

rhythmicity to less than 20% (Fig. 4K). Similar results were obtained with *dvPDF*-GAL4, which

expresses in similar neuron groups (data not shown).

To address whether other neurons have similar effects, we expressed the PER guides

elsewhere: knockout in the retina (*GMR*-GAL4), glial cells (*repo*-GAL4) or DN1s (*clk4.1M*-

GAL4 and *AstC*-GAL4) did not affect rhythmicity (Suppl. Fig. 6). These findings taken together

suggest that a clock in either of two key places, the sLNvs or the LNds, can drive rhythmic

behavior.

We also assayed the free-running DD periods of flies lacking PER in individual neuron

subgroups (Fig. 5A). These periods did not change if the PERKO was in the dorsal brain and/or

in the large PDF neurons, the lLNvs. However, a PERKO in the E cells and with drivers

expressing in these cells plus some dorsal neurons results in a slight but significant period

lengthening of approximately 0.5hr. In contrast, a PERKO in most of the lateral neuron clusters

gave rise to a short period. These two sets of period phenotypes taken together suggest that the

clocks in the two different key neuron subgroups collaborate to achieve the intermediate and

close to 24h period characteristic of wild-type flies.

Discussion

The central clock of animals is essential for dictating the myriad diurnal changes in physiology

and behavior. Knocking out core clock components such as *period* or *Clock* severely disrupts

circadian behavior as well as molecular clock properties in flies and mammals (Allada et al.,

1998; Gekakis, 1998; Konopka and Benzer, 1971). Here we show that similar behavioral effects

occur when we silence the central clock neurons and thereby abolish communication within this

network and with downstream targets, i.e., fly behavior becomes arrhythmic in LD as well as DD

conditions and resembles the phenotypes of core clock mutant strains (Konopka and Benzer,

1971).

Despite the loss of all rhythmic behavior, silencing did not impact the molecular

machinery in LD conditions: PER and PDP1 protein cycling was normal. These findings suggest

that 1) rhythmic behavior requires clock neuron output, which is uncoupled from the circadian

molecular machinery by network silencing, and 2) synchronized molecular rhythms of clock

neurons do not require neuronal activity. These findings are in agreement with previous work

showing that silencing the PDF neurons had no effect on PER cycling within these neurons

(Depetris-Chauvin et al., 2011; Nitabach et al., 2002; Wu et al., 2008). The results presumably

reflect the strong effect of the external light-dark cycle on these oscillators.

In DD however, the individual neurons change dramatically: the different neurons

desynchronize, and their protein cycling damps to different extents. Interestingly, sLNv cycling

relies most strongly on neuronal communication: these neurons cycle robustly in controls but

apparently not at all in the silenced state. sLNvs were previously shown to be essential for DD

rhythms (Grima et al., 2004; Stoleru et al., 2004). Unfortunately, the sensitivity of

immunohistochemistry precludes determining whether the molecular clock has actually stopped

or whether silencing has only (dramatically) reduced cycling amplitude. However, a simple

interpretation of the adult-specific silencing experiment favors a stopped clock: decreasing the

temperature to 18 degrees after a week at high temperature failed to rescue rhythmic behavior. A

similar experiment in mammals gave rise to the opposite result, suggesting an effect of firing on

circadian amplitude in that case (Yamaguchi et al., 2003). However, we cannot at this point

exclude a different explanation, for example a too large phase difference between the different

neuronal subgroups to reverse after a week without communication.

In either case, a stopped clock or an effect on clock protein oscillation amplitude, these

results make another link to the mammalian literature: modeling of the clock network suggests

that different neurons resynchronize more easily if the most highly-connected cells are

intrinsically weak oscillators (Webb et al., 2012). The sLNvs are essential for DD rhythms,

known to communicate with other clock neurons (Grima et al., 2004; Stoleru et al., 2004) and are

situated in the accessory medulla; this is an area of extensive neuronal interactions in many

insects (Reischig and Stengl, 2003). These considerations rationalize weak sLNv oscillators.

An important role of interneuron communication in DD is in agreement with previous

work showing that altering the speed of individual neuron groups can change the phase of

downstream target neurons (Yao and Shafer, 2014). An important signaling molecule is the

neuropeptide PDF: its absence changes the phase of downstream target neurons, and silencing

PDF neurons causes an essentially identical phenotype to the lack of PDF (Im et al., 2011; Lin et

al., 2004; Wu et al., 2008). However, the effects reported here are much stronger and show

different levels of autonomy than PDF ablation, suggesting that other signaling molecules and/or

the neuronal activity of additional clock neurons are essential to maintain proper rhythmic clock

protein expression.

To address these possibilities, we took two approaches. First, we investigated clock gene

RNA levels after silencing. The goal was to assess whether the damping of silenced neurons is

under gene expression control, likely transcriptional control. Indeed, *tim* mRNA profiles nicely

reproduced the protein cycling profiles: robust cycling of all (assayed) clock neurons was

maintained in LD even with silencing, but *tim*-mRNA levels in the sLNvs stopped cycling in

DD; in contrast, robust cycling was maintained in the LNds (Fig 2D and 2G). This suggests that

the changes in protein cycling amplitude and also possibly phase are under transcriptional

control. Importantly, the *tim* signal in the sLNvs disappeared upon silencing, suggesting that

neuronal activity promotes clock gene transcription at least in this subset of neurons. This

recapitulates for the first time in *Drosophila* the robust positive relationship between neuronal

firing and clock gene transcription in mammals (Shigeyoshi et al., 1997). To date, *Drosophila*

neuronal firing had only been connected to post-transcriptional clock protein regulation, namely

TIM degradation (Guo et al., 2014). Conceivably, these two effects are connected: TIM

degradation might be required to relieve transcriptional repression and maintain cycling.

The second approach was a cell-specific knockout strategy, applied to the clock neuron

network. We generated three guides targeting the CDS of *per* and also expressed CAS9 in a cell-

specific manner. The guides caused double strand breaks in the *per* gene, which in turn led to

cell-specific *per* mutations. This adult brain knockout strategy worked reliably and specifically,

in glial cells as well as neurons, with a greater than 90% efficiency and with no apparent

background effects (Fig. 4B-G). We have successfully used this strategy to knock out most if not

all *Drosophila* GPCRs (data not shown) and believe it will be superior to RNAi for most

purposes.Importantly, expression of the guides with the *clk856*-GAL4 driver phenocopied *per[01]*

behavior (Fig. 4C and 4H). To focus on individual clock neurons, we generated cell-specific

knockouts in different clock neurons. To our surprise, a PERKO in the PDF cells did not increase

the level of arrhythmicity. Only a PERKO in most lateral neurons, E cells as well as PDF cells,

generated high levels of arrhythmic behavior. As PDF cell ablation also causes high levels of

arrhythmicity (Stoleru et al., 2004), the data shown here suggest that the LNds can drive the

rhythmic output of key sLNv genes even in the absence of a clock in these neurons. Consistent

with this interpretation, recent data indicate that the LNds project dendritic as well as axonal

arborizations into the accessory medulla, the location of the sLNvs, indicating extensive

communication between these two important subgroups of clock neurons (Schlichting et al.,

n.d.). This interpretation is further supported by examining the free-running period of the cell-

specific knockouts: Ablating *per* in the LNds causes a long period, whereas ablating PER in most

lateral clock neurons causes a short period phenotype. These data suggest that interactions

between the sLNvs and other clock cells, perhaps within the accessory medulla, are essential for

the close to 24h speed of the overall brain and behavioral clock.

While this manuscript was being written, we became aware of two other studies

addressing the contribution of different circadian subgroups and neuronal interactions to

Drosophila rhythms. The strategy, results and conclusions in the first study overlap extensively

to what we report here (Delventhal et al., cosubmitted paper). That work exploited guides against

tim as well as against *per* and thoroughly characterized the efficacy of the cell-specific knockout

strategy, including effects on mRNA cycling.

The second study is recently published and similarly highlights the dependence of DD

rhythms on network properties (Bulthuis et al., 2019). However, they report a decrease in

rhythmicity upon knockout of the circadian clock in PDF cells, an effect that neither we nor

Delventhal observed. This difference may be due to their knockout strategy, namely,

overexpression of a dominant negative Cycle isoform within PDF cells. This protein may have

effects on gene expression beyond knocking out the circadian clock. This further suggests that

the conceptually simpler PERKO strategy has fewer side effects and is therefore superior.

Some of the communication properties described here resemble what has been found in

mammalian systems. For example, decreasing neuronal interactions by creating sparse SCN

cultures changes the free-running period and activity phase of individual neurons (Welsh et al.,

1995). This suggests that communication is also critical for circadian phase and period

determination in mammals. Nonetheless, fly clock cells may be even less cell-autonomous than

what has been described for mammals (reviewed in (Evans, 2016)). First, the fly system may be

particularly dependent on light. For example, peripheral fly clocks appear strongly light-

dependent in contrast to what has been described for mammalian liver (reviewed in (Ito and

Tomioka, 2016). Although much of the fly data could reflect cellular asynchrony in constant

darkness, circadian cycling in the periphery crashes rapidly under these conditions and resembles

the strong and rapid non-cycling that occurs in the sLNvs upon silencing in DD. Notably, fly

cryptochrome but not mammalian cryptochrome is light-sensitive (reviewed in (Michael et al.

2017) and probably contributes to the light-dependence of fly peripheral clocks. This is also

because light can directly penetrate the thin insect cuticle, which probably contributes to making

the fly brain less dependent on ocular photoreception than the mammalian brain. However, some

fly clock neurons do not express Cryptochrome, suggesting that the fly clock system is

dependent on network interactions even in a light-dark cycle (Benito et al., 2008; Yoshii et al.,

2008). These considerations suggest that the fly circadian network is an attractive object of study

not only because of its limited size of 75 neuron pairs but also because of its strong dependence

on neuronal communication.

Material and Methods

<u>Fly strains and rearing:</u>

The following fly lines were used: *clk856*-GAL4 (Gummadova et al., 2009), UAS-*Kir2.1* (BL:

6595), *pdf-GS*-GAL4 (Depetris-Chauvin et al., 2011), *per^{01} (Konopka and Benzer, 1971)*,

mai179-GAL4 (Grima et al., 2004), *MB122*-GAL4 (Guo et al., 2017), *pdfM*-GAL4 (Renn et al.,

1999) , *clk4.1M*-GAL4 (Zhang et al., 2010), UAS-*Cas9.P2* (BL 58986), AstC-GAL4 (BL:

52017), dvPDF-GAL4 (Guo et al., 2014)*, w;CyO/Sco;MKRS/TM6B* (BL: 3703), VGlut-GAL4

(BL: 60312), GMR-ss0650-GAL4, GMR-ss01038-GAL4, GMR-ss00849-GAL4, GMR-

ss00367-GAL4, GMR-ss00645-GAL4, GMR-GAL4 (BL: 1104), repo-GAL4 (BL: 7415), tub-

GAL80ts (BL: 7018). All flies were reared on standard cornmeal medium at a temperature of

25°C, with the exception of adult-specific silencing experiments for which flies were raised at

18°C.

<u>Fly line generation:</u>

We generated a UAS-*per-g* line following the protocol published by (Port and Bullock, 2016). In

short, we digested the pCFD6 Vector (addgene #73915) with BbsI, PCR amplified two PCR

fragments carrying three guides targeting the CDS of *per* and performed a Gibson Assembly to

include those in the pCFD6 backbone. Positive clones were sent for injection to Rainbow

Transgenic Flies Inc (Camarillo, CA, USA) and the transgene was inserted into the second

chromosome by phi-recombinase using BL 8621. Flies were crossed to w^{1118} for screening and

positive individuals were balanced using BL 3703. The following guide sequences were used:

per guide1: GGCAGAGCCACAACGACCTC

per guide2: CAAGATCATGGAGCACCCGG

per guide3: GAGCAAGATCATGGAGCACC

<u>Behavior recording and data analysis.</u>

Individual 2-6 days old male flies were singularly transferred into glass tubes (diameter 0.5mm)

with food (2% agar and 4% sucrose) on one end and a cotton plug to close the tube on the other

end. The tubes were placed into *Drosophila* Activity Monitors (DAM, Trikinetics) in a way that

the infrared light beam was located in the center of the tube. A computer measured the number of

light-beam interruptions caused by the movement of the fly in one-minute intervals. We recorded

the behavior of all flies at a constant temperature of 25°C for 5-7 days under standard light-dark

conditions of 12h light and 12h darkness (LD12:12) followed by constant darkness (DD) for at

least 6 days. For performing adult-specific silencing experiments, we raised the flies at 18

degrees and performed 2 separate sets of experiments: In the LD experiment, we recorded the

behavior of the flies for 3 days at 18°C and switched to 30°C to silence the neurons and follow

the behavioral change within the same sets of flies. In a second set of experiments, we raised the

two groups of flies at 18°C and then performed LD to DD experiments either at 18°Cor 30°C. In

the 30°C experiment, we decreased the temperature back to 18°C at Circadian Time (CT) 0 after

6 days in DD to investigate possible emergence of rhythmic behavior after silencing. We

continued recording the behavior for 6 more days in DD at 18°C.

We generated actograms using ActogramJ (Schmid et al., 2011). We next generated average

activity profiles of at least the last 3 days of LD condition as previously described (Schlichting

and Helfrich-Förster, 2015). Each experiment consists of at least 2 biological repeats. DD

analysis was performed using chi2-analysis. Statistical analysis was performed using a student's

t-test or one-way ANOVA followed by post-hoc Tukey analysis.

<u>Immunohistochemistry:</u>

2-6 days old male flies were entrained in LD 12:12 at 25°C for three days and collected at ZT0 to

analyze the CRISPR/Cas9 knockout strategy. To investigate clock protein cycling, 2-6 days old

male flies were entrained in LD 12:12 at 25°C for 5 days and collected in 6h intervals around the

clock. Similarly, flies were entrained for 5 days and released into DD for 5 more days to obtain

cycling data at DD5.

The whole flies were fixed for 2h 45min in 4% paraformaldehyde (PFA) in phosphate-

buffered saline (pH=7.4) including 0.5% TritonX (PBST). The flies were rinsed 5 times for 10

min each with PBST and subsequently the brains were dissected in PBST. Brains were blocked

in 5% normal goat serum (NGS) in PBST for 3h at room temperature (RT). The primary

antibody (rabbit anti-PER, 1:1000, (Stanewsky et al., 1998), mouse anti-PDF, 1:500, *Drosophila*

Studies Hybridoma Library (DSHB), C7 and guinea-pig anti PDP1, 1:2000, (Benito et al., 2007))

was applied overnight at RT and the brains subsequently rinsed 5 x 10mins with PBST.

Secondary antibodies (Alexa, Fisher Scientific, 1:200) were applied for 3h at RT. Afterwards,

the brains were rinsed 5 x 10 mins with PBST and mounted on glass slides using Vectashield

(Vector Laboratories INC., Burlingame, CA, USA) mounting medium.

Confocal microscopy was performed using a Leica SP5 microscope. Sections of 1.5 um

thickness were obtained. Laser settings were kept constant across genotypes to obtain

comparable results. Image acquisition was performed using Fiji. Staining intensity was assessed

by quantifying the brightest 3x3 pixel area of individual neurons of at least 5 brains per

timepoint. Each experiment consists of at least 2 biological repeats. Three different background

intensities were determined the same way and subtracted from the neuronal intensity. Data points

represent average and SEM.

<u>Fluorescent in-situ hybridization (fish)</u>

2-6 days old male flies were entrained for 5 days in LD 12:12 and collected in 6h intervals

around the clock. In a second set of experiments, flies were released into DD for 5 days and

collected in 6 h intervals. Flies were dissected fresh under red light to avoid phase-shifting the

molecular machinery. Brains were subsequently fixed in 4% PFA in PBS for 55 min at RT.

Afterwards, brains were washed 3 x 10min in PBST and dehydrated as described in(Long et al.,

2017). Brains were kept in 100% EtOH until all time points were collected and all further steps

were done simultaneously as described in (Long et al., 2017).

A set of 20-probe sequences were designed for the entire *pdf* mRNA sequence and

conjugated with Quasar 570 (Stellaris Probes, Biosearch Technologies, CA, USA). The *tim*

probes consist of a set of 48-probe sequences against the entire *tim* mRNA sequence, including

the 5' and 3' untranslated regions. The *tim* probes were conjugated with Quasar 670 dye

(Stellaris Probes, Biosearch Technologies, CA, USA). Probes were diluted to a stock

concentration of 25 μM and aliquoted in -20 °C. The final concentration of *pdf* probes and *tim*

probes were 250 nM and 750 nM, respectively.

Brains were mounted on glass slides using Vectashield mounting medium (Vector

Laboratories INC., Burlingame, CA, USA) and scanned using a Leica SP5 microscope in 1.5 um

sections. All samples were scanned in one session to avoid signal loss. Fluorescence intensity

was assessed by quantifying the brightest 3x3 pixel area of individual neurons of at least 5

brains. Each experiment consists of at least 2 biological repeats. Three different background

intensities were determined the same way and subtracted from the neuronal intensity. Data points

represent average and SEM.

Acknowledgements

We would like to thank Dr. Fang Guo and Dr. Katharine Abruzzi for discussions and comments

on the manuscript. Also, we are thankful to Dr. Norbert Perrimon, Dr. Paul Hardin, Dr. Gerald

Rubin and the Janelia Research Campus for providing fly lines and antibodies. Stocks obtained

from the Bloomington *Drosophila* Stock Center (NIH P40OD018537) were used in this study.

This work was supported by the Howard Hughes Medical Institute (HHMI). M.S. was sponsored

by a DFG research fellowship (SCHL2135 1/1).

Figure Legends

Figure 1 Silencing the clock network has differential effects on behavior and clock protein

cycling in LD. **A-C** Silencing most of the clock network abolishes rhythmic LD behavior. GAL4

control (**A**) and UAS control (**C**) show bimodal activity patterns with M anticipation and E

anticipation. Silenced flies (**B**) show no sign of anticipation neither in the morning nor in the

evening. Flies show short activity increases at the transitions of day/night and night/day which

are considered masking. **D** Behavior of *per^{01}* flies in LD 12:12. *per^{01}* mutants show behavior

similar to *clk856>Kir* with no M and E anticipation but short reactions to the light transitions. **E-**

F Silencing PDF neurons alters LD behavior. **E** *PDF-GS>Kir* on Vehicle food does not express

Kir. Flies show the typical bimodal activity with M and E anticipation peaks. The M peak is

close to lights-on (ZT0) whereas the E peak is close to lights-off (ZT12). **F** Silencing the PDF

neurons by adding RU 486 to the food causes an advanced E peak, similar to *pdf^{01}* flies. **G-I** PER

protein cycling is largely unaffected by neuronal silencing in LD. PER cycling in control brains

(black data points ± SEM, pooled GAL4 and UAS) is highly synchronized with peak levels

around ZT0. Silencing the clock network (red data points ± SEM) had little effect on LD PER

rhythms in sLNvs (**G**) and LNds (**H**). DN1s appear dampened after silencing (**I**). **J-L** PDP1

protein cycling is largely unaffected by neuronal silencing in LD. PDP1 cycling in control brains

(black data points ± SEM, pooled GAL4 and UAS) is highly synchronized with peak levels

around ZT18. Silencing the clock network (red data points ± SEM) had little effect on LD PDP1

rhythms in sLNvs (**J**) and LNds (**K**) and DN1s (**L**).

<u>Figure 2</u> Silencing the clock network strongly affects behavior and molecular rhythms in DD **A**

Percentage of rhythmic flies in DD. Silencing most of the clock neurons significantly reduces

rhythmicity to less than 20 percent, suggesting that clock neuron activity is essential for rhythmic

behavior output. None of the *per^{01}* flies were rhythmic as expected. Silencing the PDF neurons

slightly decreased the level of rhythmicity. **B** Free-running period of rhythmic flies from A. The

few rhythmic flies of *clk856>Kir* show a significantly longer period ($F_{(2,48)}$=14.355 p<0.001,

p<0.01 for UAS and GAL4 control). Adult-specific silencing of the PDF neurons caused a

significant period shortening (p=0.0214). **C-E** PER protein cycling in DD5. PER cycling in

control brains (black data points ± SEM, pooled GAL4 and UAS) is highly synchronized with

peak levels around CT18. Silencing the clock network (red data points ± SEM) had variable

effects on PER rhythms: The sLNvs (**C**) dampen strongly upon silencing. The LNds (**D**) show

only little differences between the groups and the DN1s strongly dampen, similar to sLNvs (**E**).

F-H PDP1 protein cycling in DD5. PER cycling in control brains (black data points ± SEM,

pooled GAL4 and UAS) is highly synchronized with peak levels around CT18. Silencing the

clock network (red data points ± SEM) had similar effects as observed in PER rhythms: The

sLNvs (**F**) dampen strongly upon silencing. The LNds (**G**) show only little differences between

the groups and the DN1s strongly dampen, but appear to have a phase-advanced PDP1 peak (**H**).

Figure 3 *tim* mRNA cycling in sLNvs and LNds shows similar trends as protein cycling observed

by FISH. **A-B** *tim* mRNA cycling in LD 12:12 in sLNvs (**A**) and LNds (**B**). Control flies (black

data points ± SEM, pooled GAL4 and UAS) show high amplitude cycling with peak levels at the

beginning of the night. Silencing the clock network (red data points ± SEM) has only little effect

on cycling amplitude or timing in LD. **C-D** *tim* mRNA cycling in DD5 in sLNvs (**C**) and LNds

(**D**). Control flies (black data points ± SEM, pooled GAL4 and UAS) show high amplitude

cycling with peak levels at the beginning of the night. Silencing the sLNvs (**C**) leads to an

overall reduction of *tim* mRNA levels and a loss of rhythmicity. In the LNds (**D**) silencing did

not decrease cycling amplitude or shifted peak mRNA expression.

Figure 4 A clock in the LNds or the sLNvs can drive rhythmic behavior **A** Schematic model of

cell-specific knockout (KO) strategy. We generated a UAS-*per-g* line using the pCFD6 vector,

allowing us to express three guides under the control of one UAS promoter (after Port and

Bullock 2016). We cloned three guides targeting the *per* CDS with guide 1 targeting the second

exon shared by all transcripts and guides 2 and 3 targeting the 4th commonly shared exon. The

guides will recruit the Cas9 protein and induce double-strand breaks and thereby cause mutations

which lead to a non-functional protein. **B-D** Behavior of perKO using *clk856*-GAL4 in LD 12:12

reproduces *per^{01}* phenotype. Flies expressing *Cas9* in the majority of the clock neurons (**B**) and

flies with both UAS-constructs (**D**) show bimodal activity with an M anticipation peak around

lights-on and an E anticipation peak around lights-off. KO of per using *clk856*-GAL4 (**C**)

abolishes M and E anticipation similar to *per^{01}* flies. **E-G** Immunocytochemistry of Control

(*clk856>Cas9*) and KO (*clk856>Cas9, perG*) staining against PER (magenta) and PDF (cyan).

Control flies show PER staining in both, sLNvs and lLNvs, whereas there is no detectable PER

signal in the PDF cells in the KO strain (**E**). Similarly, we see six LNds in the control and two

LNds in the experimental flies, showing that some neurons can escape (**F**) The number of PER+

DN1s is strongly reduced in the KO strain and we do not detect PER in the DN2 neurons (**G**). **H-**

K A clock in LNd or PDF neurons is necessary for rhythmic behavior. *Clk856*-GAL4 mediated

KO reduces rhythmicity to less than 10% (**H**). KO in PDF neurons (*pdf*-GAL4) (**I**) or in the

LNds (*MB122*-split-GAL4) (**J**) had no effect on rhythmicity. KO in both places (*Mai179*-GAL4)

significantly decreases rhythmicity (**K**).

<u>Figure 5</u> KO of PER in subsets of neurons changes free-running period. **A** Changes of free-

running period upon KO. Red bars represent the change of period between the KO line and the

GAL4 control (± SEM), green bars represent the change of period between the KO line and the

UAS control (± SEM). KO in most of the lateral neurons (*Mai179*-GAL4, *dvPDF*-GAL4) causes

a shift towards short periods. KO in the dorsal brain (*clk4.1M*-GAL4, *AstC*-GAL4, *GMR-*

ss00657-GAL4, *GMR-ss00650*-GAL4, *GMR-ss01038*-GAL4, *VGlut*-GAL4) does not affect the

period, whereas KO in the LNds (*MB122*-GAL4, *GMR-ss00849*-GAL4) lengthens the period. **B**

Model of neuronal communication and light influencing the Drosophila clock machinery.

Silencing the clock network in DD causes a damping of molecular oscillations and a drifting

apart from the common phase as indicated by the red waves. If network communication is

allowed, the different neuronal sub-clusters are mostly in sync and show robust cycling,

suggesting neuronal communication is essential for molecular oscillations. In a normal LD cycle

light drives high amplitude and synchronized cycling even in the absence of neuronal

communication, establishing a hierarchy of synchronization cues with light on the top.

Supplement Figure 1 Adult-specific silencing of most clock neurons causes arrhythmic LD

behavior. Control (*clk856*-GAL4) and experimental (*clk856*-GAL4 UAS-*Kir tub-GAL80ts*) flies

show bimodal activity at 18 degrees with an M and an E peak of activity. M anticipation is

reduced in both cases due to the low temperature. At 30 degrees, control flies (upper panel) show

a bimodal activity pattern with an M anticipation peak and an E anticipation peak. The E peak is

delayed due to high temperatures. The experimental flies (lower panel) show no M or E peak at

30 degrees.

Supplement Figure 2 Adult-specific silencing of most clock neurons causes arrhythmic DD

behavior. **A** At 18 degrees we observe no decrease in rhythmicity in *clk856>tubKir* (dark red)

compared to control (black) flies as expected. Increasing the temperature to 30 degrees (**B**)

activates GAL4 and hence silences the neurons. *Clk856>tubKir* (red) flies get arrhythmic,

whereas both controls (black) show high levels of rhythmicity. We observed no effect on free-

running period at 18 degrees (**C**) but flies experimental flies showed the tendency towards a long

period at 30 degrees (**D**), similar to *clk856>Kir* (Fig. 2B).

Supplement Figure 3 Adult-specific silencing reproduces PER cycling profiles. Flies were raised

at 18 degrees and entrained at 30 degrees to allow network silencing. Flies were collected in 6h

intervals in the fifth day of constant darkness. Control flies (*tub-GAL80ts UAS-Kir*, black data

points ± SEM) show cycling with peak expression around ZT0 in all subsets of neurons.

Silencing the network causes the sLNvs to get arrhythmic (left panel). The LNds show no

reduction in cycling amplitude and only a slight shift in protein cycling (middle panel). The

DN1s (right panel) show a reduction of PER cycling amplitude.

Supplement Figure 4 Stopping neuronal silencing after 6 days in DD does not re-establish

rhythmic behavior. Actograms of individual flies recorded at 30 degrees for 3 days in LD 12:12

(indicated by yellow boxes). Flies were then transferred into constant darkness at 30 degrees (red

area). Flies show no sign of rhythm due to the silencing of the network. We lowered the

temperature to 18 degrees (blue area) to stop neuronal silencing. None of the flies re-established

rhythmic behavior.

Supplement Figure 5 Immunolabeling of cell-specific per KO using *Mai179*-GAL4. *Mai179*-

GAL 4 is expressed in 3 out of 6 LNds, the sLNvs and shows weakly and variable expression in

DN1 and lLNv neurons. We performed PER (green) and PDF (magenta) in *Mai179>Cas9 per-g*

flies and found that 3 LNds were PER+, the sLNvs were PER- and one of the lLNvs was PER-.

This nicely reproduces the expression pattern mentioned above. Asterix represents successful

KO.

Supplement Figure 6 PER knockout in non-lateral neuron clusters does not affect rhythmicity

levels. All investigated genotypes showed high levels of rhythmicity, suggesting that the LNds

and sLNvs are the key players in determining rhythmicity.

References

Abruzzi KC, Zadina A, Luo W, Wiyanto E, Rahman R, Guo F, Shafer O, Rosbash M. 2017.

RNA-seq analysis of *Drosophila* clock and non-clock neurons reveals neuron-specific

cycling and novel candidate neuropeptides. *PLoS Genet* **13**:e1006613.

Allada R, White NE, So WV, Hall JC, Rosbash M. 1998. A mutant *Drosophila* homolog of

mammalian Clock disrupts circadian rhythms and transcription of period and timeless. *Cell*

93:791–804.

Benito J, Houl JH, Roman GW, Hardin PE. 2008. The blue-light photoreceptor

CRYPTOCHROME is expressed in a subset of circadian oscillator neurons in the

Drosophila CNS. *J Biol Rhythms* **23**:296–307.

Benito J, Zheng H, Hardin PE. 2007. PDP1 Functions Downstream of the Circadian Oscillator to

Mediate Behavioral Rhythms. *Journal of Neuroscience*. doi:10.1523/jneurosci.4870-

06.2007

Bulthuis N, Spontak KR, Kleeman B, Cavanaugh DJ. 2019. Neuronal Activity in Non-LNv

Clock Cells Is Required to Produce Free-Running Rest:Activity Rhythms in *Drosophila*. *J*

Biol Rhythms 748730419841468.

Depetris-Chauvin A, Berni J, Aranovich EJ, Muraro NI, Beckwith EJ, Ceriani MF. 2011. Adult-

specific electrical silencing of pacemaker neurons uncouples molecular clock from circadian

outputs. *Curr Biol* **21**:1783–1793.

Evans JA. 2016. Collective timekeeping among cells of the master circadian clock. *J Endocrinol*

230:R27–49.

Fernández MP, Berni J, Ceriani MF. 2008. Circadian remodeling of neuronal circuits involved in

rhythmic behavior. *PLoS Biol* **6**:e69.

Gekakis N. 1998. Role of the CLOCK Protein in the Mammalian Circadian Mechanism. *Science*.

doi:10.1126/science.280.5369.1564

Grima B, Chélot E, Xia R, Rouyer F. 2004. Morning and evening peaks of activity rely on

different clock neurons of the *Drosophila* brain. *Nature* **431**:869–873.

Gummadova JO, Coutts GA, Glossop NRJ. 2009. Analysis of the *Drosophila* Clock promoter

reveals heterogeneity in expression between subgroups of central oscillator cells and

identifies a novel enhancer region. *J Biol Rhythms* **24**:353–367.

Guo F, Cerullo I, Chen X, Rosbash M. 2014. PDF neuron firing phase-shifts key circadian

activity neurons in *Drosophila*. *Elife* **3**. doi:10.7554/eLife.02780

Guo F, Chen X, Rosbash M. 2017. Temporal calcium profiling of specific circadian neurons in

freely moving flies. *Proc Natl Acad Sci U S A* **114**:E8780–E8787.

Guo F, Holla M, Díaz MM, Rosbash M. 2018. A Circadian Output Circuit Controls Sleep-Wake

Arousal in *Drosophila*. *Neuron* **100**:624–635.e4.

Guo F, Yu J, Jung HJ, Abruzzi KC, Luo W, Griffith LC, Rosbash M. 2016. Circadian neuron

feedback controls the *Drosophila* sleep--activity profile. *Nature* **536**:292–297.

Hamada FN, Rosenzweig M, Kang K, Pulver SR, Ghezzi A, Jegla TJ, Garrity PA. 2008. An

internal thermal sensor controlling temperature preference in *Drosophila*. *Nature* **454**:217–

220.

Hardin PE. 2011. Molecular genetic analysis of circadian timekeeping in *Drosophila*. *Adv Genet*

74:141–173.

Helfrich-Förster C. 1997. Development of pigment-dispersing hormone-immunoreactive neurons

in the nervous system of *Drosophila melanogaster*. *J Comp Neurol* **380**:335–354.

Helfrich-Förster C, Yoshii T, Wülbeck C, Grieshaber E, Rieger D, Bachleitner W, Cusumano P,

Rouyer F. 2007. The lateral and dorsal neurons of *Drosophila melanogaster:* new insights

about their morphology and function. *Cold Spring Harb Symp Quant Biol* **72**:517–525.

Im SH, Li W, Taghert PH. 2011. PDFR and CRY Signaling Converge in a Subset of Clock

Neurons to Modulate the Amplitude and Phase of Circadian Behavior in *Drosophila*. *PLoS*

ONE. doi:10.1371/journal.pone.0018974

Ito C, Tomioka K. 2016. Heterogeneity of the Peripheral Circadian Systems in *Drosophila*

melanogaster: A Review. *Front Physiol* **7**:8.

Johns DC, Marx R, Mains RE, O'Rourke B, Marbán E. 1999. Inducible genetic suppression of

neuronal excitability. *J Neurosci* **19**:1691–1697.

Kistenpfennig C, Nakayama M, Nihara R, Tomioka K, Helfrich-Förster C, Yoshii T. 2018. A

Tug-of-War between Cryptochrome and the Visual System Allows the Adaptation of

Evening Activity to Long Photoperiods in *Drosophila melanogaster*. *J Biol Rhythms* **33**:24–

34.

Konopka RJ, Benzer S. 1971. Clock mutants of *Drosophila melanogaster*. *Proc Natl Acad Sci U*

S A **68**:2112–2116.

Lamaze A, Krätschmer P, Chen K-F, Lowe S, Jepson JEC. 2018. A Wake-Promoting Circadian

Output Circuit in *Drosophila*. *Curr Biol* **28**:3098–3105.e3.

Lin Y, Stormo GD, Taghert PH. 2004. The neuropeptide pigment-dispersing factor coordinates

pacemaker interactions in the *Drosophila* circadian system. *J Neurosci* **24**:7951–7957.

Li Y, Guo F, Shen J, Rosbash M. 2014. PDF and cAMP enhance PER stability in *Drosophila*

clock neurons. *Proc Natl Acad Sci U S A* **111**:E1284–90.

Long X, Colonell J, Wong AM, Singer RH, Lionnet T. 2017. Quantitative mRNA imaging

throughout the entire Drosophila brain. *Nature Methods*. doi:10.1038/nmeth.4309

McGuire SE, Le PT, Osborn AJ, Matsumoto K, Davis RL. 2003. Spatiotemporal rescue of

memory dysfunction in *Drosophila*. *Science* **302**:1765–1768.

Menegazzi P, Vanin S, Yoshii T, Rieger D, Hermann C, Dusik V, Kyriacou CP, Helfrich-Förster

C, Costa R. 2013. Drosophila clock neurons under natural conditions. *J Biol Rhythms* **28**:3–

14.

Mohawk JA, Green CB, Takahashi JS. 2012. Central and peripheral circadian clocks in

mammals. *Annu Rev Neurosci* **35**:445–462.

Ni JD, Gurav AS, Liu W, Ogunmowo TH, Hackbart H, Elsheikh A, Verdegaal AA, Montell C.

2019. Differential regulation of the sleep homeostat by circadian and arousal inputs. *Elife* **8**.

doi:10.7554/eLife.40487

Nitabach MN, Blau J, Holmes TC. 2002. Electrical silencing of *Drosophila* pacemaker neurons

stops the free-running circadian clock. *Cell* **109**:485–495.

Port F, Bullock SL. 2016. Augmenting CRISPR applications in *Drosophila* with tRNA-flanked

sgRNAs. *Nat Methods* **13**:852–854.

Reischig T, Stengl M. 2003. Ultrastructure of pigment-dispersing hormone-immunoreactive

neurons in a three-dimensional model of the accessory medulla of the cockroach

Leucophaea maderae. *Cell Tissue Res* **314**:421–435.

Renn SC, Park JH, Rosbash M, Hall JC, Taghert PH. 1999. A *pdf* neuropeptide gene mutation

and ablation of PDF neurons each cause severe abnormalities of behavioral circadian

rhythms in *Drosophila*. *Cell* **99**:791–802.

Rieger D, Stanewsky R, Helfrich-Förster C. 2003. Cryptochrome, compound eyes, Hofbauer-

Buchner eyelets, and ocelli play different roles in the entrainment and masking pathway of

the locomotor activity rhythm in the fruit fly *Drosophila melanogaster*. *J Biol Rhythms*

18:377–391.

Schlichting M, Helfrich-Förster C. 2015. Photic Entrainment in *Drosophila* Assessed by

Locomotor Activity Recordings. *Methods in Enzymology*. doi:10.1016/bs.mie.2014.10.017

Schlichting M, Menegazzi P, Rosbash M, Helfrich-Förster C. 2019. A distinct visual pathway

mediates high light intensity adaptation of the circadian clock in *Drosophila*. *The Journal of*

Neuroscience. doi:10.1523/jneurosci.1497-18.2018

Schlichting M, Weidner P, Diaz M, Menegazzi P, Dalla-Benetta E, Helfrich-Foerster C, Rosbash

M. n.d. Light-mediated circuit switching in the *Drosophila* neuronal clock network.

doi:10.1101/515478

Schmid B, Helfrich-Förster C, Yoshii T. 2011. A new ImageJ plug-in "ActogramJ" for

chronobiological analyses. *J Biol Rhythms* **26**:464–467.

Shafer OT, Taghert PH. 2009. RNA-interference knockdown of *Drosophila* pigment dispersing

factor in neuronal subsets: the anatomical basis of a neuropeptide's circadian functions.

PLoS One **4**:e8298.

Shang Y, Griffith LC, Rosbash M. 2008. Light-arousal and circadian photoreception circuits

intersect at the large PDF cells of the *Drosophila* brain. *Proc Natl Acad Sci U S A*

105:19587–19594.

Sheeba V, Fogle KJ, Kaneko M, Rashid S, Chou Y-T, Sharma VK, Holmes TC. 2008. Large

ventral lateral neurons modulate arousal and sleep in *Drosophila*. *Curr Biol* **18**:1537–1545.

Shigeyoshi Y, Taguchi K, Yamamoto S, Takekida S, Yan L, Tei H, Moriya T, Shibata S, Loros

JJ, Dunlap JC, Okamura H. 1997. Light-induced resetting of a mammalian circadian clock is

associated with rapid induction of the *mPer1* transcript. *Cell* **91**:1043–1053.

Stanewsky R, Kaneko M, Emery P, Beretta B, Wager-Smith K, Kay SA, Rosbash M, Hall JC.

1998. The *cryb* mutation identifies cryptochrome as a circadian photoreceptor in

Drosophila. *Cell* **95**:681–692.

Stoleru D, Peng Y, Agosto J, Rosbash M. 2004. Coupled oscillators control morning and evening

locomotor behaviour of *Drosophila*. *Nature* **431**:862–868.

Stoleru D, Peng Y, Nawathean P, Rosbash M. 2005. A resetting signal between *Drosophila*

pacemakers synchronizes morning and evening activity. *Nature* **438**:238–242.

Webb AB, Taylor SR, Thoroughman KA, Doyle FJ, Herzog ED. 2012. Weakly Circadian Cells

Improve Resynchrony. *PLoS Computational Biology*. doi:10.1371/journal.pcbi.1002787

Welsh DK, Logothetis DE, Meister M, Reppert SM. 1995. Individual neurons dissociated from

rat suprachiasmatic nucleus express independently phased circadian firing rhythms. *Neuron*.

doi:10.1016/0896-6273(95)90214-7

Wu Y, Cao G, Nitabach MN. 2008. Electrical silencing of PDF neurons advances the phase of

non-PDF clock neurons in *Drosophila*. *J Biol Rhythms* **23**:117–128.

Yamaguchi S, Isejima H, Matsuo T, Okura R, Yagita K, Kobayashi M, Okamura H. 2003.

Synchronization of Cellular Clocks in the Suprachiasmatic Nucleus. *Science*.

doi:10.1126/science.1089287

Yao Z, Bennett AJ, Clem JL, Shafer OT. 2016. The *Drosophila* Clock Neuron Network Features

Diverse Coupling Modes and Requires Network-wide Coherence for Robust Circadian

Rhythms. *Cell Rep* **17**:2873–2881.

Yao Z, Shafer OT. 2014. The *Drosophila* circadian clock is a variably coupled network of

multiple peptidergic units. *Science* **343**:1516–1520.

Yoshii T, Todo T, Wülbeck C, Stanewsky R, Helfrich-Förster C. 2008. Cryptochrome is present

in the compound eyes and a subset of *Drosophila's* clock neurons. *J Comp Neurol* **508**:952–

966.

Zhang L, Chung BY, Lear BC, Kilman VL, Liu Y, Mahesh G, Meissner R-A, Hardin PE, Allada

R. 2010. DN1(p) circadian neurons coordinate acute light and PDF inputs to produce robust

daily behavior in *Drosophila*. *Curr Biol* **20**:591–599.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Supplement Figure 1

Supplement Figure 2

Supplement Figure 3

Supplement Figure 4

Supplement Figure 5

Supplement Figure 6