www.Malaria.tools - comparative genomic and transcriptomic database for *Plasmodium* species

Qiao Wen Tan[1], Marek Mutwil[1*]

[1]School of Biological Sciences, Nanyang Technological University, 60 Nanyang Drive, Singapore 637551, Singapore

* To whom correspondence should be addressed. Tel: +65 6904 7503; Email: mutwil@ntu.edu.sg

ABSTRACT

Malaria is a tropical parasitic disease caused by the *Plasmodium* genus, which resulted in an estimated 219 million cases of malaria and 435,000 malaria-related deaths in 2017. Despite the availability of the *P. falciparum* genome since 2002, almost 50% of the genes remain unannotated. To remedy this paucity of functional information, we used transcriptomic data to build gene co-expression networks for two *Plasmodium* species (*P. falciparum and P. berghei*), and included genomic data of four other *Plasmodium* species, *P. yoleii, P. knowlesi, P. vivax* and *P. cynomolgi*, as well as two non-*Plasmodium* species from the Apicomplexa, *Toxoplasma gondii* and *Theileria parva.* The database is preloaded with tools that allow the identification and cross-species comparison of co-expressed gene neighborhoods, clusters, and life stage-specific expression, thus providing sophisticated tools to predict gene function. Moreover, we exemplify how the tools can be used to easily identify genes relevant for pathogenicity and various life stages of the malaria parasite. The database is freely available at www.malaria.tools.

INTRODUCTION

Malaria is a widespread infectious disease transmitted by the *Anopheles* mosquito which caused an estimated 219 million cases and 435,000 malaria-related deaths in the year 2017 (1). It is caused by various *Plasmodium* species with *P. falciparum* and *P. vivax* being the two most widespread and deadly for humans (2). Over the years, varying degrees of resistance have emerged in *Plasmodium* against all drugs used for treating malaria. Hence, it is a race against time to find new treatments, and this requires an understanding of malaria biology, and more specifically, the characterisation of genes and their functions in order to develop drugs that target genes responsible for pathogenicity.

Despite the availability of genomes of the various *Plasmodium* parasites for already more than a decade, many of the genes have unknown functions. These genes are often parasite-specific (3), thus making conventional gene function prediction methods which are based on homology to genes from other organisms less efficient. To this end, alternative methods for gene function prediction such as gene co-expression networks have been developed. The rationale behind using gene co-expression networks stems from the observation that functionally related genes have similar gene expression profiles (4). Thus, by identifying clusters of genes that have highly similar expression profiles, we attain groups of

functionally related genes where the function of uncharacterised genes can be inferred from its
neighbours.

We have built the database Malaria.tools (www.malaria.tools) based on the CoNekT database
framework (5) using publicly available RNA sequencing data of two model *Plasmodium* species. This
database calculates gene co-expression networks from gene expression data obtained from more than
800 experiments and contains multiple tools to predict gene function from co-expression network
neighbours, co-expressed clusters and specific expression profiles. Additionally, an in-built phylogenetic
tree function combined with gene expression comparison allows *Plasmodium* researchers to identify
orthologs of *P. falciparum* genes. Malaria.tools provides the malaria research community a
comprehensive and highly valuable resource for an efficient characterisation of genes based on their
expression profile, and will aid in the identification of potential drug targets *in silico* in the race for new
antimalarial drugs*.*

MATERIALS AND METHODS
RNA sequencing experiments for *P. falciparum* and *P. berghei* were downloaded as fastq files from
European Nucleotide Archive (ENA) (6) via aspera v3.8.1.160447. For paired end experiments, only the
file containing the first read, designated with "_1" was downloaded. To remove possible RNA
contaminants from the parasite's hosts, the reads were mapped first against human and mouse for *P.*
falciparum and *P. berghei*, respectively. The unmapped reads were then mapped against the mosquito
vector. Human mosquito vector was used for both *Plasmodium* species as the CDS of the mouse
mosquito vector was unavailable. Unmapped reads were mapped against the respective *Plasmodium*
species*,* and experiments with at least 1 million reads and 50% of reads mapped to the *Plasmodium*
species were used to construct the database.

The mapping was done using kallisto v0.44.0 (7). Kallisto index files were generated for *Homo*
sapiens (GCF_000001405.38)*, Anopheles gambiae* (GCF_000005575.2)*, P. falciparum* 3D7 and *P.*
berghei ANKA CDS sequences with default parameters. Both paired and single libraries were treated as
single end libraries for mapping using kallisto quant for single end library with estimated fragment length
of 200bp, estimated standard deviation of 20 and pseudobam option. Unmapped reads from the output
BAM files were written into a new fastq file using samtools v1.9-52-g651bf14 (8).

In total, 206 experiments from *P. berghei* (Figure S1A) and 620 experiments from *P. falciparum*
(Figure S1B) were included in database and annotated based on the information available from NCBI
Sequence Read Archive, ENA and literature (Supplementary Table S3 (*P. berghei)* and Supplementary
Table S4 (*P. falciparum*)). The Transcripts Per Kilobase Million (TPM) values from kallisto output of the
selected experiments were represented as an expression matrix where genes are arranged in rows and
experiments in columns. The expression matrices are available in Table S1 for *P. berghei* and Table S2
for *P. falciparum.* Highest Reciprocal Rank (HRR) co-expression networks were then constructed (9).

CDS sequences, description and associated GO terms for the 8 species in the database were
obtained from the various sources described in Table 1. Superfamily and Pfam domain annotations from
interproscan v5.32-71.0 (10) were used as sequence description when descriptions were not available.
Orthologous groups of genes (Supplementary Table S5) and phylogenetic trees were obtained from
Orthofinder v1.1.8 (11) using BLAST　for sequence similarity inference with default settings. The
database was based on the CoNekt database framework (5) with default settings, where Heuristic Cluster
Chiseling Algorithm (HCCA) (9) cluster size was limited to 100 genes. Experiments involving wild type
Plasmodium were further binned according to its various life stages (ring, trophozoite, schizont, male
gametocyte and female gametocyte) and used to calculate tissue specificity. Nucleotide and protein blast
databases for blastn and blastp were created using makeblastdb v2.9.0+ (12).

RESULTS
Malaria.tools offers a wide selection of tools to query the database. For example, the user can find the
genes of interest by using BLAST, gene IDs (e.g., *PF3D7_1223100*) and keywords (e.g., rhoptry). Genes
that work together in a specific biological process or contain a particular domain can be identified by
querying the database with Gene Ontology terms (e.g., GO:0009405) or a Pfam domain (e.g.,
VSA_Rifin), respectively. The database offers multiple comparative genomic and transcriptomic tools that
allow the user to view and compare expression profiles within and across species, and to investigate the
phylogenetic and expression relationships of gene families. A full description of the features is found at
https://malaria.sbs.ntu.edu.sg/features. To exemplify some of the features of malaria.tools, we provide
three analyses showing typical case studies.

Identification of a co-expression neighborhood important for erythrocyte invasion
A co-expression neighbourhood consists of a gene of interest and its co-expressed genes (neighbors)
calculated based on Highest Reciprocal Rank (HRR). To identify co-expression neighborhoods of interest
to malaria researchers, we calculated which genes are network neighbors to already functionally
characterized genes. We identified gene *PF3D7_1223100*, which was co-expressed with 91 other genes
of which 48 (53%) are annotated with specific GO terms indicating that experimental evidence for the
gene function already exists (e.g. evidence codes EXP, IDA, IPI). The high percentage of functionally
characterized genes in this neighborhood indicates that the corresponding biological process has
received special attention from the malaria researchers, likely due to its involvement in pathogenicity.
To gain insight into the function of this neighborhood, we first studied the expression profile of
PF3D7_1223100 (https://malaria.sbs.ntu.edu.sg/sequence/view/16011). The expression profiles in
malaria.tools are available in a detailed format that showcase expression in all annotated samples (Figure
1A), and as an average expression in the major life stages of malaria (Figure 1B,
https://malaria.sbs.ntu.edu.sg/profile/view/8699). In both expression profiles, we observed that
PF3D7_1223100 and its co-expressed genes are expressed in all major life stages with particularly high
expression in the schizont stage (Figure 1A,B). Next, we retrieved publications on the functional

characterization of the genes and we found that the highly studied genes are clearly associated with functions important for erythrocyte invasion. The genes can be classified into 3 major groups relating to motility, cytoadherence and erythrocyte invasion. Genes relating to the glideosome complex [*PF3D7_0918000* (GAP50) (13), *PF3D7_1323700* (GAPM1) (14)], the inner membrane complex (15) [*PF3D7_0109000* (PHIL1) and *PF3D7_1003600* (IMC1c)] and cytoskeleton (16, 17) [*PF3D7_1251200* (coronin), *PF3D7_0932200* (Profilin) and *PF3D7_1246200* (Actin 1)] are essential for the parasite to move towards a new red blood cell through gliding motility. Upon reaching the red blood cell, merozoite surface proteins such as *PF3D7_1035400* (MSP3), *PF3D7_1335100* (MSP7), *PF3D7_1035500* (MSP6) (18) and *PF3D7_1035700* (DBLMSP) facilitate the binding of the parasite to the red blood cell. Finally, erythrocyte invasion is enabled by various genes such as enzymes [*PF3D7_0507500* (SUB1) (19), *PF3D7_1136500* (casein kinase 1) (20) and *PF3D7_0404700* (DPAP3) (21), signalling mediators (22) [*PF3D7_0934800* (PKAc), *PF3D7_1223100* (PKAr)], rhoptry proteins (23, 24) (*PF3D7_0929400* (RhopH2), *PF3D7_0905400* (RhopH3), *PF3D7_1410400* (RAP1), *PF3D7_0414900* (ARO), *PF3D7_1017100* (RON12), *PF3D7_0501600* (RAP2) and *PF3D7_0817700* (RON5)] and others [*PF3D7_0423800* (CyRPA) (25), *PF3D7_0935800* (CLAG9) (26), *PF3D7_0612700* (P12), *PF3D7_0404900* (P41) (27)]. The schizont is a non-infective life stage during the erythrocytic cycle. However, a mature schizont contains multiple merozoites, which upon rupture of the schizont moves and invades fresh erythrocytes. In the co-expression neighbourhood of *PF3D7_1223100*, we observe an upregulation of merozoite and erythrocyte invasion-related genes during the schizont stage. In conclusion, the remaining 47% of genes in this cluster that are not yet functionally characterized are prime candidates for further studies on parasite motility, cytoadherence and erythrocyte invasion.

Comparative transcriptomic analysis of gene modules involved in male gametocyte-specific motility

Comparative transcriptomic analyses can reveal which gene modules are conserved across species (28, 29), thus enabling the identification of the core genetic components of specific biological processes (29, 30). Malaria.tools provides two methods to extract these conserved transcriptional programs by (i) identifying common gene families that are specifically expressed in a particular life stage of two malaria species or by (ii) identifying conserved clusters of co-expressed genes.

Using the first method to identify conserved transcriptional programs, we navigated to 'Tools\Compare specificities', selected species *P. falciparum* and *P. berghei*, set condition 'Gametocyte (male) for both species and clicked 'Compare specificity'. The database first identified genes that are preferentially expressed in male gametocytes in both species (specificity measure (SPM) > 0.85) (31), which revealed that 187 gene families are expressed at this life stage in the two parasites (Figure 2A, Table S6). The table below the Venn diagram shows the identity and links to the 187 gene families, and clicking on the tree links of a gene family depicted the phylogenetic and expression relationships of the genes in the family. Not surprisingly, many of the gene families show conserved male gametocyte-specific expression in the two *Plasmodium* species, as exemplified by the phylogenetic tree of gene family

OG0000502 (Figure 2B, https://malaria.sbs.ntu.edu.sg/tree/view/503), indicating that this family is male

gametocyte-specific. Interestingly, we also observed cases where only one of the clades of the

phylogenetic tree showed a male gametocyte-specific expression (Figure 2C, OG0000055,

https://malaria.sbs.ntu.edu.sg/tree/view/56), suggesting that for this particular gene family an ancient

gene duplication took place in the ancestor of the *Plasmodium* species followed by either a sub-

functionalization or neo-functionalization of the genes. The list of the conserved male-specific genes and

gene families provides a good starting point to dissect the genetic basis of male gametocyte-specific

biological processes.

The second method to identify conserved gene modules using malaria.tools is based on co-

expression network clusters. The clusters are used to identify functionally related genes based on the

topology of the networks (9), i.e. similar clusters are found by identifying which cluster pairs contain

significantly similar (P<0.05, hypergeometric test) number of gene families (expressed as Jaccard

index,(5)). To exemplify this feature, we clicked first on one of the *P. berghei* genes found in the table

(*PBANKA_0102700*, https://malaria.sbs.ntu.edu.sg/sequence/view/17914) and then on the co-expression

cluster 13 containing this gene (https://malaria.sbs.ntu.edu.sg/cluster/view/40). The 'Similar clusters' table

found on this page identified *P. falciparum* cluster 15 as being significantly similar to cluster 13 (Jaccard

index = 0.246). Clicking on the 'Compare' button revealed the co-expression networks of the two modules

(Figure 2D). As expected, the two conserved clusters show male gametocyte-specific expression profiles

(https://malaria.sbs.ntu.edu.sg/cluster/view/40,   https://malaria.sbs.ntu.edu.sg/cluster/view/124), further

reinforcing that the two clusters represent a *bona fide* conserved transcriptional program for male

gametocyte-specific motility.

In summary, the analysis of conserved gene modules resulted in 132 genes present in the co-

expression networks whereof 123 genes showed male gametocyte-specific expression. However, only 60

of these homologs were annotated and the remaining ones were conserved proteins of unknown function

(4 genes) or conserved *Plasmodium* proteins of unknown function (59 genes, Table S7). A closer

inspection of the homologs present in the clusters revealed motor proteins such as kinesin

(*PBANKA_0202700*, *PF3D7_0111000*, *PBANKA_0902400*, *PF3D7_1146700*, *PBANKA_1458800* and

PF3D7_1245600) and dynein (*PBANKA_1022400* and *PF3D7_1420800*) (32), as well as flagella-related

proteins such as the radial spoke protein 3 (*PBANKA_1039000*), growth arrest protein (33)

(*PBANKA_1455800* and *PF3D7_1242400*) and MORN repeat containing protein which localises near the

flagellar basal body in male gametocytes (34) (*PBANKA_1018200* and *PF3D7_1426400*). Taken

together, the functions of these genes suggest an overall motility and flagella-related function associated

with the clusters. Hence, the unannotated genes in these homologous networks will be of prime interest

for researchers interested in male gametocyte motility.

Identification of microneme-specific co-expressed gene clusters

Micronemes are protein rich, secretory organelles important for host-cell invasion and gliding motility in parasitic Apicomplexans (35). Proteins are being discharged to facilitate entry of the parasites into red blood cells.

To gain insight into microneme biogenesis and function using malaria.tools, we entered GO:0020009 (GO term for microneme) to arrive at the page dedicated to microneme cellular component (https://malaria.sbs.ntu.edu.sg/go/view/12866). The page revealed 270 and 299 annotated microneme-associated genes in *P. berghei* and *P. falciparum*, respectively. Furthermore, the page contains information about Pfam domains (Prot_kinase_dom, VWF_A, MORN and others) and gene families (OG_01_0000012, OG_01_0000038 and others), which may also play a role in the microneme function. Furthermore, the database identified cluster 2 (https://malaria.sbs.ntu.edu.sg/cluster/view/31) and cluster 7 (https://malaria.sbs.ntu.edu.sg/cluster/view/33) from *P. berghei* and cluster 2 (https://malaria.sbs.ntu.edu.sg/cluster/view/74) and cluster 12 (https://malaria.sbs.ntu.edu.sg/cluster/view/146) from *P. falciparum* as being significantly similar (P<0.05, Benjamini-Hochenberg corrected p-value)(36), implicating these clusters in a microneme-specific process.

To learn more about the function of these four clusters, we investigated their expression profiles. While cluster 2 from *P. berghei* and cluster 12 from *P. falciparum* show ubiquitous expression at all life stages of malaria (Figure 3A), cluster 7 from *P.berghei* and cluster 2 from *P. falciparum* show ookinete- and sporozoite-specific expression respectively. Since ookinetes and sporozoites are mosquito stage-specific, we speculate that the two *Plasmodium* species have at least two types of micronemes, one being ubiquitously expressed (clusters 2 and 12) and another being mosquito-specific (clusters 7 and 2).

We further investigated the putative function of the ookinete-specific cluster 7 from *P. berghei* (https://malaria.sbs.ntu.edu.sg/cluster/view/33), and found that the cluster is significantly enriched for GO terms such as entry into host cell (GO:0030260, Table S8), which is in line with the microneme being involved in invasion of blood cells. A closer look at the genes found in cluster 7 from *P. berghei* revealed that it contains genes that are essential for the infectivity and maturation of ookinetes (Figure 3B). Specifically, a group of genes relating to the inner membrane complex, surface ookinete protein and secreted ookinete protein is required for efficient gliding motility and midgut traversal [*PBANKA_1354600* (ISC1), *PBANKA_1025700* (IMC1l), *PBANKA_0513000* (IMC1m) (37), *PBANKA_1106900* (PIMMS2) (38), *PBANKA_0714300* (HSP20) (39) and *PBANKA_1432300* (CelTOS) (40)]. Another group of co-expressed genes contains perforins and a secreted protein, which are known to be important in midgut invasion where the parasite disrupts the membrane of the endothelial cell [*PBANKA_0824200* (PLP3), *PBANKA_0711400* (PLP4), *PBANKA_0711600* (PLP5) (41) and *PBANKA_1037800* (SOAP) (42)]. Last but not least, genes important for the transition from the ookinete to oocyst stage are also found enriched in this cluster [*PBANKA_0701900* (GAMA/PSOP9) (43), *PBANKA_1119200* (PSOP25) (44) and

PBANKA_0412900 (CTRP) (45)]. Taken together, cluster 7 from *P. berghei* contain genes that are most
likely important for microneme function and host cell invasion.

CONCLUSIONS
The lack of comparative genomic and transcriptomic resources for malaria prompted us to construct
malaria.tools, a state-of-the-art database containing a wide range of user-friendly features. The database
can be queried by BLAST, gene IDs, keywords and Pfam domains and Gene Ontology searches. To
identify novel genes relevant for a biological process of interest, the co-expression neighborhoods and
clusters can be mined for uncharacterized candidates that are connected to well-studied genes.
Alternatively, the database allows an easy identification of genes that are expressed during a specific life
stage of the malaria parasite, thus allowing researchers to dissect the transcriptome critical for
pathogenicity and other life stages. Finally, the database can compare the clusters and stage-specific
expression profiles to identify the conserved core components of various biological processes. We
envision that malaria.tools will aid malaria researchers in selecting relevant genes for experimental
functional characterisation and potential drug development for successful combating the emerging drug
resistances.
232

233 **DATA AVAILABILITY**
234 The expression matrices, RNAseq sample annotation and gene families are available from the
235 supplementary material. The co-expression networks, coding and protein sequences can be downloaded
236 from malaria.tools.
237

238 **SUPPLEMENTARY DATA**
239 The Supplementary Data are available at NAR Online.
240

ACKNOWLEDGEMENTS
Malaria.tools is hosted at Nanyang Technological University Singapore and we would like to thank Ryan
Chee Kiang Ng for excellent tech support. We would like to thank Dr. Daniela Mutwil-Anderwald for
proofreading the manuscript. Furthermore, we would like to thank Dr. Lei Zhu from Prof. Zbynek Bozdech
lab, SBS, NTU for useful discussions.
Author Contributions: Malaria.tools was implemented by Q.W.T. who also prepared the data and built
malaria.tools with input from M.M. Both Q.W.T. and M.M. wrote the manuscript.
248

FUNDING

We would like to thank Nanyang Technological University Start-Up Grant for funding.

Conflict of interest statement. None declared.

REFERENCES

1. World Health Organisation (2018) World Malaria Report 2018 Geneva.

2. Thu,A.M., Phyo,A.P., Landier,J., Parker,D.M. and Nosten,F.H. (2017) Combating multidrug-
resistant Plasmodium falciparum malaria. *FEBS J.*, **284**, 2569–2578.
https://doi.org/10.1111/febs.14127
http://www.ncbi.nlm.nih.gov/pubmed/28580606

3. Florent,I., Maréchal,E., Gascuel,O. and Bréhélin,L. (2010) Bioinformatic strategies to provide
functional clues to the unknown genes in Plasmodium falciparum genome.

4. Zhou,X., Kao,M.-C.J. and Wong,W.H. (2002) Transitive functional annotation by shortest-path
analysis of gene expression data. *Proc. Natl. Acad. Sci. U. S. A.*, **99**, 12783–12788.
https://doi.org/10.1073/pnas.192159399
http://www.ncbi.nlm.nih.gov/pubmed/12196633

5. Proost,S. and Mutwil,M. (2018) CoNekT: an open-source framework for comparative genomic
and transcriptomic network analyses. *bioRxiv*, 10.1101/255075.
https://doi.org/10.1101/255075

6. Silvester,N., Alako,B., Amid,C., Cerdeño-Tarrága,A., Clarke,L., Cleland,I., Harrison,P.W.,
Jayathilaka,S., Kay,S., Keane,T., *et al.* (2017) The European Nucleotide Archive in 2017. *Nucleic*
Acids Res., **46**, D36–D40.
https://doi.org/10.1093/nar/gkx1125

7. Bray,N.L., Pimentel,H., Melsted,P. and Pachter,L. (2016) Near-optimal probabilistic RNA-seq
quantification. *Nat. Biotechnol.*, **34**, 525–527.
https://doi.org/10.1038/nbt.3519
http://www.ncbi.nlm.nih.gov/pubmed/27043002

8. Li,H., Handsaker,B., Wysoker,A., Fennell,T., Ruan,J., Homer,N., Marth,G., Abecasis,G. and
Durbin,R. (2009) The Sequence Alignment/Map format and SAMtools. *Bioinformatics*, **25**, 2078–
2079.
https://doi.org/10.1093/bioinformatics/btp352
http://www.ncbi.nlm.nih.gov/pubmed/19505943
290
291 9. Mutwil,M., Usadel,B., Schütte,M., Loraine,A., Ebenhöh,O. and Persson,S. (2010) Assembly of an
292 interactive correlation network for the Arabidopsis genome using a novel heuristic clustering
293 algorithm. *Plant Physiol.*, **152**, 29–43.
294 https://doi.org/10.1104/pp.109.145318
295 http://www.ncbi.nlm.nih.gov/pubmed/19889879
296
297 10. Jones,P., Binns,D., Chang,H.-Y., Fraser,M., Li,W., McAnulla,C., McWilliam,H., Maslen,J.,
298 Mitchell,A., Nuka,G., *et al.* (2014) InterProScan 5: genome-scale protein function classification.
299 *Bioinformatics*, **30**, 1236–1240.
300 https://doi.org/10.1093/bioinformatics/btu031
301
302 11. Emms,D.M. and Kelly,S. (2015) OrthoFinder: solving fundamental biases in whole genome
303 comparisons dramatically improves orthogroup inference accuracy. *Genome Biol.*, **16**.
304 https://doi.org/10.1186/s13059-015-0721-2
305 http://www.ncbi.nlm.nih.gov/pubmed/26243257
306
307 12. Camacho,C., Coulouris,G., Avagyan,V., Ma,N., Papadopoulos,J., Bealer,K. and Madden,T.L.
308 (2009) BLAST+: architecture and applications. *BMC Bioinformatics*, **10**, 421.
309 https://doi.org/10.1186/1471-2105-10-421
310 http://www.ncbi.nlm.nih.gov/pubmed/20003500
311
312 13. Yeoman,J.A., Hanssen,E., Maier,A.G., Klonis,N., Maco,B., Baum,J., Turnbull,L.,
313 Whitchurch,C.B., Dixon,M.W.A. and Tilley,L. (2011) Tracking Glideosome-Associated Protein 50
314 Reveals the Development and Organization of the Inner Membrane Complex of Plasmodium
315 falciparum. *Eukaryot. Cell*, **10**, 556–564.
316 https://doi.org/10.1128/EC.00244-10
317
14. Bullen,H.E., Tonkin,C.J., O'Donnell,R.A., Tham,W.-H., Papenfuss,A.T., Gould,S., Cowman,A.F.,
Crabb,B.S. and Gilson,P.R. (2009) A Novel Family of Apicomplexan Glideosome-associated
Proteins with an Inner Membrane-anchoring Role. *J. Biol. Chem.* , **284**, 25353–25363.
https://doi.org/10.1074/jbc.M109.036772

15. Saini,E., Zeeshan,M., Brady,D., Pandey,R., Kaiser,G., Koreny,L., Kumar,P., Thakur,V.,

Tatiya,S., Katris,N.J., *et al.* (2017) Photosensitized INA-Labelled protein 1 (PhIL1) is novel

component of the inner membrane complex and is required for Plasmodium parasite development.

Sci. Rep., **7**, 15577.

https://doi.org/10.1038/s41598-017-15781-z

http://www.ncbi.nlm.nih.gov/pubmed/29138437

16. Olshina,M.A., Angrisano,F., Marapana,D.S., Riglar,D.T., Bane,K., Wong,W., Catimel,B., Yin,M.-

X., Holmes,A.B., Frischknecht,F., *et al.* (2015) Plasmodium falciparum coronin organizes arrays of

parallel actin filaments potentially guiding directional motility in invasive malaria parasites. *Malar. J.*,

14, 280.

https://doi.org/10.1186/s12936-015-0801-5

17. Moreau,C.A., Bhargav,S.P., Kumar,H., Quadt,K.A., Piirainen,H., Strauss,L., Kehrer,J.,

Streichfuss,M., Spatz,J.P., Wade,R.C., *et al.* (2017) A unique profilin-actin interface is important for

malaria parasite motility. *PLoS Pathog.*, **13**, e1006412–e1006412.

https://doi.org/10.1371/journal.ppat.1006412

18. Beeson,J.G., Drew,D.R., Boyle,M.J., Feng,G., Fowkes,F.J.I. and Richards,J.S. (2016) Merozoite

surface proteins in red blood cell invasion, immunity and vaccines against malaria. *FEMS Microbiol.*

Rev., **40**, 343–372.

https://doi.org/10.1093/femsre/fuw001

19. Withers-Martinez,C., Suarez,C., Fulle,S., Kher,S., Penzo,M., Ebejer,J.-P., Koussis,K.,

Hackett,F., Jirgensons,A., Finn,P., *et al.* (2012) Plasmodium subtilisin-like protease 1 (SUB1):

insights into the active-site structure, specificity and function of a pan-malaria drug target. *Int. J.*

Parasitol., **42**, 597–612.

https://doi.org/10.1016/j.ijpara.2012.04.005

20. Dorin-Semblat,D., Demarta-Gatsi,C., Hamelin,R., Armand,F., Carvalho,T.G., Moniatte,M. and

Doerig,C. (2015) Malaria Parasite-Infected Erythrocytes Secrete PfCK1, the Plasmodium

Homologue of the Pleiotropic Protein Kinase Casein Kinase 1. *PLoS One*, **10**, e0139591.

21. Lehmann,C., Tan,M.S.Y., de Vries,L.E., Russo,I., Sanchez,M.I., Goldberg,D.E. and Deu,E.

(2018) Plasmodium falciparum dipeptidyl aminopeptidase 3 activity is important for efficient

erythrocyte invasion by the malaria parasite. *PLoS Pathog.*, **14**, e1007031.

https://doi.org/10.1371/journal.ppat.1007031
http://www.ncbi.nlm.nih.gov/pubmed/29768491
361
362   22. Baker,D.A., Drought,L.G., Flueck,C., Nofal,S.D., Patel,A., Penzo,M. and Walker,E.M. (2017)
363   Cyclic nucleotide signalling in malaria parasites. *Open Biol.*, **7**, 170213.
364   https://doi.org/10.1098/rsob.170213
365
366   23. Counihan,N.A., Kalanon,M., Coppel,R.L. and De Koning-Ward,T.F. (2013) Plasmodium rhoptry
367   proteins: Why order is important. *Trends Parasitol.*, **29**, 228–236.
368   https://doi.org/10.1016/j.pt.2013.03.003
369
370   24. Counihan,N.A., Chisholm,S.A., Bullen,H.E., Srivastava,A., Sanders,P.R., Jonsdottir,T.K.,
371   Weiss,G.E., Ghosh,S., Crabb,B.S., Creek,D.J., *et al.* (2017) Plasmodium falciparum parasites
372   deploy RhopH2 into the host erythrocyte to obtain nutrients, grow and replicate. *Elife*, **6**.
373   https://doi.org/10.7554/eLife.23217
374   http://www.ncbi.nlm.nih.gov/pubmed/28252383
375
376   25. Favuzza,P., Guffart,E., Tamborrini,M., Scherer,B., Dreyer,A.M., Rufer,A.C., Erny,J.,
377   Hoernschemeyer,J., Thoma,R., Schmid,G., *et al.* (2017) Structure of the malaria vaccine candidate
378   antigen CyRPA and its complex with a parasite invasion inhibitory antibody. *Elife*, **6**.
379   https://doi.org/10.7554/eLife.20383
380
26. Ling,I.T., Florens,L., Dluzewski,A.R., Kaneko,O., Grainger,M., Yim Lim,B.Y.S., Tsuboi,T.,
Hopkins,J.M., Johnson,J.R., Torii,M., *et al.* (2004) The Plasmodium falciparum clag9 gene encodes
a rhoptry protein that is transferred to the host erythrocyte upon invasion. *Mol. Microbiol.*, **52**, 107–
118.
https://doi.org/10.1111/j.1365-2958.2003.03969.x
http://www.ncbi.nlm.nih.gov/pubmed/15049814
387
27. Tonkin,M.L., Arredondo,S.A., Loveless,B.C., Serpa,J.J., Makepeace,K.A.T., Sundar,N.,
Petrotchenko,E. V, Miller,L.H., Grigg,M.E. and Boulanger,M.J. (2013) Structural and biochemical
characterization of Plasmodium falciparum 12 (Pf12) reveals a unique interdomain organization and
the potential for an antiparallel arrangement with Pf41. *J. Biol. Chem.*, **288**, 12805–12817.
https://doi.org/10.1074/jbc.M113.455667
http://www.ncbi.nlm.nih.gov/pubmed/23511632
394
395   28. Mutwil,M., Klie,S., Tohge,T., Giorgi,F.M., Wilkins,O., Campbell,M.M., Fernie,A.R., Usadel,B.,

396 Nikoloski,Z. and Persson,S. (2011) PlaNet: Combined Sequence and Expression Comparisons

397 across Plant Networks Derived from Seven Species. *Plant Cell*, 10.1105/tpc.111.083667.

398 https://doi.org/10.1105/tpc.111.083667

399 http://www.ncbi.nlm.nih.gov/pubmed/21441431

401 29. Movahedi,S., Van Bel,M., Heyndrickx,K.S. and Vandepoele,K. (2012) Comparative co-

402 expression analysis in plant biology. *Plant, Cell Environ.*, **35**, 1787–1798.

403 https://doi.org/10.1111/j.1365-3040.2012.02517.x

404 http://www.ncbi.nlm.nih.gov/pubmed/22489681

406 30. Hansen,B.O., Vaid,N., Musialak-Lange,M., Janowski,M. and Mutwil,M. (2014) Elucidating gene

407 function and function evolution through comparison of co-expression networks of plants. *Front. Plant*

408 *Sci.*, **5**, 1–9.

409 https://doi.org/10.3389/fpls.2014.00394

410 http://www.ncbi.nlm.nih.gov/pubmed/25191328

412 31. Xiao,S.J., Zhang,C., Zou,Q. and Ji,Z.L. (2010) TiSGeD: A database for tissue-specific genes.

413 *Bioinformatics*, **26**, 1273–1275.

414 https://doi.org/10.1093/bioinformatics/btq109

415 http://www.ncbi.nlm.nih.gov/pubmed/20223836

417 32. Talman,A.M., Prieto,J.H., Marques,S., Ubaida-Mohien,C., Lawniczak,M., Wass,M.N., Xu,T.,

418 Frank,R., Ecker,A., Stanway,R.S., *et al.* (2014) Proteomic analysis of the Plasmodium male gamete

419 reveals the key role for glycolysis in flagellar motility. *Malar. J.*, **13**, 315.

420 https://doi.org/10.1186/1475-2875-13-315

422 33. Yeh,S.-D., Chen,Y.-J., Chang,A.C.Y., Ray,R., She,B.-R., Lee,W.-S., Chiang,H.-S., Cohen,S.N.

423 and Lin-Chao,S. (2002) Isolation and properties of Gas8, a growth arrest-specific gene regulated

424 during male gametogenesis to produce a protein associated with the sperm motility apparatus. *J.*

425 *Biol. Chem.*, **277**, 6311–7.

426 https://doi.org/10.1074/jbc.M106941200

427 http://www.ncbi.nlm.nih.gov/pubmed/11751847

429 34. Ferguson,D.J.P., Sahoo,N., Pinches,R.A., Bumstead,J.M., Tomley,F.M. and Gubbels,M.-J.

430 (2008) MORN1 Has a Conserved Role in Asexual and Sexual Development across the

431 Apicomplexa. *Eukaryot. Cell*, **7**, 698–711.

432 https://doi.org/10.1128/EC.00021-08

35. Black,M.W. and Boothroyd,J.C. (2000) Lytic cycle of Toxoplasma gondii. *Microbiol. Mol. Biol.*

Rev., **64**, 607–623.

36. Benjamini,Y. and Hochberg,Y. (1995) Controlling the False Discovery Rate: A Practical and

Powerful Approach to Multiple Testing. *J. R. Stat. Soc. Ser. B*, **57**, 289–300.

37. Harding,C.R. and Meissner,M. (2014) The inner membrane complex through development of

Toxoplasma gondii and Plasmodium. *Cell. Microbiol.*, **16**, 632–641.

https://doi.org/10.1111/cmi.12285

38. Ukegbu,C. V, Akinosoglou,K.A., Christophides,G.K. and Vlachou,D. (2017) Plasmodium berghei

PIMMS2 Promotes Ookinete Invasion of the Anopheles gambiae Mosquito Midgut. *Infect. Immun.*,

85, e00139-17.

https://doi.org/10.1128/IAI.00139-17

39. Montagna,G.N., Buscaglia,C.A., Münter,S., Goosmann,C., Frischknecht,F., Brinkmann,V. and

Matuschewski,K. (2012) Critical Role for Heat Shock Protein 20 (HSP20) in Migration of Malarial

Sporozoites. *J. Biol. Chem.*, **287**, 2410–2422.

https://doi.org/10.1074/jbc.M111.302109

40. Kariu,T., Ishino,T., Yano,K., Chinzei,Y. and Yuda,M. (2006) CelTOS, a novel malarial protein

that mediates transmission to mosquito and vertebrate hosts. *Mol. Microbiol.*, **59**, 1369–1379.

https://doi.org/10.1111/j.1365-2958.2005.05024.x

41. Deligianni,E., Silmon de Monerri,N.C., McMillan,P.J., Bertuccini,L., Superti,F., Manola,M.,

Spanos,L., Louis,C., Blackman,M.J., Tilley,L., *et al.* (2018) Essential role of Plasmodium perforin-

like protein 4 in ookinete midgut passage. *PLoS One*, **13**, e0201651.

https://doi.org/10.1371/journal.pone.0201651

42. Dessens,J.T., Siden-Kiamos,I., Mendoza,J., Mahairaki,V., Khater,E., Vlachou,D., Xu,X.-J.,

Kafatos,F.C., Louis,C., Dimopoulos,G., *et al.* (2003) SOAP, a novel malaria ookinete protein

involved in mosquito midgut invasion and oocyst development. *Mol. Microbiol.*, **49**, 319–329.

http://www.ncbi.nlm.nih.gov/pubmed/12828632

43. Ecker,A., Bushell,E.S.C., Tewari,R. and Sinden,R.E. (2008) Reverse genetics screen identifies

six proteins important for malaria development in the mosquito. *Mol. Microbiol.*, **70**, 209–220.

https://doi.org/10.1111/j.1365-2958.2008.06407.x

44. Zheng,W., Liu,F., He,Y., Liu,Q., Humphreys,G.B., Tsuboi,T., Fan,Q., Luo,E., Cao,Y. and Cui,L.

(2017) Functional characterization of Plasmodium berghei PSOP25 during ookinete development

and as a malaria transmission-blocking vaccine candidate. *Parasit. Vectors*, **10**, 8.

https://doi.org/10.1186/s13071-016-1932-4

45. Ramakrishnan,C., Dessens,J.T., Armson,R., Pinto,S.B., Talman,A.M., Blagborough,A.M. and

Sinden,R.E. (2011) Vital functions of the malarial ookinete protein, CTRP, reside in the A domains.

Int. J. Parasitol., **41**, 1029–1039.

https://doi.org/10.1016/j.ijpara.2011.05.007

TABLE AND FIGURES LEGENDS

Table 1. Source of data for species in database.

Species	RNA-seq samples	CDS	CDS description	GO annotation
Theilera parva Muguga	N/A	NCBI Ref-seq	NCBI Ref-seq	Interproscan
Toxoplasma gondii ME49	N/A	NCBI Ref-seq	NCBI Ref-seq	Interproscan
Plasmodium berghei ANKA	206	PlasmoDB	GeneDB	PlasmoDB
Plasmodium cynomolgi B	N/A	PlasmoDB	Interproscan	Interproscan
Plasmodium falciparum 3D7	620	PlasmoDB	GeneDB	PlasmoDB
Plasmodium knowlesi Malayan Pk1A	N/A	PlasmoDB	Interproscan	Interproscan
Plasmodium vivax P01	N/A	PlasmoDB	GeneDB	Interproscan
Plasmodium yoelii YM	N/A	PlasmoDB	Interproscan	Interproscan

Figure 1. Expression profiles and co-expression neighborhood of gene *PF3D7_1223100*. A) Full

expression profile of the gene. The x-axis represents the different RNA-seq experiments capturing the life

stages and genetic perturbations of *P. falciparum*, while the y-axis indicates the expression level

(Transcripts Per Million, TPM). The different life stages are color-coded by blue (ring), purple

(trophozoite), green (schizont), light blue (male gametocyte), pink (female gametocyte) and brown

(sporozoite). The bars indicate the mean expression value, while the dots show the expression values of

the individual samples. For brevity, only the general descriptions of the samples are shown. B) Simplified

expression profile of *PF3D7_1223100*, showing the average expression in the five major life stages. C)

Co-expression network neighborhood containing functionally characterized genes. Nodes represent
genes, edges (lines) connect co-expressed genes, while colored shapes indicate orthogroups. For
brevity, only genes with experimentally verified function supported by at least two publications are shown
in the figure.

Figure 2. Comparative analysis of male gametocyte-specific gene expression in *P. berghei* and *P.*
falciparum. A) Venn diagram showing the overlap of the male gametophyte-specific gene families in the
two malaria species. B) Phylogenetic gene tree of gene family OG0109400. The different species are
color coded and represent *Toxoplasma gondii* in red (gene IDs XP_NNNNNNNNN, NP_XXXXXX),
Theilera parva in orange (gene IDs XP_NNNNNN), *P. falciparum* in olive (gene IDs PF3D7_XXXXXXX),
P. berghei in dark green (gene IDs PBANKA_XXXXXXX), *P. yoleii* in light green (gene IDs
PYYM_XXXXXXX), *P. knowlesi* in blue (gene IDs PKNOH_SXXXXXXXXX-t35_1), *P. vivax* in dark blue
(gene IDs PVP01_XXXXXXX) and *P. cynomolgi* in purple (gene IDs PCYB_XXXXXX-t26_1). The colored
boxes to the right of the gene IDs show the average gene expression in five major life stages of the
malaria parasite, where yellow and blue color indicates low and high gene expression. C) Phylogenetic
gene tree of gene family OG0000055. D) Comparison of cluster 13 (left blue box) and cluster 15 (right
green box) from *P. berghei* and *P. falciparum*, respectively. Nodes represent, genes, solid edges connect
co-expressed genes, dashed edges connect orthologs, while colored shapes indicate genes belonging to
the same gene families. E) Average expression profiles of the genes found in cluster 13 (left) and 15
(right).

Figure 3. Expression profiles and co-expression network of the ookinete-enriched clusters in *P.*
falciparum and *P. berghei*. A) Expression profiles of clusters 2 (first) and 7 (second) from *P. berghei* and
clusters 2 (third) and 12 (fourth) from *P. falciparum*. The different life stages are color-coded. For brevity,
the sample annotations are abbreviated to the major life stages. B) Co-expression cluster 7 from *P.*
berghei. Nodes represent genes, co-expressed genes are connected by gray edges, while colored
shapes indicate orthogroups. For brevity, only the discussed genes are highlighted.

520
Table S1. TPM expression matrix for *Plasmodium berghei*. Genes are found in rows, while samples are
found in columns.

Table S2. TPM expression matrix for *Plasmodium falciparum*. Genes are found in rows, while samples
are found in columns.

Table S3. Annotation of RNAseq samples for *Plasmodium berghei*.

Table S4. Annotation of RNAseq samples for *Plasmodium falciparum*.

Table S5. Orthogroups of the eight Apicomplexa species included in this study. Each row represents one

orthogroup and the genes belonging to the orthogroup.

Table S6. Male gametocyte specific genes of *Plasmodium falciparum* and *Plasmodium berghei*.

Table S7. Annotation of comparative clusters *Plasmodium berghei* cluster 13 and *Plasmodium falciparum*

cluster 15

Table S8. Enriched GO terms of *P. berghei* HCCA cluster 7 and genes associated with it.

Figure S1. Overview of the sample quality of the RNA-seq samples that contain at least 1 million reads.

A) *Plasmodium berghei*. For each sample, the percentage of reads mapping to the parasite (grey), mouse

or human (blue), mosquito (orange) and not mapping to either (yellow) is shown. The sample cutoff is

indicated by the red line. B) Overview of the sample quality of *Plasmodium falciparum*.

Figure 1

Figure 2

Figure 3

A

Percentage mapped (*P. berghei*)

B

Percentage mapped (*P. falciparum*)