1 **Gene knock-ins in *Drosophila* using homology-independent insertion of universal**

2 **donor plasmids**

4 Justin A. Bosch[1], Ryan Colbeth[1], Jonathan Zirin[1], Norbert Perrimon[1,2]

6 [1] Department of Genetics, Harvard Medical School, Boston, MA, 02115, USA

7 [2] Howard Hughes Medical Institute, Boston, MA, 02115, USA

Short Running Title: Homology-independent knock-ins in flies

Keywords: CRISPR-Cas9, knock-in, homology-independent, gene function, CRISPaint

100-word article summary: We report a new homology-independent genomic knock-in

method in *Drosophila* to insert large DNA elements into any target gene. Using

CRISPR-Cas9 and non-homologous end joining (NHEJ), an entire donor plasmid is

inserted into the genome without the need for homology arms. This approach eliminates

the burden associated with designing and constructing traditional donor plasmids. We

demonstrate its usefulness in cultured cells and in vivo to fluorescently tag endogenous

proteins, generate reporters of gene expression, and disrupt gene function.

Corresponding author:

Norbert Perrimon

Harvard Medical School

77 Avenue Louis Pasteur

Dept. of Genetics, NRB 336

Boston, MA 02115

617-432-7672

Email: perrimon@genetics.med.harvard.edu

Abstract

Site-specific insertion of DNA into endogenous genes (knock-in) is a powerful method to

study gene function. However, traditional methods for knock-in require laborious cloning

of long homology arms for homology-directed repair. Here, we report a simplified

method in *Drosophila melanogaster* to insert large DNA elements into any gene using

homology-independent repair. This method, known as CRISPaint, employs CRISPR-

Cas9 and non-homologous end joining (NHEJ) to linearize and insert donor plasmid

DNA into a target genomic cut site. The inclusion of commonly used elements such as

GFP on donor plasmids makes them universal, abolishing the need to create gene-

specific homology arms and greatly reducing user workload. Using this method, we

show robust gene-specific integration of donor plasmids in cultured cells and the fly

germ line. Furthermore, we use this method to analyze gene function by fluorescently

tagging endogenous proteins, disrupting gene function, and generating reporters of

gene expression. Finally, we assemble a collection of donor plasmids for germ line

knock-in that contain commonly used insert sequences. This method simplifies the

generation of site-specific large DNA insertions in *Drosophila* cell lines and fly strains,

and better enables researchers to dissect gene function in vivo.

Introduction

Insertion of DNA into the animal genome is a powerful method to study gene function. This approach is multipurpose, and can be used to visualize protein localization (Crivat and Taraska 2012; Hasse *et al.* 2016; Kanca *et al.* 2017), to disrupt gene function (Housden *et al.* 2017), to assay gene expression (Brand and Perrimon 1993; Ivics *et al.* 2009; Bouabe and Okkenhaug 2013), or to purify endogenous proteins (Kimple *et al.* 2013). Furthermore, the ability to insert large DNA elements such as promoters, protein coding sequences, or entire genes into the genome offers researchers endless options for genome modification. *Drosophila melanogaster* is an excellent animal model to analyze gene function because of its many genetic tools, fast generation time, and in vivo analysis (Venken *et al.* 2016; Korona *et al.* 2017; Bier *et al.* 2018).

The two most commonly used methods in *Drosophila* to knock-in DNA into endogenous genes involve either transposable elements or homology directed-repair (HDR). Transposable DNA elements insert randomly in the genome by a Transposase enzyme (Bellen *et al.* 2011), and cannot be used to target a user-specified gene. In contrast, HDR is used to insert DNA into a specific genomic location by cleavage at the genomic locus and precise homologous recombination of the DNA insert into the genome (Bier *et al.* 2018). Circular plasmids are commonly used as donor DNA for HDR because they can carry a large DNA insert (≤10kb) and homology arms corresponding to the target locus are added by traditional cloning techniques. While HDR is a useful method, the design and construction of unique plasmid donors for each gene is laborious. As a cloning-free alternative, synthesized single-stranded DNA (ssDNA) with short homology arms (~50-100 bp each) (Bier *et al.* 2018) or long

ssDNAs of ≤2kb (QUADROS *et al.* 2017) can be used as donors. However, ssDNA donors

are limited to relatively small insertions such as epitope tags, and, like HDR plasmid

donors, must be designed and produced for each gene that is targeted. Therefore, there

is a need for easier, faster, and cheaper alternatives to knock-in large DNA elements

into the *Drosophila* genome.

It was recently shown that large DNA elements could be knocked into a specific

target locus without homology arms, known as homology-independent insertion

(CRISTEA *et al.* 2013; MARESCA *et al.* 2013; AUER *et al.* 2014; KATIC *et al.* 2015; LACKNER

et al. 2015; SCHMID-BURGK *et al.* 2016; SUZUKI *et al.* 2016; KATOH *et al.* 2017). In this

method, simultaneous cutting of a circular donor plasmid and a genomic target-site by a

nuclease such as Cas9 results in integration of the linearized donor plasmid into the

genomic cut site by non-homologous end joining (NHEJ). This removes the need to

construct homology arms, and only requires cloning or synthesizing a gene-specific

single guide RNA (sgRNA). Furthermore, this approach is modular, since donor

plasmids containing common insert sequences (e.g. GFP) can be targeted to different

genomic locations and are thus "universal". Generating knock-ins by homology-

independent insertion has been successfully applied in human cell lines (CRISTEA *et al.*

2013; MARESCA *et al.* 2013; LACKNER *et al.* 2015; SCHMID-BURGK *et al.* 2016; KATOH *et*

al. 2017), mouse somatic cells (SUZUKI *et al.* 2016), zebrafish (AUER *et al.* 2014), and *C.*

elegans (KATIC *et al.* 2015). However, this approach has not yet been applied in

Drosophila.

Here, we show that homology-independent insertion functions effectively in

Drosophila by using the CRISPaint method. We first characterize this method in

cultured S2R+ cells, showing that a universal mNeonGreen donor plasmid can be used

to fluorescently tag endogenous proteins at their C-terminus. We then demonstrate that

this approach works *in vivo*, using a universal *T2A-Gal4* donor plasmid in the fly germ

line to obtain fly lines with insertions in a number of characterized genes. We show that

these insertions can be used as expression reporters for the target gene and to

generate loss-of-function phenotypes. Finally, we present a collection of different

universal donor plasmids for the purpose of enabling the *Drosophila* research

community to employ this method for their specific uses.

Materials and Methods

Plasmid cloning

pCFD3-frame_selector_(0, 1, or 2) plasmids (Addgene #s 127553-127555) were

cloned by ligating annealed oligos encoding sgRNAs that target the CRISPaint target

site (SCHMID-BURGK *et al.* 2016) into *pCFD3* (PORT *et al.* 2014), which contains the

Drosophila U6:3 promoter.

Additional sgRNA-encoding plasmids were generated by the TRiP

(https://fgr.hms.harvard.edu/) or obtained from Filip Port (PORT *et al.* 2015). sgRNA

plasmids targeting CDS close to the stop codon were *GP07595* (*Act5c*), *GP07596*

(*His2Av*), *GP07609* (*alphaTub84B*), and *GP07612* (*Lam*). sgRNA plasmids targeting

CDS close to the start codon were *GP06461* (*wg*), *GP02894* (*FK506-bp2*), *GP05054*

(*alphaTub84B*), *GP00225* (*esg*), *GP00364* (*Myo1a*), *GP00400* (*btl*), *GP00583* (*Mhc*),

GP01881 (*hh*), *GP03252* (*Desat1*), *GP05302* (*ap*), *pFP545* (*ebony*), and *pFP573*

(*ebony*). These sgRNAs were cloned into *pCFD3*, with the exception of those targeting

esg, *Myo1a*, *btl*, and *Mhc*, which were cloned into *pl100* (KONDO AND UEDA 2013).

pCRISPaint-T2A-Gal4-3xP3-RFP (Addgene # 127556) was constructed using

Gibson assembly (E2611, NEB) of three DNA fragments: 1) *Gal4-SV40-3xP3-RFP* was

PCR amplified from *pHD-Gal4-DsRed* (Xu et al. 2019 submitted PNAS, (GRATZ *et al.*

2014); 2) linear plasmid backbone generated by digesting *pWalium10-roe* (PERKINS *et*

al. 2015) with AscI/SacI; and 3) a synthesized double-stranded DNA fragment (gBlock,

IDT) encoding the CRISPaint target site, linker sequence, T2A, and ends that overlap

the other two fragments.

pCRISPaint-T2A-ORF-3xP3-RFP donor plasmids (Addgene #s 127557-127565)

were cloned by PCR amplifying the ORFs and Gibson cloning into *CRISPaint-T2A-*

Gal4-3xP3-RFP cut with NheI/KpnI. ORF sequences were amplified from templates as

follows: *sfGFP* [amplified from *pUAS-TransTimer* (He et al. 2019 submitted)], *LexGAD*

[amplified from p*CoinFLP-LexGAD/Gal4* (BOSCH *et al.* 2015)], *QF2* amplified from

Addgene #80274, *Cas9-T2A-GFP* (amplified from template kindly provided by Raghuvir

Viswanatha), *FLPo* (amplified from Addgene #24357), *Gal80* (amplified from Addgene

#17748), *Nluc* (amplified from Addgene #62057), *Gal4DBD*, (amplified from Addgene

#26233), and *p65* (amplified from Addgene #26234).

pCRISPaint-sfGFP-3xP3-RFP (Addgene # 127566) was cloned by PCR

amplifying *sfGFP* coding sequence and Gibson cloning into *CRISPaint-T2A-Gal4-3xP3-*

RFP cut with NotI/KpnI.

pCRISPaint-CRIMIC_phase_(0,1, or 2) (Addgene #s 127567-127569) donor

plasmids were cloned by ligating annealed oligos containing the CRISPaint target site

into *CRIMIC* [*pM37*, (LEE *et al.* 2018)] (frames 0,1,2) cut with NsiI.

pCRISPaint-TGEM_phase_(0,1,or 2) (Addgene #s 127570-127572) donor

plasmids were cloned by ligating annealed oligos containing the CRISPaint target site

into *T-GEM* (DIAO *et al.* 2015) (frames 0,1,2) cut with AgeI/NotI.

See Supplemental Table 6 for oligo and dsDNA sequences and Addgene for

plasmid sequences.

Cell culture

Drosophila S2R+ cells stably expressing Cas9 and a mCherry protein trap in *Clic*

(known as PT5/Cas9) (VISWANATHA *et al.* 2018) were cultured at 25°C using Schneider's

media (21720-024, ThermoFisher) with 10% FBS (A3912, Sigma) and 50 U/ml penicillin

strep (15070-063, ThermoFisher). S2R+ cells were transfected using Effectene

(301427, Qiagen) following the manufacturer's instructions. Plasmid mixes were

composed of sgRNA-expressing plasmids (see above) and *pCRISPaint-mNeon-PuroR*

(SCHMID-BURGK *et al.* 2016). Cells were transfected with plasmid mixes in 6-well dishes

at 1.8×10^6 cells/ml, split at a dilution of 1:6 after 3-4 days, and incubated with 2 μg/ml

Puromycin (540411 Calbiochem). Every 3-5 days, the media was replaced with fresh

Puromycin until the cultures became confluent (~12-16 days). For single-cell cloning

experiments, cultures were split 1:3 two days before sorting. Cells were resuspended in

fresh media, triturated to break up cell clumps, and pipetted into a cell straining FACS

tube (352235 Corning). Single cells expressing mNeonGreen were sorted into single

wells of a 96 well plate containing 50% conditioned media 50% fresh media using an

Aria-594 instrument at the Harvard Medical School Division of Immunology's Flow

Cytometry Facility. Once colonies were visible by eye (3-4 weeks), they were expanded

and screened for mNeonGreen fluorescence.

Fly genetics and embryo injections

Flies were maintained on standard fly food at 25°C. Fly stocks were obtained

from the Perrimon lab collection or Bloomington Stock center (indicated with BL#).

Stocks used in this study are as follows: *yw* (Perrimon Lab), *yw/Y hs-hid* (BL8846), *yw;*

nos-Cas9attP40/CyO (derived from BL78781), *yw;; nos-Cas9attP2* (derived from

BL78782), *yw; Sp hs-hid/CyO* (derived from BL7757), *yw;; Dr hs-hid/TM3,Sb* (derived

from BL7758), *UAS-2xGFP* (BL6874), *wg1-17/CyO* (BL2980), *wg1-8/CyO* (BL5351),

Df(2L)BSC291/CyO (BL23676), *Mhc[k10423]/CyO* (BL10995), Df(2L)H20/CyO

(BL3180), *Df(2L)ED8142/SM6a* (BL24135), *hh[AC]/TM3 Sb* (BL1749),

Df(3R)ED5296/TM3, Sb (BL9338), *esgG66/CyO UAS-GFP* (BL67748),

Df(2R)Exel6069/CyO (BL7551).

For embryo injections, each plasmid was column purified (Qiagen) twice, eluted

in injection buffer (100 μM NaPO4, 5 mM KCl), and adjusted to 200ng/μl. Plasmids were

mixed equally by volume, and mixes were injected into *Drosophila* embryos using

standard procedures. For targeting genes on Chr. 2, plasmid mixes were injected into

yw;; nos-Cas9attP2 embryos. For targeting genes on Chr. 3, plasmid mixes were

injected into *yw; nos-Cas9attP40/CyO* embryos. Approximately 500 embryos were

injected for each targeted gene.

Injected G0 flies were crossed with *yw*. We used *yw/Y hs-hid* to facilitate

collecting large numbers of virgin flies by incubating larvae and pupae at 37°C for 1hr.

G1 flies were screened for RFP expression in the adult eye on a Zeiss Stemi SVII

fluorescence microscope. G1 RFP+ flies were crossed with the appropriate balancer

stock (*yw; Sp hs-hid/CyO* or *yw;; Dr hs-hid/TM3,Sb)*. G2 RFP+ males that were *yellow-*

(to remove the *nos-Cas9* transgene) and balancer+ were crossed to virgins of the

appropriate balancer stock (*yw; Sp hs-hid/CyO* or *yw;; Dr hs-hid/TM3,Sb*). G3 larvae

and pupae were heat shocked at 37°C for 1hr to eliminate the *hs-hid* chromosome,

which generates a balanced stock (e.g. yw; [RFP+]/CyO).

Imaging

S2R+ cells expressing mNeonGreen were plated into wells of a glass-bottom 384

well plate (6007558, PerkinElmer). For fixed cell images, cells were incubated with 4%

paraformaldehyde for 30min, washed with PBS with .1% TritonX-100 (PBT) 3x 5min

each, stained with 1:1000 DAPI (D1306, ThermoFisher) and 1:1000 phalloidin-TRITC

(P1951, Sigma), and washed with PBS. Plates were imaged on an IN Cell Analyzer

6000 (GE) using a 20x or 60x objective. Time-lapse videos of live mNeonGreen

expressing single cell cloned lines were obtained by taking an image every minute using

a 60x objective. Images were processed using Fiji software.

Wing imaginal discs from 3[rd] instar larvae were dissected in PBS, fixed in 4%

paraformaldehyde, and permeabilized in PBT. For Wg staining, carcasses were blocked

for 1hr in 5% normal goat serum (S-1000, Vector Labs) at room temp, and incubated

with 1:50 mouse anti-wg (4D4, DSHB) primary antibody and 1:500 anti-mouse 488 (A-

21202, Molecular Probes) secondary antibody. Primary and secondary antibody

incubations were performed at 4°C overnight. All carcasses were stained with DAPI and

phalloidin-TRITC, and mounted on glass slides with vectashield (H-1000, Vector

Laboratories Inc.) under a coverslip. Images of mounted wing discs were acquired on a

Zeiss 780 confocal microscope.

Larvae, pupae, and adult flies were imaged using a Zeiss Axio Zoom V16

fluorescence microscope.

Quantification of mNeonGreen expressing S2R+ cells

For FACs-based cell counting, we collected cultures from each gene knock-in experiment before and after puromycin selection. Pre-selection cultures were obtained by collecting of 500ul of culture 3-4 days after transfection. Post-selection cultures were obtained after at least 2 weeks of puromycin incubation. Non-transfected cells were used as a negative control. 100,000 cells were counted for each sample and FlowJo software was used to analyze and graph the data. FSC-A vs GFP-A was plotted and we defined mNeonGreen+ cells by setting a signal intensity threshold where <0.02% of negative controls are counted due to autofluorescence.

For microscopy-based cell counting, the number of mNeonGreen cells was quantified by analyzing confocal images in Fiji using the manual Cell Counter Plugin (model). For transfected cells, 6 fields containing at least 200 cells were quantified (i.e. n=6). For puro-selected cells, 3 fields containing at least 200 cells were quantified (i.e. n=3).

Western blotting

Single cell-cloned cell lines were grown until confluent and 1ml of resuspended cells was centrifuged at 250g for 10min. The cell pellet was resuspended in 1ml ice cold PBS, re-centrifuged, and the pellet was lysed in 250ul 2x SDS-Sample buffer and boiled for 5min. 10ul was loaded on a 4-20% Mini-Protean TGX SDS-Page gel (4561096, BioRad), transferred to PVDF membrane (IPFL00010, Millipore), blocked in 5% non-fat dry milk, primary blotting using anti-mNeonGreen (1:1000, Chromtek 32F6) or hFAB™ Rhodamine Anti-Actin (12004164 BioRad), and secondary blotting using 1:3000 anti-

mouse HRP (NXA931, Amersham), imaging using ECL (34580, ThermoFisher) on a

ChemiDoc MP Imaging System (BioRad)**.**

PCR, sequencing, and sgRNA cutting assays

S2R+ cell genomic DNA was isolated using QuickExtract (QE09050, Lucigen).

Fly genomic DNA was isolated by grinding a single fly in 50µl squishing buffer (10 mM

Tris-Cl pH 8.2, 1 mM EDTA, 25 mM NaCl) with 200µg/ml Proteinase K (3115879001,

Roche), incubating at 37˚C for 30 min, and 95˚C for 2 minutes. PCR was performed

using Taq polymerase (TAKR001C, ClonTech) when running DNA fragments on a gel,

and Phusion polymerase (M-0530, NEB) was used when DNA fragments were

sequenced. DNA fragments corresponding to *mNeonGreen* or *T2A-Gal4* insertion sites

were amplified using primer pairs where one primer binds to genomic sequence and the

other primer binds to the insert. For amplifying non-knock-in sites, we used primers that

flank the sgRNA target site. Primer pairs used for gel analysis and/or Sanger

sequencing were designed to produce DNA fragments <1kb. Primer pairs used for next-

generation sequencing of the insertion site were designed to produce DNA fragments

200-280bp. DNA fragments were run on a 1% agarose gel for imaging or purified on

QIAquick columns (28115, Qiagen) for sequencing analysis. See Supplemental Table 6

for oligo sequences.

Sanger sequencing was performed at the DF/HCC DNA Resource Core facility

and chromatograms were analyzed using Lasergene 13 software. Next-generation

sequencing was performed at the MGH CCIB DNA Core. Fastq files were analyzed

using CRISPresso2 (CLEMENT *et al.* 2019) by entering the PCR fragment sequence into

the exon specification window and setting the window size to 10 bases. Quantification of

insertion types (seamless, in-frame in/del, and frameshift in/del) was taken from the

allele plot and frame shift analysis outputs of CRISPresso2. The small proportion of

"unmodified" reads that were not called by frameshift analysis were not included in the

quantification.

T7 endonuclease assays (M0302L, NEB) were performed following the

manufacturer instructions.

Data availability:

Donor plasmids and frame selector sgRNA plasmids will be deposited at Addgene. Fly

strains, S2R+ cell lines, and sequence data are available on request. Oligo and dsDNA

sequences are listed in Supplemental Table 6.

Results

To test homology-independent knock-in in *Drosophila*, we implemented a

strategy known as CRISPaint (SCHMID-BURGK *et al.* 2016). This system is used to insert

a protein tag or reporter gene into the coding sequence of an endogenous gene.

Although it was originally designed for mammalian cell culture, CRISPaint has several

advantages for use in *Drosophila*. First, this system uses CRISPR-Cas9 to induce

double-strand breaks (DSBs), which is known to function efficiently in *Drosophila*

cultured cells (BOTTCHER *et al.* 2014; VISWANATHA *et al.* 2018) and the germ line (KONDO

AND UEDA 2013; REN *et al.* 2013; YU *et al.* 2013; BASSETT *et al.* 2014). Second, its use of

a frame-selector gRNA target site makes insertion into the appropriate translation frame

simple and modular (see below). Third, a collection of existing CRISPaint donor

plasmids (SCHMID-BURGK et al. 2016) containing common tags (e.g. GFP, RFP,

Luciferase) are seemingly compatible for expression in *Drosophila*.

The CRISPaint system works by introducing three components into Cas9-

expressing cells: 1) a single guide RNA (sgRNA) targeting a genomic locus; 2) a donor

plasmid containing an insert sequence; and 3) a frame selector sgRNA targeting the

donor plasmid (Figure 1A). This causes simultaneous cleavage of the genomic locus

and donor plasmid, leading to the integration of linearized donor into the genomic cut

site by non-homologous end joining (NHEJ). To ensure that the donor plasmid inserts

in-frame with the endogenous gene, one of three frame selector sgRNAs are used.

Importantly, these frame-selector sgRNAs do not target the *Drosophila* genome.

Homology-independent insertion functions efficiently in *Drosophila* S2R+ cells to

produce endogenous protein tags

To test if the CRISPaint method functions in *Drosophila*, we set out to replicate

the findings of (SCHMID-BURGK et al. 2016) in cultured S2R+ cells by genetically tagging

endogenous proteins at their C-terminus. To accomplish this, we generated plasmids

expressing frame-selector sgRNAs (frame 0,1, or 2) under the control of *Drosophila* U6

sequences (PORT et al. 2015) (Figure 1A). In addition, we generated plasmids

expressing sgRNAs that target the 3' coding sequence of endogenous *Drosophila*

genes. We chose to target *Actin5c*, *His2Av*, *alphaTub84B*, and *Lamin* because these

genes are expressed in S2R+ cells (HU et al. 2017) and encode proteins with known

subcellular localization (actin filaments, chromatin, microtubules, nuclear envelope,

respectively). For donor plasmid, we used *pCRISPaint-mNeonGreen-T2A-PuroR*

(SCHMID-BURGK et al. 2016), which contains a frame-selector sgRNA target site

upstream of coding sequence for the fluorescent mNeonGreen protein and Puromycin

resistance protein (PuroR) linked by a cleavable T2A peptide sequence. Importantly,

only integration of the donor plasmid in-frame with the target gene coding sequence will

result in translation of mNeonGreen-T2A-PuroR (Figure 1A).

We transfected Cas9-expressing S2R+ cells (VISWANATHA *et al.* 2018) with a mix

of three plasmids: *pCRISPaint-mNeonGreen-T2A-PuroR* donor, target-gene sgRNA,

and the appropriate frame-selector sgRNA (Figure 1A, Supplemental Table 1). As an

initial method to detect knock-in events, we used PCR to amplify the predicted insertion

sites from transfected cells. Using primers that are specific to the target gene and

mNeonGreen sequence, we successfully amplified *gene-mNeonGreen* DNA fragments

for all four genes (Figure 1B). Furthermore, next-generation sequencing of these

amplified fragments revealed that 34-50% of sense-orientation insertions are in frame

with the target gene (Figure 1B, Supplemental Figure 1, Supplemental Table 1).

Next, we measured mNeonGreen fluorescence in transfected S2R+ cells as a

more direct method of quantifying the frequency of in-frame knock-ins. Flow cytometry-

based cell counting of transfected cells revealed that the number of mNeonGreen+ cells

range from 0.19-2.4% (Figure 1C, Supplemental Table 1), in agreement with published

results in human cultured cells (SCHMID-BURGK *et al.* 2016). These results were

confirmed by confocal analysis of transfected cells, which showed mNeonGreen

fluorescence in a small subset of cells (Figure 1C). Analysis of confocal images of *Act5c*

and *His2Av* samples showed that 3.2% and 2.4% of transfected cells expressed

mNeonGreen (Figure 1C, Supplemental Table 1), which roughly agreed with flow

cytometry cell counting. Finally, mNeonGreen localized to the expected subcellular

compartments, most obviously observed by His2Av-mNeonGreen and Lam-

mNeonGreen co-localization with the nucleus, and Act5c-mNeonGreen and alphatub-

mNeonGreen exclusion from the nucleus (Figure 1C). These results suggest that a

significant number of transfected S2R+ cells received in-frame insertion of mNeonGreen

at their C-terminus using the CRISPaint homology-independent insertion method.

For knock-in cells to be useful in experiments, it is important to derive cultures

where most cells, if not all, carry the insertion. Therefore, we enriched for in-frame

insertion events using Puromycin selection (Figure 1D). After a two-week incubation of

transfected S2R+ cells with Puromycin, flow-cytometry and confocal analysis revealed

that most cells express mNeonGreen and exhibit correct subcellular localization (Figure

1E, Supplemental Table 1). For *alphaTub84B*, cell counting by flow-cytometry greatly

underestimated the number of mNeonGreen+ cells counted by confocal analysis, likely

because the mNeonGreen expression level was so low. Therefore, Puromycin selection

is a fast and efficient method of selecting for mNeonGreen expressing knock-in cells

after transfection.

A subset of cells in Puro-selected cultures had no mNeonGreen expression or

unexpected localization (Figure 1E). Since each culture is composed of different cells

with independent insertion events, we used FACs to derive single-cell cloned lines

expressing mNeonGreen for further characterization (Figure 2A). At least 14 single-cell

cloned lines were isolated for each target gene and imaged by confocal microscopy.

Within a given clonal culture, every cell exhibited the same mNeonGreen localization

(Figure 2B), confirming our single-cell cloning approach and demonstrating that the

insertion is genetically stable over many cell divisions. Importantly, while many clones

exhibited the predicted mNeonGreen localization, a subset of the clonal cell lines

displayed an unusual localization pattern (Figure 2B). For example, three *Act5c-*

mNeonGreen clones had localization in prominent rod structures, and 12 *Lamin-*

mNeonGreen clones had asymmetric localization in the nuclear envelope (Figure 2B).

In addition, some clones had diffuse mNeonGreen localization in the cytoplasm and

nucleus (Figure 2B).

To better characterize the insertions in single cell-cloned lines, we further

analyzed three clones per gene (12 total), selecting different classes when possible

(correct localization, unusual localization, and diffuse localization) (Supplemental Table

2). Using PCR amplification of the predicted insertion site (Figure 1A, Figure 2C) and

sequencing of amplified fragments (Supplemental Table 2), we determined that all

clones with correct or unusual mNeonGreen localization contained an in-frame insertion

of mNeonGreen with the target gene. In contrast, we were unable to amplify DNA

fragments from the expected insertion site in clones with diffuse mNeonGreen

localization (Figure 2C). Western blotting of cell lysates confirmed that only clones with

in-frame mNeonGreen insertion express fusion proteins that match the predicted

molecular weights (Figure 2D). All together, these results suggest that clones with

correct mNeonGreen localization are likely to contain an in-frame insert in the correct

target gene.

Since S2R+ cells are polyploid (LEE *et al.* 2014), clones expressing mNeonGreen

could bear one or more insertions. Furthermore, in/dels induced at the non-insertion

locus could disrupt protein function. To explore these possibilities, we amplified the non-

insertion locus in our single-cell cloned lines and used Sanger and next-generation

sequencing to analyze the DNA fragments (Figure 2C, Supplemental Table 2). For each

gene, we could find in/dels occurring at the non-insertion sgRNA cut site. For example,

we could distinguish four distinct alleles in clone B11: a 3bp deletion, a 2bp deletion, a

1bp deletion, and a 27bp deletion. In addition, we identified an unusual mutation in

clone C6, where a 1482bp DNA fragment inserted at the sgRNA cut site, which

corresponds to a region from *alphatub84D*. We assume that this large insertion was

caused by homologous recombination, since *alphatub84D* and *alphatub84B* share 92%

genomic sequence identity (Flybase). For *Act5c-mNeonGreen* clones A5 and A19,

numerous in/del sequences were found, suggesting this region has an abnormal

number of gene copies. We were unable to amplify a DNA fragment from *Lam-*

mNeonGreen D9, despite follow-up PCRs using primers that bind genomic sequence

further away from the insertion site (not shown).

One useful application of cell lines with fluorescently tagged endogenous

proteins is to track their localization over time. Therefore, we used live confocal imaging

of our single-cell cloned lines to capture mNeonGreen localization during cell division

(Figure 2E, Supplemental Videos). Time-lapse images of dividing cells showed that

Act5c-mNeonGreen localized to rod structures that asymmetrically or symmetrically

distribute into daughter cells, His2Av-mNeonGreen localized to chromosomes that

segregate into daughter cells, Lam-mNeonGreen showed disassembly and reassembly

at the nuclear envelope, and alphaTub84B-mNeonGreen localized to mitotic spindles.

These results demonstrate the usefulness of knock-in *Drosophila* cell lines to track the

dynamic localization of endogenous proteins.

***In vivo* germ line knock-in of T2A-Gal4 into endogenous genes using homology-**

independent insertion

We next tested if homology-independent insertion could function in the

Drosophila germ line for the purpose of generating knock-in fly strains. Compared to

cultured cells, the isolation of flies bearing insertions that are in-frame with endogenous

genes required additional considerations. As opposed to antibiotic selection, visible

markers are commonly used to identify transgenic animals. In addition, since some

genes are expressed at low levels, target gene expression of an inserted reporter

element may be insufficient to identify in-frame insertion events.

To overcome these issues, we constructed the donor plasmid *pCRISPaint-T2A-*

Gal4-3xP3-RFP (Figure 3A). This donor contains a frame-selector sgRNA target site

upstream of the reporter gene *T2A-Gal4*, which encodes a form of the transcription

factor Gal4 that is cleaved from tagged endogenous protein (DIAO AND WHITE 2012).

Insertion of this element in-frame with genomic coding sequence would result in Gal4

translation, which can be detected using a *UAS-reporter* transgene (BRAND AND

PERRIMON 1993). In addition, this donor plasmid contains a *3xP3-RFP* selectable marker

gene that expresses bright red fluorescence in *Drosophila* larval tissues and the adult

eye (BERGHAMMER *et al.* 1999; GRATZ *et al.* 2014) (Figure 3B). Importantly, the

expression of *3xP3-RFP* is not dependent on in-frame insertion with the target gene.

Next, we tested for *pCRISPaint-T2A-Gal4-3xP3-RFP* insertion into 11

endogenous genes (Supplemental Table 3). These genes were selected based on their

known expression pattern, expression levels, or loss of function phenotype.

Furthermore, we targeted *pCRISPaint-T2A-Gal4-3xP3-RFP* to insert into the 5' portion

of the coding sequence (Figure 3A). This insertion location is designed to disrupt the

protein product by premature truncation. Plasmid mixes were injected into *nos-Cas9*

embryos, the resulting G0 progeny were outcrossed to *yw*, and G1 adults were

screened for RFP fluorescence (Figure 3C). Each RFP+ founder fly was outcrossed to

an appropriate balancer stock to establish a stable line. Figure 3D and Supplemental

Table 3 shows the integration efficiency results for each gene and Supplemental Table

4 has information on each balanced RFP+ line. From this data, we find that the

frequency of G0 crosses yielding RFP+ G1 progeny varies between 5% and 21%

(Figure 3D, Supplemental Table 3). For example, when targeting *ebony* with *pFP545*, 3

out of 16 G0 crosses produced ≥1 RFP+ G1 flies. Therefore, the *pCRISPaint-T2A-Gal4-*

3xP3-RFP donor can insert into the genome of germ line cells in a homology-

independent manner.

477   To gain insight into the genomic location of the insertions in our RFP+ lines, we

first analyzed them by simple genetic crosses. During the fly stock balancing process,

we determined that each insertion was located on the intended chromosome

(Supplemental Table 4). In addition, flies that were homozygous for the insertion

exhibited known phenotypes. For example, homozygous insertions in *ebony* produced

flies with dark cuticle pigment (Figure 3E). Furthermore, flies with insertions targeting

wg, *Mhc*, *hh*, and *esg* were homozygous lethal, which is consistent with known loss of

function mutations in these genes (Supplemental Table 4). To test if the lethality of flies

with homozygous insertions was due to on- or off-target gene disruption, we performed

complementation tests by crossing RFP+ insertion lines with lines containing a known

loss of function allele or genomic deletion spanning the gene. In all cases tested, trans-

heterozygous combinations were lethal (Supplemental Table 4). Together, these results

suggest that the *pCRISPaint-T2A-Gal4-3xP3-RFP* donor plasmid inserted into the

intended target genes.

491   For T2A-Gal4 to be expressed by the target gene, the linearized donor plasmid

must insert in the sense orientation relative to the target gene and in-frame with the

coding sequence. As an initial screen for such events, we crossed RFP+ lines to a *UAS-*

GFP line and assayed progeny for fluorescence. Through this approach, we identified

Gal4-expressing lines for *ebony*, *myo1a*, *wg*, and *Mhc* (Figure 4A, Supplemental Tables

3, 4). *wg-T2A-Gal4* (#1 and 4), *Mhc-T2A-Gal4* (#1 and 2), and *Myo1a-T2A-Gal4* (#1)

insertions express in the imaginal disc, larval muscle, and larval gut (Figure 4A),

respectively, which matches the known expression patterns for these genes.

Furthermore, *wg-T2A-Gal4 #1 and #4* insertions were expressed in a distinctive Wg

pattern in the wing disc pouch (Figure 4B). The expression pattern of *ebony* is less well

understood. We find that *ebony-T2A-Gal4 pFP545 #2* is expressed in the larval brain

(Figure 4A) and throughout the pupal body (Figure 4C), which is consistent with a

previous study (HOVEMANN *et al.* 1998). However, *ebony-T2A-Gal4 pFP545 #2* is also

expressed in the larval tracheal openings (Figure 4A), indicating that *ebony* may play a

role in this tissue.

Next, we analyzed the insertion orientation and sequence structure in the RFP+

lines that express Gal4. We PCR amplified a region flanking the predicted insertion site

from genomic DNA using primer pairs to distinguish sense and anti-sense insertions

(Supplemental Figure 2). All RFP+ lines with Gal4 expression had insertions that were

in the sense orientation (Supplemental Table 4, Supplemental Figure 2). Sequencing

the resulting PCR fragments showed that the insert was present at the sgRNA cut site

and each insertion contained an in/del between the target gene and *T2A-Gal4*

sequence (Figure 4D). For example, *ebony-T2A-Gal4 pFP545 #2* contains a 15bp

genomic deletion that is predicted to keep T2A-Gal4 in-frame with ebony. Similarly, *wg-*

T2A-Gal4 #1 contains an in-frame 45bp deletion and 21bp insertion. Remarkably, *wg-*

T2A-Gal4 #4 contains a frameshift in/del (Figure 4D), yet still expresses Gal4 in the Wg

pattern, albeit at significantly lower levels than *wg-T2A-Gal4 #1* (Figure 4B). In similar

cases, *Mhc-T2A-Gal4* lines #1, #2, and *Myo1a-T2A-Gal4* #1, each have in/dels that put

T2A-Gal4 out of frame with the target gene coding sequence. These findings confirm

that our Gal4-expressing lines have *T2A-Gal4* inserted in the correct gene and

orientation, but that in-frame insertion with the target gene is not necessarily a

requirement.

To better characterize the insertion events in our collection of RFP+ lines, we

analyzed those that did not produce fluorescence when crossed with *UAS-GFP*

(Supplemental Table 4, Supplemental Figures 2,3). Using PCR and sequencing

analysis, we found that some lines contained insertions in the correct target site but in

the anti-sense orientation. In addition, we identified lines with insertions in the sense

orientation, but were out of frame relative to the target gene. Unexpectedly, we found

that *wg-T2A-Gal4* #6 contained a sense orientation in-frame insertion. Yet, unlike *wg-*

T2A-Gal4 #1, *wg-T2A-Gal4* #6 does not express Gal4. Importantly, our molecular

analysis of every independently isolated RFP+ line (20 in total) revealed that each

contained an insertion in the intended target site (Supplemental Table 4, Supplemental

Figures 2,3).

We did not obtain RFP+ insertions when targeting *ap*, *alphaTub84B*, *btl*, or

Desat1. Therefore, we investigated whether the sgRNAs targeting these genes were

functional. All 4 sgRNAs used for germ line knock-ins have an acceptable efficiency

score of >5, with the exception of the sgRNA targeting *btl* (Supplemental Table 5). We

tested whether the sgRNAs were functional in transfected S2R+ cells by performing a

T7 endonuclease assay that detects in/dels at the cut site. This test revealed that

sgRNAs targeting *ap*, *alphaTub84B*, *btl* can cut at the target site, whereas the results

with *desat1* were inconclusive (Supplemental Figure 4A). As an alternative functional

test, we used PCR to detect knock-in events in S2R+ cells transfected with the

pCRISPaint-T2A-Gal4-3xP3-RFP donor plasmid. This showed that sgRNAs targeting

ap, *alphaTub84B*, *btl* and *desat1* can successfully knock-in *pCRISPaint-T2A-Gal4-*

3xP3-RFP (Supplemental Figure 4B). Finally, we sequenced the sgRNA target sites in

the *nos-Cas9* fly strains and found a SNP in the *btl* sgRNA binding site (not shown). The

10 remaining sgRNAs had no SNPs in the target site. In summary, we conclude that the

sgRNAs targeting *ap*, *alphaTub84B*, *btl*, and *Desat1* are able to induce cleavage at their

target site in S2R+ cells, but that the sgRNA targeting *btl* will not function in the germ

line using our *nos-Cas9* strains.

A resource of CRISPaint donor plasmids for germ line knock-ins in Drosophila

554   To facilitate the insertion of other sequences using the CRISPaint insertion

method, we generated 10 additional donor plasmids based on the same architecture as

pCRISPaint-T2A-Gal4-3xP3-RFP (Figure 5A). These include T2A-containing donors

with sequence encoding the alternative binary reporters LexGAD, QF2, and split-Gal4,

as well as Cas9 nuclease, FLP recombinase, Gal80 repression protein, NanoLuc

luminescence reporter, and super-folder GFP. Like *T2A-Gal4*, these can be used to

insert at 5' coding sequence, capturing endogenous gene expression and generating a

loss-of function. In addition, we generated *pCRISPaint-sfGFP-3xP3-RFP*, which can be

used to insert into 3' coding sequence, generating a C-terminal GFP fusion protein.

563   Several groups have demonstrated that coding sequence containing a splice

acceptor (SA) and inserted in a gene intron can produce a protein trap with the

preceding coding exon (MORIN *et al.* 2001; VENKEN *et al.* 2011). Recently, two studies

produced *SA-T2A-Gal4* donor plasmids for intron insertion by HDR, called CRIMIC and

T-GEM (DIAO *et al.* 2015; LEE *et al.* 2018). Therefore, we modified these two plasmids to

contain a CRISPaint target site upstream of the splice acceptor (Figure 5B).

Discussion

The insertion of large DNA elements into the genome by HDR requires a great

deal of expertise and labor for the design and construction of donor plasmids. Some

groups have developed strategies to improve the efficiency and scale at which

homology arms are cloned into donor plasmids (HOUSDEN *et al.* 2014; GRATZ *et al.*

2015), but the root problem still remains. Furthermore, each new gene-specific donor

plasmid requires the same amount of investment for their construction but is only used

once to achieve the desired knock-in. For these reasons, we believe that the current

methods for knock-in by HDR may act as a barrier to achieving widespread use by the

Drosophila community.

In this study, we addressed these challenges by demonstrating that large DNA

elements can insert into the *Drosophila* genome by a homology-independent

mechanism, using the previously established CRISPaint system. This approach has two

major advantages over HDR. 1) No construction of a donor plasmid is necessary, as

long as a suitable CRISPaint-compatible donor plasmid already exists. The only unique

reagent needed is an sgRNA that targets the endogenous gene (also required for HDR).

Cloning sgRNAs into expression plasmids, such as *pCFD3* (PORT *et al.* 2014), is simple,

fast, inexpensive, and works nearly every time. Furthermore, the availability of sgRNA-

encoding plasmids from public resources (e.g. TRiP, Addgene), and synthesized

sgRNA from commercial companies, means that researchers can increasingly order

their sgRNAs. 2) CRISPaint-compatible donor plasmids are "universal" and thus

modular. For example, different genes can be targeted by the same CRISPaint donor

plasmid, and different CRISPaint donor plasmids can be targeted to the same gene.

Publicly available collections of CRISPaint donor plasmids [(SCHMID-BURGK *et al.* 2016),

this study] ensure that researchers only need to select their insert of choice. Indeed, the

CRISPaint donor plasmids originally used for mammalian cell culture also function in

Drosophila S2R+ cells (Figure 1) and the *3xP3-RFP* marker in our germ line donor

plasmids is compatible with other insects (BERGHAMMER *et al.* 1999).

An important step in obtaining correctly targeted knock-ins is molecularly

validating the candidate insertions. Confirming an HDR insertion requires amplifying a

large DNA fragment (~1.5kb-2kb) that encompasses part of the insert, an entire

homology arm, and a portion of genomic sequence flanking the homology arm. This is

necessary to verify that the donor did not insert off-target. These PCRs can sometimes

fail or give inconclusive results due to the large fragment size. In contrast, CRISPaint

knock-ins are easier to characterize by PCR analysis and sequencing because the

amplified region is relatively small (~200-800bp) (Figure 1B, Figure 2C, Supplemental

Figures 2,4). However, CRISPaint knock-ins require more work to screen since they can

insert in two directions and in/dels occur at the insertion site. When possible, we

recommend that researchers select for insert expression before molecular validation.

In this study, we generated knock-ins by inserting the entire linearized CRISPaint

donor plasmid into the target gene. Since the backbone contains bacterial sequences, it

may cause transgene silencing or impact neighboring gene expression (CHEN *et al.*

2004; SUZUKI *et al.* 2016). However, we note that thousands of transgenic fly lines

contain bacterial sequences from phiC31 integration (PERKINS *et al.* 2015) with no

reports of ill effects. Another issue is that insertion of the entire plasmid restricts the

design of gene-tagging events to only append the insert 3' to the target insertion site.

Different groups have used approaches that address these issues, such as providing

donor plasmids as mini-circles (SCHMID-BURGK *et al.* 2016; SUZUKI *et al.* 2016), cutting

donor plasmid twice to liberate the insert fragment (LACKNER *et al.* 2015; SUZUKI *et al.*

2016), or using PCR amplified inserts (Manna et al. 2019 BioRxiv). The first two

modifications could in theory be made to our germ-line donor plasmids (e.g.

pCRISPaint-T2A-Gal4-3xP3-RFP), but for this study we opted to establish the simplest

protocol possible. Furthermore, we reasoned that cutting the donor twice would give rise

to two donor fragments and this could reduce knock-in efficiency.

625   Using the CRISPaint method in S2R+ cells, we readily identified cell lines with

endogenous proteins tagged with mNeonGreen at their C-terminus (Figure 2,

Supplemental Table 2). However, some lines exhibited unusual or unexpected protein

localization. In clones D6 and D9, Lamin-mNeonGreen localizes to the nuclear

envelope, but in D9 this localization is enriched asymmetrically in the direction of the

previous plane of cell division. Since these two clones contain the same seamless

mNeonGreen insertion, we speculate that mutations at non-knock-in loci account for this

difference. Indeed, clone D6 contained an in-frame 3bp deletion at the non-knock-in

locus, likely retaining wild-type function, whereas D9 had no remaining non-knock-in

locus. We saw a similar pattern for clones A3 and A5, where both had seamless

mNeonGreen insertions in *Actin5c*, but clone A5 exhibited distinct rod structures.

Finally, alphaTub84B-mNeonGreen fluorescence and protein levels were extremely low

in all cell lines, despite *alphaTub84B* being highly expressed in S2R+ cells (HU *et al.*

2017). We speculate that the alphaTub84B-mNeonGreen fusion protein is unstable and

previous studies in other organisms have highlighted problems with C-terminal tagging

of alpha-Tubulin (CARMINATI AND STEARNS 1997). Similarly, C-terminal tags can disrupt

Lamin and Actin function (DAVIES *et al.* 2009; NAGASAKI *et al.* 2017). These findings

illustrate the need for experimenters to consider the existing knowledge of the protein

when generating C-terminal protein fusions, and to carefully screen individual single cell

cloned lines.

We constructed a CRISPaint-compatible *T2A-Gal4* donor plasmid for use in the

fly germ line and successfully identified insertion lines. Our knock-in efficiency, defined

by the percentage of injected G0 flies that give RFP+ progeny, ranged from 5-21%

(Figure 3B, Supplemental Table 3), which is roughly similar to knock-in efficiencies

observed when using HDR [5-22% (GRATZ *et al.* 2014), 46-88% (PORT *et al.* 2015), 7-

42% (GRATZ *et al.* 2015)]. In addition, all 20 of our RFP+ fly lines, which encompass 8

different sgRNA target sites, contain an insertion at the correct location. Though, we do

not rule out the possibility of a second-site off-target event on the same chromosome.

To obtain *T2A-Gal4* insertions that express Gal4 under the control of the target

gene (Figure 4), it was necessary to screen multiple independently derived insertions,

due to the in/dels that occur at the insertion site and the two insertion orientations.

However, we found for some genes the overall efficiencies were too low to obtain a

successful Gal4-expressing line (*hh, esg, FK506-bp2*), or we did not obtain any RFP+

insertions (*ap, alphaTub84B, btl,* and *Desat1*). Additional steps could be taken to

improve insertion efficiency, such as optimization of the injected plasmid concentrations,

increasing the number of injected embryos, or simply reattempting with a different

sgRNA. It is also possible that certain insertions are toxic to cells/animals during G0

germ-line development or in G1 progeny.

There were three unexpected findings with our germ-line insertions. First, some

Gal4-expressing lines had *T2A-Gal4* inserted out of frame relative to the target gene.

We speculate that it may be the result of ribosome frameshifting (KETTELER 2012), an

internal ribosome entry site (IRES) (KOMAR AND HATZOGLOU 2005), or the presence of

alternative open reading frames (altORFs) (MOUILLERON *et al.* 2016). However, we find

no obvious evidence of these mechanisms by analyzing the sequence flanking *wg, Mhc,*

and *Myo1a* insertion sites (not shown). Ultimately, we consider this a fortuitous effect as

long as Gal4 is expressed in the correct pattern. Second, we found an in-frame insertion

in *wg* (#6) that that does not express Gal4. This finding highlights the importance of

screening RFP+ insertions for Gal4 expression when possible. Third, we found that,

unlike in cell culture, all of our RFP+ fly lines contain in/dels at the insertion site. Germ

cells are known to differ in their NHEJ mechanisms compared to somatic cells (PRESTON

et al. 2006; AHMED *et al.* 2015), but it is not clear why this would reduce the frequency of

seamless insertions. Perhaps genetic or chemical manipulation of NHEJ regulators

during embryo injection could address this issue in the future. This finding also suggests

that the CRISPaint frame-selector approach may not be as useful in the fly germ line as

it is in cell culture.

Our collection of donor plasmids (Figure 5) provides many options for inserting

protein-coding sequence into target genes. However, other uses for homology-

independent knock-in can be imagined, such as inserting enhancer sequences (e.g.

UAS) upstream of endogenous genes to induce their overexpression (RORTH 1996), a

reporter gene near non-coding regulatory sequences to capture the transcriptional

expression pattern of neighboring genes (BRAND AND PERRIMON 1993), entire genes into

intergenic sequence (SADELAIN *et al.* 2011), or sequences to be used for labeling DNA

loci (ROBINETT *et al.* 1996). Furthermore, the donor plasmids described in this study

could be used to simply knock out endogenous genes with a selectable marker. Indeed,

all of our mNeonGreen-expressing single cell cloned lines contain mutations in the non-

knock-in locus and our fly germ line insertions produced loss of function phenotypes.

This approach could greatly increase the efficiency of selecting knock-out alleles, which

are traditionally done by laborious PCR-based screening of frameshift in/dels. We also

note that, similar to the *T2A-Gal4* reporters in vivo, cell lines could be targeted with

translational reporters such as NanoLuciferase or GFP. Finally, since our collection of

CRISPaint donor plasmids contain enzyme restriction sites that flank the insert

sequence, they are also useful as parental vectors for constructing traditional HDR

donor plasmids.

In summary, our homology-independent knock-in approach enables researchers

to focus more effort on screening for correct insertions in cells or flies than on designing

and constructing donor plasmids. Furthermore, the techniques required for screening

knock-ins are less specialized than those for constructing donor plasmids, making this

trade off potentially attractive for labs with less molecular biology expertise or resources.

Therefore, we hope that this method will put knock-in technology into the hands of more

researchers due to its simplicity.

Acknowledgements

We thank Jonathan Schmid-Burgk for advice and the *CRISPaint-mNeonGreen* donor

plasmid, Claire Hu and the TRiP for sgRNA design and construction, Ben Ewen-

Campen for valuable comments on the manuscript, Stephanie Mohr, Oguz Kanca, and

Hugo Bellen for helpful discussions, Raghuvir Viswanatha for the *Cas9-T2A-EGFP*

template sequence, and Rich Binari and Cathryn Murphy for general assistance. J.A.B.

was supported by the Damon Runyon Foundation. This work was supported by NIH

grants R01GM084947, R01GM067761, R24OD019847. N.P. is an investigator of the

Howard Hughes Medical Institute.

References:

Ahmed, E. A., H. Scherthan and D. G. de Rooij, 2015 DNA Double Strand Break
Response and Limited Repair Capacity in Mouse Elongated Spermatids. Int J
Mol Sci 16**:** 29923-29935.
Auer, T. O., K. Duroure, A. De Cian, J. P. Concordet and F. Del Bene, 2014 Highly
efficient CRISPR/Cas9-mediated knock-in in zebrafish by homology-independent
DNA repair. Genome Res 24**:** 142-153.
Bassett, A. R., C. Tibbit, C. P. Ponting and J. L. Liu, 2014 Mutagenesis and homologous
recombination in Drosophila cell lines using CRISPR/Cas9. Biol Open 3**:** 42-49.
Bellen, H. J., R. W. Levis, Y. He, J. W. Carlson, M. Evans-Holm *et al.*, 2011 The
Drosophila gene disruption project: progress using transposons with distinctive
site specificities. Genetics 188**:** 731-743.
Berghammer, A. J., M. Klingler and E. A. Wimmer, 1999 A universal marker for
transgenic insects. Nature 402**:** 370-371.
Bier, E., M. M. Harrison, K. M. O'Connor-Giles and J. Wildonger, 2018 Advances in
Engineering the Fly Genome with the CRISPR-Cas System. Genetics 208**:** 1-18.
Bosch, J. A., N. H. Tran and I. K. Hariharan, 2015 CoinFLP: a system for efficient
mosaic screening and for visualizing clonal boundaries in Drosophila.
Development 142**:** 597-606.
Bottcher, R., M. Hollmann, K. Merk, V. Nitschko, C. Obermaier *et al.*, 2014 Efficient
chromosomal gene modification with CRISPR/cas9 and PCR-based homologous
recombination donors in cultured Drosophila cells. Nucleic Acids Res 42**:** e89.
Bouabe, H., and K. Okkenhaug, 2013 Gene targeting in mice: a review. Methods Mol
Biol 1064**:** 315-336.
Brand, A. H., and N. Perrimon, 1993 Targeted gene expression as a means of altering
cell fates and generating dominant phenotypes. Development 118**:** 401-415.
Carminati, J. L., and T. Stearns, 1997 Microtubules orient the mitotic spindle in yeast
through dynein-dependent interactions with the cell cortex. J Cell Biol 138**:** 629-
641.
Chen, Z. Y., C. Y. He, L. Meuse and M. A. Kay, 2004 Silencing of episomal transgene
expression by plasmid bacterial DNA elements in vivo. Gene Ther 11**:** 856-864.
Clement, K., H. Rees, M. C. Canver, J. M. Gehrke, R. Farouni *et al.*, 2019
CRISPResso2 provides accurate and rapid genome editing sequence analysis.
Nat Biotechnol 37**:** 224-226.
Cristea, S., Y. Freyvert, Y. Santiago, M. C. Holmes, F. D. Urnov *et al.*, 2013 In vivo
cleavage of transgene donors promotes nuclease-mediated targeted integration.
Biotechnol Bioeng 110**:** 871-880.
Crivat, G., and J. W. Taraska, 2012 Imaging proteins inside cells with fluorescent tags.
Trends Biotechnol 30**:** 8-16.
Davies, B. S., L. G. Fong, S. H. Yang, C. Coffinier and S. G. Young, 2009 The
posttranslational processing of prelamin A and disease. Annu Rev Genomics
Hum Genet 10**:** 153-174.
Diao, F., H. Ironfield, H. Luan, F. Diao, W. C. Shropshire *et al.*, 2015 Plug-and-play
genetic access to drosophila cell types using exchangeable exon cassettes. Cell
Rep 10**:** 1410-1421.
Diao, F., and B. H. White, 2012 A novel approach for directing transgene expression in
Drosophila: T2A-Gal4 in-frame fusion. Genetics 190**:** 1139-1144.
Gratz, S. J., C. D. Rubinstein, M. M. Harrison, J. Wildonger and K. M. O'Connor-Giles,
2015 CRISPR-Cas9 Genome Editing in Drosophila. Curr Protoc Mol Biol 111**:** 31
32 31-20.

Gratz, S. J., F. P. Ukken, C. D. Rubinstein, G. Thiede, L. K. Donohue *et al.*, 2014 Highly
specific and efficient CRISPR/Cas9-catalyzed homology-directed repair in
Drosophila. Genetics 196**:** 961-971.
Hasse, S., A. A. Hyman and M. Sarov, 2016 TransgeneOmics--A transgenic platform for
protein localization based function exploration. Methods 96**:** 69-74.
Housden, B. E., S. Lin and N. Perrimon, 2014 Cas9-based genome editing in
Drosophila. Methods Enzymol 546**:** 415-439.
Housden, B. E., M. Muhar, M. Gemberling, C. A. Gersbach, D. Y. Stainier *et al.*, 2017
Loss-of-function genetic tools for animal models: cross-species and cross-
platform differences. Nat Rev Genet 18**:** 24-40.
Hovemann, B. T., R. P. Ryseck, U. Walldorf, K. F. Stortkuhl, I. D. Dietzel *et al.*, 1998
The Drosophila ebony gene is closely related to microbial peptide synthetases
and shows specific cuticle and nervous system expression. Gene 221**:** 1-9.
Hu, Y., A. Comjean, N. Perrimon and S. E. Mohr, 2017 The Drosophila Gene
Expression Tool (DGET) for expression analyses. BMC Bioinformatics 18**:** 98.
Ivics, Z., M. A. Li, L. Mates, J. D. Boeke, A. Nagy *et al.*, 2009 Transposon-mediated
genome manipulation in vertebrates. Nat Methods 6**:** 415-422.
Kanca, O., H. J. Bellen and F. Schnorrer, 2017 Gene Tagging Strategies To Assess
Protein Expression, Localization, and Function in Drosophila. Genetics 207**:** 389-
412.
Katic, I., L. Xu and R. Ciosk, 2015 CRISPR/Cas9 Genome Editing in Caenorhabditis
elegans: Evaluation of Templates for Homology-Mediated Repair and Knock-Ins
by Homology-Independent DNA Repair. G3 (Bethesda) 5**:** 1649-1656.
Katoh, Y., S. Michisaka, S. Nozaki, T. Funabashi, T. Hirano *et al.*, 2017 Practical
method for targeted disruption of cilia-related genes by using CRISPR/Cas9-
mediated, homology-independent knock-in system. Mol Biol Cell 28**:** 898-906.
Ketteler, R., 2012 On programmed ribosomal frameshifting: the alternative proteomes.
Front Genet 3**:** 242.
Kimple, M. E., A. L. Brill and R. L. Pasker, 2013 Overview of affinity tags for protein
purification. Curr Protoc Protein Sci 73**:** Unit 9 9.
Komar, A. A., and M. Hatzoglou, 2005 Internal ribosome entry sites in cellular mRNAs:
mystery of their existence. J Biol Chem 280**:** 23425-23428.
Kondo, S., and R. Ueda, 2013 Highly improved gene targeting by germline-specific
Cas9 expression in Drosophila. Genetics 195**:** 715-721.
Korona, D., S. A. Koestler and S. Russell, 2017 Engineering the Drosophila Genome for
Developmental Biology. J Dev Biol 5.
Lackner, D. H., A. Carre, P. M. Guzzardo, C. Banning, R. Mangena *et al.*, 2015 A
generic strategy for CRISPR-Cas9-mediated gene tagging. Nat Commun 6**:**
10237.
Lee, H., C. J. McManus, D. Y. Cho, M. Eaton, F. Renda *et al.*, 2014 DNA copy number
evolution in Drosophila cell lines. Genome Biol 15**:** R70.
Lee, P. T., J. Zirin, O. Kanca, W. W. Lin, K. L. Schulze *et al.*, 2018 A gene-specific T2A-
GAL4 library for Drosophila. Elife 7.
Maresca, M., V. G. Lin, N. Guo and Y. Yang, 2013 Obligate ligation-gated
recombination (ObLiGaRe): custom-designed nuclease-mediated targeted
integration through nonhomologous end joining. Genome Res 23**:** 539-546.
Morin, X., R. Daneman, M. Zavortink and W. Chia, 2001 A protein trap strategy to
detect GFP-tagged proteins expressed from their endogenous loci in Drosophila.
Proc Natl Acad Sci U S A 98**:** 15050-15055.

Mouilleron, H., V. Delcourt and X. Roucou, 2016 Death of a dogma: eukaryotic mRNAs
can code for more than one protein. Nucleic Acids Res 44**:** 14-23.
Nagasaki, A., T. K. S, T. Yumoto, M. Imaizumi, A. Yamagishi *et al.*, 2017 The Position
of the GFP Tag on Actin Affects the Filament Formation in Mammalian Cells. Cell
Struct Funct 42**:** 131-140.
Perkins, L. A., L. Holderbaum, R. Tao, Y. Hu, R. Sopko *et al.*, 2015 The Transgenic
RNAi Project at Harvard Medical School: Resources and Validation. Genetics
201**:** 843-852.
Port, F., H. M. Chen, T. Lee and S. L. Bullock, 2014 Optimized CRISPR/Cas tools for
efficient germline and somatic genome engineering in Drosophila. Proc Natl Acad
Sci U S A 111**:** E2967-2976.
Port, F., N. Muschalik and S. L. Bullock, 2015 Systematic Evaluation of Drosophila
CRISPR Tools Reveals Safe and Robust Alternatives to Autonomous Gene
Drives in Basic Research. G3 (Bethesda) 5**:** 1493-1502.
Preston, C. R., C. C. Flores and W. R. Engels, 2006 Differential usage of alternative
pathways of double-strand break repair in Drosophila. Genetics 172**:** 1055-1068.
Quadros, R. M., H. Miura, D. W. Harms, H. Akatsuka, T. Sato *et al.*, 2017 Easi-CRISPR:
a robust method for one-step generation of mice carrying conditional and
insertion alleles using long ssDNA donors and CRISPR ribonucleoproteins.
Genome Biol 18**:** 92.
Ren, X., J. Sun, B. E. Housden, Y. Hu, C. Roesel *et al.*, 2013 Optimized gene editing
technology for Drosophila melanogaster using germ line-specific Cas9. Proc Natl
Acad Sci U S A 110**:** 19012-19017.
Robinett, C. C., A. Straight, G. Li, C. Willhelm, G. Sudlow *et al.*, 1996 In vivo localization
of DNA sequences and visualization of large-scale chromatin organization using
lac operator/repressor recognition. J Cell Biol 135**:** 1685-1700.
Rorth, P., 1996 A modular misexpression screen in Drosophila detecting tissue-specific
phenotypes. Proc Natl Acad Sci U S A 93**:** 12418-12422.
Sadelain, M., E. P. Papapetrou and F. D. Bushman, 2011 Safe harbours for the
integration of new DNA in the human genome. Nat Rev Cancer 12: 51-58.
Schmid-Burgk, J. L., K. Honing, T. S. Ebert and V. Hornung, 2016 CRISPaint allows
modular base-specific gene tagging using a ligase-4-dependent mechanism. Nat
Commun 7**:** 12338.
Suzuki, K., Y. Tsunekawa, R. Hernandez-Benitez, J. Wu, J. Zhu *et al.*, 2016 In vivo
genome editing via CRISPR/Cas9 mediated homology-independent targeted
integration. Nature 540**:** 144-149.
Venken, K. J., A. Sarrion-Perdigones, P. J. Vandeventer, N. S. Abel, A. E. Christiansen
et al., 2016 Genome engineering: Drosophila melanogaster and beyond. Wiley
Interdiscip Rev Dev Biol 5**:** 233-267.
Venken, K. J., K. L. Schulze, N. A. Haelterman, H. Pan, Y. He *et al.*, 2011 MiMIC: a
highly versatile transposon insertion resource for engineering Drosophila
melanogaster genes. Nat Methods 8**:** 737-743.
Viswanatha, R., Z. Li, Y. Hu and N. Perrimon, 2018 Pooled genome-wide CRISPR
screening for basal and context-specific fitness gene essentiality in Drosophila
cells. Elife 7.
Yu, Z., M. Ren, Z. Wang, B. Zhang, Y. S. Rong *et al.*, 2013 Highly efficient genome
modifications mediated by CRISPR/Cas9 in Drosophila. Genetics 195**:** 289-291.
864

Figure 1

Figure 1 – Knock-in of *mNeonGreen-T2A-PuroR* into Drosophila S2R+ cells using homology-independent insertion. (**A**) Schematic of CRISPaint knock-in approach. *mNeonGreen-T2A-PuroR* is inserted into 3' coding sequence. (**B**) Analysis of knock-in efficiency of transfected cells by diagnostic PCR (DNA gel image) and next-generation sequencing (pie charts). (**C**) Analysis of knock-in efficiency of transfected cells by FACs and confocal microscopy. Numbers indicate percentage of cells with fluorescence. F-actin stained using Phalloidin-TRITC (red), nuclei labeled with DAPI (blue), mNeonGreen signal is in green. Scale bar 10µm. (**D**) Schematic of Puromycin selection of mNeonGreen-expressing cells. (**E**) Analysis of knock-in frequency of puromycin-selected cells using FACs and confocal microscopy. Numbers indicate percentage of cells with green fluorescence. Scale bar 10µm.

Figure 2. Analysis of S2R+ mNeonGreen-expressing single-cell cloned lines. (**A**) Schematic of FACS isolation of single-cell clones expressing mNeonGreen. (**B**) Confocal images of live mNeonGreen-expressing cell lines, categorized into three clone types. Numbers indicate the frequency of each clone type for each gene targeted. Images show fluorescence from Clic-mCherry (red) and mNeonGreen (green). Scale bar 25μm. (**C**) Agarose gel with PCR fragments amplified from knock-in (Gene_F/mNeonGreen_R) and non-knock-in loci (Gene_F/R). Positive control bands were amplified from *Rp49* genomic sequence. (**D**) Western blot detecting mNeonGreen protein fusions. Arrowheads indicate expected molecular weight. X's indicate incorrect molecular weight. (**E**) Confocal images of live S2R+ cells expressing mNeonGreen protein fusions during cell division at three timepoints. Arrowheads indicate cells before/after cell division.

Figure 3

A.

CRISPaint target site

Donor plasmid

pCRISPaint-T2A-Gal4-3xP3-RFP

Target gene

Knock-in (sense)

Knock-in (anti-sense)

B.

C.

nos-Cas9 Fertilized egg

Injection Mix:
1. *pCRISPaint-T2A-Gal4-3xP3RFP*
2. frame selector sgRNA
3. target gene sgRNA

Injected Adult   *yw*

G₀   x

G₁   RFP+

Establish line
Cross with *UAS-GFP*
Sequence insertion

D.

E.

ebony+   *ebony-*   *ebony FP545 #1*

Figure 3 – Knock-in of *T2A-Gal4* into the *Drosophila* germ line using homology-independent insertion. (**A**) Schematic of knock-in approach. *pCRISPaint-T2A-Gal4-3xP3-RFP* is inserted into 5' coding sequence. (**B**) Images of adult flies with 3xP3-RFP fluorescence in the eye. Top panel is brightfield, bottom panel is fluorescence. (**C**) Schematic of plasmid injections, fly crosses, and analysis of insertions. (**D**) Graph with results of knock-in efficiency for 12 sgRNA target sites and 11 genes. (**E**) Image of adult flies. Homozygous *ebony-T2A-Gal4 FP545* #1 flies have dark cuticle pigment.

Figure 4

Figure 4 – Germ line insertions that express *T2A-Gal4* under the control of the target gene. (**A**) Fluorescence images of 3rd instar larvae with indicated genotypes. Expression of Gal4 under control of the target gene drives expression of the *UAS-GFP* reporter. (**B**) Confocal images of wing imaginal discs showing protein staining of Wg protein (anti-wg, green) or *UAS-GFP* expression (green). GFP fluorescence was recorded at identical exposure settings for lines *wg-T2A-Gal4* #1 and #4. Inset shows digitally increased GFP signal. Scale bar 50μm. (**C**) Fluorescence image of pupae, *ebony-T2A-Gal4 pFP545* #2 expression is visible throughout the cuticle. (**D**) Sequence structure of *T2A-Gal4* insertions that express Gal4.

Figure 5

A

Figure 5 – A collection of CRISPaint-compatible donor plasmids for germ line knock-ins. (**A**) Donor plasmids for insertion into 5' coding sequence. (**B**) Donor plasmids for insertion into intronic sequence, modified from CRIMIC and T-GEM HDR donor plasmids.

Supplemental Figure 1

Supplemental Figure 1 – Next-generation sequencing analysis of mNeonGreen insertion sites in transfected S2R+ cells using CRISPresso2

Supplemental Figure 2 – Diagnostic PCR of *CRISPaint-T2A-Gal4* insertions in RFP+ fly lines to confirm their insertion site and orientation.

Supplemental Figure 3

Supplemental Figure 3 – Sequence structure of *CRISPaint-T2A-Gal4* insertions in RFP+ fly lines

Supplemental Figure 4

A

B

Supplemental Figure 4 – Cutting efficiency of 12 sgRNAs in transfected S2R+ cells. (**A**) T7 endonuclease assay. (**B**) Diagnostic PCR to detect for presence of sense orientation *CRISPaint-T2A-Gal4* insertion events.

	Gene and sgRNA information			
Gene Name	**Protein localization**	**sgRNA plasmid (TRiP)**	**sgRNA target site with PAM**	**frame selector sgRNA**
Act5c	actin	GP07595	GCGGTGCACAATGGAGGGGCCGG	2
His2Av	chromatin	GP07596	GGTGCAGGATCCGCAGCGGAAGG	2
αTub84B	microtubules	GP07609	CGGCGAGGGTGAGGGCGCTGAGG	2
Lamin	nuclear envelope	GP07612	CGCAGCGTGACCGCCGTGGACGG	1
Untransfected				

	CRISPresso2 output			
Gene Name	**seamless**	**in-frame in/del**	**frameshift in/del**	**% in-frame**
Act5c	5543	1343	13068	34.50937155
His2Av	2384	2620	8279	37.6722126
αTub84B	726	3034	6901	35.26873652
Lamin	971	801	1757	50.21252479
Untransfected				

	FACs Transfected			FACs Puro-selected		
Gene Name	**mNeonGreen+**	**Single cells total**	**% mNeonGreen+**	**mNeonGreen+**	**Single cells total**	**% mNeonGreen**
Act5c	1963	80589	2.435816302	45355	50883	89.1358607
His2Av	1554	80291	1.935459765	32780	46298	70.80219448
αTub84B	150	78752	0.190471353	16682	52643	31.6889235
Lamin	385	81688	0.471305455	34826	49234	70.73567047
Untransfected	2	83869	0.002384671	10	55656	0.017967515

	Cell counting transfected (n=6)			Cell counting Puro-selected (n=3)		
Gene Name	**mNeonGreen+**	**Total cells**	**Average % mNeonGreen+**	**mNeonGreen+**	**Total cells**	**Average % mNeonGreen+**
Act5c	17/17/26/18/22/25	571/679/577/599/574/613	3.5	342/280/318	397/398/377	80.3
His2Av	26/21/12/19/13/19	634/634/568/525/669/612	3	314/327/315	397/402/400	79.7
αTub84B				412/400/403	412/400/403	100
Lamin				270/282/259	382/388/374	70.9
Untransfected						

Supplemental Table 1 – Quantification of CRISPaint-mNeonGreen insertion events in transfected and puro-selected S2R+ cells by CRISPresso2, FACs, and cell counting of confocal images.

| Clone name | Gene-mNeonGreen | predicted mNeonGreen fusion (kDa) | Observed Localization | mNeon insertion site | endogenous gene | | | |
					allele 1	allele 2	allele 3	allele 4
A3	Act5c	71.4	very infrequent small rods	seamless	1bp deletion	1bp insertion		
A5	Act5c	71.4	very frequent long rods	seamless	numerous in/dels			
A19	Act5c	71.4	no rods, cyto and nuclear	N/A	numerous in/dels			
B1	His2Av	44.5	cyto and nuclear	N/A	4bp deletion	19bp deletion		
B11	His2Av	44.5	nuclear	seamless	3bp deletion	2bp deletion	1bp deletion	27bp deletion
B14	His2Av	44.5	nuclear	3 bp deletion, 1bp change	19bp deletion	3bp deletion	1bp deletion	
C2	alphatub84B	79.5	mitotic spindles	seamless	wt	6bp deletion	1bp deletion	
C6	alphatub84B	79.5	mitotic spindles	seamless	insertion of 1482bp from alphatub84D			
C13	alphatub84B	79.5	mitotic spindles	3bp deletion	wt	6bp insertion		
D1	Lam	100.9	cytoplasmic, nuclear	N/A	8bp deletion	4bp deletion	3bp insertion	
D6	Lam	100.9	nuclear lamina	seamless	3bp deletion			
D9	Lam	100.9	nuclear lamina, polarized	seamless	N/A			

Supplemental Table 2 – Molecular characterization of single cell cloned mNeonGreen-expressing S2R+ lines.

Gene	Chr.	target gene sgRNA	target site (no PAM)	Frame selector sgRNA	# G0 crosses	# G0 founders	% founders/total	# G1 progeny screened	# G1 RFP+ progeny	% RFP G1/total G1
Mhc	2	GP00583	CTGAAGCCCCGCATCAAGGT	1	29	6	21	2086	21	1.01
ebony	3	pFP545	TGGCCATCTGGAAGGCTGG	1	16	3	19	1298	11	0.85
FK506-bp2	2	GP02894	TGTAGTGGACCGTGACCTTT	1	13	2	15	901	2	0.22
wg	2	GP06461	GGGGCCGGGGCTCCATGTGG	0	40	6	15	4423	26	0.59
ebony	3	pFP573	TCTACACCTCGGGCAGTAC	1	23	3	13	2121	27	1.27
esg	2	GP00225	CTCCACCAACATGTCTTCCA	2	15	1	7	772	1	0.13
Myo1a	2	GP00364	ATGGCTATGCAACGGGAAGC	1	17	1	6	860	1	0.12
hh	3	GP01881	TAACCACAGCTCAGTGCCTT	2	39	2	5	1975	3	0.15
alphaTub84B	3	GP05054	GCTGACGGTAGGTTCCGGTA	1	6	0	0	506	0	0.00
btl	2	GP00400	GGCAAAAGTGCCGATCACGC	2	26	0	0	2085	0	0.00
Desat1	3	GP03252	AAGCTGCAGGAGGACTCCAC	1	7	0	0	386	0	0.00
ap	2	GP05302	CACCACCTGTAGCACATCAA	0	29	0	0	2928	0	0.00

Supplemental Table 3 – Germ line knock-in efficiency of *CRISPaint-T2A-Gal4*

#	Gene targeted	sgRNA	G0 cross #	Insert orientation	Insert frame	Insertion site sequence	Gal4 expression	Homozygous phenotype	Complementation test
1	ebony	pFP545	1	Sense	Out	8bp del	none	dark cuticle pigment	fail to complement e1 on TM3
2	ebony	pFP545	2	Sense	In	15bp del	anterior and posterior trachae in larvae, whole body cuticle in pupae and adult	dark cuticle pigment	fail to complement e1 on TM3
3	ebony	pFP573	1	Sense	Out	7bp del	none	dark cuticle pigment	fail to complement e1 on TM3
4	ebony	pFP573	2	Antisense	N/A	5bp del, 25bp ins	none	dark cuticle pigment	fail to complement e1 on TM3
5	ebony	pFP573	3	Sense	Out	8bp deletion	none	dark cuticle pigment	fail to complement e1 on TM3
6	hh	GP01881	1	Antisense	N/A	1897bp del, 8bp ins	none	lethal	fail to complement hh[AC], Df(3R)ED5296
7	Mhc	GP00583	1	Sense	Out	10bp del, 3bp ins	muscle	lethal	fail to complement P{lacW}Mhck10423, Df(2L)H20
8	Mhc	GP00583	2	Sense	Out	9bp del, 13bp ins	muscle	lethal	fail to complement P{lacW}Mhck10423, Df(2L)H20
9	Mhc	GP00583	3	Antisense	N/A	27bp del, 1bp ins	none	lethal	fail to complement P{lacW}Mhck10423, Df(2L)H20
10	Mhc	GP00583	4	Sense	Out	2bp del, 4bp ins	none	lethal	fail to complement P{lacW}Mhck10423, Df(2L)H20
11	Mhc	GP00583	5	Antisense	N/A	2bp deletion	none	lethal	fail to complement P{lacW}Mhck10423, Df(2L)H20
12	Myo1a	GP00364	1	Sense	Out	4bp del, 2bp ins	gut	viable	complement Df(2L)ED8142
13	esg	GP00225	1	Antisense	N/A	25bp del, 10bp	none	lethal	fail to complement P{enG}esg[G66]
14	FK506-bp2	GP02894	1	Antisense	N/A	30bp del	none	viable	complement Df(2R)Exel6069
15	wg	GP06461	1	Sense	In	45bp del, 21bp ins	wingless, strong expression	lethal	fail to complement wg[l-17], wg[l-8], and Df(2L)BSC291
16	wg	GP06461	2	Antisense	N/A	33bp del, 11bp ins	none	lethal	fail to complement wg[l-17], wg[l-8], and Df(2L)BSC291
17	wg	GP06461	3	Sense	Out	21bp del, 11bp ins	none	lethal	fail to complement wg[l-17], wg[l-8], and Df(2L)BSC291
18	wg	GP06461	4	Sense	Out	32bp del, 7bp ins	wingless, weak expression	lethal	fail to complement wg[l-17], wg[l-8], and Df(2L)BSC291
19	wg	GP06461	5	Sense	Out	28bp del, 5bp ins	none	in progress	fail to complement wg[l-17], wg[l-8], and Df(2L)BSC291
20	wg	GP06461	6	Sense	In	21bp insertion	none	in progress	in progress

Supplemental Table 4 – Molecular and phenotypic characterization of 20 RFP+ fly strains, each carrying a distinct *CRISPaint-T2A-Gal4* insertion.

gene	sgRNA name	sgRNA sequence	DRSC efficiency score
ebony	pFP545	gTGGCCATCTGGAAGGCTGG	5.56968
ebony	pFP573	gTCTACACCTCGGGCAGTAC	5.08877
wg	GP06461	GGGGCCGGGGCTCCATGTGG	5.61857
FK506-bp2	GP02894	TGTAGTGGACCGTGACCTTT	6.5446
alphaTub84B	GP05054	GCTGACGGTAGGTTCCGGTA	6.20359
esg	GP00225	CTCCACCAACATGTCTTCCA	9.35069
Myo1a	GP00364	ATGGCTATGCAACGGGAAGC	7.98579
btl	GP00400	GGCAAAAGTGCCGATCACGC	3.52487
Mhc	GP00583	CTGAAGCCCCGCATCAAGGT	8.00559
hh	GP01881	TAACCACAGCTCAGTGCCTT	6.99912
Desat1	GP03252	AAGCTGCAGGAGGACTCCAC	6.60024
ap	GP05302	CACCACCTGTAGCACATCAA	5.73411

Supplemental Table 5 – Efficiency scores for 12 sgRNAs used in germ line knock-ins

Name	Sequence	How used
JB880_frame+0_gRNA_top	GTCGgccagtacccaaaaagcggg	for cloning pCFD3-CRISPaint_frameselector
JB881_frame+0_gRNA_bot	AAACcccgcttttttgggtactggc	for cloning pCFD3-CRISPaint_frameselector
JB882_frame+1_gRNA_top	GTCGggccagtacccaaaaagcgg	for cloning pCFD3-CRISPaint_frameselector
JB883_frame+1_gRNA_bot	AAACccgcttttttgggtactggcc	for cloning pCFD3-CRISPaint_frameselector
JB884_frame+2_gRNA_top	GTCGgggccagtacccaaaaagcg	for cloning pCFD3-CRISPaint_frameselector
JB885_frame+2_gRNA_bot	AAACcgcttttttgggtactggccc	for cloning pCFD3-CRISPaint_frameselector
JB886_target-T2A-Gal4overlap_gBlock	ttcacgacctgaggcgcgccgggccagtacccaaaaagcggggggtccggtggaagcggagg tagcgcggccgccgagggccgcgcagcctgctgacctgcggcgatgtggaggagaacccg ggcccGCTAGCatgaagctactgtcttctatcgaacaagcatgcgatatttgccgacttaaa aagctcaagtgctccaaagaaaaaccgaagtgcgccaagtgtct	gBlock containing CRISPaint site and T2A and overlap sequence
JB877_Gal4_F	atgaagctactgtcttctatcgaaca	to amplify Gal4-3xP3RFP
JB878_target-T2A-Gal4_P3RFP_R	gggaacaaaagctggagctcataacttcgtatagcatacattatacgaagttatCGTATGGG CCTTCGCTGCTTACAG	to amplify Gal4-3xP3RFP
JB915_sfGFP_HIKI_T2A_F	tgtggaggagaaccccgggcccgctagcGTGTCCAAGGGCGAGGAG	for cloning into JAB290 cut with NheI/KpnI
JB916_sfGFP_HIKI_R	caaaagatcctctagaggtaccCTACTTGTACAGCTCATCCATGC	for cloning into JAB290 cut with NheI/KpnI
JB1000_LexGAD_HIKI_T2A_F	tgtggaggagaaccccgggcccgctagcATGCCACCCAAGAAGAAGC	for cloning into JAB290 cut with NheI/KpnI
JB1001_LexGAD_HIKI_R	caaaagatcctctagaggtaccCTACTCCTTCTTTGGGTTCGG	for cloning into JAB290 cut with NheI/KpnI
JB998_QF2_HIKI_T2A_F	tgtggaggagaaccccgggcccgctagcATGCCACCCAAGCGCAAA	for cloning into JAB290 cut with NheI/KpnI
JB999_QF2_HIKI_R	caaaagatcctctagaggtaccTCACTGTTCGTATGTATTAATGTCG	for cloning into JAB290 cut with NheI/KpnI
JB1055_Cas9-T2A-EGFP_HIKI_T2A_F	tgtggaggagaaccccgggcccgctagcATGGATTACAAGGATCACGATG	for cloning into JAB290 cut with NheI/KpnI
JB1056_Cas9-T2A-EGFP_HIKI_R	caaaagatcctctagaggtaccTTAGTACAGCTCGTCCATGCC	for cloning into JAB290 cut with NheI/KpnI
JB1002_FLPo_HIKI_T2A_F	tgtggaggagaaccccgggcccgctagcatgagccagttcgacatcct	for cloning into JAB290 cut with NheI/KpnI
JB1003_FLPo_HIKI_R	caaaagatcctctagaggtacctcagatccgcctgttgatgtagc	for cloning into JAB290 cut with NheI/KpnI
JB1004_Gal80_HIKI_T2A_F	tgtggaggagaaccccgggcccgctagcATGGACTACAACAAGAGATCTTCG	for cloning into JAB290 cut with NheI/KpnI
JB1005_Gal80_HIKI_R	caaaagatcctctagaggtaccTTATAAACTATAATGCGAGATATTGCTAA	for cloning into JAB290 cut with NheI/KpnI
JB1008_Nluc_HIKI_T2A_F	tgtggaggagaaccccgggcccgctagcATGGTCTTCACACTCGAAGATTT	for cloning into JAB290 cut with NheI/KpnI
JB1009_Nluc_HIKI_R	caaaagatcctctagaggtaccTTACGCCAGAATGCGTTC	for cloning into JAB290 cut with NheI/KpnI
JB1032_Gal4DBD_HIKI_T2A_F	tgtggaggagaaccccgggcccgctagcATGCTGGAGATCCGCGCC	for cloning into JAB290 cut with NheI/KpnI
JB1033_Gal4DBD_HIKI_R	caaaagatcctctagaggtaccTTACGATACCGTCAGTTGCC	for cloning into JAB290 cut with NheI/KpnI
JB1030_p65_HIKI_T2A_F	tgtggaggagaaccccgggcccgctagcATGGATAAAGCGGAATTAATTCC	for cloning into JAB290 cut with NheI/KpnI
JB1031_p65_HIKI_R	caaaagatcctctagaggtaccTTACTTGCCGCCGCCCAG	for cloning into JAB290 cut with NheI/KpnI
JB917_sfGFP_HIKI_F	ggtggaagcggaggtagcgcggccgccGTGTCCAAGGGCGAGGAG	for cloning into JAB290 cut with NotI/KpnI
JB916_sfGFP_HIKI_R	caaaagatcctctagaggtaccCTACTTGTACAGCTCATCCATGC	for cloning into JAB290 cut with NotI/KpnI
JB969_crispaintsite_NsiI_top	agggccagtacccaaaaagcgggggggTGCA	for cloning into CRIMIC pM37
JB970_crispaintsite_NsiI_bottom	accccccgcttttttgggtactggccctTGCA	for cloning into CRIMIC pM37
JB971_crispaintsite_AgeI-Not1_top	ccggtgggccagtacccaaaaagcggggggggc	for cloning into T-GEM
JB972_crispaintsite_AgeI-Not1_bottom	ggccgcccccccgcttttttgggtactggccca	for cloning into T-GEM
JB1355_FK506-bp2_geno_1F	acgcgccaaaatacaaaaac	amplification of endogenous target gene and knock-in
JB1356_FK506-bp2_geno_1R	GCCTATTCGACCTTGAGCAG	amplification of endogenous target gene
JB1357_alphaTub84B_geno_1F	tttgtgtgggcaaaattcaa	amplification of endogenous target gene and knock-in
JB1358_alphaTub84B_geno_1R	GCTTGGACTTCTTGCCGTAG	amplification of endogenous target gene
JB1359_esg_geno_1F	CGTTTGGTATTTGTGCATCG	amplification of endogenous target gene and knock-in
JB1360_esg_geno_1R	GTAGGGCGACATGTGGAAGT	amplification of endogenous target gene
JB1361_btl_geno_1F	aactaagggaggggcaaaaa	amplification of endogenous target gene and knock-in
JB1362_btl_geno_1R	CGTCCACCAAGGATTTGAGT	amplification of endogenous target gene
JB1363_Desat1_geno_1F	aatccacctggtgcttgttc	amplification of endogenous target gene and knock-in
JB1364_Desat1_geno_1R	GTAACCGAAGGCGATGATGT	amplification of endogenous target gene
JB1365_ap_geno_1F	ttgcaaatctgtcaggaacg	amplification of endogenous target gene and knock-in
JB1366_ap_geno_1R	ATCTGGACACGAGGATGAGG	amplification of endogenous target gene
JB959_ebony_geno_F	gcattagcctgcattgcata	amplification of endogenous target gene and knock-in
JB960_ebony_geno_R	CACGCCCTCATCGAAATAGT	amplification of endogenous target gene
JB961_wg_geno_F	CAGTTAAGCGTTGGCACTGA	amplification of endogenous target gene and knock-in
JB962_wg_geno_R	ttgttgcatctctgcggtag	amplification of endogenous target gene
JB963_Myo1a_geno_F	TCGTCGTCATCAACAGAAGC	amplification of endogenous target gene and knock-in
JB964_Myo1a_geno_R	tctggagtggaaccgaaaac	amplification of endogenous target gene
JB965_Mhc_geno_F	cggctaaagactgacccaaa	amplification of endogenous target gene and knock-in
JB966_Mhc_geno_R	CTCTTGCTCCATGACGAACA	amplification of endogenous target gene
JB967_hh_geno_F	TCGTACTCGCACTCGAACAC	amplification of endogenous target gene and knock-in
JB1388_hh_geno_6F	aaaatcaaagctggaccaaatc	amplification of endogenous target gene and knock-in, used with hh #1 insertion
JB968_hh_geno_R	GTTGTAGTTGGGCACGAGGT	amplification of endogenous target gene
JB900_T2A_R	cgggggttctcctccacat	reverse primer for amplifying T2A-Gal4 insert in sense orientation
JB958_T2A-Gal4_3'_F	gttttcccagtcacgacgtt	reverse primer for amplifying T2A-Gal4 insert in antisense orientation
JB659_3P3dsred_seq1F	ACTCCAAGCTGGACATCACC	to amplify 3xP3-dsred as a control
JB660_3P3dsred_seq1R	CGAGGGTTCGAAATCGATAA	to amplify 3xP3-dsred as a control
JB713_Rp49_F	ATCGGTTACGGATCGAACAA	to amplify endogenous Rp49 gene as a control
JB714_Rp49_R	GACAATCTCCTTGCGCTTCT	to amplify endogenous Rp49 gene as a control
JB1051_alphaTub84B_C-term_F	CCTTCGTCCACTGGTACGTT	amplification of endogenous target gene and knock-in
JB1052_Act5c_C-term_F	CGTCGACCATGAAGATCAAG	amplification of endogenous target gene and knock-in
JB1053_His2Av_C-term_F	CTCCTCGCCACTTACAGCTC	amplification of endogenous target gene and knock-in
JB1054_Lam_C-term_F	GCCGACAACACTAGGACGAT	amplification of endogenous target gene and knock-in
JB1050_mNeonGreen_R	GGGAGAGAGGCGTTATCCTC	reverse primer for amplifying gene-mNeonGreen insert
JB1120_act5c_seqnextgenR	CGACTTCTCCTCCTCCTCCT	amplification of endogenous target gene
JB1121_His2av_seqnextgenR	TCGTCGGTGTTTTAGCTTGTC	amplification of endogenous target gene
JB1122_alphatub_seqnextgenR	GCGATTGGAAGCGTAAACAC	amplification of endogenous target gene
JB1123_Lam_seqnextgenR	GTGTTGTGCTGCGTTTGATT	amplification of endogenous target gene
JB1192_Lam_2F	GCGGCTAATCAACGAGAAAG	amplification of non-knock-in Lamin gene
JB1193_Lam_2R	TCTGTTGTCAGGAGCGTTTG	amplification of non-knock-in Lamin gene
JB1194_Lam_3F	ACGAGGAGCAGATGCAGATT	amplification of non-knock-in Lamin gene
JB1195_Lam_3R	GGTCTAAACCGGGAGAAAGC	amplification of non-knock-in Lamin gene
JB1196_Lam_4F	AGCTGCAGAACCTGAACGAT	amplification of non-knock-in Lamin gene
JB1197_Lam_4R	ACTAGCCGAACCCAGGATTT	amplification of non-knock-in Lamin gene
JB1198_Lam_5F	cccattacaagcgacgattt	amplification of non-knock-in Lamin gene
JB1199_Lam_5R	GAACAGCTCCACTCCTCCAG	amplification of non-knock-in Lamin gene
JB1200_Lam_6F	GTCTCGGTCTGCTCCTCATC	amplification of non-knock-in Lamin gene
JB1201_Lam_6R	GAATGGCATAGCCACCACTT	amplification of non-knock-in Lamin gene
JB1202_Lam_7F	CATTTGCAAGATGGTGGTTG	amplification of non-knock-in Lamin gene
JB1203_Lam_7R	ccaattaggccaacactgct	amplification of non-knock-in Lamin gene
JB1050_mNeonGreen_R	GGGAGAGAGGCGTTATCCTC	amplification and sequencing primer, 5' end of mNeonGreen

Supplemental Table 6 – Oligo and dsDNA sequences