1 **Gene knock-ins in *Drosophila* using homology-independent insertion of universal**
2 **donor plasmids**
3

4 Justin A. Bosch[1], Ryan Colbeth[1], Jonathan Zirin[1], Norbert Perrimon[1,2]
5
6 [1] Department of Genetics, Harvard Medical School, Boston, MA, 02115, USA
7 [2] Howard Hughes Medical Institute, Boston, MA, 02115, USA

8
9 **Short Running Title:** Homology-independent knock-ins in flies
10
11 **Keywords:** CRISPR-Cas9, knock-in, homology-independent, gene function, CRISPaint
12
100-word article summary: We report a fast and simple genomic knock-in method in
Drosophila to insert large DNA elements into any target gene. Using CRISPR-Cas9 and
non-homologous end joining (NHEJ), an entire donor plasmid is inserted into the
genome without the need for homology arms. We demonstrate its usefulness in cultured
cells to fluorescently tag endogenous proteins and in the fly germ line to generate
heritable insertions that disrupt gene function and can act as expression reporters.
19

Corresponding author:
Norbert Perrimon
Harvard Medical School
77 Avenue Louis Pasteur
Dept. of Genetics, NRB 336
Boston, MA 02115
617-432-7672
27
28 Email: perrimon@genetics.med.harvard.edu

29
30 **Abstract**
31

Targeted genomic knock-ins are a valuable tool to probe gene function. However,
knock-in methods involving homology-directed repair (HDR) can be laborious. Here, we
adapt the mammalian CRISPaint homology-independent knock-in method for
Drosophila melanogaster, which uses CRISPR/Cas9 and non-homologous end joining
(NHEJ) to insert universal donor plasmids into the genome. This method is a simple and
fast alternative to HDR for certain strategies such as C-terminal tagging and gene
disruption. Using this method in cultured S2R+ cells, we efficiently tagged four
endogenous proteins with the bright fluorescent protein mNeonGreen, thereby
demonstrating that an existing collection of CRISPaint universal donor plasmids is
compatible with insect cells. In addition, we inserted the transgenesis marker *3xP3-RFP*
into seven genes in the fly germ line, producing heritable loss of function alleles that
were isolated by simple fluorescence screening. Unlike in cultured cells, indels always
occurred at the genomic insertion site, which prevents predictably matching the insert
coding frame to the target gene. Despite this effect, we were able to isolate *T2A-Gal4*
insertions in four genes that serve as in vivo expression reporters. Finally, we apply this
fast knock-in method to uncharacterized small open reading frame (smORF) genes.
Therefore, homology-independent insertion is a useful genome editing technique in
Drosophila that will better enable researchers to dissect gene function.

50
51

Introduction

Insertion of DNA into the animal genome is a powerful method to study gene function. This approach is multipurpose, and can be used, for example, to alter gene function, assay gene expression, visualize protein localization, and purify endogenous proteins (AUER AND DEL BENE 2014; SINGH et al. 2015). Furthermore, the ability to insert large DNA elements such as promoters, protein coding sequences, or entire genes into the genome offers researchers endless options for genome modification. *Drosophila melanogaster* is an excellent animal model to analyze gene function because of its many genetic tools, fast generation time, and in vivo analysis (VENKEN et al. 2016; KORONA et al. 2017; BIER et al. 2018).

Two commonly used methods in *Drosophila* to insert DNA into endogenous genes are transposable DNA elements and homology directed-repair (HDR). Transposable elements insert randomly in the genome (BELLEN et al. 2011) and thus cannot be used to target a user-specified gene. In contrast, HDR is used to insert DNA into a precise genomic location by homologous recombination (BIER et al. 2018). Circular plasmids are commonly used as donor DNA for HDR because they can carry a large DNA insert (≤10kb) and homology arms corresponding to the target locus are added by molecular cloning. However, the design and construction of plasmid donors for each gene can be laborious and prone to troubleshooting. As a cloning-free alternative, synthesized single-stranded DNA (ssDNA) with short homology arms (~50-100 bp each) (BIER et al. 2018) or long ssDNAs of ≤2kb (QUADROS et al. 2017) can be used as donors. However, ssDNA donors are limited to small insertions, and, like plasmid donors, must be designed and generated for each gene that is targeted by HDR. The burden of generating unique donor DNAs for HDR makes this method inefficient for targeting many genes in parallel, such as hits from a genetic or proteomic screen, or the same gene in divergent species. Therefore, there is a need for simpler and more scalable alternatives to knock-in large DNA elements into the *Drosophila* genome.

Large DNA elements can also be inserted in target genes without homology arms, known as homology-independent insertion (CRISTEA et al. 2013; MARESCA et al. 2013; AUER et al. 2014; KATIC et al. 2015; LACKNER et al. 2015; SCHMID-BURGK et al. 2016; SUZUKI et al. 2016; KATOH et al. 2017; KUMAGAI et al. 2017; ZHANG et al. 2018; GAO et al. 2019). In this method, simultaneous cutting of a circular donor plasmid and a genomic target-site by a nuclease results in integration of linearized insert DNA into the genomic cut site by non-homologous end joining (NHEJ). Such donor plasmids are "universal" because they lack gene-specific homology arms, allowing insert sequences (e.g. GFP) to be targeted to any genomic location. Despite this advantage, homology-independent insertion is less precise than HDR. For example, donor DNA can insert in two directions, insertion and deletion mutations (indels) at the integration site can affect the insert translation frame, the entire plasmid can integrate, and inserts can form concatemers. Nevertheless, for some targeting strategies such as C-terminal protein tagging or gene disruption, homology-independent insertion is a fast, simple, and effective alternative to HDR. While homology-independent insertion has been successfully applied in human cell lines (CRISTEA et al. 2013; MARESCA et al. 2013; LACKNER et al. 2015; SCHMID-BURGK et al. 2016; KATOH et al. 2017; ZHANG et al. 2018), mouse somatic cells (SUZUKI et al. 2016; GAO et al. 2019), zebrafish (AUER et al. 2014), *C. elegans* (KATIC et al. 2015), and *Daphnia* (KUMAGAI et al. 2017), it has not yet been utilized in *Drosophila*.

Here, we use the CRISPaint strategy to show that homology-independent insertion functions effectively in *Drosophila*. First, we inserted an mNeonGreen

universal donor plasmid into the C-terminus of four target proteins in S2R+ cells and used Puromycin-resistance to select for in-frame insertions, demonstrating that this is an efficient method to produce cell lines with tagged endogenous proteins. Second, we inserted a *T2A-Gal4 3xP3-RFP* universal donor plasmid into the fly germ line and isolated knock-in lines for seven genes by simple fluorescent screening. By targeting insertions to 5' coding sequence, we demonstrate that this is an effective method to disrupt gene function and generate loss of function fly strains. Furthermore, by screening insertions for Gal4 expression, we identified four that are in vivo reporters of their target gene. Finally, we use this knock-in method to quickly generate insertions in uncharacterized small open reading frame (smORF) genes.

Materials and Methods

Plasmid cloning

    pCFD3-frame_selector_(0, 1, or 2) plasmids (Addgene #127553-127555, DGRC #1482-1484) were cloned by ligating annealed oligos encoding sgRNAs that target the CRISPaint target site (SCHMID-BURGK *et al.* 2016) into *pCFD3* (PORT *et al.* 2014), which contains the *Drosophila U6:3* promoter.
    Additional sgRNA-encoding plasmids were generated by the TRiP (https://fgr.hms.harvard.edu/) or obtained from Filip Port (PORT *et al.* 2015). sgRNA plasmids targeting CDS close to the stop codon were *GP07595* (*Act5c*) (Addgene #130278, DGRC #1492), *GP07596* (*His2Av*), *GP07609* (*alphaTub84B*), and *GP07612* (*Lam*). sgRNA plasmids targeting CDS close to the start codon were *GP06461* (*wg*), *GP02894* (*FK506-bp2*), *GP05054* (*alphaTub84B*), *GP00225* (*esg*), *GP00364* (*Myo1a*), *GP00400* (*btl*), *GP00583* (*Mhc*), *GP01881* (*hh*), *GP03252* (*Desat1*), *GP05302* (*ap*), *pFP545* (*ebony*), and *pFP573* (*ebony*). These sgRNAs were cloned into *pCFD3*, with the exception of those targeting *esg*, *Myo1a*, *btl*, and *Mhc*, which were cloned into *pl100* (KONDO AND UEDA 2013).
    pCRISPaint-T2A-Gal4-3xP3-RFP (Addgene #127556, DGRC #1481) was constructed using Gibson assembly (E2611, NEB) of three DNA fragments: 1) *Gal4-SV40-3xP3-RFP* was PCR amplified from *pHD-Gal4-DsRed* (Xu et al. 2019 in preparation, (GRATZ *et al.* 2014); 2) linear plasmid backbone generated by digesting *pWalium10-roe* (PERKINS *et al.* 2015) with AscI/SacI; and 3) a synthesized double-stranded DNA fragment (gBlock, IDT) encoding the CRISPaint target site, linker sequence, T2A, and ends that overlap the other two fragments.
    pCRISPaint-3xP3-RFP (Addgene # 130276, DGRC #1490) was constructed by digesting *pCRISPaint-T2A-Gal4-3xP3-RFP* with Not1/BamHI to remove *T2A-Gal4* sequences and using Gibson to add a gBlock with a multiple cloning site for future manipulation.
    pCRISPaint-3xP3-GFP (Addgene # 130277, DGRC #1491) was constructed by Gibson assembly of PCR-amplified backbone from *pCRISPaint-3xP3-RFP* and PCR-amplified EGFP sequence from pM37 (LEE *et al.* 2018).
    pCRISPaint-T2A-ORF-3xP3-RFP donor plasmids (Addgene #127557-127565, DGRC #1485-1489,1493-1497) were cloned by PCR amplifying the ORFs and Gibson cloning into *CRISPaint-T2A-Gal4-3xP3-RFP* cut with NheI/KpnI. ORF sequences were amplified from templates as follows: *sfGFP* [amplified from *pUAS-TransTimer* (HE *et al.* 2019)], *LexGAD* [amplified from p*CoinFLP-LexGAD/Gal4* (BOSCH *et al.* 2015)], *QF2* amplified from Addgene #80274, *Cas9-T2A-GFP* (amplified from template kindly provided by Raghuvir Viswanatha, Cas9 from *Streptococcus pyogenes*), *FLPo* (amplified from Addgene #24357), *Gal80* (amplified from Addgene #17748), *Nluc*

(amplified from Addgene #62057), *Gal4DBD*, (amplified from Addgene #26233), and
p65 (amplified from Addgene #26234).
pCRISPaint-sfGFP-3xP3-RFP (Addgene #127566, DGRC #1486) was cloned by
PCR amplifying *sfGFP* coding sequence and Gibson cloning into *CRISPaint-T2A-Gal4-*
3xP3-RFP cut with NotI/KpnI.
pCFD5-frame_selectors_0,1,2 (Addgene #131152) was cloned by PCR
amplifying two fragments encoding the three sgRNAs that cut the CRISPaint target site
and Gibson cloning into *pCFD5* (PORT AND BULLOCK 2016). This plasmid allows
expression of sgRNAs that are separated by tRNA sequences, and thus *pCFD5-*
frame_selectors_0,1,2 expresses all three frame-selector sgRNAs simultaneously.
pLHA-T2A-Gal4-3xP3-RFP-RHA_ebony was cloned by digesting *pCRISPaint-*
T2A-Gal4-3xP3-RFP with AscI/SacI and purifying insert and backbone DNA together
(QIAQuick, Qiagen). Homology arms were amplified from *Drosophila* genomic DNA (*yw*,
single fly) using Phusion polymerase (LHA: 1592bp, RHA: 1544bp). Insert, backbone,
and both homology arms were assembled by Gibson cloning.
See Supplemental Table 7 for oligo and dsDNA sequences and Addgene and
DGRC for plasmid sequences.
171
172   **Cell culture**
173
Drosophila S2R+ cells stably expressing Cas9 and a mCherry protein trap in *Clic*
(known as PT5/Cas9) (VISWANATHA *et al.* 2018) were cultured at 25°C using Schneider's
media (21720-024, ThermoFisher) with 10% FBS (A3912, Sigma) and 50 U/ml penicillin
strep (15070-063, ThermoFisher). S2R+ cells were transfected using Effectene
(301427, Qiagen) following the manufacturer's instructions. Plasmid mixes were
composed of sgRNA-expressing plasmids (see above) and *pCRISPaint-mNeon-PuroR*
(SCHMID-BURGK *et al.* 2016). Cells were transfected with plasmid mixes in 6-well dishes
at 1.8×10^6 cells/ml, split at a dilution of 1:6 after 3-4 days, and incubated with 2 µg/ml
Puromycin (540411 Calbiochem). Every 3-5 days, the media was replaced with fresh
Puromycin until the cultures became confluent (~12-16 days). For single-cell cloning
experiments, cultures were split 1:3 two days before sorting. Cells were resuspended in
fresh media, triturated to break up cell clumps, and pipetted into a cell straining FACS
tube (352235 Corning). Single cells expressing mNeonGreen were sorted into single
wells of a 96 well plate containing 50% conditioned media 50% fresh media using an
Aria-594 instrument at the Harvard Medical School Division of Immunology's Flow
Cytometry Facility. Once colonies were visible by eye (3-4 weeks), they were expanded
and screened for mNeonGreen fluorescence.
191
192   **Fly genetics and embryo injections**
193
Flies were maintained on standard fly food at 25°C. Fly stocks were obtained
from the Perrimon lab collection or Bloomington Stock center (indicated with BL#).
Stocks used in this study are as follows: *yw* (Perrimon Lab), *yw/Y hs-hid* (BL8846), *yw;*
nos-Cas9attP40/CyO (derived from BL78781), *yw;; nos-Cas9attP2* (derived from
BL78782), *yw; Sp hs-hid/CyO* (derived from BL7757), *yw;; Dr hs-hid/TM3,Sb* (derived
from BL7758), *UAS-2xGFP* (BL6874), *wg1-17/CyO* (BL2980), *wg1-8/CyO* (BL5351),
Df(2L)BSC291/CyO (BL23676), *Mhc[k10423]/CyO* (BL10995), Df(2L)H20/CyO
(BL3180), *Df(2L)ED8142/SM6a* (BL24135), *hh[AC]/TM3 Sb* (BL1749),
Df(3R)ED5296/TM3, Sb (BL9338), *esgG66/CyO UAS-GFP* (BL67748),
Df(2R)Exel6069/CyO (BL7551), ywCre; D/TM3, Sb (BL851), Dp(2;1)Sco[rv23];
Df(2L)Sco[rv23], b[1] pr[1]/CyO (BL6230).

For embryo injections, each plasmid was column purified (Qiagen) twice, eluted
in injection buffer (100 µM NaPO4, 5 mM KCl), and adjusted to 200ng/µl. Plasmids were
mixed equally by volume, and mixes were injected into *Drosophila* embryos using
standard procedures. For targeting genes on Chr. 2, plasmid mixes were injected into
yw;; nos-Cas9attP2 embryos. For targeting genes on Chr. 3, plasmid mixes were
injected into *yw; nos-Cas9attP40/CyO)* embryos. Approximately 500 embryos were
injected for each targeted gene.
Injected G0 flies were crossed with *yw*. We used *yw/Y hs-hid* to facilitate
collecting large numbers of virgin flies by incubating larvae and pupae at 37°C for 1hr.
G1 flies were screened for RFP expression in the adult eye on a Zeiss Stemi SVII
fluorescence microscope. G1 RFP+ flies were crossed with the appropriate balancer
stock (*yw; Sp hs-hid/CyO* or *yw;; Dr hs-hid/TM3,Sb)*. G2 RFP+ males that were *yellow-*
(to remove the *nos-Cas9* transgene) and balancer+ were crossed to virgins of the
appropriate balancer stock (*yw; Sp hs-hid/CyO* or *yw;; Dr hs-hid/TM3,Sb*). G3 larvae
and pupae were heat shocked at 37°C for 1hr to eliminate the *hs-hid* chromosome,
which generates a balanced stock (e.g. yw; [RFP+]/CyO).
221
222 **Imaging**
223
S2R+ cells expressing mNeonGreen were plated into wells of a glass-bottom 384
well plate (6007558, PerkinElmer). For fixed cell images, cells were incubated with 4%
paraformaldehyde for 30min, washed with PBS with .1% TritonX-100 (PBT) 3x 5min
each, stained with 1:1000 DAPI (D1306, ThermoFisher) and 1:1000 phalloidin-TRITC
(P1951, Sigma), and washed with PBS. Plates were imaged on an IN Cell Analyzer
6000 (GE) using a 20x or 60x objective. Time-lapse videos of live mNeonGreen
expressing single cell cloned lines were obtained by taking an image every minute using
a 60x objective. Images were processed using Fiji software.
Wing imaginal discs from 3rd instar larvae were dissected in PBS, fixed in 4%
paraformaldehyde, and permeabilized in PBT. For Wg staining, carcasses were blocked
for 1hr in 5% normal goat serum (S-1000, Vector Labs) at room temp, and incubated
with 1:50 mouse anti-wg (4D4, DSHB) primary antibody and 1:500 anti-mouse 488 (A-
21202, Molecular Probes) secondary antibody. Primary and secondary antibody
incubations were performed at 4°C overnight. All carcasses were stained with DAPI and
phalloidin-TRITC, and mounted on glass slides with vectashield (H-1000, Vector
Laboratories Inc.) under a coverslip. Images of mounted wing discs were acquired on a
Zeiss 780 confocal microscope.
Larvae, pupae, and adult flies were imaged using a Zeiss Axio Zoom V16
fluorescence microscope.
243
244 **Quantification of mNeonGreen expressing S2R+ cells**
245
For FACs-based cell counting, we collected cultures from each gene knock-in
experiment before and after puromycin selection. Pre-selection cultures were obtained
by collecting of 500ul of culture 3-4 days after transfection. Post-selection cultures were
obtained after at least 2 weeks of puromycin incubation. Non-transfected cells were
used as a negative control. 100,000 cells were counted for each sample and FlowJo
software was used to analyze and graph the data. FSC-A vs GFP-A was plotted and we
defined mNeonGreen+ cells by setting a signal intensity threshold where <0.02% of
negative controls are counted due to autofluorescence.
For microscopy-based cell counting, the number of mNeonGreen cells was
quantified by analyzing confocal images in Fiji using the manual Cell Counter Plugin

(model). For transfected cells, 6 fields containing at least 200 cells were quantified (i.e.
n=6). For puro-selected cells, 3 fields containing at least 200 cells were quantified (i.e.
n=3).
259
260 **Western blotting**
261
Single cell-cloned cell lines were grown until confluent and 1ml of resuspended
cells was centrifuged at 250g for 10min. The cell pellet was resuspended in 1ml ice cold
PBS, re-centrifuged, and the pellet was lysed in 250ul 2x SDS-Sample buffer and boiled
for 5min. 10ul was loaded on a 4-20% Mini-Protean TGX SDS-Page gel (4561096,
BioRad), transferred to PVDF membrane (IPFL00010, Millipore), blocked in 5% non-fat
dry milk, primary blotting using mouse anti-mNeonGreen (1:1000, Chromtek 32F6) or
hFAB™ Rhodamine Anti-Actin (12004164 BioRad), and secondary blotting using
1:3000 anti-mouse HRP (NXA931, Amersham), imaging using ECL (34580,
ThermoFisher) on a ChemiDoc MP Imaging System (BioRad).
271
272 **PCR, sequencing, and sgRNA cutting assays**
273
S2R+ cell genomic DNA was isolated using QuickExtract (QE09050, Lucigen).
Fly genomic DNA was isolated by grinding a single fly in 50µl squishing buffer (10 mM
Tris-Cl pH 8.2, 1 mM EDTA, 25 mM NaCl) with 200µg/ml Proteinase K (3115879001,
Roche), incubating at 37°C for 30 min, and 95°C for 2 minutes. PCR was performed
using Taq polymerase (TAKR001C, ClonTech) when running DNA fragments on a gel,
and Phusion polymerase (M-0530, NEB) was used when DNA fragments were
sequenced. DNA fragments corresponding to *mNeonGreen* or *T2A-Gal4* insertion sites
were amplified using primer pairs where one primer binds to genomic sequence and the
other primer binds to the insert. For amplifying non-knock-in sites, we used primers that
flank the sgRNA target site. Primer pairs used for gel analysis and/or Sanger
sequencing were designed to produce DNA fragments <1kb. Primer pairs used for next-
generation sequencing of the insertion site were designed to produce DNA fragments
200-280bp. DNA fragments were run on a 1% agarose gel for imaging or purified on
QIAquick columns (28115, Qiagen) for sequencing analysis. See Supplemental Table 7
for oligo sequences.
Sanger sequencing was performed at the DF/HCC DNA Resource Core facility
and chromatograms were analyzed using Lasergene 13 software (DNASTAR). Next-
generation sequencing was performed at the MGH CCIB DNA Core. Fastq files were
analyzed using CRISPresso2 (CLEMENT *et al.* 2019) by entering the PCR fragment
sequence into the exon specification window and setting the window size to 10 bases.
Quantification of insertion types (seamless, in-frame indel, and frameshift indel) was
taken from the allele plot and frameshift analysis outputs of CRISPresso2. The small
proportion of "unmodified" reads that were not called by frameshift analysis were not
included in the quantification.
T7 endonuclease assays (M0302L, NEB) were performed following the
manufacturer instructions.
Splinkerette sequencing was performed as previously described (POTTER AND
LUO 2010). Briefly, genomic DNA was isolated by single fly squishing (described above)
and digested using enzyme BstYI (R0523S, NEB) at 60°C overnight and heat
inactivated for 20min at 80°C. Digested DNA was ligated with annealed splinkerette
oligonucleotides overnight at 16°C. PCR annealing temperatures were 60°C (round 1)
and 64°C (round 2). Round 2 PCR products were either run on an agarose gel and gel
fragments purified using QIAquick Gel Extraction Kit (28704, Qiagen), or purified using

ExoSAP-IT PCR clean up reagent (78200.200.UL, Applied Biosystems). Oligo
sequences used for splinkerette sequencing are listed in Supplemental Table 7. 3'
splinkerette sequence traces, using the *M13R* sequencing primer, frequently mapped to
vermillion, even when genomic DNA was isolated from yw flies. We believe this resulted
from contamination, since commonly used plasmids and fly lines (e.g. *pCFD3* and
pValium20) have *M13R* sequence adjacent to the *vermillion+* marker transgene.
313
314 **Data availability**
315
Plasmids generated in this paper will be available from Addgene and the Drosophila
Genomics Resource Center (DGRC). Gal4-expressing fly strains will be available at the
Bloomington Drosophila Stock Center (BDSC). Two mNeonGreen-expressing S2R+ cell
lines are available at the DGRC (His2Av-mNeonGreen clone B11, DGRC #295; Lamin-
mNeonGreen clone D6, DGRC #296). The remaining flies, cells, and sequence data are
available from the Perrimon lab on request. Oligo and dsDNA sequences are listed in
Supplemental Table 7.
323
324 **Results**
325
To test if homology-independent insertion works in *Drosophila*, we implemented
a strategy known as CRISPaint (SCHMID-BURGK *et al.* 2016). This system is used to
insert sequence encoding a protein tag or reporter gene into the coding sequence of an
endogenous gene. Although it was originally developed for mammalian cell culture,
CRISPaint has several advantages for use in *Drosophila*. First, this system uses
CRISPR-Cas9 to induce double-strand breaks (DSBs), which is known to function
efficiently in *Drosophila* cultured cells (BOTTCHER *et al.* 2014; VISWANATHA *et al.* 2018)
and the germ line (KONDO AND UEDA 2013; REN *et al.* 2013; YU *et al.* 2013; BASSETT *et*
al. 2014). Second, its use of a modular frame-selector system makes it simple to obtain
insertions that are translated with the target gene. Third, a collection of existing
CRISPaint donor plasmids (SCHMID-BURGK *et al.* 2016) containing common tags (e.g.
GFP, RFP, Luciferase) are seemingly compatible for expression in *Drosophila*.
The CRISPaint system works by introducing three components into Cas9-
expressing cells: 1) a single guide RNA (sgRNA) targeting a genomic locus; 2) a donor
plasmid containing an insert sequence; and 3) a frame selector sgRNA targeting the
donor plasmid (Figure 1A). This causes cleavage of both the genomic locus and the
donor plasmid, leading to the integration of the entire linearized donor plasmid into the
genomic cut site by non-homologous end joining (NHEJ). This insertion site destroys
both sgRNA target sites and will no longer be cut. While indels can occur at the
insertion site due to error-prone NHEJ, a remarkably high percentage of insertions in
human cells are seamless (48-86%) (SCHMID-BURGK *et al.* 2016). Therefore, users have
some control over their insertion frame by using one of three frame-selector sgRNAs.
Importantly, these frame-selector sgRNAs do not target the *Drosophila* genome.
349
350 **Homology-independent insertion functions efficiently in *Drosophila* S2R+ cells to**
351 **tag endogenous proteins**
352
To test the CRISPaint method in *Drosophila*, we set out to replicate the findings
of (SCHMID-BURGK *et al.* 2016) in cultured S2R+ cells by tagging endogenous proteins at
their C-terminus. To accomplish this, we generated plasmids expressing frame-selector
sgRNAs (frame 0,1, or 2) under the control of *Drosophila U6* sequences (PORT *et al.*
2015) (Figure 1A). In addition, we generated plasmids expressing sgRNAs that target

the 3' coding sequence of *Drosophila* genes. We chose to target *Actin5c*, *His2Av*,
alphaTub84B, and *Lamin* because these genes are expressed in S2R+ cells (HU *et al.*
2017) and encode proteins with known subcellular localization (actin filaments,
chromatin, microtubules, nuclear envelope, respectively). For donor plasmid, we used
pCRISPaint-mNeonGreen-T2A-PuroR (SCHMID-BURGK *et al.* 2016), which contains a
frame-selector sgRNA target site upstream of coding sequence for the fluorescent
mNeonGreen protein and Puromycin resistance protein (PuroR) linked by a cleavable
T2A peptide sequence. Only integration of the donor plasmid in-frame with the target
coding sequence should result in translation of mNeonGreen-T2A-PuroR (Figure 1A).
We transfected Cas9-expressing S2R+ cells (VISWANATHA *et al.* 2018) with a mix
of three plasmids: *pCRISPaint-mNeonGreen-T2A-PuroR* donor, target-gene sgRNA,
and the appropriate frame-selector sgRNA (Figure 1A, Supplemental Table 1). As an
initial method to detect knock-in events, we used PCR to amplify the predicted insertion
sites from transfected cells. Using primers that are specific to the target gene and
mNeonGreen sequence, we successfully amplified sense-orientation *gene-*
mNeonGreen DNA fragments for all four genes (Figure 1B). Furthermore, next-
generation sequencing of these amplified fragments revealed that 34-50% of sense-
orientation insertions are in frame with the target gene (Figure 1B, Supplemental Figure
1, Supplemental Table 1). 7-27% of sense-orientation insertions are seamless, which is
slightly lower than previously observed (SCHMID-BURGK *et al.* 2016).
Next, we quantified in-frame knock-in frequency by measuring mNeonGreen
fluorescence in transfected S2R+ cells. Flow cytometry-based cell counting of
transfected cells revealed that the number of mNeonGreen+ cells range from 0.19-2.4%
(Figure 1C, Supplemental Table 1), in agreement with human cultured cells (SCHMID-
BURGK *et al.* 2016). These results were confirmed by confocal analysis of transfected
cells, which showed mNeonGreen fluorescence in a small subset of cells (Figure 1C).
3.2% (*Act5c*) and 2.4% (*His2Av*) of transfected cells expressed mNeonGreen (Figure
1C, Supplemental Table 1), which roughly agreed with flow cytometry cell counting.
Finally, mNeonGreen localized to the expected subcellular compartments, most
obviously observed by His2Av-mNeonGreen and Lam-mNeonGreen co-localization with
the nucleus, and Act5c-mNeonGreen and alphatub-mNeonGreen exclusion from the
nucleus (Figure 1C). These results suggest that a significant number of transfected
S2R+ cells received in-frame insertion of mNeonGreen at their C-terminus using the
CRISPaint homology-independent insertion method.
For knock-in cells to be useful in experiments, it is important to derive cultures
where most cells, if not all, carry the insertion. Therefore, we enriched for in-frame
insertion events using Puromycin selection (Figure 1D). After a two-week incubation of
transfected S2R+ cells with Puromycin, flow-cytometry and confocal analysis revealed
that 31.7-89.1% of cells expressed mNeonGreen and exhibited correct subcellular
localization (Figure 1E, Supplemental Table 1). For *alphaTub84B*, cell counting by flow-
cytometry greatly underestimated the number of mNeonGreen+ cells counted by
confocal analysis, likely because mNeonGreen expression level was so low. These
results demonstrate that Puromycin selection is a fast and efficient method of selecting
for mNeonGreen expressing knock-in cells.
A subset of cells in Puro-selected cultures had no mNeonGreen expression or
unexpected localization (Figure 1E). Since each culture is composed of different cells
with independent insertion events, we used FACs to derive single-cell cloned lines
expressing mNeonGreen for further characterization (Figure 2A). At least 14 single-cell
cloned lines were isolated for each target gene and imaged by confocal microscopy.
Within a given clonal culture, each cell exhibited the same mNeonGreen localization
(Figure 2B), confirming our single-cell cloning approach and demonstrating that the

insertion is genetically stable over many cell divisions. Importantly, while many clones
exhibited the predicted mNeonGreen localization, a subset of the clonal cell lines
displayed an unusual localization pattern (Figure 2B). For example, 3/21 Act5c-
mNeonGreen clones had localization in prominent rod structures, and 12/14 Lamin-
mNeonGreen clones had asymmetric localization in the nuclear envelope (Figure 2B).
In addition, some clones had diffuse mNeonGreen localization in the cytoplasm and
nucleus (Figure 2B).
To better characterize the insertions in single cell-cloned lines, we further
analyzed three clones per gene (12 total), selecting different classes when possible
(correct localization, unusual localization, and diffuse localization) (Figure 2C,
Supplemental Table 2). Using PCR amplification of the predicted insertion site (Figure
1A, Figure 2D) and sequencing of amplified fragments (Figure 2E, Supplemental Table
2), we determined that all clones with correct or unusual mNeonGreen localization
contained an in-frame insertion of mNeonGreen with the target gene. In contrast, we
were unable to amplify DNA fragments from the expected insertion site in clones with
diffuse mNeonGreen localization (Figure 2D). Western blotting of cell lysates confirmed
that only clones with in-frame mNeonGreen insertion express fusion proteins that match
the predicted molecular weights (Figure 2F). All together, these results suggest that
clones with correct mNeonGreen localization are likely to contain an in-frame insert in
the correct target gene.
S2R+ cells are polyploid (LEE *et al.* 2014), and clones expressing mNeonGreen
could bear one or more insertions. Furthermore, indels induced at the non-insertion
locus could disrupt protein function. To explore these possibilities, we amplified the non-
insertion locus in our single-cell cloned lines and used Sanger and next-generation
sequencing to analyze the DNA fragments (Figure 2D, Supplemental Table 2). For each
gene, we could find indels occurring at the non-insertion sgRNA cut site. For example,
we could distinguish four distinct alleles in clone B11: a 3bp deletion, a 2bp deletion, a
1bp deletion, and a 27bp deletion. In addition, we identified an unusual mutation in
clone C6, where a 1482bp DNA fragment inserted at the sgRNA cut site, which
corresponds to a region from *alphatub84D*. We assume that this large insertion was
caused by homologous recombination, since *alphatub84D* and *alphatub84B* share 92%
genomic sequence identity (FlyBase, http://flybase.org/). For *Act5c-mNeonGreen* clones
A5 and A19, numerous indel sequences were found, suggesting this region has an
abnormal number of gene copies. We were unable to amplify a DNA fragment from
Lam-mNeonGreen D9, despite follow-up PCRs using primers that bind genomic
sequence further away from the insertion site (not shown).
445
446 **Loss of function knock-in fly lines by homology-independent insertion in the**
447 **germ line**
448
We next explored whether homology-independent insertion could function in the
Drosophila germ line for the purpose of generating knock-in fly strains. As opposed to
antibiotic selection in cultured cells, we wanted to identify transgenic flies using a visible
body marker that was not dependent on target gene expression. In addition, we wanted
to target insertions to 5' coding sequence to create loss of function alleles by premature
protein truncation. Finally, we wanted to test if insertion of a reporter could capture the
expression of the target gene. To accomplish these goals, we constructed a new
universal donor plasmid called *pCRISPaint-T2A-Gal4-3xP3-RFP* (Figure 3A). This
donor contains a frame-selector sgRNA target site upstream of *T2A-Gal4*, which
encodes a self-cleaving form of the transcription factor (DIAO AND WHITE 2012). This

donor plasmid also contains the transgenesis marker *3xP3-RFP*, which expresses red
fluorescence in larval tissues and the adult eye (BERGHAMMER *et al.* 1999) (Figure 3A).
To test if homology-independent insertion functions in the germ line and
determine its approximate efficiency, we targeted 11 genes using *pCRISPaint-T2A-*
Gal4-3xP3-RFP (Supplemental Table 3). These genes were selected based on their
known loss of function phenotypes, expression patterns, and availability of sgRNA
plasmids from the TRiP (https://fgr.hms.harvard.edu/). Plasmid mixes of donor, frame-
selector sgRNA, and target gene sgRNA were injected into *nos-Cas9* embryos, and the
resulting G0 progeny were crossed to *yw*. G1 progeny were screened for RFP
fluorescence in adult eyes, and each RFP+ founder fly was crossed to balancer flies to
establish a stable stock. Figure 3C and Supplemental Table 3 shows the integration
efficiency results for each gene and Supplemental Table 4 has information on 20 RFP+
lines, each derived from a different G0 founder. For injections that produced RFP+
animals, the frequency of G0 crosses yielding RFP+ G1 progeny ranges from 5-21%
(Figure 3C, Supplemental Table 3). For example, when targeting *ebony* with sgRNA
pFP545, 3 out of 16 G0 crosses produced ≥1 RFP+ G1 flies. In one month following
embryo injection, we obtained balanced RFP+ lines for seven out of 11 genes targeted
(64%) (Figure 3C, Supplemental Table 3).
Next, we used PCR and sequencing to confirm the insertion sites in our RFP+
lines. For each target site, we used two primer pairs to detect the presence of an on-
target insertion as well as its orientation (Supplemental Figure 2). Gel images of PCR
amplified DNA showed that all 20 of our RFP+ lines contained an insertion in the correct
target site (Supplemental Table 4, Supplemental Figure 2), where 13 insertions were in
the sense orientation and seven were antisense (Figure 3D). Sequence analysis of PCR
fragments confirmed that the insertions were present at the target site (Supplemental
Table 4, Supplemental Figure 3). Unexpectedly, all 20 lines contained genomic
sequence indels at the insertion site (Supplemental Figure 3), unlike the frequent
seamless insertions observed in cultured cells. Germ line indels were also
predominantly longer than indels in cultured cells. In particular, a remarkably long
1896bp genomic deletion was found at the *hh* insertion site. Regardless, these results
demonstrate that all 20 independently derived RFP+ lines contained *pCRISPaint-T2A-*
Gal4-3xP3-RFP at the target site.
Next, we analyzed insertion lines for loss of function phenotypes. Flies with
insertions in *wg*, *Mhc*, *hh*, and *esg* were homozygous lethal (Supplemental Table 4),
which is consistent with characterized mutations in these genes (FlyBase). Similarly,
complementation tests revealed that insertions in *wg*, *Mhc*, *hh*, and *esg* were lethal in
trans with loss of function alleles or genomic deletions spanning the target gene
(Supplemental Table 4). In addition, homozygous insertions in *hh* and *wg* cause a "lawn
of denticles" phenotype in embryos (BEJSOVEC AND MARTINEZ ARIAS 1991) (Figure 3F)
and flies with heterozygous insertions in *Mhc* were flightless (MOGAMI *et al.* 1986) (data
not shown), all reminiscent of classical allele phenotypes. Four insertions in *ebony*
produced flies with dark cuticle pigment when homozygous (Figure 3E, Supplemental
Table 4) or in trans with *ebony*[1] (not shown). In one case (*ebony pFP545 #2),* the
insertion was homozygous lethal, but was viable over *ebony*[1] and trans-het flies
exhibited dark cuticle pigment. An insertion in *myo1a* (also known as *Myo31DF*) was
viable and flies exhibited genital rotation phenotypes observed in previously reported
loss-of-function alleles (SPEDER *et al.* 2006) (Figure 3G). An insertion in *FK506-bp2* was
homozygous viable, though this gene has not been well characterized in *Drosophila*. In
all, these results indicate that insertion of *pCRISPaint-T2A-Gal4-3xP3-RFP* into 5'
coding sequence can disrupt gene function.

An important consideration when using genome editing is off-target effects. For
example, in addition to on-target insertions, our RFP+ fly lines could contain off-target
insertions or other mutations that disrupt non-target genes. We noticed that RFP
fluorescence always co-segregated with the target chromosome when balancing
insertions (not shown), suggesting that multiple insertions may be uncommon.
Furthermore, we did not detect evidence of off-target insertions by splinkerette PCR
(POTTER AND LUO 2010), which largely reconfirmed the on-target sites (Supplemental
Table 4). However, this analysis was problematic because some sequence traces
aligned to the donor plasmid or to genomic sequence on the opposite expected side of
the target site, suggesting that insertions could be concatemers. Others have observed
concatemers when performing NHEJ knock-ins (AUER *et al.* 2014; KUMAGAI *et al.* 2017),
and we found that seven of our RFP+ insertions were head-to-tail concatemers as
determined by PCR (Supplemental Figure 4). We could rescue the lethality of our *esg*
insertion using a duplication chromosome (stock # BL6230), suggesting there were no
other second site lethal mutations on this chromosome. Finally, one of five *ebony*
insertions (*ebony pFP545 #2*) was homozygous lethal, suggesting it contains a second
site lethal mutation.
We did not obtain insertions when targeting *ap*, *alphaTub84B*, *btl*, or *Desat1*.
Therefore, we investigated whether the sgRNAs targeting these genes were functional.
4 sgRNAs used for germ line knock-ins have an acceptable efficiency score of >5, with
the exception being the sgRNA targeting *btl* (Supplemental Table 5). A T7
endonuclease assay from transfected cells revealed that sgRNAs targeting *ap*,
alphaTub84B, *btl* can cut at the target site, whereas the results with *desat1* were
inconclusive (Supplemental Figure 5A). As an alternative functional test, we used PCR
to detect knock-in events in S2R+ cells transfected with the *pCRISPaint-T2A-Gal4-*
3xP3-RFP donor plasmid. This showed that sgRNAs targeting *ap*, *alphaTub84B*, *btl* and
desat1 can lead to successful knock-in of *pCRISPaint-T2A-Gal4-3xP3-RFP* and
suggests that the sgRNAs are functional (Supplemental Figure 5B). Finally, we
sequenced the sgRNA target sites in the *nos-Cas9* fly strains and found a SNP in the *btl*
sgRNA binding site (not shown), whereas all 10 remaining sgRNAs had no SNPs in the
target site. In summary, we conclude that the sgRNAs targeting *ap*, *alphaTub84B*, *btl*,
and *Desat1* are able to induce cleavage at their target site in S2R+ cells, but that the
sgRNA targeting *btl* may not function in the germ line using our *nos-Cas9* strains.
Collectively, our results demonstrate that the *pCRISPaint-T2A-Gal4-3xP3-RFP*
universal donor plasmid inserts into target loci in the fly germ line and can be used to
generate loss of function fly lines. Therefore, we generated two simplified universal
donor plasmids that were more tailored to this purpose, *pCRISPaint-3xP3-RFP* and
pCRISPaint-3xP3-GFP (Figure 3H). These donor plasmids contain green or red
fluorescent markers and a useful multiple cloning site for adding additional insert
sequence.
549
550   **Knock-ins that express Gal4 under the control of the target gene**
551
We next determined if any of our 20 RFP+ lines expressed Gal4 from the target
gene. Since the *T2A-Gal4* reporter gene in *pCRISPaint-T2A-Gal4-3xP3-RFP* is
promoterless, we reasoned that only insertions in the sense orientation and in-frame
with the target gene should express Gal4. Nonetheless, we took an unbiased approach
by crossing all 20 RFP+ lines to *UAS-GFP* and screening progeny for GFP fluorescence
throughout development. From this effort, we identified 6 Gal4-expressing lines (Figure
4A, Supplemental Tables 3,4). *wg-T2A-Gal4* (#1 and 4), *Mhc-T2A-Gal4* (#1 and 2), and
Myo1a-T2A-Gal4 #1 insertions are expressed in the imaginal disc, larval muscle, and

larval gut (Figure 4B, Supplemental Table 4), respectively, which matches the known
expression patterns for these genes. *wg-T2A-Gal4 #1 and #4* insertions are expressed
in a distinctive Wg pattern in the wing disc pouch (Figure 4C). *ebony-T2A-Gal4 pFP545*
#2 is expressed in the larval brain and throughout the pupal body, consistent with a
previous study (HOVEMANN *et al.* 1998). Interestingly, it is also expressed in the larval
trachea, with particularly strong expression in the anterior and posterior spiracles
(Figure 4D). Indeed, classical *ebony* mutations are known to cause dark pigment in
larval spiracles (BREHME 1941).
Next, we compared the Gal4 expression screening results with our sequence
analysis of the insertion sites. As expected, all lines that expressed Gal4 had insertions
that were in the sense orientation (Supplemental Table 4, Supplemental Figure 2). For
example, *ebony-T2A-Gal4 pFP545 #2* contains a 15bp genomic deletion that is
predicted to keep *T2A-Gal4* in-frame with *ebony* coding sequence. Similarly, *wg-T2A-*
Gal4 #1 contains an in-frame 45bp deletion and 21bp insertion. Remarkably, *wg-T2A-*
Gal4 #4 contains a frameshift indel (Supplemental Figure 3), yet still expresses Gal4 in
the Wg pattern, albeit at significantly lower levels than *wg-T2A-Gal4* #1 (Figure 4C). In
addition, *Mhc-T2A-Gal4* lines #1, #2, and *Myo1a-T2A-Gal4* #1, each have indels that
put *T2A-Gal4* out of frame with the target gene coding sequence. These findings
confirm that our Gal4-expressing lines have *T2A-Gal4* inserted in the sense orientation,
but that in-frame insertion with the target gene coding sequence is not necessarily a
requirement for Gal4 expression.
Artifacts due to homology-independent insertion, such as indels at the insertion
site, could conceivably interfere with Gal4 expression. For example, *ebony* expression
has not been well studied, so we did not know if *ebony-T2A-Gal4 pFP545* #2 ("*ebony-*
Gal4 CRISPaint") was an accurate reporter. To address this issue, we generated a
precise in-frame HDR insertion in *ebony* ("*ebony-Gal4* HDR") using the same insert
sequence (*T2A-Gal4-3xP3-RFP*) and target gene sgRNA (*pFP545*). *ebony-Gal4* HDR
knock-in fly lines were validated by PCR (Supplemental Figure 6) and *ebony-Gal4* HDR
homozygous flies were viable and had dark cuticle pigment (not shown), suggesting on-
target knock-in. By crossing CRISPaint and HDR knock-in alleles with *UAS-GFP*, we
found that their expression patterns were similar at larval, pupal, and adult stages
(Figure 4D,E). However, *ebony-Gal4* CRISPaint had higher levels of RFP fluorescence
and expressed Gal4 in the larval anal pad and gut, which is coincident with expression
of *3xP3-RFP*. Since this insertion is a concatemer (Supplemental Figure 4,
Supplemental Table 4), we speculated that multiple copies of *3xP3-RFP* lead to
increased RFP fluorescence and ectopic expression of Gal4. Unfortunately, we could
not test this because our efforts to remove *3xP3-RFP* using Cre/*loxP* excision were
unsuccessful; progeny expressing Cre and *ebony-Gal4* CRISPaint were apparently
lethal, whereas we easily generated RFP-free derivatives of *ebony-Gal4* HDR using the
same crossing scheme. In contrast, *Mhc-T2A-Gal4* lines #1, #2, and *Myo1a-T2A-Gal4*
#1 do not express in the anal pad, and gut and anal pad expression in *wg-T2A-Gal4* #1
likely corresponds to normal *wg* expression in these tissues (TAKASHIMA AND MURAKAMI
2001) (Figure 4B). In conclusion, these results suggest that Gal4-expressing CRISPaint
insertions can capture the endogenous expression pattern of the target gene, albeit with
caveats that we demonstrated (also see discussion).
To facilitate the insertion of other sequences into the germ line to generate
expression reporters, we generated 10 additional universal donor plasmids (Figure 4F).
These include T2A-containing donors with sequence encoding the alternative binary
reporters LexGAD, QF2, and split-Gal4, as well as Cas9 nuclease, FLP recombinase,
Gal80 repression protein, NanoLuc luminescence reporter, and super-folder GFP. In
addition, we generated *pCRISPaint-sfGFP-3xP3-RFP*, which can be used to insert into

3' coding sequence, generating a C-terminal GFP fusion protein. Finally, we created a
single plasmid expressing all three frame-selector sgRNAs (*pCFD5-frame-*
selectors_0,1,2). Since indels at the insertion site in the germ line effectively
randomizes the coding frame, we reasoned that simultaneous expression of all three
sgRNAs would maximize cutting and linearization of a CRISPaint universal donor
plasmid.
617
618 **Knock-ins targeting uncharacterized small open reading frame (smORF) genes**
619
To further demonstrate that this method can quickly generate knock-in insertions
for any gene, we targeted a class of uncharacterized small open reading frame
(smORF) genes. smORF genes encode peptides <100 amino acids and have been
historically understudied (COUSO AND PATRAQUIM 2017). For example, these genes may
have been missed in mutagenesis and proteomics screens because of their small size.
In addition, they frequently lack long introns useful for isolating exon trap insertions
(BUSZCZAK *et al.* 2007).
Using sequence alignment tools (BLASTP, NCBI; DIOPT, www.flyrnai.org/diopt)
and gene databases (FlyBase; Alliance of Genome Resources,
www.alliancegenome.org/) we identified uncharacterized smORF genes that have clear
human homologues and targeted them for knock-in using sgRNAs that cut 5' coding
sequence and the *pCRISPaint-T2A-Gal4-3xP3-RFP* donor plasmid. Using the same
injection and crossing scheme described above, we isolated 12 insertions in 5 genes as
balanced stocks in 1 month (Supplemental Table 6). Diagnostic PCR analysis confirmed
that all the insertions integrated in the correct target site (Supplemental Figure 7). These
knock-in alleles will be described elsewhere. These results demonstrate that this is a
fast method for producing knock-in lines for any gene.
637
638 **Discussion**
639
Inserting large DNA elements into the genome by HDR requires a great deal of
expertise and labor for the design and construction of donor plasmids. Some groups
have developed strategies to improve the efficiency and scale at which homology arms
are cloned into donor plasmids (HOUSDEN *et al.* 2014; GRATZ *et al.* 2015), but the root
problem still remains. In this study, we showed that homology-independent insertion can
be used in *Drosophila* as an alternative to HDR for certain gene targeting strategies in
cultured cells and in vivo. With no donor design and construction, knock-in lines are
simpler and faster to obtain (see Table 1), making it easier to target multiple genes in
parallel.
To perform homology-independent insertion in *Drosophila*, we adapted the
mammalian CRISPaint system (SCHMID-BURGK *et al.* 2016) because of its user-friendly
design. CRISPaint donor plasmids are "universal" because they lack homology arms,
and publicly available collections enable researchers to "mix and match" insert
sequences and target genes. Indeed, we showed that *pCRISPaint-mNeonGreen-T2A-*
PuroR, originally used in human cells, functions in *Drosophila* S2R+ cells (Figure 1). To
linearize donor plasmids in *Drosophila* in three coding frames, we generated plasmids
expressing the three CRISPaint frame-selector sgRNAs under the *Drosophila U6*
promoter. Similarly, we also generated a plasmid to express all three sgRNAs
simultaneously for maximum cutting efficiency. Therefore, to perform a knock-in
experiment, only an sgRNA that targets a gene of interest is needed, which are simple
to produce by molecular cloning or ordering. In fact, all target gene sgRNAs plasmids

used in this study were obtained from the TRiP facility and an ever-growing number of
additional sgRNAs are being generated.
Unlike HDR, which can seamlessly insert any DNA into a target genomic
location, homology-independent insertion is restricted to specific targeting scenarios.
For example, since the entire *pCRISPaint* donor plasmid integrates into the target site, it
cannot be used to replace genes, make amino acid changes, or tag proteins at their N-
terminus. In addition, potential concatemerization of the insertion (AUER *et al.* 2014;
KUMAGAI *et al.* 2017) prevents its use for strategies where a single copy of the insertion
is required, such as artificial exon traps that exploit target gene splicing (BUSZCZAK *et al.*
2007). Despite these limitations, homology-independent insertion is still well suited to
generate loss of function alleles by premature truncation, as well as C-terminal gene
tagging when a 3' UTR is included.
Using CRISPaint in the fly germ line, we rapidly generated loss of function alleles
by inserting the selectable marker *3xP3-RFP* into target gene coding sequence. This
method has several advantages over isolating frameshift alleles using CRISPR-induced
indels (BIER *et al.* 2018). First, identifying founder flies with RFP fluorescence is easier
and faster than PCR genotyping/sequencing for frameshift indels. Second, insertion into
5' coding sequence is more likely to disrupt the target gene, whereas frameshift alleles
can sometimes retain protein activity (TULADHAR *et al.* 2019). Though, insertions could
also have unintended consequences, such as disrupting neighboring gene expression.
Third, genetic crosses are simplified by tracking insertion alleles with a fluorescent
marker, which could be especially useful in non-*melanogaster* species that do not have
balancer chromosomes. Indeed, the combined use of *3xP3-RFP* and *3xP3-GFP*
universal donor plasmids (Figure 3H) could facilitate generating double mutant lines.
While we did not find evidence for off-target insertions, knock-in lines should be vetted
using traditional techniques, including molecular verification of the insertion site,
comparing independent insertions, outcrossing to wild type, and performing
complementation tests.
Using homology-independent insertion for gene tagging requires screening for
properly expressed inserts, due to the unpredictable and error-prone nature of NHEJ.
For example, donor plasmids can integrate in two orientations and indels at the insertion
site can shift the coding frame. In cultured S2R+ cells, we used antibiotic resistance to
efficiently select for mNeonGreen protein fusions (Figure 1D). However, such an assay
is not feasible in flies, and so we screened germ line *T2A-Gal4* insertions for expression
by crossing with *UAS-GFP* (Figure 3B, Figure 4). This yielded Gal4-expressing lines for
ebony, *wg*, *Mhc*, and *myo1a*. In contrast, we obtained no insertions for *alpthaTub84B*,
btl, *Desat*, *ap*, and one non-Gal4 expressing insertion each for *FK506-bp2*, *esg*, and *hh*.
This highlights the importance of obtaining multiple independently derived germ line
insertions to screen for insert expression. Additional steps could be taken to obtain
more insertions per gene, such as increasing the number of injected embryos or
reattempting with a different sgRNA. Since our knock-in efficiency is roughly similar to
HDR (5-21% (Figure 3C) vs. 5-22% (GRATZ *et al.* 2014), 46-88% (PORT *et al.* 2015), and
7-42% (GRATZ *et al.* 2015)), the limited number of independent germ line insertions may
simply be a constraint of embryo injection-based transgenesis.
Sequence analysis of insertion sites in cultured cells and in vivo revealed
expected and unexpected results. Single cell cloned S2R+ lines expressing
mNeonGreen fusion proteins had insertion sites that were seamless (7/9) or caused an
in-frame 3bp deletion (2/9), which is consistent with human cell data (SCHMID-BURGK *et*
al. 2016). However, all sequenced germ line insertions (20) contained indels at the
insertion site, and indels were larger than in S2R+ cells (Supplemental Figure 3). Germ
cells are known to differ in their NHEJ mechanisms compared to somatic cells (PRESTON

et al. 2006; AHMED *et al.* 2015), but it is not clear why this would prevent seamless
insertions. In addition, 4/6 of Gal4-expressing insertions were out of frame relative to the
target gene (Supplemental Figure 3). We speculate that this could result from ribosome
frameshifting (KETTELER 2012), an internal ribosome entry site (IRES) (KOMAR AND
HATZOGLOU 2005), alternative open reading frames (altORFs) (MOUILLERON *et al.* 2016),
or Gal4 translation initiation from its own start codon. Ultimately, we consider this a
fortuitous effect as long as Gal4 is expressed in the correct pattern. Conversely, *wg #6*
is in-frame but does not express Gal4, perhaps due to a mutation of Gal4 sequences.
Indeed, splinkerette sequencing results from 3/20 of our RFP+ insertions suggest that
portions of the donor plasmid may be deleted during insertion (Supplemental Table 4).
These findings again highlight the importance of screening insertions for insert
expression.
For gene-tagged cell lines and fly lines generated using homology-independent
insertion, potential artifacts that could impact faithful protein fusion or reporter
expression. First, indels at the insertion site could introduce cryptic splice sites, delete
an important sequence region, or cause non-optimal codons that impact translation of
the fusion protein. Second, bacterial sequences in the inserted plasmid may cause
transgene silencing or impact neighboring gene expression (CHEN *et al.* 2004; SUZUKI *et*
al. 2016). Though, we note that thousands of transgenic fly lines contain bacterial
sequences from phiC31 integration (PERKINS *et al.* 2015) with no reports of ill effects.
Third, concatemer insertions could affect gene expression in unpredictable ways.
Indeed, insertion *ebony-T2A-Gal4 pFP545 #2* was a head-to-tail concatemer
(Supplemental Figure 4) and had ectopic Gal4 expression from the *3xP3* enhancer
(Figure 4D,E). Finally, as noted above, the insertion frame may be important for
expression, since *wg-T2A-Gal4 #4* line is expressed weakly and in a non-faithful patchy
pattern in the wing disc. While these are important considerations, we note that 9/9 of
our mNeonGreen tagged S2R+ lines were localized properly, and 5/6 Gal4 expressing
lines appear to be faithful reporters of their target genes, with the above exceptions
noted.
Some of our S2R+ mNeonGreen protein fusion lines exhibited unusual or
unexpected protein localization. In clones D6 and D9, Lamin-mNeonGreen localized to
the nuclear envelope, but in D9 this localization was enriched asymmetrically in the
direction of the previous plane of cell division. Since both clones had seamless
mNeonGreen insertions, we believe the localization difference is caused by the lack of a
non-knock-in locus in D9, whereas D6 contained an in-frame 3bp deletion at the non-
knock-in locus, likely retaining wild-type function. We saw a similar pattern for clones A3
and A5, both of which had seamless insertions in *Actin5c*, but clone A5 exhibited
distinct rod structures. alphaTub84B-mNeonGreen fluorescence and protein levels were
low in all cell lines, despite *alphaTub84B* being highly expressed in S2R+ cells (HU *et al.*
2017). We speculate that the alphaTub84B-mNeonGreen fusion protein is unstable and
previous studies in other organisms have highlighted problems with C-terminal tagging
of alpha-Tubulin (CARMINATI AND STEARNS 1997). Similarly, C-terminal tags can disrupt
Lamin and Actin function (DAVIES *et al.* 2009; NAGASAKI *et al.* 2017). Therefore, it is
important to consider the existing knowledge of the protein when generating C-terminal
protein fusions and examine multiple single cell cloned lines. Importantly, this level of
scrutiny is also required when using HDR to produce knock-ins. Further validation steps
could be done, such as comparing the tagged protein localization to antibody staining of
the untagged wild type protein.
Additional applications of homology-independent insertion in the *Drosophila* germ
line can be imagined. For example, new insert sequences could be added to
pCRISPaint-3xP3-RFP or *pCRISPaint-3xP3-GFP* starting plasmids. In addition, a

modified donor plasmid could be constructed that does not integrate the bacterial plasmid backbone, such as providing donor plasmids as mini-circles (SCHMID-BURGK *et al.* 2016; SUZUKI *et al.* 2016), cutting donor plasmid twice to liberate the insert fragment (LACKNER *et al.* 2015; SUZUKI *et al.* 2016; GAO *et al.* 2019), or using PCR amplified inserts (MANNA *et al.* 2019). Protein tagging could be performed in vivo, but the large indels observed in germ line may limit its use to proteins that have C-terminal tails that can be deleted without altering protein function. Regardless, we generated a donor plasmid for in vivo sfGFP C-terminal fusions (Figure 4F). Finally, we note that most of our germ line donor plasmids contain enzyme restriction sites that can be used to insert genomic homology arms by molecular cloning, making them "dual use" reagents for HDR and homology-independent insertion. Indeed, we used this approach to generate the *ebony-Gal4* HDR allele (see Materials and Methods).

In summary, we demonstrated that homology-independent insertion can be used as an alternative to HDR in *Drosophila*, enabling researchers to rapidly obtain knock-ins without donor design and construction. The most practical application of this approach is to perform C-terminal protein tagging in cultured cells and gene knockout by insertion in vivo. While we obtained in vivo *T2A-Gal4* gene-tagged insertions, it is low efficiency and thus less appealing compared to HDR. However, the techniques required for screening insertions for *T2A-Gal4* are less specialized than those for constructing donor plasmids, making this trade off potentially attractive for those with less molecular biology expertise or who previously failed using HDR. Finally, our in vivo donor plasmids are immediately useable in other arthropod species because of the *3xP3*-fluorescent marker (BERGHAMMER *et al.* 1999), a testament to the modularity of this knock-in system and the benefits of a community of researchers creating and sharing universal donor plasmids.

Acknowledgements

We thank Jonathan Schmid-Burgk for advice and the *CRISPaint-mNeonGreen* donor plasmid, Claire Hu and the TRiP for sgRNA design and construction, Ben Ewen-Campen for valuable comments on the manuscript, Stephanie Mohr, Oguz Kanca, and Hugo Bellen for helpful discussions, Raghuvir Viswanatha for the *Cas9-T2A-EGFP* template sequence, Rich Binari for help with mounting embryos on slides, and Cathryn Murphy for general assistance. We also thank the HMS MicRoN (Microscopy Resources on the North Quad) Core and the HMS Department of Immunology's Flow Cytometry Facility for technical assistance. J.A.B. was supported by the Damon Runyon Foundation. This work was supported by NIH grants R01GM084947, R01GM067761, R24OD019847. N.P. is an investigator of the Howard Hughes Medical Institute.

References:

Ahmed, E. A., H. Scherthan and D. G. de Rooij, 2015 DNA Double Strand Break Response and Limited Repair Capacity in Mouse Elongated Spermatids. Int J Mol Sci 16**:** 29923-29935.

Auer, T. O., and F. Del Bene, 2014 CRISPR/Cas9 and TALEN-mediated knock-in approaches in zebrafish. Methods 69**:** 142-150.

Auer, T. O., K. Duroure, A. De Cian, J. P. Concordet and F. Del Bene, 2014 Highly efficient CRISPR/Cas9-mediated knock-in in zebrafish by homology-independent DNA repair. Genome Res 24**:** 142-153.

Bassett, A. R., C. Tibbit, C. P. Ponting and J. L. Liu, 2014 Mutagenesis and homologous recombination in Drosophila cell lines using CRISPR/Cas9. Biol Open 3**:** 42-49.

Bejsovec, A., and A. Martinez Arias, 1991 Roles of wingless in patterning the larval
epidermis of Drosophila. Development 113**:** 471-485.
Bellen, H. J., R. W. Levis, Y. He, J. W. Carlson, M. Evans-Holm *et al.*, 2011 The
Drosophila gene disruption project: progress using transposons with distinctive
site specificities. Genetics 188**:** 731-743.
Berghammer, A. J., M. Klingler and E. A. Wimmer, 1999 A universal marker for
transgenic insects. Nature 402**:** 370-371.
Bier, E., M. M. Harrison, K. M. O'Connor-Giles and J. Wildonger, 2018 Advances in
Engineering the Fly Genome with the CRISPR-Cas System. Genetics 208**:** 1-18.
Bosch, J. A., N. H. Tran and I. K. Hariharan, 2015 CoinFLP: a system for efficient
mosaic screening and for visualizing clonal boundaries in Drosophila.
Development 142**:** 597-606.
Bottcher, R., M. Hollmann, K. Merk, V. Nitschko, C. Obermaier *et al.*, 2014 Efficient
chromosomal gene modification with CRISPR/cas9 and PCR-based homologous
recombination donors in cultured Drosophila cells. Nucleic Acids Res 42**:** e89.
Brehme, K. S., 1941 The Effect of Adult Body Color Mutations upon the Larva of
Drosophila Melanogaster. Proc Natl Acad Sci U S A 27**:** 254-261.
Buszczak, M., S. Paterno, D. Lighthouse, J. Bachman, J. Planck *et al.*, 2007 The
carnegie protein trap library: a versatile tool for Drosophila developmental
studies. Genetics 175**:** 1505-1531.
Carminati, J. L., and T. Stearns, 1997 Microtubules orient the mitotic spindle in yeast
through dynein-dependent interactions with the cell cortex. J Cell Biol 138**:** 629-
641.
Chen, Z. Y., C. Y. He, L. Meuse and M. A. Kay, 2004 Silencing of episomal transgene
expression by plasmid bacterial DNA elements in vivo. Gene Ther 11**:** 856-864.
Clement, K., H. Rees, M. C. Canver, J. M. Gehrke, R. Farouni *et al.*, 2019
CRISPResso2 provides accurate and rapid genome editing sequence analysis.
Nat Biotechnol 37**:** 224-226.
Couso, J. P., and P. Patraquim, 2017 Classification and function of small open reading
frames. Nat Rev Mol Cell Biol 18**:** 575-589.
Cristea, S., Y. Freyvert, Y. Santiago, M. C. Holmes, F. D. Urnov *et al.*, 2013 In vivo
cleavage of transgene donors promotes nuclease-mediated targeted integration.
Biotechnol Bioeng 110**:** 871-880.
Davies, B. S., L. G. Fong, S. H. Yang, C. Coffinier and S. G. Young, 2009 The
posttranslational processing of prelamin A and disease. Annu Rev Genomics
Hum Genet 10**:** 153-174.
Diao, F., and B. H. White, 2012 A novel approach for directing transgene expression in
Drosophila: T2A-Gal4 in-frame fusion. Genetics 190**:** 1139-1144.
Gao, Y., E. Hisey, T. W. A. Bradshaw, E. Erata, W. E. Brown *et al.*, 2019 Plug-and-Play
Protein Modification Using Homology-Independent Universal Genome
Engineering. Neuron.
Gratz, S. J., C. D. Rubinstein, M. M. Harrison, J. Wildonger and K. M. O'Connor-Giles,
2015 CRISPR-Cas9 Genome Editing in Drosophila. Curr Protoc Mol Biol 111**:** 31
32 31-20.
Gratz, S. J., F. P. Ukken, C. D. Rubinstein, G. Thiede, L. K. Donohue *et al.*, 2014 Highly
specific and efficient CRISPR/Cas9-catalyzed homology-directed repair in
Drosophila. Genetics 196**:** 961-971.
He, L., R. Binari, J. Huang, J. Falo-Sanjuan and N. Perrimon, 2019 In vivo study of gene
expression with an enhanced dual-color fluorescent transcriptional timer. Elife 8.
Housden, B. E., S. Lin and N. Perrimon, 2014 Cas9-based genome editing in
Drosophila. Methods Enzymol 546**:** 415-439.

Hovemann, B. T., R. P. Ryseck, U. Walldorf, K. F. Stortkuhl, I. D. Dietzel *et al.*, 1998
The Drosophila ebony gene is closely related to microbial peptide synthetases
and shows specific cuticle and nervous system expression. Gene 221**:** 1-9.

Hu, Y., A. Comjean, N. Perrimon and S. E. Mohr, 2017 The Drosophila Gene
Expression Tool (DGET) for expression analyses. BMC Bioinformatics 18**:** 98.

Katic, I., L. Xu and R. Ciosk, 2015 CRISPR/Cas9 Genome Editing in Caenorhabditis
elegans: Evaluation of Templates for Homology-Mediated Repair and Knock-Ins
by Homology-Independent DNA Repair. G3 (Bethesda) 5**:** 1649-1656.

Katoh, Y., S. Michisaka, S. Nozaki, T. Funabashi, T. Hirano *et al.*, 2017 Practical
method for targeted disruption of cilia-related genes by using CRISPR/Cas9-
mediated, homology-independent knock-in system. Mol Biol Cell 28**:** 898-906.

Ketteler, R., 2012 On programmed ribosomal frameshifting: the alternative proteomes.
Front Genet 3**:** 242.

Komar, A. A., and M. Hatzoglou, 2005 Internal ribosome entry sites in cellular mRNAs:
mystery of their existence. J Biol Chem 280**:** 23425-23428.

Kondo, S., and R. Ueda, 2013 Highly improved gene targeting by germline-specific
Cas9 expression in Drosophila. Genetics 195**:** 715-721.

Korona, D., S. A. Koestler and S. Russell, 2017 Engineering the Drosophila Genome for
Developmental Biology. J Dev Biol 5.

Kumagai, H., T. Nakanishi, T. Matsuura, Y. Kato and H. Watanabe, 2017 CRISPR/Cas-
mediated knock-in via non-homologous end-joining in the crustacean Daphnia
magna. PLoS One 12**:** e0186112.

Lackner, D. H., A. Carre, P. M. Guzzardo, C. Banning, R. Mangena *et al.*, 2015 A
generic strategy for CRISPR-Cas9-mediated gene tagging. Nat Commun 6**:**
10237.

Lee, H., C. J. McManus, D. Y. Cho, M. Eaton, F. Renda *et al.*, 2014 DNA copy number
evolution in Drosophila cell lines. Genome Biol 15**:** R70.

Lee, P. T., J. Zirin, O. Kanca, W. W. Lin, K. L. Schulze *et al.*, 2018 A gene-specific T2A-
GAL4 library for Drosophila. Elife 7.

Manna, P. T., L. J. Davis and M. S. Robinson, 2019 Fast and cloning-free
CRISPR/Cas9-mediated genomic editing in mammalian cells. bioRxiv.

Maresca, M., V. G. Lin, N. Guo and Y. Yang, 2013 Obligate ligation-gated
recombination (ObLiGaRe): custom-designed nuclease-mediated targeted
integration through nonhomologous end joining. Genome Res 23**:** 539-546.

Mogami, K., P. T. O'Donnell, S. I. Bernstein, T. R. Wright and C. P. Emerson, Jr., 1986
Mutations of the Drosophila myosin heavy-chain gene: effects on transcription,
myosin accumulation, and muscle function. Proc Natl Acad Sci U S A 83**:** 1393-
1397.

Mouilleron, H., V. Delcourt and X. Roucou, 2016 Death of a dogma: eukaryotic mRNAs
can code for more than one protein. Nucleic Acids Res 44**:** 14-23.

Nagasaki, A., T. K. S, T. Yumoto, M. Imaizumi, A. Yamagishi *et al.*, 2017 The Position
of the GFP Tag on Actin Affects the Filament Formation in Mammalian Cells. Cell
Struct Funct 42**:** 131-140.

Perkins, L. A., L. Holderbaum, R. Tao, Y. Hu, R. Sopko *et al.*, 2015 The Transgenic
RNAi Project at Harvard Medical School: Resources and Validation. Genetics
201**:** 843-852.

Port, F., and S. L. Bullock, 2016 Augmenting CRISPR applications in Drosophila with
tRNA-flanked sgRNAs. Nat Methods 13**:** 852-854.

Port, F., H. M. Chen, T. Lee and S. L. Bullock, 2014 Optimized CRISPR/Cas tools for
efficient germline and somatic genome engineering in Drosophila. Proc Natl Acad
Sci U S A 111**:** E2967-2976.

915   Port, F., N. Muschalik and S. L. Bullock, 2015 Systematic Evaluation of Drosophila
916   CRISPR Tools Reveals Safe and Robust Alternatives to Autonomous Gene
917   Drives in Basic Research. G3 (Bethesda) 5**:** 1493-1502.
918   Potter, C. J., and L. Luo, 2010 Splinkerette PCR for mapping transposable elements in
919   Drosophila. PLoS One 5**:** e10168.
920   Preston, C. R., C. C. Flores and W. R. Engels, 2006 Differential usage of alternative
921   pathways of double-strand break repair in Drosophila. Genetics 172**:** 1055-1068.
922   Quadros, R. M., H. Miura, D. W. Harms, H. Akatsuka, T. Sato *et al.*, 2017 Easi-CRISPR:
923   a robust method for one-step generation of mice carrying conditional and
924   insertion alleles using long ssDNA donors and CRISPR ribonucleoproteins.
925   Genome Biol 18**:** 92.
926   Ren, X., J. Sun, B. E. Housden, Y. Hu, C. Roesel *et al.*, 2013 Optimized gene editing
927   technology for Drosophila melanogaster using germ line-specific Cas9. Proc Natl
928   Acad Sci U S A 110**:** 19012-19017.
929   Schmid-Burgk, J. L., K. Honing, T. S. Ebert and V. Hornung, 2016 CRISPaint allows
930   modular base-specific gene tagging using a ligase-4-dependent mechanism. Nat
931   Commun 7**:** 12338.
932   Singh, P., J. C. Schimenti and E. Bolcun-Filas, 2015 A mouse geneticist's practical
933   guide to CRISPR applications. Genetics 199**:** 1-15.
934   Speder, P., G. Adam and S. Noselli, 2006 Type ID unconventional myosin controls left-
935   right asymmetry in Drosophila. Nature 440**:** 803-807.
936   Suzuki, K., Y. Tsunekawa, R. Hernandez-Benitez, J. Wu, J. Zhu *et al.*, 2016 In vivo
937   genome editing via CRISPR/Cas9 mediated homology-independent targeted
938   integration. Nature 540**:** 144-149.
939   Takashima, S., and R. Murakami, 2001 Regulation of pattern formation in the
940   Drosophila hindgut by wg, hh, dpp, and en. Mech Dev 101**:** 79-90.
941   Tuladhar, R., Y. Yeu, J. Tyler Piazza, Z. Tan, J. Rene Clemenceau *et al.*, 2019
942   CRISPR-Cas9-based mutagenesis frequently provokes on-target mRNA
943   misregulation. Nat Commun 10**:** 4056.
944   Venken, K. J., A. Sarrion-Perdigones, P. J. Vandeventer, N. S. Abel, A. E. Christiansen
945   *et al.*, 2016 Genome engineering: Drosophila melanogaster and beyond. Wiley
946   Interdiscip Rev Dev Biol 5**:** 233-267.
947   Viswanatha, R., Z. Li, Y. Hu and N. Perrimon, 2018 Pooled genome-wide CRISPR
948   screening for basal and context-specific fitness gene essentiality in Drosophila
949   cells. Elife 7.
950   Yu, Z., M. Ren, Z. Wang, B. Zhang, Y. S. Rong *et al.*, 2013 Highly efficient genome
951   modifications mediated by CRISPR/Cas9 in Drosophila. Genetics 195**:** 289-291.
952   Zhang, C., X. He, Y. K. Kwok, F. Wang, J. Xue *et al.*, 2018 Homology-independent
953   multiallelic disruption via CRISPR/Cas9-based knock-in yields distinct functional
954   outcomes in human cells. BMC Biol 16**:** 151.
955

Figure 1

Figure 1 – Knock-in of *mNeonGreen-T2A-PuroR* into Drosophila S2R+ cells using homology-independent insertion. (**A**) Schematic of CRISPaint knock-in approach. *mNeonGreen-T2A-PuroR* is inserted into 3' coding sequence. (**B**) Analysis of knock-in efficiency of transfected cells by diagnostic PCR (DNA gel image) and next-generation sequencing (pie charts). (**C**) Analysis of knock-in efficiency of transfected cells by FACs and confocal microscopy. Numbers indicate percentage of cells with fluorescence. F-actin stained using Phalloidin-TRITC (red), nuclei labeled with DAPI (blue), mNeonGreen signal is in green. Scale bar 10µm. (**D**) Schematic of Puromycin selection of mNeonGreen-expressing cells. (**E**) Analysis of knock-in frequency of puromycin-selected cells using FACs and confocal microscopy. Numbers indicate percentage of cells with green fluorescence. Scale bar 10µm.

Figure 2

Figure 2. Analysis of S2R+ mNeonGreen-expressing single-cell cloned lines. (**A**) Schematic of FACS isolation of single-cell clones expressing mNeonGreen. (**B**) Confocal images of live mNeonGreen-expressing cell lines, categorized into three clone types. Numbers indicate the frequency of each clone type for each gene targeted. Images show fluorescence from Clic-mCherry (red) and mNeonGreen (green). Scale bar 25μm. (**C**) Single cell cloned lines retained for further analysis. (**D**) Agarose gel with PCR fragments amplified from knock-in (Gene_F/mNeonGreen_R) and non-knock-in loci (Gene_F/R). Positive control bands were amplified from *Rp49* genomic sequence. (**E**) Example sequence results of *His2Av-mNeonGreen* clones. sgRNA target site and PAM sequence shown as grey bars, Cas9 cut sites shown with arrowheads. (**F**) Western blot detecting mNeonGreen protein fusions. Arrowheads indicate expected molecular weight. X's indicate incorrect molecular weight.

Figure 3

Figure 3 – Germ line knock-ins using homology-independent insertion. (**A**) Schematic of knock-in approach. *pCRISPaint-T2A-Gal4-3xP3-RFP* is inserted into 5' coding sequence. Inset shows images of adult flies with 3xP3-RFP fluorescence in the eye. Top panel is brightfield, bottom panel is fluorescence. (**B**) Schematic of plasmid injections, fly crosses, and analysis of insertions. (**C**) Graph with results of knock-in efficiency for 12 sgRNA target sites and 11 genes. (**D**) Insert orientation and frame of 20 RFP+ fly lines. (**E**) Image of adult flies. Homozygous *ebony-T2A-Gal4 FP545* #1 flies have dark cuticle pigment. (**F**) Darkfield images of *hh* #1 and *wg* #1 homozygous embryo "lawn of denticles" phenotypes. (**G**) Quantification of genital rotation in wild type (*yw*) and a *myo1a* insertion line. Rotation is determined from the direction of wrapping of the adult male spermiduct around the gut from dissected live abdomens. (**H**). Diagram of two simplified universal donor plasmids for germ line knock-ins.

Figure 4

Figure 4 – Germ line insertions that express *T2A-Gal4* under the control of the target gene. (**A**) Proportion of 20 RFP+ knock-in lines that express Gal4. (**B**) Fluorescence images of 3rd instar larvae with indicated genotypes. Two larvae of the same genotype are shown as rotated in different angles. Expression of Gal4 under control of the target gene drives expression of the *UAS-GFP* reporter. (**C**) Confocal images of wing imaginal discs showing protein staining of Wg protein (anti-wg, green) or *UAS-GFP* expression (green). GFP fluorescence was recorded at identical exposure settings for lines *wg-T2A-Gal4* #1 and #4. Inset shows digitally increased GFP signal. Scale bar 50µm. (**D,E**) Fluorescence images of 3rd instar larvae and pupae. Fluorescence is from *3xP3-RFP* (red) and Gal4 expression with *UAS-GFP* (green). (**D**) shows results with *ebony-T2A-Gal4 pFP545* #2 (*ebony-Gal4* CRISPaint) and (**E**) shows results with *ebony-Gal4* HDR. (**F**) Additional universal donor plasmids for insertion into 5' coding sequence.

S2R+ gene tagging (e.g. mNeonGreen)	Donor design and construction	sgRNA cloning	Transfection -> single cell clones	Molecular characterization	Screening expression
HDR knock-in	Variable, 1-3 weeks	3 days	2 months	1-2 days (PCR, Gel image)	1 day, confocal microscopy
CRISPaint knock-in	N/A	3 days	2 months	3 days (PCR, Gel image, Sequence)	1 day, confocal microscopy

In vivo KO	Donor design and construction	sgRNA cloning	Injection -> fly stock	Molecular characterization
HDR knock-in	Variable, 1-3 weeks	3 days	1 month	1-2 days (PCR, Gel image)
In/del frameshift	N/A	3 days	1 month	3-7 days (PCR, Gel image, Sequence)
CRISPaint knock-in	N/A	3 days	1 month	3 days (PCR, Gel image, Sequence)

In vivo gene tagging (e.g. T2A-Gal4)	Donor design and construction	sgRNA cloning	Injection -> fly stock	Molecular characterization	Screening expression
HDR knock-in	Variable, 1-3 weeks	3 days	1 month	1-2 days (PCR, Gel image)	Optional though recommended: Cross with UAS-GFP, 5-14 days
CRISPaint knock-in	N/A	3 days	1 month	3 days (PCR, Gel image, Sequence)	Cross with UAS-GFP, 5-14 days

Table 1 – Knock-in timeline comparison in S2R+ cells and in vivo for CRISPaint and HDR

Supplemental Figure 1

Supplemental Figure 1 – Next-generation sequencing analysis of mNeonGreen insertion sites in transfected S2R+ cells using CRISPresso2

Supplemental Figure 2 – Diagnostic PCR of *CRISPaint-T2A-Gal4* insertions in RFP+ fly lines to confirm their insertion site and orientation.

Supplemental Figure 3

Supplemental Figure 3 – Sequence structure of *CRISPaint-T2A-Gal4* insertions in RFP+ fly lines

Supplemental Figure 4

Supplemental Figure 4 – Diagnostic PCR of *CRISPaint-T2A-Gal4* insertions in RFP+ fly lines to detect concatemers

Supplemental Figure 5

A

B

Supplemental Figure 5 – Cutting efficiency of 12 sgRNAs in transfected S2R+ cells. (**A**) T7 endonuclease assay. (**B**) Diagnostic PCR to detect for presence of sense orientation *CRISPaint-T2A-Gal4* insertion events.

A

ebony HDR Donor plasmid

B

Supplemental Figure 6 – Construction of *ebony-Gal4* HDR insertion. (**A**) Targeting strategy of *ebony* HDR donor plasmid and *ebony* genomic locus. (**B**) Diagnostic PCR of *ebony-Gal4* HDR insertion. Asterisks indicate correct size band.

Supplemental Figure 7 – Diagnostic PCR of CRISPaint-T2A-Gal4 insertions in smORF genes

	Gene and sgRNA information			
Gene Name	Protein localization	sgRNA plasmid (TRiP)	sgRNA target site with PAM	frame selector sgRNA
Act5c	actin	GP07595	GCGGTGCACAATGGAGGGGCCGG	2
His2Av	chromatin	GP07596	GGTGCAGGATCCGCAGCGGAAGG	2
αTub84B	microtubules	GP07609	CGGCGAGGGTGAGGGCGCTGAGG	2
Lamin	nuclear envelope	GP07612	CGCAGCGTGACCGCCGTGGACGG	1
Untransfected				

	CRISPresso2 output			
Gene Name	seamless	in-frame in/del	frameshift in/del	% in-frame
Act5c	5543	1343	13068	34.50937155
His2Av	2384	2620	8279	37.6722126
αTub84B	726	3034	6901	35.26873652
Lamin	971	801	1757	50.21252479
Untransfected				

	FACs Transfected			FACs Puro-selected		
Gene Name	mNeonGreen+	Single cells total	% mNeonGreen+	mNeonGreen+	Single cells total	% mNeonGreen
Act5c	1963	80589	2.435816302	45355	50883	89.1358607
His2Av	1554	80291	1.935459765	32780	46298	70.80219448
αTub84B	150	78752	0.190471353	16682	52643	31.6889235
Lamin	385	81688	0.471305455	34826	49234	70.73567047
Untransfected	2	83869	0.002384671	10	55656	0.017967515

	Cell counting transfected (n=6)			Cell counting Puro-selected (n=3)		
Gene Name	mNeonGreen+	Total cells	Average % mNeonGreen+	mNeonGreen+	Total cells	Average % mNeonGreen+
Act5c	17/17/26/18/22/25	571/679/577/599/574/613	3.5	342/280/318	397/398/377	80.3
His2Av	26/21/12/19/13/19	634/634/568/525/669/612	3	314/327/315	397/402/400	79.7
αTub84B				412/400/403	412/400/403	100
Lamin				270/282/259	382/388/374	70.9
Untransfected						

Supplemental Table 1 – Quantification of CRISPaint-mNeonGreen insertion events in transfected and puro-selected S2R+ cells by CRISPresso2, FACs, and cell counting of confocal images.

Clone name	Gene-mNeonGreen	predicted mNeonGreen fusion (kDa)	Observed Localization	mNeon insertion site	endogenous gene			
					allele 1	allele 2	allele 3	allele 4
A3	Act5c	71.4	very infrequent small rods	seamless	1bp deletion	1bp insertion		
A5	Act5c	71.4	very frequent long rods	seamless	numerous in/dels			
A19	Act5c	71.4	no rods, cyto and nuclear	N/A	numerous in/dels			
B1	His2Av	44.5	cyto and nuclear	N/A	4bp deletion	19bp deletion		
B11	His2Av	44.5	nuclear	seamless	3bp deletion	2bp deletion	1bp deletion	27bp deletion
B14	His2Av	44.5	nuclear	3 bp deletion, 1bp change	19bp deletion	3bp deletion	1bp deletion	
C2	alphatub84B	79.5	mitotic spindles	seamless	wt	6bp deletion	1bp deletion	
C6	alphatub84B	79.5	mitotic spindles	seamless	insertion of 1482bp from alphatub84D			
C13	alphatub84B	79.5	mitotic spindles	3bp deletion	wt	6bp insertion		
D1	Lam	100.9	cytoplasmic, nuclear	N/A	8bp deletion	4bp deletion	3bp insertion	
D6	Lam	100.9	nuclear lamina	seamless	3bp deletion			
D9	Lam	100.9	nuclear lamina, polarized	seamless	N/A			

Supplemental Table 2 – Molecular characterization of single cell cloned mNeonGreen-expressing S2R+ lines.

Gene	Chr.	target gene sgRNA	target site (no PAM)	Frame selector sgRNA	# G0 crosses	# G0 founders	% founders/total	# G1 progeny screened	# G1 RFP+ progeny	% RFP G1/total G1
Mhc	2	GP00583	CTGAAGCCCCGCATCAAGGT	1	29	6	21	2086	21	1.01
ebony	3	pFP545	TGGCCATCTGGAAGGCTGG	1	16	3	19	1298	11	0.85
FK506-bp2	2	GP02894	TGTAGTGGACCGTGACCTTT	1	13	2	15	901	2	0.22
wg	2	GP06461	GGGGCCGGGGCTCCATGTGG	0	40	6	15	4423	26	0.59
ebony	3	pFP573	TCTACACCTCGGGCAGTAC	1	23	3	13	2121	27	1.27
esg	2	GP00225	CTCCACCAACATGTCTTCCA	2	15	1	7	772	1	0.13
Myo1a	2	GP00364	ATGGCTATGCAACGGGAAGC	1	17	1	6	860	1	0.12
hh	3	GP01881	TAACCACAGCTCAGTGCCTT	2	39	2	5	1975	3	0.15
alphaTub84B	3	GP05054	GCTGACGGTAGGTTCCGGTA	1	6	0	0	506	0	0.00
btl	2	GP00400	GGCAAAAGTGCCGATCACGC	2	26	0	0	2085	0	0.00
Desat1	3	GP03252	AAGCTGCAGGAGGACTCCAC	1	7	0	0	386	0	0.00
ap	2	GP05302	CACCACCTGTAGCACATCAA	0	29	0	0	2928	0	0.00

Supplemental Table 3 – Germ line knock-in efficiency of *CRISPaint-T2A-Gal4*

#	Gene targeted	sgRNA	G0 cross #	Head-tail insertion by PCR?	Splinkerette 5' (PCR Purification)	Splinkerette 5' (Gel Purification)	Splinkerette 3' (PCR Purification)	Splinkerette 3' (Gel Purification)	Insert orientation	Insert frame	Insertion site sequence	Gal4 expression	Homozygous phenotype	Complementation test
1	ebony	pFP545	1	no	ebony 5' insertion site	no bands	ebony 3' insertion site	ebony 3' insertion site	Sense	Out	8bp del	none	dark cuticle pigment	dark cuticle pigment over e1 on TM3
2	ebony	pFP545	2	yes	ebony 3' insertion site	Band 1: ebony 3' insertion site Band 2: 3' end of donor plasmid	5' end of donor plasmid	5' end of donor plasmid	Sense	In	15bp del	anterior and posterior trachae in larvae, whole body cuticle in pupae and adult	lethal	dark cuticle pigment over e1 on TM3
3	ebony	pFP573	1	no	internal donor plasmid	internal donor plasmid	vermillion	vermillion	Sense	Out	7bp del	none	dark cuticle pigment	dark cuticle pigment over e1 on TM3
4	ebony	pFP573	2	no	ebony 3' insertion site	ebony 3' insertion site	vermillion	vermilllion	Antisense	N/A	5bp del, 25bp ins	none	dark cuticle pigment	dark cuticle pigment over e1 on TM3
5	ebony	pFP573	3	yes	3' end of donor plasmid	3' end of donor plasmid	vermillion	bad sequence data	Sense	Out	8bp deletion	none	dark cuticle pigment	dark cuticle pigment over e1 on TM3
6	hh	GP01881	1	no	bad sequence data	no bands	vermillion	Band 1: hh 5' insertion site Band 2: hh 3' insertion site Band 3: vermillion	Antisense	N/A	1897bp del, 8bp ins	none	lethal	fail to complement hh[AC] and Df(3R)ED5296
7	Mhc	GP00583	1	no	Mhc 5' insertion site	no bands	vermillion	Band 1: Mhc 5' insertion site Band 2: vermillion	Sense	Out	10bp del, 3bp ins	muscle	lethal	fail to complement P{lacW}Mhck10423 and Df(2L)H20
8	Mhc	GP00583	2	no	Mhc 5' insertion site	no bands	bad sequence data	bad sequence data	Sense	Out	9bp del, 13bp ins	muscle	lethal	fail to complement P{lacW}Mhck10423 and Df(2L)H20
9	Mhc	GP00583	3	no	Mhc 3' insertion site	Mhc 3' insertion site	bad sequence data	bad sequence data	Antisense	N/A	27bp del, 1bp ins	none	lethal	fail to complement P{lacW}Mhck10423 and Df(2L)H20
10	Mhc	GP00583	4	yes	Mhc 3' insertion site	Band 1: 3' end of donor plasmid Band 2: 3' end of donor plasmid	bad sequence data	Band 1: 5' end of donor plasmid Band 2: internal donor plasmid	Sense	Out	2bp del, 4bp ins	none	lethal	fail to complement P{lacW}Mhck10423 and Df(2L)H20
11	Mhc	GP00583	5	yes	Mhc 3' insertion site	Band 1: Mhc 3' insertion site Band 2: 3' end of donor plasmid	5' end of donor plasmid	5' end of donor plasmid	Antisense	N/A	2bp deletion	none	lethal	fail to complement P{lacW}Mhck10423 and Df(2L)H20
12	Myo1a	GP00364	1	no	Myo1a 5' insertion site	Myo1a, 5' insertion site	internal donor plasmid	internal donor plasmid	Sense	Out	4bp del, 2bp ins	gut	viable	N/A
13	esg	GP00225	1	yes	esg, 3' insertion site	Band 1: esg, 3' insertion site Band 2: bad sequence data	vermillion	Band 1: 5' end of donor plasmid Band 2: vermillion	Antisense	N/A	25bp del, 10bp	none	lethal	fail to complement P{enG}esg[G66]
14	FK506-bp2	GP02894	1	no	FK506-bp2 3' insertion site	FK506-bp2 3' insertion site	FK506-bp2 5' insertion site	Band 1: FK506-bp2 5' insertion site Band 2: vermillion	Antisense	N/A	30bp del	none	viable	N/A
15	wg	GP06461	1	yes	wg 3' insertion site	Band 1: wg 3' insertion site Band 2: 3' end of donor plasmid	vermillion	Band 1: wg 5' insertion site Band 2: 5' end of donor plasmid	Sense	In	45bp del, 21bp ins	wingless, strong expression	lethal	fail to complement wg[l-17], wg[l-8], and Df(2L)BSC291
16	wg	GP06461	2	no	3' end of donor plasmid	3' plasmid	wg 5' insertion site	5' end of donor plasmid	Antisense	N/A	33bp del, 11bp ins	none	lethal	fail to complement wg[l-17], wg[l-8], and Df(2L)BSC291
17	wg	GP06461	3	yes	wg 5' insertion site	Band 1: wg 5' insertion site Band 2: wg 3' insertion site Band 3: 3' end of donor plasmid	vermillion	wg 3' insertion site	Sense	Out	21bp del, 11bp ins	none	lethal	fail to complement wg[l-17], wg[l-8], and Df(2L)BSC291
18	wg	GP06461	4	no	wg 5' insertion site	Band 1: wg 5' insertion site Band 2: 3' end of donor plasmid	wg 3' insertion site	Band 1: wg 3' insertion site Band 2: attB sequence?	Sense	Out	32bp del, 7bp ins	wingless, weak expression	lethal	fail to complement wg[l-17], wg[l-8], and Df(2L)BSC291
19	wg	GP06461	5	no	wg 5' insertion site	wg 5' insertion site	wg 3' insertion site	vermillion	Sense	Out	28bp del, 5bp ins	none	lethal	fail to complement wg[l-17], wg[l-8], and Df(2L)BSC291
20	wg	GP06461	6	no	wg 5' insertion site	Band 1: wg 5' insertion site Band 2: wg 3' insertion site	vermillion	no band	Sense	In	21bp insertion	none	lethal	fail to complement wg[l-17], wg[l-8], and Df(2L)BSC291

Supplemental Table 4 – Molecular and phenotypic characterization of 20 RFP+ fly strains, each carrying a distinct *CRISPaint-T2A-Gal4* insertion.

gene	sgRNA name	sgRNA sequence	DRSC efficiency score
ebony	pFP545	gTGGCCATCTGGAAGGCTGG	5.56968
ebony	pFP573	gTCTACACCTCGGGCAGTAC	5.08877
wg	GP06461	GGGGCCGGGGCTCCATGTGG	5.61857
FK506-bp2	GP02894	TGTAGTGGACCGTGACCTTT	6.5446
alphaTub84B	GP05054	GCTGACGGTAGGTTCCGGTA	6.20359
esg	GP00225	CTCCACCAACATGTCTTCCA	9.35069
Myo1a	GP00364	ATGGCTATGCAACGGGAAGC	7.98579
btl	GP00400	GGCAAAAGTGCCGATCACGC	3.52487
Mhc	GP00583	CTGAAGCCCCGCATCAAGGT	8.00559
hh	GP01881	TAACCACAGCTCAGTGCCTT	6.99912
Desat1	GP03252	AAGCTGCAGGAGGACTCCAC	6.60024
ap	GP05302	CACCACCTGTAGCACATCAA	5.73411

Supplemental Table 5 – Efficiency scores for 12 sgRNAs used in germ line knock-ins

Gene targeted	FBgn	Size (aa)	Human homolgoue	sgRNA	sgRNA target site	G0 cross #	Insert orientation
bc10	FBgn0040239	79	BLCAP	GP01409	GGGTTTCGGTATCAGTAGTA	1	Antisense
CG42308	FBgn0259204	61	C12orf73	GP01629	ATGCCCGCCGGAGTTTCCTG	1	Sense
CG18336	FBgn0040763	94	FAM166A	GP02853	GGGTAGCACCATCCCTCATC	1	Sense
						2	Sense
						3	Antisense
						4	Antisense
						5	Antisense
						6	Sense
CG42381	FBgn0259727	76	CMC4	GP01437	TGGTCAGACAGGCTTGTATG	1	Sense
CG42518	FBgn0260392	85	NCBP2-AS2	GP06479	GCAGCTCATCCAGCGCATGG	1	Sense
						2	Sense
						3	Sense

Supplemental Table 6 – smORF genes targeted for knock-in with pCRISPaint-T2A-Gal4-3xP3-RFP

Name	Sequence	How used
JB880_frame+0_gRNA_top	GTCGgccagtacccaaaaagcggg	for cloning pCFD3-CRISPaint_frameselector
JB881_frame+0_gRNA_bot	AAACcccgcttttttgggtactggc	for cloning pCFD3-CRISPaint_frameselector
JB882_frame+1_gRNA_top	GTCGggccagtacccaaaaagcgg	for cloning pCFD3-CRISPaint_frameselector
JB883_frame+1_gRNA_bot	AAACccgcttttttgggtactggcc	for cloning pCFD3-CRISPaint_frameselector
JB884_frame+2_gRNA_top	GTCGgggccagtacccaaaaagcg	for cloning pCFD3-CRISPaint_frameselector
JB885_frame+2_gRNA_bot	AAACcgcttttttgggtactggccc	for cloning pCFD3-CRISPaint_frameselector
JB886_target-T2A-Gal4overlap_gBlock	ttcacgacctgaggcgcgccgggccagtacccaaaaagcggggggtccggtggaagcggaggtagcgcggccgccgagg gccgcggcagcctgctgacctgcggcgatgtggaggagaaccccgggcccGCTAGCatgaagctactgtcttctatcga acaagcatgcgatatttgccgacttaaaaagctcaagtgctccaaagaaaaaccgaagtgcgccaagtgtct	gBlock containing CRISPaint site and T2A and overlap sequence
JB877_Gal4_F	atgaagctactgtcttctatcgaaca	to amplify Gal4-3xP3RFP
JB878_target-T2A-Gal4_P3RFP_R	gggaacaaaagctggagctcataacttcgtatagcatacattatacgaagttatCGTATGGGCCTTCGCTGCTTACAG	to amplify Gal4-3xP3RFP
JB915_sfGFP_HIKI_T2A_F	tgtggaggagaaccccgggcccgctagcGTGTCCAAGGGCGAGGAG	for cloning into JAB290 cut with NheI/KpnI
JB916_sfGFP_HIKI_R	caaagatcctctagaggtaccCTACTTGTACAGCTCATCCATGC	for cloning into JAB290 cut with NheI/KpnI
JB1000_LexGAD_HIKI_T2A_F	tgtggaggagaaccccgggcccgctagcATGCCACCCAAGAAGAAGC	for cloning into JAB290 cut with NheI/KpnI
JB1001_LexGAD_HIKI_R	caaagatcctctagaggtaccCTACTCCTTCTTTGGGTTCGG	for cloning into JAB290 cut with NheI/KpnI
JB998_QF2_HIKI_T2A_F	tgtggaggagaaccccgggcccgctagcATGCCACCCAAGCGCAAA	for cloning into JAB290 cut with NheI/KpnI
JB999_QF2_HIKI_R	caaagatcctctagaggtaccTCACTGTTCGTATGTATTAATGTCG	for cloning into JAB290 cut with NheI/KpnI
JB1055_Cas9-T2A-EGFP_HIKI_T2A_F	tgtggaggagaaccccgggcccgctagcATGGATTACAAGGATCACGATG	for cloning into JAB290 cut with NheI/KpnI
JB1056_Cas9-T2A-EGFP_HIKI_R	caaagatcctctagaggtaccTTAGTACAGCTCGTCCATGCC	for cloning into JAB290 cut with NheI/KpnI
JB1002_FLPo_HIKI_T2A_F	tgtggaggagaaccccgggcccgctagcatgagccagttcgacatcct	for cloning into JAB290 cut with NheI/KpnI
JB1003_FLPo_HIKI_R	caaagatcctctagaggtacctcagatccgcctgttgatgtagc	for cloning into JAB290 cut with NheI/KpnI
JB1004_Gal80_HIKI_T2A_F	tgtggaggagaaccccgggcccgctagcATGGACTACAACAAGAGATCTTCG	for cloning into JAB290 cut with NheI/KpnI
JB1005_Gal80_HIKI_R	caaagatcctctagaggtaccTTATAAACTATAATGCGAGATATTGCTAA	for cloning into JAB290 cut with NheI/KpnI
JB1008_Nluc_HIKI_T2A_F	tgtggaggagaaccccgggcccgctagcATGGTCTTCACACTCGAAGATTT	for cloning into JAB290 cut with NheI/KpnI
JB1009_Nluc_HIKI_R	caaagatcctctagaggtaccTTACGCCAGAATGCGTTC	for cloning into JAB290 cut with NheI/KpnI
JB1032_Gal4DBD_HIKI_T2A_F	tgtggaggagaaccccgggcccgctagcATGCTGGAGATCCGCGCC	for cloning into JAB290 cut with NheI/KpnI
JB1033_Gal4DBD_HIKI_R	caaagatcctctagaggtaccTTACGATACCGTCAGTTGCC	for cloning into JAB290 cut with NheI/KpnI
JB1030_p65_HIKI_T2A_F	tgtggaggagaaccccgggcccgctagcATGGATAAAGCGGAATTAATTCC	for cloning into JAB290 cut with NheI/KpnI
JB1031_p65_HIKI_R	caaagatcctctagaggtaccTTACTTGCCGCCGCCCAG	for cloning into JAB290 cut with NheI/KpnI
JB917_sfGFP_HIKI_F	ggtggaagcggaggtagcgcggccgccGTGTCCAAGGGCGAGGAG	for cloning into JAB290 cut with NotI/KpnI
JB916_sfGFP_HIKI_R	caaagatcctctagaggtaccCTACTTGTACAGCTCATCCATGC	for cloning into JAB290 cut with NotI/KpnI
JB1355_FK506-bp2_geno_1F	acgcgccaaaatacaaaaac	amplification of endogenous target gene and knock-in
JB1356_FK506-bp2_geno_1R	GCCTATTCGACCTTGAGCAG	amplification of endogenous target gene
JB1357_alphaTub84B_geno_1F	tttgtgtgggcaaaattcaa	amplification of endogenous target gene and knock-in
JB1358_alphaTub84B_geno_1R	GCTTGGACTTCTTGCCGTAG	amplification of endogenous target gene
JB1359_esg_geno_1F	CGTTTGGTATTTGTGCATCG	amplification of endogenous target gene and knock-in
JB1360_esg_geno_1R	GTAGGGCGACATGTGGAAGT	amplification of endogenous target gene
JB1361_btl_geno_1F	aactaagggaggggcaaaaa	amplification of endogenous target gene and knock-in
JB1362_btl_geno_1R	CGTCCACCAAGGATTTGAGT	amplification of endogenous target gene
JB1363_Desat1_geno_1F	aatccacctggtgcttgttc	amplification of endogenous target gene and knock-in
JB1364_Desat1_geno_1R	GTAACCGAAGGCGATGATGT	amplification of endogenous target gene
JB1365_ap_geno_1F	ttgcaaatctgtcaggaacg	amplification of endogenous target gene and knock-in
JB1366_ap_geno_1R	ATCTGGACACGAGGATGAGG	amplification of endogenous target gene
JB959_ebony_geno_F	gcattagcctgcattgcata	amplification of endogenous target gene and knock-in
JB960_ebony_geno_R	CACGCCCTCATCGAAATAGT	amplification of endogenous target gene
JB961_wg_geno_F	CAGTTAAGCGTTGGCACTGA	amplification of endogenous target gene and knock-in
JB962_wg_geno_R	ttgttgcatctctgcggtag	amplification of endogenous target gene
JB963_Myo1a_geno_F	TCGTCGTCATCAACAGAAGC	amplification of endogenous target gene and knock-in
JB964_Myo1a_geno_R	tctggagtggaaccgaaaac	amplification of endogenous target gene
JB965_Mhc_geno_F	cggctaaagactgacccaaa	amplification of endogenous target gene and knock-in
JB966_Mhc_geno_R	CTCTTGCTCCATGACGAACA	amplification of endogenous target gene
JB967_hh_geno_F	TCGTACTCGCACTCGAACAC	amplification of endogenous target gene and knock-in
JB1388_hh_geno_6F	aaatcaaagctggaccaaatc	amplification of endogenous target gene and knock-in, used with hh #1 insertion
JB968_hh_geno_R	GTTGTAGTTGGGCACGAGGT	amplification of endogenous target gene
JB900_T2A_R	cggggttctcctccacat	reverse primer for amplifying T2A-Gal4 insert in sense orientation
JB958_T2A-Gal4_3'_F	gttttcccagtcacgacgtt	reverse primer for amplifying T2A-Gal4 insert in antisense orientation
JB659_3P3dsred_seq1F	ACTCCAAGCTGGACATCACC	to amplify 3xP3-dsred as a control
JB660_3P3dsred_seq1R	CGAGGGTTCGAAATCGATAA	to amplify 3xP3-dsred as a control
JB713_Rp49_F	ATCGGTTACGGATCGAACAA	to amplify endogenous Rp49 gene as a control
JB714_Rp49_R	GACAATCTCCTTGCGCTTCT	to amplify endogenous Rp49 gene as a control
JB1051_alphaTub84B_C-term_F	CCTTCGTCCACTGGTACGTT	amplification of endogenous target gene and knock-in
JB1052_Act5c_C-term_F	CGTCGACCATGAAGATCAAG	amplification of endogenous target gene and knock-in
JB1053_His2Av_C-term_F	CTCCTCGCCACTTACAGCTC	amplification of endogenous target gene and knock-in
JB1054_Lam_C-term_F	GCCGACAACACTAGGACGAT	amplification of endogenous target gene and knock-in
JB1050_mNeonGreen_R	GGGAGAGAGGCGTTATCCTC	reverse primer for amplifying gene-mNeonGreen insert
JB1120_act5c_seqnextgenR	CGACTTCTCCTCCTCCTCCT	amplification of endogenous target gene
JB1121_His2av_seqnextgenR	TCGTCGGTGTTTTAGCTTGTC	amplification of endogenous target gene
JB1122_alphatub_seqnexgenR	GCGATTGGAAGCGTAAACAC	amplification of endogenous target gene
JB1123_Lam_seqnextgenR	GTGTTGTGCTGCGTTTGATT	amplification of endogenous target gene
JB1192_Lam_2F	GCGGCTAATCAACGAGAAAG	amplification of non-knock-in Lamin gene
JB1193_Lam_2R	TCTGTTGTCAGGAGCGTTTG	amplification of non-knock-in Lamin gene
JB1194_Lam_3F	ACGAGGAGCAGATGCAGATT	amplification of non-knock-in Lamin gene
JB1195_Lam_3R	GGTCTAAACCGGGAGAAAGC	amplification of non-knock-in Lamin gene
JB1196_Lam_4F	AGCTGCAGAACCTGAACGAT	amplification of non-knock-in Lamin gene
JB1197_Lam_4R	ACTAGCCGAACCCAGGATTT	amplification of non-knock-in Lamin gene
JB1198_Lam_5F	cccattacaagcgacgattt	amplification of non-knock-in Lamin gene
JB1199_Lam_5R	GAACAGCTCCACTCCTCCAG	amplification of non-knock-in Lamin gene
JB1200_Lam_6F	GTCTCGGTCTGCTCCTCATC	amplification of non-knock-in Lamin gene
JB1201_Lam_6R	GAATGGCATAGCCACCACTT	amplification of non-knock-in Lamin gene
JB1202_Lam_7F	CATTTGCAAGATGGTGGTTG	amplification of non-knock-in Lamin gene
JB1203_Lam_7R	ccaattaggccaacactgct	amplification of non-knock-in Lamin gene
JB1050_mNeonGreen_R	GGGAGAGAGGCGTTATCCTC	amplification and sequencing primer, 5' end of mNeonGreen
JB1423_Not1-MCS-BamHI_gBlock	agggtttttcccagtcacgacgttgtaaaacgacggccagtgaattgtaatacgactcactataggggcgaattgggtaca agcttcacgacctgaggcgcgccgggccagtacccaaaaagcggggggtccggtggaagcggaggtagcgcggccgcac GCTAGCtgCTCGAGacGGTACCtaGAATTCacAGATCTataacttcgtataatgtatgctatacgaagttatcgtacgG GATCTAATTCAATTAGAGACTAATTCAATTAGAGCTAATTCAATTAGGATCCAAGCTTATCGATTTCGAACCCTCGACC GCCGGAGTATAAATAGAGGCGCTTCGTCTACGGAGCGACAATTCAATTCAAACAA	gBlock containing multiple cloning site to construct pCRISPaint-3xP3-RFP
JB1426_pCRISPaint_3xP3RFP_backbone_F	gggccgcgactctagatcataatc	to amplify pCRISPaint-3xP3-RFP backbone to swap out RFP sequence
JB1427_pCRISPaint_3xP3RFP_backbone_R	ggtgagcgaccggcttcga	to amplify pCRISPaint-3xP3-RFP backbone to swap out RFP sequence
JB1428_GFP_overlapCRISPaint3xP3RFP_F	gctcgaagccggtcgccaccatggtgtccaagggcgag	to amplify EGFP to Gibson clone into pCRISPaint-3xP3-RFP
JB1429_GFP_overlapCRISPaint3xP3RFP_R	atgatctagagtcgcggccccctacttgtacagctcatccatgc	to amplify EGFP to Gibson clone into pCRISPaint-3xP3-RFP
JB939_PCR1fwrd_frame0_1_2_pCFD5	GCGGCCCGGGTTCGATTCCCGGCCGATGCAgcccagtacccaaaaagcgggGTTTTAGAGCTAGAAATAGCAAG	to construct pCFD5-frame_selector_0,1,2
JB940_PCR1rev_frame0_1_2_pCFD5	ccgcttttttgggtactggccTGCACCAGCCGGGAATCGAACCC	to construct pCFD5-frame_selector_0,1,2
JB941_PCR2fwrd_frame0_1_2_pCFD5	ggccagtacccaaaaagcggGTTTTAGAGCTAGAAATAGCAAG	to construct pCFD5-frame_selector_0,1,2
JB942_PCR2rev_frame0_1_2_pCFD5	ATTTTAACTTGCTATTTCTAGCTCTAAAACcgcttttttgggtactggcccTGCACCAGCCGGGAATCGAACCC	to construct pCFD5-frame_selector_0,1,2
JB1294_ebony_LHA_T2A-Gal4_2F	tgggtacaagcttcacgacctgaggggcacgggtaattgtcgttctt	to amplify ebony left homology arm
JB1295_ebony_LHA_T2A-Gal4_2R	ccccgcttttttgggtactggcccggtGCCTTCCAGATGGCCAGC	to amplify ebony left homology arm
JB975_ebony_RHA_T2A-Gal4_F	atgtatgctatacgaagttatgagctTGGCGGCGCGTATTTGCC	to amplify ebony right homology arm
JB976_ebony_RHA_T2A-Gal4_R	accctcactaaagggaacaaaagctggagctCCTCAGGAACCTGCGAGTAG	to amplify ebony right homology arm
JB1484_ebony_HDRgeno_F	gggcaaacatagtgcaggtt	use with to verify ebony-T2A-Gal4 HDR knock-in
JB1485_ebony_HDRgeno_R	AGACCCTTGGGCAGGTAGTT	to verify ebony-T2A-Gal4 HDR knock-in
JB1422_bc10_CRISPaint_F	TAAACGACCCGAAAATCGAG	amplification of endogenous target gene and knock-in
JB1413_CG42308_CRISPaint_F	AAACACGGCACCTTTGAATC	amplification of endogenous target gene and knock-in
JB1414_CG18336_CRISPaint_F	TCTCCTGTCAGGCCAGCTAT	amplification of endogenous target gene and knock-in
JB1416_CG42381_CRISPaint_F	CAACCGCCAAGAAGAAGAAG	amplification of endogenous target gene and knock-in
JB1417_CG42518_CRISPaint_F	CCACAAGgtgagtccttggt	amplification of endogenous target gene and knock-in
JB1045_SPLNK-GATC-TOP	GATCCCACTAGTGTCGACACCAGTCTCTAATTTTTTTTTTCAAAAAAA	splinkerette oligos for annealing
JB1046_SPLNK-BOT	CGAAGAGTAACCGTTGCTAGGAGAGACCCGTGGCTGAATGAGACTGGTGTCGACACTAGTGG	splinkerette oligos for annealing
JB1047_SPLNK#1	CGAAGAGTAACCGTTGCTAGGAGAGACC	splinkerette round 1 primer
JB1048_SPLNK#2	GTGGCTGAATGAGACTGGTGTCGAC	splinkerette round 2 primer
JB1057_T2A-Gal4_5'#1	agcggagacctttttggtttt	5' splinkerette round 1 primer
JB1058_T2A-Gal4_5'#2	tcggtttttcttttggagcac	5' splinkerette round 2 primer
JB1059_T2A-Gal4_5'Seq	cggggttctcctccacat	5' splinkerette sequencing primer
JB1060_T2A-Gal4_3'#1	gtgctgcaaggcgattaagt	3' splinkerette round 1 primer
JB1061_T2A-Gal4_3'#2	gttttcccagtcacgacgtt	3' splinkerette round 2 primer
JB1062_T2A-Gal4_3'Seq	GTAAAACGACGGCCAG	3' splinkerette sequencing primer

Supplemental Table 7 – Oligo and dsDNA sequences