Neural tube development depends on notochord-derived Sonic hedgehog released into the sclerotome

Nitza Kahane and Chaya Kalcheim[*]

Department of Medical Neurobiology, Institute of Medical Research Israel-Canada (IMRIC) and the Edmond and Lily Safra Center for Brain Sciences (ELSC)

Hebrew University of Jerusalem-Hadassah Medical School

Jerusalem 9112102, P.O.Box 12272, ISRAEL

FAX# 972-2-6757451, Phone # 972-2-6758438

* Corresponding author- Email: kalcheim@cc.huji.ac.il

Short title: Neuro-mesodermal interactions mediated by Sonic hedgehog.

Abstract

Sonic hedgehog (Shh) is necessary both for neural and mesodermal development. By loss and gain of Shh function, and floor plate deletions, we report that Shh released from notochord into sclerotome constitutes a dynamic pool of ligand serving both tissues. Depletion of Shh in sclerotome by membrane-tethered hedgehog-interacting protein or by Patched1, but not by a dominant active version of Patched, reduced motoneuron numbers and also compromised myotome differentiation. These effects were a specific and direct consequence of reducing Shh. In addition, grafting notochords in a basal, but not apical location vis-a-vis the tube, profoundly affected motoneuron development, suggesting that initial ligand presentation occurs at the basal side of epithelia corresponding to the sclerotome-neural tube interface.
Collectively, our results reveal the sclerotome as a previously unknown common pool and potential site of a Shh gradient that coordinates development of mesodermal and neural progenitors.

Introduction

The morphogen Sonic hedgehog (Shh) plays fundamental roles in the development of both neural tube (NT) and somites [1-6]. Its signaling is initiated by binding of the proteolytically processed and lipid modified ligand to the transmembrane receptor Patched (Ptc), that represses the pathway in the absence of ligand [7-10]. Ligand binding to Ptc abrogates its repressive effect on Smoothened (Smo), a key effector essential for canonical Hedgehog signal transduction [11]. The repressive role of Ptc correlates with its localization in the primary apically located cilium, that functions as a signal transduction compartment [12, 13]. Binding of Shh to Ptc removes Smo from the cilium, thereby allowing Smo to enter and propagate the signal further downstream [14, 15] to regulate Gli transcription factor activity [16, 17].

Shh signaling is highly regulated by negative and positive modulators. Ptc1, Hedgehog interacting protein (Hhip) and Gli1 are direct targets of Shh and the former two also inhibit its activity [18, 19]. Sulfatase1 (Sulf1) is a known Shh target [20] and the co-receptors Boc, Gas and Cdo [21, 22] all enhance ligand activities and are expressed in the NT and/or developing mesoderm [23].

It is well established that at early stages following neurulation, Shh secreted by the notochord (No) induces distinct ventral cell identities in the overlying NT by a mechanism that depends on relative concentrations and duration of exposure [24-26]. No-derived Shh is also involved in mesoderm patterning [4, 5]. A ventro-dorsal activity gradient of Shh/Gli signaling in sclerotome was directly visualized using an *in vivo* reporter in mice [23]. In addition, in chick embryos, Shh activity spreads from the midline through the sclerotome to reach the dermomyotome (DM). There it

promotes terminal myogenic differentiation of both epaxial and hypaxial DM-derived progenitors and maintains epitheliality of DM cells [23]. Notably, in both floor plate (FP) and myotome, the activities of Shh are transient and cells become refractory to the ligand, a mechanism that allows dynamic phase transitions to take place within these systems [23, 27].

Because Shh is important for the development of both NT and mesoderm, two functionally interconnected systems, the question arises whether the effects of Shh on either tissue are independent of each other or interrelated. Answering this question is of utmost significance both for better understanding the mechanism of Shh activity and for achieving an integrated molecular view of regional development.

In this study we report that, in addition to affecting muscle development, depletion of Shh in the sclerotome by Hhip1 also significantly reduces motoneuron numbers in NT. Notably, similar effects are monitored when a membrane-tethered version of Hhip1 (Hhip:CD4), unable to be released from cells, is locally electroporated into the sclerotome. The observed phenotypes are a specific and direct consequence of Shh depletion as they are rescued by excess Shh, direct Shh targets such as *Hhip1* and *Gli1* mRNAs are reduced, and its effects are not mediated through other signaling pathways. Most importantly, the effects of Hhip:CD4 are phenocopied by electroporation of the transmembrane receptor Ptch1 but not by constitutively active PTC$^{\Delta loop2}$ which does not recognize the ligand. In addition, by gain and loss of Shh function, and by FP deletions, we show that the sclerotome, but not the NT itself, constitutes a dynamic pool of No-derived Shh that acts both on motoneuron as well as on myotome development. Furthermore, grafting No fragments adjacent to the basal,

sclerotomal side of the NT profoundly affects its development when compared to apical grafts. A similar basal grafting with respect to the DM significantly enhances myotome formation, suggesting a general need for initial ligand presentation at the basal side of epithelia. Together, our results uncover the sclerotome as a hitherto unknown common pool of Shh that promotes development of both mesodermal and neural progenitors.

Results

Reduction of Shh in sclerotome by Hhip1 affects both myotome and motoneuron differentiation

To investigate possible interactions between neural and mesodermal progenitors mediated by Shh, electroporations were performed in avian embryos aged 23-25 somite pairs at the level of epithelial somites prior to myotome development. In this region, the NT is composed of proliferative cells [28] and neural patterning is already apparent and ongoing, as evidenced by expression of the positive Shh targets *Nkx2.2, Olig2, Nkx6.2 and Nkx6.1* (Supplem. Fig. S1, A-D). However, differentiation into Hb9-expressing motoneurons has not yet occurred at this stage (Supplem. Fig. S1, E) and only starts about 10 hr later at the level of somites 11-12 located rostral to the last segmented pair of somites (Supplem. Fig.S1, F, arrows). Hence, the timing of manipulations of Shh activity corresponds to the transition of specified proliferative progenitors into differentiated motoneurons [29].

In a previous study, we reported that Shh traversing the sclerotome is necessary for myotome differentiation, as missexpression of the high affinity and selective Shh antagonist Hhip1 in the sclerotome resulted in smaller myotomes expressing desmin accompanied by a corresponding accumulation of Pax7-positive progenitors [[23] and Fig. 1, A,B]. Here, we report that the hemi-NT facing the transfected mesoderm was also affected, exhibiting a significant reduction in the number of Hb9+ motoneurons when compared to controls that received GFP alone (asterisks in Fig. 1A',B', C, p<0.001, N= 12 and in both control and treated embryos). Moreover, a ventral expansion of the Pax7-positive domain was frequently observed (arrows in Fig.

1A,B), likely due to a reduced size of the ventral extent of the transfected hemi-NT (see Fig.2).

These results can be explained by the sclerotome constituting a common pool of Shh acting on both mesodermal and neural progenitors. Alternatively, they might result from Hhip1 moving towards the NT. Indeed, in spite of the initial findings that Hhip1 is a transmembrane glycoprotein with cell autonomous functions [18, 30], it has recently been reported that Hhip1 is a secreted Shh antagonist able to exert long range effects on Shh signaling [31, 32]. This is compatible with the possibility that sclerotomal Hhip1 crosses into the NT or at least localizes to its surrounding basement membrane [31].

Differential behavior of secreted Hhip1 compared to membrane-tethered Hhip1:CD4 vis-à-vis Shh

To discriminate between the above possibilities, we produced a Hhip:CD4 plasmid encoding for a membrane-tethered version of Hhip1, unable to undergo secretion [31, 32]. First, we asked whether NT or sclerotomal cells missexpressing either Hhip1 or Hhip:CD4 are able to sequester Shh. To this end, we implemented a plasmid encoding the N-terminus of Shh fused in frame to YFP (ShhN:YFP). ShhN:YFP is able to undergo palmitoylation but not addition of a cholesterol moiety, a property that enables free movement of the mutant protein when compared to native Shh [33]. When electroporated to the sclerotome, secreted ShhN:YFP was apparent along the basement membrane of the NT where it co-localized with laminin, yet no fluorescent signal was detected along the neuroepithelial cells (Supplem. Fig.S2, A-C, arrows).

When double electroporations of ShhN:YFP to the NT and Hhip:CD4 to sclerotome or vice-versa were performed,   the Hhip:CD4-transfected sclerotome or NT progenitors, respectively, were decorated with ShhN:YFP, demonstrating that Hhip:CD4 binds and immobilizes the ligand in the expressing cells (Fig. 1D-D",F-F"). In contrast, no such co-localization could be observed in either sclerotome or NT upon   double electroporation of   ShhN:YFP and Hhip1 (Fig. 1, E-E", G-G") consistent with the notion that native Hhip1 is a secreted protein.

In addition, we monitored the expression of endogenous Shh protein upon transfection of Hhip1 or Hhip:CD4 to the sclerotome. Shh immunoreactive protein was evident both intracellularly as well as associated with the cell membranes of the FP and No likely exposed to their external surface (Supplem. Fig. S2, D-F'). Electroporation of control GFP and of Hhip:CD4 had no effect on Shh immunoreactive protein in either the No or FP (Supplem. Fig.S2, D-E'). In contrast, missexpression of Hhip1 markedly reduced Shh levels in both structures unilaterally adjacent to the transfected cells; this reduction was mainly apparent at their basal sides closer to the source of Hhip1 (Supplem. Fig.S2, arrows in F'). This effect may result from Hhip1 masking antibody binding to the ligand as the 5E1 Shh antibody and Hhip1 bind to the same pseudo-active site on the Shh molecule [30, 34]. Together, the above data confirm that Hhip1 is secreted to adsorb Shh far from the producing cells, whereas Hhip:CD4 acts cell-autonomously.

The effects of Hhip:CD4 in the NT resemble those observed with other Shh inhibitors
Next, we employed the NT to examine the specificity of Hhip:CD4 relative to other, known inhibitors of the Shh pathway. Electroporation of Hhip:CD4, like that of Hhip1, Ptc1 or PTC$^{\Delta loop2}$ to hemi-NTs, significantly reduced the number of Hb9+

motoneurons and that of pH3+ mitotic nuclei while enhancing cellular apoptosis. Furthermore, the total area of the transfected hemi-NTs, that reflects overall changes in both proliferation and survival, was significantly smaller in all treatments when compared to control GFP (Supplem. Fig. S3, N=4 for all treatments, *p<0.05, **p<0.03, ***p<0.01). The observed effects on motoneuron numbers could thus result from reduced cell differentiation or, indirectly, from effects on progenitor proliferation or survival. These data confirm that Shh acts both as a mitogen and survival factor in the NT [35, 36]. Most importantly, they demonstrate that Hhip:CD4, which acts like Hhip1, Ptc1 or PTC$^{\Delta loop2}$, is a specific tool to abrogate Shh activity.

Loss of Shh activity in either NT or sclerotome inhibits cell differentiation in the adjacent tissue

Local depletion of Shh activity by Hhip:CD4 or Ptc1 in sclerotome inhibits motoneuron differentiation in the NT

Next, we addressed the question whether the effects originally observed across tissues (e.g; between NT and mesoderm) with secreted Hhip1 (Fig.1), can be mimicked by missexpression of two different Shh inhibitors, Hhip:CD4 and the Shh receptor Ptc1, both membrane associated molecules.

Electroporation of Hhip1 to sclerotome caused significant effects in the NT adjacent to the transfected mesoderm, as expected from a secreted molecule. These included a reduction in the number of Hb9+ motoneurons (Fig. 2, N=8, p<0.01), a decreased number of pH3+ mitotic nuclei (N=4, p<0.03), enhanced apoptosis (N=4) and an overall decrease in the area of the respective hemi-NT (N=4, p<0.03) when compared to controls (N=5, Fig.2 B,B',G,L compared to A,A',F and K, and see quantification in P).

Notably, transfection of both Hhip:CD4 and Ptc1 significantly affected motoneuron numbers (N=4, p<0.03 and N=7, p<0.01, respectively, Fig. 2C,C'D,D',P), yet had a mild but non statistically significant effect on cell proliferation (N=4 and 7), survival or total hemi-NT area (N=4 and 6, respectively) opposite the treated sclerotomes (Fig.2 H,M,I,N,P). Thus, inhibition of Shh activity in mesoderm mainly affects motoneuron differentiation, contrasting with Shh abrogation in the NT where all parameters were considerably compromised (Supplem. Fig.S3). The finding that motoneuron differentiation is more sensitive to a reduced amount of ligand, indicates that progenitor proliferation, survival and motoneuron differentiation are separable processes that depend upon different Shh concentrations.

As a control for Ptc1 activity, we electroporated PTC$^{\Delta loop2}$ that is unable to bind circulating Shh and acts cell autonomously to inhibit its signaling. PTC$^{\Delta loop2}$, like Hhip1, Hhip:CD4 and Ptc1 adversely affected the size of the electroporated sclerotomes when compared to the intact contralateral ones (Fig.2Q, N=5, p<0.01), altogether demonstrating that Shh signaling is necessary for proliferation and/or survival of sclerotomal progenitors. In contrast, PTC$^{\Delta loop2}$ had no significant effect on either proliferation, survival or total area of adjacent NT cells (Fig.2E,E',J,O,P). As expected, unlike Ptc1, PTC$^{\Delta loop2}$ had no effect on motoneurons (Fig.2E,E',P) suggesting that reduced sclerotomal mass is not sufficient to account for the observed loss of motoneurons.

It is worth mentioning that Hhip1 also caused a slight ventralization of Pax7 expression (arrows in B) in association with a reduced size of transfected hemi-NTs. This effect was less apparent or not evident upon electroporation of Hhip:CD4, Ptc1 or PTC$^{\Delta loop2}$, which did not significantly affect hemi-NT size (Fig.2, C,D and E).

The specificity of Hhip:CD4 was further tested by examining *Olig2*, an earlier marker of motoneuron progenitors. The expression domain of *Olig2* was reduced adjacent to the electroporated sclerotome (Supplem. Fig. S4). This indicates that depletion of Shh in sclerotome also affects ongoing specification of motoneuron progenitors, a process that already begun by the time electroporations were performed (Supplem. Fig.S1).

In addition, while sclerotomal missexpression of Hhip:CD4 significantly affected the number of Hb9+ cells, Shh alone moderately but not significantly (p=0.056), enhanced their differentiation, and co-treatment of Hhip:CD4 with Shh rescued the effect of Hhip:CD4 back to control levels (Fig. 3, N=12, 26, 8 and 7, for control GFP, Hhip:CD4, Hhip:CD4+Shh and Shh alone, respectively, p<0.001).

Local depletion of Shh activity by Hhip:CD4 or Ptc1 in NT inhibits myotome differentiation in mesoderm

Next, we examined whether depletion of Shh in the NT influences myotomal size. Electroporation of control GFP to hemi-NTs had no effect of myotome size (N=14) whereas Hhip1 and Hhip:CD4 caused a significant decrease in the size of the desmin+ myotomes adjacent to the transfected side of the neuroepithelium (N=3,10, respectively, p<0.001, Fig. 4, A-C, F). Electroporation of Ptc1 exhibited a similar effect (N=16, p<0.001); in contrast, $PTC^{\Delta loop2}$ revealed no reduction in myotome size (N=6, Fig. 4D,E,F).

Together, our data suggest that depletion of Shh in the sclerotome promotes a concomitant loss of effective Shh in the NT and vice-versa. Since Hhip:CD4 and Ptc1 are membrane-bound and not secreted, the present results could be explained by the existence of a common pool that supplies Shh to both tissues.

The effects of Hhip:CD4 are a direct consequence of Shh depletion

The similarity between the effects of Hhip:CD4 and Ptc1 (Fig. 2) and the rescue of motoneuron numbers by co-electroporation of Shh along with Hhip:CD4 (Fig. 3) suggests that the effects of Hhip:CD4 are specifically mediated by ligand depletion. To further investigate the possibility of direct versus indirect effects, we examined the expression of *Gli1* and *Hhip1*, two transcriptional targets of Shh and compared it to *Gli3* mRNA expression, which is not directly regulated by Shh [37]. Electroporation of control GFP or of Hhip:CD4 to sclerotome or NT had no effect on expression of *Gli3* mRNA in the same or adjacent tissue (Supplem. Fig. S5, A-C). In contrast, similar transfections of Hhip:CD4 to NT or sclerotome reduced *Gli1* and *Hhip1* mRNAs in both the transfected and in adjacent tissues when compared to the respective contralateral sides (Supplem. Fig. S5, D-I). These results further support the notion that the observed effects on motoneuron and myotome development specifically and directly result from ligand depletion.

Next, we examined the possibility of secondary effects across tissues mediated by Shh depletion. Two candidates are the BMP pathway which acts antagonistically to Shh [1, 38, 39] and retinoic acid from the somite that affects NT development [40, 41]. To this end, control GFP or Hhip:CD4 were electroporated into the sclerotome. If the effects on the NT of Shh deprivation in sclerotome are mediated by BMP, it is predicted that the extent of BMP signaling is expanded. No such change of pSmad 1,5,8 immunoreactivity, a readout of BMP activity, was measured upon Hhip:CD4 treatment (Supplem. Fig.S6, N=4 per treatment). As an internal control for Hhip:CD4

activity, we observed a reduced number of Hb9+ motoneurons in the ventral NT adjacent to the transfected side (Supplem. Fig.S6B, arrow).

In addition, control GFP or Hhip:CD4 were electroporated into the sclerotome or NT and RARE-AP, a specific reporter of retinoic acid activity was transfected into the adjacent tissue, respectively. The specificity of RARE-AP was first tested by co-electroporating it with a dominant negative receptor plasmid, that abolished RARE-AP signal (Supplem. Fig. S7A-B").   No apparent changes in RARE-AP were observed in either tissue upon depletion of Shh in the neighboring one when compared to control GFP (N= 4 for each treatment, Supplem. Fig. S7, C-F).

Taken together, these results show that the observed effects are a direct consequence of ligand depletion. Thus, inhibition of Shh in the sclerotome by Hip:CD4 or Ptc1 results in a corresponding reduction of Shh ligand in the NT and vice-versa.

Gain of Shh function in sclerotome enhances motoneuron differentiation but Shh missexpression in NT has no effect on myotome

Our loss of function results would be consistent with the possibility that Shh can translocate bidirectionally between mesoderm and NT. To gain additional insight into the directionality of Shh effects across NT and mesoderm, we adopted a complementary gain of function approach. Control GFP or full length Shh were electroporated into sclerotome. A day later, an increase of approximately 40% in the number of Hb9+ motoneurons was monitored in the NT of Shh-treated embryos ipsilateral to the treated side when compared to controls (p<0.001, N=9 for each

treatment, Fig. 5A-C). In addition, desmin+ myotomes were enlarged at the expense of Pax+ DM progenitors (Fig. 5B).

Reciprocally, control GFP or full length Shh were electroporated into hemi-NTs and the size of adjacent desmin+ myotomes was examined. Although the overall size of the transfected hemi-NT increased, no significant change in myotomal size was measured and no apparent effects on DM or sclerotome were observed (N=6 and 5 in control and treated embryos, respectively, Fig. 5D-F).

Hence, our gain of function data are inconsistent with the simpler possibility that Shh moves bidirectionally between mesoderm and NT, that is based solely on data from loss of Shh activity. While excess Shh in mesoderm profoundly affects NT development, the observation that gain of Shh in NT has no effect on myotome development is in line with Shh being transported into the neuroepithelium but not outside into mesoderm. In this regard, our finding that Shh depletion in the NT inhibits myotome differentiation (Fig. 4) is consistent with this procedure causing a corresponding lower effective concentration in mesoderm. This could be accounted for by enhanced uptake of the ligand into NT cells via directional baso-apical transport from the sclerotome. This suggests that the sclerotome constitutes a common pool of No-derived Shh that serves both tissues.

Sclerotome, but not FP-derived Shh is necessary for motoneuron differentiation

Classical data support the view that the development of various cell types in the NT depends upon a local ventrodorsal gradient of Shh emanating from the No and FP [42, 43]. Our present results, show that No-derived Shh secreted into the sclerotome is also

necessary for motoneuron development (Figs. 4,5). Hence, we examined the relative contribution of the FP compared to sclerotomal Shh to the differentiation of Hb9+ neurons.

Control GFP or Hhip:CD4 were electroporated dorsoventrally to attain the FP of the NT. Control GFP had no effect on FP integrity or on expression of Shh protein (N= 7, Fig.6A). Surprisingly, electroporation of Hhip:CD4 caused the total disintegration of the FP and consequent loss of FP-derived Shh (N= 10, Fig. 6, D,E); an effect that enabled us to accurately monitor the contribution of the FP to motoneuron development. Because the effect is bilateral, in order to prevent variability between embryos and obtain a reliable measurement of motoneuron numbers, the proportion of Hb9+ neurons was measured as the ratio between the neurons present at the flank level lacking a FP to the intact neck level of the same embryos. In spite of FP disappearance, the proportion of flank-level motoneurons was unaltered when compared to control embryos that received GFP only (N=4 for each treatment, Fig. 6, B,E,H). Similar to what was observed with Hhip:CD4, dorsoventral electroporation of Ptc1 also compromised FP integrity while having no apparent effect on motoneurons (N=5, Fig.6G). In contrast, inhibition of Shh in the sclerotome by Hhip:CD4 exhibited a visible reduction in ventral motoneurons (Fig. 6, C,F arrow, and see also Figs. 2 and 3).

Sclerotomal Shh is not likely to act by affecting ligand levels in the producing cells because inhibition of Shh by Hhip:CD4 in sclerotome had no effect on Shh expression in either FP or No (Supplem. Fig. 2, E,E'). Thus, we conclude that Shh traversing the sclerotome plays a significant part in motoneuron differentiation whereas the FP itself

has no apparent contribution at least at the stages examined in our study (see Discussion).

<u>A basal, but not apical, presentation of Shh is required for ligand activity on both NT and DM/myotome</u>

Based on our finding that Shh transiting through the sclerotome is needed for NT development, we predicted that the NT would be more sensitive to Shh presented from its basal side abutting the sclerotome than from its apical side. To examine this hypothesis, fragments of No were grafted into either the lumen of the NT (apical) or between somites and NT (basal grafts). A day later, the expression of Pax7 in the dorsal NT was not affected by the luminal grafts yet was dorsally restricted when facing the basal grafts (Fig. 7A,B). The latter also caused a characteristic bending of the NT, previously reported to represent an ectopic FP-like structure [44](N= 6 and 6 for apical vs. basal grafts, Fig. 7, B,D). In addition, the amount of Hb9+ motoneurons was unchanged by the apical grafts, yet was markedly increased in the basal grafts in which the No's were similarly localized in a dorsal position vis-vis the NT (N= 6 and 6 for apical vs. basal grafts, Fig. 7, C,D), in agreement with classical No graft experiments [45].

Likewise, implantation of No fragments at epithelial somite levels and at an apical position with respect to the DM had only a mild effect on the subsequent development of desmin-positive myotomes with no apparent alteration in Pax7 expression in the DM. In striking contrast, equivalent grafts performed basal to the prospective DM, produced large myotomes expressing desmin and a radical *in situ* differentiation of the DM into muscle at the expense of Pax7-positive progenitors (N= 5 and 7 for

apical vs. basal grafts, Fig. 7, E,F). Therefore, an initial basal presentation of the ligand vis-à-vis the target epithelium is required for the activity of Shh. This is consistent with our results showing that Shh emanating from the sclerotome and reaching the NT from its basal side, is necessary and sufficient for aspects of NT differentiation.

Discussion

In the present study we uncover a previously unknown domain, the sclerotome, as being an important "en passant" pool of Shh that influences not only DM and myotome development, as previously shown [23] but also aspects of NT differentiation (Fig. 7G). Our loss of function data show that local depletion of Shh in either NT or sclerotome results in major defects across tissues such as reduced myotomal size and less motoneurons, respectively. Reciprocally, only gain of Shh function in the sclerotome significantly enhances motoneuron differentiation while missexpression of the ligand in NT has no effect on myotomal size. Taken together, these results suggest that there is a common pool of Shh ligand present in sclerotome that supplies both tissues.

In contrast to the reduction in motoneurons observed upon inhibition of Shh in sclerotome, we show that Hhip:CD4-mediated ablation of the FP has no short-term effects on motoneuron numbers. This is consistent with the development of a normal ventral pattern in Gli2 mutants that lack a FP [46, 47]. Likewise, loss of Shh in FP did not dramatically alter ventral neural patterning, yet altered gliogenesis at a later stage [48]. Furthermore, abrogating Shh in FP under the regulation of Brn4, revealed a normal short-term expression of both *Nkx2.2* and *Olig2*, but a reduction at later stages suggesting a continuous need for FP-derived Shh in development of the ventral NT [24]. This initial phenotype is consistent with our results yet we did not analyze later stages. Moreover, it is possible that in the absence of a FP, the No which although separated from the NT is still closely adjacent to it, compensates for the loss of FP-derived Shh as suggested for Gli2 mutants [reviewed in [49]]. In comparison, abrogation of Shh ligand in mesoderm, performed at the same stage and for a similar

duration, revealed a significant NT phenotype even in the presence of both Shh-producing axial structures. Taken together, we propose that a major fraction of Shh operating on the neuroepithelium stems from the sclerotome which, at the stages examined, seems more active that the FP in promoting motoneuron development.

Furthermore, how can Shh from sclerotome be more important than Shh from FP, given that the ligand is readily detected in FP but not in mesoderm by immunostaining? Shh in the synthesizing cells of the FP and No is intracellular and membrane-bound [[46] and Supplem. Fig. S2)] whereas in the sclerotome it is expected to be extracellular and/or included in organelles (e.g, exosomes, etc). Most protocols used for tissue processing are likely to keep only the former type of immunoreactive protein and wash away the extracellular ligand in sclerotome. Notably, when using a method that allowed proteoglycan/glycosaminoglycan preservation and/or perhaps also a different antibody, a previous study showed a sclerotomal localization of Shh immunoreactive protein [50].

It is noteworthy that different procedures that perturb the production of Shh protein or its signaling have different effects on FP integrity. Whereas Gli2 mutants lack a FP [51], deletion of Shh in FP does not appear to be necessary for maintenance of this structure [24]. In both chick and mouse, Shh was suggested to be necessary for initial FP induction but later on, during somitogenesis, the FP becomes refractory to the ligand [52]. In our experiments, we implemented a membrane tethered version of the high affinity Shh inhibitor Hhip and also the Shh receptor Ptc1, both resulting in the death of FP cells. This might be accounted for by a combination of ligand depletion

with accumulation of Shh-Hhip or Shh-Ptc1 complexes at the cell membrane altogether compromising the structural integrity of this epithelium.

One open question stemming from our results is how is Shh transported through the sclerotome. Possible models could be packaging of the ligand in No-derived exosomes [53], diffusion of Shh released by matrix metalloproteinases in a lipid-free form [54]; secretion of Shh as multimeric complexes of various molecular compositions [see for example [55]] and/or via carrier-mediated transport through the extracellular space [56]. The precise mechanism responsible for Shh transport in the present context remains to be unraveled.

If provided to the NT from the sclerotomal domain, it is inferred that neurepithelial cells sense Shh from their basal pole that faces the mesoderm. Consistent with this notion, grafting No fragments in a basal, but not apical position with respect to the NT, profoundly affects NT shape, motoneuron differentiation and Pax7 expression. Indeed, during normal development, the No underlies the basal domain of the NT, further supporting the idea that No-derived Shh can only reach the NT via a basal route. In line with the above, it was reported that lipidated Shh enters the cells of the imaginal disc in Drosophila only through its basolateral surface [57]. A similar phenomenon was observed in high density human gastruloids, that self organize into an epithelium. In these cultures, cells were responsive to BMP4 or Activin ligands only when presented from the basal side and this correlated with the localization of BMP receptors at the basolateral domain of the cells [58]. Although it is unclear if Shh receptors are also localized at the basal aspect of neuroepithelial cells, or if Shh

needs first to be transported baso-apically in order to act on apically localized receptors, this and our results suggest that cell polarization controls ligand response.

The above findings are interesting in light of the proposed concept that apically-localized cilia serve as antennae to sense and transduce a Shh signal [59, 60]. Based on our data, we suggest that, initially, Shh is presented from the basal side of epithelial cells from which it may be transported to the apical domain where cilia are localized. Since an apical presentation of the No and associated Shh was without a significant effect in our implant experiments, we propose that cilia act primarily as transducers of a Shh signal, but not as the primary antennae sensing the presence of the ligand. Moreover, the observation that a direct apical presentation of Shh is without effect, further suggests that basal reception followed by baso-apical transport might be necessary for the activity of Shh arriving at the cilia. Along this line, growing evidence suggests cilia-independent Shh reception that occurs through basally localized cytonemes [reviewed in [61] and refs therein]. Likewise, in the retina neuroepithelium, Shh and its coreceptor Cdo colocalize at the basal side of the cells where filopodia-like structures are present [62]. An extreme example is provided by some cell types where Shh signaling takes place even in the absence of cilia [reviewed in [61] and refs. therein].

In the NT, direct visualization of the behavior of labeled Shh, revealed that the ligand from the No concentrates in association with the apically localized basal bodies from which cilia stem, while forming a dynamic gradient in the ventral NT [46]. In light of the present results, the above observed graded distribution of ligand could be explained as being the end point of a transport process that begins at the No, travels

through the sclerotome forming there a ventro-dorsal gradient, then reaches the neuroepithelium through its basal side finally concentrating in the apical cilia (Fig. 7G). In such a model, the mechanisms mediating the baso-apical transport of the ligand associated with possible biochemical changes of the protein that make it availabile to cilia for signaling remain to be unraveled. A microtubular network spanning the extent of neuroepithelial cells could be involved in this process, as previously suggested [46].

An initial presentation of Shh from the basal side of an epithelium seems to be of general significance as basal grafting of a No also elicited robust *in situ* differentiation of DM progenitors into myocytes when compared to an apical implant that exhibited only a mild phenotype. How can this differential effect be explained given that the endogenous ligand apparently arrives from the apical sclerotomal direction?. Initially, pioneer myotomal cells in the epithelial somite face the No from their basal aspect [63]. With ongoing development and upon sclerotome dissociation, the epithelial DM is consolidated and becomes composed of a central sheet and four inwardly curved lips pointing towards the sclerotome. In previous studies, we and others showed that the four lips of the DM are the main sources of myotomal cells [[64] and refs therein] and it is their basal domain that points towards the sclerotome from which Shh arrives. Next, myotomal progenitors enter the nascent myotome and a basement membrane begins assembling at the interface between myotome and sclerotome [65]. At this time, the central DM sheet also contributes myotomal progenitors by direct translocation into the myotome [66], which then differentiate in a Shh-dependent manner [23]. These precursors were shown to exhibit an inverse apicobasal polarity

when compared to the central DM cells [66]. Hence, both the DM lips and these translocating progenitors point with their basal surfaces towards the source of endogenous Shh, likely being the main targets for its activity and also accounting for the partial effect of the apical grafts.

As discussed above, the basal domain of the DM/myotome and NT epithelia are characterized by the presence of a surrounding basement membrane. An association between Shh and the basal lamina has been shown. For instance, in cerebellar granule cell precursors, the laminin-containing basement membrane binds and locally enhances Shh signaling [67]. Similarly, in the mouse somite, Shh induces the activation of Myf5 in DM. Myf5+ cells then translocate to the myotome and upregulate α6β1 integrin and dystroglycan, allowing a myotomal basement membrane to be assembled using primarily laminin α1 produced by sclerotomal cells, and laminin α5 produced by the dorsomedial lip of the DM [reviewed in [65]]. This is further confirmed in Shh-deficient mice, which fail to form a myotomal basement membrane, and in which myotomal cells do not properly exit the cell cycle, maintain Pax3 expression and delay the differentiation program [68, 69]. Additionally, Shh immunoreactive protein was found to localize in the basement membrane surrounding the NT [50]. Taken together, these results suggest that in both NT and DM/myotome, a feedforward mechanism may exist whereby Shh controls laminin expression and the assembly of a basement membrane. This could allow a local concentration of the ligand and/or signal stabilization. However, the basement membrane alone is unlikely to serve as the common pool, as much of the ligand is present in the mesoderm itself (our data and see also [50]).

Our present findings provide an additional argument in support of the importance of NT-somite interactions which are pivotal for the normal patterning of trunk components. For instance, opposite gradients of retinoic acid and Fgf8 in mesoderm are required for early NT development [40]. In addition, the nascent DM controls the timing of neural crest delamination by modulating noggin mRNA and BMP activity levels in the NT [70]. Reciprocally, Bmp4 and/or Wnt1 from the dorsal NT pattern the somite-derived DM and affect myogenesis [71, 72]. Formation of the dorsal dermis from the DM is influenced by NT-derived Wnt1 and by neurotrophin-3 [71, 73, 74]. Furthermore, interactions between neural and somitic cells control neural crest migration and segmentation of peripheral ganglia and nerves as well as specific aspects of myogenesis [75, 76]. The present results raise the intriguing possibility that the dual activity on both motoneurons and myotome of Shh released into sclerotome, serves to couple and coordinate development of the neuromuscular system.

Materials and Methods

Embryos

Chick (Gallus gallus) and quail (Coturnix japonica) eggs were from commercial sources (Moshav Orot and Moshav Mata, respectively).

Expression vectors and electroporation

Expression vectors were: pCAGGS-GFP, pCAGGS-RFP [77], Ptc1 [78], PTC$^{\Delta loop2}$ [23, 78], a retinoic acid reporter fused to alkaline phosphatase (pRARE-AP, from J. Sen) [79], a dominant negative pan-retinoic acid receptor (RAR403dn-IRES-GFP, from S. Sockanathan) that abrogates retinoic acid signaling [80], full length Shh [23], cholesterol deficient Shh (mShh-N-YFP, from V. Wallace) [33] that was subcloned into pCAGGS, and Hhip1 [23]. To produce Hhip1:CD4, a membrane-tethered version of Hhip1, the transmembrane and intracellular domains of mouse CD4 were fused to the C-terminal domain of Hhip1 lacking amino acids A679-V700, as previously described [31, 32] and further subcloned into pCAGGS for electroporation.

For electroporations, DNA (1-4 μg/μl) was microinjected into the center of flank-level epithelial somites (somites 20-25) of 23-25 somite-stage embryos. Electroporations were performed to the ventral half of epithelial somites (prospective sclerotome). To this end, the positive tungsten electrode was placed under the blastoderm in a location corresponding to the ventro-medial portion of epithelial somites on a length of about 7 segments, and the negative electrode was placed in a superficial, dorso-lateral position with respect to the same somites. [23, 66, 81, 82].

For hemi-NT electroporations, DNA was microinjected into the lumen of the NT. One tungsten electrode was placed underneath the blastoderm on one side of the embryo

and the other electrode was placed in a superficial position at the contralateral side. For FP electroporations, the positive electrode was inserted under the blastoderm near the midline and the negative electrode was placed over the dorsal NT. In some cases double electroporations to the hemi NT and sclerotome were performed sequentially. A square wave electroporator (ECM 830, BTX, Inc.) was used. One pulse x12V for 5msec was applied.

No grafts

No fragments comprising a length of 7-8 segments were enzymatically excised from donor embryos aged 25 somite pairs as previously described [35] and kept in cold phosphate buffered saline until grafting. For grafting at the apical side of the NT, the ectoderm and dorsal NT of host embryos were cut and the No piece placed in the NT lumen. A day following implantation, the grafts were usually found in the dorsal portion of the NT facing its cavity. For basal grafting with respect to the NT, a slit was performed between somites and NT and the No fragments were inserted. A similar, but more profound slit, was performed to reach the ventral sclerotome abutting the apical side of the DM. To reach the basal domain of the DM, the ectoderm was cut to precisely accommodate the length of the No fragment. Following microsurgery, embryos were reincubated for additional 24 hours.

Immunohistochemistry

Embryos were fixed overnight at 4°C with 4% formaldehyde in phosphate-buffered saline (PBS) (pH 7.4) followed by washings in PBS. Most immunostainings except for desmin were performed on whole embryo fragments. Immunolabeling for desmin was performed on tissue sections, as described [83, 84].

For wholemount immunostaining, antibodies were diluted in PBS containing 1% Triton X-100 and 5% newborn calf serum and tissues were incubated overnight at 4°C on a rotatory shaker. Next, they were washed twice in a large volume of PBS/1% Triton X-100 first for 10 min and then for 2 hours at room temperature. Secondary antibodies were similarly diluted in PBS/ 1% Triton X-100/ 5% newborn calf serum and incubated overnight followed by repetitive washings. Embryo fragments were dehydrated in increasing ethanol solutions (30%,70%, 90% and 100%, 10 minutes each) followed by toluene (2 times, 10 minutes each), then embedded in paraffin wax and sectioned at 8μm. Paraffin was removed in Xylene and slides were rehydrated in decreasing ethanol solutions.

The following antibodies were used: rabbit anti GFP (1:2000, Invitrogen, Thermo Fisher Scientific, A6455) and mouse anti-desmin (1:200, Molecular Probes, 10519). Monoclonal antibodies against Pax7 (PAX7-s, 1:20), Shh (5E1, 1:20) and Hb9 (1:200) were from DSHB, University of Iowa). Phosphorylated Smad 1-5-8 (PSmad, 1:1000) was a gift from Ed Laufer. Anti-Histone H3 (phospho S10) was from Abcam (mAbcam 14955, 1:500). Detection of DNA fragmentation was done by TUNEL (ab66110, Abcam) according to manufacturer's instructions. Nuclei were visualized with Hoechst.

In situ hybridization

Embryos were fixed in Fornoy (60% ethanol, 30% formaldehyde, 10% acetic acid), then dehydrated in ethanol/toluene, processed for paraffin wax embedding and sectioned at 10 μm. Slides were rehydrated in toluene/ethanol/PBS, treated with proteinase K (1μg/ml, Sigma Aldrich P2308) at 37°C for 7 minutes, and then fixed in 4% formaldehyde at room temperature for 20 minutes. Next, slides were washed in PBS followed by 2X SSC and hybridized in hybridization buffer (1X salt solution composed

of 2M NaCl, 0.12M Tris, 0.04M NaH_2PO_42H2O, 0.05M Na_2HPO_4, 0.05M EDTA, pH7.5], 50% formamide, 10% dextran sulfate, 1mg/ml Yeast RNA, 1X Denhardt solution) containing 1μg/ml DIG labeled RNA probes (prepared with a DIG RNA labeling mix, Roche, 11277073910) for overnight at 65°C in a humid chamber. Post-hybridization, slides were rinsed in a rotating incubator with 50% formamide, 1X SSC, 0.1% Tween 20, until coverslips dropped and then an additional wash for 1 hour followed by 2 washes in MABT (10% Maleic acid 1M pH 7.5, 3% NaCl 5M, 0.1% Tween 20) and preincubation in MABT/ 2.5% FCS. Anti-DIG-AP antibody (1/1000, Roche 11093274910) diluted in MABT+2% BBR+20% FCS was then added for overnight at room temperature. This was followed by rinsing in MABT and then in NTMT (2% NaCl 5M, 10% Tris HCl 1M pH9.5, 5% MgCl2 1M, 0.1% Tween20), and then incubation in NTMT + 1:200 NBT/BCIP Stock Solution (Sigma-Aldrich, 11681451001) at 37°C until the AP reaction was completed.

The following probes were employed: *Hhip1* (from J. Briscoe) , *Nkx*2.2, *Nkx* 6.1, *Nkx* 6.2, and *Olig2* (from J. Ericson), and *Gli1*, *Gli3* from A.G. Borycki.

Data analysis and statistics

Four to 26 embryos were analyzed per experimental treatment. The number of Hb9-positive motoneurons or phospho-histone H3 (pH3) + nuclei was counted in 5-10 alternate sections per embryo.

Myotomes were defined by desmin staining. The area occupied by desmin+ myotomes was measured in alternate sections of 3 to 16 embryos per experimental treatment. Sclerotomes of 4 to 6 embryos were defined in the mediolateral aspect as the tissue between myotome and NT, and in the dorsoventral extent as the mesenchyme between the dorsomedial lip of the DM up to the dorsal border of the

cardinal vein and aorta. The surface area of hemi-NTs was monitored in 4 sections per embryo. Myotomal, sclerotomal and hemi-NT areas were measured using Image J software (NIH) and expressed as the mean ratio between treated and control sides±SEM. All results are expressed as the mean proportion of positive cells or area in treated compared to control contralateral sides±SEM.

Images were photographed using a DP73 (Olympus) cooled CCD digital camera mounted on a BX51 microscope (Olympus) with Uplan FL-N 20x/0.5 and 40x/0.75 dry objectives (Olympus). For figure preparation, images were exported into Photoshop CS6 (Adobe). If necessary, the levels of brightness and contrast were adjusted to the entire image and images were cropped without color correction adjustments or γ adjustments. Final figures were prepared using Photoshop CS6.

Significance of results was determined using the non-parametric Mann–Whitney test. All tests applied were two-tailed, and a P-value of 0.05 or less was considered statistically significant. Data were analyzed using the IBM SPSS software version 25. The number of embryos analyzed for each treatment (N) is detailed in the Results Section. P-values can be found both in the Results Section and in the corresponding Legends.

Acknowledgements

We thank Mordechai Applebaum and Dina Rekler for help with figure preparation and Tali Bdolach for assistance with statistics. We thank James Briscoe and Johan Ericson for helpful comments during the course of this study, and Joel Yisraeli for critical reading of the manuscript. This work was supported by the Israel Science Foundation (#97/13) to CK.

Author Contributions- NK and CK conceived the project and designed the experiments; NK performed the experiments; CK and NK wrote the paper.

Declaration of Interests- The authors declare no competing interests.

References

1.   Ericson J, Briscoe J, Rashbass P, Van Heyningen V, Jessell TM. Graded sonic hedgehog signaling and the specification of cell fate in the ventral neural tube. Cold Spring Harbor SympQuantBiol. 1997;62:451-66.

2.   Ericson J, Rashbass P, Schedl A, Brenner-Morton S, Kawakami A, Van Heyningen V, et al. Pax6 controls progenitor cell identity and neuronal fate in response to graded shh signaling. Cell. 1997;90:169-80.

3.   Briscoe J. Making a grade: Sonic Hedgehog signalling and the control of neural cell fate. EMBO J. 2009;28(5):457-65. Epub 2009/02/07. doi: emboj200912 [pii]10.1038/emboj.2009.12. PubMed PMID: 19197245; PubMed Central PMCID: PMC2647768.

4.   Borycki AG, Mendham L, Emerson CP, Jr. Control of somite patterning by Sonic hedgehog and its downstream signal response genes. Development. 1998;125(4):777-90. Epub 1998/04/04. PubMed PMID: 9435297.

5.   Cairns DM, Sato ME, Lee PG, Lassar AB, Zeng L. A gradient of Shh establishes mutually repressing somitic cell fates induced by Nkx3.2 and Pax3. Dev Biol. 2008;323:152-65. Epub 2008/09/18. doi: S0012-1606(08)01154-8 [pii]10.1016/j.ydbio.2008.08.024. PubMed PMID: 18796301.

6.   Gustafsson MK, Pan H, Pinney DF, Liu YL, Lewandowski A, Epstein DJ, et al. *Myf5* is a direct target of long-range Shh signaling and Gli regulation for muscle specification. Genes Dev. 2002;16:114-26.

7.   Goodrich LV, Milenkovic L, Higgins KM, Scott MP. Altered neural cell fates and medulloblastoma in mouse patched mutants. Science. 1997;277:1109-13.

8.   Hidalgo A, Ingham P. Cell patterning in the Drosophila segment: spatial regulation of the segment polarity gene patched. Development. 1990;110(1):291-301. PubMed PMID: 2081466.

9.   Ingham PW, Taylor AM, Nakano Y. Role of the Drosophila patched gene in positional signalling. Nature. 1991;353(6340):184-7. doi: 10.1038/353184a0. PubMed PMID: 1653906.

10.   Johnson RL, Rothman AL, Xie J, Goodrich LV, Bare JW, Bonifas JM, et al. Human homolog of patched, a candidate gene for the basal cell nevus syndrome. Science. 1996;272:1668-71.

11.   van den Heuvel M, Ingham PW. smoothened encodes a receptor-like serpentine protein required for hedgehog signalling. Nature. 1996;382(6591):547-51. doi: 10.1038/382547a0. PubMed PMID: 8700230.

12.   Rohatgi R, Milenkovic L, Scott MP. Patched1 regulates hedgehog signaling at the primary cilium. Science. 2007;317(5836):372-6. doi: 10.1126/science.1139740. PubMed PMID: 17641202.

13.   Caspary T, Larkins CE, Anderson KV. The graded response to Sonic Hedgehog depends on cilia architecture. Dev Cell. 2007;12(5):767-78. doi: 10.1016/j.devcel.2007.03.004. PubMed PMID: 17488627.

14.   Rohatgi R, Milenkovic L, Corcoran RB, Scott MP. Hedgehog signal transduction by Smoothened: pharmacologic evidence for a 2-step activation process. Proc Natl Acad Sci U S A. 2009;106(9):3196-201. doi: 10.1073/pnas.0813373106. PubMed PMID: 19218434; PubMed Central PMCID: PMCPMC2642660.

15.   Milenkovic L, Scott MP, Rohatgi R. Lateral transport of Smoothened from the plasma membrane to the membrane of the cilium. J Cell Biol. 2009;187(3):365-74. doi: 10.1083/jcb.200907126. PubMed PMID: 19948480; PubMed Central PMCID: PMCPMC2779247.

16.   Ribes V, Briscoe J. Establishing and interpreting graded Sonic Hedgehog signaling during vertebrate neural tube patterning: the role of negative feedback. Cold Spring Harb Perspect Biol. 2009;1:1-16. Epub 2010/01/13. doi: 10.1101/cshperspect.a002014. PubMed PMID: 20066087; PubMed Central PMCID: PMC2742090.

17.   Briscoe J, Therond PP. The mechanisms of Hedgehog signalling and its roles in development and disease. Nat Rev Mol Cell Biol. 2013;14(7):416-29. doi: 10.1038/nrm3598. PubMed PMID: 23719536.

18.   Chuang PT, McMahon AP. Vertebrate Hedgehog signalling modulated by induction of a Hedgehog-binding protein. Nature. 1999;397:617-21.

19.   Ingham PW, McMahon AP. Hedgehog signaling in animal development: paradigms and principles. Genes Dev. 2001;15:3059-87. Epub 2001/12/04. doi: 10.1101/gad.938601. PubMed PMID: 11731473.

20.   Dhoot GK, Gustafsson MK, Ai X, Sun W, Standiford DM, Emerson CP, Jr. Regulation of Wnt signaling and embryo patterning by an extracellular sulfatase. Science 2001;293:1663-166.

21.   Allen BL, Song JY, Izzi L, Althaus IW, Kang JS, Charron F, et al. Overlapping roles and collective requirement for the coreceptors GAS1, CDO, and BOC in SHH pathway function. Dev Cell. 2011;20:775-87. Epub 2011/06/15. doi: S1534-5807(11)00172-9 [pii]10.1016/j.devcel.2011.04.018. PubMed PMID: 21664576; PubMed Central PMCID: PMC3121104.

22.   Izzi L, Levesque M, Morin S, Laniel D, Wilkes BC, Mille F, et al. Boc and Gas1 each form distinct Shh receptor complexes with Ptch1 and are required for Shh-mediated cell proliferation. Dev Cell. 2011;20:788-801. Epub 2011/06/15. doi: S1534-5807(11)00171-7 [pii]10.1016/j.devcel.2011.04.017. PubMed PMID: 21664577.

23.   Kahane N, Ribes V, Kicheva A, Briscoe J, Kalcheim C. The transition from differentiation to growth during dermomyotome-derived myogenesis depends on temporally restricted hedgehog signaling. Development. 2013;140:1740-50. doi: 140/8/1740 [pii]10.1242/dev.092726. PubMed PMID: 23533174.

24.   Dessaud E, Ribes V, Balaskas N, Yang LL, Pierani A, Kicheva A, et al. Dynamic assignment and maintenance of positional identity in the ventral neural tube by the morphogen sonic hedgehog. PLoS Biol. 2010;8:e1000382. doi: 10.1371/journal.pbio.1000382. PubMed PMID: 20532235; PubMed Central PMCID: PMC2879390.

25.   Briscoe J, Small S. Morphogen rules: design principles of gradient-mediated embryo patterning. Development. 2015;142(23):3996-4009. doi: 10.1242/dev.129452. PubMed PMID: 26628090; PubMed Central PMCID: PMCPMC4712844.

26.   Stamataki D, Ulloa F, Tsoni SV, Mynett A, Briscoe J. A gradient of Gli activity mediates graded Sonic Hedgehog signaling in the neural tube. Genes Dev. 2005;19:626-41. Epub 2005/03/03. doi: 19/5/626 [pii]10.1101/gad.325905. PubMed PMID: 15741323; PubMed Central PMCID: PMC551582.

27.   Cruz C, Ribes V, Kutejova E, Cayuso J, Lawson V, Norris D, et al. Foxj1 regulates floor plate cilia architecture and modifies the response of cells to sonic hedgehog signalling. Development. 2010;137:4271-82. Epub 2010/11/26. doi: 137/24/4271 [pii]10.1242/dev.051714. PubMed PMID: 21098568; PubMed Central PMCID: PMC2990214.

28.   Kahane N, Kalcheim C. Identification of early postmitotic cells in distinct embryonic sites and their possible roles in morphogenesis. Cell Tissue Res. 1998;294:297-307.

29.   Ericson J, Morton S, Kawakami A, Roelink H, Jessell TM. Two critical periods of Sonic Hedgehog signaling required for the specification of motor neuron identify. Cell. 1996;87:661-73.

30.   Bishop B, Aricescu AR, Harlos K, O'Callaghan CA, Jones EY, Siebold C. Structural insights into hedgehog ligand sequestration by the human hedgehog-interacting protein HHIP. Nat Struct Mol Biol. 2009;16(7):698-703. doi: 10.1038/nsmb.1607. PubMed PMID: 19561611; PubMed Central PMCID: PMCPMC2709225.

31.   Holtz AM, Griffiths SC, Davis SJ, Bishop B, Siebold C, Allen BL. Secreted HHIP1 interacts with heparan sulfate and regulates Hedgehog ligand localization and function. J Cell Biol. 2015;209(5):739-57. doi: 10.1083/jcb.201411024. PubMed PMID: 26056142; PubMed Central PMCID: PMCPMC4460154.

32.   Kwong L, Bijlsma MF, Roelink H. Shh-mediated degradation of Hhip allows cell autonomous and non-cell autonomous Shh signalling. Nature communications. 2014;5:4849. doi: 10.1038/ncomms5849. PubMed PMID: 25215859; PubMed Central PMCID: PMCPMC4163939.

33.   Beug ST, Parks RJ, McBride HM, Wallace VA. Processing-dependent trafficking of Sonic hedgehog to the regulated secretory pathway in neurons. Mol Cell Neurosci. 2011;46(3):583-96. doi: 10.1016/j.mcn.2010.12.009. PubMed PMID: 21182949.

34.   Maun HR, Wen X, Lingel A, de Sauvage FJ, Lazarus RA, Scales SJ, et al. Hedgehog pathway antagonist 5E1 binds hedgehog at the pseudo-active site. J Biol Chem. 2010;285(34):26570-80. doi: 10.1074/jbc.M110.112284. PubMed PMID: 20504762; PubMed Central PMCID: PMCPMC2924095.

35.   Charrier JB, Lapointe F, Le Douarin NM, Teillet MA. Anti-apoptotic role of Sonic hedgehog protein at the early stages of nervous system organogenesis. Development. 2001;128:4011-20.

36.   Cayuso J, Ulloa F, Cox B, Briscoe J, Marti E. The Sonic hedgehog pathway independently controls the patterning, proliferation and survival of neuroepithelial cells by regulating Gli activity. Development. 2006;133(3):517-28. Epub 2006/01/18. doi: 133/3/517 [pii]10.1242/dev.02228. PubMed PMID: 16410413.

37.   Sasaki H, Nishizaki Y, Hui CC, Nakafuku M, Kondoh H. Regulation of Gli2 and Gli3 activities by an amino-terminal repression domain: implication of Gli2 and Gli3 as primary mediators of Shh signaling. Development. 1999;126:3915-24.

38.   McMahon JA, Takada S, Zimmerman LB, Fan CM, Harland RM, McMahon AP. Noggin-mediated antagonism of BMP signaling is required for growth and patterning of the neural tube and somite. Genes Dev. 1998;12:1438-52.

39.   Dessaud E, Yang LL, Hill K, Cox B, Ulloa F, Ribeiro A, et al. Interpretation of the sonic hedgehog morphogen gradient by a temporal adaptation mechanism. Nature. 2007;450(7170):717-20. doi: 10.1038/nature06347. PubMed PMID: 18046410.

40.   Diez del Corral R, Olivera-Martinez I, Goriely A, Gale E, Maden M, Storey K. Opposing FGF and retinoid pathways control ventral neural pattern, neuronal differentiation, and segmentation during body axis extension. Neuron. 2003;40(1):65-79. Epub 2003/10/07. doi: S0896627303005658 [pii]. PubMed PMID: 14527434.

41.   Wilson L, Gale E, Maden M. The role of retinoic acid in the morphogenesis of the neural tube. J Anat. 2003;203(4):357-68. Epub 2003/11/19. PubMed PMID: 14620376; PubMed Central PMCID: PMC1571178.

42.   Smith JC. Dorso-ventral patterning in the neural tube   specification of cell type in the neuraltube depends on signals derived from both dorsal cells and the floor plate , suggesting the existence of a double -gradient system. CurrBiol. 1993;3:582-5.

43.   Trousse F, Giess MC, Soula C, Ghandour S, Duprat A-M, Cochard P. Notochord and floor plate stimulate oligodendrocyte differentiation in cultures of the chick dorsal neural tube. JNeurosciRes. 1995;41:552-60.

44.   van Straaten HW, Hekking JW, Wiertz-Hoessels EJ, Thors F, Drukker J. Effect of the notochord on the differentiation of a floor plate area in the neural tube of the chick embryo. Anat Embryol (Berl). 1988;177(4):317-24. PubMed PMID: 3354847.

45.   van Straaten HW, Hekking JW, Beursgens JP, Terwindt-Rouwenhorst E, Drukker J. Effect of the notochord on proliferation and differentiation in the neural tube of the chick embryo. Development. 1989;107(4):793-803. PubMed PMID: 2632234.

46.   Chamberlain CE, Jeong J, Guo C, Allen BL, McMahon AP. Notochord-derived Shh concentrates in close association with the apically positioned basal body in neural target cells and forms a dynamic gradient during neural patterning. Development. 2008;135:1097-106. Epub 2008/02/15. doi: dev.013086 [pii]10.1242/dev.013086. PubMed PMID: 18272593.

47.   Matise MP, Epstein DJ, Park HL, Platt KA, Joyner AL. *Gli2* is required for induction of floor plate and adjacent cells, but not most ventral neurons in the mouse central nervous system. Development. 1998;125:2759-70.

48.   Yu K, McGlynn S, Matise MP. Floor plate-derived sonic hedgehog regulates glial and ependymal cell fates in the developing spinal cord. Development. 2013;140(7):1594-604. doi: 10.1242/dev.090845. PubMed PMID: 23482494; PubMed Central PMCID: PMCPMC3596997.

49.   Danesin C, Soula C. Moving the Shh Source over Time: What Impact on Neural Cell Diversification in the Developing Spinal Cord? J Dev Biol. 2017;5(2). doi: 10.3390/jdb5020004. PubMed PMID: 29615562; PubMed Central PMCID: PMCPMC5831764.

50.   Gritli-Linde A, Lewis P, McMahon AP, Linde A. The whereabouts of a morphogen: Direct evidence for short- and graded long-range activity of hedgehog signaling peptides. DevBiol. 2001;236:364-86.

51.   Matise MP, Epstein DJ, Park HL, Platt KA, Joyner AL. Gli2 is required for induction of floor plate and adjacent cells, but not most ventral neurons in the mouse central nervous system. Development. 1998;125(15):2759-70. PubMed PMID: 9655799.

52.   Ribes V, Balaskas N, Sasai N, Cruz C, Dessaud E, Cayuso J, et al. Distinct Sonic Hedgehog signaling dynamics specify floor plate and ventral neuronal progenitors in the vertebrate neural tube. Genes Dev. 2010;24:1186-200. doi: 24/11/1186 [pii]10.1101/gad.559910. PubMed PMID: 20516201; PubMed Central PMCID: PMC2878655.

53.   Vyas N, Walvekar A, Tate D, Lakshmanan V, Bansal D, Lo Cicero A, et al. Vertebrate Hedgehog is secreted on two types of extracellular vesicles with different signaling properties. Sci Rep. 2014;4:7357. doi: 10.1038/srep07357. PubMed PMID: 25483805; PubMed Central PMCID: PMCPMC4258658.

54.   Dierker T, Dreier R, Petersen A, Bordych C, Grobe K. Heparan sulfate-modulated, metalloprotease-mediated sonic hedgehog release from producing cells. J Biol Chem. 2009;284(12):8013-22. doi: 10.1074/jbc.M806838200. PubMed PMID: 19176481; PubMed Central PMCID: PMCPMC2658095.

55.   Chen MH, Li YJ, Kawakami T, Xu SM, Chuang PT. Palmitoylation is required for the production of a soluble multimeric Hedgehog protein complex and long-range signaling in vertebrates. Genes Dev. 2004;18(6):641-59. doi: 10.1101/gad.1185804. PubMed PMID: 15075292; PubMed Central PMCID: PMCPMC387240.

56.   Parchure A, Vyas N, Mayor S. Wnt and Hedgehog: Secretion of Lipid-Modified Morphogens. Trends Cell Biol. 2018;28(2):157-70. doi: 10.1016/j.tcb.2017.10.003. PubMed PMID: 29132729.

57.   Callejo A, Torroja C, Quijada L, Guerrero I. Hedgehog lipid modifications are required for Hedgehog stabilization in the extracellular matrix. Development. 2006;133(3):471-83. doi: 10.1242/dev.02217. PubMed PMID: 16396909.

58.   Etoc F, Metzger J, Ruzo A, Kirst C, Yoney A, Ozair MZ, et al. A Balance between Secreted Inhibitors and Edge Sensing Controls Gastruloid Self-Organization. Dev Cell. 2016;39(3):302-15. doi: 10.1016/j.devcel.2016.09.016. PubMed PMID: 27746044; PubMed Central PMCID: PMCPMC5113147.

59.   Singla V, Reiter JF. The primary cilium as the cell's antenna: signaling at a sensory organelle. Science. 2006;313(5787):629-33. doi: 10.1126/science.1124534. PubMed PMID: 16888132.

60.   Corbit KC, Aanstad P, Singla V, Norman AR, Stainier DY, Reiter JF. Vertebrate Smoothened functions at the primary cilium. Nature. 2005;437(7061):1018-21. doi: 10.1038/nature04117. PubMed PMID: 16136078.

61.   Gradilla AC, Sanchez-Hernandez D, Brunt L, Scholpp S. From top to bottom: Cell polarity in Hedgehog and Wnt trafficking. BMC Biol. 2018;16(1):37. doi: 10.1186/s12915-018-0511-x. PubMed PMID: 29636056; PubMed Central PMCID: PMCPMC5894156.

62.   Cardozo MJ, Sanchez-Arrones L, Sandonis A, Sanchez-Camacho C, Gestri G, Wilson SW, et al. Cdon acts as a Hedgehog decoy receptor during proximal-distal patterning of the optic vesicle. Nature communications. 2014;5:4272. doi: 10.1038/ncomms5272. PubMed PMID: 25001599; PubMed Central PMCID: PMCPMC4102123.

63.   Kahane, N., Cinnamon, Y., Kalcheim, C. The origin and fate of pioneer myotomal cells in the avian embryo. MechDev. 1998;74:59-73.

64.   Kalcheim C, Cinnamon Y, Kahane N. Myotome formation: A multistage process. Cell Tissue Res. 1999;296:161-73.

65.   Borycki AG. The myotomal basement membrane: insight into laminin-111 function and its control by Sonic hedgehog signaling. Cell Adh Migr. 2013;7(1):72-81. doi: 10.4161/cam.23411. PubMed PMID: 23287393; PubMed Central PMCID: PMCPMC3544789.

66.   Ben-Yair R, Kahane N, Kalcheim C. LGN-dependent orientation of cell divisions in the dermomyotome controls lineage segregation into muscle and dermis. Development. 2011;138:4155-66. Epub 2011/08/20. doi: dev.065169 [pii]10.1242/dev.065169. PubMed PMID: 21852400.

67.   Blaess S, Graus-Porta D, Belvindrah R, Radakovits R, Pons S, Littlewood-Evans A, et al. Beta1-integrins are critical for cerebellar granule cell precursor proliferation. J Neurosci. 2004;24(13):3402-12. doi: 10.1523/JNEUROSCI.5241-03.2004. PubMed PMID: 15056720; PubMed Central PMCID: PMCPMC2693074.

68.   Anderson C, Thorsteinsdottir S, Borycki AG. Sonic hedgehog-dependent synthesis of laminin alpha1 controls basement membrane assembly in the myotome. Development. 2009;136:3495-504. Epub 2009/09/29. doi: 136/20/3495 [pii]10.1242/dev.036087. PubMed PMID: 19783738; PubMed Central PMCID: PMC2752398.

69.   Borycki AG, Brunk B, Tajbakhsh S, Buckingham M, Chiang C, Emerson CP, Jr. Sonic hedgehog controls epaxial muscle determination through Myf5 activation. Development. 1999;126:4053-63. Epub 1999/08/24. PubMed PMID: 10457014.

70.   Sela-Donenfeld D, Kalcheim C. Inhibition of noggin expression in the dorsal neural tube by somitogenesis: a mechanism for coordinating the timing of neural crest emigration. Development. 2000;127:4845-54.

71.   Marcelle C, Stark MR, Bronner-Fraser M. Coordinate actions of BMPs, Wnts, Shh and noggin mediate patterning of the dorsal somite. Development. 1997;124:3955-63.

72.   Abu-Elmagd M, Robson L, Sweetman D, Hadley J, Francis-West P, Munsterberg A. Wnt/Lef1 signaling acts via Pitx2 to regulate somite myogenesis. Dev Biol. 2010;337:211-9. doi: S0012-1606(09)01282-2 [pii]10.1016/j.ydbio.2009.10.023. PubMed PMID: 19850024.

73.   Sela-Donenfeld D, Kalcheim C. Localized BMP4-noggin interactions generate the dynamic patterning of noggin expression in somites. DevBiol. 2002;246:311-28.

74.   Brill G, Kahane N, Carmeli C, Von Schack D, Barde Y-A, Kalcheim C. Epithelial-mesenchymal conversion of dermatome progenitors requires neural tube-derived signals: Characterization of the role of Neurotrophin-3. Development. 1995;121:2583-94.

75.   Kalcheim C. Regulation of trunk myogenesis by the neural crest: a new facet of neural crest-somite interactions. Dev Cell. 2011;21:187-8. Epub 2011/08/16. doi: S1534-5807(11)00301-7 [pii]10.1016/j.devcel.2011.07.009. PubMed PMID: 21839914.

76.   Kalcheim C, Goldstein R. Segmentation of sensory and sympathetic ganglia: Interaction between neural crest and somite cells. JPhysiol(Lond). 1991;85:110-7.

77.   Krispin S, Nitzan E, Kassem Y, Kalcheim C. Evidence for a dynamic spatiotemporal fate map and early fate restrictions of premigratory avian neural crest. Development. 2010;137:585-95. Epub 2010/01/30. doi: 137/4/585 [pii]10.1242/dev.041509. PubMed PMID: 20110324.

78.   Briscoe J, Chen Y, Jessell TM, Struhl G. A hedgehog-insensitive form of patched provides evidence for direct long-range morphogen activity of sonic hedgehog in the neural tube. Mol Cell. 2001;7:1279-91. Epub 2001/06/30. doi: S1097-2765(01)00271-4 [pii]. PubMed PMID: 11430830.

79.   Gupta S, Sen J. Retinoic acid signaling regulates development of the dorsal forebrain midline and the choroid plexus in the chick. Development. 2015;142(7):1293-8. doi: 10.1242/dev.122390. PubMed PMID: 25758461.

80.   Novitch BG, Wichterle H, Jessell TM, Sockanathan S. A requirement for retinoic acid-mediated transcriptional activation in ventral neural patterning and motor neuron specification. Neuron. 2003;40(1):81-95. Epub 2003/10/07. doi: S0896627303005294 [pii]. PubMed PMID: 14527435.

81.   Halperin-Barlev O, Kalcheim C. Sclerotome-derived Slit1 drives directional migration and differentiation of Robo2-expressing pioneer myoblasts. Development. 2011;138:2935-45. Epub 2011/06/10. doi: dev.065714 [pii]10.1242/dev.065714. PubMed PMID: 21653616.

82.   Kahane N, Ben-Yair R, Kalcheim C. Medial pioneer fibers pattern the morphogenesis of early myoblasts derived from the lateral somite. Dev Biol. 2007;305:439-50. PubMed PMID: 17382923.

83.   Kahane, N., Cinnamon, Y., Bachelet, I., Kalcheim, C. The third wave of myotome colonization by mitotically competent progenitors: regulating the balance between differentiation and proliferation during muscle development. Development. 2001;128:2187-98.

84.   Burstyn-Cohen T, Kalcheim C. Association between the Cell Cycle and Neural Crest Delamination through Specific Regulation of G1/S Transition. DevCell 2002;3:383-95.

Legends to Figures

Fig. 1- Reduction of Shh in sclerotome by Hhip1 affects both myotome and motoneuron differentiation.

 (A, B) Electroporation of control GFP (A) or Hhip1/GFP (B) to the prospective sclerotome. A day later a reduction in myotome size (blue desmin staining) is apparent adjacent to the transfected cells in B when compared to the control side and to control GFP (A). In addition, note the ventral expansion of the Pax7+ domain in B (small arrows) when compared to the control side and to control GFP.

(A', B') Higher magnification of the insets in A,B, respectively depicting a reduction of Hb9+ motoneurons in B upon Hhip1 treatment when compared to the control side and to control GFP (A'). Asterisks (*) denote motoneurons adjacent to the electroporated sclerotomes. (C) Quantification of Hb9+ motoneurons (***, p<0.001).

(D-D") Double electroporation of ShhN:YFP to NT and Hhip1:CD4 to sclerotome. Note that sclerotomal cells missexpressing Hhip:CD4 (red) are decorated by Shh (green) immunolabeling (D', D" arrows).

(E-E") Double electroporation of ShhN:YFP to NT and Hhip1 to sclerotome. No labeling of Shh (green) is apparent in sclerotomal cells missexpressing Hhip1 (red). D" and E" are high magnifications of insets in D and E.

(F-F") Double electroporation of ShhN:YFP to sclerotome and Hhip1:CD4 to NT. Note labeling of Shh (green) in both the basement membrane as well as along neuroepithelial cells missexpressing Hhip:CD4 (F', F"). F" is a higher magnification of the inset in F/F' where only ShhN:YFP (green) is shown to highlight that Shh co-localizes with the red, Hhip1:CD4-missexpressing cells, in the neuroepithelium (F).

(G-G") Double electroporation of ShhN:YFP to sclerotome and Hhip1 to NT. Note labeling of Shh (green) in basement membrane but no co-staining of neuroepithelial cells missexpressing Hhip1. G" is a high magnifications of the inset in G'. Abbreviations, DM, dermomyotome; M, myotome; NT, neural tube; No, notochord; Scl, sclerotome. Bars=50 μM.

Fig.2-Electroporation of Hhip:CD4 and Ptc1 into sclerotome inhibit motoneuron differentiation without significantly affecting progenitor proliferation or survival.

(A-E') Electroporation of depicted plasmids (green) to the sclerotome followed a day later by immunostaining for Pax7 and Hb9. A'-E' represent higher magnifications of the boxed regions in A-E. Note that all plasmids except $PTC^{\Delta loop2}$ reduced the number of motoneurons adjacent to the transfected sclerotomes. Note in panel B that Hhip1 caused a slight ventralization of Pax7 expression (arrows in B) in association with a reduced hemi-NT.

(F-J) Tunel staining (red). Note weak immunostaining of electroporated plasmids (green) because no anti-GFP antibodies were implemented to enable better visualization of apoptotic nuclei. Only Hhip1 exhibited numerous Tunel+ nuclei adjacent to the transfected sclerotome.

(K-O) staining of mitotic nuclei with anti pH3 (red).

(P) Quantification of motoneurons, mitotic nuclei and area. (Q) Quantification of the relative area of electroporated sclerotomes. Abbreviations, NT, neural tube; Scl, sclerotome. *p<0.05, **p<0.03, ***p<0.01. Bar=50μM.

Fig. 3- Depletion of Shh activity by Hhip:CD4 in sclerotome inhibits motoneuron differentiation and is rescued by co-treatment with Shh

(A-D) Electroporation of the depicted plasmids to sclerotome (Scl) (green). Hhip:CD4 to Scl reduces motoneuron numbers compared to control GFP, whereas co-transfection with Shh rescues the effect. A'-D' are higher magnifications of the insets in A-D. (E) Quantification of Hb9+ motoneurons, ***$p<0.001$. Bar=50 μM.

Fig. 4- Electroporations of Hhip, Hhip:CD4 or Ptc1, but not of PTC$^{\Delta loop2}$ into the neural tube, reduce myotome size.

(A-E) Electroporation of the depicted plasmids (green). Hhip1, Hhip:CD4 and Ptc1 reduce the size of adjacent desmin+ myotomes (arrows, red) compared to control GFP yet PTC$^{\Delta loop2}$has no significant effect. (F) Quantification of myotome size. ***$p<0.001$. Bar=50 μM.

Fig.5- Gain of Shh function in sclerotome enhances motoneuron differentiation but Shh missexpression in NT has no effect on myotome

(A-C) Electroporation of Shh (green) to the sclerotome (Scl) enhances the number of Hb9+ motoneurons in the ventral neural tube (NT) compared to control (arrow in B, red). Note in inset the enlarged desmin+ myotome (blue) when compared to the control side (B). Quantification in C, ***$p<0001$. (D-F) Electroporation of Shh (green) to the NT has no effect on the size of desmin+ myotomes (blue) or on expression of Pax7 (red) in the dermomyotme. Quantification in F.
Abbreviations, No, notochord; NT, neural tube, Scl, sclerotome. Bar=50 mM.

Fig. 6- Sclerotome, but not FP-derived Shh is necessary for motoneuron differentiation

(A,B) Dorsoventral electroporation of control GFP showing in A, the presence of the labeled floor plate (FP) that co-expresses Shh protein. In B, note Hb9+ motoneurons dorsal to the labeled FP. (C) Hb9+ motoneurons in control GFP electroporation to sclerotome.

(D,E) Loss of FP tissue upon Hhip:CD4 electroporation. Asterisks (**) in D and E denote absence of a FP and panel D shows concomitant absence of Shh expression. Hhip:CD4 electroporation has no effect on ventral Hb9+ motoneurons or dorsal Pax7 expression (both in red). (F) In contrast, less motoneurons are apparent adjacent to the sclerotome transfected with Hhip:CD4 (arrow). (G) Loss of the FP upon electroporation of Ptc1 (**) shows no apparent effect on motoneurons. (H) Quantification showing no effect of FP deletion on the proportion of Hb9+ motoneurons in flank (electroporated)/neck (intact region). Abbreviations, No, notochord; NT, neural tube. Bar=50 μM.

Fig. 7- A basal, but not apical, presentation of Shh is required for ligand activity on both NT and DM/myotome. A model accounting for the effects of Shh traversing the sclerotome on both NT and myotome development.

Implantation of No fragments (dotted circles) in apical (A,C,E) or basal (B,D,F) positions vis-à-vis the dorsal NT, or the dermomyotome (DM) (E,F).

(A,C) A No piece was grafted inside the NT facing its luminal side. Note that a day later it localizes in the dorsal NT. No effect on Pax7 (arrows) or Hb9+ motoneurons (arrowheads) is apparent.

(B,D) A No piece was grafted in a basal position vis-à-vis the dorsal NT. Note a dorsal restriction of Pax7 (arrows in B) as well as a dorsalward expansion of Hb9+ motoneurons (arrowheads in D).

(E,F) A No fragment localized apical to the DM causes a mild change in the adjacent desmin+ myotome (green, E) whereas the basally located No strongly increases myotomal size and promotes an in situ differentiation of the DM into myotome with concomitant loss of Pax7 expression.

(G) A proposed model for the activity of No-derived Shh. The No secretes Shh that acts on the ventral NT and also traverses the sclerotome (Scl). We propose that a ventral to dorsal gradient of ligand is created in Scl and plays a pivotal role both on myotome differentiation as well as on motoneuron development. Shh is thus presented to the target cells via its basal domain, likely by initial association with the laminin-containing basement membrane. See text for details. Bar=50 μM.

Legends to Supplementary Figures

Supplementary. Fig.S1- Expression of neural tube markers by the time of *in ovo* electroporation

(A-D) In situ hybridization of embryos aged 25 somite pairs showing expression of ventral neural tube (NT) markers at the epithelial somite (ES) stage.

(E,F) Co-immunolabeling of Pax7 and Hb9 antibodies. Note in E that only Pax7 is expressed in both dorsal NT and dorsal aspect of the ES, whereas the first Hb9+ ventral motoneurons appear at a later stage following somite dissociation (arrows in F). Both dorsal Pax7 and ventral Hb9 immunostainings are in red as the respective antibodies are monoclonal of the same subtype. Abbreviations, No, notochord. Bar=50μM.

Supplementary Fig. S2- Differential behavior of secreted Hhip1 compared to membrane-tethered Hhip:CD4

(A-C) ShhN:YFP, electroporated into sclerotome (Scl, green), colocalizes with laminin (red) in the basement membrane around the neural tube (NT) (arrows).

(D-F') Immunolabeling with Shh and Hb9 antibodies of embryos electroporated with control GFP (D,D'), Hhip:CD4 (E,E') or Hhip1 (F,F'). Note that Hhip1 caused a reduction in ipsilateral Shh in both notochord (No) and FP (FP) (arrows in F') whereas GFP or Hhip:CD4 were without effect. Bars=50 μM.

Supplementary. Fig.S3- Depletion of Shh in the NT by Hhip:CD4 phenocopy those exerted by Hhip, Ptc1 and PTC$^{\Delta loop2}$

(A-E') Electroporation of depicted plasmids (green) to hemi-NTs followed a day later by immunostaining for Pax7 and Hb9. A'-E' represent higher magnifications of the boxed regions in A-E.

(F-J) staining of mitotic nuclei with anti pH3 (red) following electroporation with depicted plasmids.

(K-O) Tunel staining (red). Note weak immunostaining of electroporated plasmids (green) because no anti-GFP antibodies were implemented to enable better visualization of apoptotic nuclei.

(P) Quantification of motoneurons, mitotic nuclei and area. $*p<0.05$, $**p<0.03$, $***p<0.01$. Bar=50μM.

Supplementary Figure S4- Electroporation of Hhip:CD4 into the sclerotome reduces the extent of *Olig2* expression in NT.

(A,B) Electroporation of control GFP or Hhip:CD4, respectively. Asterisks denote the transfected sclerotomes. The extent of *Olig2* mRNA expression is delimited by black lines. Bar=50μM.

Supplementary. Fig. S5- The effects of Hhip:CD4 on NT and myotome are a direct consequence of Shh depletion

(A-I) In situ hybridization for *Gli3, Gli1* and *Hhip1* as depicted following electroporation of control GFP or Hhip:CD4. Asterisks (*) mark the electroporated sites in each panel. Control GFP had no effect on expression of either gene. In contrast, Hhip:CD4 transfected to NT reduced both *Gli1* and *Hhip1* mRNAs in NT and sclerotome (Scl) (arrowheads in E and H). Likewise, Hhip:CD4 transfected to sclerotome reduced both *Gli1* and *Hhip1* mRNAs in NT and sclerotome (Scl)

(arrowheads in F and I). No effect was apparent on expression of *Gli3* (A-C). Bar=50 μM.

Supplementary Figure S6- Electroporation of Hhip:CD4 to sclerotome has no effect on BMP signaling in dorsal NT.

(A,B) Missexpression of Hhip:CD4 in sclerotome (green) had no effect on expression of phospho-Smad 1,5,8 in the dorsal NT compared to control GFP (arrowheads), yet reduced the number of Hb9+ motoneurons (arrow in B). (C) Quantification of the area of phospho-Smad 1,5,8 expression. Bar=50 μM.

Supplementary Figure S7- Inhibition of Shh activity in sclerotome or NT has no effect on retinoic signaling in the adjacent tissue.

(A-B") Specificity of the retinoic acid reporter (pRARE-AP). Note in A-A" the expression of pRARE-AP in control GFP-electroporated NT. In contrast, no signal is apparent when retinoic acid activity is abolished by a dnRAR plasmid (B-B"). A" and B" are overlays of the precedent panels, respectively.

(C-F) Double electroporations of Hhip1 and RARE-AP to sclerotome and NT and vice-versa. (C,D) Missexpression of Hhip1 in sclerotome (*) had no effect on RARE-AP activity in NT. (E,F) Missexpression of Hhip1 in NT (*) had no effect on RARE-AP activity in sclerotome. Bars=50 μM.

Fig.1 Kahane and Kalcheim

Fig. 2 Kahane and Kalcheim

Fig. 3 Kahane and Kalcheim

Fig. 4 Kahane and Kalcheim

Fig.5 Kahane and Kalcheim

Fig. 6   Kahane and Kalcheim

Fig. 7   Kahane and Kalcheim

Suppl. Fig.1 Kahane and Kalcheim

Suppl. Fig. 2 Kahane and Kalcheim

Suppl. Fig. 3 Kahane and Kalcheim

GFP | **Hhip:CD4**

A

NT

*

Olig 2

B

*

Suppl. Fig. 4 Kahane and Kalcheim

Suppl. Fig. 5 Kahane and Kalcheim

Suppl. Fig. 6 Kahane and Kalcheim

Suppl. Fig.7 Kahane and Kalcheim