Community structure – ecosystem function relationships in the Congo Basin

methane cycle depend on the physiological scale of function

Kyle M. Meyer[1#*+], Anya M. Hopple[1#], Ann M. Klein[1], Andrew H. Morris[1], Scott

Bridgham[1], Brendan J. M. Bohannan[1]

1. Institute of Ecology and Evolution, University of Oregon, Eugene, OR, USA

# Authors contributed equally to the preparation of this manuscript

*Corresponding author: kmmeyer@berkeley.edu

+ Present address: Department of Integrative Biology, University of California –

Berkeley, Berkeley, CA, USA

17 Keywords: Biodiversity-Ecosystem Function, Congo Basin, Methane, Microbial

18 ecology

20 Running title: Community structure – function relationships

ABSTRACT

Belowground ecosystem processes can be highly variable and difficult to predict using microbial community data. Here we argue that this stems from at least three issues: 1) complex covariance structure of samples (with environmental conditions or spatial proximity) can make distinguishing biotic drivers a challenge, 2) communities can control ecosystem processes through multiple mechanisms, making the identification of these controls a challenge and 3) ecosystem function assessments can be broad in physiological scale, encapsulating multiple processes with unique microbially mediated controls. We test these assertions using methane (CH_4)-cycling processes in soil samples collected along a wetland-to-upland habitat gradient in the Congo Basin. We perform our measurements of function under controlled laboratory conditions and include environmental covariates in statistical analyses to aid in identifying biotic drivers. We divide measurements of microbial communities into four attributes (abundance, activity, composition, and diversity) that represent different forms of community control. Lastly, our process measurements differ in physiological scale, including broader processes (gross methanogenesis and methanotrophy) that involve more mediating groups, to finer processes (hydrogenotrophic methanogenesis and high-affinity CH_4 oxidation) with fewer mediating groups. We observed that finer scale processes can be more readily predicted from microbial community structure than broader scale processes. In addition, the nature of those relationships differed, with broad processes limited by abundance while fine-scale processes were associated with diversity and composition. These findings demonstrate the importance of carefully defining the

47   physiological scale of ecosystem function and performing community

48   measurements that represent the range of possible controls on ecosystem

49   processes.

INTRODUCTION

Belowground ecosystem processes can be highly variable, and there have

been numerous attempts to explain or predict this variation using microbial

community data. However, our ability to identify microbial community drivers of

processes remains in its infancy. This challenge stems from at least three issues.

First, community variables often have complex covariance structure (e.g. spatial or

environmental autocorrelation) (Legendre, 1993), which can complicate the task of

distinguishing the biotic signal underlying ecosystem process rate variation. Second,

communities can control processes through a number of mechanisms, making the

identification of these controls a challenge (Bier et al., 2015). For example,

communities can influence the rate of a process through abundance (of particular

groups or the community as a whole), composition (i.e. the identity of community

members), diversity, and activity, each of which would require the quantification of

different community attributes. Lastly, function is often assessed at a broad

physiological scale that encapsulates many processes that could have unique

controls (Inkpen et al., 2017). For example, gross decomposition of organic matter

or net flux of methane (CH_4) into the atmosphere are commonly measured

belowground ecosystem functions, yet these measurements represent the aggregate

result of many different physiological processes.

We tested this idea using the two counteracting microbial processes that

largely control atmospheric CH_4 emissions from soils: CH_4 production

(methanogenesis) and CH_4 consumption (methanotrophy). Each of these processes

can be further divided into underlying processes that involve fewer mediating

microbial taxa. For instance, gross methanogenesis can have direct inputs from at

least three processes: hydrogenotrophic (which uses CO_2 and H_2 to produce CH_4),

acetoclastic (which uses acetate), and/or methylotrophic (which uses methanol,

methyl-amides, or methyl-sulfides) methanogenesis. The process of methanotrophy

can be subdivided as well. Methanotrophy at high CH_4 concentrations (typical of

those found in waterlogged soil and sediment) is performed by a wide diversity of

methanotrophs found in the Gammaproteobacteria, Alphaproteobacteria, and

Verrucomicrobia (Knief, 2015), whereas the consumption of CH_4 at atmospheric

concentrations (a.k.a. high-affinity CH_4 oxidation) appears to be much more

phylogenetically restricted and is likely performed by many fewer taxa (Ho et al.,

2013; Knief, 2015; Kolb, 2009). Taken together, these observations suggest that

gross measurements of either methanogenesis or methanotrophy may mask subtle

variations in these underlying processes and could thereby blur the connection

between community variation and ecosystem function.

A major challenge in the study of ecosystem function lies in identifying

underlying community controls. Group abundance is one such form of control,

whereby the rate of a process only proceeds as quickly as the total number of

capable individuals. There are examples of this control for both methanogenesis

(Ma, Conrad, & Lu, 2012) and methanotrophy (Freitag & Prosser, 2009), but many

studies have found no such relationship (see Rocca et al., 2014). While this could be

driven in part by technical issues (e.g. abundance being inaccurately assessed due to

primer bias), processes may be controlled by other community attributes. Most soil

organisms are in an inactive state at any given time (Lennon and Jones, 2011). Low

activity levels can limit the rate of ecosystem processes either by decreasing

syntrophic interactions or preventing substrate uptake. Measurements of

methanotroph and methanogen transcriptional activity levels have been shown to

positively correlate with process rates (Freitag & Prosser, 2009; Freitag, Toet,

Ineson, & Prosser, 2010), suggesting that the activity levels of key groups can

constrain these processes. Moreover, community assessments using RNA may more

closely approximate the active fraction of the community (Jones & Lennon, 2010;

Kamke et al., 2010; Baldrian et al., 2012; although see Papp et al., 2018a) and

therefore provide a means to approximate activity-based community controls.

The composition of a community can impose another constraint on

ecosystem processes. Composition control (a.k.a. species identity effects (Bannar -

Martin et al., 2018)) could occur if certain community members perform a process

more efficiently or rapidly than others, and those taxa/traits are dispersal limited.

In this case, certain individuals would be expected to more strongly correlate with

process rate variation than others. Distinguishing such a compositional signal can be

a challenge because communities tend to co-vary depending on their spatial

proximity or similarity in environmental controls (Legendre, 1993). There is

precedent to suggest that community composition is an important determinant of

CH_4-cycling rates. For instance, even closely related methanogen/methanotroph

taxa can differ in traits such as substrate affinity, substrate preference,

environmental tolerance, or competitive ability (Garcia, Patel, & Ollivier, 2000; Ho et

al., 2013; Knief, 2015; Nazaries, Murrell, Millard, Baggs, & Singh, 2013). Therefore,

differences in community composition have the potential to influence CH_4-cycling

processes, which has been shown in several systems (Bodelier et al., 2013; McCalley

et al., 2014; Nazaries, Pan, et al., 2013; Sierocinski et al., 2017, 2018). Membership of

other functional guilds within a community is also an important consideration,

particularly for processes that require metabolic byproducts as the primary

substrate. Methanogens depend on the fermentative byproducts (*e.g.* H^+ and

acetate) of other community members and must compete for these substrates with

other community members. This is also the case for methanotrophs, where

competition for O_2 and/or other soil nutrients can limit rates of methanotrophy

(Bodelier, Roslev, Henckel, & Frenzel, 2000). Thus, *who is there* may be a

particularly important consideration for predicting the rates of ecosystem

processes, particularly if efficient traits are dispersal limited or competition for

substrates is common.

Lastly, a growing body of work suggests that biodiversity may affect rates of

ecosystem processes (Tilman, Wedin, & Knops, 1996), as well as the resistance,

resilience, or stability of a process through time (Loreau et al., 2001; Tilman &

Downing, 1994). This pattern could arise through multiple ecological mechanisms

including: 1) niche complementarity, 2) sampling effects, or 3) facilitation. It has

been argued that ecosystem functions mediated by a wide assortment of taxa may

be less influenced by levels of diversity than those mediated by relatively fewer taxa

(Schimel, 1995; Schimel & Gulledge, 1998). This would suggest that as one narrows

the physiological scale of ecosystem function, the diversity of participants should

become a stronger determinant of process rates. The abilities to produce and

consume CH_4 are both highly phylogenetically conserved (Martiny, Treseder, &

Pusch, 2013), and relatively few taxa possess the ability to carry out these

processes. There are even fewer taxa that perform underlying processes such as

hydrogenotrophic methanogenesis and high-affinity methanotrophy (Knief, 2015;

Kolb, 2009). Positive diversity – ecosystem function relationships have been

observed for methanotrophy experimentally using intact soil cores from the field

(Bodelier et al., 2013), as well as in artificially assembled methanotroph

communities (Schnyder, Bodelier, Hartmann, Henneberger, & Niklaus, 2018). This

has also been observed for methanogenesis in anaerobic digesters (Sierocinski et al.,

2018). Although several studies have tested whether diversity and process rates are

related, few have asked whether adjusting the physiological scale at which function

is assessed improves this relationship.

Here we present results from a set of experiments designed to identify the

biological drivers of variation in CH_4-cycling dynamics. Our goals were to determine

(1) whether we could identify significant community controls on CH_4 cycling after

accounting for covariance structure, and (2) whether the strength and nature of

those controls changes with the physiological scale of the process. To do this, we

collected soils along a diverse wetland-to-upland habitat gradient in the Congo

Basin, a region that has received very little scientific attention for its role in the

global CH_4 cycle (Bridgham, Cadillo-Quiroz, Keller, & Zhuang, 2013; Kirschke et al.,

2013) and has been shown to exhibit highly dynamic spatial and temporal CH_4

cycling dynamics (R. A. Delmas, Tathy, & Cros, 1992; R. Delmas, Servant, Tathy, Cros,

& Labat, 1992; MacDonald et al., 1999; Tathy et al., 1992). We estimate process rate

potentials under controlled laboratory conditions, which minimizes the impact of

environmental variation and enhances the biotic signal underlying process rate

variation. We varied the physiological scale at which we assessed ecosystem

function, ranging from broader scales including gross methanogenesis and

methanotrophy, to finer scales including hydrogenotrophic methanogenesis and

high-affinity methanotrophy. We paired RNA- and DNA-based microbial community

measurements with each process measurement and calculated four sets of

community attributes (abundance, activity, composition, and diversity) to aid in the

identification of putative community controls. We hypothesized that there would be

a connection between microbial community structure and CH_4-cycling processes

after accounting for environmental variation, and that finer scale processes (*i.e.*

those involving a lower diversity of mediating taxa) would exhibit a stronger

relationship with community attributes than broader processes, as exemplified by a

higher proportion of variance explained. Our results suggest that communities are

an important driver of CH_4-cycling processes in tropical ecosystems, and that the

strength and nature of the relationship between community structure and

ecosystem function can depend on the physiological scale by which function is

assessed.

METHODS AND MATERIALS

Site selection and sampling

We selected a diverse set of 15 sites in Southwestern Gabon in and around

the Gamba Complex of Protected Areas (Lee, Alonso, Dallmeier, Campbell, &

Pauwels, 2006). Gabonese ecosystems belong to the Guineo-Congolian regional

center of endemism (White, 1979), making them broadly representative of the

Congo Basin. Our sites included 4 mineral-soil forested wetlands, 2 peatlands, 2

seasonally wet mineral-soil forests, 3 upland grasslands (2 with termite mounds), 2

upland forests, a 1-year old plantation, and 1 abandoned plantation. Seasonal

wetlands were identified by tree species composition, thick O and/or A soil horizons

with mottling, and tree watermarks. Water tables were at least 40 cm below the soil

surface in seasonal wetlands. Our sampling took place at the beginning of the

Gabonese rainy season, from October to November 2014.

Potential gross and high-affinity CH_4 oxidation rates

We collected three intact soil cores from above the water table at the

beginning, middle, and end of 35 m field transects established at each site. The

samples were stored in PVC tubes (5 cm dia., 8 cm height) to maintain soil structure

and kept at *in situ* air temperatures (28 °C) until the end of the sampling trip (~3

weeks). One week after returning to the University of Oregon, each PVC core was

placed into a gas-tight Mason jar that was retro-fitted with a headspace sampling

port and incubated at 28 °C in the dark. Rates of aerobic CH_4 oxidation were

determined under initially low (5 ppm CH_4/mL in headspace) and high (1000 ppm

CH_4/mL headspace) CH_4 concentrations, to estimate rates of high-affinity and gross

methanotrophy, respectively. CH_4 concentration measurements were taken at 0.33,

3, 6, and 9 hours for high-affinity methanotrophy and at 0.33, 3, 6, 9, 24, and 48

hours for gross methanotrophy. The same soil cores were used to determine high-

affinity methane oxidation and gross methanotrophy in quick succession. We

applied a first-order exponential decay function to determine the rate constant (k,

units = d^{-1}; *i.e.* $dCH_4/dt = k[CH_4]$) of the exponential decrease in CH_4 when at low

initial CH_4 concentrations (*i.e.* high-affinity methanotrophy). At high CH_4

concentrations we measured the linear decreases in CH_4 to determine the maximum

velocity rates (V_{max}, units = $\mu g\ CH_4\ cm^{-3}\ d^{-1}$) (*i.e.* gross methanotrophy). During our

high-affinity CH_4 oxidation incubations, CH_4 concentrations exponentially decreased

in all samples, except for one, across our four measurements; thus, all time points

were included in subsequent calculations. While one sample did not immediately

exhibit CH_4 oxidation at the first timepoint (0.33 hours), it did show exponential

decreases in CH_4 concentrations following the 3 hour timepoint; thus, only three

timepoints were used to calculate the k-value for this sample. During our gross

methanotrophy incubations, we included only timepoints that encompassed the

linear portion of the exponential decrease in CH_4 concentration, which ranged from

3-5 timepoints depending on the sample.

Potential gross methanogenesis and hydrogenotrophic pathway predominance

Across all wetland sites, we collected three intact soil cores in PVC tubes

from 0 – 10 cm below the water table at the beginning, middle, and end of 35 m

transects at each site. The samples were topped with site water and tightly sealed to

ensure an anaerobic environment and transported to the University of Oregon.

Anaerobic incubations took place at field temperatures (28 °C) within two days

following arrival at University of Oregon. In a glove box filled with a N_2 atmosphere

(<5% H_2 in the presence of palladium catalyst; Coy Laboratory Products, Grass Lake,

Michigan, USA), approximately 10 g of wet weight soil were added to 120 mL serum

bottles and mixed with 10 mL of deoxygenated, deionized water. Sample bottles

were then flushed with N_2 for 15 minutes to begin the incubation. Headspace

samples were analyzed over the course of three days (0, 1, 2, and 3 days). Total CH_4

was calculated using Henry's Law, adjusting for solubility, temperature, and pH

(Bridgham & Ye, 2013). CH_4 production rates were calculated using the linear

accumulation ($r^2 > 0.90$ in all cases) of gas through time. To measure

hydrogenotrophic methanogenesis, we used a $^{14}CO_2$ tracer method (Keller &

Bridgham, 2007), which allowed us to distinguish the percent of total CH_4 that had

been produced using CO_2/H_2 as the substrate (i.e. hydrogenotrophic

methanogensis). This was done with a gas chromatograph fitted with a radioactive

gas proportional counter (LabLogic Systems) simultaneously with the gross

methanogenesis measurements over three days.

Soil physical and chemical analysis

We measured a suite of abiotic variables from each soil sample to

incorporate into our community model as a covariate (see below). For the

methanogenesis study, we recorded pH, total % nitrogen (N), organic carbon (C),

moisture content, and bulk density. For the methanotrophy study, we measured N,

C, moisture content, and bulk density. Soil pH was determined from a 1:1 soil to

deionized water solution. Total % N and organic C were measured on a Costech ECS

4010 Elemental Analyzer (Valencia, CA), with each sample analyzed in duplicate.

Moisture content (gravimetric water content) and bulk density were measured by

the change in weight of a soil sub-sample following 48 hours of drying at 60 °C.

Additionally, material from the three soil cores at each site was combined and

homogenized for texture analysis using the hydrometer method (Gavlak, Horneck,

Miller, & Kotuby-Amarcher, 2003), with 5% sodium hexametaphosphate as the

dispersing solution. Because soil texture measurements required pooling samples

from each site- and all other variables were measured on a per sample basis, texture

was not included in the environmental covariate calculation.

Soil RNA/DNA co-extraction and sequencing

286  Soil DNA and RNA were co-extracted from Lifeguard-preserved soil samples

using MoBio's Powersoil RNA Isolation kit with the DNA Elution Accessory Kit

(MoBio, California, USA) following manufacturer's instructions. RNA was reverse

transcribed to cDNA using Superscript III first-strand reverse transcriptase and

random hexamer primers (Life Technologies, USA). Extractions were quantified

using Qubit (Life Technologies, USA). We amplified the V4 region of the 16S SSU

rRNA gene in sample DNA and cDNA using the primers 515F and 806R (Caporaso et

al., 2011). Sequencing libraries were prepped using a dual-indexing approach

(Fadrosh et al., 2014; Kozich, Westcott, Baxter, Highlander, & Schloss, 2013). In

short, each PCR reaction was performed using 12.5 µl NEBNext Q5 Hot Start HiFi

PCR master mix (New England Biolabs, USA), 11.5 µl gene-specific primer mix (1.09

µM of each primer), and 1 µl template (DNA or cDNA). A subset of samples was used

to find 1) the optimal primer annealing temperature, and 2) the minimum number

of cycles for adequate target amplification. For the 16S rRNA gene target this was

61° C and 20 cycles. The final reaction conditions were: 98° C 30 seconds

(initialization), 98° C 10 seconds (denaturation), gene-specific annealing step for 20

seconds (see above), and 72° C for 20 seconds (final extension). Reactions were

followed by magnetic bead purification using 20 μl Mag-Bind RxnPure Plus isolation

beads (Omega Bio-Tek, USA). Reactions were quantified using Qubit (Life

Technologies, USA) then were multiplexed at equimolar concentration. Final pooled

amplicon libraries were sequenced using the Illumina MiSeq (300 paired-end)

platform at the Oregon State University Center for Genome Research and

Biocomputing facility.

Bioinformatic processing

Demultiplexed reads were joined using PEAR (version 0.9.10) with default

parameters (Zhang, Kobert, Flouri, & Stamatakis, 2014) and quality filtered using

Prinseq (version 0.20.4) (Schmieder & Edwards, 2011). Sequences with a mean

quality score ≥ 30 and length 250-350 bp were retained. Sequences were

dereplicated, denoised, and checked for chimeras using the DADA2 pipeline

(Version 1.6) (Callahan et al., 2016) implemented in QIIME2 (Caporaso et al., 2010)

(https://qiime2.org). For the methanogenesis study 17 of the 18 DNA-based

samples, and 13 of the 18 RNA-based samples had sufficient quality to be included.

For the methanotrophy study 41 of the 44 DNA-based samples, and 34 of the 44

RNA-based samples passed our quality filter. Taxonomy was assigned to the

resulting amplicon sequence variants (ASVs) using the Ribosomal Database Project

online classifier (Wang, Garrity, Tiedje, & Cole, 2007) with database release 11.5

(Cole et al., 2014).

Quantitative PCR

We quantified the abundance of methanogens in each sample from which

CH_4 production and hydrogenotrophic pathway predominance were measured. To

do so we performed qPCR of the *mcrA* gene on sample DNA using the mlas-mcrA-rev

primer combination (Steinberg & Regan, 2008). Samples were run on an ABI

StepOnePlus thermocycler (ABI, USA), using Kapa SYBR reagents (Kapa Biosystems,

USA) according to manufacturer recommendations. For each sample, 8 ng DNA was

used, and the following amplification conditions were applied following

optimization: 98° C 10 minutes, 98° C 15 seconds, 55.6° 15 seconds, 72° 60 seconds.

A melt curve analysis was performed to verify target amplification. Not all samples

reliably amplified; in the end we obtained useable *mcrA* quantifications for 12 of the

18 samples. Although attempted, the transcriptional activity of *mcrA* from soil cDNA

did not produce reliable results, and thus is not included in this study. We used a

similar approach to quantify methanotroph abundance and transcriptional activity

by targeting the *pmoA* gene in the sample DNA and cDNA, respectively, using the

A189 – mb661 primer combination (Bourne, Donald, & Murrell, 2001). Reactions

were performed on a Bio-Rad CFX96 real-time qPCR instrument, using SsoAdvanced

Universal SYBR Green supermix (Bio-Rad, USA) following manufacturer

instructions. For each sample, 2 ng template were used with the following reaction

conditions: 98° C 10 minutes, 98° C 15 seconds, 55.6° 15 seconds, 72° 60 seconds.

All samples were amplified in triplicate. In both cases, sample amplification was

compared to a standard positive control to quantify total gene (or transcript) copy

number. We obtained useable *pmoA* quantifications for 42 of the 44 DNA samples

and 32 of the 44 RNA samples. In the case of *mcrA*, the positive control was an *mcrA*

plasmid and, in the case of *pmoA*, we used purified DNA from strain *Methylococcus*

capsulatus Foster and Davis (ATCC 33009D-5). We used LinRegPCR (Ramakers,

Ruijter, Deprez, & Moorman, 2003; Ruijter et al., 2009) to process amplification data

which allows for the calculation of individual PCR efficiencies. Individual PCR

efficiencies did not significantly differ between habitat types, so gene copy was

calculated using the average PCR efficiency of all reactions. Finally, gene copy (or

transcript) numbers were normalized to the total ng DNA (or cDNA) used in the

reaction.

Microbial community attribute calculations, statistical analysis, and

comprehensive modeling

All statistical analyses were performed using R (version 3.4.4) (R Core Team,

2018). Community matrices for the methanotrophy experiments were rarefied ten

times at 28,000 counts per sample, then averaged, and community matrices for the

methanogenesis experiments were rarefied to 30,400 counts per sample ten times

and averaged, in order to account for differences in sampling extent across samples.

Additionally, these rarefied community matrices were subsetted for known

methanogens/methanotrophs (Supplementary Table 1) to create functional group

community matrices. We categorized community data into the following four

attributes (Table 1): 1) *Abundance* of methanogens/methanotrophs using qPCR of

mcrA and *pmoA* genes (described above) respectively, as well as the relative

abundance of known methanogens/methanotrophs in the DNA-inferred 16S-based

prokaryotic community, 2) *Activity* of methanogens/methanotrophs using the

relative abundance of methanogens and methanotrophs in the RNA-inferred

prokaryotic community, and qPCR of *pmoA* using soil cDNA (described above), 3)

Composition of the broader prokaryotic community or of subsets of known

methanogen/methanotroph taxa from the 16S-inferred community matrix

(described below), and 4) *Diversity* of the prokaryotic community as well as

methanogen/methanotroph community subsets using ASV-level richness and

Shannon diversity. Species richness and Shannon diversity were calculated using

the 'vegan' package (Oksanen et al., 2015) in R. Regression plots were created using

the package 'ggplot2' (Wickham, 2009), and 95% confidence interval bands were

plotted around each linear fit using the 'geom_smooth' function (option = 'lm').

Difference in process rates among wetlands and uplands was assessed using a

Kruskal-Wallis test.

Associations between community attributes and CH_4-cycling processes

Because communities tend to have complex covariance structure, we took

several measures to avoid spurious associations when distinguishing biotic

correlates of CH_4-cycling processes. First, our process measurements were

performed under constant temperatures similar to those experienced in the field

(described above). By doing so, we minimized the influence of certain

environmental factors (*e.g.* temperature fluctuations or precipitation events) on our

process rate measurements. Secondly, we added two covariates to our model when

assessing associations with process rate measurements: 1) a soil physicochemical

covariate, and 2) and a sample (community) covariate. The soil physicochemical

covariate was calculated using principle components analysis (PCA) on soil

physicochemical measurements (described above). This was done using the prcomp

function in the 'stats' package of R (R Core Team, 2018), with variables scaled to

unit variance to account for differences in units of measurement. Sample scores for

the first principle component (PC1) were used as the environmental covariate. We

applied a similar approach to calculate the community covariate. In this case, the

rarefied community matrix was Hellinger transformed, then ordinated using the

prcomp function while scaling to unit variance, in order to calculate the community-

based sample PC1 scores. By including these covariates in our model we are

regressing each community attribute against the residual variation of the CH_4-

cycling process of interest after accounting for the variation explained by

environment and underlying sample structure.

To assess associations between community membership and process rate

variation we used the regression approach described above on the relative

abundance of individual taxa (ASVs). Since performing this type of analysis on

thousands of individual taxa increases the risk of false positives, we adjusted our

significance threshold (alpha = 0.05) using a Bonferroni correction, i.e. by dividing

by the number of of taxa being tested. In cases where significant taxa were

identified, those taxa were subsetted from the rarefied community matrix, then

ordinated (following the above procedure) to reduce them to a single variable. The

sample scores of the first principle component were subsequently used for

regression against CH_4-cycling process rates. Linear regressions were performed

using the lm function in the 'stats' package in R. Gross methanotrophy and

methanogenesis rate data were $\log_2$-transformed in order to satisfy assumptions of

normality. For each relationship we report the adjusted R^2 and *p* value.

RESULTS

Identifying biotic drivers of CH_4-cycling processes

Methanogenesis processes

Gross Methanogenesis

Gross methanogenesis rates varied from 0.06 to 6.7 µmol CH_4 g soil^{-1} d^{-1}

(average 1.1 ± 1.5 µmol CH_4 g soil^{-1} d^{-1}, Fig. 1A), with rates in mineral soil wetlands

on average 3.8 times higher than peatlands. For breakdown of soil physico-chemical

variables by habitat see Supplementary Table 2. We identified two community

attributes that were associated with gross methanogenesis rate after accounting for

sample and environmental covariance structure: methanogen abundance (*i.e.* the

abundance of *mcrA* genes, Adj. R^2 = 0.47, *p* = 0.008, Fig. 1B, Table 2), and the alpha

diversity of methanogens (Richness: Adj. R^2 = 0.25, *p* = 0.023, Shannon: Adj. R^2 =

0.23, *p* = 0.029). Both relationships were positive, suggesting that more abundant

and/or diverse populations of methanogens are associated with higher rates of

gross methanogenesis. We did not identify any individual prokaryotic taxa that were

significantly associated with gross methanogenesis after accounting for

environmental and sample covariance and adjusting for multiple comparisons. Soil

pH was the only significant abiotic predictor of methanogenesis (Adj. R^2 = 0.24, p =

0.023), showing a positive relationship whereby values approaching neutral pH had

higher rates of methanogenesis.

Hydrogenotrophic methanogenesis

445   The most pronounced difference in the predominance of the

hydrogenotrophic pathway was between mineral soil wetlands and peatlands,

where it accounted for 30.7 ± 11.2 % and 76.7 ± 23.0 % of total methanogenesis,

respectively (Fig. 1C). We identified a negative relationship between the DNA-

inferred alpha diversity of the methanogens and the predominance of the

hydrogenotrophic pathway (Shannon: Adj. R^2 = 0.710, p < 0.001 Fig. 1D, Table 2,

Richness: Adj. R^2 = 0.455, p = 0.002). We also identified a negative relationship with

the RNA-inferred alpha diversity of the prokaryotic community (Shannon: Adj. R^2 =

0.609, p = 0.001). The only abiotic variables that were significantly associated with

hydrogenotrophic methanogenesis were a negative relationship with soil pH (Adj.

R^2=0.32, p < 0.01), and a positive relationship with soil C (Adj. R^2=0.341, p < 0.01).

Methanotrophy processes

Gross methanotrophy

Average rates of gross CH_4 consumption (V_{max}) were higher in wetland sites

(including seasonal wetlands) (0.3 ± 0.05 µmol CH_4 g soil^{-1} d^{-1}) than upland sites

(0.1 ± 0.02 µmol CH_4 g soil^{-1} d^{-1}) (Fig. 2A, Kruskal-Wallis: Chi-squared = 6.85, df = 1,

$p = 0.008$), and significantly correlated with every abiotic variable measured. Gross

methanotrophy rates were negatively correlated with soil bulk density (Adj. R^2=

0.135, p =0.008). Note that the inverse of V_{max} is presented, i.e. higher uptake rates

are represented by higher values. Conversely gross methanotrophy rates were

positively correlated with soil moisture (Adj. R^2= 0.22, $p < 0.001$), soil C (Adj. R^2=

0.094, p =0.024), and soil N (Adj. R^2= 0.105, p =0.02). The only community attribute

that correlated with gross methanotrophy after accounting for variation due to soil

physicochemical variables was methanotroph abundance (i.e. *pmoA* copy number,

Adj. R^2=0.139, p=0.01, Fig. 2B, Table 2).

High-affinity methanotrophy

Of the four processes measured, high-affinity methanotrophy was the only

process to exhibit composition-based control involving the broader prokaryotic

community beyond known methanotrophs. Rates of high-affinity methane oxidation

varied considerably across our samples (range -9.15 to 2.08) (Fig. 2C) and were

higher in upland sites relative to wetland sites (including seasonal wetlands)

(Kruskal-Wallis: Chi-squared = 11.52, df = 1, $p < 0.001$). After accounting for sample

and environmental covariance, we identified 11 taxa in the DNA-inferred

community and 5 taxa in the RNA-inferred community whose relative abundance

significantly corresponded to high-affinity methanotrophy rates. The finest level of

taxonomic assignment for these taxa varied from order to genus. DNA-inferred taxa

identified included members of the genera *Nitrosphaera* (Thaumarchaeota),

Roseiarcus (Alphaproteobacteria), *Aminiphilus* (Synergistetes), and *Acidisoma*

(Alphaproteobacteria), a member of the family Thermosporaceae (Actinobacteria),

four members of Acidobacteria Group 2, and two members of Acidobacteria Group 6

(Supplementary Table 3). There was no ASV-level overlap with the RNA-inferred

taxa that were identified. The RNA-inferred taxa included members of the genera

Beijerinckia (Alphaproteobacteria), *Deferrisoma* (Deltaproteobacteria),

Nitrospirillum (Alphaproteobacteria), the family Thermonosporaceae

(Actinobacteria), and a member of Acidobacteria Group 1 (Supplementary Table 3).

We next ordinated the relative abundances of these taxa to reduce them to a

single variable. The first principle component of both of these community subsets

significantly correlated with high-affinity methanotrophy (RNA-inferred PC1 axis:

Adj. R^2 = 0.647, p < 0.001, Fig. 2D, Table 2, DNA-inferred PC1 axis: Adj. R^2 = 0.583, p

< 0.001). Lastly, we tested if high-affinity methanotrophy correlated with any other

community attributes. After accounting for covariance structure, there were no

other biotic correlates of high-affinity methanotrophy beyond composition-based

variables.

DISCUSSION

The utility of microbial community attributes in predicting function

The utility of conventional microbial community measurements for

predicting ecosystem functions has been recently called into question (Graham et

al., 2014; Inkpen et al., 2017; Louca et al., 2018; Rocca et al., 2014). Here we argue

that many of the difficulties associated with detecting community structure –

ecosystem function relationships could be overcome with careful consideration of

the community attributes chosen, the physiological scale of functional assessment,

and how environmental variation is controlled. In this study we have addressed

each of these concerns. Our work demonstrates 1) that there is a connection

between communities and ecosystem processes after accounting for variation due

to external factors, and 2) that the strength, as well as the nature, of the relationship

between communities and ecosystem function depends on the physiological scale of

the function.

Ecosystem function measurements face issues of physiological scale (Inkpen

et al., 2017), i.e. they represent the aggregate result of numerous underlying

processes. Detecting a causal link between community measurements and

ecosystem function thus becomes increasingly difficult for functions that include

inputs from multiple processes. This suggests that as one narrows the functional

scale to a level that more closely approximates individual physiologies, the strength

of the relationship with the community should increase. Our study supports this

hypothesis. For both methanogenesis- and methanotrophy-related processes, a

similar trend emerged; broader-scale functions had a weaker relationship with

community attributes than finer-scale functions. For instance, the best predictor of

gross methanogenesis was methanogen abundance (i.e. *mcrA* copy number),

explaining 47% of the variance, whereas hydrogenotrophic methanogensis – which

represents a finer physiological scale– showed the strongest association with

methanogen alpha diversity, explaining 71% of the variance (Table 2). For

methanotrophy processes, gross methanotrophy was best predicted by

methanotroph abundance (i.e. *pmoA* copy number), but this connection was quite

weak and only explained ~14% of the variance. By contrast, much more of the

variance in high-affinity methanotrophy could be explained by RNA-inferred

community composition (~64%, Table 2). While our specific hypothesis only

represents two processes for each type of function, our results provide support for

the idea that the physiological scale of function is an important consideration for

community structure-ecosystem function studies. Future work could extend the

generality of our findings by assessing an even wider assortment of physiological

scales (e.g. net CH_4 flux) or by extending this framework to other ecosystem

functions (e.g. soil respiration).

Developing community variables that represent putative process controls

542   Communities can control processes in several distinct ways. It is therefore

important that the variables we choose to represent communities reflect these types

of control. One key finding from our work was that broader scale processes had

distinctly different predictors than the underlying finer scale processes. Hence,

several of the relationships uncovered by our work would have gone undetected

had we focused solely on a single attribute. Both gross methanogenesis and

methanotrophy were best predicted by abundance-based variables (i.e. *mcrA* and

pmoA copy number, respectively), suggesting that these processes could be limited

by the total number of individuals capable of performing them. Similar relationships

between abundance and methanotrophy/methanogenesis rates have been reported

(Freitag & Prosser, 2009; Ma et al., 2012), although a number of studies have

demonstrated no such relationship (see Rocca et al., 2014) – suggesting other

important controls. Hydrogenotrophic methanogenesis and high-affinity

methanotrophy – the finer-scale processes – by contrast were best predicted by

diversity and composition measures, respectively. Hydrogenotrophic predominance

was inversely related to methanogen diversity. One possible explanation for this

relationship is that more diverse methanogen consortia would be more likely to

contain other methanogen functional groups (*e.g.* acetoclastic or methylotrophic

taxa), which would thus affect the relative contribution of the hydrogenotrophic

pathway to gross methanogenesis.

The strongest – and only – biotic variables associated with high-affinity

methanotrophy were based on RNA- and DNA-inferred composition (i.e. taxon

identity), suggesting that trait differences among taxa could be especially important

for this process. Interestingly, none of the associated taxa we identified were known

methanotrophs. This insight highlights the importance of not restricting our search

to taxa that are canonically associated with methanotrophy. While there are issues

with the commensurability of taxonomic composition and ecosystem function

(Inkpen et al., 2017), these taxa could still be causally linked to high-affinity

methanotrophy in at least two ways. First, at least some could be non-canonical

methanotrophs that are consuming CH_4. Even though the ability to consume CH_4 is

conserved phylogenetically (Martiny et al., 2013), canonical methanotrophs (as well

as high affinity methanotrophs) are nevertheless polyphyletic (Knief, 2015),

suggesting that methanotrophy has evolved multiple times or has been laterally

transferred. Another possibility is that these taxa influence methanotrophy through

ecological interactions with methanotrophs (e.g. competition or facilitation). A final

possibility is that these taxa are simply a marker for high-affinity methanotrophy,

co-varying with a variable that is causally linked. One of the DNA-inferred taxa

identified, *Nitrosphaera sp.* (Thaumarchaeota), is a known ammonia oxidizer.

Ammonia monooxygenase and methane monooxygenase are phylogenetically

related (Holmes, Costello, Lidstrom, & Murrell, 1995), and it has been suggested that

ammonia oxidizers may be able to bind CH_4 (Arp, Sayavedra-Soto, & Hommes, 2002;

Ross & Rosenzweig, 2017). Another taxon, *Roseiarcus* (Alphaproteobacteria), is

closely related to alphaproteobacterial methanotrophs and has been isolated from a

methanotrophic consortium enriched from *Sphagnum* peat (Kulichevskaya,

Danilova, Tereshina, Kevbrin, & Dedysh, 2014). Thus, *Roseiarcus* could be either

directly consuming CH_4, or interacting with methanotrophs. Two of the RNA-

inferred taxa, *Nitrospirillum* and *Beijerinckia* (both Alphaproteobacteria), are

diazotrophs and could thus influence methanotroph activity by fixing atmospheric

N_2. The influence of other correlated taxa on methanotrophy is less clear, but

competition for soil nutrients/O_2, facilitation, or simply acting as a marker remain

possibilities.

While we attempted to equally represent each attribute in our modeling,

activity-based attributes were under-represented in this study. Transcriptional

levels of *pmoA* and *mcrA* have been used as proxies for methanotroph/methanogen

activity in a variety of studies (Chen, Dumont, Cébron, & Murrell, 2007; Freitag &

Prosser, 2009; Freitag et al., 2010). While we successfully quantified *pmoA*

transcriptional activity, our attempt to do so for *mcrA* was problematic. This was not

likely due to a lack of activity of methanogens (since CH_4 was continuously produced

throughout the experiment), but rather due to PCR inhibitors such as humic

substances present in organic-rich soils. We took our community inference methods

a step further than many other studies by using both RNA- and DNA-based 16S

rRNA profiling. It has been established that the relative abundances of taxa in the

RNA-inferred community should not be used as a direct assessment of activity levels

(Papp, Mau, et al., 2018; Papp, Hungate, et al., 2018). However, it is likely that the

RNA-inferred community is at least enriched for the active fraction of soil taxa

relative to the DNA-inferred community (which could contain higher proportions of

inactive, dormant, or dead individuals) (Lennon & Jones, 2011). Thus, by including

RNA-based community inference we were able to use an additional window onto

the community that could have functional relevance.

Gaining a better understanding of tropical CH_4 dynamics

Despite the established importance of tropical ecosystems as global CH_4

sources (Kirschke et al., 2013; Sjögersten et al., 2014), our understanding of the

ecological controls over CH_4-cycling throughout these regions remains limited

(Bridgham et al., 2013). The majority of studies aimed at identifying controls on

CH_4-cycling dynamics have been performed in higher latitude ecosystems such as

northern peatlands (Bridgham et al., 2013). The contribution of biotic variables to

CH_4-cycling dynamics may be a more important determinant of variation across

tropical environments relative to higher latitudes. Methanogen population growth

rates are slow because all methanogenic pathways have very low thermodynamic

yields that are often barely above the threshold for growth under *in-situ* conditions

(Conrad, 1999; Megonigal, Mines, & Visscher, 2004). Thus, methanogens inhabiting

high latitude environments must recover each season from freeze-thaw events and

will exhibit low levels of activity/growth until warmer temperatures are reached. In

tropical ecosystems where prolonged freezing conditions are essentially absent,

trait-level distributions of CH_4-cycling taxa could play a stronger role regulating

spatial differences in CH_4 dynamics. While our study was not designed to compare

differences between high- and low-latitude ecosystems, an interesting trend that

emerged from our results was that abiotic variables tended to be less predictive of

CH_4 cycling relative to biotic variables (with the exception of gross methanotrophy).

Importantly, our laboratory incubations deliberately downplayed the influence of

several key environmental factors (*e.g.* temperature fluctuations or rainfall events)

that could influence CH_4-cycling rates. Nevertheless, most of our environmental

variables were independent from our incubation conditions (*e.g.* %C, %N, pH, soil

bulk density), and each only explained a relatively small fraction of the variance.

Thus, an important outstanding question is whether the strength and nature of

biotic controls on CH_4-cycling dynamics differs between high- and low-latitude

ecosystems.

Conclusion

How communities map onto ecosystem function is a widely debated topic.

Central to this debate are issues related to the physiological scale of function, the

identification of useful community attributes, and accounting for the underlying

influence of the environment. Our study provides two unique perspectives to this

debate. First, we suggest that broad definitions of function could be obscuring the

connections with community attributes. Secondly, by compiling community

attributes according to their putative role in controlling processes, we show that

finer scale processes may face controls that differ from those of the coarser

processes to which they contribute. Collectively, these findings demonstrate the

importance of carefully defining the physiological scale of ecosystem function and

performing community measurements that represent putative controls on

ecosystem processes.

ACKNOWLEDGEMENTS

We thank the Government of Gabon, Centre National de la Recherche

Scientifique et Technologique for permission (Permit No

AR0035/14/MESRS/CENAREST/CG/CST/CSAR) to conduct this study. We thank F.

Bivigou, H. Memiaghe, L. Tchignoumba, E. Tobi, I. Akendengue for their help

collecting samples. The University of Oregon, the Gabon-Oregon Transnational

Center on Environment and Development, the Smithsonian Conservation Biology

Institute, and Shell Gabon provided financial and logistical support. BB and KM are

grateful for support by the National Science Foundation – Dimensions of

Biodiversity program (DEB 14422214), and SB and AH are grateful for support by

A

B

C

D

689
Fig. 1: The rates and biotic drivers of methanogenesis-related processes across
Congo Basin wetlands. A) Methanogenesis rate (μmol CH_4 g soil^{-1} d^{-1}) across the two
wetland habitats. MSW = mineral soil wetland. B) The log$_2$ methanogenesis rate is
best predicted by log$_2$ methanogen abundance (qPCR of *mcrA* gene). C) The
predominance (%) of the hydrogenotrophic pathway in gross methanogenesis
across wetland habitats. D) Hydrogenotrophic pathway predominance is inversely
related to the alpha diversity (Shannon index) of the methanogen community. Each
point represents an individual soil sample. Gray bands represent 95% confidence
intervals of the linear model.
699

Fig. 2 The rates and biotic drivers of methanotrophy-related processes across
Congo Basin habitats. Note the inverse of V_{max} and k and presented so that higher
positive values correspond to higher rates of uptake. A) Gross methanotrophy rate
(-1* V_{max}) across habitats. MSW = mineral soil wetland, Seasonal. Wet for. =
seasonally inundated forest, Upland for. = upland forest, Ab. plantation = abandoned
plantation, Grassland w/ termite = grassland habitat dominated by termite mounds.
B) Gross methanotrophy rate is positively related to the $\log_2$ abundance of
methanotrophs (*pmoA* gene copy number). C) High-affinity methanotrophy rate
across habitats. D) High-affinity methanotrophy rates are best predicted by the first
principle component (PC1) of a subset of significantly associated taxa (Sig. taxa)
identified in the RNA-inferred prokaryotic community. Gray bands represent 95%
confidence intervals of the linear model.

715 **Table 1:** The community attributes calculated in this study used to identify putative
716 community controls on process rates.

Attribute	Variables
Abundance	qPCR (of *mcrA* & *pmoA* genes)
	Relative abundance (in DNA-inferred prokaryotic community) of methanogens/methanotrophs
Activity	cDNA qPCR (pmoA)
	Relative abundance (in RNA-inferred prokaryotic community) of methanogens/methanotrophs
Composition	Matrix of 16S-based prokaryotic community inferred from RNA
	Matrix of 16S-based prokaryotic community (DNA-inferred)
	Matrix of methanogen/methanotroph taxa subsetted from 16S-based matrix (DNA-inferred)
	Species (ASV) richness of RNA/DNA-inferred prokaryotic community
Diversity	ASV richness of methanogen/methanotroph community subsets (RNA & DNA)
	Shannon (ASV) diversity of RNA/DNA-inferred prokaryotic community
	Shannon diversity of methanogen/methanotroph community subsets (RNA & DNA)

728 **Table 2:** Narrow scale processes have a higher association with community
729 attributes. Process assessment scale: whether a process assessment is the aggregate
730 result of several processes (broad) or only one (fine). Adj. R^2 = the proportion of the
731 variance explained by the attribute.
732

	Physiological scale of process	Best fit community attribute	Adj. R^2	*P* value
Methanotrophy processes				
Gross Methanotrophy (V_{max})	Broad	Abundance (*pmoA* copy)	0.14	0.01
High affinity methanotrophy (k)	Fine	Composition (RNA-inferred)	0.64	< 0.001
Methanogenesis processes				
Gross Methanogenesis (μmol CH_4 g soil^{-1} d^{-1})	Broad	Abundance (*mcrA* copy)	0.47	0.008
Hydrogenotrophic methanogenesis (% gross CH_4 production)	Fine	Diversity (methanogens)	0.79	< 0.001

DATA ACCESSIBILITY

Sequence data, sample meta data, biogeochemical data, ASV community matrices

and R script for analysis will be made available upon acceptance of publication.

AUTHOR CONTRIBUTIONS

KM, AH, SB and BB designed and performed research. KM, AH, AK, and AM analyzed

data. KM wrote the paper with contributions from all authors.

REFERENCES

Arp, D. J., Sayavedra-Soto, L. A., & Hommes, N. G. (2002). Molecular biology and
biochemistry of ammonia oxidation by Nitrosomonas europaea. *Archives of*
Microbiology, *178*(4), 250–255. http://doi.org/10.1007/s00203-002-0452-0
Baldrian, P., Kolařík, M., Stursová, M., Kopecký, J., Valášková, V., Větrovský, T., …
Voříšková, J. (2012). Active and total microbial communities in forest soil are
largely different and highly stratified during decomposition. *The ISME Journal*,
6(2), 248–58. http://doi.org/10.1038/ismej.2011.95
Bannar-Martin, K. H., Kremer, C. T., Ernest, S. M., Leibold, M. A., Auge, H., Chase, J., …
Supp, S. R. (2018). Integrating community assembly and biodiversity to better
understand ecosystem function: the Community Assembly and the Functioning
of Ecosystems (CAFE) approach. *Ecology Letters*, *21*, 167–180.
http://doi.org/10.1111/ele.12895
Bier, R. L., Bernhardt, E. S., Boot, C. M., Graham, E. B., Hall, E. K., Lennon, J. T., …
Wallenstein, M. D. (2015). Linking microbial community structure and
microbial processes: An empirical and conceptual overview. *FEMS Microbiology*
Ecology, *91*(10), 1–11. http://doi.org/10.1093/femsec/fiv113
Bodelier, P. L. E., Meima-Franke, M., Hordijk, C. a, Steenbergh, A. K., Hefting, M. M.,
Bodrossy, L., … Seifert, J. (2013). Microbial minorities modulate methane
consumption through niche partitioning. *The ISME Journal*, *7*(11), 2214–28.
http://doi.org/10.1038/ismej.2013.99
Bodelier, P. L. E., Roslev, P., Henckel, T., & Frenzel, P. (2000). Stimulation by
ammonium-based fertilizers of methane oxidation in soil around rice roots.
Nature, *403*, 421–424.
Bourne, D. G., Donald, I. A. N. R. M. C., & Murrell, J. C. (2001). Comparison of pmoA
PCR Primer Sets as Tools for Investigating Methanotroph Diversity in Three
Danish Soils. *Applied and Environmental Microbiology*, *67*(9), 3802–3809.
http://doi.org/10.1128/AEM.67.9.3802
Bridgham, S. D., Cadillo-Quiroz, H., Keller, J. K., & Zhuang, Q. (2013). Methane
emissions from wetlands: Biogeochemical, microbial, and modeling
perspectives from local to global scales. *Global Change Biology*, *19*(5), 1325–
1346. http://doi.org/10.1111/gcb.12131
Bridgham, S. D., & Ye, R. (2013). Organic Matter Mineralization and Decomposition.
In R. D. DeLaune, K. R. Reddy, C. J. Richardson, & J. P. Megonigal (Eds.), *Methods*
in Biogeochemistry of Wetlands (pp. 385–406). Madison, WI: SSSA.
Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A., & Holmes, S. P.
(2016). DADA2: High-resolution sample inference from Illumina amplicon data.
Nature Methods, *13*, 581. Retrieved from
http://dx.doi.org/10.1038/nmeth.3869
Caporaso, J. G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F. D., Costello, E.
K., … Knight, R. (2010). QIIME allows analysis of high- throughput community
sequencing data Intensity normalization improves color calling in SOLiD
sequencing. *Nature Methods*, *7*(5), 335–336.
http://doi.org/10.1038/NMETH.F.303
Caporaso, J. G., Lauber, C. L., Walters, W. A., Berg-lyons, D., Lozupone, C. A.,

791 Turnbaugh, P. J., … Knight, R. (2011). Global patterns of 16S rRNA diversity at a
792  depth of millions of sequences per sample. *Proceedings of the National Academy*
793  *of Sciences*, *108*, 4516–4522. http://doi.org/10.1073/pnas.1000080107/-
794  /DCSupplemental.www.pnas.org/cgi/doi/10.1073/pnas.1000080107
795 Chen, Y., Dumont, M. G., Cébron, A., & Murrell, J. C. (2007). Identification of active
796  methanotrophs in a landfill cover soil through detection of expression of 16S
797  rRNA and functional genes. *Environmental Microbiology*, *9*(11), 2855–69.
798  http://doi.org/10.1111/j.1462-2920.2007.01401.x
799 Cole, J. R., Wang, Q., Fish, J. A., Chai, B., McGarrell, D. M., Sun, Y., … Tiedje, J. M. (2014).
800  Ribosomal Database Project: data and tools for high throughput rRNA analysis.
801  *Nucleic Acids Research*, *42*(D1), D633–D642.
802  http://doi.org/10.1093/nar/gkt1244
803 Conrad, R. (1999). Contribution of hydrogen to methane production and control of
804  hydrogen concentrations in methanogenic soils and sediments. *FEMS*
805  *Microbiology Ecology*, *28*(3), 193–202. http://doi.org/10.1016/S0168-
806  6496(98)00086-5
807 Delmas, R. A., Tathy, J. P., & Cros, B. (1992). Atmospheric methane budget in Africa.
808  *Journal of Atmospheric Chemistry*, *14*(1), 395–409.
809  http://doi.org/10.1007/BF00115247
810 Delmas, R., Servant, J., Tathy, J., Cros, B., & Labat, M. (1992). Sources and sinks of
811  methane and carbone dioxide exchanges in mountain forest in Equatorial
812  Africa. *Journal of Geophysical Research*, *97*(90), 6169–6179.
813 Fadrosh, D. W., Ma, B., Gajer, P., Sengamalay, N., Ott, S., Brotman, R. M., & Ravel, J.
814  (2014). An improved dual-indexing approach for multiplexed 16S rRNA gene
815  sequencing on the Illumina MiSeq platform. *Microbiome*, *2*(6), 1–7.
816 Freitag, T. E., & Prosser, J. I. (2009). Correlation of methane production and
817  functional gene transcriptional activity in a peat soil. *Applied and Environmental*
818  *Microbiology*, *75*(21), 6679–87. http://doi.org/10.1128/AEM.01021-09
819 Freitag, T. E., Toet, S., Ineson, P., & Prosser, J. I. (2010). Links between methane flux
820  and transcriptional activities of methanogens and methane oxidizers in a
821  blanket peat bog. *FEMS Microbiology Ecology*, *73*, 157–165.
822  http://doi.org/10.1111/j.1574-6941.2010.00871.x
823 Garcia, J. L., Patel, B. K., & Ollivier, B. (2000). Taxonomic, phylogenetic, and ecological
824  diversity of methanogenic Archaea. *Anaerobe*, *6*(4), 205–26.
825  http://doi.org/10.1006/anae.2000.0345
826 Gavlak, R., Horneck, D., Miller, R. O., & Kotuby-Amarcher, J. (2003). Soil, Plant, and
827  Water Reference Methods for the Western Region. *Western Region Extension*
828  *Publication*, *WREP-125*.
829 Graham, E. B., Wieder, W. R., Leff, J. W., Weintraub, S. R., Townsend, A. R., Cleveland,
830  C. C., … Nemergut, D. R. (2014). Do we need to understand microbial
831  communities to predict ecosystem function ? A comparison of statistical models
832  of nitrogen cycling processes. *Soil Biology and Biochemistry*, *68*, 279–282.
833  http://doi.org/10.1016/j.soilbio.2013.08.023
834 Ho, A., Kerckhof, F.-M., Luke, C., Reim, A., Krause, S., Boon, N., & Bodelier, P. L. E.
835  (2013). Conceptualizing functional traits and ecological characteristics of
836  methane-oxidizing bacteria as life strategies. *Environmental Microbiology*

837 *Reports*, *5*(3), 335–45. http://doi.org/10.1111/j.1758-2229.2012.00370.x

838 Holmes, A. J., Costello, A., Lidstrom, M. E., & Murrell, J. C. (1995). Evidence that
839  particulate methane monooxygenase and ammonia monooxygenase may be
840  evolutionarily related. *FEMS Microbiology Letters*, *132*, 203–208.

841 Inkpen, S. A., Douglas, G. M., Brunet, T. D. P., Leuschen, K., Doolittle, W. F., & Langille,
842  M. G. I. (2017). The coupling of taxonomy and function in microbiomes. *Biology*
843  *& Philosophy*. http://doi.org/10.1007/s10539-017-9602-2

844 Jones, S. E., & Lennon, J. T. (2010). Dormancy contributes to the maintenance of
845  microbial diversity. *Proceedings of the National Academy of Sciences of the*
846  *United States of America*, *107*(13), 5881–6.
847  http://doi.org/10.1073/pnas.0912765107

848 Kamke, J., Taylor, M. W., & Schmitt, S. (2010). Activity profiles for marine sponge-
849  associated bacteria obtained by 16S rRNA vs 16S rRNA gene comparisons. *The*
850  *ISME Journal*, *4*(4), 498–508. http://doi.org/10.1038/ismej.2009.143

851 Keller, J., & Bridgham, S. (2007). Pathways of anaerobic carbon cycling across an
852  ombrotrophic-minerotrophic peatland gradient. *Limnology and Oceanography*,
853  *52*(1), 96–107. http://doi.org/10.4319/lo.2007.52.1.0096

854 Kirschke, S., Bousquet, P., Ciais, P., Saunois, M., Canadell, J. G., Dlugokencky, E. J., …
855  Zeng, G. (2013). Three decades of global methane sources and sinks. *Nature*
856  *Geoscience*, *6*(10), 813–823. Retrieved from
857  http://dx.doi.org/10.1038/ngeo1955

858 Knief, C. (2015). Diversity and habitat preferences of cultivated and uncultivated
859  aerobic methanotrophic bacteria evaluated based on pmoA as molecular
860  marker. *Frontiers in Microbiology*, *6*, 1–38.
861  http://doi.org/10.3389/fmicb.2015.01346

862 Kolb, S. (2009). The quest for atmospheric methane oxidizers in forest soils.
863  *Environmental Microbiology Reports*, *1*(5), 336–346.
864  http://doi.org/10.1111/j.1758-2229.2009.00047.x

865 Kozich, J. J., Westcott, S. L., Baxter, N. T., Highlander, S. K., & Schloss, P. D. (2013).
866  Development of a dual-index sequencing strategy and curation pipeline for
867  analyzing amplicon sequence data on the MiSeq Illumina sequencing platform.
868  *Applied and Environmental Microbiology*, *79*(17), 5112–20.
869  http://doi.org/10.1128/AEM.01043-13

870 Kulichevskaya, I. S., Danilova, O. V, Tereshina, V. M., Kevbrin, V. V, & Dedysh, S. N.
871  (2014). Descriptions of Roseiarcus fermentans gen. nov., sp. nov., a
872  bacteriochlorophyll a-containing fermentative bacterium related
873  phylogenetically to alphaproteobacterial methanotrophs, and of the family
874  Roseiarcaceae fam. nov. *International Journal of Systematic and Evolutionary*
875  *Microbiology*, *64*(8), 2558–2565.
876  https://ijs.microbiologyresearch.org/content/journal/ijsem/10.1099/ijs.0.064
877  576-0

878 Lee, M. E., Alonso, A., Dallmeier, F., Campbell, P., & Pauwels, O. S. G. (2006). The
879  Gamba Complex of Protected Areas : An Illustration of Gabon's Biodiversity.
880  *Bulletin of the Biological Society of Washington*, (12), 229–242.

881 Legendre, P. (1993). Spatial autocorrelation: trouble or new paradigm? *Ecology*,
882  *74*(6), 1659–1673.

Lennon, J. T., & Jones, S. E. (2011). Microbial seed banks: the ecological and
evolutionary implications of dormancy. *Nature Reviews Microbiology*, *9*(2),
119–30. http://doi.org/10.1038/nrmicro2504
Loreau, M., Naeem, S., Inchausti, P., Bengtsson, J., Grime, J. P., Hector, A., ... Wardle, D.
A. (2001). Biodiversity and Ecosystem Functioning: Current Knowledge and
Future Challenges. *Science*, *294*(5543), 804 LP-808. Retrieved from
http://science.sciencemag.org/content/294/5543/804.abstract
Louca, S., Polz, M. F., Mazel, F., Albright, M. B. N., Huber, J. A., Connor, M. I. O., ...
Parfrey, L. W. (2018). Function and functional redundancy in microbial
systems. *Nature Ecology & Evolution*. http://doi.org/10.1038/s41559-018-
0519-1
Ma, K., Conrad, R., & Lu, Y. (2012). Responses of methanogen mcrA genes and their
transcripts to an alternate dry/wet cycle of paddy field soil. *Applied and
Environmental Microbiology*, *78*(2), 445–54.
http://doi.org/10.1128/AEM.06934-11
MacDonald, J. A., Jeeva, D., Eggleton, P., Davies, R., Bignell, D. E., Fowler, D., ...
Maryati, M. (1999). The effect of termite biomass and anthropogenic
disturbance on the CH4 budgets of tropical forests in Cameroon and Borneo.
Global Change Biology, *5*(8), 869–879.
Martiny, A. C., Treseder, K., & Pusch, G. (2013). Phylogenetic conservatism of
functional traits in microorganisms. *The ISME Journal*, *7*(4), 830–8.
http://doi.org/10.1038/ismej.2012.160
McCalley, C. K., Woodcroft, B. J., Hodgkins, S. B., Wehr, R. a., Kim, E.-H., Mondav, R., ...
Saleska, S. R. (2014). Methane dynamics regulated by microbial community
response to permafrost thaw. *Nature*, *514*(7523), 478–481.
http://doi.org/10.1038/nature13798
Megonigal, J. P., Mines, M. E., & Visscher, P. T. (2004). Anaerobic Metabolism:
Linkages to Trace Gases and Aerobic Processes. In *Biogeochemistry* (Vol. 8, pp.
350–362). Gulf Professional Publishing.
Nazaries, L., Murrell, J. C., Millard, P., Baggs, L., & Singh, B. K. (2013). Methane,
microbes and models: fundamental understanding of the soil methane cycle for
future predictions. *Environmental Microbiology*, *15*(9), 2395–417.
http://doi.org/10.1111/1462-2920.12149
Nazaries, L., Pan, Y., Bodrossy, L., Baggs, E. M., Millard, P., Murrell, J. C., & Singh, B. K.
(2013). Evidence of microbial regulation of biogeochemical cycles from a study
on methane flux and land use change. *Applied and Environmental Microbiology*,
79(13), 4031–40. http://doi.org/10.1128/AEM.00095-13
Oksanen, J., Blanchet, F. G., Roeland, K., Legendre, P., Minchin, P., O'Hara, R. B., ...
Wagner, H. (2015). vegan: Community ecology package. Retrieved from
http://cran.r-project.org
Papp, K., Hungate, B. A., & Schwartz, E. (2018). Microbial rRNA Synthesis and
Growth Compared through Quantitative Stable Isotope Probing with H218O.
Applied and Environmental Microbiology, *84*(8), 1–11.
Papp, K., Mau, R. L., Hayer, M., Koch, B. J., Hungate, B. A., & Schwartz, E. (2018).
Quantitative stable isotope probing with H218O reveals that most bacterial
taxa in soil synthesize new ribosomal RNA. *The ISME Journal*, 18–20.

http://doi.org/10.1038/s41396-018-0233-7

R Core Team. (2018). R: A language and environment for statistical computing.
Vienna, Austria: R Foundation for statistical computing. Retrieved from
http://cran.r-project.org

Ramakers, C., Ruijter, J. M., Deprez, R. H. L., & Moorman, A. F. . (2003). Assumption-
free analysis of quantitative real-time polymerase chain reaction (PCR) data.
Neuroscience Letters, *339*(1), 62–66. http://doi.org/10.1016/S0304-
3940(02)01423-4

Rocca, J. D., Hall, E. K., Lennon, J. T., Evans, S. E., Waldrop, M. P., Cotner, J. B., …
Wallenstein, M. D. (2014). Relationships between protein-encoding gene
abundance and corresponding process are commonly assumed yet rarely
observed. *The ISME Journal*, *9*(8), 1693–1699.
http://doi.org/10.1038/ismej.2014.252

Ross, M. O., & Rosenzweig, A. C. (2017). A tale of two methane monooxygenases.
Journal of Biological Inorganic Chemistry, *22*(2–3), 307–319.
http://doi.org/10.1007/s00775-016-1419-y

Ruijter, J. M., Ramakers, C., Hoogaars, W. M. H., Karlen, Y., Bakker, O., van den Hoff, M.
J. B., & Moorman, a F. M. (2009). Amplification efficiency: linking baseline and
bias in the analysis of quantitative PCR data. *Nucleic Acids Research*, *37*(6), e45.
http://doi.org/10.1093/nar/gkp045

Schimel, J. P. (1995). Ecosystem Consequences of Microbial Diversity and
Community Structure. In F. S. Chapin III & C. Körner (Eds.), *Arctic and Alpine*
Biodiversity: Patterns, Causes and Ecosystem Consequences (pp. 239–254).
Berlin: Springer-Verlag.

Schimel, J. P., & Gulledge, J. (1998). Microbial community structure and global trace
gases. *Global Change Biology*, *4*, 745–758.

Schmieder, R., & Edwards, R. (2011). Quality control and preprocessing of
metagenomic datasets. *Bioinformatics*, *27*(6), 863–864.
http://doi.org/10.1093/bioinformatics/btr026

Schnyder, E., Bodelier, P., Hartmann, M., Henneberger, R., & Niklaus, P. (2018).
Positive diversity-functioning relationships in model communities of
methanotrophic bacteria. *Ecology*, *99*(3), 714–723.
http://doi.org/10.1002/ecy.2138

Sierocinski, P., Bayer, F., Melia, G. Y., Großkopf, T., Alston, M., Swarbreck, D., …
Buckling, A. (2018). Biodiversity – function relationships in methanogenic
communities. *Molecular Ecology*, *27*, 4641–4651.
http://doi.org/10.1111/mec.14895

Sierocinski, P., Milferstedt, K., Bayer, F., Großkopf, T., Alston, M., Bastkowski, S., …
Buckling, A. (2017). A Single Community Dominates Structure and Function of a
Mixture of Multiple Methanogenic. *Current Biology*, 3390–3395.

Sjögersten, S., Black, C. R., Evers, S., Hoyos-Santillan, J., Wright, E. L., & Turner, B.
(2014). Tropical wetlands: A missing link in the global carbon cycle? *Global*
Biogeochemical Cycles, *28*, 1371–1386.
http://doi.org/10.1002/2014GB004844.Received

Steinberg, L. M., & Regan, J. M. (2008). Phylogenetic comparison of the methanogenic
communities from an acidic, oligotrophic fen and an anaerobic digester treating

municipal wastewater sludge. *Applied and Environmental Microbiology*, *74*(21),

6663–71. http://doi.org/10.1128/AEM.00553-08

Tathy, J. P., Cros, B., Delmas, R. A., Marenco, A., Servant, J., & Labat, M. (1992).

Methane emission from flooded forest in central Africa. *Journal of Geophysical*

Research: Atmospheres, *97*(D6), 6159–6168.

http://doi.org/10.1029/90JD02555

Tilman, D., & Downing, J. A. (1994). Biodiversity and stability in grasslands. *Nature*,

367, 363–365.

Tilman, D., Wedin, D., & Knops, J. (1996). Productivity and sustainability influenced

by biodiversity in grassland ecosystems. *Nature*, *379*, 718. Retrieved from

http://dx.doi.org/10.1038/379718a0

Wang, Q., Garrity, G. M., Tiedje, J. M., & Cole, J. R. (2007). Naive Bayesian Classifier for

Rapid Assignment of rRNA Sequences into the New Bacterial Taxonomy.

Applied and Environmental Microbiology, *73*(16), 5261–5267.

http://doi.org/10.1128/AEM.00062-07

White, F. (1979). The Guineo-Congolian Region and Its Relationships to Other

Phytochoria. *Bulletin Du Jardin Botanique National de Belgique / Bulletin van de*

National Plantentuin van Belgie, *49*(1/2), 11–55.

http://doi.org/10.2307/3667815

Wickham, H. (2009). *ggplot2: elegant graphics for data analysis*. Springer Science &

Business Media.

Zhang, J., Kobert, K., Flouri, T., & Stamatakis, A. (2014). PEAR: a fast and accurate

Illumina Paired-End reAd mergeR. *Bioinformatics*, *30*(5), 614–620.

http://doi.org/10.1093/bioinformatics/btt593