TITLE
Fungal communities living within leaves of native Hawaiian dicots are structured by landscape-scale variables as well as by host plants

RUNNING TITLE
Hawaiian leaf fungi at the landscape scale

AUTHORS
Darcy JL[1,2], Swift SOI[1], Cobian GM[1], Zahn G[3], Perry BA[4], Amend AS[1]

[1] Department of Botany, University of Hawaii, Manoa
[2] Division of Biomedical Informatics and Personalized Medicine, University of Colorado Anschutz Medical Campus
[3] Department of Biology, Utah Valley University
[4] Department of Biological Sciences, California State University East Bay

CORRESPONDING AUTHOR
John L. Darcy – +1 303 887 0477 – darcyj@colorado.edu

ABSTRACT

A phylogenetically diverse array of fungi live within healthy leaf tissue of dicotyledonous plants. Many studies

have examined these endophytes within a single plant species and/or at small spatial scales, but landscape-scale

variables that determine their community composition are not well understood, either across geographic space,

across climatic conditions, or in the context of host plant phylogeny. Here, we evaluate the contributions of these

variables to endophyte beta-diversity using a survey of foliar endophytic fungi in native Hawaiian dicots

sampled across the Hawaiian archipelago. We used Illumina technology to sequence fungal ITS1 amplicons to

characterize foliar endophyte communities across 5 islands and 80 host plant genera. We found that communities

of foliar endophytic fungi showed strong geographic structuring between distances of 7 and 36 kilometers.

Endophyte community structure was most strongly associated with host plant phylogeny and evapotranspiration,

and was also significantly associated with NDVI, elevation, and solar radiation. Additionally, our bipartite

network analysis revealed that the five islands we sampled each harbored significantly specialized endophyte

communities. These results demonstrate how the interaction of factors at large and small spatial and

phylogenetic scales shape fungal symbiont communities.

KEYWORDS
Fungi, Phylogeography, Endophytes, Hawai'i, High-throughput sequencing, Island Biogeography, Generalized Dissimilarity Modeling

INTRODUCTION

Foliar endophytic fungi (FEF), defined here as all fungi living within leaf tissue but not causing any outward

signs of disease (sensu (Stone, Bacon, & White, 2000)) are effectively invisible and represent a "hotspot" of

undescribed fungal diversity (Arnold & Lutzoni, 2007; Porras-Alfaro & Bayman, 2011). Indeed, a large portion

of the vast undescribed fungal diversity on earth (Meredith Blackwell, 2011) are presumed to live cryptic

lifestyles in association with plant hosts (Hawksworth & Rossman, 1997), particularly in the tropics where plant

diversity is highest. Due to their cryptic lifestyles and high species richness (M. Blackwell, Hibbett, Taylor, &

Spatafora, 2006; Rodriguez, White, Arnold, & Redman, 2009), many questions remain about which factors

determine how FEF are distributed throughout nature.

Although all available evidence suggests that most eudicot FEF are horizontally transmitted and not inherited via

seed (Bayman, Angulo-Sandoval, Báez-Ortiz, & Lodge, 1998), it is unclear which factors structure FEF

community composition, and how those factors rank in their relative importance. Previous studies that have

examined foliar fungal communities in this ecological and biogeographic context have noted several different

drivers of community composition and biogeography. Temperature (Coince et al., 2014), geographic distance

(U'Ren, Lutzoni, Miadlikowska, Laetsch, & Arnold, 2012), elevation and rainfall (Zimmerman & Vitousek,

2012), vegetation density/urbanization (Massimo et al., 2015; Unterseher, Petzold, & Schnittler, 2012; Vincent,

Weiblen, & May, 2016) are all putatively important variables for FEF community composition. We might

attribute this diversity of results to idiosyncrasies of experimental design in which specific variables or host taxa

were held constant to address focused hypotheses.

In fact, the bulk of FEF community research is represented by studies focusing on one or two specific host plants

to address hypotheses about how distance or climate impacts composition (Felber et al., 2016; González-Teuber,

2016; Kato, Fukasawa, & Seiwa, 2017; Mejía et al., 2014; Oono, Lefèvre, Simha, & Lutzoni, 2015; Polonio et

al., 2015; Saucedo-García, Anaya, Espinosa-García, & González, 2014; Zimmerman & Vitousek, 2012). Studies

that have surveyed FEF or other leaf-associated fungi among multiple plant species typically consider small

geographic scales in order to isolate host identity as a variable. Many have found that host identity strongly

determines fungal composition (Huang, Devan, U'Ren, Furr, & Arnold, 2016; Kembel & Mueller, 2014;

Massimo et al., 2015; U'Ren & Arnold, 2016; Unterseher et al., 2012; Vincent et al., 2016). Despite these

examples of studies at broad phylogenetic or spatial scales, comparatively few combine these into a single

analytical framework (but see (U'Ren et al., 2012, 2019)). By simultaneously examining a wide diversity of

hosts over a wide diversity of habitats we might explicitly address how host identity interacts with the abiotic

environment to structure FEF community composition.

Here, we use the breadth of environmental and evolutionary diversity in the Hawaiian Archipelago to determine

which factors structure FEF beta-diversity in native Hawaiian dicots. Specifically, we test the hypotheses that

elevation, rainfall, geographic distance, and host plant phylogeny are the strongest predictors of FEF community

composition of native plants across Hawaii (Figure 1). We include additional potential explanatory variables in

our analysis such as evapotranspiration (Kivlin, Hawkes, & Treseder, 2011; Lewis, Ravel, Naffaa, Astier, &

Charmet, 1997) and geographic distance (Higgins, Arnold, Coley, & Kursar, 2014), known to be important

determinants in other fungal systems. We found that FEF community structure was most strongly associated with

host plant phylogeny and evapotranspiration, and was also significantly associated with NDVI, elevation, and

solar radiation. Spatial structuring of FEF communities was strongest at distances between 7 and 36 kilometers.

MATERIALS AND METHODS

Sample Collection

General sampling locations were selected to maximize habitat, phylogenetic and spatial diversity. Location

selection prioritized access, a reasonable permitting process, and the known presence of ten or more native plant

species. Sample collection took place between July 2014 and October 2016. Access was the most limiting factor

in comparatively fewer sampling locations on Molokai and Maui Islands. We chose not to collect plants that are

federally listed as threatened or endangered. At each location, the first occurrence of an apparently healthy,

naturally recruited individual was selected for sampling. We only sampled native plants. Because some native

species are comparatively common, and others rare, we limited our sampling a single individual per location,

because we did not wish to confound our data with uneven spatial heterogeneity among sub-sampled plants. In

most cases more than a single individual was collected per island since it occurred at more than one location.

Numerous mature sun leaves were collected such that when combined, they covered a surface area roughly

equivalent to two adult-sized hands. Life form and stature differed markedly across our sampled plants, so it was

difficult to precisely standardize collection location within plants. When possible, leaves from trees and shrubs

were collected at eye level and from at least four aspects of the tree canopy. We targeted dicots because they

have horizontally transmitted FEF (compared to many vertically transmitted FEF in some monocots). Although

not all host plant genera were collected at every sample location, the three most common genera we collected

(*Metrosideros,* n=119; *Leptecophylla*, n=89; and *Vaccinium, n=*147) were each collected across elevations

ranging from sea level to over 2000 meters above sea level. The location of each plant was recorded with a GPS

and plants were positively identified in the field and/or vouchered for subsequent identification (vouchers

deposited at Joseph F. Rock Herbarium at the University of Hawaii, Manoa; HAW). Leaves were refrigerated

until subsequent processing (within 72 hours of collection). A total of 1099 samples were collected in this way,

although not all were used in the analyses presented here (see results section).

We surface sterilized leaves to exclude fungi present on leaf surfaces. After rinsing in water, forty leaf discs were

extracted per individual host by punching leaves with a sterile standard paper single hole punch (approximately

0.5 cm diameter). For plants with very small leaves, entire leaves were used such that the area of those leaves

was the same as the area of leaf discs. Leaf discs (or aforementioned collections of small leaves) were then

placed into loose-leaf tea bags that were subsequently stapled shut, submerged in 1% NaOCl for 2 minutes, then

70% EtOH for 2 minutes, followed by two rinses with sterile water for 2 minutes each. Rinse water was included

in extraction controls to verify sterility of surface water.

DNA Isolation

Ten leaf discs per DNA extraction were placed in MP Biomedical Lysing Matrix A tubes (MP Biomedical, Santa

Ana, CA, USA) containing DNA isolation solutions from the MoBio PowerPlant Pro DNA Isolation kit

(Solution PD1, Solution PD2, Phenolic Separation Solution, and RNase A Solution; MO Bio, Carlsbad, CA,

USA). Leaf discs were homogenized using a Mini-Beadbeater 24 (BioSpecs Inc. OK) at 3,000 oscillations per

min for 2 minutes. Lysate was centrifuged at 13,000 RPM for 2 minutes and transferred to individual wells of a

MoBio PowerPlant Pro DNA 96-well Isolation kit for subsequent extraction following the manufacturer's

protocol.

PCR Amplification and Illumina Library Preparation

We amplified the ITS1 region of the ribosomal cistron using fungal specific primers ITS1f and ITS2, along with

Illumina adaptors and Golay barcodes incorporating a dual indexing approach, using previously published

thermal cycling parameters (Smith & Peay, 2014). PCRs were carried out in 25 µl reactions using the KAPA3G

Plant PCR kit (KAPA Biosystems, Wilmington, MA, USA), 9 ul of DNA extraction (concentration not

measured) and 0.2 µM each of the forward and reverse primers. Negative PCR and extraction controls were

included. PCR products were purified and normalized using just-a-plate 96 PCR Purification and Normalization

Kit (Charm Biotech, San Diego, California, USA). Normalized PCR products were pooled and concentrated

using a streptavidin magnetic bead solution. Pooled PCR products were sequenced on three separate reactions in

order of processing completion, using the 2 x 300 paired-end (PE) sequencing protocol on an Illumina MiSeq

sequencing platform (Illumina Inc., Dan Diego, CA, USA). We did not include controls for sequencing run

variation, and we acknowledge this as a limitation of our study. Due to low sequencing success, negative

controls were not accounted for in sample processing.

DNA Sequence data processing and bioinformatics

QIIME (Caporaso et al., 2010) was used to demultiplex raw DNA sequence data into individual fastq files for

each sample. Although paired-end sequencing was used, only the R1 read (corresponding to primer ITS1f) was

used for downstream analysis, since sequencing quality of reverse reads was generally poor after the first 150

nucleotides. VSEARCH (Rognes, Flouri, Nichols, Quince, & Mahé, 2016) was used to discard reads with an

average quality score below 25 (Illumina Q+33 format), then ITSx (Bengtsson-Palme et al., 2013) was used to

extract the ITS1 region from quality-filtered files. Sequence data were filtered by average score rather than

maximum score, because nucleotides with poor quality scores that were not within the ITS1 region could

disqualify sequences that would be otherwise high-quality within the ITS1 region. In order to eliminate

variability in sequences caused by sequencing and PCR error, ITSx-processed sequences were de-noised using

DADA2 (Callahan et al., 2016) using the authors' recommended settings (maximum expected errors = 2,

maximum Ns = 0). Separate DADA2 error rates models were used for each sequencing run in our data set,

although the models appeared qualitatively similar. De-noised sequences were then clustered into 95% OTUs

using VSEARCH (Rognes et al., 2016).

Taxonomy was assigned to each OTU using the UNITE database (v 7) (Nilsson, 2011) and QIIME's

assign_taxonomy.py script (Caporaso et al., 2010) with the BLAST method using the default maximum e-value

of 0.001. OTUs not belonging to kingdom Fungi were discarded. Our OTU table was then rarefied (i.e.

randomly downsampled) to 1500 sequences per sample (Supplementary figure S1), discarding samples with

fewer than 1500 sequences or samples for which host plants could not be satisfactorily identified.

GhostTree (Fouquier et al., 2016) was used to construct a phylogenetic tree for the remaining OTUs. Briefly,

GhostTree allows phylogenetic trees to be made from ITS1 sequence data, which are often un-alignable at high

taxonomic ranks. This is done using a backbone tree created with the 18S rRNA gene, and then ITS1 sequences

are used to refine the tree at a phylogenetic scale where those sequences can be meaningfully aligned (e.g. genus

level). When used in an analysis of real and simulated ITS1 data, the GhostTree+UniFrac approach resulted in

higher amounts of variance explained than using phylogeny-independent metrics (Fouquier et al., 2016), and

this approach was also successfully used to study differences in fungal community composition in the human

infant microbiome (Ward et al., 2018) and in a study of FEF in grass as well (Lumibao et al., 2019). A new

GhostTree was made using the SILVA database (v 128) for the 18S backbone, and the UNITE database (v 7).

Tips of the GhostTree were renamed with OTU identifiers where OTUs were assigned taxonomy to a UNITE

entry in the GhostTree. In cases where multiple OTUs were assigned to the same UNITE entry, a polytomy was

created to fit those OTUs into the tree. This approach may exclude novel taxa that are not present in the UNITE

reference database, and this is a limitation of all studies using such a database, including this one.

The tree was used with weighted UniFrac (Lozupone & Knight, 2005) (hereafter referred to as "UniFrac") to

construct a beta-diversity matrix for the samples. UniFrac was used because it is downweighs additional branch

length contributed by slight intragenomic variation among tandem repeats. Even if an individual fungus contains

several OTUs, each will only contribute a negligible amount of branch length to a sample, as compared to non-

phylogenetic metrics (e.g. Bray-Curtis), which would consider those OTUs as different as any other pair.

Because UniFrac community dissimilarity considers the shared phylogenetic branch-lengths between two

communities, it is robust to the case where an OTU is only found within one sample, and is similarly robust to

the case where samples do not share any OTUs, which can be problematic for other community dissimilarity

metrics where overall diversity is much higher than within sample diversity. This is important for our analysis of

FEF communities, because many previous studies have shown that FEF are hyperdiverse even at local scales

(e.g. within one host or within one hundred meters) (Jumpponen & Jones, 2009, 2010; Rodriguez et al., 2009;

Zimmerman & Vitousek, 2012). We suspected that this large amount of diversity would result in many pairs of

samples that shared few or zero OTUs, which would result in an inflation of 1-values (maximum dissimilarity)

when using non-phylogenetic beta-diversity metrics such as Bray-Curtis or Jaccard community dissimilarity.

Indeed, this same issue has been discussed by previous users of GDM as a saturation of maximal dissimilarities,

and was solved by using phylogenetic dissimilarty metrics (Warren, Cardillo, Rosauer, & Bolnick, 2014). We

confirmed that this was indeed the case, and that UniFrac distance values were normally distributed but Bray-

Curtis dissimilarities were severely 1-inflated (Supplementary figure S2). For this reason, we used the UniFrac

beta-diversity matrix for the remainder of our analysis.

Spatiotemporal data

Using sample geographic coordinates and collection dates, environmental and climatic data for each sample

were extracted from GIS layers using the R packages 'raster' (Hijmans et al., 2014) and 'rgdal' (Pebesma,

Rowlingson, & Rogerbivandnhhno, 2012). Data were extracted from rasters generated for the same month that

samples were collected (except for elevational data). Supplementary table S1 shows the sources of each GIS

layer. Many explanatory variables were obtained from the Rainfall of Hawai'i and Evapotranspiration of Hawai'i

websites (T W Giambelluca et al., 2014; Thomas W. Giambelluca et al., 2013), and elevational data were

obtained using the USGS EarthExplorer online tool (http://earthexplorer.usgs.gov), courtesy of NASA EOSDIS

Land Processes Distributed Active Archive Center and the United States Geological Survey's Earth Resources

Observation and Science Center. These explanatory variables were chosen either because previous studies of

Hawaiian FEF had identified them as important (elevation, rainfall)(Zimmerman & Vitousek, 2012), or because

they were significant drivers of Hawaiian plant composition (slope, aspect), or because they made intuitive sense

in the context of fungi that live within leaves (solar radiation, transpiration, evapotranspiration, leaf area index,

NDVI). Slope and aspect of each sampling location were calculated from elevation raster data using the terrain

function in the 'raster' package (Hijmans et al., 2014). Data for aspect (the direction a sampling location faces)

were converted into a distance matrix using the smallest arc-difference between any two given aspects. This was

done because Euclidean distance is unsuitable for a measurement like aspect, where 355° is closer to 1° than it is

to 340°. All variables are mean monthly values. NDVI (normalized difference vegetation index), is an index

calculated from the amount of infra-red light reflected by plants, which is normalized using multiple

wavelengths of visible light. This allows for discrimination between habitat types that are differentially

vegetated. Similarly, leaf area index is a measure of surface area of leaves (one-sided) per unit area of ground,

and while it does not discriminate between different types of vegetation as does NDVI, unlike NDVI it measures

the density of leaf surface area (habitat for FEF).

Host plant phylogeny

A distance matrix of host plant phylogenetic distances was created using the angiosperm phylogeny of Qian and

Jin (Qian & Jin, 2016). This distance matrix was made because the modeling approach we use here (GDM,

below) can accommodate distance matrices as explanatory variables, allowing for a phylogenetic distance matrix

of hosts to be used instead of a simplified data structure such as a principal components vector or an array of

taxon identities. For each pair-wise comparison between two samples, pair-wise host plant phylogenetic distance

was calculated as the mean cophenetic (branch-length) distance between members of the plant genera that were

sampled, using the function cophenetic.phylo in the R package 'ape' (Paradis & Schliep, 2019). In cases where

host plant genus was not included in the phylogeny, the genus was substituted for the most closely-related genus

available. Four genera were substituted in this way out of 80 total genera: Labordia → Logania, Touchardia →

Urtica, Waltheria → Hermannia, Nothocestrum → Withania. We chose to resolve host plants at the genus level

because species-level taxonomy can be unstable for recent adaptive radiations, and many Hawaiian species-level

phylogenies are not resolved (Ziegler, 2002).

Geospatial analyses

A geographic semivariogram (Bachmaier & Backes, 2008) was constructed in R using a matrix of $\log_{10}$

transformed geographic distances between samples, and the UniFrac beta-diversity matrix of FEF community

differences between samples. Distance classes in the variogram were constructed using a sliding window

approach, where each class shared two-thirds of its points with the previous class (John L. Darcy, King,

Gendron, & Schmidt, 2017). A logistic curve was fit to semivariance values and mean geographic distances

within semivariogram distance classes using the Nelder-Mead algorithm, minimizing the sum of squared

differences in beta-diversity semivariance. The curve's derivative was then used to find the geographic distance

class that accounts for 99% of geographic distance where beta diversity is spatially autocorrelated. Mantel

correlations were then performed on data within each of the three resulting distance classes.

Generalized dissimilarity modeling (GDM) (Ferrier, Manion, Elith, & Richardson, 2007) was used to model FEF

beta diversity based on climatic factors (Supplementary table S1), geographic distance, host plant phylogeny,

and DNA sequencing run. GDM is a form of non-linear matrix regression that is well-suited to statistical

questions involving dissimilarity matrices (e.g. our beta-diversity matrix, host plant cophenetic distance matrix,

geographic distance matrix, and aspect arc-difference matrix). Unlike pair-wise Mantel tests or PerMANOVA,

which make use of similar data, GDM can quantify the relative importance of environmental and geographic

variables on community dissimilarity, even when the functional relationship between community dissimilarity

and the environment is nonlinear (Fitzpatrick et al., 2013; Warren et al., 2014). Furthermore, GDM is effective

because it models community dissimilarity associated with a given predictor variable while holding all other

predictor variables constant (Fitzpatrick et al., 2013; Landesman, Nelson, & Fitzpatrick, 2014). For example, this

enables GDM to model the effect of elevation while accounting for the effect of host plant phylogenetic distance.

Because our data were generated using three separate DNA sequencing runs, we included run as a categorical

variable in our model by creating a distance matrix where samples from the same run had distance 0, and

samples of different runs had distance 1, as recommended by GDM's creators (Ferrier et al., 2007).

We used backward elimination as implemented in the GDM R package to build a model, and then to simplify the

model by removing minimally predictive variables. We began this process with a GDM model excluding

variables that were correlated with another variable in the analysis at rho > 0.70 (Spearman). Leaf area index,

wet canopy evaporation, temperature, cloud frequency, and relative humidity were removed this way

(Supplementary figure S3). Log-transformed geographic distance between samples, host plant phylogenetic

distance, sequencing run, and aspect arc-difference matrix were included as predictor matrices in the GDM, with

FEF UniFrac distance as the response matrix. We then used GDM to test each variable within the model for

significance using a permutation test. During this iterative process, the variable with the highest P-value was

eliminated, and then the model was recalculated. This process was repeated until all remaining variables were

statistically significant ($P < 0.05$). Sequencing run and transpiration were eliminated this way. The final GDM

model was also analyzed using Bray-Curtis dissimilarities instead of UniFrac distances, to illustrate how the

severe 1-inflation observed with Bray-Curtis was an impediment to model fitting.

Principal coordinate analysis (PCoA) was used to validate GDM results using an alternate statistical approach.

PCoA vectors were calculated from the UniFrac distance matrix using the R package 'vegan' (Oksanen et al.,

2016). PCNM (Borcard & Legendre, 2002) was used on geographic and host phylogenetic distance matrices, for

dimensionality reduction and to account for nonlinear features within those data. The same data used in our final

GDM, except with geography and host represented by their first 6 PCNM vectors, were fit to FEF PCoA and

tested using the 'envfit' function (Oksanen et al., 2016).

Island FEF specificity analysis

Bipartite network analysis was used to test the extent to which each island (Figure 1) harbored specific FEF. The

d' ("d prime") statistic was calculated for each island using the OTU table using the Bipartite package in R

(Dormann et al., 2008). d' is a measure of network specialization that ranges from 0 to 1, where 0 is perfect

cosmopolitanism (all species are evenly shared among islands) and 1 is perfect specialization (each species is

specific to only one island). d' is calculated using a contingency matrix where each row is a unique lower-level

group (island) and each column is a unique higher-level group (OTU), but in our OTU table each island contains

multiple samples, so the number of observations per island is not consistent. To remedy this, we calculated d' by

aggregating all samples from the same island into one large sample (column sums), rarefied this aggregated table

using the same depth that samples were rarefied to above (1500 observations), then calculated d' values.

Empirical d' values were compared to null d' values generated with the Vázquez null model, which is a fixed-

connectance null model for bipartite networks (Vázquez et al., 2007), implemented as the vaznull function of the

R package 'bipartite' (Dormann, Gruber, & Fruend, 2008). The procedure of aggregating, calculating an

empirical d', and calculating a Vázquez null d' was repeated 1000 times in order to obtain bootstrapped

distributions of empirical and null d' values for each island. Statistical significance for d' values for each island

was tested using Welch's unequal variance t-test to compare empirical and island null d' values. This test was 2-

tailed, since d' could be significantly lower than the null distribution indicating cosmopolitanism, or significantly

higher indicating specificity.

A similar procedure was applied to host plant genera in place of fungal OTUs. Since host is only observed once

per sample, d' was not bootstrapped. This was done in order to evaluate the extent to which our selection of host

plants was significantly specialized to islands.

To evaluate whether island specificity of FEF could be attributed to island-specific plant selections, we also

tested FEF-island specificity under the constraint of a generically identical plant community (at genus level) for

each island. The same procedure for analyzing FEF-island specificity above was carried out, but during the

aggregation step of each bootstrap, random samples for each island belonging to a fixed plant community were

chosen to the exclusion of all others. This fixed community was defined by the plant identification overlap

between all five islands. All d' values calculated this way were calculated from the same plant community from

each island. Analysis of these d' values was calculated as above.

RESULTS

Sequence data

Samples were collected across a wide range of climatic conditions (Figure 2), representative of the distributions

of those conditions in Hawaiʻi. Our data set comprised 896 samples that passed quality-filtering and ITS1

extraction, consisting of 7002 OTUs. After non-fungal OTUs were discarded, samples were rarefied to 1500

sequences per sample, resulting in 3458 OTUs across 760 samples. Of those samples, 417 were from Hawai'i,

84 from Kaua'i, 56 from Maui, 72 from Moloka'i, and 131 from O'ahu. Mean richness (number of OTUs

observed) per sample was 27.3 with a standard deviation of 18.9 (Supplementary Figure S1). Geographic and

botanical information for each sample used in our analysis is available as Supplementary Figure S4 and is

available from our FigShare repository (see Data Accessibility section) . Of 27 sequencing negative controls

included in the study, 13 resulted in sequence data, with an average of 123 reads and 2.2 OTUs per sample. This

sequencing depth was less than 1% of that found in the average biological sample (12,860). Because we were

unable to trace the source of these contaminants (whether tag switching, cross contamination or reagent

contamination) and because they amplified so poorly compared to biological samples, we did not attempt to

adjust our data set to account for these.

Geospatial results

Our semivariogram analysis showed that FEF beta-diversity is spatially structured between 7.6 and 36.3

kilometers, where it reaches its autocorrelation distance (Figure 3). This plateau and subsequent noise in beta-

diversity semivariance occurs when FEF beta-diversity is no longer spatially autocorrelated (Robeson et al.,

2011). In our analysis, this occurs at geographic distances of 36.3 kilometers. Within the distance class where

beta diversity was spatially autocorrelated, r_M (Mantel correlation) was 0.16. In the other two distance classes, r_M

was closer to zero as expected (Figure 3).

After GDM backward elimination, many variables were left as statistically significant: All *P*-values were the

minimum possible *P*-value given that GDM's backward elimination was run using 50 permutations. This means

that GDM always predicted a smaller proportion of FEF beta-diversity when any given predictor variable was

permuted, except for sequencing run and transpiration which were eliminated this way. Host plant phylogenetic

distance and evapotranspiration explained the most compositional dissimilarity in FEF communities, as given by

their GDM coefficients (the maximum height of their splines) (Ferrier et al., 2007; Fitzpatrick et al., 2013) which

were 0.090 and 0.078, respectively (Figure 4). These values indicate the relative importances of variables in the

model. The shape of GDM *i*-splines reflects FEF community turnover over the range of differences in a predictor

variable, and both host and evapotranspiration were related to FEF beta-diversity over their entire ranges.

Sample collection date was related to FEF beta-diversity only at temporally proximate values, while solar

radiation and NDVI were related to FEF beta-diversity only at large differences (Figure 4). Although all

variables shown in Figure 4 are statistically significant and the whole model fit nicely to the data with relatively

symmetrical residuals (Figure 4, top), GDM was only able to predict 6% of FEF beta-diversity. In contrast to the

GDM fit using the UniFrac distance matrix, GDM run with Bray-Curtis dissimilarity fit poorly as a result of

severe 1-inflation of distances (Supplementarey figures S2 and S5).

Principal component analysis (PCoA) was used to confirm GDM results using an alternative statistical approach.

Similar to GDM, host (PCNM 3 of host phylogenetic distance matrix) had the strongest relationship with FEF

beta-diverity with an R^2 of 0.106 ($P = 0.001$), and evapotranspiration had the second strongest relationship with

an R^2 of 0.073 ($P = 0.001$).

Bipartite network analysis

Each of the 5 islands we sampled showed a statistically significant (P < 0.001, Welch's unequal variance t-test)

pattern of FEF specialization (Figure 5), since the d' values for each island were higher than those generated

using a null model (null d' values centered around d'=0.25 for each island). Specialization in this case means that

each island harbors more OTUs that are uniquely associated with that island than would be expected by chance.

We found that our sampling of host plant genera was significantly specialized to island as well (Supplementary

figure S6), however host-island d' values were not significantly related to OTU-island d' values (linear

regression, *P* = 0.45). We also analyzed FEF island specificity while keeping plant taxonomy identical across

islands by randomly drawing a pre-set plant community from each island in each permutation of the d' analysis.

This analysis was done to remove the effect of plant community on FEF specificity. We still found significant

specificity for all islands, although in this analysis all 5 islands had very similar d' values (Supplementary figure

S7).

Endophyte Taxonomic composition

At the Class level, FEF composition was consistent across islands, and dominated by Dothideomycetes and

Sordariomycetes (members of the Pezizomycotina)(Supplementary figure S8). 12.14% of endophyte OTUs were

unidentified at the class level (corresponding to 11.29% of rarefied sequence data), perhaps reflective of

Hawaii's isolation and comparative paucity of taxonomic research.

DISCUSSION

The most striking pattern we found in our analysis of FEF communities in native Hawaiian plants was that

evapotranspiration, a comparatively coarse measure of plant community traits at the landscape level, is

meaningfully predictive for the community composition of microscopic fungi living within plant leaves. These

results suggest that aside from environmental or host determinants, endophyte community composition results

from a microbial "neighborhood", in addition to the traits of individual hosts. Evapotranspiration was the

second-most important variable in our analysis in terms of community composition (Figure 4); and it was more

important in the model than elevation, which was measured at a much finer spatial resolution (Supplementary

table S1). This was supported by our principal coordinate analysis as well. This result makes the interpretation of

our hypothesis regarding the effect of climate on FEF community composition surprising, since previous studies

(Coince et al., 2014; Zimmerman & Vitousek, 2012) suggested that temperature, elevation, and rainfall would be

the most important factors structuring community composition instead, and evapotranspiration has not been

previously considered as an important variable for FEF. While elevation (tightly correlated with temperature),

host plant phylogeny, rainfall, and Julian date were significant explanatory variables in our GDM analysis, each

of their effects were smaller than that of evapotranspiration (Figure 4).

Very few studies measure fungal community response to evapotranspiration, and to our knowledge none yet

have included FEF. In a grass system spanning 15 European countries, the response of endophytic fungi to a

transpiration gradient was substantial (Lewis et al., 1997). Association between native FEF and transpiration has

also been observed in Theobroma cacao (Arnold & Engelbrecht, 2007). Evapotranspiration could also drive FEF

community composition by changing the leaf interior habitat, and thereby select for different FEF communities

at high vs. low evapotranspiration. Indeed, evapotranspiration is strongly correlated with the moisture content of

leaves (Lambers, Chapin, & Pons, 2008). Evapotranspiration encompasses both plant transpiration and the

evaporation of water from soil and other surfaces, and both soil water content and stomatal conductance affect

leaf interior moisture (Lambers et al., 2008; Tardieu, Lafarge, & Simonneau, 1996). In light of previous studies

suggesting a link between fungal endophytes and evapotranspiration (Arnold & Engelbrecht, 2007; Kivlin et al.,

2011; Lewis et al., 1997), our finding that evapotranspiration is a significant predictor of differences between

FEF communities makes sense, and future physiological experiments at the scale of individual leaves might

shine light on the nature of this interaction.

The association between FEF beta diversity and host plant phylogenetic distance was mostly expected, and it

was the most important predictor in our GDM model. This result could either mean that FEF, host plants, or both

exhibit a degree phylogenetic of niche conservatism (Wiens et al., 2010). For example, FEF community

preference may be phylogenetically conserved among closely related plant species, or perhaps host preference is

conserved among closely related FEF. While many previous studies of FEF have focused on single host species

or genera, those that have surveyed multiple hosts have found that host identity is commonly associated with

fungal endophyte community composition (U'Ren et al., 2012; Unterseher et al., 2012; Vincent et al., 2016).

Thus, it is not surprising that we found a significant host effect in both our GDM and principal coordinate

analyses. The GDM *i*-spline for host steadily increases across the whole range of host phylogenetic distance

(Figure 4), although the abrupt increase at low distances indicates rapid turnover in FEF community composition

with only small changes in host phylogenetic distance. Indeed, host plant phylogeny may have been a more

robust predictor of FEF community dissimilarity if our host plants had been classified to species level instead of

genus level, however species level phylogenies are not fully resolved for many Hawaiian lineages, particularly

those that have undergone rapid and recent adaptive radiations. Nevertheless, the significant association we

observed from genus-level all the way up to higher taxonomic ranks agrees with a study of broad-scale host

associations among tropical foliar fungal epiphytes, in which family level host taxonomy is most predictive

(Kembel & Mueller, 2014).

NDVI (normalized difference vegetation index) was the most robust predictor of FEF community dissimilarity in

our data set, suggesting that areas that are differentially vegetated harbor different communities of FEF. This

difference may be related to the total percent land cover of vegetation, which is a component of NDVI

(Purevdorj, Tateishi, Ishiyama, & Honda, 1998), or related to the type of plant cover, i.e. different plant

community compositions (Lunetta, Knight, Ediriwickrema, Lyon, & Worthy, 2006), which is also encompassed

within the NDVI metric. Indeed, foliar fungal communities potentially respond to both land cover and habitat

type in the host plant *Quercus macrocarpa* (Jumpponen & Jones, 2009, 2010). The pixel size for the NDVI data

we used was 250 m^2 (Supplementary table S1), meaning that the value at any given location is an aggregate

value for a large plant community. Thus, the 9.3% of FEF community dissimilarity that was significantly

explained by NDVI in our analysis is related to the density and composition neighboring plant communities and

not specifically the host individual, a dynamic suggested in others studies as well (Kato et al., 2017).

The other significant drivers of FEF community composition (Figure 4) were mostly expected, particularly

elevation which explained 6.1% of FEF community dissimilarity when all other variables in our analysis were

held constant. This is similar to a result reported by Zimmerman and Vitousek (2012), who used PerMANOVA

to test the effects of rainfall, elevation, and substrate age on FEF beta-diversity patterns in *Metrosideros*

polymorpha (O'hia) trees. They found that elevation explained roughly 17% of compositional dissimilarity

between FEF communities, varying slightly depending on which dissimilarity metric was used. That study also

took place in Hawai'i, and the area sampled overlaps partially with the area of Hawai'i island that we sampled

(Figure 1). Zimmerman and Vitousek (2012) also found a large effect of rainfall, which was much smaller (but

still significant) in our data set. Rainfall is also a significant driver of FEF community structure in grasses

(Giauque & Hawkes, 2013), although in a larger continental-scale analysis of cultured FEF isolates, rainfall was

not a strong predictor of FEF diversity (U'Ren et al., 2012). Solar radiation was also a significant variable in our

analysis, and indeed previous research has shown that light levels are important for FEF colonization, likely

because different intensities of light exposure can affect the chemical makeup of the phyllosphere (Bahnweg et

al., 2005).

Patterns of host specificity among FEF communities have been thoroughly documented (Cobian, Egan, &

Amend, 2019; Huang et al., 2016; Kembel & Mueller, 2014; Massimo et al., 2015; Unterseher et al., 2012;

Vincent et al., 2016). Thus, it is not surprising that phylogenetic differences among host plants was a statistically

significant predictor of FEF community dissimilarity in our analysis (Figure 4). Unlike most previous studies

that found host associations of FEF (although see (Davis & Shaw, 2008)), we used the phylogeny of host plants

as an explanatory variable in place of their identity, meaning that under our hypothesis that FEF communities are

structured by host phylogenetic difference, more phylogenetically similar plants are expected to harbor similar

FEF communities, and conversely, phylogenetically distant plants are expected to harbor less similar FEF

communities. This pattern of phylosymbiosis is widespread throughout natural symbiotic systems, although the

phylogenetic signal can be lost in the presence of environmental factors that simultaneously impact symbiotic

community composition (Brooks, Kohl, Brucker, van Opstal, & Bordenstein, 2016).

Geographic distance was of middling importance in our GDM analysis, especially since importance in GDM is

calculated with all other variables accounted for. However, in our semivariogram and subsequent Mantel

correlogram analyses, we found that FEF beta diversity is significantly spatially structured by geographic

distance, especially between geographic distances of 7.6 and 36.3 kilometers (Figure 3). Communities are not

strongly spatially autocorrelated at distance of less than 7.6 kilometers, and reach their autocorrelation range at

36.3 kilometers. The lag before 7.6 kilometers indicates that species are sampled from the same local community

before that point (Robeson et al., 2011). After the curve plateaus around 36.3 kilometers, FEF beta diversity

reaches its "autocorrelation range", after which autocorrelation becomes noisy. At distances greater than the

autocorrelation range, communities are not spatially autocorrelated; *i.e.* community composition is no longer a

function of geographic distance (King et al., 2010; Robeson et al., 2011). However, from our semivariogram we

can qualitatively see that there may be spatial structuring at greater distances as well, perhaps up to 100 km,

indicating perhaps that community structure is autocorrelated within islands but not between them. The nature of

Hawaii's volcanic origins might partially account for these patterns. Easterly trade winds and orographic lift

drive much of the precipitation patterns in the archipelago, such that each island is neatly divided between

comparatively hot and dry leeward, and cool wet windward sides. Nearly all island flora reflect this division and

it is unsurprising, therefore, to find spatial autocorrelation at within, rather than among, island scales.

However, even though beta-diversity may not be spatially autocorrelated at the inter-island scale, individual

species may still be unique to islands. Said another way, islands may harbor unique cohorts of FEF in addition to

cosmpolitan FEF, even though geographic space has little or no bearing on beta-diversity. We tested this, and

found a significant effect of FEF specificity to islands (Figure 5). Our bipartite network analysis showed that

each island harbors a significant composition of fungi that are more regionally specific than would be expected

due to chance alone. Beta-diversity approaches like GDM and our semivariogram are very different analytical

approaches than bipartite network analysis, and answer different questions. Our GDM analysis focused on

patterns in FEF beta-diversity, which is aggregate dissimilarity of community composition and does not take into

account patterns at the OTU level, especially because of UniFrac's down-weighting of phylogenetically similar

OTUs. Our bipartite network analysis instead focused on the distribution of FEF OTUs within versus among

islands, testing the extent to which a given island harbors OTUs that are exclusive to that island.

We presumed that the likeliest drivers of island specificity are differences in habitat, dispersal limitation among

islands and/or differences in host availability (whether due to plant distributions or sampling bias). Since our

beta-diversity analysis revealed a significant effect of host plant phylogeny, and because Hawaiian island floras

are disharmonious (Ziegler, 2002), and because this disharmony was reflected in our sampling, differences in

sampled plant hosts might account for specialization patterns. We used two separate approaches to evaluate

whether this could account for specialization patterns alone. A bipartite analysis of host-island specificity

(Supplementary figure S6) indicated that in our experimental design, hosts are indeed specific to islands.

However, we expected that if host-island specificity bias within our experimental design was driving the FEF-

island specificity pattern we observed, d' values from the two would correlate strongly. There was no significant

relationship between host-island specificity and FEF-island specificity, and host-island d' values were much

lower than those observed for OTUs. Further, when we restricted our analyses to only those plant genera that

were sampled on all islands, the FEF specialization patterns remained.

How dispersal limitation shapes island specificity is an open question worthy of future research. There is some

evidence for restricted gene flow among populations of a putative native mushroom, *Rhodocollybia laulaha*

(Keirle, Avis, Feldheim, Hemmes, & Mueller, 2011) among islands, although these patterns might also be

explained by selection. Similarly, the lag in our semivariogram (Figure 3) could be the distance at which

dispersal limitation begins to structure FEF communities, or simply the distance at which un-measured

environmental variables begin to vary enough to structure FEF communities. A long-term study of aerobiota at

the summit of Hawaii's peak, Mauna Loa, found evidence of likely long-distance dispersal via air mass (Tipton

et al., 2019), although it is unclear how frequent and ecologically relevant such events are to endophyte systems.

The epidemiology and rapid inter-island spread of fungal pathogens such as rapid Ohia death (*Ceratocystis* spp.)

or Ohia rust (*Puccinia psidii*) suggest that at least some species are capable of crossing ocean channels, whether

naturally or with the unwitting help of humans. Of course not all species are so easily dispersed, and the requisite

and complex interaction between FEF, host and habitat likely reinforce island specialization patterns.

In summary, our analysis highlights that the various factors contributing to Hawaiian FEF community structure

do so at the landscape scale. We analyzed these important plant symbionts across both a large geographic scale

(Figure 1) and across a large host phylogenetic scale (80 plant genera), while using high-throughput sequencing

to thoroughly inventory FEF community composition. We examined the effects of climate, geographic distance,

and host identity using this system, and for the most part, found them to be strong predictors of differences in

FEF communities between samples – even when the measurements were taken at relatively coarse resolution

(Supplementary table S1). We found that elevation (Zimmerman & Vitousek, 2012), host plant phylogenetic

difference (Cobian et al., 2019; Huang et al., 2016; Kembel & Mueller, 2014; Massimo et al., 2015; Unterseher

et al., 2012), geographic distance (U'Ren et al., 2012), and habitat type (Jumpponen & Jones, 2009, 2010; Kato

et al., 2017) hypotheses were robust in our analysis at these large scales.

AUTHOR CONTRIBUTIONS

JLD wrote the paper, performed bioinformatic and statistical analyses, and created figures. ASA and BAP

conceived of the experiment. SOS, GC, and GLZ collected data and performed laboratory work. All authors

contributed to the preparation of the manuscript.

ACKNOWLEDGEMENTS

This work was supported by NSF award #1255972 to Amend, NSF award #1256128 to Perry and an NSF GRFP

award to Cobian. The authors thank D Hemmes, J Adams, V Costello, E Datlof, S Walsh, K Kimball, M Purell, T

Flynn, A Williams, R O'Rorke, S Perlman, P Biley and K Coelho for help in the field, and L Tooman for

assistance in the lab. The authors additionally thank C Lozupone and J Foquier for helpful discussions. The

authors declare that no conflict of interest exists.

FIGURES
557
558

560
Figure 1: Map of sample locations across the Hawaiian Archipelago. Samples were collected from the 5
major islands in the Hawaiian archipelago: Hawaiʻi, Maui, Molokaʻi, Oʻahu, and Kauaʻi. Sampling was most
dense on Hawaiʻi and on Oʻahu islands, where accessibility was easiest. Our sampling strategy was to use
elevational transects where possible, in order to capture elevational and climatic variation. Elevations in
figure legend are meters above sea level. This is visible in the transects on Hawaiʻi, Molokaʻi, Kauaʻi, and
Oʻahu, which run orthogonally to the topographic lines (white). Transects are less pronounced on Maui
because of limited accessibility.
568

570
571 Figure 2: Ranges and distributions of explanatory variables. In this figure, each variable's distribution across 722
572 samples is shown as a smoothed density curve between its range (numbers to left and right of curves, height of
573 curve is relative frequency of observation). Units for range values are shown in Supplementary table S1. Each
574 variable included in our analysis covered a wide range of environmental heterogeneity.
575
576

Figure 3: Semivariogram and Mantel correlations of FEF beta diversity. This semivariogram (Bachmaier & Backes, 2008) was constructed using geographic distance classes, each containing the same number of points and each sharing two-thirds of its points with the previous class. Dots indicate the mean geographic distance within each class, and horizontal lines indicate its geographic range. The black curve is a logistic function fit to the data. Increasing semivariance indicates spatial autocorrelation. The lag in spatial autocorrelation before 7.6 kilometers (first dashed line) indicates a lack of spatial structuring, and that samples at that range may be drawn from the same community. The plateau and subsequent noise at 36.3 kilometers (second dashed line) is called the "autocorrelation distance", and is a typical feature of semivariogram analysis (King et al., 2010; Robeson et al., 2011). It indicates the distance after which spatial structuring becomes noisy or random, and geographic space is no longer a meaningful variable for community structure. The plot at bottom shows the data used to create the semivariogram. It also shows the Mantel correlation (r_M) of the three distance classes, as determined by 99% intervals of the logistic fit's derivative.

Figure 4: Model fit and coefficients for GDM model of FEF community dissimilarity. The observed community dissimilarity (UniFrac distance) between pairwise samples exhibited a linear but noisy relationship with the community dissimilarity predicted by the GDM model (top), which roughly corresponded to a 1:1 line (dashed line). In the bottom plot, GDM i-splines are shown for statistically significant predictor variables. Spline height indicates the relative importances of predictor variables, and the spline's slope corresponds to the rate of change in compositional dissimilarity over the range of pairwise dissimilarities within the variable. Host phylogeny and Evapotranspiration were the strongest predictor variables in our analysis. These two variables were also the strongest correlates of beta diversity principal coordinate vectors in our confirmatory PCoA analysis.

Figure 5: FEF specialization of each island. *d'* is a measure of how unique or cosmopolitan are the OTUs
found on that island. In this violin plot, distributions shown in as filled violins with black borders are
bootstrapped empirical *d'* values for each island. Unfilled distributions with a solid gray border are from
the Vázquez null model for bipartite networks, which is a fixed-connectance null model for bipartite
networks. Welch's unequal variance *t*-tests show that each island's FEF community is significantly
specialized as compared to the island null, meaning that FEF OTUs are more specific to their island of
origin than expected by chance.

REFERENCES
612
613
614 Arnold, A. E., & Engelbrecht, B. M. J. (2007). Fungal endophytes nearly double minimum leaf conductance
615   in seedlings of a neotropical tree species. *Journal of Tropical Ecology*, *23*(3), 369–372. doi:
616   10.1017/S0266467407004038

617 Arnold, A. E., & Lutzoni, F. (2007). Diversity and host range of foliar fungal endophytes: Are tropical leaves
618   biodiversity hotspots? *Ecology*, *88*(3), 541–549. doi: 10.1890/05-1459

619 Bachmaier, M., & Backes, M. (2008). Variogram or semivariogram? Understanding the variances in a
620   variogram. *Precision Agriculture*, *9*(3), 173–175. doi: 10.1007/s11119-008-9056-2

621 Bahnweg, G., Heller, W., Stich, S., Knappe, C., Betz, G., Heerdt, C., … Sandermann, H. (2005). Beech leaf
622   colonization by the endophyte Apiognomonia errabunda dramatically depends on light exposure and
623   climatic conditions. *Plant Biology*, *7*(6), 659–669. doi: 10.1055/s-2005-872943

624 Bayman, P., Angulo-Sandoval, P., Báez-Ortiz, Z., & Lodge, D. J. (1998). Distribution and dispersal of Xylaria
625   endophytes in two tree species in Puerto Rico. *Mycological Research*, *102*(8), 944–948. doi:
626   10.1017/S095375629700590X

627 Bengtsson-Palme, J., Ryberg, M., Hartmann, M., Branco, S., Wang, Z., Godhe, A., … Nilsson, R. H. (2013).
628   Improved software detection and extraction of ITS1 and ITS2 from ribosomal ITS sequences of fungi
629   and other eukaryotes for analysis of environmental sequencing data. *Methods in Ecology and*
630   *Evolution*, *4*(10), 914–919. doi: 10.1111/2041-210X.12073

631 Blackwell, M., Hibbett, D. S., Taylor, J. W., & Spatafora, J. W. (2006). Research Coordination Networks: a
632   phylogeny for kingdom Fungi (Deep Hypha). *Mycologia*, *98*(6), 829–837. doi:
633   10.3852/mycologia.98.6.829

634 Blackwell, Meredith. (2011). The fungi: 1, 2, 3 … 5.1 million species? *American Journal of Botany*, *98*(3),
635   426–438. doi: 10.3732/ajb.1000298

636 Borcard, D., & Legendre, P. (2002). All-scale spatial analysis of ecological data by means of principal
637   coordinates of neighbour matrices. *Ecological Modelling*. doi: 10.1016/S0304-3800(01)00501-4

638 Brooks, A. W., Kohl, K. D., Brucker, R. M., van Opstal, E. J., & Bordenstein, S. R. (2016). Phylosymbiosis:
639   Relationships and Functional Effects of Microbial Communities across Host Evolutionary History.
640   *PLoS Biology*, *14*(11), e2000225. doi: 10.1371/journal.pbio.2000225

641 Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A., & Holmes, S. P. (2016). DADA2: High-
642   resolution sample inference from Illumina amplicon data. *Nature Methods*, *13*, 581–583. doi:
643   10.1038/nmeth.3869

644 Caporaso, J. G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F. D., Costello, E. K., … Knight, R. (2010).
645   QIIME allows analysis of high-throughput community sequencing data. *Nature Methods*, *7*(5), 335–
646   336. doi: 10.1038/nmeth.f.303

Cobian, G. M., Egan, C. P., & Amend, A. S. (2019). Plant-microbe specificity varies as a function of elevation.
The ISME Journal, (13), 2778–2778.

Coince, A., Cordier, T., Lengell, J., Defossez, E., Vacher, C., Robin, C., … Marais, B. (2014). Leaf and root-
associated fungal assemblages do not follow similar elevational diversity patterns. *PLoS ONE*, *9*(6),
e100668. doi: 10.1371/journal.pone.0100668

[dataset] Darcy, J.L., Swift, S. O. I., Cobian, G. M., Zahn, G., Perry, B. A., & Amend, A. S. (2017). Foliar
endophytic fungi of native Hawaiian plants. *NCBI SRA*, PRJNA470970. Retrieved from
https://www.ncbi.nlm.nih.gov/bioproject/PRJNA470970/

Darcy, John L., King, A. J., Gendron, E. M. S., & Schmidt, S. K. (2017). Spatial autocorrelation of microbial
communities atop a debris-covered glacier is evidence of a supraglacial chronosequence. *FEMS*
Microbiology Ecology, *93*(8), 7287–7290. doi: 10.1093/femsec/fix095

Davis, E. C., & Shaw, A. J. (2008). Biogeographic and phylogenetic patterns in diversity of liverwort-
associated endophytes. *American Journal of Botany*, *95*(8), 914–924. doi: 10.3732/ajb.2006463

Dormann, C. F., Gruber, B., & Fruend, J. (2008). Introducing the bipartite Package: Analysing Ecological
Networks. *R News*, *8*(2), 8–11.

Felber, A. C., Orlandelli, R. C., Rhoden, S. A., Garcia, A., Costa, A. T., Azevedo, J. L., & Pamphile, J. A. (2016).
Bioprospecting foliar endophytic fungi of Vitis labrusca Linnaeus, Bordô and Concord cv. *Annals of*
Microbiology, *66*(2), 765–775. doi: 10.1007/s13213-015-1162-6

Ferrier, S., Manion, G., Elith, J., & Richardson, K. (2007). Using generalized dissimilarity modelling to
analyse and predict patterns of beta diversity in regional biodiversity assessment. *Diversity and*
Distributions, *13*(3), 252–264. doi: 10.1111/j.1472-4642.2007.00341.x

Fitzpatrick, M. C., Sanders, N. J., Normand, S., Svenning, J. C., Ferrier, S., Gove, A. D., & Dunn, R. R. (2013).
Environmental and historical imprints on beta diversity: Insights from variation in rates of species
turnover along gradients. *Proceedings of the Royal Society B: Biological Sciences*, *280*(1768). doi:
10.1098/rspb.2013.1201

Fouquier, J., Rideout, J. R., Bolyen, E., Chase, J., Shiffer, A., McDonald, D., … Kelley, S. T. (2016). Ghost-tree:
Creating hybrid-gene phylogenetic trees for diversity analyses. *Microbiome*, *4*(11), 1–10. doi:
10.1186/s40168-016-0153-6

Giambelluca, T W, Shuai, X., Barnes, M. L., Alliss, R. J., Longman, R. J., Miura, T., … Businger, A. D. (2014).
Evapotranspiration of Hawai'i. Final report submitted to the U.S. Army Corps of Engineers—Honolulu
District, and the Commission on Water Resource Management, State of Hawaii. Retrieved from
http://evapotranspiration.geography.hawaii.edu/

Giambelluca, Thomas W., Chen, Q., Frazier, A. G., Price, J. P., Chen, Y. L., Chu, P. S., … Delparte, D. M. (2013).
Online rainfall atlas of Hawai'i. *Bulletin of the American Meteorological Society*, *94*(3), 313–316. doi:
10.1175/BAMS-D-11-00228.1

Giauque, H., & Hawkes, C. V. (2013). Climate affects symbiotic fungal endophyte diversity and performance. *American Journal of Botany*, *100*(7), 1435–1444. doi: 10.3732/ajb.1200568

González-Teuber, M. (2016). The defensive role of foliar endophytic fungi for a South American tree. *AoB Plants*, *8*, plw050. doi: 10.1093/aobpla/plw050

Hawksworth, D. L., & Rossman, A. Y. (1997). Where Are All the Undescribed Fungi? *Phytopathology*, *87*(9), 888–891. doi: 10.1094/PHYTO.1997.87.9.888

Higgins, K. L., Arnold, A. E., Coley, P. D., & Kursar, T. A. (2014). Communities of fungal endophytes in tropical forest grasses: Highly diverse host- and habitat generalists characterized by strong spatial structure. *Fungal Ecology*, *8*(1), 1–11. doi: 10.1016/j.funeco.2013.12.005

Hijmans, R. J., Etten, J. van, Mattiuzzi, M., Sumner, M., Greenberg, J. A., Lamigueiro, O. P., … Shortridge, A. (2014). Package "raster." *R*, (2008), 1–27. doi: 10.1103/PhysRevE.74.016110.

Huang, Y.-L., Devan, M. M. N., U'Ren, J. M., Furr, S. H., & Arnold, A. E. (2016). Pervasive Effects of Wildfire on Foliar Endophyte Communities in Montane Forest Trees. *Microbial Ecology*, *71*(2), 452–468. doi: 10.1007/s00248-015-0664-x

Jumpponen, A., & Jones, K. L. (2009). Massively parallel 454 sequencing indicates hyperdiverse fungal communities in temperate *Quercus macrocarpa* phyllosphere. *New Phytologist*, *184*(2), 438–448. doi: 10.1111/j.1469-8137.2009.02990.x

Jumpponen, A., & Jones, K. L. (2010). Seasonally dynamic fungal communities in the Quercus macrocarpa phyllosphere differ between urban and nonurban environments. *New Phytologist*, *186*(2), 496–513. doi: 10.1111/j.1469-8137.2010.03197.x

Kato, S., Fukasawa, Y., & Seiwa, K. (2017). Canopy tree species and openness affect foliar endophytic fungal communities of understory seedlings. *Ecological Research*, *32*(2), 157–162. doi: 10.1007/s11284-016-1426-z

Keirle, M. R., Avis, P. G., Feldheim, K. A., Hemmes, D. E., & Mueller, G. M. (2011). Investigating the allelic evolution of an imperfect microsatellite locus in the Hawaiian mushroom rhodocollybia laulaha. *Journal of Heredity*. doi: 10.1093/jhered/esr099

Kembel, S. W., & Mueller, R. C. (2014). Plant traits and taxonomy drive host associations in tropical phyllosphere fungal communities. *Botany*, *92*(4), 303–311. doi: 10.1139/cjb-2013-0194

King, A. J., Freeman, K. R., McCormick, K. F., Lynch, R. C., Lozupone, C., Knight, R., & Schmidt, S. K. (2010). Biogeography and habitat modelling of high-alpine bacteria. *Nature Communications*, *1*(5), 53. doi: 10.1038/ncomms1055

Kivlin, S. N., Hawkes, C. V., & Treseder, K. K. (2011). Global diversity and distribution of arbuscular mycorrhizal fungi. *Soil Biology and Biochemistry*, *43*(11), 2294–2303. doi: 10.1016/j.soilbio.2011.07.012

Lambers, H., Chapin, F. S., & Pons, T. L. (2008). Plant Water Relations. In *Plant Physiological Ecology* (pp. 163–223). doi: 10.1007/978-0-387-78341-3_5

Landesman, W. J., Nelson, D. M., & Fitzpatrick, M. C. (2014). Soil properties and tree species drive ß-
diversity of soil bacterial communities. *Soil Biology and Biochemistry*, *76*, 201–209. doi:
10.1016/j.soilbio.2014.05.025

Lewis, G. C., Ravel, C., Naffaa, W., Astier, C., & Charmet, G. (1997). Occurrence of Acremonium endophytes in
wild populations of Lolium spp. in European countries and a relationship between level of infection
and climate in France. *Annals of Applied Biology*, *130*(2), 227–238. doi: 10.1111/j.1744-
7348.1997.tb06828.x

Lozupone, C., & Knight, R. (2005). UniFrac : a New Phylogenetic Method for Comparing Microbial
Communities UniFrac : a New Phylogenetic Method for Comparing Microbial Communities. *Applied
and Environmental Microbiology*, *71*(12), 8228–8235. doi: 10.1128/AEM.71.12.8228

Lumibao, C. Y., Borer, E. T., Condon, B., Kinkel, L., May, G., & Seabloom, E. W. (2019). Site-specific responses
of foliar fungal microbiomes to nutrient addition and herbivory at different spatial scales. *Ecology
and Evolution*. doi: 10.1002/ece3.5711

Lunetta, R. S., Knight, J. F., Ediriwickrema, J., Lyon, J. G., & Worthy, L. D. (2006). Land-cover change detection
using multi-temporal MODIS NDVI data. *Remote Sensing of Environment*, *105*(2), 142–154. doi:
10.1016/j.rse.2006.06.018

Massimo, N. C., Nandi Devan, M. M., Arendt, K. R., Wilch, M. H., Riddle, J. M., Furr, S. H., … Arnold, A. E.
(2015). Fungal endophytes in aboveground tissues of desert plants: infrequent in culture, but highly
diverse and distinctive symbionts. *Microbial Ecology*, *70*(1), 61–76. doi: 10.1007/s00248-014-0563-6

Mejía, L. C., Herre, E. A., Sparks, J. P., Winter, K., García, M. N., Van Bael, S. A., … Maximova, S. N. (2014).
Pervasive effects of a dominant foliar endophytic fungus on host genetic and phenotypic expression
in a tropical tree. *Frontiers in Microbiology*, *5*, 479. doi: 10.3389/fmicb.2014.00479

Nilsson, R. H. (2011). Molecular Identification of Fungi: Rationale, Philosophical Concerns, and the UNITE
Database. *The Open Applied Informatics Journal*, *5*(1), 81–86. doi: 10.2174/1874136301105010081

Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGinn, D., … Wagner, H. (2016). *vegan:
Community Ecology Package*. Retrieved from https://github.com/vegandevs/vegan

Oono, R., Lefèvre, E., Simha, A., & Lutzoni, F. (2015). A comparison of the community diversity of foliar
fungal endophytes between seedling and adult loblolly pines (Pinus taeda). *Fungal Biology*, *119*(10),
917–928. doi: 10.1016/J.FUNBIO.2015.07.003

Paradis, E., & Schliep, K. (2019). Ape 5.0: An environment for modern phylogenetics and evolutionary
analyses in R. *Bioinformatics*, *35*(3), 526–528. doi: 10.1093/bioinformatics/bty633

Pebesma, E., Rowlingson, B., & Rogerbivandnhhno, M. R. B. (2012). Package ' rgdal .' *R-CRAN*, *4*, 1–41. doi:
10.1353/lib.0.0050

Polonio, J. C., Almeida, T. T., Garcia, A., Mariucci, G. E. G., Azevedo, J. L., Rhoden, S. A., & Pamphile, J. A.
(2015). Biotechnological prospecting of foliar endophytic fungi of guaco (Mikania glomerata Spreng.)

with antibacterial and antagonistic activity against phytopathogens. *Genetics and Molecular Research*,
14(3), 7297–7309. doi: 10.4238/2015.July.3.5

Porras-Alfaro, A., & Bayman, P. (2011). Hidden Fungi, Emergent Properties: Endophytes and Microbiomes.
Annual Review of Phytopathology, *49*(1), 291–315. doi: 10.1146/annurev-phyto-080508-081831

Purevdorj, T. S., Tateishi, R., Ishiyama, T., & Honda, Y. (1998). Relationships between percent vegetation
cover and vegetation indices. *International Journal of Remote Sensing*, *19*(18), 3519–3535. doi:
10.1080/014311698213795

Qian, H., & Jin, Y. (2016). An updated megaphylogeny of plants, a tool for generating plant phylogenies and
an analysis of phylogenetic community structure. *Journal of Plant Ecology*, *9*(2), 233–239. doi:
10.1093/jpe/rtv047

Robeson, M. S., King, A. J., Freeman, K. R., Birky, C. W., Martin, A. P., & Schmidt, S. K. (2011). Soil rotifer
communities are extremely diverse globally but spatially autocorrelated locally. *Proceedings of the*
National Academy of Sciences of the United States of America, *108*(11), 4406–4410. doi:
10.1073/pnas.1012678108

Rodriguez, R. J., White, J. F., Arnold, a E., & Redman, R. S. (2009). Fungal endophytes: diversity and
functional roles. *The New Phytologist*, *182*(2), 314–330. doi: 10.1111/j.1469-8137.2009.02773.x

Rognes, T., Flouri, T., Nichols, B., Quince, C., & Mahé, F. (2016). VSEARCH: a versatile open source tool for
metagenomics. *PeerJ Preprints*, *4*, e2409v1. doi: 10.7287/peerj.preprints.2409v1

Saucedo-García, A., Anaya, A. L., Espinosa-García, F. J., & González, M. C. (2014). Diversity and Communities
of Foliar Endophytic Fungi from Different Agroecosystems of Coffea arabica L. in Two Regions of
Veracruz, Mexico. *PLoS ONE*, *9*(6), e98454. doi: 10.1371/journal.pone.0098454

Smith, D. P., & Peay, K. G. (2014). Sequence depth, not PCR replication, improves ecological inference from
next generation DNA sequencing. *PLoS ONE*, *9*(2). doi: 10.1371/journal.pone.0090234

Stone, J. K., Bacon, C. W., & White, J. F. (2000). An overview of endophytic microbes: endophytism defined.
Microbial Endophytes, (January 2000), 3–29. doi: 10.1163/_q3_SIM_00374

Tardieu, F., Lafarge, T., & Simonneau, T. (1996). Stomatal control by fed or endogenous xylem ABA in
sunflower: interpretation of correlations between leaf water potential and stomatal conductance in
anisohydric species. *Plant, Cell and Environment*, *19*(1), 75–84. doi: 10.1111/j.1365-
3040.1996.tb00228.x

Tipton, L., Zahn, G., Datlof, E., Kivlin, S. N., Sheridan, P., Amend, A. S., & Hynson, N. A. (2019). Fungal
aerobiota are not affected by time nor environment over a 13-year time series at the Mauna Loa
Observatory. *PNAS*, *in press*.

U'Ren, J. M., & Arnold, A. E. (2016). Diversity, taxonomic composition, and functional aspects of fungal
communities in living, senesced, and fallen leaves at five sites across North America. *PeerJ*, *4*, e2768.
doi: 10.7717/peerj.2768

U'Ren, J. M., Lutzoni, F., Miadlikowska, J., Laetsch, A. D., & Arnold, A. E. (2012). Host and geographic
structure of endophytic and endolichenic fungi at a continental scale. *American Journal of Botany*,
99(5), 898–914. doi: 10.3732/ajb.1100459

U'Ren, J. M., Lutzoni, F., Miadlikowska, J., Zimmerman, N. B., Carbone, I., May, G., & Arnold, A. E. (2019). Host
availability drives distributions of fungal endophytes in the imperilled boreal realm. *Nature Ecology*
& Evolution, *3*, pages1430–1437. doi: 10.1038/s41559-019-0975-2

Unterseher, M., Petzold, A., & Schnittler, M. (2012). Xerotolerant foliar endophytic fungi of Populus
euphratica from the Tarim River basin, Central China are conspecific to endophytic ITS phylotypes of
Populus tremula from temperate Europe. *Fungal Diversity*, *54*(1), 133–142. doi: 10.1007/s13225-
012-0167-8

Vázquez, D. P., Melián, C. J., Williams, N. M., Blüthgen, N., Krasnov, B. R., & Poulin, R. (2007). Species
abundance and asymmetric interaction strength in ecological networks. *Oikos*, *116*(7), 1120–1127.
doi: 10.1111/j.0030-1299.2007.15828.x

Vincent, J. B., Weiblen, G. D., & May, G. (2016). Host associations and beta diversity of fungal endophyte
communities in New Guinea rainforest trees. *Molecular Ecology*, *25*(3), 825–841. doi:
10.1111/mec.13510

Ward, T. L., Dominguez-Bello, M. G., Heisel, T., Al-Ghalith, G., Knights, D., & Gale, C. A. (2018). Development
of the Human Mycobiome over the First Month of Life and across Body Sites. *MSystems*, *3*(3), e00140-
17. doi: 10.1128/mSystems.00140-17

Warren, D. L., Cardillo, M., Rosauer, D. F., & Bolnick, D. I. (2014). Mistaking geography for biology: Inferring
processes from species distributions. *Trends in Ecology and Evolution*, Vol. 29, pp. 572–580. doi:
10.1016/j.tree.2014.08.003

Wiens, J. J., Ackerly, D. D., Allen, A. P, Anacker, B. L., Buckley, L. B., Cornell, H. V., … Stephens, P. R. (2010).
Niche conservatism as an emerging principle in ecology and conservation biology. *Ecology Letters*,
13(10), 1310–1324. doi: 10.1111/j.1461-0248.2010.01515.x

Ziegler, A. C. (2002). Hawaiian Natural History, Ecology, and Evolution. In *Hawaiian Natural History,*
Ecology, and Evolution. doi: 10.1515/9780824842437

Zimmerman, N. B., & Vitousek, P. M. (2012). Fungal endophyte communities reflect environmental
structuring across a Hawaiian landscape. *Proceedings of the National Academy of Sciences*, *109*(32), 13022–
13027. doi: 10.1073/pnas.1209872109

818
819
DATA ACCESSIBILITY STATEMENT