1 **TITLE**
2 Fungal communities living within leaves of native Hawaiian dicots are structured by landscape-scale variables as
3 well as by host plants
4
5 **RUNNING TITLE**
6 Hawaiian leaf fungi at the landscape scale
7

8 **AUTHORS**
9 Darcy JL[1,2], Switf SOI[1], Cobian GM[1], Zahn G[3], Perry BA[4], Amend AS[1]
10
11 [1] Department of Botany, University of Hawaii, Manoa
12 [2] Division of Biomedical Informatics and Personalized Medicine, University of Colorado Anschutz Medical
13 Campus
14 [3] Department of Biology, Utah Valley University
15 [4] Department of Biological Sciences, California State University East Bay
16
17 **CORRESPONDING AUTHOR**
18 John L. Darcy – +1 303 887 0477 – darcyj@colorado.edu

19
ABSTRACT

A phylogenetically diverse array of fungi live within healthy leaf tissue of dicotyledonous plants. Many studies

have examined these endophytes within a single plant species and/or at small spatial scales, but landscape-scale

variables that determine their community composition are not well understood, either across geographic space,

across climatic conditions, or in the context of host plant phylogeny. Here, we evaluate the contributions of these

variables to endophyte beta diversity using a survey of foliar endophytic fungi in native Hawaiian dicots

sampled across the Hawaiian archipelago. We used Illumina technology to sequence fungal ITS1 amplicons to

characterize foliar endophyte communities across five islands and 80 host plant genera. We found that

communities of foliar endophytic fungi showed strong geographic structuring between distances of seven and 36

km. Endophyte community structure was most strongly associated with host plant phylogeny and

evapotranspiration, and was also significantly associated with NDVI, elevation, and solar radiation. Additionally,

our bipartite network analysis revealed that the five islands we sampled each harbored significantly specialized

endophyte communities. These results demonstrate how the interaction of factors at large and small spatial and

phylogenetic scales shape fungal symbiont communities.

34
KEYWORDS
Fungi, Phylogeography, Endophytes, Hawai'i, High-throughput sequencing, Island Biogeography, Generalized
Dissimilarity Modeling

INTRODUCTION

Foliar endophytic fungi (FEF), defined here as all fungi living within leaf tissue but not causing any outward

signs of disease (sensu (Stone, Bacon, & White, 2000)) are effectively invisible and represent a "hotspot" of

undescribed fungal diversity (Arnold & Lutzoni, 2007; Porras-Alfaro & Bayman, 2011). Indeed, a large portion

of the vast undescribed fungal diversity on earth (Blackwell, 2011) are presumed to live cryptic lifestyles in

association with plant hosts (Hawksworth & Rossman, 1997), particularly in the tropics where plant diversity is

highest. Due to their cryptic lifestyles and high species richness (Blackwell, Hibbett, Taylor, & Spatafora, 2006;

Rodriguez, White, Arnold, & Redman, 2009), many questions remain about which factors determine how FEF

are distributed throughout nature.

Although all available evidence suggests that most eudicot FEF are horizontally transmitted and not inherited via

seed (Bayman, Angulo-Sandoval, Báez-Ortiz, & Lodge, 1998), it is unclear which factors structure FEF

community composition, and how those factors rank in their relative importance. Previous studies that have

examined foliar fungal communities in this ecological and biogeographic context have noted several different

drivers of community composition and biogeography. Temperature (Coince et al., 2014), geographic distance

(U'Ren, Lutzoni, Miadlikowska, Laetsch, & Arnold, 2012), elevation and rainfall (Zimmerman & Vitousek,

2012), and vegetation density/urbanization (Massimo et al., 2015; Unterseher, Petzold, & Schnittler, 2012;

Vincent, Weiblen, & May, 2016) are all putatively important variables for FEF community composition. We

might attribute this diversity of results to idiosyncrasies of experimental design in which specific variables or

host taxa were held constant to address focused hypotheses, and to the simultaneous influence of many

possible factors that may not all be analyzed in any given study.

In fact, the bulk of FEF community research is represented by studies focusing on one or two specific host plants

(Felber et al., 2016; González-Teuber, 2016; Kato, Fukasawa, & Seiwa, 2017; Mejía et al., 2014; Oono, Lefèvre,

Simha, & Lutzoni, 2015; Polonio et al., 2015; Saucedo-García, Anaya, Espinosa-García, & González, 2014;

Zimmerman & Vitousek, 2012), often to address hypotheses about how distance or climate impacts composition.

Still, studies that have surveyed FEF or other leaf-associated fungi among multiple plant species have found that

host identity strongly determines fungal composition (Huang, Devan, U'Ren, Furr, & Arnold, 2016; Kembel &

Mueller, 2014; Massimo et al., 2015; U'Ren & Arnold, 2016; Unterseher et al., 2012; Vincent et al., 2016).

Despite examples of studies at either broad phylogenetic scales or broad spatial scales, comparatively few

studies combine these into a single analytical framework (but see (U'Ren et al., 2012, 2019)). By simultaneously

examining a wide diversity of hosts over a wide diversity of habitats at the landscape scale, we might explicitly

address how host identity interacts with the abiotic environment to structure FEF community composition, and

the extent to which FEF are geographically restricted either by distance or by island barriers.

Here, we use the breadth of environmental and evolutionary diversity in the Hawaiian Archipelago to determine

which factors structure FEF beta diversity in native Hawaiian dicots. Specifically, we test the hypotheses that

elevation, rainfall, and host plant phylogeny are the strongest predictors of FEF community composition of

native plants across Hawaii (Figure 1). We include additional potential explanatory variables in our analysis such

as evapotranspiration (Kivlin, Hawkes, & Treseder, 2011; Lewis, Ravel, Naffaa, Astier, & Charmet, 1997) and

geographic distance (Higgins, Arnold, Coley, & Kursar, 2014), known to be important determinants in other

fungal systems. We found that FEF community structure was most strongly associated with host plant phylogeny

and evapotranspiration, and was also significantly associated with NDVI, elevation, and solar radiation. Spatial

structuring of FEF communities was strongest at distances between seven and 36 km.

MATERIALS AND METHODS

Sample Collection

General sampling locations were selected to maximize habitat, phylogenetic and spatial diversity. Location

selection prioritized access, a reasonable permitting process, and the known presence of ten or more native plant

species. Sample collection took place between July 2014 and October 2016. Access was the most limiting factor

in comparatively fewer sampling locations on Molokai and Maui Islands. We chose not to collect plants that are

federally listed as threatened or endangered. At each location, the first occurrence of an apparently healthy,

naturally recruited individual was selected for sampling. We only sampled native plants. Because some native

species are comparatively common, and others rare, we limited our sampling to a single individual from each

naturally occurring non-listed dicot at each location (*e.g.* a trail or a park), because we did not wish to confound

our data with uneven spatial heterogeneity among sub-sampled plants. In most cases more than a single

individual was collected per island since it occurred at more than one location. Numerous mature sun leaves

were collected such that when combined, they covered a surface area roughly equivalent to two adult-sized

hands. Life form and stature differed markedly across our sampled plants, so it was difficult to precisely

standardize the height at which leaves were collected. When possible, leaves from trees and shrubs were

collected at eye level and from at least four aspects of the tree canopy. We targeted dicots because they have

horizontally transmitted FEF (compared to many vertically transmitted FEF in some monocots). Although not all

host plant genera were collected at every sample location, the three most common genera we collected

(*Metrosideros*, n=119; *Leptecophylla*, n=89; and *Vaccinium, n=*147) were each collected across elevations

ranging from sea level to over 2000 meters above sea level. The location of each plant was recorded with a GPS

and plants were positively identified in the field and/or vouchered for subsequent identification (vouchers

deposited at Joseph F. Rock Herbarium at the University of Hawaii, Manoa; HAW). Leaves were refrigerated

until subsequent processing (within 72 hours of collection). A total of 1099 samples were collected in this way,

although not all were used in the analyses presented here (see results section).

We surface sterilized leaves to exclude fungi present on leaf surfaces. After rinsing in water, forty leaf discs were

extracted per individual host by punching leaves with a sterile standard paper single hole punch (approximately

0.5 cm diameter). Hole punches were sterilized by washing with 70% ethanol and flamed between uses. For

plants with very small leaves, entire leaves were used such that the area of those leaves was the same as the area

of leaf discs. Leaf discs (or aforementioned collections of small leaves) were then placed into loose-leaf tea bags

that were subsequently stapled shut, submerged in 1% NaOCl for two minutes, then 70% EtOH for two minutes,

followed by two rinses with sterile water for two minutes each. Rinse water was included in extraction controls

to verify sterility of surface water.

DNA Isolation

Ten leaf discs per DNA extraction were placed in MP Biomedical Lysing Matrix A tubes (MP Biomedical, Santa

Ana, CA, USA) containing DNA isolation solutions from the Mo Bio PowerPlant Pro DNA Isolation kit

(Solution PD1, Solution PD2, Phenolic Separation Solution, and RNase A Solution; Mo Bio, Carlsbad, CA,

USA, a subsidiary of QIAGEN). Leaf discs were homogenized using a Mini-Beadbeater 24 (BioSpecs Inc. OK)

at 3,000 oscillations per min for two minutes. Lysate was centrifuged at 13,000 RPM for two minutes and

transferred to individual wells of a MoBio PowerPlant Pro DNA 96-well Isolation kit for subsequent extraction

following the manufacturer's protocol.

PCR Amplification and Illumina Library Preparation

We amplified the ITS1 region of the ribosomal cistron using fungal specific primers ITS1F (5'-

CTTGGTCATTTAGAGGAAGTAA-3') and ITS2 (5'-GCTGCGTTCTTCATCGATGC-3'), along with Illumina

adapters and Golay barcodes incorporating dual indexing, using previously published thermal cycling parameters

(Smith & Peay, 2014). PCRs were carried out in 25 µl reactions using the KAPA3G Plant PCR kit (KAPA

Biosystems, Wilmington, MA, USA), nine ul of DNA extract (concentration not measured) and 0.2 µM each of

the forward and reverse primers. Negative PCR and extraction controls were included. PCR products were

purified and normalized using just-a-plate 96 PCR Purification and Normalization Kit (Charm Biotech, San

Diego, California, USA). Normalized PCR products were pooled and concentrated using a streptavidin magnetic

bead solution. Pooled PCR products were sequenced on three separate reactions in order of processing

completion, using the 2x300 paired-end (PE) sequencing protocol on an Illumina MiSeq sequencing platform

(Illumina Inc., Dan Diego, CA, USA). We did not include controls for sequencing run variation, and we

acknowledge this as a limitation of our study. We also did not quantify DNA concentration of extracts, but the

amplification-normilization experimental design allows for characterization of microbial communities even from

inefficient DNA extractions. We do not think this is a significant source of bias in our analysis, since negative

controls (which contained too little DNA to amplify) produced a paucity of sequence data when compared to

samples, almost all of which produced many thousands of sequence reads. Still, it is possible that variation in

extraction efficiency could result in biased alpha diversity estimates. Due to low sequencing success, negative

controls were not accounted for in sample processing.

DNA Sequence data processing and bioinformatics

QIIME v1.9.1 (Caporaso et al., 2010) was used to demultiplex raw DNA sequence data into individual fastq files

for each sample. Although paired-end sequencing was used, only the R1 read (corresponding to primer ITS1f)

was used for downstream analysis, since sequencing quality of reverse reads was generally poor after the first

150 nucleotides. VSEARCH v2.14.2 (Rognes, Flouri, Nichols, Quince, & Mahé, 2016) was used to discard reads

with an average quality score below 25 (Illumina Q+33 format), then ITSx v1.1 (Bengtsson-Palme et al.,

2013) was used to extract the ITS1 region from quality-filtered files. Sequence data were filtered by average

score rather than maximum score, because nucleotides with poor quality scores that were not within the ITS1

region could disqualify sequences that would be otherwise high-quality within the ITS1 region. In order to

eliminate variability in sequences caused by sequencing and PCR error, ITSx-processed sequences were de-

noised using DADA2 v1.12 (Callahan et al., 2016) using the authors' recommended settings (maximum

expected errors = 2, maximum Ns = 0). Separate DADA2 error rates models were used for each sequencing run

in our data set, although the models appeared qualitatively similar. De-noised sequences were then clustered into

95% OTUs using VSEARCH (Rognes et al., 2016).

Taxonomy was assigned to each OTU using the UNITE database (v7) (Nilsson, 2011) and QIIME's

assign_taxonomy.py script (Caporaso et al., 2010) with the BLAST method using the default maximum e-value

of 0.001. OTUs not belonging to kingdom Fungi were discarded. Our OTU table was then rarefied (*i.e.*

randomly downsampled) to 1500 sequences per sample, discarding samples with fewer than 1500 sequences or

samples for which host plants could not be satisfactorily identified. We chose the depth of 1500 sequences as a

compromise between a depth at which almost all samples had plateaued on our collector's curves

(Supplementary figure S1), and a level that resulted in the fewest discarded samples. Filtering and rarefaction

were performed in R v3.5.0 (R Core Team, 2014) using custom software written by the authors.

Ghost Tree v2019 (Fouquier et al., 2016) was used to construct a phylogenetic tree for the remaining OTUs.

Briefly, Ghost Tree allows phylogenetic trees to be made from ITS1 sequence data, which are often un-alignable

at high taxonomic ranks. This is done using a backbone tree created with the 18S rRNA gene, and then ITS1

sequences are used to refine the tree at a phylogenetic scale where those sequences can be meaningfully aligned

(*e.g.* genus level). In an analysis of real and simulated ITS1 data, Fouquier et al. (2016) found that using UniFrac

distances calculated using a tree made with Ghost Tree were better explained by environmental variables in

ANOSIM models than were distances calculated using phylogeny-independent metrics. (Fouquier et al., 2016).

This method was also successfully used to study differences in fungal community composition in the human

infant microbiome (Ward et al., 2018) and in a study of FEF in grass as well (Lumibao et al., 2019). We used a

Ghost Tree that was made using the SILVA database (v132) for the 18S backbone, and the UNITE database (v8).

was used for ITS refinement. Tips of the Ghost Tree were renamed with OTU identifiers where OTUs were

assigned taxonomy to a UNITE entry in the Ghost Tree. In cases where multiple OTUs were assigned to the

same UNITE entry, a polytomy was created to fit those OTUs into the tree. This method may exclude novel taxa

that are not present in the UNITE reference database, and this is a limitation of all studies using such a database,

including this one.

The Ghost Tree was used with weighted UniFrac (Lozupone & Knight, 2005) (hereafter referred to as

"UniFrac") to construct a beta diversity distance matrix for the samples. UniFrac was used because it is

downweights additional branch length contributed by slight intragenomic variation among tandem repeats. Even

if an individual putative fungal species contains several OTUs, each will only contribute a negligible amount of

branch length to a sample, as compared to non-phylogenetic metrics (*e.g.* Bray-Curtis), which would consider

those OTUs as different as any other pair. Because UniFrac community dissimilarity considers the shared

phylogenetic branch-lengths between two communities, it is robust to the case where an OTU is only found

within one sample, and is similarly robust to the case where samples do not share any OTUs, which can be

problematic for other community dissimilarity metrics where overall diversity is much higher than within sample

diversity. This is important for our analysis of FEF communities, because many previous studies have shown that

FEF are hyperdiverse even at local scales (*e.g.* within one host or within one hundred meters) (Jumpponen &

Jones, 2009, 2010; Zimmerman & Vitousek, 2012). We suspected that this large amount of diversity would result

in many pairs of samples that shared few or zero OTUs, which would result in an inflation of 1-values

(maximum dissimilarity) when using non-phylogenetic beta diversity metrics such as Bray-Curtis or Jaccard

community dissimilarity. Indeed, this same issue has been discussed by previous users of Generalized

Dissimilarity Modeling (GDM) as a saturation of maximal dissimilarities, and was resolved by using

phylogenetic dissimilarty metrics (Warren, Cardillo, Rosauer, & Bolnick, 2014). We confirmed that this was

indeed the case, and that UniFrac distance values were normally distributed but Bray-Curtis dissimilarities were

severely 1-inflated (Supplementary figure S2). For this reason, we used the UniFrac distance matrix for the

remainder of our analysis.

Spatiotemporal data

Using sample geographic coordinates and collection dates, environmental and climatic data for each sample

were extracted from GIS layers using the R packages '*raster*' v2.9 (Hijmans et al., 2014) and '*rgdal*' v1.4

(Pebesma, Rowlingson, & Rogerbivandnhhno, 2012). Data were extracted from rasters generated for the same

month that samples were collected (except for elevational data). Supplementary table S1 shows the sources of

each GIS layer. Many explanatory variables were obtained from the Rainfall of Hawai'i and Evapotranspiration

of Hawai'i websites (T W Giambelluca et al., 2014; Thomas W. Giambelluca et al., 2013), and elevation data

were obtained using the USGS EarthExplorer online tool (http://earthexplorer.usgs.gov), courtesy of NASA

EOSDIS Land Processes Distributed Active Archive Center and the United States Geological Survey's Earth

Resources Observation and Science Center. These explanatory variables were chosen either because previous

studies of Hawaiian FEF had identified them as important (elevation, rainfall)(Zimmerman & Vitousek, 2012),

because they were significant drivers of Hawaiian plant composition (slope, aspect), or because they made

intuitive sense in the context of fungi that live within leaves (solar radiation, transpiration, evapotranspiration,

leaf area index, NDVI). Slope and aspect of each sampling location were calculated from elevation raster data

using the terrain function in '*raster*'. Data for aspect (the direction a sampling location faces) were converted

into a distance matrix using the smallest arc-difference between any two given aspects. This was done because

Euclidean distance is unsuitable for a measurement like aspect, where 355° is closer to 1° than it is to 340°. All

variables are mean monthly values. NDVI (normalized difference vegetation index) is an index calculated from

the amount of infra-red light reflected by plants, which is normalized using multiple wavelengths of visible light.

This allows for discrimination between habitat types that are deferentially vegetated. Similarly, leaf area index is

a measure of surface area of leaves (one-sided) per unit area of ground, and while it does not discriminate

between different types of vegetation as does NDVI, unlike NDVI it measures the density of leaf surface area

(habitat for FEF).

Host plant phylogeny

A distance matrix of host plant phylogenetic distances was created using the angiosperm phylogeny of Qian and

Jin (Qian & Jin, 2016). This distance matrix was made because the modeling method we use here (GDM, below)

can accommodate distance matrices as explanatory variables, allowing for a phylogenetic distance matrix of

hosts to be used instead of a simplified data structure such as a principal components vector or an array of

categorical taxon identities. For each pair-wise comparison between two samples, pair-wise host plant

phylogenetic distance was calculated as the mean cophenetic (branch-length) distance between members of the

plant genera that were sampled, using the function cophenetic.phylo in the R package '*ape*' v5.2 (Paradis &

Schliep, 2019). In cases where host plant genus was not included in the phylogeny, the genus was substituted for

the most closely-related genus available. Four genera were substituted in this way out of 80 total genera:

Labordia → Logania, Touchardia → Urtica, Waltheria → Hermannia, Nothocestrum → Withania. We chose to

resolve host plants at the genus level because species-level taxonomy can be unstable for recent adaptive

radiations, and many Hawaiian species-level phylogenies are not resolved (Ziegler, 2002).

Geospatial analyses

In order to test and visualize the spatial structuring of FEF beta diversity, we constructed a geographic

semivariogram (Bachmaier & Backes, 2008) in R using a matrix of log_{10} transformed geographic distances

between samples, and the UniFrac distance matrix of FEF community differences between samples.

Semivariograms are useful tools in spatial ecology because they visualize the extent to which beta diversity

exhibits spatial connectedness. They can show at what geographic range community composition is

homogeneous, and at what range community composition is structured by distance. See Robeson et al. (2011) for

a thorough discussion of the applicability of Semivariograms in this context, models that can be fit to them, and

what hypotheses they can be used to test. Distance classes in the semivariogram were constructed using a sliding

window methods, where each class shared two-thirds of its points with the previous class (Darcy, King,

Gendron, & Schmidt, 2017). A logistic curve was fit to semivariance values and mean geographic distances

within semivariogram distance classes using the Nelder-Mead algorithm, minimizing the sum of squared

differences in beta diversity semivariance. The curve's derivative was then used to find the geographic distance

class that accounts for 99% of geographic distance where beta diversity is spatially autocorrelated. Since the

derivative function of a logistic curve is a unimodal distribution, the 99% intervals of that function yield three

distance classes for the logistic function: the first plateau, the class corresponding to 99% of the change in slope

for the curve, and the second plateau. Mantel correlations were then performed on data within each of the three

resulting distance classes.

Generalized dissimilarity modeling (GDM) (Ferrier, Manion, Elith, & Richardson, 2007) was used to model FEF

beta diversity based on climatic factors (Supplementary table S1), geographic distance, host plant phylogeny,

and DNA sequencing run. The goal of this analysis was to discover the relative importances of environmental

variables on FEF community composition, and at what scales the variables explain FEF community composition

(*e.g.* are small changes in a variable important, or only large changes). GDM is a form of non-linear matrix

regression that is well-suited to statistical questions involving dissimilarity matrices (*e.g.* our UniFrac distance

matrix, host plant cophenetic distance matrix, geographic distance matrix, and aspect arc-difference matrix).

Unlike pair-wise Mantel tests or PerMANOVA, which make use of similar data, GDM can quantify the relative

importance of environmental and geographic variables on community dissimilarity, even when the functional

relationship between community dissimilarity and the environment is nonlinear (Fitzpatrick et al., 2013; Warren

et al., 2014). Furthermore, GDM is effective because it models community dissimilarity associated with a given

predictor variable while holding all other predictor variables constant (Fitzpatrick et al., 2013; Landesman,

Nelson, & Fitzpatrick, 2014). For example, this enables GDM to model the effect of elevation while accounting

for the effect of host plant phylogenetic distance. Because our data were generated using three separate DNA

sequencing runs, we included "run" as a categorical variable in our model by creating a distance matrix where

samples from the same run had distance=0, and samples of different runs had distance=1, as recommended by

GDM's creators (Ferrier et al., 2007).

We used backward elimination as implemented in the '*gdm*' v1.3.11 R package to build a model, and then to

simplify the model by removing minimally predictive variables. We began this process with a GDM model

excluding variables that were correlated with another variable in the analysis at rho > 0.70 (Spearman). Leaf area

index, wet canopy evaporation, temperature, cloud frequency, and relative humidity were removed this way

(Supplementary figure S3). Log-transformed geographic distance between samples, host plant phylogenetic

distance, sequencing run, and aspect arc-difference matrix were included as predictor matrices in the GDM, with

FEF UniFrac distance as the response matrix. We then used GDM to test each variable within the model for

significance using a permutation test. During this iterative process, the variable with the highest P-value was

eliminated, and then the model was recalculated. This process was repeated until all remaining variables were

statistically significant ($P < 0.05$). Sequencing run and transpiration were eliminated this way. The final GDM

model was also analyzed using Bray-Curtis dissimilarities instead of UniFrac distances, to illustrate how the

severe 1-inflation observed with Bray-Curtis was an impediment to model fitting.

Principal coordinate analysis (PCoA) was used to validate GDM results using an alternate statistical approach.

PCoA vectors were calculated from the UniFrac distance matrix using the R package '*vegan*' v2.5 (Oksanen et

al., 2016). Principal Coordinates of Neighbor Matrices (PCNM)(Borcard & Legendre, 2002) was used on

geographic and host phylogenetic distance matrices, for dimensionality reduction and to account for nonlinear

features within those data. The same data used in our final GDM, except with geography and host represented by

their first six PCNM vectors, were fit to FEF PCoA and tested using the 'envfit' function (Oksanen et al., 2016).

FEF-island specialization analysis

Bipartite network analysis was used to test the extent to which each island (Figure 1) harbored specific FEF

(FEF-island specialization). The d' ("d prime") statistic was calculated for each island using the OTU table using

the '*bipartite*' package v2.11 in R (Dormann, Gruber, & Fruend, 2008). d' is a measure of network specialization

that ranges from zero to one, where zero is perfect cosmopolitanism (all species are evenly shared among

islands) and one is perfect specialization (each species is specific to only one island). d' is calculated using a

contingency matrix where each row is a unique lower-level group (island) and each column is a unique higher-

level group (OTU), but in our OTU table each island contains multiple samples, so the number of observations

per island is not consistent. To remedy this, we calculated d' by aggregating all samples from the same island

into one large sample (column sums), rarefied this aggregated table using the same depth to which samples were

rarefied above (1500 observations), then calculated d' values. Empirical d' values were compared to null d'

values generated with the Vázquez null model, which is a fixed-connectance null model for bipartite networks

(Vázquez et al., 2007), implemented as the vaznull function of '*bipartite*'. The procedure of aggregating,

calculating an empirical d', and calculating a Vázquez null d' was repeated 1000 times in order to obtain

bootstrapped distributions of empirical and null d' values for each island. Statistical significance for d' values for

each island was tested using Welch's unequal variance *t*-test to compare empirical and island null d' values. This

test was 2-tailed, since d' could be significantly lower than the null distribution indicating cosmopolitanism, or

significantly higher indicating specialization.

A similar procedure was applied to host plant genera in place of fungal OTUs (plant-island specialization). Since

host is only observed once per sample, d' was not bootstrapped. This was done in order to evaluate the extent to

which our selection of host plants was significantly specialized to islands.

To evaluate whether FEF-island specialization could be attributed to island-specific plant selections, we also

tested FEF-island specialization under the constraint of a generically identical plant community (at genus level)

for each island. The same procedure for analyzing FEF-island specialization above was carried out, but during

the aggregation step of each bootstrap, random samples for each island belonging to a fixed plant community

were chosen to the exclusion of all others. This fixed community was defined by the plant identification overlap

among all five islands. All d' values calculated this way were calculated from the same plant community from

each island. Analysis of these d' values was calculated as above.

RESULTS

Sequence data

Samples were collected across a wide range of climatic conditions (Figure 2), representative of the distributions

of those conditions in Hawai'i. Our data set comprised 896 samples that passed quality-filtering and ITS1

extraction, consisting of 7002 OTUs. After non-fungal OTUs were discarded, samples were rarefied to 1500

sequences per sample, resulting in 3458 OTUs across 760 samples. Of those samples, 417 were from Hawai'i,

84 from Kaua'i, 56 from Maui, 72 from Moloka'i, and 131 from O'ahu. Mean richness (number of OTUs

observed) per sample was 27.3 with a standard deviation of 18.9 (Supplementary Figure S1). Geographic and

botanical information for each sample used in our analysis is available as Supplementary Figure S4 and is

available from our FigShare repository (see Data Accessibility section) . Of 27 sequencing negative controls

included in the study, 13 resulted in sequence data, with an average of 123 reads and 2.2 OTUs per sample. This

sequencing depth was less than 1% of that found in the average biological sample (12,860). Because we were

unable to trace the source of these contaminants (whether tag switching, cross contamination or reagent

contamination) and because they amplified so poorly compared to biological samples, we did not attempt to

adjust our data set to account for these. A data table containing geographic and host taxonomic information for

each of the 760 samples used in our analysis is available as Supplementary Table S2.

Geospatial results

Our semivariogram analysis showed that FEF beta diversity is spatially structured between 7.6 and 36.3 km,

where it reaches its autocorrelation distance (Figure 3). This plateau and subsequent noise in beta diversity

semivariance occurs when FEF beta diversity is no longer spatially autocorrelated (Robeson et al., 2011). In our

analysis, this occurs at geographic distances of 36.3 km. Within the distance class where beta diversity was

spatially autocorrelated, r_M (Mantel correlation) was 0.16. In the other two distance classes, r_M was closer to zero

as expected (Figure 3).

After GDM backward elimination, many variables were left as statistically significant: All *P*-values were the

minimum possible *P*-value given that GDM's backward elimination was run using 50 permutations. This means

that GDM always predicted a smaller proportion of FEF beta diversity when any given predictor variable was

permuted, except for sequencing run and transpiration which were eliminated this way. Host plant phylogenetic

distance and evapotranspiration explained the most compositional dissimilarity in FEF communities, as given by

their GDM coefficients (the maximum height of their splines) (Ferrier et al., 2007; Fitzpatrick et al., 2013) which

were 0.090 and 0.078, respectively (Figure 4). These values indicate the relative importances of variables in the

model. The shape of GDM i-splines reflects FEF community turnover over the range of differences in a predictor

variable, and both host and evapotranspiration were related to FEF beta diversity over their entire ranges.

Sample collection date was related to FEF beta diversity only at temporally proximate values, while solar

radiation and NDVI were related to FEF beta diversity only at large differences (Figure 4). Although all

variables shown in Figure 4 are statistically significant and the whole model fit nicely to the data with relatively

symmetrical residuals (Figure 4, top), GDM was only able to predict 6% of FEF beta diversity. In contrast to the

GDM fit using the UniFrac distance matrix, GDM run with Bray-Curtis dissimilarity fit poorly as a result of

severe 1-inflation of distances (Supplementary figures S2 and S5).

Principal coordinate analysis (PCoA) was used to confirm GDM results using an alternative statistical approach.

Similar to GDM, host (PCNM3 of host phylogenetic distance matrix) had the strongest relationship with FEF

beta diverity with an R^2 of 0.106 (P = 0.001), and evapotranspiration had the second strongest relationship with

an R^2 of 0.073 (P = 0.001). A table of PCoA R^2 and P values is available as Supplementary Table S3.

Bipartite network analysis

Each of the five islands we sampled showed a statistically significant (P < 0.001, Welch's unequal variance t-

test) pattern of FEF specialization (Figure 5), since FEF-island d' values were higher than those generated using

a null model (null d' values centered around d'=0.25 for each island). Specialization in this case means that each

island harbors more FEF OTUs that are uniquely associated with that island than would be expected by chance.

We found that our sampling of host plant genera was significantly specialized to island as well (Supplementary

figure S6), however plant-island d' values were not significantly related to OTU-island d' values (linear

regression, $P = 0.45$). We also analyzed FEF-island specialization while keeping plant taxonomy identical across

islands by randomly drawing a pre-set plant community from each island in each permutation of the d' analysis.

This analysis was done to remove the effect of plant community on FEF-island specialization. We still found

significant specialization for all islands, although in this analysis all five islands had very similar d' values

(Supplementary figure S7).

Endophyte Taxonomic composition

At the Class level, FEF composition was consistent across islands, and dominated by Dothideomycetes and

Sordariomycetes (members of the Pezizomycotina)(Supplementary figure S8). 12.14% of endophyte OTUs were

unidentified at the class level (corresponding to 11.29% of rarefied sequence data), perhaps reflective of

Hawaii's isolation and comparative paucity of taxonomic research.

DISCUSSION

The most striking pattern we found in our analysis of FEF communities in native Hawaiian plants was that

evapotranspiration, a comparatively coarse measure of plant community traits at the landscape level, is

meaningfully predictive for the community composition of microscopic fungi living within plant leaves. These

results suggest that aside from environmental or host determinants, endophyte community composition results

from a microbial "neighborhood", in addition to the traits of individual hosts. Evapotranspiration was the

second-most important variable in our analysis in terms of community composition (Figure 4); and it was more

important in the model than elevation, which was measured at a much finer spatial resolution (Supplementary

table S1). This was supported by our principal coordinate analysis as well. This result makes the interpretation of

our hypothesis regarding the effect of climate on FEF community composition surprising, since previous studies

(Coince et al., 2014; Zimmerman & Vitousek, 2012) suggested that temperature, elevation, and rainfall would be

the most important factors structuring community composition instead, and evapotranspiration has not been

previously considered as an important variable for FEF. While elevation (tightly correlated with temperature),

host plant phylogeny, rainfall, and Julian date were significant explanatory variables in our GDM analysis, each

of their effects were smaller than that of evapotranspiration (Figure 4).

Evapotranspiration encompasses both plant transpiration and the evaporation of water from soil and other

surfaces, and both soil water content and stomatal conductance affect leaf interior moisture (Lambers, Chapin, &

Pons, 2008). As such, evapotranspiration could drive FEF community composition by changing the leaf interior

habitat, and thereby select for different FEF communities at high vs. low evapotranspiration. Indeed,

evapotranspiration is strongly correlated with the moisture content of leaves (Lambers et al., 2008; Tardieu,

Lafarge, & Simonneau, 1996). Alternatively, if plant stomata are open for longer periods, this may affect

probabilities of different FEF successfully entering the leaf interior. To our knowledge, no studies have measured

fungal community response to evapotranspiration, although associations between endophytic fungi and

transpiration (a component of evapotranspiration) have been strong. In a grass system spanning 15 European

countries, the response of endophytic fungi to a transpiration gradient was substantial (Lewis et al., 1997).

Association between native FEF and transpiration has also been observed in *Theobroma cacao* (Arnold &

Engelbrecht, 2007). In light of previous studies suggesting a link between fungal endophytes and

evapotranspiration (Arnold & Engelbrecht, 2007; Kivlin et al., 2011; Lewis et al., 1997), our finding that

evapotranspiration is a significant predictor of differences between FEF communities makes sense, and future

physiological experiments at the scale of individual leaves might shine light on the nature of this interaction.

The association we found between FEF beta diversity and host plant phylogenetic distance was the most

important predictor in our GDM model. Previous studies of FEF that have surveyed multiple host species, and

have found that host identity is commonly associated with fungal endophyte community composition (Liu, Zhao,

Wang, & Chen, 2019; U'Ren et al., 2012; Vincent et al., 2016), even among closely-related hosts (Christian et

al., 2020), so our result was mostly expected. But unlike most previous studies that found host associations of

FEF (although see (Davis & Shaw, 2008)), we used the phylogeny of host plants as an explanatory variable in

place of their identity. Under the hypothesis that FEF communities are structured by host phylogenetic

difference, more phylogenetically similar plants are expected to harbor similar FEF communities, and

conversely, phylogenetically distant plants are expected to harbor less similar FEF communities. Following this

hypothesis, in our analysis the GDM i-spline for host steadily increased across the whole range of differences in

host phylogenetic distance (Figure 4), although the abrupt increase at low distances indicates rapid turnover in

FEF community composition even with small changes in host phylogenetic distance. The significant association

we observed from genus-level all the way up to higher taxonomic ranks agrees with a study of broad-scale host

associations among tropical foliar fungal epiphytes, in which family level host taxonomy is most predictive

(Kembel & Mueller, 2014).

This relationship between FEF beta diversity and host plant phylogeny could either mean that FEF, host plants,

or both, exhibit a degree phylogenetic niche conservatism (Wiens et al., 2010). For example, FEF community

preference may be phylogenetically conserved among closely related plant species, or perhaps host preference is

conserved among closely related FEF. Patterns of phylosymbiosis are widespread throughout natural symbiotic

systems, including plant-pathogen systems. However, phylogenetic signal can be lost in the presence of

environmental factors that simultaneously impact symbiotic community composition (Brooks, Kohl, Brucker,

van Opstal, & Bordenstein, 2016). Furthermore, in our analysis, host plant phylogeny may have been a more

robust predictor of FEF community dissimilarity if our host plants had been classified to species level instead of

genus level; however species level phylogenies are not fully resolved for many Hawaiian lineages, particularly

those that have undergone rapid and recent adaptive radiations.

NDVI (normalized difference vegetation index) was a significant predictor of FEF community dissimilarity in

our data set, suggesting that areas that are differentially vegetated harbor different communities of FEF. This

difference may be related to the total percent land cover of vegetation, which is a component of NDVI

(Purevdorj, Tateishi, Ishiyama, & Honda, 1998), or related to the type of plant cover, *i.e.* different plant

community compositions (Lunetta, Knight, Ediriwickrema, Lyon, & Worthy, 2006), which is also encompassed

within the NDVI metric. Indeed, foliar fungal communities potentially respond to both land cover and habitat

type in the host plant *Quercus macrocarpa* (Jumpponen & Jones, 2009, 2010). This association between FEF and

NDVI means that FEF conservation efforts would likely need to consider differential vegetation cover and

urbanization at the landscape scale. The pixel size for the NDVI data we used was 250 m^2 (Supplementary table

S1), meaning that the value at any given location is an aggregate value for a large plant community. Thus, the

proportion of FEF community dissimilarity that was significantly explained by NDVI in our analysis is related to

the density and composition neighboring plant communities and not specifically the host individual, a dynamic

suggested in others studies as well (Kato et al., 2017).

The other statistically significant drivers of FEF community composition (Figure 4) were mostly expected,

particularly elevation (statistically significant in both GDM and PCoA analyses). This is similar to a result

reported by Zimmerman and Vitousek (2012), who used PerMANOVA to test the effects of rainfall, elevation,

and substrate age on FEF beta diversity patterns in *Metrosideros polymorpha* (ʻŌʻhia) trees. They found that

elevation explained roughly 17% of compositional dissimilarity between FEF communities, varying slightly

depending on which dissimilarity metric was used. That study also took place in Hawaiʻi, and the area sampled

overlaps partially with the area of Hawaiʻi island that we sampled (Figure 1). Zimmerman and Vitousek (2012)

also found a large effect of rainfall, which was also statistically significant in our GDM analysis. Rainfall is also

a significant driver of FEF community structure in grasses (Giauque & Hawkes, 2013), although in a larger

continental-scale analysis of cultured FEF isolates, rainfall was not a strong predictor of FEF diversity (U'Ren et

al., 2012). Solar radiation was also a significant variable in our GDM and PCoA analyses, and was much more

important in our GDM model than either elevation or rainfall. Indeed, previous research has shown that light

levels are important for FEF colonization, likely because different intensities of light exposure can affect the

chemical makeup of the phyllosphere (Bahnweg et al., 2005).

Geographic distance was of middling importance in our GDM analysis, especially since importance in GDM is

calculated with all other variables accounted for. However, in our semivariogram and subsequent Mantel

correlogram analyses, we found that FEF beta diversity is significantly spatially structured by geographic

distance, especially between geographic distances of 7.6 and 36.3 km (Figure 3). Communities are not strongly

spatially autocorrelated at distance of less than 7.6 km, and reach their autocorrelation range at 36.3 km. The lag

before 7.6 km indicates that species are sampled from the same local community before that point (Robeson et

al., 2011). After the curve plateaus around 36.3 km, FEF beta diversity reaches its "autocorrelation range", after

which autocorrelation becomes noisy. At distances greater than the autocorrelation range, communities are not

spatially autocorrelated; *i.e.* community composition is no longer a function of geographic distance (King et al.,

2010; Robeson et al., 2011). From our semivariogram we can qualitatively see that there may be spatial

structuring at greater distances as well, perhaps up to 100 km, indicating that community structure is

autocorrelated within islands, but not between them. Said another way, this indicates that the microbial

"neighborhood" for FEF is at the sub-island scale. The Hawaiian islands' volcanic origins might partially

account for these patterns. Easterly trade winds and orographic lift drive much of the precipitation patterns in the

archipelago, such that each island is neatly divided between comparatively hot and dry leeward, and cool wet

windward sides. Nearly all island flora reflect this division, therefore it is unsurprising to find spatial

autocorrelation at within, rather than among, island scales.

However, even though beta diversity may not be spatially autocorrelated at the inter-island scale, individual

species may still be unique to islands. Said another way, islands may harbor unique cohorts of FEF in addition to

cosmopolitan FEF, even though geographic space has little or no bearing on beta diversity at that scale. We

tested this, and found a significant effect of FEF specialization to islands (Figure 5). Our bipartite network

analysis showed that each island harbors a significant composition of fungi that are more regionally specific than

would be expected due to chance alone. Beta diversity tools like GDM and our semivariogram are very different

analytical methods than bipartite network analysis, and answer different questions. Our GDM analysis focused

on patterns in FEF beta diversity, which is aggregate dissimilarity of community composition and does not take

into account patterns at the OTU level, especially because of UniFrac's down-weighting of phylogenetically

similar OTUs. Our bipartite network analysis instead focused on the distribution of FEF OTUs within versus

among islands, testing the extent to which a given island harbors OTUs that are exclusive to that island.

We presumed that the likeliest drivers of FEF-island specialization are differences in habitat, dispersal limitation

among islands or differences in host availability (whether due to plant distributions or sampling bias). Since our

beta diversity analysis revealed a significant effect of host plant phylogeny, because Hawaiian island floras are

disharmonious (Ziegler, 2002), and because this disharmony was reflected in our sampling, differences in

sampled plant hosts might account for specialization patterns. We used two separate analyses to evaluate whether

this could account for specialization patterns alone. A bipartite analysis of plant-island specialization

(Supplementary figure S6) indicated that in our experimental design, hosts are indeed specific to islands.

However, we expected that if plant-island specialization bias within our experimental design was driving the

FEF-island specialization pattern we observed, d' values from the two would correlate strongly. There was no

significant relationship between plant-island specialization and FEF-island specialization, and plant-island d'

values were much lower (less specialized) than those observed for OTUs. Further, when we restricted our

analyses to only those plant genera that were sampled on all islands, the FEF specialization patterns remained.

How dispersal limitation shapes FEF-island specialization is an open question worthy of future research. There is

some evidence for restricted gene flow among populations of a putative native mushroom, *Rhodocollybia*

laulaha (Keirle, Avis, Feldheim, Hemmes, & Mueller, 2011) among islands, although these patterns might also

be explained by selection. Similarly, the lag in our semivariogram (Figure 3) could be the distance at which

dispersal limitation begins to structure FEF communities, or simply the distance at which un-measured

environmental variables begin to vary enough to structure FEF communities. A long-term study of aerobiota at

the summit of Hawaii's peak, Mauna Loa, found evidence of likely long-distance dispersal via air mass (Tipton

et al., 2019), although it is unclear how frequent and ecologically relevant such events are to endophyte systems.

The epidemiology and rapid inter-island spread of fungal pathogens such as rapid ʻŌʻhia death (*Ceratocystis*

spp.) or ʻŌʻhia rust (*Puccinia psidii*) suggest that at least some species are capable of crossing ocean channels,

whether naturally or with the unwitting help of humans. Of course not all species are so easily dispersed, and the

requisite and complex interaction between FEF, host and habitat likely reinforce island specialization patterns.

In summary, our analysis highlights that the various factors contributing to Hawaiian FEF community structure

do so at the landscape scale. We analyzed these important plant symbionts across both a large geographic scale

(Figure 1) and across a large host phylogenetic scale (80 plant genera), while using high-throughput sequencing

to thoroughly inventory FEF community composition. We examined the effects of climate, geographic distance,

and host identity using this system, and for the most part, found them to be strong predictors of differences in

FEF communities between samples – even when the measurements were taken at relatively coarse resolution

(Supplementary table S1). We found that elevation (Zimmerman & Vitousek, 2012), host plant phylogenetic

difference (Cobian, Egan, & Amend, 2019; Huang et al., 2016; Kembel & Mueller, 2014; Massimo et al., 2015;

Unterseher et al., 2012), geographic distance (U'Ren et al., 2012), and habitat type (Jumpponen & Jones, 2009,

2010; Kato et al., 2017) hypotheses were robust in our analysis at these large scales.

AUTHOR CONTRIBUTIONS

JLD wrote the paper, performed bioinformatic and statistical analyses, and created figures. ASA and BAP

conceived of the experiment. SOS, GC, and GLZ collected data and performed laboratory work. All authors

contributed to the preparation of the manuscript.

ACKNOWLEDGEMENTS

This work was supported by NSF award #1255972 to Amend, NSF award #1256128 to Perry and an NSF GRFP

award to Cobian. The authors thank D Hemmes, J Adams, V Costello, E Datlof, S Walsh, K Kimball, M Purell, T

Flynn, A Williams, R O'Rorke, S Perlman, P Biley and K Coelho for help in the field, and L Tooman for

assistance in the lab. The authors additionally thank C Lozupone and J Foquier for helpful discussions. The

authors declare that no conflict of interest exists.

FIGURES

586
587
588

590
Figure 1: Map of sample locations across the Hawaiian Archipelago. Samples were collected from the five
major islands in the Hawaiian archipelago: Hawaiʻi, Maui, Molokaʻi, Oʻahu, and Kauaʻi. Sampling was most
dense on Hawaiʻi and on Oʻahu islands, where accessibility was easiest. Our sampling strategy was to use
elevational transects where possible, in order to capture elevational and climatic variation. Elevations in
figure legend are meters above sea level. This is visible in the transects on Hawaiʻi, Molokaʻi, Kauaʻi, and
Oʻahu, which run orthogonally to the topographic lines (white). Transects are less pronounced on Maui
because of limited accessibility.
598

600
601 Figure 2: Ranges and distributions of explanatory variables. In this figure, each variable's distribution across 722
602 samples is shown as a smoothed density curve between its range (numbers to left and right of curves, height of
603 curve is relative frequency of observation). Units for range values are shown in Supplementary table S1. Each
604 variable included in our analysis covered a wide range of environmental heterogeneity.
605
606

Figure 3: Semivariogram and Mantel correlations of FEF beta diversity. This semivariogram (Bachmaier & Backes, 2008) was constructed using geographic distance classes, each containing the same number of points and each sharing two-thirds of its points with the previous class. Dots indicate the mean geographic distance within each class, and horizontal lines indicate its geographic range. The black curve is a logistic function fit to the data. Increasing semivariance indicates spatial autocorrelation. The lag in spatial autocorrelation before 7.6 km (first dashed line) indicates a lack of spatial structuring, and that samples at that range may be drawn from the same community. The plateau and subsequent noise at 36.3 km (second dashed line) is called the "autocorrelation distance", and is a typical feature of semivariogram analysis (King et al., 2010; Robeson et al., 2011). It indicates the distance after which spatial structuring becomes noisy or random, and geographic space is no longer a meaningful variable for community structure. The plot at bottom shows the data used to create the semivariogram. It also shows the Mantel correlation (r_M) of the three distance classes, as determined by 99% intervals of the logistic fit's derivative.

Figure 4: Model fit and coefficients for GDM model of FEF community dissimilarity. The observed
community dissimilarity (UniFrac distance) between pairwise samples exhibited a linear but noisy
relationship with the community dissimilarity predicted by the GDM model (top), which roughly
corresponded to a 1:1 line (dashed line). In the bottom plot, GDM i-splines are shown for statistically
significant predictor variables. Spline height indicates the relative importances of predictor variables, and
the spline's slope corresponds to the rate of change in compositional dissimilarity over the range of
pairwise dissimilarities within the variable. Host phylogeny and Evapotranspiration were the strongest
predictor variables in our analysis. These two variables were also the strongest correlates of beta diversity
principal coordinate vectors in our confirmatory PCoA analysis.

Figure 5: FEF specialization of each island. *d'* is a measure of how unique or cosmopolitan are the OTUs found on that island. In this violin plot, distributions shown in as filled violins with black borders are bootstrapped empirical *d'* values for each island. Unfilled distributions with a solid gray border are from the Vázquez null model , which is a fixed-connectance null model for bipartite networks. Welch's unequal variance *t*-tests show that each island's FEF community is significantly specialized as compared to the island null, meaning that FEF OTUs are more specific to their island of origin than expected by chance.

REFERENCES

Arnold, A. E., & Engelbrecht, B. M. J. (2007). Fungal endophytes nearly double minimum leaf conductance in seedlings of a neotropical tree species. *Journal of Tropical Ecology*, *23*(3), 369–372. doi: 10.1017/S0266467407004038

Arnold, A. E., & Lutzoni, F. (2007). Diversity and host range of foliar fungal endophytes: Are tropical leaves biodiversity hotspots? *Ecology*, *88*(3), 541–549. doi: 10.1890/05-1459

Bachmaier, M., & Backes, M. (2008). Variogram or semivariogram? Understanding the variances in a variogram. *Precision Agriculture*, *9*(3), 173–175. doi: 10.1007/s11119-008-9056-2

Bahnweg, G., Heller, W., Stich, S., Knappe, C., Betz, G., Heerdt, C., … Sandermann, H. (2005). Beech leaf colonization by the endophyte Apiognomonia errabunda dramatically depends on light exposure and climatic conditions. *Plant Biology*, *7*(6), 659–669. doi: 10.1055/s-2005-872943

Bayman, P., Angulo-Sandoval, P., Báez-Ortiz, Z., & Lodge, D. J. (1998). Distribution and dispersal of Xylaria endophytes in two tree species in Puerto Rico. *Mycological Research*, *102*(8), 944–948. doi: 10.1017/S095375629700590X

Bengtsson-Palme, J., Ryberg, M., Hartmann, M., Branco, S., Wang, Z., Godhe, A., … Nilsson, R. H. (2013). Improved software detection and extraction of ITS1 and ITS2 from ribosomal ITS sequences of fungi and other eukaryotes for analysis of environmental sequencing data. *Methods in Ecology and Evolution*, *4*(10), 914–919. doi: 10.1111/2041-210X.12073

Blackwell, M., Hibbett, D. S., Taylor, J. W., & Spatafora, J. W. (2006). Research Coordination Networks: a phylogeny for kingdom Fungi (Deep Hypha). *Mycologia*, *98*(6), 829–837. doi: 10.3852/mycologia.98.6.829

Blackwell, Meredith. (2011). The fungi: 1, 2, 3 … 5.1 million species? *American Journal of Botany*, *98*(3), 426–438. doi: 10.3732/ajb.1000298

Borcard, D., & Legendre, P. (2002). All-scale spatial analysis of ecological data by means of principal coordinates of neighbour matrices. *Ecological Modelling*. doi: 10.1016/S0304-3800(01)00501-4

Brooks, A. W., Kohl, K. D., Brucker, R. M., van Opstal, E. J., & Bordenstein, S. R. (2016). Phylosymbiosis: Relationships and Functional Effects of Microbial Communities across Host Evolutionary History. *PLoS Biology*, *14*(11), e2000225. doi: 10.1371/journal.pbio.2000225

Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A., & Holmes, S. P. (2016). DADA2: High-resolution sample inference from Illumina amplicon data. *Nature Methods*, *13*, 581–583. doi: 10.1038/nmeth.3869

Caporaso, J. G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F. D., Costello, E. K., … Knight, R. (2010). QIIME allows analysis of high-throughput community sequencing data. *Nature Methods*, *7*(5), 335–336. doi: 10.1038/nmeth.f.303

Christian, N., Sedio, B. E., Florez-Buitrago, X., Ramírez-Camejo, L. A., Rojas, E. I., Mejía, L. C., … Herre, E. A.
(2020). Host affinity of endophytic fungi and the potential for reciprocal interactions involving host
secondary chemistry. *American Journal of Botany*. doi: 10.1002/ajb2.1436

Cobian, G. M., Egan, C. P., & Amend, A. S. (2019). Plant-microbe specificity varies as a function of elevation.
The ISME Journal, (13), 2778–2778.

Coince, A., Cordier, T., Lengell??, J., Defossez, E., Vacher, C., Robin, C., … Mar??ais, B. (2014). Leaf and root-
associated fungal assemblages do not follow similar elevational diversity patterns. *PLoS ONE*, *9*(6),
e100668. doi: 10.1371/journal.pone.0100668

Darcy, J. L., King, A. J., Gendron, E. M. S., & Schmidt, S. K. (2017). Spatial autocorrelation of microbial
communities atop a debris-covered glacier is evidence of a supraglacial chronosequence. *FEMS*
Microbiology Ecology. doi: 10.1093/femsec/fix095

Darcy, J. L., Swift, S. O. I., Cobian, G. M., Zahn, G., Perry, B. A., & Amend, A. S. (2017). Foliar endophytic fungi
of native Hawaiian plants. *NCBI SRA*, PRJNA470970.

Davis, E. C., & Shaw, A. J. (2008). Biogeographic and phylogenetic patterns in diversity of liverwort-
associated endophytes. *American Journal of Botany*, *95*(8), 914–924. doi: 10.3732/ajb.2006463

Dormann, C. F., Gruber, B., & Fruend, J. (2008). Introducing the bipartite Package: Analysing Ecological
Networks. *R News*, *8*(2), 8–11.

Felber, A. C., Orlandelli, R. C., Rhoden, S. A., Garcia, A., Costa, A. T., Azevedo, J. L., & Pamphile, J. A. (2016).
Bioprospecting foliar endophytic fungi of Vitis labrusca Linnaeus, Bordô and Concord cv. *Annals of*
Microbiology, *66*(2), 765–775. doi: 10.1007/s13213-015-1162-6

Ferrier, S., Manion, G., Elith, J., & Richardson, K. (2007). Using generalized dissimilarity modelling to
analyse and predict patterns of beta diversity in regional biodiversity assessment. *Diversity and*
Distributions, *13*(3), 252–264. doi: 10.1111/j.1472-4642.2007.00341.x

Fitzpatrick, M. C., Sanders, N. J., Normand, S., Svenning, J. C., Ferrier, S., Gove, A. D., & Dunn, R. R. (2013).
Environmental and historical imprints on beta diversity: Insights from variation in rates of species
turnover along gradients. *Proceedings of the Royal Society B: Biological Sciences*, *280*(1768). doi:
10.1098/rspb.2013.1201

Fouquier, J., Rideout, J. R., Bolyen, E., Chase, J., Shiffer, A., McDonald, D., … Kelley, S. T. (2016). Ghost-tree:
Creating hybrid-gene phylogenetic trees for diversity analyses. *Microbiome*, *4*(11), 1–10. doi:
10.1186/s40168-016-0153-6

Giambelluca, T W, Shuai, X., Barnes, M. L., Alliss, R. J., Longman, R. J., Miura, T., … Businger, A. D. (2014).
Evapotranspiration of Hawai'i. Final report submitted to the U.S. Army Corps of Engineers—Honolulu
District, and the Commission on Water Resource Management, State of Hawaii.

Giambelluca, Thomas W., Chen, Q., Frazier, A. G., Price, J. P., Chen, Y. L., Chu, P. S., … Delparte, D. M. (2013).
Online rainfall atlas of Hawai'i. *Bulletin of the American Meteorological Society*, *94*(3), 313–316. doi:
10.1175/BAMS-D-11-00228.1

Giauque, H., & Hawkes, C. V. (2013). Climate affects symbiotic fungal endophyte diversity and performance.
American Journal of Botany, *100*(7), 1435–1444. doi: 10.3732/ajb.1200568

González-Teuber, M. (2016). The defensive role of foliar endophytic fungi for a South American tree. *AoB*
Plants, *8*, plw050. doi: 10.1093/aobpla/plw050

Hawksworth, D. L., & Rossman, A. Y. (1997). Where Are All the Undescribed Fungi? *Phytopathology*, *87*(9),
888–891. doi: 10.1094/PHYTO.1997.87.9.888

Higgins, K. L., Arnold, A. E., Coley, P. D., & Kursar, T. A. (2014). Communities of fungal endophytes in tropical
forest grasses: Highly diverse host- and habitat generalists characterized by strong spatial structure.
Fungal Ecology, *8*(1), 1–11. doi: 10.1016/j.funeco.2013.12.005

Hijmans, R. J., Etten, J. van, Mattiuzzi, M., Sumner, M., Greenberg, J. A., Lamigueiro, O. P., … Shortridge, A.
(2014). Package "raster." *R*, (2008), 1–27. doi: 10.1103/PhysRevE.74.016110.

Huang, Y.-L., Devan, M. M. N., U'Ren, J. M., Furr, S. H., & Arnold, A. E. (2016). Pervasive Effects of Wildfire on
Foliar Endophyte Communities in Montane Forest Trees. *Microbial Ecology*, *71*(2), 452–468. doi:
10.1007/s00248-015-0664-x

Jumpponen, A., & Jones, K. L. (2009). Massively parallel 454 sequencing indicates hyperdiverse fungal
communities in temperate *Quercus macrocarpa* phyllosphere. *New Phytologist*, *184*(2), 438–448. doi:
10.1111/j.1469-8137.2009.02990.x

Jumpponen, A., & Jones, K. L. (2010). Seasonally dynamic fungal communities in the Quercus macrocarpa
phyllosphere differ between urban and nonurban environments. *New Phytologist*, *186*(2), 496–513.
doi: 10.1111/j.1469-8137.2010.03197.x

Kato, S., Fukasawa, Y., & Seiwa, K. (2017). Canopy tree species and openness affect foliar endophytic fungal
communities of understory seedlings. *Ecological Research*, *32*(2), 157–162. doi: 10.1007/s11284-
016-1426-z

Keirle, M. R., Avis, P. G., Feldheim, K. A., Hemmes, D. E., & Mueller, G. M. (2011). Investigating the allelic
evolution of an imperfect microsatellite locus in the Hawaiian mushroom rhodocollybia laulaha.
Journal of Heredity. doi: 10.1093/jhered/esr099

Kembel, S. W., & Mueller, R. C. (2014). Plant traits and taxonomy drive host associations in tropical
phyllosphere fungal communities. *Botany*, *92*(4), 303–311. doi: 10.1139/cjb-2013-0194

King, A. J., Freeman, K. R., McCormick, K. F., Lynch, R. C., Lozupone, C., Knight, R., & Schmidt, S. K. (2010).
Biogeography and habitat modelling of high-alpine bacteria. *Nature Communications*, *1*(5), 53. doi:
10.1038/ncomms1055

Kivlin, S. N., Hawkes, C. V., & Treseder, K. K. (2011). Global diversity and distribution of arbuscular
mycorrhizal fungi. *Soil Biology and Biochemistry*, *43*(11), 2294–2303. doi:
10.1016/j.soilbio.2011.07.012

Lambers, H., Chapin, F. S., & Pons, T. L. (2008). Plant Water Relations. In *Plant Physiological Ecology* (pp.
163–223). doi: 10.1007/978-0-387-78341-3_5

Landesman, W. J., Nelson, D. M., & Fitzpatrick, M. C. (2014). Soil properties and tree species drive ß-
diversity of soil bacterial communities. *Soil Biology and Biochemistry*, *76*, 201–209. doi:
10.1016/j.soilbio.2014.05.025

Lewis, G. C., Ravel, C., Naffaa, W., Astier, C., & Charmet, G. (1997). Occurrence of Acremonium endophytes in
wild populations of Lolium spp. in European countries and a relationship between level of infection
and climate in France. *Annals of Applied Biology*, *130*(2), 227–238. doi: 10.1111/j.1744-
7348.1997.tb06828.x

Liu, J., Zhao, J., Wang, G., & Chen, J. (2019). Host identity and phylogeny shape the foliar endophytic fungal
assemblages of Ficus. *Ecology and Evolution*. doi: 10.1002/ece3.5568

Lozupone, C., & Knight, R. (2005). UniFrac: a New Phylogenetic Method for Comparing Microbial
Communities UniFrac: a New Phylogenetic Method for Comparing Microbial Communities. *Applied*
and Environmental Microbiology, *71*(12), 8228–8235. doi: 10.1128/AEM.71.12.8228

Lumibao, C. Y., Borer, E. T., Condon, B., Kinkel, L., May, G., & Seabloom, E. W. (2019). Site-specific responses
of foliar fungal microbiomes to nutrient addition and herbivory at different spatial scales. *Ecology*
and Evolution. doi: 10.1002/ece3.5711

Lunetta, R. S., Knight, J. F., Ediriwickrema, J., Lyon, J. G., & Worthy, L. D. (2006). Land-cover change detection
using multi-temporal MODIS NDVI data. *Remote Sensing of Environment*, *105*(2), 142–154. doi:
10.1016/j.rse.2006.06.018

Massimo, N. C., Nandi Devan, M. M., Arendt, K. R., Wilch, M. H., Riddle, J. M., Furr, S. H., … Arnold, A. E.
(2015). Fungal endophytes in aboveground tissues of desert plants: infrequent in culture, but highly
diverse and distinctive symbionts. *Microbial Ecology*, *70*(1), 61–76. doi: 10.1007/s00248-014-0563-6

Mejía, L. C., Herre, E. A., Sparks, J. P., Winter, K., García, M. N., Van Bael, S. A., … Maximova, S. N. (2014).
Pervasive effects of a dominant foliar endophytic fungus on host genetic and phenotypic expression
in a tropical tree. *Frontiers in Microbiology*, *5*, 479. doi: 10.3389/fmicb.2014.00479

Nilsson, R. H. (2011). Molecular Identification of Fungi: Rationale, Philosophical Concerns, and the UNITE
Database. *The Open Applied Informatics Journal*, *5*(1), 81–86. doi: 10.2174/1874136301105010081

Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGinn, D., … Wagner, H. (2016). *vegan:*
Community Ecology Package.

Oono, R., Lefèvre, E., Simha, A., & Lutzoni, F. (2015). A comparison of the community diversity of foliar
fungal endophytes between seedling and adult loblolly pines (Pinus taeda). *Fungal Biology*, *119*(10),
917–928. doi: 10.1016/J.FUNBIO.2015.07.003

Paradis, E., & Schliep, K. (2019). Ape 5.0: An environment for modern phylogenetics and evolutionary
analyses in R. *Bioinformatics*, *35*(3), 526–528. doi: 10.1093/bioinformatics/bty633

Pebesma, E., Rowlingson, B., & Rogerbivandnhhno, M. R. B. (2012). Package ' rgdal .' *R-CRAN*, *4*, 1–41. doi:
10.1353/lib.0.0050

Polonio, J. C., Almeida, T. T., Garcia, A., Mariucci, G. E. G., Azevedo, J. L., Rhoden, S. A., & Pamphile, J. A.
(2015). Biotechnological prospecting of foliar endophytic fungi of guaco (Mikania glomerata Spreng.)
with antibacterial and antagonistic activity against phytopathogens. *Genetics and Molecular Research*,
14(3), 7297–7309. doi: 10.4238/2015.July.3.5

Porras-Alfaro, A., & Bayman, P. (2011). Hidden Fungi, Emergent Properties: Endophytes and Microbiomes.
Annual Review of Phytopathology, *49*(1), 291–315. doi: 10.1146/annurev-phyto-080508-081831

Purevdorj, T. S., Tateishi, R., Ishiyama, T., & Honda, Y. (1998). Relationships between percent vegetation
cover and vegetation indices. *International Journal of Remote Sensing*, *19*(18), 3519–3535. doi:
10.1080/014311698213795

Qian, H., & Jin, Y. (2016). An updated megaphylogeny of plants, a tool for generating plant phylogenies and
an analysis of phylogenetic community structure. *Journal of Plant Ecology*, *9*(2), 233–239. doi:
10.1093/jpe/rtv047

R Core Team. (2014). R: A language and environment for statistical computing. *R Foundation for Statistical*
Computing, Vol. 1, p. 409. doi: 10.1007/978-3-540-74686-7

Robeson, M. S., King, A. J., Freeman, K. R., Birky, C. W., Martin, A. P., & Schmidt, S. K. (2011). Soil rotifer
communities are extremely diverse globally but spatially autocorrelated locally. *Proceedings of the*
National Academy of Sciences of the United States of America, *108*(11), 4406–4410. doi:
10.1073/pnas.1012678108

Rodriguez, R. J., White, J. F., Arnold, a E., & Redman, R. S. (2009). Fungal endophytes: diversity and
functional roles. *The New Phytologist*, *182*(2), 314–330. doi: 10.1111/j.1469-8137.2009.02773.x

Rognes, T., Flouri, T., Nichols, B., Quince, C., & Mahé, F. (2016). VSEARCH: a versatile open source tool for
metagenomics. *PeerJ Preprints*, *4*, e2409v1. doi: 10.7287/peerj.preprints.2409v1

Saucedo-García, A., Anaya, A. L., Espinosa-García, F. J., & González, M. C. (2014). Diversity and Communities
of Foliar Endophytic Fungi from Different Agroecosystems of Coffea arabica L. in Two Regions of
Veracruz, Mexico. *PLoS ONE*, *9*(6), e98454. doi: 10.1371/journal.pone.0098454

Smith, D. P., & Peay, K. G. (2014). Sequence depth, not PCR replication, improves ecological inference from
next generation DNA sequencing. *PLoS ONE*, *9*(2). doi: 10.1371/journal.pone.0090234

Stone, J. K., Bacon, C. W., & White, J. F. (2000). An overview of endophytic microbes: endophytism defined.
Microbial Endophytes, (January 2000), 3–29. doi: 10.1163/_q3_SIM_00374

Tardieu, F., Lafarge, T., & Simonneau, T. (1996). Stomatal control by fed or endogenous xylem ABA in
sunflower: interpretation of correlations between leaf water potential and stomatal conductance in
anisohydric species. *Plant, Cell and Environment*, *19*(1), 75–84. doi: 10.1111/j.1365-
3040.1996.tb00228.x

Tipton, L., Zahn, G., Datlof, E., Kivlin, S. N., Sheridan, P., Amend, A. S., & Hynson, N. A. (2019). Fungal
aerobiota are not affected by time nor environment over a 13-year time series at the Mauna Loa
Observatory. *PNAS*, *in press*.

U'Ren, J. M., & Arnold, A. E. (2016). Diversity, taxonomic composition, and functional aspects of fungal
communities in living, senesced, and fallen leaves at five sites across North America. *PeerJ*, *4*, e2768.
doi: 10.7717/peerj.2768

U'Ren, J. M., Lutzoni, F., Miadlikowska, J., Laetsch, A. D., & Arnold, A. E. (2012). Host and geographic
structure of endophytic and endolichenic fungi at a continental scale. *American Journal of Botany*,
99(5), 898–914. doi: 10.3732/ajb.1100459

U'Ren, J. M., Lutzoni, F., Miadlikowska, J., Zimmerman, N. B., Carbone, I., May, G., & Arnold, A. E. (2019). Host
availability drives distributions of fungal endophytes in the imperilled boreal realm. *Nature Ecology &*
Evolution, *3*, pages1430–1437. doi: 10.1038/s41559-019-0975-2

Unterseher, M., Petzold, A., & Schnittler, M. (2012). Xerotolerant foliar endophytic fungi of Populus
euphratica from the Tarim River basin, Central China are conspecific to endophytic ITS phylotypes of
Populus tremula from temperate Europe. *Fungal Diversity*, *54*(1), 133–142. doi: 10.1007/s13225-
012-0167-8

Vázquez, D. P., Melián, C. J., Williams, N. M., Blüthgen, N., Krasnov, B. R., & Poulin, R. (2007). Species
abundance and asymmetric interaction strength in ecological networks. *Oikos*, *116*(7), 1120–1127.
doi: 10.1111/j.0030-1299.2007.15828.x

Vincent, J. B., Weiblen, G. D., & May, G. (2016). Host associations and beta diversity of fungal endophyte
communities in New Guinea rainforest trees. *Molecular Ecology*, *25*(3), 825–841. doi:
10.1111/mec.13510

Ward, T. L., Dominguez-Bello, M. G., Heisel, T., Al-Ghalith, G., Knights, D., & Gale, C. A. (2018). Development
of the Human Mycobiome over the First Month of Life and across Body Sites. *MSystems*, *3*(3), e00140-
17. doi: 10.1128/mSystems.00140-17

Warren, D. L., Cardillo, M., Rosauer, D. F., & Bolnick, D. I. (2014). Mistaking geography for biology: Inferring
processes from species distributions. *Trends in Ecology and Evolution*, Vol. 29, pp. 572–580. doi:
10.1016/j.tree.2014.08.003

Wiens, J. J., Ackerly, D. D., Allen, A. P., Anacker, B. L., Buckley, L. B., Cornell, H. V., … Stephens, P. R. (2010).
Niche conservatism as an emerging principle in ecology and conservation biology. *Ecology Letters*,
13(10), 1310–1324. doi: 10.1111/j.1461-0248.2010.01515.x

Ziegler, A. C. (2002). Hawaiian Natural History, Ecology, and Evolution. In *Hawaiian Natural History,*
Ecology, and Evolution. doi: 10.1515/9780824842437

Zimmerman, N. B., & Vitousek, P. M. (2012). Fungal endophyte communities reflect environmental
structuring across a Hawaiian landscape. *Proceedings of the National Academy of Sciences*, *109*(32), 13022–
13027. doi: 10.1073/pnas.1209872109

851
852
853 **DATA ACCESSIBILITY STATEMENT**
854
DNA sequence data have been submitted to the NCBI Sequence Read Archive (SRA) database under BioProject
accession number PRJNA470970 (Darcy, Swift, et al., 2017). Computer code and input files that can be used to

replicate this analysis are available at FigShare: https://figshare.com/s/34a59ebd8051589734c1 (temporary

private link until publication). Additionally, a simple summary of the four most commonly sampled plant genera

for each island is available as Supplementary table S4. DNA sequence data are available from both NCBI SRA

and FigShare repositories.