Three-dimensional morphological analysis of the dynamic digestive system in the green brittle

star

Daiki Wakita[1], Keisuke Naniwa[2], and Hitoshi Aonuma[1,2*]

[1]Graduate School of Life Science, Hokkaido University, Sapporo 0600810, Japan

[2]Research Institute for Electronic Science, Hokkaido University, Sapporo 0600812, Japan

[*]Corresponding author: Hitoshi Aonuma, D.Sc.,

Research Institute for Electronic Science, Hokkaido University, Sapporo 0600812, Japan.

Tel/Fax: +81-117063832, Email: aon@es.hokudai.ac.jp

ORCID:

Daiki Wakita, 0000-0001-8653-4094

Keisuke Naniwa, 0000-0003-3171-958X

Hitoshi Aonuma, 0000-0001-8380-1820

Abstract

Brittle stars (Echinodermata: Ophiuroidea) digest a great diversity of food in their stomach, which

widely lies in the central disk. As for a possible digestive activity, the green brittle star *Ophiarachna*

incrassata (Lamarck, 1816) is known to show a dynamic movement at the disk. This phenomenon

would deeply involve the morphological structure of the stomach. However, past anatomical studies

have shown the digestive system in two dimensions after wide incision of the body wall anchoring the

stomach. This methodology restrains us from understanding how the stomach actually shapes inside a

brittle star. We aim to visualize the morphology of brittle stars' digestive system in a non-destructive

and three-dimensional way, with a comparison between a relaxed specimen and a specimen fixed at

the very moment of the disk's movement. Employing X-ray micro-computed tomography (micro-CT)

and introducing an instant freezing method with cryogenic ethanol, we found the stomach wholly

transformed during the movement. We here brought transparency to the *in vivo* position of gut

contents to hint the mechanism and digestive function of the movement. Our outcome spotlights a

dynamic digestive process in echinoderms and a widely applicable method for probing into its relation

with body structure.

Introduction

More than 2,300 species of brittle stars (Echinodermata: Ophiuroidea) are known worldwide and

constitute the largest class among extant echinoderms (Stöhr et al. 2019). Diversity in feeding habits

could be an explanatory factor for the current success of this group (Fontaine 1965). Their food varies

in sort and scale, from sediment or small organisms including diatoms, dinoflagellates, foraminifera,

and copepods, to the whole or part of large organisms such as polychaetes, bivalves, crabs, fish, other

echinoderms, and sessile algae (Nagabhushanam and Colman 1959; Fontaine 1965; Hendler and

Miller 1984; Pearson and Gage, 1984; Ambrose 1993). Their radially extending arms play a role in

capturing food (Fontaine 1965; Warner 1971; Reimer and Reimer 1975; Hendler and Miller 1984), and

then internal digestive organs take center stage.

Contrary to the various targets in feeding, brittle stars share the general structure of the

digestive system. Its mouth is followed in order by buccal cavity, pharynx, esophagus, and stomach

(Schechter and Lucero 1968). Lacking intestine and anus, it terminates with the stomach, which

occupies a large space inside the disk—central part of the body (Smith 1940; Schechter and Lucero

1968; Pentreath 1971; Uchida and Irimura 1974). The stomach consists of a single sac radially

dividing into 10 pouches: five radial (ambulacral) pouches and five interradial (interambulacral)

pouches (Pentreath 1969). We also read a contradicting statement that brittle stars' stomach totally has

15 swellings in a common context (Uchida and Irimura 1974). Interradial pouches are well-folded

structure deeply lying between the bases of arms, whereas radial pouches are limited in narrow spaces

over arms (Pentreath 1969, 1971; Uchida and Irimura 1974). These pouches never extend into arms

except the species *Ophiocanops fugiens* Koehler, 1922 (Fell 1963).

Over and above anatomical studies, a dynamic perspective of digestion has been discussed

based on live observation. In the green brittle star *Ophiarachna incrassata*, Wakita et al. (2018)

reported a rhythmic movement termed "*pumping*" (Online Resource 1; c.f. Fig. 1), which is frequently

observed at the disk after feeding. The series of expansion and shrinkage can be recognized as a sort of

peristaltic movement of the stomach. Its unique coordinated patterns are explainable by assuming

internal fluid flows, the way of which depends on the morphology of the disk. In particular, five-armed

individuals make unsynchronized movements between five body parts, whereas those are well

synchronized in a six-armed case—peculiar individual difference in brittle stars. Thus, in this

phenomenon, the morphological structure of the stomach may be of great importance, although it is

unclear how pumping transforms it.

In previous studies, the morphological structure of the digestive system in brittle stars has

been visualized only by two-dimensional sketches or tissue sections (Smith 1940; Pentreath 1969,

1971; Schechter and Lucero 1968; Uchida and Irimura 1974; Frolova and Dolmatov 2006, 2010).

Moreover, traditional anatomy has employed a wide dissection of the body wall, which considerably

distorts the morphology in focus. This issue arises from the fragileness of digestive organs as well as

its deep attachment to the body wall by collagen strands (Schechter and Lucero 1968; Uchida and

Irimura 1974). The aim of our study is to visualize non-destructive and three-dimensional (3D)

morphological structure of the digestive system in brittle stars, in comparative terms of a relaxed

condition and a condition during pumping. For this purpose, we employ X-ray micro-computed

tomography (micro-CT) while introducing an instant freezing method using cryogenic ethanol for

making a *snapshot* of the dynamically moving body (Fig. 1). We also probe into the internal

morphology of the six-armed specimen studied by Wakita et al. (2018), so as to internally validate the

prior assumption that this specific individual has six symmetrical units—previously it was made up

merely in external terms. The primary conclusion in our study is that pumping could transform the

entire stomach to help digestion in large brittle stars, with its coordinated patterns apparently reflecting

five- or six-fold symmetrical arrangement in internal morphology.

Fig. 1   Frozen individual at the moment of the dynamic movement "pumping" in the green brittle star

Ophiarachna incrassata. (a) Oral view. (b) Aboral view. Photos were taken just after pouring –80°C ethanol

while we observed pumping at the disk. Arrowheads denote well-expanding portions. Scale bars represent 10

mm. Scanned and segmented images of this specimen are shown in Fig. 2. Video of another individual's

pumping is shown in Online Resource 1

91   **Materials and methods**

Animals

Individuals of the green brittle star *Ophiarachna incrassata* were obtained commercially (Aqua Shop

Saien, Sapporo, Japan) and reared in aquariums ($600 \times 600 \times 600$ mm) filled with artificial seawater at

25–28°C with the salinity of 32–35‰ (TetraMarin Salt Pro, Tetra Japan Co, Tokyo, Japan). They were

fed with dried krill (Tetra Krill-E, Tetra Japan Co, Tokyo, Japan).

X-ray micro-computed tomography

With the aim to investigate relaxed and pumping bodies as well as five- and six-armed bodies, we

chose three individuals: (1) a five-armed individual a day after feeding, with the disk diameter of 15

mm in a living anesthetized condition; (2) a five-armed individual a week after feeding, with the disk

diameter of 18 mm; (3) a six-armed individual with the diameter of 25 mm, which had hardly shown

feeding behavior for a few months. The six-armed one corresponds to that studied by Wakita et al.

(2018). For the relaxed case, the animals (1) and (3) each were anaesthetized in 3% $MgCl_2$ solution for

an hour at room temperature and then fixed with Bouin solution—(1) for 10 days and (3) for five

months—at 3°C with their arms cut near the bases. For looking into the pumping body, the animal (2)

was put in a styrofoam box ($127 \times 157 \times 100$ mm) with 100–200 ml of artificial seawater and then fed

with a dried krill. While we observed the rhythmic movement (pumping; c.f. Online Resource 1),

−80°C ethanol was poured onto the disk so that the animal (2) kept a momentary shape with expansion

and shrinkage (Fig. 1). After slowly shaking the box for 10 min, the sample was put in Bouin solution

for a day at 3°C with their arms cut.

After fixation, all the samples (1)–(3) were dehydrated with ethanol series (70%, 80%, and

90%) for two days each, and stained with 1% iodine diluted in ethanol for three days at 3°C to

enhance the contrast of tissues in later X-ray exposure (Metscher 2009). They were rinsed with 100%

ethanol for a day at room temperature and then moved into *t*-butyl alcohol liquidized with a water bath

above 40°C. After immersion in *t*-butyl alcohol for a day twice at 26°C, samples were superficially

dried on tissues for several seconds and then put at −20°C for 10 min so that instantly frozen *t*-butyl

alcohol would keep the original morphology as possible. They were freeze-dried by using a vacuum

evaporator (PX-52, Yamato Ltd., Japan) with a cold alcohol trap (H2SO5, AS ONE, Japan). All

chemicals were purchased from Kanto Chemical Co. (Tokyo, Japan).

Samples were scanned on an X-ray micro-CT system (inspeXio, SMX-100CT, Shimadzu

Corporation, Kyoto, Japan), where X-ray source was operated at 75 kV and 40 µA. Scanned images

were reconstructed and rendered by using VGStudio MAX ver. 2.2.6 (Volume Graphics, Heidelberg,

Germany) with the voxel size of 10–50 µm. For segmentation of each sample using Amira ver. 2019.1

(Thermo Scientific, Waltham, USA), we traced the inner surface—boundary with contrasting X-ray

absorptivity—of the digestive cavity beginning from the mouth. Note that segmentation was not

available in regions where the inner wall stuck to each other so that the cavity was too flat to be

identified. We also segmented gut contents, which were recognizable as highly absorptive areas inside

the cavity. 3D animations were created with VGStudio MAX for slice images and Amira for

segmented images, which are given in Online Resources 2–4.

134   **Results**

The morphology of the digestive system, particularly the stomach, was well visualized by segmenting

the inner wall of the digestive cavity (Figs. 2–4). In the five-armed specimen fixed after anesthesia

(Online Resource 2), skeletal structure comprised five symmetrical sectors in appearance (Fig. 2a).

The cavity's surface was smoothly defined from the mouth to the stomach (Fig. 2b), so we required

less subjectivity in the segmentation (Fig. 2c–e). The stomach comprised five larger interradial

pouches and five smaller radial pouches (Fig. 2c). As noted in "Materials and methods", no region in a

totally flat cavity was segmented, hence the gaps forming a cobweb-like structure could be interpreted

as the missing flattened parts of the stomach, not representing there were many holes in morphology

(Fig. 2c). The stomach was plain in the aboral side (Fig. 2c, d) but well wrinkled in the oral (Fig. 2e).

Viewed orally, a ridge could be traced along each midline of two sorts of pouches, which descended

into several branches (Fig. 2e). Interradial pouches narrowed at the bases with their breadth increasing

distally, with the end being round so that we could trace smoothly between the oral and aboral surfaces

(Fig. 2c–e). Radial pouches were more flat, shaped along arm skeletal plates, and made distal ends

with rough and sharp edges (Fig. 2d, e). The distal parts of interradial pouches extended until near the

oral wall, whereas their bases and radial pouches were restricted aborally (Fig. 2d). The oral room at

the center contained a jaw apparatus and its peripheral organs, which would include the circumoral

nerve ring and the water vascular system (Fig. 2a, b). In this specimen, two large pieces of food were

respectively observed at the aboral base of an interradial pouch—might be partially shared by one

adjacent radial pouch—and the oral end of another interradial (Fig. 2c–e).

Fig. 2 Three-dimensional visualization of the digestive system in a five-armed relaxed individual of the green

brittle star *Ophiarachna incrassata*. Body structure was scanned with X-ray micro-computed tomography

(micro-CT) and reconstructed in three dimensions, which is displayed in grayscale. The inner surface of the

digestive cavity beginning from the mouth is colored blue. Contents in the stomach are colored red. Regions

where the inward cavity was totally flat were not segmented (not colored blue) with a technical limitation, which

reflects the apparent holes of the stomach and the apparent exposure of gut contents. (a) Reconstructed images

viewed from the oral side, sectioned at a plane shown in (b) by the dotted line. (b) Oral-aboral section on a plane

indicated in (a) by the dotted line; the bottom is oral; slab thickness is 0.38 µm. (c) Aboral view of the

segmented model. (d) Lateral view of the segmented model from the side indicated in (c) by the arrow; the

bottom is oral; the grayscale images are truncated in the front for clarity. (e) Enlarged oral view of the segmented

model. Abbreviations: a, arm skeleton; b, body wall; e, esophagus; f, food (gut content); m, mouth; ir, interradial

pouch; r, radial pouch. Scale bars represent 5 mm. Three-dimensional animation is shown in Online Resource 2

In the specimen frozen during pumping (Online Resource 3), we found no noticeable

damage due to the instant freezing method in external and internal morphology (Figs. 1, 3a). Besides,

there seemed to be no large difference in the texture of the digestive wall in scanned images, compared

to those observed in the relaxed one. We recognized a well-defined separation between the mouth and

the stomach and a tight closure of the mouth (Fig. 3b; see the esophagus "e" and the mouth "m"), so

we did not clearly identify the continuous space from the mouth opening. However, we had no

ambiguity in segmenting the internal surface which was recognizable as the stomach's one (Fig. 3c–e)

when comparing to the relaxed case (Fig. 2c–e). The stomach wall during pumping was smoothly

fitted to the distorted body wall with less obvious folding (Fig. 3c). The globular shaping of the

stomach was also upheld from the observation that we saw no network-like segmentation denoting

flattened patches (Fig. 3c), contrasting to the relaxed one (Fig. 2c). Expansions were remarkable in

interradial pouches, which directed toward the aboral and lateral sides (Fig. 3d). Meanwhile, radial

pouches at their neighbors also became more or less open (Fig. 3d). The oral surface likewise showed

an entire extension, where we barely found sharp structure such as branched ridges (Fig. 3e). The

cavity between the body and stomach walls (perivisceral coelom) appeared to be narrowly limited (Fig.

3b, d), not largely differing from the relaxed case (Fig. 2b, d). A food lied almost at the center of the

stomach in this specimen (Fig. 3c, e).

Fig. 3 Three-dimensional visualization of the digestive system in a five-armed individual during the dynamic movement "pumping" in the green brittle star *Ophiarachna incrassata*. Body structure of an instantly frozen individual shown in Fig. 1 was scanned with X-ray micro-computed tomography (micro-CT) and reconstructed in three dimensions, which is displayed in grayscale. The inner wall of the digestive cavity is colored blue, which was identified in comparison with the relaxed specimen (Fig. 2). Contents in the stomach are colored red. Regions where the inward cavity was totally flat were not segmented (not colored blue) with a technical limitation, which reflects the apparent exposure of gut contents. Arrowheads denote well-expanding portions. (a) Reconstructed images viewed from the oral side, sectioned at a plane shown in (b) by the dotted line. (b) Oral-aboral section on a plane indicated in (a) by the dotted line; the bottom is oral; slab thickness is 0.36 µm. (c) Aboral view of the segmented model. (d) Lateral view of the segmented model from the side indicated in (c) by the arrow; the bottom is oral; the grayscale images are truncated in the front for clarity. (e) Enlarged oral view of the segmented model. Abbreviations: a, arm skeleton; b, body wall; e, esophagus; f, food (gut content); m, mouth; ir, interradial pouch; r, radial pouch. Scale bars represent 5 mm. Three-dimensional animation is shown in Online Resource 3

In the specimen with six arms (Online Resource 4), a jaw apparatus and arm skeletal plates

apparently arranged in six-fold radial symmetry (Fig. 4a). As in the five-armed relaxed case, the

digestive cavity was smoothly traceable from the mouth to the stomach (Fig. 4b). Though the

segmentation, we realized that the inner openings of interradial pouches were almost as flat as those of

radial ones (Fig. 4b). This reduction was probably because this individual had not fed for a few

months before fixation. Although the boundaries between radial and interradial pouches were less

conspicuous (Fig. 4c) than the five-armed ones (Fig. 2c), the two types could be distinguished when

we saw the model from several angles (Fig. 4b, c). In particular, interradial pouches hung down more

orally (Fig. 4b) and showed ridged midlines on the oral surface (Fig. 4c). Here we could count 12

pouches—six interradial and six radial pouches (Fig. 4c).

Fig. 4 Three-dimensional visualization of the digestive system in a six-armed relaxed individual of the green

brittle star *Ophiarachna incrassata*. Body structure was scanned with X-ray micro-computed tomography

(micro-CT) and reconstructed in three dimensions, which is displayed in grayscale. The inner wall of the

digestive cavity beginning from the mouth is colored blue. Regions where the inward cavity was totally flat were

not segmented (not colored blue) with a technical limitation. (a) Reconstructed images viewed from the oral side,

sectioned at a plane indicated in (b) by the dotted line. (b) Lateral view of the segmented model from the side

indicated in (c) by the arrow; the bottom is oral; the grayscale images are truncated in the front for clarity. (c)

Oral view of the segmented model. Abbreviations: a, arm skeleton; b, body wall; ir, interradial pouch; r, radial

pouch. Scale bars represent 5 mm. Three-dimensional animation is shown in Online Resource 4

Discussion

Our study has three main achievements. The first is 3D visualization of the uninjured stomach in

brittle stars (Fig. 2, Online Resource 2). The schematics were directly reconstructed from micro-CT

scanned images, including less imagination than previous sketches. The second is introduction of a

fresh methodology in micro-CT scanning, where we investigated a moving body's momentary shape

made by instant freezing with cryogenic ethanol (Fig. 3, Online Resource 3). The resultant snapshot

provides a further understanding of the previously reported phenomenon, *pumping* (Wakita et al.,

2018), as a possible digestive activity. The last is internal inspection of the specific six-armed

individual studied by Wakita et al. (2018) (Fig. 4, Online Resource 4). A supernumerary seemed to be

simply a member of six equivalents, which helps a prebuilt assumption on pumping coordination.

233   In a five-armed representative, we counted 10 major swellings in the stomach (Fig. 2c) as

Pentreath (1969) reported, not supporting another literature writing the total number 15 (Uchida and

Irimura 1974). Since the latter also mentions radial and interradial pouches, it might be miswriting or

another interpretation given the complex folding. The wrinkles running on the oral surface would be a

reflection in part of skeletal morphology and other organs' anchoring, while probably being capable of

a large amount of food by its extension. In fact, the specimen shown in Fig. 3 accommodates a

relatively large food ranging across several pouches. Although the segmented stomach structure in Fig.

2 looks like a web network, we can interpret that these holes as untraceable flat parts of the cavity so

that pouches connect with each other throughout the central space.

242   Comparison between the relaxed and pumping specimens in scanned images (Figs. 2, 3)

indicates that pumping dynamically transforms the stomach, where it is natural to suppose internal

fluid flows to some extent. The outer space of the stomach (perivisceral coelom) would make

insignificant flows for pumping, considering the constant narrowness even in the pumping body (Fig.

3b, d). The mouth-stomach separation emerging in the pumping specimen (Fig. 3b) can be defined as

the constriction of esophagus, referring to Schechter and Lucero (1968). This observation and the tight

closure of the jaw apparatus—surrounding buccal cavity—(Fig. 3b) both would reinforce Wakita et

al.'s (2018) assumption that the total fluid volume is constant during pumping; there is no outward

leakage. Although their study built a water-connecting network with five nodes for the five-armed case,

there could be 10 rooms given the number of pouches (Fig. 2c). However, the slits between the

pouches were actually not deep as depicted in the pumping specimen (Fig. 3c). With this texture, the

five-fold arm skeleton would rather work as partitions (Fig. 3a, b), making it reasonable to represent

five rooms influencing the major behavior of internal flows. This explanation could also apply to the

six-armed case. The rigid skeletal partitions symmetrically made by six arms (Fig. 4a) would be more

dominant in fluid flow, compared to the flexibly transformable stomach with 12 pouches (Fig. 4c).

Our scan also supports Wakita et al.'s (2018) explanation with six symmetrical nodes for this

six-armed specimen. We thus retain the explanatory power of the pumping network with five or six

symmetrical nodes—not 10 or 12 nodes in two different sizes.

Visibility in the original position of gut contents gives two suggestions for the phenomenon

pumping. The first is about its initiation. After a brittle star eats something, food fragments would be

seated at some pouches (Fig. 2c–e); even if a prey stays at the center, its body shape would never

weight equally among all the pouches (Fig. 3c). The contents thus make the stomach morphology

more asymmetric, which might trigger the initiation of a pumping series as Wakita et al. (2018) let one

interradial volume unequal at the beginning of simulation. The second involves the purpose of

pumping. In many animals including humans, food transfers from the mouth to the anus in one

direction; the linear structure guarantees that nutrients are absorbed point by point. On the other hand,

the digestive cavity of brittle stars has neither unidirectional tracts nor the anus, so a piece might easily

stick to a dead end—just as shown in Fig. 2. Transformation of the stomach by pumping would give

more opportunities to displace the piece with it spreading a nutritious flow. This strategy would be

effective in large-sized species, where gut contents travel longer distances piece by piece. Therefore,

although we used the single species *Ophiarachna incrassata*, other large brittle stars are supposed to

exhibit pumping in a similar manner.

CT scanning technique for 3D visualization has been employed by several studies on brittle

stars. Landschoff and Griffiths (2015) revealed how a brooding brittle star accommodates several

juveniles inside its body, comparing two species; another was similarly examined later (MacKinnon et

al. 2017). In a taxonomic context, Okanishi et al. (2017) described skeletal structure of a euryalid

brittle star without dissolving its thick skin. Clark et al. (2018) paid attention to the joint connection of

vertebrae to understand the mobility of arms in two species. Our study would carry a novelty in (1)

focusing on the digestive system, (2) comparing two behavioral conditions (relaxed v.s. pumping), and

(3) comparing two morphologically different individuals within a species (five-armed v.s. six-armed)

in CT scanned brittle stars. Notably, the snapshotting method for (2), where cryogenic ethanol is

poured onto a living animal, is widely applicable for the scanning purpose of body structure during

dynamic movements in echinoderms. These approaches give prominence to a dynamic digestive

process and its relation with body structure, which would be a hot clue to ethology, ecology, and

evolution in echinoderms.

Acknowledgments   This work was partly supported by JSPS KAKENHI (Grant Number 16KT0099),

JST CREST (Grant Number JPMJCR14D5), and Hokkaido University Frontier Foundation (Nitobe

School Financial Assistance), Japan.

Funding   This study was funded by JSPS KAKENHI (grant number 16KT0099), JST CREST (grant

number JPMJCR14D5), and Hokkaido University Frontier Foundation (Nitobe School Financial

Assistance), Japan.

Data availability   The datasets obtained during the current study are available from the

corresponding author on reasonable request.

301   **Compliance with ethical standards**

303   **Conflict of interest**   The authors declare that they have no conflict of interest.

305   **Ethical approval**   This article does not contain any studies with human participants performed by

306   any of the authors.

309   **References**

311   Ambrose WG Jr (1993) Effects of predation and disturbance by ophiuroids on soft-bottom community

312   structure in Oslofjord: results of a mesocosm study. Mar Ecol Prog Ser 97:225–236.

313 https://doi.org/10.3354/meps097225

314 Clark EG, Hutchinson JR, Darroch SA, Mongiardino Koch N, Brady TR, Smith SA, Briggs DE (2018)

315  Integrating morphology and in vivo skeletal mobility with digital models to infer function in

316  brittle star arms. J Anat 233:696–714. https://doi.org/10.1111/joa.12887

317 Fell HB (1963) The phylogeny of sea-stars. Philos T R Soc B 246:381–435.

318  https://doi.org/10.1098/rstb.1963.0010

319 Fontaine AR (1965) The feeding mechanisms of the ophiuroid *Ophiocomina nigra*. J Mar Biol Assoc

320  UK 45:373–385. https://doi.org/10.1017/S0025315400054904

321 Frolova LT, Dolmatov IY (2006) Regeneration of the epithelial lining of the stomach after autotomy of

322  a disk in the brittle star *Amphipholis kochii* (Lütken) (Echinodermata: Ophiuroidea). Russ J Mar

323  Biol 32:68–70. https://doi.org/10.1134/S106307400601010X

324 Frolova LT, Dolmatov IY (2010) Microscopic anatomy of the digestive system in normal and

325  regenerating specimens of the brittlestar *Amphipholis kochii*. Biol Bull 218:303–316.

326  https://doi.org/10.1086/BBLv218n3p303

327 Hendler G, Miller JE (1984) Feeding behavior of *Asteroporpa annulata*, a gorgonocephalid brittlestar

328  with unbranched arms. B Mar Sci 34:449–460. https://doi.org/10088/8751

329 Lamarck JD (1816) Ordre second: radiaires échinodermes. In: Lamarck JD (ed) Histoire Naturelle des

330  Animaux sans Vertèbres 2. Verdière, Libraire, quai des Augustins, No. 27, Paris, pp. 522–568.

331  https://doi.org/10.5962/bhl.title.12712

332 Landschoff J, Griffiths CL (2015) Three-dimensional visualisation of brooding behaviour in two

333  distantly related brittle stars from South African waters. Afr J Mar Sci 37:533–541.

334  https://doi.org/10.2989/1814232X.2015.1095801

335 MacKinnon RB, Landschoff J, Griffiths CL (2017) Seasonality and 3D-visualization of brooding in

336   the hermaphroditic ophiuroid *Amphiura capensis*. Invertebr Biol 136:146–158.

337   https://doi.org/10.1111/ivb.12164

338 Metscher BD (2009) MicroCT for comparative morphology: simple staining methods allow

339   high-contrast 3D imaging of diverse non-mineralized animal tissues. BMC Physiol 9:11.

340   https://doi.org/10.1186/1472-6793-9-11

341 Nagabhushanam AK, Colman JS (1959) Carrion-eating by ophiuroids. Nature 184:285.

342   https://doi.org/10.1038/184285a0

343 Okanishi M, Fujita T, Maekawa Y, Sasaki T (2017) Non-destructive morphological observations of the

344   fleshy brittle star, *Asteronyx loveni* using micro-computed tomography (Echinodermata,

345   Ophiuroidea, Euryalida). ZooKeys 663:1–19. https://doi.org/10.3897/zookeys.663.11413

346 Pearson M, Gage JD (1984) Diets of some deep-sea brittle stars in the Rockall Trough. Mar Biol

347   82:247–258. https://doi.org/10.1007/BF00392406

348 Pentreath RJ (1969) The morphology of the gut and a qualitative review of digestive enzymes in some

349   New Zealand ophiuroids. J Zool 159:413–423.

350   https://doi.org/10.1111/j.1469-7998.1969.tb03898.x

351 Pentreath RJ (1971) Respiratory surfaces and respiration in three New Zealand intertidal ophiuroids. J

352   Zool 163:397–412. https://doi.org/10.1111/j.1469-7998.1971.tb04540.x

353 Reimer RD, Reimer AA (1975) Chemical control of feeding in four species of tropical ophiuroids of

354   the genus *Ophioderma*. Comp Biochem Phys A 51:915–927.

355   https://doi.org/10.1016/0300-9629(75)90075-4

356 Schechter J, Lucero J (1968) A light and electron microscopic investigation of the digestive system of

357   the ophiuroid *Ophiuroiderma panamensis* (brittle star). J Morphol 124:451–481.

358   https://doi.org/10.1002/jmor.1051240405

Smith J (1940) The reproductive system and associated organs of the brittle-star *Ophiothrix fragilis*. J

Cell Sci 2:267–309

Stöhr S, O'Hara T, Thuy B (2019) World Ophiuroidea Database.

http://www.marinespecies.org/ophiuroidea. Accessed 29 April 2019.

https://doi.org/10.14284/358

Uchida T, Irimura S (1974) Ophiuroidea. In: Uchida T (ed) The Systematic Zoology 8B. Nakayama

Book Company, Tokyo, pp 142–207

Wakita D, Hayase Y, Aonuma H (2018) Five breaks synchrony while six keeps synchrony: individual

difference in the coordinated pattern of five- and six-armed brittle stars. BioRxiv 340471.

https://doi.org/10.1101/340471

Warner GF (1971) On the ecology of a dense bed of the brittle-star *Ophiothrix fragilis*. J Mar Biol

Assoc UK 51:267–282. https://doi.org/10.1017/S0025315400031775

373   **Electronic supplementary material**

375 **Online Resource 1** Rhythmic movement "pumping" in a five-armed individual of the green brittle

376 star *Ophiarachna incrassata*.

378 **Online Resource 2** 3D animation of the images shown in Fig. 2 (five-armed relaxed individual).

380 **Online Resource 3** 3D animation of the images shown in Fig. 3 (five-armed pumping individual).

382   **Online Resource 4**   3D animation of the images shown in Fig. 4 (six-armed relaxed individual).