Complex ecological phenotypes on phylogenetic trees: a hidden Markov model for comparative

analysis of multivariate count data

Michael C. Grundler[1,*] and Daniel L. Rabosky[1]

¹Museum of Zoology and Department of Ecology and Evolutionary Biology, University of

Michigan, Ann Arbor, MI USA 48109

Corresponding author: mgru@umich.edu

ABSTRACT

The evolutionary dynamics of complex ecological traits – including multistate

representations of diet, habitat, and behavior – remain poorly understood. Reconstructing the

tempo, mode, and historical sequence of transitions involving such traits poses many challenges

for comparative biologists, owing to their multidimensional nature and intraspecific variability.

Continuous-time Markov chains (CTMC) are commonly used to model ecological niche

evolution on phylogenetic trees but are limited by the assumption that taxa are monomorphic and

that states are univariate categorical variables. Thus, a necessary first step when using standard

CTMC models is to categorize species into a pre-determined number of ecological states. This

approach potentially confounds interpretation of state assignments with effects of sampling

variation because it does not directly incorporate empirical observations of resource use into the

statistical inference model. The neglect of sampling variation, along with univariate

representations of true multivariate phenotypes, potentially leads to the distortion and loss of

information, with substantial implications for downstream macroevolutionary analyses. In this

study, we develop a hidden Markov model using a Dirichlet-multinomial framework to model

resource use evolution on phylogenetic trees. Unlike existing CTMC implementations, states are

unobserved probability distributions from which observed data are sampled. Our approach is

expressly designed to model ecological traits that are intra-specifically variable and to account

for uncertainty in state assignments of terminal taxa arising from effects of sampling variation.

The method uses multivariate count data for individual species to simultaneously infer the

number of ecological states, the proportional utilization of different resources by different states,

and the phylogenetic distribution of ecological states among living species and their ancestors.

The method is general and may be applied to any data expressible as a set of observational

counts from different categories.

Keywords: Ecological niche evolution; intraspecific variation; hidden Markov model;

macroevolution; comparative methods; Dirichlet-multinomial

Most species in the natural world make use of multiple, categorically-distinct types of

ecological resources. Many butterfly species use multiple host plants, for example (Ehrlich &

Raven 1964; Robinson 1999). Insectivorous warblers in temperate North America use multiple

distinct microhabitats and foraging behaviors (MacArthur 1958), as do honeyeaters in mesic and

arid Australia (Miller et al. 2017). The evolution of novel patterns of resource use can impact

phenotypic evolution (Martin & Wainwright 2011; Davis et al. 2016), diversification (Mitter et

al. 1988; Givnish et al. 2014), community assembly (Losos et al. 2003; Gillespie 2004), and

ecosystem function (Harmon et al. 2009; Bassar et al. 2010). Consequently, there has been

substantial interest in understanding how ecological traits related to resource use evolve and for

exploring their impacts on other evolutionary and ecological phenomena (Vrba 1987; Futuyma &

Moreno 1988; Forister et al. 2012; Price et al. 2012; Burin et al. 2016).

Making inferences about the evolutionary dynamics of resource use, however, first

requires summarizing the complex patterns of variation observed among taxa into traits that can

be modeled on phylogenetic trees. It is widely recognized that the real-world complexities of

resource use are not adequately described by a set of categorical variables (Hardy & Linder

2005; Hardy 2006). Nonetheless, it is also true that major differences in resource use can

sometimes be summed up in a small set of ecological states, a point made by Mitter et al. (1988)

in their study of phytophagy and insect diversification. For this reason, continuous-time Markov

chain (CTMC) models, which require classifying species into a set of character states, have

become commonplace in macroevolutionary studies of ecological trait evolution (Kelley &

Farrell 1998; Nosil 2002; Price et al. 2012; Hardy & Otto 2014; Cantalapiedra et al. 2014; Burin

et al. 2016). CTMC models describe a stochastic process for evolutionary transitions among a set

of character states and are used to infer ancestral states and evolutionary rates, and to perform

model-based hypothesis tests (O'Meara 2012).

The utility of continuous-time Markov chains for studying the evolutionary dynamics of

resource use is limited by the modeling assumption that taxa are monomorphic for ecological

states (Hardy & Linder 2005; Hardy 2006). As a practical solution, most empirical studies define

one or more generalized states to accommodate species that use multiple resource types and that

therefore cannot be characterized as specialists for a particular resource (Alencar et al. 2013;

Price et al. 2012; Burin et al. 2016; Gajdzik et al. 2019). Another solution, rather than classifying

each species as a specialist or a generalist, represents each resource category with a binary score

of present or absent (Janz et al. 2001; Colston et al. 2010; Hardy 2017). In this case, the

ecological state of a species is the set resources scored as present. Each of these approaches is

one solution to the modeling challenge posed by intraspecific variation in resource use, but both

solutions neglect variation in the relative importance of different resources for different taxa.

Consequently, species classified in single state can nonetheless exhibit substantial differences in

patterns of resource use, creating challenges for interpreting evolutionary transitions among

character states as well as for understanding links between character state evolution and

diversification.

Another limitation of continuous-time Markov chains for modeling resource use

evolution emerges from the fact that species are classified into ecological states without regard

for the quality and quantity of information available to perform the classification exercise. As an

example, species with few ecological observations might be classified as specialists for a

particular resource, when their apparent specialization is strictly a function of the small number

of ecological observations available for the taxon. More generally, by failing to use a statistical

model for making resource state assignments, we neglect a major source of uncertainty in our

data: the uneven and incomplete knowledge of resource use across different taxa. This

uncertainty, in turn, has substantial implications for how we project patterns of resource use onto

a set of resource states. By failing to account for uneven and finite sample sizes characteristic of

empirical data on resource use we cannot be certain if state assignments reflect true similarities

or differences in resource use or are merely the expected outcome of sampling variation.

Consider the simple four-species example in Figure 1. Panels (i) and (ii) illustrate the true

resource states and their phylogenetic distribution across a set of four species and their ancestors.

Here, an ancestral specialist evolved a generalist diet via a single transition (panel ii), such that

there are two extant species with the ancestral specialist diet (species X and Y) and two with the

derived generalist diet (species P and Q). In panels (iii) and (iv) the relative importance estimates

of three food resource categories in the diets of four species are used to classify each species into

one of two diet states. These relative importance estimates are based on uneven, and in some

cases quite small, sample sizes, consistent with many empirical datasets (Vitt & Vangilder 1983;

Shine 1994; Alencar et al. 2013). In panel (v) we imagine repeating the state assignment process

on independent datasets while holding the samples sizes fixed to those in panel (iii), which

reveals that both the initial state assignments and the number of states from (iv) are highly

sensitive to real-world levels of sampling variation. This has obvious implications for

downstream macroevolutionary analyses. There is a serious risk of conflating different (similar)

state assignments with different (similar) diets when the differences (similarities) are expected,

even in the absence of true differences (similarities), from sampling variation alone. In the

analyses that underlie Figure 1, we find that more than 70 percent of tip state classifications do

not match the true pattern of resource use.

Is this a problem in practice? This issue is difficult to assess because few studies provide

information about the sample sizes that underlie state assignments. In most cases, ecological

states are simply asserted as known. It is also important to emphasize that the specific problem in

Figure 1 is an outcome of a more general problem: standard CTMC models have a limited ability

to model complex ecological phenotypes because of the assumption that states in the model are

categorical variables. While it is true that CTMC models work with a countable state space, it is

not true that the states of the system must represent categorical variables. In a hidden Markov

model, the observed data are assumed to be the outcome of a CTMC where the states are not

directly observable. Instead, states are probability distributions from which observed data are

sampled. Although hidden state CTMC models already used in macroevolution (Marazzi et al.

2012; Beaulieu et al. 2013; Beaulieu & O'Meara 2016; Caetano et al. 2018) can be interpreted

like this they generally are not because in these cases the observed data are categorical variables

and the hidden states are indistinguishable from the observed categories. Instead, hidden states

are interpreted as unobserved factors that affect rates of change or rates of diversification.

Because of this, the potential flexibility of hidden state models for modeling complex phenotypes

remains poorly explored.

In this paper we use the formulation of hidden states as probability distributions to

develop a CTMC model for studying the evolutionary dynamics of ecological resource use on

phylogenetic trees. Our approach is explicitly designed to model resource traits that are intra-

specifically variable and to account for uncertainty in ecological state assignments of terminal

taxa arising from effects of sampling variation. We assume that each state is an unobserved

(latent) multinomial distribution and that observed data are sampled outcomes from these latent

distributions (see panels (i) through (iii) of Fig. 1). The number of states in the model and the

states themselves are not directly observed and are estimated from the data. Using simulations

and an empirical dataset of snake diets, we show how the method can use observational counts to

simultaneously infer the number of resource states, the proportional utilization of resources by

different states, and the phylogenetic distribution of ecological states among living species and

their ancestors. The method is general and applicable to any data expressible as a set of

observational counts from different resource categories.

MATERIALS & METHODS

Model description

We assume that the data for each species are represented by a vector of J category counts.

Each category is a resource (e.g. a diet or habitat component), and each count represents the

number of observations of a species utilizing a particular resource. Each node in a phylogeny is

to be placed into one of K distinct resource states. States are unobserved, even at the tips of a

phylogeny: the observed data consist of sampled outcomes from an underlying (latent)

multinomial distribution that represents the state for each species. All count data are drawn

independently from their respective states and counts from each state are also independent of one

another. We assume that the multinomial parameters for each state are drawn from a common

Dirichlet distribution with parameter β. This parameterization allows us to analytically

marginalize over the unknown multinomial parameters underlying each state so that the

likelihood of the observed data is the product of K independent Dirichlet-multinomial

distributions (Appendix). The parameter β acts as a vector of pseudo-counts. Higher values

require more data for the model to discriminate two samples as having originated from different

states. Letting X denote the resource state assignments for nodes in the phylogeny, the likelihood

of the set of count data D_k generated from state k is,

$$p(D_k|X,\beta) = \frac{\Gamma(J\beta)}{\Gamma(n_k + J\beta)} \frac{\prod_j \Gamma(n_k^j + \beta)}{\Gamma(\beta)^J} \tag{1}$$

168   where n_k is the total number of observations generated from state k and n_k^j is the subset

169   of those observations that represent utilization of resource category j. The full likelihood for the

170   count data is just,

$$p(D|X,\beta) = \prod_k p(D_k|X,\beta) \tag{2}$$

This model for count data is closely related to topic models of word composition in a

collection of text documents (Blei et al. 2003; Yin and Wang 2014) and to population genetic

models of allele frequency composition in a set of populations (e.g., program STRUCTURE:

Pritchard et al. 2000). The key difference here is that the state assigned to a taxon is the outcome

of evolution and is not independent of the states of other lineages. Conceptually this is similar to

phylogenetic threshold models, where the full likelihood combines a probability model for the

evolution of an unobserved variable and a probability model for sampling the observed data

conditioned on the set of unobserved variables (Felsenstein 2012; Revell 2014). We model

evolution as a Poisson process where the rate of change is the same between all states (i.e. there

is no evolutionary trend in the model) but varies among lineages. We introduce two mechanisms

for accommodating this rate variation.

The first mechanism takes advantage of the random local clocks model introduced by

Drummond and Suchard (2010). In this framework, there is an overall rate of evolution Λ and an

unknown number of lineages that deviate from this rate by a set of multiplicative constants.

Specifically, the root node is defined to have a relative rate of 1 while the relative rates of all

other nodes are equal to the relative rate of their ancestor multiplied by a branch-specific positive

rate multiplier. Complexity is controlled through the use of a prior that makes it unlikely for

many of these multipliers to differ from unity. Under the fully symmetric Poisson model with

random local clocks the probability of change across an ancestral-descendant branch is,

191
$$p\big(X_i \big| X_{pa(i)}, \Lambda, r_i\big) = (1 - e^{-K\Lambda r_i t_i})\frac{1}{K} + \delta_{X_i X_{pa(i)}} e^{-K\Lambda r_i t_i} \tag{3}$$

192 where X_i is the state of node i, $X_{pa(i)}$ is the state of i's ancestor, r_i is the branch-specific

193 normalized relative rate of evolution, t_i is the length of the branch in units of time, and $\delta_{X_i X_{pa(i)}}$

194 is the Kronecker delta. For simplicity, we may occasionally notate transition probabilities by p_{ii}

195 and p_{ij} to indicate the probability that a descendant's state is the same as, or different than, the

196 state of its ancestor. The likelihood of the node states is just,

197
$$p(X|\Lambda, r) = \prod_i p\big(X_i \big| X_{pa(i)}, \Lambda, r_i\big) \tag{4}$$

where the product is taken over all nodes and $p\big(X_i \big| X_{pa(i)}, \Lambda, r_i\big) \equiv \frac{1}{K}$ if node i is the root.

The normalized rates effectively expand and contract the temporal durations of the branches.

They are derived by scaling the relative rates in such a way that the total absolute time in which

evolution has had to occur is held constant even as the effective time is allowed to vary over

phylogeny (that is, $\sum_i t_i = \sum_i r_i t_i$).

The second mechanism for accommodating rate heterogeneity is essentially a saturated

version of the random local clocks model where each branch has a unique rate of evolution.

Following Huelsenbeck et al. (2008), this allows us to model branch-specific rates as nuisance

parameters drawn independently from a Gamma distribution with parameter vector $(\alpha, 1)$. This

model induces the same distribution of node states as a model where the number of expected

character state changes along a branch is the same for all branches (Appendix). This has

elsewhere been termed the ultra-common mechanism model (Steel 2011) to mark its contrast

with the no-common mechanism model (Tuffley and Steel 1997) from which it derives. In this

case the probability of change across an ancestral-descendant branch is,

212
$$p\left(X_i \middle| X_{pa(i)}, \alpha\right) = \left(1 - \frac{1}{(K/(K-1)+1)^\alpha}\right)\frac{1}{K} + \delta_{X_i X_{pa(i)}}\frac{1}{(K/(K-1)+1)^\alpha} \qquad (5)$$

Phylogenetic signal is controlled by the parameter α, which is equal to the expected number of

state changes that occur from ancestor to descendant. As $\alpha \rightarrow 0$, phylogenetic signal approaches

1 because descendants almost surely resemble their ancestors. As $\alpha \rightarrow \infty$, phylogenetic signal

approaches 0 because a descendant's state becomes independent of its ancestor's state and

resembles a random draw from a discrete uniform distribution. The likelihood of the node states

is just,

219
$$p(X|\alpha) = \frac{1}{K}p_{ii}{}^n p_{ij}{}^m \qquad (6)$$

where n is the number of nodes with the same state as their ancestor, m is the number of

nodes with a different state than their ancestor, and the factor $\frac{1}{K}$ accounts for the root state

probability.

Bayesian inference

We simulated the posterior distribution of node states and model parameters using the

Metropolis-Hastings algorithm (Hastings 1970). The different proposal mechanisms are

described below.

Updating node states.— A Gibbs update mechanism is used for proposing changes to

node states. The full conditional distribution for the state of a node can be written as,

230
$$p(X_i|X_{\neg i}, D, \beta, \theta) = \frac{p(D, X|\beta, \theta)}{p(D, X_{\neg i}|\beta, \theta)}$$

231
$$\propto \frac{p(D, X|\beta, \theta)}{p(D_{\neg i}, X_{\neg i}|\beta, \theta)}$$

232
$$= \frac{p(D|X, \beta)}{p(D_{\neg i}|X_{\neg i}, \beta)} \frac{p(X|\theta)}{p(X_{\neg i}|\theta)} \tag{7}$$

233   where the symbol $\neg i$ denotes the exclusion of node i. Here, depending on whether (4) or

234   (6) is used as the likelihood model for node states, θ is equal to (Λ, r) or α, respectively. Because

235   changing a node state only affects the branches incident to the affected node, all terms in the

236   ratio $\frac{p(X|\theta)}{p(X_{\neg i}|\theta)}$ not involving those branches cancel and it simplifies to,

237
$$\frac{p(X|\theta)}{p(X_{\neg i}|\theta)} = p\big(X_i|X_{pa(i)}, \theta\big) \prod_k p\big(X_{ch(i,k)}|X_i, \theta\big)$$

238   where $X_{ch(i,k)}$ is the state of the k-th immediate descendant of node i. Similarly, altering

239   the state of a node only affects the likelihood of count data associated with a single state so that

240   factors in the ratio $\frac{p(D|\beta)}{p(D_{\neg i}|\beta)}$ not involving, say, state k cancel, and it simplifies to,

241
$$\frac{p(D|X, \beta)}{p(D_{\neg i}|X_{\neg i}, \beta)} = \frac{p(D_k|X, \beta)}{p(D_{k\neg i}|X_{\neg i}, \beta)}$$

242
$$= \frac{\prod_j \Gamma\big(n^j_{k\neg i} + N^j_i + \beta\big)}{\prod_j \Gamma\big(n^j_{k\neg i} + \beta\big)} \frac{\Gamma(n_{k\neg i} + J\beta)}{\Gamma(n_{k\neg i} + N_i + J\beta)}$$

where N_i is the total number of observations for node i and N_i^j is the subset of those

observations that represent utilization of resource category j. Note that for nodes with no

associated count data (which includes all internal nodes), this factor is equal to 1. To perform the

Gibbs update in practice, we calculate the conditional likelihood for each diet state according to

equation (7) and choose a state with probability proportional to its conditional likelihood, using

the sum of all conditional likelihoods as a normalizing constant. The marginal posterior

probability that a node is in a given diet state is simply the fraction of posterior samples where it

appears in that state.

Once a state is sampled for a node any count data associated with that node are added to

the set of count data generated from the sampled state. Because the Dirichlet distribution is

conjugate to the multinomial distribution, the posterior distribution of the multinomial

distribution underlying each state is also Dirichlet distributed with parameter $\left(n_k^1 + \beta, \dots, n_k^J + \right.$

$\left. \beta \right)$. During the course of updating node states we keep track of the average expected proportional

utilization of each resource by each state. The expected proportional utilization of resources is

simply the mean of the posterior distribution which is $\left(\frac{n_k^1 + \beta}{\sum_j n_k^j + J\beta}, \dots, \frac{n_k^J + \beta}{\sum_j n_k^j + J\beta} \right)$.

Updating β.—The symmetric hyperparameter β controls the shape of the Dirichlet prior

distribution on the latent multinomial distributions underlying each resource state. When $\beta = 1$

the distribution is uniform over the J-dimensional simplex of resources. When $\beta < 1$ the

distribution concentrates toward the corners of the simplex, and when $\beta > 1$ the distribution

concentrates toward the center. Because empirical datasets are typically sparse with many zeros,

we assume that β is uniformly distributed on the interval $(0, 1)$ and update its value using a

sliding window proposal mechanism. The prior and proposal ratios are 1.

Updating α.—When equation (6) is used to compute the likelihood the hyperparameter α

controls phylogenetic signal. Although it can take on any positive value the likelihood surface

plateaus relatively quickly as its magnitude increases and phylogenetic signal decays. By solving

the logarithm of (6) for the maximum likelihood estimate of α we find that

269
$$\hat{\alpha} = -\frac{\log\left(\frac{fK-1}{K-1}\right)}{\log(K/(K-1)+1)}$$

where f is the fraction of nodes that have the same state as their ancestor. Values of $f \leq \frac{1}{K}$

are consistent with infinite values of α. We therefore bound α above by the value

$-\log\left(\frac{\frac{\lfloor N/K\rfloor+1}{N}K-1}{K-1}\right)\Big/\log(K/(K-1)+1)$, where N is the number of nodes (not including the

root) in the phylogeny. We assume that α is uniformly distributed between zero and this upper

value and update its value using a sliding window proposal mechanism. The prior and proposal

ratios are 1.

Updating Λ.—When equation (4) is used to compute the likelihood the parameter Λ

controls phylogenetic signal. As for α, we want a reasonable upper bound for Λ because the

likelihood plateaus relatively quickly as the rate increases. By replacing branch lengths in

equation (4) with the average branch length and solving for the maximum likelihood estimate of

Λ under the assumption of no rate variation we find that

281
$$\hat{\Lambda} = -\frac{\log\left(\frac{fK-1}{K-1}\right)}{K\bar{t}}$$

where $\bar{t}$ is the average branch length. As before, values of $f \leq \frac{1}{K}$ are consistent with

infinite values of Λ. We therefore bound Λ above by the value $-\log\left(\frac{\frac{\lfloor N/K\rfloor+1}{N}K-1}{K-1}\right)\Big/K\bar{t}$. We

assume that Λ is uniformly distributed between zero and this upper value and update its value

using a sliding window proposal mechanism. The prior and proposal ratios are 1. We use the

same prior distributions and proposal mechanisms detailed by Drummond and Suchard (2010)

for updating the number of local random clocks and the rate multipliers associated with local

clocks.

Implementation

Functions for fitting the model to data are provided as an R package available from

github.com/blueraleigh/phyr. The package includes two R functions that call compiled C

programs implementing the random local clocks and ultra-common mechanism models.

Simulation study

To illustrate application of the method we designed a simulation study using an empirical

dataset on pseudoboine snake diets (Alencar et al. 2013). Our rationale for basing simulations on

an empirical dataset is to ensure that properties of the data used to evaluate performance of the

method are consistent with real studies, especially the distribution of observations per taxon and

the distribution of resource specialization. Pseudoboine snakes are common members of the

squamate communities found in lowland rainforests of South America. Predominantly terrestrial

or semi-arboreal, these snakes mainly eat small mammals, lizards, and other snakes. The dataset

includes 606 observations of prey items from 8 prey categories for 32 species of pseudoboine

snakes. Per species sample sizes range from 1 to 56 observations (or 0.125-fold to seven-fold

coverage, where coverage is the number of observations divided by the number of resource

categories). We reanalyzed these data using a 33-species pseudoboine phylogeny extracted from

the posterior distribution of trees in Tonini et al. (2016). The dataset is illustrated in Figure 2.

The original publication coded each species with at least 8 diet observations into a set of 5

specialist diet states and 1 generalist diet state. Species were considered specialists if the

resource represented at least 70 percent of recorded prey items (as in our Figure 1). When we

applied the resampling procedure illustrated in Figure 1 to this empirical dataset under the

assumption that the original state assignments represented the "truth", we found that in

approximately 20 percent of resampled datasets at least one original state (not always the same

state) was not present and that in about 84 percent of cases at least one species was coded

incorrectly (although overall coding accuracy was high, ranging from 0.77 to 1). Thus, this

dataset illustrates some of the concerns raised in our introduction but is also well-sampled

enough and shows enough variation to facilitate the estimation of separate multinomial

distributions.

Simulated datasets were generated from $K = 2, 3, 4,$ and 5 diet states using the empirical

sample size distribution with the original 8 food resource categories. For each K we first

performed Bayesian inference under the ultra-common mechanism model to estimate the

unobserved multinomial distributions. The estimated multinomial distributions were

subsequently used to simulate diet observations. For each K we simulated 20 datasets at each of

7 different levels of phylogenetic signal $(0.1, 0.3, 0.5, 0.6, 0.7, 0.8,$ and $0.9)$ using the transition

probabilities in both equations (3) and (5), resulting in 560 datasets for each model and 1,120

datasets altogether. We defined phylogenetic signal as $p_{ii} - p_{ji}$, which ranges from 0 to 1 and

quantifies how much information a descendant's state provides about the state of its ancestor

(Royer-Carenzi et al. 2013). Using equation (5) for transition probabilities results in phylogenetic

signal equal to $\left(\frac{1}{K/(K-1)+1}\right)^{\alpha}$. We used this result to calculate the value of α for each simulation.

When equation (3) is used for transition probabilities each branch has a unique phylogenetic

signal. Because phylogenetic signal is a convex function of branch length, the average

phylogenetic signal of all branches is greater than or equal to the phylogenetic signal of the

average branch, which is $e^{-K\Lambda\bar{t}}$. We used the phylogenetic signal of the average branch to

calculate the value of Λ for each simulation, which we applied to all branches (i.e., datasets did

not include random local clock variation). Interestingly, for a given branch length (measured as

expected number of state changes) phylogenetic signal with equation (5) is always greater than

phylogenetic signal with equation (3), suggesting that estimating the rate of evolution trades off

with estimating ancestral node states (Gascuel and Steel 2018). For each simulated dataset we

ran a set of Markov chains with 1, 2, …, up to K+3 diet states. Each chain was run for 160,000

iterations after a burnin of 30,000 iterations, sampling every 128 iterations to yield 1,250

posterior samples.

Determining the number of resource states

Because the model does not include a process for generating the number of states, we

must perform analyses across multiple values of K and apply an *a posteriori* inference procedure

to choose between them. A similar problem is encountered when trying to infer the number of

demes from multi-locus genotype data with the program STRUCTURE (Pritchard et al. 2000).

Our approach is to choose the largest value of K for which all states are unambiguously assigned

to at least one terminal taxon. Specifically, by examining the terminal nodes of the phylogeny we

can determine the maximum marginal posterior probability assigned to each state across all

terminal nodes. Call the smallest of these q_K. A low value of q_K implies that at least one state is

not assigned to any individuals (terminal taxa) with high probability. As K varies from low to

high there will come a point when additional states become redundant to previous states. When

that occurs, q_K will drop well below 1. In other words, a point will be reached when at least one

state is never unambiguously assigned to at least one terminal node. To choose the best value for

K we first identify the K with the steepest drop in q_K with respect to q_{K-1}. $K^* = K - 1$ then

becomes a candidate best choice. If $q_{K^*} = 1$ (or nearly so) we treat K^* as the best choice.

Otherwise, we keep setting $K^* = K^* - 1$ until $q_{K^*} = 1$. This procedure, which we call the q_K

rule, is illustrated in Figure 3 on the empirical dataset.

Assessing model adequacy

In practice, the number of states identified by the q_K rule will depend on the underlying

data and may change as data are updated. This is because the model may be unable to distinguish

truly different samples as having arisen from separate distributions if sample sizes are small.

Therefore, empirical applications of the method require a way of assessing how well the inferred

multinomial distributions explain the empirical resource observations. Our approach is to

compute a per-species adequacy score that measures the similarity of each taxon, with respect to

sampled observations, to other taxa assigned to the same state. The procedure we describe

effectively measures the compositional heterogeneity of the model-inferred states with respect to

sampled diet observations. If the model is fully adequate and accurately describes patterns of

resource use, then all species assigned to a given state will have similar sampled diets (e.g., the

observed data). However, some species may be assigned to states even where they have

dissimilar sampled observations from other species in the same state, reflecting overdispersed

diet distributions for the state. Such overdispersion might arise if species are assigned to the

"wrong" state. These incorrect state assignments might be preferred under the model if a species

does not have enough observations to provide information about the existence of a distinct state,

and so the species is conservatively assigned to the immediately ancestral diet state. To compute

the adequacy score, we first draw a state for each terminal node from its array of marginal

posterior probabilities computed using equation (7) (which utilizes count data and evolutionary

history). Then we visit each terminal node in turn and perform the following exercise. Given the

configuration of states for all terminal nodes, we identify the set of count data generated by the

state of the current node and compute the likelihood of these data using equation (1) (which only

utilizes count data). Next, we compute the likelihood of these same data, using equation (1)

together with equation (2), but assuming that they were generated from two states by placing the

current node in its own unique state. Finally, we take the negative log likelihood ratio of these

two values. We repeat this procedure for a thousand independent configurations and record the

average negative log likelihood ratio (adequacy score) for each terminal node. Large negative

adequacy scores highlight terminal nodes that were assigned to a state whose other members

have a strongly dissimilar pattern of resource utilization. This procedure is illustrated in Figure 4

on the empirical dataset (see also Figure S4).

RESULTS

Overall, the q_K rule correctly identified the number of resource states in 492 of 560

simulations from the ultra-common mechanism model (Fig. 5). In the 68 cases where the method

incorrectly identified the number of states, it underestimated the number of states by one (61

instances), two (4 instances), and three states (2 instance) and overestimated the number of states

by one state in one instance. When the q_K rule was used with the random local clocks model it

correctly identified the number of states in 475 of 560 simulations (Fig. S1). In the 85 cases

where the method incorrectly identified the number of states, it underestimated the number of

states by one (77 instances) and two states (8 instances). Failure to correctly identify the number

of states commonly occurs when the number of observations generated by a state is small

relative to the number of observations from other states. This happens when the terminal nodes

representing a state have poorly sampled diets causing the state to be subsumed into the state of

near relatives.

Estimation of latent multinomial probabilities underlying resource states was highly

accurate (Figs. 5, S1, S5). In simulations from both the ultra-common mechanism model and the

random local clocks model, the correlation between true and estimated probabilities ranged from

0.92 to 0.999 with a mean of 0.99. Such high values occur because even when the number of

states is underestimated, the number of sampled observations from the missing states are few

enough in number that they do not appreciably alter the estimated proportions of the other states

(Fig S6).

Across all levels of phylogenetic signal, and with simulations from both the ultra-

common mechanism model and the random local clocks model, the method consistently

classified greater than ninety-percent of terminal nodes in the correct state. For internal nodes,

reconstruction accuracy was comparable to reconstruction accuracy for terminal nodes at the

highest level of phylogenetic signal but, in accordance with expectation, decayed toward the

expected accuracy of a random guess as phylogenetic signal declined toward zero (Figs. 6, S2).

This behavior was mirrored in the posterior distributions of α and Λ (Figs. 7, S3). When

phylogenetic signal was low, posterior distributions of rate estimates were diffuse and resembled

their uniform prior distributions except that they were shifted away from the lower bound. As

phylogenetic signal increased, posterior distributions of rate estimates concentrated around the

true values used to generate simulated datasets.

DISCUSSION

We developed a comparative method for macroevolutionary analysis of multivariate

count data. The method is general and may be applied to any data expressible as a set of

observational counts from different categories. Such datatypes arise frequently in community

ecology and behavior. Potential applications include the comparative analysis of diet, foraging

behavior, activity patterns, and habitat preferences. The method is similar to standard

continuous-time Markov chain models of phenotypic evolution but differs in several important

respects. First, the number of states in the model and the states themselves are unobserved and

must be estimated from empirical data on resource use. Second, each state is an unobserved

multinomial distribution rather than a categorical variable. This latter property enables

researchers to model ecological traits that show intraspecific variation and to account for

uncertainty in the state assignments of terminal taxa that arises from the effects of sampling

variation.

Simulations revealed that the new method is generally able to determine the correct

number of states and that it provides accurate estimates of the underlying (unobserved)

multinomial distributions, both for terminal taxa as well as internal nodes. We designed

simulations around empirical patterns of resource use in a dataset on snake diets (Alencar et al.

2013). Therefore, caution is warranted in generalizing the good performance observed in the

current study to other datasets. In particular, performance of the model will depend on the

idiosyncrasies of individual datasets, including the distribution of sample sizes and the

distribution of overlaps in resource use among species. We expect that states represented by few

observations will be difficult to infer, especially if those states show appreciable overlap with

other states.

Our empirical analysis identified at least two feeding modalities among the set of species

Alencar et al. (2013) recognize as "generalists": species that feed predominantly on snakes but

that regularly eat lizards and mammals, and species that feed predominantly on mammals and

lizards. Ancestral state estimates strongly suggest that each of these feeding modalities arose

from a more specialized diet comprised almost entirely of lizards. This is in contrast to the

results of Alencar et al. (2013), which imply that nearly all origins of specialized feeding

modalities occurred from a generalist ancestor, although direct comparison of results is made

difficult by the use of different phylogenies in the two studies.

As currently implemented, the approach described here does not directly model gains and

losses or substitutions of different resources. Indeed, no resource is ever truly absent from the

reconstructed states (although its proportional representation may approach zero as β becomes

small). This contrasts with biogeographic-type models that explicitly model resource use

expansions, contractions, and substitutions (e.g. see Hardy (2017) for application of Ree and

Smith's (2008) dispersal-extinction-cladogenesis model to binary encoded diet data). Although

these types of changes are implicit in the sequence of reconstructed states derived from the

model, future studies might want to explore how to combine more complex evolutionary models

with the current model for count data. Nonetheless, the advantage of a simple evolutionary

model is that it has broad scope. It would be possible, for example, to apply our method to

continuous characters by keeping the same evolutionary model but changing the model for

observations from a Dirichlet-multinomial to a multivariate-normal distribution, which could

then be applied to other data types used for quantifying resource use such as stable isotope ratios

of carbon and nitrogen.

One challenge for comparative methods is their limited ability to model ecological

phenotypes that cannot be neatly summarized by a single value (Hardy & Linder 2005; Hardy

2006). Recent years have seen progress in this direction for continuous traits, including models

that accommodate intraspecific variation, function-valued traits, and other non-gaussian data

(Ives et al. 2007; Felsenstein 2008; Evans et al. 2009; Jones & Moriarty 2013; Goolsby 2015;

Quintero et al. 2015). The general approach developed here, where each state is a multinomial

distribution rather than a categorical variable, extends this progress to traits like diet and habitat

that are typically treated as categorical variables. By placing an emphasis on individual natural

history observations, the method draws attention to the central role such observations play in

evolutionary biology (Greene 1986) and to the many remaining opportunities for developing

comprehensive ecological databases that advance our understanding of biodiversity (Hortal et al.

2015).

SUMMARY

We described a novel methodological framework for studying the evolutionary dynamics

of complex ecological traits on phylogenetic trees. Previous approaches to this problem have

assumed that ecological states are categorical variables and that species are monomorphic for

particular states. We relaxed this assumption through the use of a hidden Markov model that

treats ecological states as unobserved probability distributions from which observed data are

sampled. Although our method is designed for the analysis of multivariate count data, we suggest

that the approach of treating states as hidden probability distributions has wide applicability and

will greatly facilitate the comparative analysis of novel sources of ecological data.

ACKNOWLEDGEMENTS

This research was supported in part through computational resources and services provided by

Advanced Research Computing at the University of Michigan, Ann Arbor. This material is based

upon work supported by the National Science Foundation Graduate Research Fellowship and by

the University of Michigan Department of Ecology and Evolutionary Biology Block Grant.

LITERATURE CITED

Alencar L.R.V., Gaiarsa M.P., Martins M. 2013. The evolution of diet and microhabitat use in
500   pseudoboine snakes. South American Journal of Herpetology. 8:60–66.

501   Bassar R.D., Marshall M.C., López-Sepulcre A., Zandona E., Auer S.K., Travis J., Pringle C.M.,
502   Flecker A.S., Thomas S.A., Fraser D.F., Reznick D.N. 2010. Local adaptation in
503   Trinidadian guppies alters ecosystem processes. PNAS. 107:3616–3621.

504   Beaulieu J.M., O'Meara B.C. 2016. Detecting hidden diversification shifts in models of trait-
505   dependent speciation and extinction. Systematic Biology. 65:583–601.

506   Beaulieu J.M., O'Meara B.C., Donoghue M.J. 2013. Identifying hidden rate changes in the
507   evolution of a binary morphological character: the evolution of plant habit in Campanulid
508   angiosperms. Systematic Biology. 62:725–737.

509   Blei D.M., Ng A.Y., Jordan M.I. 2003. Latent Dirichlet Allocation. Journal of Machine Learning
510   Research. 3:993–1022.

511   Burin G., Kissling W.D., Guimarães P.R., Sekercioglu C.H., Quental T.B. Omnivory in birds is a
512   macroevolutionary sink. Nature Communications. 7:11250.

513   Caetano D.S., Beaulieu J.M., O'Meara B.C. 2018. Hidden state models improve state-dependent
514   diversification approaches, including biogeographical models. Evolution. 72:2308–2324.

515   Cantalapiedra J.L., FitzJohn R.G., Kuhn T.S., Hernández Fernández M., DeMiguel D., Azanza
516   B., Morales J., Mooers A.Ø. 2014. Dietary innovations spurred the diversification of
517   ruminants during the Caenozoic. Proceedings of the Royal Society B: Biological
518   Sciences. 281:20132746.

Colston T.J., Costa G.C., Vitt L.J. 2010. Snake diets and the deep history hypothesis. Biological
Journal of the Linnean Society. 101:476–486.

Davis A.M., Unmack P.J., Vari R.P., Betancur-R R. 2016. Herbivory promotes dental
disparification and macroevolutionary dynamics in grunters (Teleostei: Terapontidae), a
freshwater adaptive radiation. The American Naturalist. 187:320–333.

Drummond A.J., Suchard M.A. 2010. Bayesian random local clocks, or one rate to rule them all.
BMC Biology. 8:114.

Ehrlich P.R., Raven P.H. 1964. Butterflies and plants: a study in coevolution. Evolution. 18:586–
608.

Evans M.E.K., Smith S.A., Flynn R.S., Donoghue M.J. 2009. Climate, niche evolution, and
diversification of the "bird-cage" evening primroses (*Oenothera*, Sections *Anogra* and
Kleinia). The American Naturalist. 173:225–240.

Felsenstein J. 2008. Comparative methods with sampling error and within-species variation:
contrasts revisited and revised. The American Naturalist. 171:713–725.

Felsenstein J. 2012. A comparative method for both discrete and continuous characters using the
threshold model. The American Naturalist. 179:145–156.

Forister M.L., Dyer L.A., Singer M.S., Stireman III J.O., Lill J.T. 2012. Revisiting the evolution
of ecological specialization, with emphasis on insect-plant interactions. Ecology. 93:981–
991.

Futuyma D.J., Moreno G. 1988. The evolution of ecological specialization. Annual Review of
Ecology and Systematics. 19:207–233.

Gajdzik L., Aguilar-Medrano R., Frédérich B. 2019. Diversification and functional evolution of
reef fish feeding guilds. Ecology Letters. 22:572–582.

Gascuel O., Steel M. 2018. A Darwinian uncertainty principle. bioRxiv.
https://doi.org/10.1101/506535

Gillespie R. 2004. Community assembly through adaptive radiation in Hawaiian spiders.
Science. 303:356–359.

Givnish T.J., Barfuss M.H.J., Van Ee B., Riina R., Schulte K., Horres R., Gonsiska P.A., Jabaily
R.S., Crayn D.M., Smith J.A., Winter K., Brown G.K., Evans T.M., Holst B.K., Luther
H., Till W., Zizka G., Berry P.E., Sytsma K.J. 2014. Adaptive radiation, correlated and
contingent evolution, and net diversification in Bromeliaceae. Molecular Phylogenetics
and Evolution. 71:55–78.

Goolsby E.W. 2015. Phylogenetic Comparative Methods for Evaluating the Evolutionary
History of Function-Valued Traits. Systematic Biology. 64:568–578.

Greene H.W. 1986. Natural history and evolutionary biology. In: Feder M.E., Lauder G.V.,
editors. Predator–prey relationships: perspectives and approaches from the study of lower
vertebrates. University of Chicago Press. p. 198.

Hardy C.R. 2006. Reconstructing ancestral ecologies: challenges and possible solutions.
Diversity and Distributions. 12:7–19.

Hardy C.R., Linder H.P. 2005. Intraspecific variability and timing in ancestral ecology
reconstruction: a test case from the cape flora. Systematic Biology. 54:299–316.

Hardy N.B. 2017. Do plant-eating insect lineages pass through phases of host-use generalism
during speciation and host switching? Phylogenetic evidence. Evolution. 71:2100–2109.

Hardy N.B., Otto S.P. 2014. Specialization and generalization in the diversification of
phytophagous insects: tests of the musical chairs and oscillation hypotheses. Proceedings
of the Royal Society B: Biological Sciences. 281:20132960.

Harmon, L. J., Matthews B., Des Roches S., Chase J.M., Shurin J.B., Schluter D. 2009.
Evolutionary diversification in stickleback affects ecosystem functioning. Nature.
458:1167–1170.

Hastings W.K. 1970. Monte Carlo Sampling Methods Using Markov Chains and Their
Applications. Biometrika. 57:97–109.

Hortal J., de Bello F., Diniz-Filho J.A.F., Lewinsohn T.M., Lobo J.M., Ladle R.J. 2015. Seven
shortfalls that beset large-scale knowledge of biodiversity. Annual Review of Ecology,
Evolution, and Systematics. 46:523–549.

Huelsenbeck J.P., Ané C., Larget B., Ronquist F. 2008. A Bayesian perspective on a non-
parsimonious parsimony model. Systematic Biology. 57:406–419.

Ives A.R., Midford P.E., Garland Jr T. 2007. Within-species variation and measurement error in
phylogenetic comparative methods. Systematic Biology. 56:252–270.

Janz N., Nyblom K., Nylin S. 2001. Evolutionary dynamics of host-plant specialization: a case
study of the tribe Nymphalini. Evolution. 55:783–796.

Jones N.S., Moriarty J. 2013. Evolutionary inference or function-valued traits: Gaussian process
regression on phylogenies. Journal of the Royal Society Interface. 10:20120616.

Kelley S.T., Farrell B.D. 1998. Is specialization a dead end? The phylogeny of host use in
Dendroctonus bark beetles (Scolytidae). Evolution. 52:1731–1743.

Losos J.B., Leal M., Glor R.E., de Queiroz K., Hertz P.E., Rodríguez Schettino L., Chamizo Lara
A., Jackman T.R., Larson A. 2003. Niche lability in the evolution of a Caribbean lizard
community. Nature. 424:542–545.

MacArthur R.H. 1958. Population ecology of some warblers of northeastern coniferous forests.
Ecology. 39:599–619.

Marazzi B., Ané C., Simon M.F., Delgado-Salinas A., Luckow M., Sanderson M.J. 2012.
Locating evolutionary precursors on a phylogenetic tree. Evolution. 66:3918–3930.

Martin C.H., Wainwright P.C. 2011. Trophic novelty is linked to exceptional rates of
morphological diversification in two adaptive radiations of *Cyprinodon* pupfish.
Evolution. 65:2197–2212.

Miller E.T., Wagner S.K., Harmon L.J., Ricklefs R.E. 2017. Radiating despite a lack of
character: ecological divergence among closely related, morphologically similar
honeyeaters (Aves: Meliphagidae) co-occurring in arid Australian environments. The
American Naturalist. 189:E14–E30.

Mitter C., Farrell B., Wiegmann B. 1988. The phylogenetic study of adaptive zones: has
phytophagy promoted insect diversification? The American Naturalist. 132:107–128.

Nosil P. 2002. Transition rates between specialization and generalization in phytophagous
insects. Evolution. 56:1701–1706.

O'Meara B.C. 2012. Evolutionary inferences from phylogenies: a review of methods. Annual
Review of Ecology and Systematics. 43:267–285.

Price S.A., Hopkins S.S.B., Smith K. K., Roth V.L. 2012. Tempo of trophic evolution and its
impact on mammalian diversification. PNAS. 109:7008–7012.

Pritchard J.K., Stephens M., Donnelly P. 2000. Inference of Population Structure Using
Multilocus Genotype Data. Genetics. 155:945–959.

Quintero I., Keil P.K., Jetz W., Crawford F.W. 2015. Historical biogeography using species
geographical ranges. Systematic Biology. 64:1059–1073.

Ree R.H., Smith S.A. 2008. Maximum likelihood inference of geographic range evolution by
dispersal, local extinction, and cladogenesis. Systematic Biology. 57:4–14.

Revell L.J. 2014. Ancestral character estimation under the threshold model from quantitative
genetics. Evolution. 68:743–759.

Robinson G.S. 1999. HOSTS - a database of the hostplants of the world's Lepidoptera. Nota
Lepidopterologica. 22:35–47.

Royer-Carenzi M., Pontarotti P., Didier G. 2013. Choosing the best ancestral character state
reconstruction method. Mathematical Biosciences. 242:95–109.

Shine R. 1994. Allometric patterns in the ecology of Australian snakes. Copeia. 1994:851–867.

Steel M. 2011. Can we avoid "SIN" in the house of "no common mechanism"? Systematic
Biology. 60:96–109.

Tonini J., Beard K.H., Barbosa Ferreira R., Jetz W., Pyron R.A. 2016. Fully-sampled
phylogenies of squamates reveal evolutionary patterns in threat status. Biological
Conservation. 204:23–31.

Tuffley C., Steel M. 1997. Links between maximum likelihood and maximum parsimony under a
simple model of site substitution. Bulletin of Mathematical Biology. 59:581–607.

Vitt L.J., Vangilder L.D. 1983. Ecology of a snake community in northeastern Brazil. Amphibia-
Reptilia. 4:273–296.

Vrba E.S. 1987. Ecology in relation to speciation rates: some case histories of Miocene-Recent
mammal clades. Evolutionary Ecology. 1:283–300.

Yin J., Wang J. 2014. A dirichlet multinomial mixture model-based approach for short text
clustering. Proceedings of the 20th ACM SIGKDD international conference on
Knowledge discovery and data mining - KDD '14.:233–242.

Figure 1. Distribution and representation of multivariate ecological phenotypes (i, ii, iii), data as sampled by researchers (iii), and sampled states as typically represented by univariate categorical traits (iv, v). Loss and distortion of information associated with complex phenotypes motivates the development of the Dirichlet-multinomial model described in this article. (i) True resource states are unobserved multinomial distributions that determine the proportional utilization of three dietary resource categories by four species. (ii) The resource state of a species is the outcome of evolution via a hidden Markov process where the hidden states are the unobserved multinomial distributions. Here, the multinomial distributions from (i) are represented as rose plots: the direction of a spoke identifies the resource category and the length of a spoke is equal to the proportional utilization of that category. The phylogeny depicts the true evolutionary history of change. (iii) Observed data are sampled outcomes from these latent multinomial distributions. (iv) Sampled outcomes are projected onto a set of resource states. Here, a species is considered a "specialist" on a particular category if the sampled proportion of the category

exceeds 0.7. Otherwise, it is considered a "generalist". In this case, the dataset and cutoff value

align to match each species with its correct modal resource category. (v) The same state

assignment process is repeated with different datasets while keeping the sample sizes for each

species identical to (iii). State assignments are sorted along the x-axis according to their

frequency of occurrence in 1,000 independent datasets. Datasets were generated by sampling

from the two multinomial distributions in (i). Note that the procedure correctly matches all

species with their modal food resource in a minority of cases and results in a variable number of

states across datasets. The implication for macroevolutionary studies using this state assignment

procedure is that we cannot be certain whether state assignments are reflective of true patterns of

resource use or are merely the expected outcome of sampling variation.

Figure 2. Summary attributes of the snake dietary dataset used to parameterize the simulation

study, including phylogeny of pseudoboine snakes (left), relative prey frequencies (middle), and

total numbers of food observations per snake species (right). Dark colors in the prey frequency

matrix indicate higher sampled proportions of a particular prey item in a given diet. The

phylogeny and sampled diet observations were used to infer the "true" (unobserved) diet states

and their evolutionary history on the phylogeny (colored circles). Each color in a pie chart is a

specific model-inferred state and the size of a slice represents the marginal posterior probability

that a node belongs to that state. Note that the terminal node marked with an asterisk is missing

data. It is treated like an internal node and information about its probable diet state is drawn only

from what the model has learned about the states of its neighbors and the likelihood of

evolutionary change. Here, the Dirichlet-multinomial model inferred 5 states, corresponding to 3

specialist (> 70% specificity for a single prey group) and 2 generalist diets. Note that the diet

states are not observed directly, even at the tips of the tree; rather, all observed data are assumed

to be sampled from a set of unknown multinomial distributions.

Figure 3. Illustration of the *a posteriori* criterion for determining the number of states in the

model. Panel (a) shows the average log likelihood of the empirical data as a function of the

number of diet states. Panel (b) depicts how q_K, the smallest maximum marginal posterior

probability with which a state is assigned to terminal taxa, changes as a function of the number

of states. Inspection of the marginal posterior probabilities reveals that the sixth state is never

unambiguously assigned to a terminal node (panels b and c). For this reason, a model with five

resource states is considered optimal. The proportional utilization of different food resources by

those five states is illustrated by the rose plot in panel (d).

Figure 4. Goodness of fit of inferred multinomial distributions to sampled observations in the

empirical snake diet dataset when the inference model assumes $K = 2, 3, 4$, and 5 states.

Adequacy scores are computed on a per-species basis; negative scores imply that sampled

observations for a species are different from other taxa assigned to the same state. Such negative

scores might arise when a species has sampled diet that is distinct from that of close relatives but

where insufficient data (few observations) are available to inform the model as to the existence

of a new and distinct dietary state. Each line segment is a species in the empirical dataset. Each

pie chart depicts the marginal posterior probabilities that a species is in a given diet state. Thus,

fitting a model with only 2 states results in a third of the species sharing a diet state with

ecologically dissimilar species. In a model with 5 states nearly all species in the empirical dataset

share a similar pattern of resource utilization with the other species assigned to the same diet

state. The two species with negative scores have a relatively high sampled percentage of

amphibian prey items or a relatively broad diet compared to other species in the dataset but only

modest sample sizes (compare to Fig. 2).

Figure 5. Performance of the method at identifying the true number of states (top) and the latent

multinomial distributions underlying states (bottom). Each column summarizes the results of a

set of 140 simulated datasets where the number of resource states varied from 2 (leftmost

column) to 5 (rightmost column). *Top:* each bar chart shows the frequency of the number of

resource states estimated using the q_K rule discussed in the main text; black bars are the number

of states in the generating model. The method correctly identifies the number of generating states

in most cases. *Bottom*: each set of solid colored spokes represents the latent multinomial

distribution underlying a resource state in the generating model. The direction of a spoke

identifies its corresponding resource and its length equals the proportional utilization of that

resource. Directions are slightly offset between diets states so that spokes with contacting edges

represent the same resource. Spokes with colored outlines but unfilled centers are the model

estimated multinomial distributions for the corresponding state. The method correctly identifies

the major and minor resources of each state in most cases.

Figure 6. Accuracy of model-inferred resource state assignments for terminal and internal nodes across varying levels of phylogenetic signal. Each point is the reconstruction accuracy for a single simulation. Reconstruction accuracy is the probability that a randomly selected node was classified in the correct resource state. The horizontal dashed lines correspond to the expected reconstruction accuracy of a random guess. The reconstruction accuracy of terminal nodes is relatively constant across levels of phylogenetic signal. The reconstruction accuracy of internal nodes is comparable to the accuracy of terminal nodes when phylogenetic signal is high and, as expected, decays toward the random expectation as phylogenetic signal declines. This decline is expected because, with low signal, the tip data provide little information about states at internal nodes. Light weight dashed lines connect the median accuracy at each level of phylogenetic signal.

Figure 7. Posterior distributions of the expected number of character state changes per branch

for simulated datasets with $K = 2, 3, 4$, and 5 states. Each violin plot summarizes the posterior

distributions of 20 simulations at each of 7 different levels of phylogenetic signal. The black

hatch mark within each violin is the expected number of changes per branch that was used to

generate simulated datasets. The width of a violin measures how frequently estimated values

appear in the posterior distributions. Phylogenetic signal increases as the expected number of

changes decreases. At low phylogenetic signals posterior distributions largely recapitulate their

uniform prior distributions but as phylogenetic signal increases they become concentrated around

the true values.

Figure S1. As for Figure 5 in the main text except that simulations were made using transition

probabilities from equation (3) rather than from equation (5).

Figure S2. As for Figure 6 in the main text except that simulations were made using transition

probabilities from equation (3) rather than from equation (5).

Figure S3. As for Figure 7 in the main text except that simulations were made using transition

probabilities from equation (3) rather than from equation (5) and phylogenetic signal refers to the

phylogenetic signal of the average branch length.

Per–species adequacy score example

Figure S4. The main text defines a per-species adequacy score to assess how well the estimated

multinomial distributions explain sampled observations for terminal taxa. The toy example in

this figure depicts how such a score is calculated for species 2. In the main text, we use the

negative logarithm of the likelihood ratio calculation depicted in the figure. Thus, positive values

for the log likelihood ratio imply that the estimated multinomial is a good fit to sampled

observations; negative values imply the opposite.

Figure S5. Correlation between true and estimated multinomial probabilities for datasets

simulated under the ultra-common mechanism model and random local clocks model.

True

Estimated

Figure S6. Two examples illustrating how high accuracy of estimated multinomial proportions is

maintained even when the number of states is underestimated. In the left plot, the blue state only

generated 1 sampled observation and the brown state only generated 5 sampled observations.

In the right plot, the brown state only generated 8 observations; the blue state, 12; and the green

state, 2. In both cases, these observations are few enough in number that they do not substantially

alter the estimated multinomial proportions when they are incorporated into the count data

generated from the other states.

899 APPENDIX

900 *Derivation of $p(D|X,\beta)$*

901   We assume that conditional on the resource state the data for each species are sampled

902 independently from the latent multinomial distribution underlying the state. That is,

$$903 \quad p(D_k|X,\theta) \propto \prod_i p(d_i|X_i = k,\theta)$$

$$904 \quad = \prod_i \prod_j \theta_j^{N_i^j}$$

905   where we have omitted terms involving multinomial coefficients as well as the

906 dependence of θ on k for ease of notation. We also assume that for each k the multinomial

907 parameter $\theta \sim \text{Dirichlet}(\beta_1, \dots, \beta_J)$ so that,

$$908 \quad p(D_k|X,\beta) \propto \int_\theta p(D_k|X,\theta)\, p(\theta|\beta)d\theta$$

$$909 \quad = \int_\theta \left(\prod_i \prod_j \theta_j^{N_i^j} \right) \frac{\Gamma(\sum_j \beta_j)}{\prod_j \Gamma(\beta_j)} \prod_j \theta_j^{\beta_j - 1}\, d\theta$$

$$910 \quad = \int_\theta \left(\prod_j \theta_j^{n_k^j} \right) \frac{\Gamma(\sum_j \beta_j)}{\prod_j \Gamma(\beta_j)} \prod_j \theta_j^{\beta_j - 1}\, d\theta$$

$$911 \quad = \int_\theta \frac{\Gamma(\sum_j \beta_j)}{\prod_j \Gamma(\beta_j)} \prod_j \theta_j^{n_k^j + \beta_j - 1}\, d\theta$$

$$912 \quad = \frac{\Gamma(\sum_j \beta_j)}{\Gamma(n_k + \sum_j \beta_j)} \frac{\prod_j \Gamma(n_k^j + \beta_j)}{\prod_j \Gamma(\beta_j)} \int_\theta \frac{\Gamma(n_k + \sum_j \beta_j)}{\prod_j \Gamma(n_k^j + \beta_j)} \prod_j \theta_j^{n_k^j + \beta_j - 1}\, d\theta$$

$$913 \quad = \frac{\Gamma(\sum_j \beta_j)}{\Gamma(n_k + \sum_j \beta_j)} \frac{\prod_j \Gamma(n_k^j + \beta_j)}{\prod_j \Gamma(\beta_j)}$$

where the last equality follows because the integrand is the density function of a Dirichlet

distribution with parameter $\left(n_k^1 + \beta_1, \ldots, n_k^J + \beta_J\right)$. By assuming that $\beta_j = \beta$ for all j this reduces

to equation (1) in the main text.

Derivation of $p(X|\alpha)$

Under the fully symmetric Poisson model the probability of change across an ancestral-

descendant branch is,

$$p\left(X_i|X_{pa(i)}, \lambda_i, t_i\right) = \left(1 - e^{-K\lambda_i t_i}\right)\frac{1}{K} + \delta_{X_i X_{pa(i)}} e^{-K\lambda_i t_i}$$

where λ_i is the branch-specific rate of evolution, t_i is the length of the branch in units of

time, and $\delta_{X_i X_{pa(i)}}$ is the Kronecker delta. Because only the product $\lambda_i t_i$ matters, we can set λ_i

equal to $\frac{1}{K-1}$ (meaning that the average rate is normalized to 1) so that that $t_i = v_i$ now measures

time in units of expected number of changes. We assume that each $v_i \sim \text{Gamma}(\alpha, 1)$. Then,

$$p\left(X_i|X_{pa(i)} \neq X_i, \alpha\right)$$

$$= \int_0^\infty p\left(X_i|X_{pa(i)}, v\right)p(v|\alpha)dv$$

$$= \int_0^\infty \left(1 - e^{-\frac{K}{K-1}v}\right)\frac{1}{K}\frac{1}{\Gamma(\alpha)}v^{\alpha-1}e^{-v}dv$$

$$= \frac{1}{K} - \frac{1}{K}\int_0^\infty \frac{1}{\Gamma(\alpha)}v^{\alpha-1}e^{-v\left(\frac{K}{K-1}+1\right)}dv$$

$$= \frac{1}{K} - \frac{1}{K}\left(\frac{1}{\frac{K}{K-1}+1}\right)^\alpha \int_0^\infty \frac{\left(\frac{K}{K-1}+1\right)^\alpha}{\Gamma(\alpha)}v^{\alpha-1}e^{-v\left(\frac{K}{K-1}+1\right)}dv$$

$$= \frac{1}{K} - \frac{1}{K}\left(\frac{1}{\frac{K}{K-1}+1}\right)^\alpha$$

where the last equality follows because the integrand is the density function of a Gamma

distribution with parameters $\left(\alpha, \frac{K}{K-1} + 1\right)$. A similar calculation shows that

$p\left(X_i \middle| X_{pa(i)} = X_i, \alpha\right) = \frac{1}{K} + \frac{K-1}{K}\left(\frac{1}{\frac{K}{K-1}+1}\right)^{\alpha}.$