Visual pathways in the brain of the jumping spider *Marpissa muscosa*

Running title: **Jumping spider visual pathways**

Philip O.M. Steinhoff[1*], Gabriele Uhl[1], Steffen Harzsch[2] & Andy Sombke[3]

[1] Zoological Institute and Museum, General and Systematic Zoology, University of Greifswald, Loitzer Straße 26, 17489 Greifswald, Germany

[2] Zoological Institute and Museum, Cytology and Evolutionary Biology, University of Greifswald, Soldmannstraße 23, 17489 Greifswald, Germany

[3] Department of Integrative Zoology, University of Vienna, Althanstraße 14, 1090 Vienna, Austria

Corresponding Author. E-Mail: philipsteinhoff@gmail.com

ACKNOWLEDGEMENTS

The authors wish to thank Georg Brenneis and Matthes Kenning (University of Greifswald) as well as Nicholas J. Strausfeld (University of Arizona, Tucson) and Friedrich G. Barth (University of Vienna) for helpful discussions. We are grateful to Kerstin Stejskal and Thomas Hummel (Department of Neurobiology, University of Vienna) for providing *Cupiennius salei* and Elvira Lutjanov, Christian Müller, Sabine Ziesemer and Jan-Peter Hildebrandt (all University of Greifswald) for help with western blot analysis. Tim M. Dederichs and Peter Michalik (both University of Greifswald) helped to obtain pictures with the Visionary Digital camera system. Monica M. Sheffer provided helpful linguistic comments that improved the manuscript. This study was financially supported by the German Research Foundation DFG UH 87/11-1 and DFG INST 292/119-1 FUGG, DFG INST 292/120-1 FUGG. POMS was additionally supported by a Bogislaw scholarship from the University of Greifswald and a Laudier Histology cooperation & travel grant.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author upon reasonable request.

Abstract

Some animals have evolved task differentiation among their eyes. A particular example is spiders, where most species have eight eyes, of which two (the principal eyes) are used for object discrimination, whereas the other three pairs (secondary eyes) detect movement. In the spider species *Cupiennius salei* these two eye types correspond to two visual pathways in the brain. Each eye is associated with its own first and second order visual neuropil. The

second order neuropils of the principal eyes are connected to the arcuate body, whereas the second order neuropils of the secondary eyes are linked to the mushroom body. However, eye size and visual fields are considerably different in jumping spiders. We explored the principal- and secondary eye visual pathways of the jumping spider *Marpissa muscosa*. We found that the connectivity of the principal eye pathway is the same as in *C. salei*, while there are differences in the secondary eye pathways. In *M. muscosa,* all secondary eyes are connected to their own first order visual neuropils. The first order visual neuropils of the anterior lateral and posterior lateral eyes are further connected with two second order visual neuropils, whereas the posterior median eyes lack second order visual neuropils and their axons project only to the arcuate body. This suggests that the posterior median eyes probably do not serve movement detection in *M. muscosa.* Furthermore, the second order visual neuropil (L2) in *Marpissa muscosa* potentially integrates information from the secondary eyes and might thus enable faster movement decisions.

Keywords

Secondary eyes, principal eyes, visual pathway, jumping spider, visual neuropils, brain, *Marpissa muscosa*, RRID AB_528479, RRID AB_2637882, RRID AB_477585, RRID AB_1541510, RRID AB_2338680, RRID AB_2534074, RRID AB_2315147

1 Introduction

In some animal species with multiple eyes, different eyes serve different specific tasks, such as compound eyes and ocelli in insects or the rhopalia in box jellyfishes (Garm et al., 2007; O'Connor et al., 2009; Paulus, 1979). Spiders vary strongly in their visual abilities, as well as in number, arrangement and size of their eyes (reviewed in Morehouse et al. 2017). Most species possess four pairs of eyes, of which one pair differs from the three other pairs in its anatomy and developmental origin (Land, 1985a; Morehouse et al., 2017). The principal eyes (or anterior median eyes) possess a movable retina and their spectral sensitivities allow for color vision (Barth et al. 1993; Land, 1985b; Schmid, 1998; Yamashita & Tateda, 1976, 1978). The other three pairs of eyes are the so-called secondary eyes (posterior median, anterior lateral and posterior lateral eyes). Their morphology and anatomy differs considerably among spider families (Homann, 1952, 1950), but they generally do not have a movable retina. In several groups of cursorial hunting spiders, the secondary eyes are used for movement detection (Land, 1985a; Morehouse et al., 2017; Schmid, 1998). Most secondary eyes possess

a light-reflecting tapetum (but absent in e.g. jumping spiders), which led to the assumption

that they play a major role for night or dim light vision (Foelix, 1996).

Jumping spiders (Salticidae) are cursorial hunters with excellent vision. Their eyes have been

studied in much greater detail compared to any other spider group (e.g. Harland et al., 2012;

Land, 1985a; Morehouse et al., 2017). The principal eyes of jumping spiders are large and face

forward; their field of view is small, but their spatial acuity is one of the highest among

invertebrate eyes (Harland et al., 2012; Land, 1985b, 1985a). The secondary eyes of salticids

possess only one photoreceptor type and thus cannot discriminate colors (Yamashita and

Tateda, 1976). Among the secondary eyes, the anterior lateral eyes (ALE) and posterior lateral

eyes (PLE) are large, able to detect motion, and elicit an orienting response of the principal

eyes towards the moving object (Duelli, 1978; Jakob et al., 2018; Spano et al., 2012; Zurek et

al., 2010; Zurek and Nelson, 2012a, 2012b). The posterior median eyes (PME) of most salticids

(except for the non-salticoids) are very small and positioned between ALE and PLE (Land,

1985b; Maddison and Hedin, 2003). Since the fields of view of ALE and PLE overlap, the PME

do not have an exclusive field of view (Land, 1985b).

While the eyes of jumping spiders have been studied in detail, less is known about the brain

structures that process visual information (Duelli, 1980; but see Hanström, 1921; Hill, 1975;

Nagata et al., 2019; Steinhoff et al., 2017; Strausfeld, 2012). Our knowledge on the visual

processing pathways in the brain of spiders is mostly based on investigations of the spider

model species *Cupiennius salei* (Keyserling, 1877) (Babu and Barth, 1984; Becherer and

Schmid, 1999; Schmid and Duncker, 1993; Strausfeld et al., 1993; Strausfeld and Barth, 1993).

C. salei possesses two separate visual pathways, one for the principal eyes and one for all

secondary eyes (Strausfeld et al., 1993; Strausfeld and Barth, 1993). In *C. salei*, every eye is

connected to its own first and second order visual neuropil (Barth, 2002). While the second

order visual neuropils of the principal eyes are connected to the arcuate body (Strausfeld et

al., 1993), the second order visual neuropils of the secondary eyes are associated with the

mushroom bodies (Strausfeld and Barth, 1993). Additionally, the first order visual neuropils

of the secondary eyes are directly connected to the arcuate body (Babu and Barth, 1984).

In *C. salei,* similar to salticids, the principal eyes are used for object recognition, while the

secondary eyes serve the purpose of movement detection (Fenk, Hoinkes, & Schmid, 2010;

Schmid, 1998). The secondary eyes of *C. salei* have a high spectral sensitivity, similar to their

principal eyes (Barth et al., 1993) and similar to jumping spiders are capable of acute motion

detection (Fenk & Schmid, 2010; Zurek & Nelson, 2012b). However, they are unable to detect

color in moving objects (Orlando and Schmid, 2011) and the relative size of different eyes and

thus fields of view in *C. salei* differ considerably from jumping spiders (Land, 1985b; Land and

Barth, 1992). Furthermore, Steinhoff et al. (2017) showed that the structure and arrangement

of the secondary eye visual neuropils in the jumping spider *Marpissa muscosa* (Clerck, 1757)

differ from those in *C. salei.* We thus hypothesize, that the secondary eye visual pathways of

jumping spiders also differ from that in the model species *C. salei.* Here, we describe the

principal and secondary eye visual pathways in the brain of the jumping spider *M. muscosa.*

We use paraffin histology, immunohistochemistry, and microCT analysis to visualize tracts in

the visual system. We compare our findings in *M. muscosa* with results of earlier studies and

new immunohistochemical data on the principal and secondary eye pathways in the brain of

C. salei. We discuss the similarities and differences in the principal and secondary eye

pathways of these two spider species in the light of different functional roles of their eyes.

2 Material and Methods

2.1 Animals

123 Adult female *Marpissa muscosa* were collected in and near Greifswald (Germany). Spiders

124 were fed on *Drosophila* sp. weekly and kept individually in plastic boxes of 145 x 110 x 68 mm

125 size that were enriched with paper tissue.

2.2 Paraffin-histology

The prosomata of two females of *M. muscosa* were fixed in Gregory's artificially aged Bouin

solution (Gregory, 1980). After four days in fixative, the prosomata were dehydrated through

a graded ethanol series (80%, 90%, and 3x 96% ethanol for 20 minutes each) and were then

transferred for two hours into a 1:1 solution of 96% ethanol:tetrahydrofuran (Carl Roth,

CP82.1) at room temperature. Samples were kept for 24 hours in pure tetrahydrofuran

followed by 24 hours in a solution of 1:1 tetrahydrofuran and paraffin (Carl Roth, 6643.1) at

60°C. Afterwards, samples were transferred to 100% paraffin at 60°C for 48 hours and then

embedded in fresh paraffin. Sagittal and transversal sections (5 µm) were cut with a

motorized rotary microtome (Microm HM 360). Sections were stained with Azan according to

Geidies (Schulze and Graupner, 1960), and mounted in Roti-Histokitt II (Carl Roth, T160.1).

2.3 MicroCT analysis

Prosomata of two individuals of *M. muscosa* and one individual of *C. salei* were fixed in

Dubosq–Brazil solution (1 g saturated alcoholic picric acid, 150 ml 80% ethanol, 15 ml pure

acetic acid, and 60 ml 37% formaldehyde). Prior to fixation, legs and opisthosomata were cut

off and cuticle and musculature were removed from the dorsocaudal part of the prosomata

to allow faster penetration of the fixative. After four days in fixative, samples were washed

six times for 20 minutes in 0.1 M phosphate buffered saline (PBS, pH 7.4) followed by

dehydration in a graded ethanol series (80%, 90%, 96% and 3x 99,8% ethanol for 20 minutes

each). Samples were then transferred to an 1% iodine solution (iodine, resublimated [Carl

Roth, X864.1] in 99.8% ethanol) over 48 h to enhance tissue contrast (Sombke et al., 2015)

and subsequently critical point dried with an automated dryer (Leica EM CPD300). The

protocol applied was: slow CO_2 admittance with a delay of 120 seconds, 30 exchange cycles,

followed by a slow heating process and slow gas discharge. Dried prosomata were mounted

using a conventional glue gun onto aluminum rods, so that the anterior median eyes were

oriented upwards.

MicroCT scans were performed using a Zeiss XradiaXCT-200 (Sombke et al., 2015). Scans were

performed with a macro-, a 4x- and a 20x objective lens unit and the following settings: 40 kV

and 8 W or 30 kV and 6 W, 200 µA, exposure times between 2 and 4 s. Tomographic

projections were reconstructed with the XMReconstructor software (Zeiss), resulting in image

stacks (TIFF format). Scans were performed using Binning 2 for noise reduction and were

reconstructed with full resolution (using Binning 1).

2.4 Immunohistochemistry

Four different combinations of markers were used to visualize neuropils and connecting

neurites in the visual system (see Table 1 for a list of marker combinations used, and Table 2

for specifications of labeling reagents).

After anaesthetization with CO_2, brains of 31 females of *M. muscosa* (details Tab. 2) and four

females of *C. salei* were dissected in PBS and fixed in 4% paraformaldehyde over night at room

temperature. Specimens for histamine immunohistochemistry were prefixed overnight at 4°C

in 4% N-(3-Dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride (EDAC, Sigma–Aldrich

E6383). Brains were washed six times for 15 minutes each in PBS and subsequently immersed

in Poly-L-Lysin (Sigma-Aldrich, P4707) for one minute. Samples were then embedded in 4%

agarose (Sigma-Aldrich, A9414) and sectioned (50 or 80 µm) with a Microm HM 650 V

vibratome. After permeabilization in PBS-TX (PBS, 0.3% Triton X-100 [Sigma-Aldrich, X100], 1

% bovine serum albumin [Sigma-Aldrich, A2153], 0.05% Na-acide; or alternatively PBS, 0.5%

Triton, 1.5% DMSO) for 1 h at room temperature, incubation in primary antibodies took place

over night at 4°C or room temperature. Sections were washed in several changes of PBS-TX

after the incubation. Incubation in secondary antibodies or counterstains took place

overnight at 4°C or room temperature. After washing in several changes of PBS, sections were

mounted in Mowiol (Merck 475904).

The brains of five additional females of *M. muscosa* were processed as whole-mounts, using

an adjusted protocol after Ott (2008) (see also Steinhoff et al., 2017). Brains were dissected

in TRIS buffer (0.1 M, pH 7.4 [Carl Roth, 4855]), and fixed in zinc-formaldehyde at room

temperature under soft agitation for several days. Afterwards, brains were washed in TRIS

buffer three times for 15 minutes each and incubated in a 4:1 mixture of methanol and DMSO

(Dimethylsulfoxide [Carl Roth, 4720]) for 2 h. After incubation in 100% methanol for 1 h,

brains were rehydrated in a graded series of methanol (90%, 70%, 50%, 30%) for 10 minutes

each and washed in TRIS buffer. This was followed by permeabilization in PBS-DMSO (PBS, 5%

bovine serum albumin, 1% DMSO, 0.05% Na-acide) for 2 h at room temperature and

incubation in primary antibody for 4 days at 4°C. Brains were then washed several times for

2 h in PBS-DMSO and incubated in secondary antibody for 4 days at 4°C, and transferred to a

graded series of glycerol, with 1%, 2%, 4% (for two hours each) and 8%, 15%, 30%, 50%, 60%,

70%, 80% (for one hour each) glycerol in TRIS buffer and 1% DMSO. After five changes in pure

ethanol for 30 minutes each, brains were mounted in fresh methyl salicylate (Sigma, 76631).

In control experiments, in which we replaced the primary antibodies with PBS-TX, no neuronal

labeling was detected.

2.5 Western Blot

In western blots of *Drosophila melanogaster* Meigen, 1830 head homogenates (Klagges et al.,

1996), the monoclonal antibody mouse anti-*Drosophila* synapsin ''Synorf 1'' (provided by E.

Buchner, Universität Würzburg, Germany; raised against a *Drosophila* Glutathione S-

Transferase(GST)-synapsin fusion protein) recognizes at least 4 synapsin isoforms (ca. 70, 74,

80, and 143 kDa). In western blot analyses in Crustacea (Harzsch and Hansson, 2008; Sullivan

et al., 2007) and Chilopoda (Sombke et al., 2011), isoforms in the same range (75-90 kDA)

were recognized. Furthermore, the Synorf 1 antibody labels synaptic neuropil in taxa as

diverse as Araneae (Fabian-Fine et al., 1999; Nagata et al., 2019; Steinhoff et al., 2017;

Widmer et al., 2005), Chaetognatha (Harzsch and Müller, 2007) and Plathelminthes (Cebrià,

2008), suggesting that the epitope that this antiserum recognizes is highly conserved. To test

this, we conducted a western blot analysis as well, in which we compared brain tissue of *D.*

melanogaster and *M. muscosa*. The antibody provided similar results for both species,

staining one strong band around 70 kDa, and in *M. muscosa* another weak band around 60

kDa. This result suggests that the epitope that Synorf 1 recognizes is conserved between *D.*

melanogaster and *M. muscosa*.

2.6 Microscopy, image processing and nomenclature

Immunohistochemically labelled sections and whole-mounts were examined and

photographed with a Leica SP5 II confocal microscope (cLSM). Paraffin sections were

examined and photographed with a customized Visionary Digital BK Plus Lab System

(duninc.com/bk-plus-lab-system.html). 3D-visualization of microCT image stacks was

prepared in AMIRA 6.0.0 (ThermoFisher). An interactive 3D-visualization of the microCT

reconstructions was created using ImageJ 1.51n and Adobe Acrobat Pro 9.0 (see supporting

information; Figure S1). Images were processed in Corel PaintShop Pro using global contrast

and brightness adjustment features as well as black and white inversion. Schematic drawings

were generated using Corel Draw 20.1 and figure plates were assembled in Corel PaintShop

Pro 21.0. A color code for all depicted structures, as used in Loesel et al. (2013) and Steinhoff

et al. (2017), was used for 3D visualizations and the schematic drawings. The terminology

used for brain structures follows Richter et al. (2010). For structures specific to spiders, we

use terms according to Lehmann et al. (2016), Strausfeld et al. (1993) and Strausfeld and Barth

(1993). Positional information is given with respect to the body axis. We are referring to

results from immunohistochemical experiments as immunoreactivity, e.g. synapsin-

immunoreactivity.

231
232 **Abbreviations**
233
AB   Arcuate body
ALE   Anterior lateral eyes
AL1   First order visual neuropils of ALE
AL2/PL2   Second order visual neuropils of ALE and PLE
AL1x   Neuropilar subunit of the AL1
AME   Anterior median eyes
AM1   First order visual neuropils of AME
AM2   Second order visual neuropils of AME
L2   Second order visual neuropils of anterior and posterior lateral eyes
MB   Mushroom bodies (neuropil with attributes of a mushroom body-like organization)
MBbr   Mushroom body bridge

245	**MBn**	Neurites between pedunculus and AL2/PL2
246	**MBh**	Mushroom body haft
247	**MBp**	Pedunculus of the mushroom body
248	**MBs**	Mushroom body shaft
249	**PLE**	Posterior lateral eyes
250	**PL1**	First order visual neuropils of the PLE
251	**PL1x**	Neuropilar subunit of the PL1
252	**PME**	Posterior median eyes
253	**PM1**	First order visual neuropils of the PME
254	**PM2**	Second order visual neuropils of the PME
255	**SC1-3**	Protocerebral soma cluster 1-3

3 Results

3.1 General organization of the protocerebrum

In *Marpissa muscosa,* the protocerebrum comprises bilaterally paired neuropils of the visual

system and the mushroom body as well as the unpaired arcuate body and the protocerebral

neuropil (Figs. 1, 2; for details see Steinhoff et al., 2017). All neuropils of the visual system are

recognizable in paraffin sections (Fig. 3) and microCT analysis (Fig. 4; supporting information,

Fig. S1), and are also characterized by a strong synapsin-immunoreactivity (Figs. 1; 5; 6; 7a,

b). Tracts and individual neurites exhibit prominent tubulin-immunoreactivity (Figs. 1; 2; 5; 6;

7). Furthermore, the optic nerves (between retinae and first order visual neuropils) and the

corresponding first order visual neuropils display histamine-   immunoreactivity (Fig. 8),

which is known to consistently label retinula cell axons and their terminals in the visual

systems of arthropods (Battelle et al., 1991; Harzsch et al., 2006; Harzsch, Wildt, Battelle, &

Waloszek, 2005; Nässel, 1999; Schmid & Duncker, 1993; Sombke & Harzsch, 2015). A soma

cortex including three discernable protocerebral clusters surrounds most of the central

nervous system (CNS) (Figs. 2; 5a; 6a-b, d; 7a–d). Soma cluster 1 (SC1) is located between the

first order visual neuropils of the lateral eyes and the second order visual neuropil of anterior

lateral and posterior lateral eyes (AL2/PL2), cluster 2 (SC2) medially to the AL2/PL2, and

cluster 3 (SC3) anteromedially to the mushroom body bridge (MBbr) (Fig. 2).

3.2 The principal eye pathway

The retina of the anterior median eyes (AME) is connected to its first order visual neuropil

(AM1), which consists of at least two, but most likely four tightly adjoining subunits (Fig.

1a,e,f). The AM1 is situated close to the second order visual neuropil of the AME (AM2) and

connected to it *via* parallel neurites (Fig. 1e,f). The AM2 is large, oval shaped and its posterior

margin within the protocerebral neuropil is not clearly recognizable in anti-synapsin labeling

(Fig. 1). Anti-tubulin labeling reveals a tract of neurites that connects the AM2 with the lateral

flanges of the AB, as well as some neurites with unclear termination that penetrate the

protocerebral neuropil (Fig. 1c-e).

3.3 Connectivity of the first order visual neuropils of the anterior and posterior lateral eyes

The retinae of the anterior lateral eyes (ALE) and posterior lateral eyes (PLE) are connected

with their respective first order visual neuropils (AL1 and PL1) *via* short, thick optic nerves

(Figs. 3b, c; 4a-b; 5c). A subpopulation of histaminergic neurites, which are interpreted here

as retinula cell axons, terminates in distinct distal layers of these neuropils (AL1x and PL1x;

Fig. 8a–c). Both the AL1 and PL1 are connected to the second order visual neuropil of the

lateral eyes (L2) and the AL2/PL2 *via* prominent tracts of parallel ordered neurites that exhibit

strong tubulin-immunoreactivity (Figs. 2; 5c; 6a-b, d; 7a-b). This connectivity is also evident in

both, histological sections (Fig. 3) and microCT analysis (Fig. 4). The tracts are predominantly

formed by neurites that originate in SC1, which surrounds this region of the anterior

protocerebrum laterally, dorsally and medially (Figs. 2; 6a-b; 7b, d). The L2 and the AL2/PL2

are bi-lobed, which corresponds to the separate inputs from AL1 and PL1 (Figs. 3a; 3b, d; 5a;

6a, C; 8a, b). In histological sections, a thin but conspicuous tract is detectable that passes

between L2 and AL2/PL2 and likely terminates in the arcuate body (AB) (Figs. 3b, c; 5a, b; 6d;

7b). The neurites forming this tract might originate in SC1 and further innervate the AL1x and

PL1x, as well as the PM1 (see below; Figs. 3; 5a, b; 6d).

3.4   Connectivity of the first order visual neuropils of the posterior median eyes

The posterior median eyes (PME) are much smaller than all three other pairs of secondary

eyes, and so is their retina (Fig. 4c, e). Because of the small retina, only a thin but long nerve,

which enters the protocerebrum between the AL1 and PL1, connects the PME to its first order

visual neuropil (PM1) (Figs. 4e; 6c, c'). The PM1 is situated between AL1 and PL1 (Figs. 3a-b;

4e, 5b; 6c-d; 7b; 8a). A prominent tract projects from the PM1 to the dorsoventral part of the

protocerebrum and seems to terminate in the anterolateral apices of the AB (Fig. 3b; 5b; 6d).

3.5   The second order visual neuropil of ALE and PLE and the mushroom body

The second order visual neuropil of ALE and PLE (AL2/PL2) is of glomerular organization and

composed of two lobes that tightly adjoin dorsally and are fused in their ventral part (Figs. 7a,

c, d; 9a-b). Thus, only a uniform array of glomeruli is visible in most sections (Figs. 3b; 4a, c,

e; 5; 6). Each glomerulus is composed of a high number of synaptic complexes (Fig. 9c–e).

These synaptic complexes can be visualized by anti-synapsin labeling and phalloidin-labeling
(Fig. 9d, e). Co-localization of synapsin and phalloidin is almost absent (Fig. 9). Individual
glomeruli are innervated by neurites originating in the AL1 and PL1 (Figs. 5c; 6a, b, d; 7d).
Anti-tubulin labeling reveals that neurites interweave the array of glomeruli mostly in parallel
from proximal to distal (Fig. 5c; 6a). Within the core-region of the AL2/PL2, tubulin-
immunoreactive neurites are more abundant (Fig. 6a,c. Two neurite bundles (MBn) connect
the AL2/PL2 and the mushroom body pedunculus (MBp) and form a distinct chiasm close to
the AL2/PL2 (Figs. 7a, c, d; 9a). Neurites from the medial SC2 contribute to these bundles by
projecting into the AL2/PL2 and into the MBp (Figs. 2; 6a-c; 7a, d). Other neurites from SC2
pass between AL2/PL2 and MBp towards the L2 (Fig. 6b, d). Thus, a large number of
interweaving neurites characterizes the region of the MBn, which makes tracing of individual
trajectories difficult (Fig. 7a, d).

Some long neurites that appear to originate in SC2 project along the periphery of the MBp
and join with neurites arising from the mushroom body haft (MBh). Together they form the
mushroom body bridge (MBbr) (Figs. 2; 6c; 7a, c, d). Other peripheral neurites from SC2 enter
the MBh, where they seem to terminate (Figs. 2; 4b, d, f; 7d)

3.6   Visual neuropils and the mushroom body in *Cupiennius salei*

The dorsalmost neuropils in the protocerebrum of *C. salei* are the AM1 and AM2, which are
connected by numerous intertwining neurites (Fig. 10a). The AM2 is connected to the AB *via*
a thin tract (Fig. 10a). Located ventrally and somewhat anterior to the AM1 is the PM1, which
is large and spans over one hemisphere of the protocerebrum at its broadest part (Fig.
10a,b,e). Its shape is cup-like, but somewhat irregular (Fig. 10e). The PL1 is situated
anterioventral of the PM1, is similar in size, but horseshoe shaped (Fig. 10b). Both the PM1
and the PL1 are subdivided into an upper, strongly synapsin-immunoreactive part where the
retinula cell axons terminate and a lower part where the synapsin-immunoreactivity is more
patchily distributed (PM1x and PL1x; Fig. 10b,e). Tubulin-immunoreactivity is strongest in the
lower part, where the neurites that connect the PM1 and PM2 with their respective second
order neuropils bundle together. The AL1 is the smallest first-order neuropil and located close
to the midline, between PM1 and PL1 (Fig. 10b). Each first order visual neuropil is connected
to its second order visual neuropil. These are of glomerular organization and situated close to
each other (Fig. 10b,c). While the AL2 is horseshoe shaped, the PL2 and PM2 are more
elongate in shape and also larger (Fig. 10b,c).

The mushroom body is characterized by strong synapsin-immunoreactivity and consists of a
pedunculus (MBp), a shaft (MBs) and a haft (MBh) (Fig. 10b). The MBs and the MBh are fused
with the MBp, and together form one continuous synapsin-immunoreactive domain (Fig.
10b).

4 Discussion

Comparative neuroanatomical studies on different spider species show that while general
brain anatomy is similar, the structure of different neuropils especially within the
protocerebrum varies (Hanström, 1921; Long, 2016; Steinhoff et al., 2017; Weltzien and
Barth, 1991). The previous investigation by Steinhoff et al. (2017) on the general brain
anatomy of *Marpissa muscosa* suggested that the brains of jumping spiders and the spider
species *Cupiennius salei* differ in structure and arrangement of neuropils. Although the
general architecture is similar, we have shown here that the connectivity of neuropils in the
secondary eye pathways differ between these two species, which suggests differences in
information processing and function of secondary eye visual neuropils.

The relative size and position of the secondary eyes differs between *M. muscosa* and *C. salei.*
While in *M. muscosa* the PME is very small and oriented sideways, it is the largest eye in *C.*
salei and oriented forward (Morehouse et al., 2017; Fig. 11). The PLE are similar in relative
size and position, but the ALE are considerably smaller in *C. salei,* albeit also oriented forward
(Fig. 11). The relative size of the retinae as well as the brain is smaller in *C. salei*, which is why
the optic nerves associated with the eyes are much longer in *C. salei* (Fig. 11).

4.1 Connectivity of neuropils in the principal eye pathway

The connection of the AME retinula cell axons to the AM1 has recently been described in
detail by Nagata et al. (2019) for the jumping spider *Hasarius adansoni*. These authors
revealed that there are four photoreceptor layers in the retina, which each connect to one of
four subregions in the AM1 (Nagata et al., 2019). We show here, that *in M. muscosa* the AM1
is further connected to the AM2, which is the same as described for *C. salei* (Strausfeld et al.,
1993). In *C. salei*, two types of neurons have been identified whose axons connect the AM2
with the flanges and the inner part of the AB (Strausfeld et al., 1993). In *M. muscosa* this
pattern is likely the same, although we were only able to unambiguously detect neurites that
connect the AM2 with the flanges of the AB. There are, however, neurites that project from

the AM2 into the protocerebral neuropil and some of those may well proceed towards the

inner part of the AB.

4.2 Connectivity of the first order visual neuropils of the anterior and posterior lateral eyes

In *M. muscosa*, all retinula cell axons from ALE, PLE and PME seem to terminate in their

associated first order visual neuropils (Fig. 8). This is in accordance to what has been found in

other Araneae (Becherer and Schmid, 1999; Duelli, 1980; Schmid and Duncker, 1993;

Strausfeld et al., 1993; Strausfeld and Barth, 1993). Nevertheless, this pattern is different in

all other investigated taxa of Chelicerata (e.g. Scorpiones, Xiphosura, or Pycnogonida), where

a subset of retinula cell axons also projects into the second order visual neuropils (Battelle,

Sombke, & Harzsch, 2016; Lehmann & Melzer, 2013, 2018). It should be noted, however, that

this discrepancy between spiders and all other arachnids still needs to be scrutinized using

cobalt fills, which have not been employed in spiders yet (Lehmann and Melzer, 2018). In *C.*

salei, a few histaminergic neurites terminate in all second order visual neuropils, and have

been interpreted as interneurons (Schmid and Duncker, 1993). However, as long as no

neuronal tracings have been carried out, it cannot be ruled out that these neurites might

instead represent retinula cell axons. In *M. muscosa*, we did not detect any histaminergic

neurites that terminate in the second order visual neuropils. There are, however, numerous

histaminergic neurites that terminate in AL1x and PL1x and we interpret those as retinula cell

axons.

In our histological sections, it appears as if some neurites that contribute to the tract between

AL1, PL1 and the AB, arise in AL1x and PL1x respectively (Fig. 3b, c; 5a-b). Hill (1975) described

in *Phidippus sp.* that neurites from the AL1x domain contribute to a tract projecting towards

the posterior protocerebrum. However, Hill (1975) did not mention terminations of retinula

cell axons in the AL1x and PL1x, but detected neurites connecting the AL1 with the AL1x. Thus,

it is possible that Hill (1975) either misinterpreted retinula cell axons as interneurons, or that

we missed connecting neurites between AL1 and AL1x. In our immunohistological

investigations, we found that synapsin-immunoreactive substructures of first order visual

neuropils are also present in *C. salei*, the PM1x and the PL1x. In both, *M. muscosa* and *C. salei,*

this region is additionally characterized by strong tubulin-immunoreactivity. This can be

explained by the tight packing of neurites that converge at the level of the AL1x and PL1x in

M. muscosa and the PM1x and PL1x in *C. salei* to form the tracts to the respective second

order visual neuropils. A subdivision of PM1 and PL1 is also visible in histamine-

immunolabeled material from the study by Schmid & Duncker (1993; compare Figs. 2c,d and

3a,b), although not discussed by the authors. It remains open whether retinula cell axons or

interneurons terminate at the synapsin immunoreactive part of the PL1x and PM1x in *C. salei*

and the AL1x and PL1x in *M. muscosa*.

In *M. muscosa*, the convoluted first order visual neuropils of the ALE and the PLE are both

connected with the AL2/PL2 and the L2, as well as with the AB, which is only partially

consistent with Hill's (1975) study on the nervous system in *Phidippus* spp., where he found

that axons of the AL1, and not the PL1, join the tract that projects towards the AB. Duelli

(1980), did not detect connections between PL1 and L2 in the salticids *Evarcha arcuta* (Clerck,

1757) and *Pseudamycus validus* (Thorell, 1877). He described "large field horizontal neurons,"

closely bypassing the AL2/PL2 and L2 and connecting the PL1 with the posterior

protocerebrum (Duelli, 1980). Judging from their structure and position, these neurons might

be identical with the neurites described in our study which contribute to the tract that

connects the first order visual neuropils of the secondary eyes with the arcuate body. This

discrepancy might be due to incomplete staining in the earlier studies where reduced-silver

staining was employed. Our approach combines specific and broader markers with specific

staining techniques. It should also be noted that the studies on the secondary eye pathways

in jumping spiders were conducted on different species. However, it seems unlikely that the

secondary eye pathways differ considerably within salticoid jumping spiders (Maddison and

Hedin, 2003; Morehouse et al., 2017).

Based on our results, AL2/PL2 and L2 are shared second order visual neuropils of both AL1

and PL1 in *M. muscosa*. The situation in *C. salei* is, however, different. Here, the first order

visual neuropils of all three secondary eyes are connected to individual second order visual

neuropils (compare Fig. 11a and b) (Strausfeld and Barth, 1993). Each of these second order

visual neuropils (AL2, PL2, PM2) is composed of small neuropilar glomeruli (compare Fig. 11).

An unstructured second order visual neuropil (without glomeruli), such as the L2, is absent in

C. salei (Strausfeld and Barth, 1993). Similar to the "giant interneurons" described for jumping

spiders by Duelli (1980), Strausfeld & Barth (1993) detected "lamina wide-field neurons" that

connect all first order visual neuropils with a region close to the AB. It is very likely that these

neurons correspond to the axons forming the tract between first order visual neuropils and

the AB that we described in *M. muscosa*.

4.3   First order visual neuropils of the posterior median eyes

The PM1 is by far the smallest first order visual neuropil of all secondary eyes in *M. muscosa*.
This correlates with the much smaller overall size of the PME (compared to ALE and PLE). The
PME have previously been considered vestigial (Land, 1985a, 1972), but since there is an optic
nerve and a corresponding first order visual neuropil, we conclude that these eyes are
functional. Previously, Steinhoff et al. (2017) considered the PM1 to be a second order visual
neuropil ("PM2"), based on its close spatial relationship with the AL1x. However, we now
demonstrate that the PM1 is innervated only by retinula cell axons originating in the PME,
and connections between PM1 and AL1x seem to be absent. This coincides with Hill's (1975)
description for *Phidippus* spp.*,* although he described additional neurites connecting PL1 and
PM1. It is likely that these neurites contribute to the tract that connects the first order visual
neuropils with the AB (compare Fig. 11a), instead of actually forming synapses in the PM1.
The PM1 differs from all other eyes by being connected to the AB only. Consequently, the
pathway of the PME lacks a second order visual neuropil. Information from this eye is not
processed within the mushroom bodies, but exclusively in the arcuate body.
461

4.4   Second order visual neuropils of the ALE and PLE
Visual neuropils with a glomerular structure have been found in the protocerebrum of several
spider species (e.g. Long, 2016). They have been termed "lame glomérulée" by Saint-Remy
(1887), "glomeruli" by Hanström (1921) and Babu & Barth (1984), "spherules" by Strausfeld
& Barth (1993) and "microglomeruli" by Steinhoff et al. (2017). The presence and
arrangement of these glomerular neuropils differs between spider species (Hanström, 1921;
Long, 2016) and some authors have considered them as a substructure of the MB (Babu and
Barth, 1984; Hill, 1975), whereas others have considered them to be separate second order
visual neuropils (Long, 2016; discussed in Steinhoff et al., 2017; Strausfeld and Barth, 1993).
Our results demonstrate, that in *M. muscosa*, the AL2/PL2 represents two closely adjoining
second order visual neuropils of the ALE and PLE, respectively. In *M. muscosa*, the AL2/PL2
was also found to react less plastically to different environmental conditions than the MBp
(Steinhoff et al., 2018), which was interpreted as further evidence for their function as
discrete visual neuropils. Within the AL2/PL2, each glomerulus seems to consist of numerous
synaptic complexes and Duelli (1980) estimated roughly 400 synapses per glomerulus in the
jumping spider *Evarcha arcuta*. Studies in insects used anti-synapsin labeling to detect
presynaptic sites, and phalloidin (which labels f-actin) to highlight postsynaptic sites of
synaptic complexes (Groh and Rössler, 2011). To investigate the AL2/PL2 of *M. muscosa*, we

have employed the same double-labeling approach (Fig. 9d-e), but the anti-synapsin and
phalloidin labeled areas in each synaptic complex were not clearly differentiated, as is the
case in social hymenopterans (Groh and Rössler, 2011), despite co-localization being minimal.
This might indicate that phalloidin does not consistently label postsynaptic sites in *M.*
muscosa, or that the organization of the pre- and postsynaptic sites within the AL2/PL2 of *M.*
muscosa differs compared to the glomeruli in mushroom body calyces of investigated insect
species. To clarify synaptic connectivity within the AL2/PL2 in *M. muscosa* and spiders in
general, a detailed transmission electron microscopic study is needed.

4.5   Mushroom body pedunculus, haft and bridge

The somata within SC2 of the neurons that connect the second order visual neuropils to the
MBp have been referred to as "globuli cells" (Hanström, 1921; Steinhoff et al., 2017) and the
neurons including these somata as "T-cells" (Strausfeld and Barth, 1993). In *C. salei*, globuli
cells have a smaller diameter of the soma and nucleus than other cell bodies in the
protocerebrum (Babu and Barth, 1984). We could not find such size differences in *M.*
muscosa. In fact, all three nuclear markers revealed that somata of the three identified
clusters as well the somata of the surrounding cortex exhibit a similar size (Figs. 2; 3; 5; 6; 7).
Thus, according to their definition (Kenyon, 1896; Richter et al., 2010), somata of SC2 are no
globuli cells in *M. muscosa*. The long neurites that connect the SC2 with the AL2/PL2 and the
L2 are probably primary neurites (Richter et al., 2010), from which dendrites branch into the
AL2/PL2 and L2 respectively, and axons project into the MBp, MBh and MBbr. A similar
arrangement has been described by Babu and Barth (1984) for *C. salei* and by Hill (1975) for
two *Phidippus* species.

The structure and organization of the MB differs between *M. muscosa* and *C. salei*, as already
reported by Steinhoff et al. (2017). The MBh is a separate structure of the mushroom body in
M. muscosa, connected to the MBp by neurites (Steinhoff et al., 2017), whereas in *C. salei*, it
is fused with the MBp (Strausfeld and Barth, 1993). Furthermore, the mushroom body shaft,
which in *C. salei* gives rise to the MBbr (Strausfeld and Barth, 1993) is absent or fused with
the MBp in *M. muscosa.*

4.5 Functional considerations

The jumping spider *Marpissa muscosa* and the wandering spider *Cupiennius salei* differ
strongly in their lifestyle. While *M. muscosa* is a diurnal hunter that actively pursues its prey

and relies on vision for prey capture, *C. salei* is an exclusively nocturnal sit-and-wait predator,

which detects its prey based on vibratory cues (Barth & Seyfarth, 1979). Although attack

behavior of *C. salei* can be elicited by visual cues under experimental conditions (Fenk et al.,

2010), mechanosensing is the major modality for hunting prey, since *C. salei* also captures

prey when its eyes are occluded (Hergenröder and Barth, 1983). Given these obvious

differences in lifestyle and relative importance of sensory modalities, it is surprising how

similar the general brain anatomy of *M. muscosa* and *C. salei* is. Apart from differences in size

and structure of neuropils, the visual pathway of the principal eyes is identical in *C. salei* and

M. muscosa. Parts of the visual pathways of the secondary eyes in the jumping spider

Marpissa muscosa differ, however, from those in the wandering spider *Cupiennius salei.*

These differences might reflect the differences in the visual fields and corresponding

functions of their secondary eyes (Land, 1985b; Land and Barth, 1992). The most conspicuous

difference between the visual systems in the brains of *M. muscosa* and *C. salei* is the pathway

of the PME. While the two larger secondary eyes (ALE and PLE) are connected to the

mushroom bodies in *M. muscosa,* the PME are not (Fig. 11). This suggests that the PME do

not play a role in motion detection (Duelli, 1978; Land, 1971; Zurek et al., 2010). In *C. salei,*

however, the PME are important movement detectors (Schmid, 1998) connected to the

mushroom bodies (Strausfeld and Barth, 1993).

Another interesting difference between the two species is the presence of the L2 in *M.*

muscosa (Steinhoff et al., 2017), which is absent in *C. salei*. The structure of this neuropil

differs from the AL2/PL2 and all second order visual neuropils in *C. salei,* since the L2 in *M.*

muscosa lacks a glomerular organization. Glomeruli might ensure a retinotopic organization

of neurites within the second order visual neuropils. This has been hypothesized for both *C.*

salei (Strausfeld and Barth, 1993) and for the PLE pathway of jumping spiders (Duelli, 1980),

where no lateral connections between individual glomeruli of the AL2/PL2 were found. We

hypothesize that while a retinotopic organization is kept in the AL2/PL2, it is lost in the L2.

Information from the movement-detecting ALE and PLE in *M. muscosa* might be integrated

not only in the MB, but also in the L2, and enable faster movement decisions. Alternatively, a

glomerular organization might be the most space-saving structure for the processing of input

from differentially tuned groups of axons (Eisthen, 2002; Hildebrand and Shepherd, 1997),

and although the L2 is not composed of glomeruli, information might still be processed in

parallel without lateral connections. This would also indicate that a higher number of neurites

synapse in the AL2/PL2 than in the L2 in *M. muscosa*.

Some early-branched off salticid species possess a large PME that has its own field of view

(e.g. *Portia*; Land, 1985b; Maddison and Hedin, 2003). Land (1985b) suggested that in jumping

spiders with a small PME, such as *M. muscosa*, the fields of view of ALE and PLE have expanded

to fill the space not covered by the PME. The evolutionary transformation to a smaller PME

and larger ALE and PLE might have led to visual pathways that also incorporated the L2. Thus

in early-branched off jumping spiders with large PME, it could be hypothesized that the L2

might serve as a second order visual neuropil of the PME. Future studies should thus analyze

the brain architecture of early-branched off salticid species.

CONFLICT OF INTEREST STATEMENT

The authors have no conflicts of interests.

AUTHOR CONTRIBUTIONS

All authors had full access to all the data in the study and take responsibility for the integrity of the

data and the accuracy of data analysis. Study concept and design: POMS, AS, GU. Acquisition of data:

POMS. Analysis and interpretation of data: POMS, AS. Wrote the manuscript: POMS, AS. Contributed

to the writing of the manuscript: GU, SH. Obtained funding: GU, SH, POMS.

LITERATURE CITED

Babu, K.S., Barth, F.G., 1984. Neuroanatomy of the central nervous system of the wandering

566   spider, *Cupiennius salei* (Arachnida, Araneida). Zoomorphology 104, 344–359.

567   https://doi.org/10.1007/BF00312185

568 Barth, F., Seyfarth, E.-A., 1979. Cupiennius salei Keys. (Araneae) in the highlands of Central

569   Guatemala. J. Arachnol. 7, 255–263.

570 Barth, F.G., 2002. The central nervous system, in: Barth, Friedrich G. (Ed.), A Spiders World;

571   Senses and Behavior. Springer, Berlin Heidelberg, pp. 161–174.

572 Barth, F.G., Nakagawa, T., Eguchi, E., 1993. Vision in the ctenid spider *Cupiennius salei*:

573   spectral range and absolute sensitivity. J. Exp. Biol. 79, 63–79. https://doi.org/ghjjagsj

574 Battelle, B.A., Calman, B.G., Andrews, A.W., Grieco, F.D., Mleziva, M.B., Callaway, J.C.,

575   Stuart, A.E., 1991. Histamine: A putative afferent neurotransmitter in *Limulus* eyes. J.

576   Comp. Neurol. 305, 527–542. https://doi.org/10.1002/cne.903050402

577 Battelle, B.A., Sombke, A., Harzsch, S., 2016. Xiphosura, in: Schmidt-Rhaesa, A., Harzsch,

578   Steffen, Purschke, G. (Eds.), Structure and Evolution of Invertebrate Nervous Systems.

579   Oxford University Press, pp. 428–442.

Becherer, C., Schmid, A., 1999. Distribution of g-aminobutyric acid-, proctolin-, *Periplaneta*

hypertrehalosaemic hormone- and FMRFamide-like immunoreactivity in the visual

ganglia of the spider *Cupiennius salei* Keys. Comp. Biochem. Physiol. Part A 122, 267–

275.

Cebrià, F., 2008. Organization of the nervous system in the model planarian *Schmidtea*

mediterranea: An immunocytochemical study. Neurosci. Res. 61, 375–384.

https://doi.org/10.1016/j.neures.2008.04.005

Duelli, P., 1980. The neuronal organization of the posterior lateral eyes of jumping spiders

(Salticidae). Zool. Jahrbücher Abteilung für Anat. und Ontog. der Tiere 103, 17–40.

Duelli, P., 1978. Movement detection in the posterolateral eyes of jumping spiders (*Evarcha*

arcuata, Salticidae). J. Comp. Physiol. A 124, 15–26.

https://doi.org/10.1007/BF00656387

Eisthen, H.L., 2002. Why are olfactory systems of different animals so similar? Brain. Behav.

Evol. 59, 273–293. https://doi.org/10.1159/000063564

Fabian-Fine, R., Volknandt, W., Seyfarth, E.-A., 1999. Peripheral synapses at identifiable

mechanosensory neurons in the spider *Cupiennius salei*: synapsin-like

immunoreactivity. Cell Tissue Res. 295, 13–19.

Fenk, L.M., Hoinkes, T., Schmid, A., 2010. Vision as a third sensory modality to elicit attack

behavior in a nocturnal spider. J. Comp. Physiol. A 196, 957–961.

https://doi.org/10.1007/s00359-010-0575-8

Fenk, L.M., Schmid, A., 2010. The orientation-dependent visual spatial cut-off frequency in a

spider. J. Exp. Biol. 213, 3111–7. https://doi.org/10.1242/jeb.041939

Foelix, R.F., 1996. Biology of Spiders. Oxford University Press.

https://doi.org/10.2307/1535947

Garm, A., O'Connor, M., Parkefelt, L., Nilsson, D.-E., 2007. Visually guided obstacle

avoidance in the box jellyfish *Tripedalia cystophora* and *Chiropsella bronzie*. J. Exp. Biol.

210, 3616–3623. https://doi.org/10.1242/jeb.004044

Gregory, G.E., 1980. The bodian protargol technique, in: Strausfeld, N.J., Miller, T.A. (Eds.),

Neuroanatomical Techniques Insect Nervous System. Springer Berlin Heidelberg, pp.

75–95.

Groh, C., Rössler, W., 2011. Comparison of microglomerular structures in the mushroom

body calyx of neopteran insects. Arthropod Struct. Dev. 40, 358–367.

https://doi.org/10.1016/j.asd.2010.12.002

Hanström, B., 1921. Über die Histologie und vergleichende Anatomie der Sehganglien und

Globuli der Araneen. K. Sven. Vetenskapsakademiens Handl. 61, 1–39.

Harland, D.P., Li, D., Jackson, R.R., 2012. How jumping spiders see the world, in: Lazareva,

O.F., Shimizu, T., Wasserman, E.A. (Eds.), How Animals See the World: Comparative

Behavior, Biology, and Evolution of Vision. Oxford University Press, pp. 133–164.

https://doi.org/10.1093/acprof:oso/9780195334654.003.0010

Harzsch, S., Hansson, B.S., 2008. Brain architecture in the terrestrial hermit crab *Coenobita*

clypeatus (Anomura, Coenobitidae), a crustacean with a good aerial sense of smell.

BMC Neurosci. 9, 1–35. https://doi.org/10.1186/1471-2202-9-58

Harzsch, S., Müller, C.H., 2007. A new look at the ventral nerve centre of *Sagitta*:

implications for the phylogenetic position of Chaetognatha (arrow worms) and the

evolution of the bilaterian nervous system. Front. Zool. 4, 14.

Harzsch, S., Vilpoux, K., Blackburn, D.C., Platchetzki, D., Brown, N.L., Melzer, R., Kempler,

K.E., Battelle, B.A., 2006. Evolution of arthropod visual systems: Development of the

eyes and central visual pathways in the horseshoe crab *Limulus polyphemus* Linnaeus,

1758 (Chelicerata, Xiphosura). Dev. Dyn. 235, 2641–2655.

https://doi.org/10.1002/dvdy.20866

Harzsch, S., Wildt, M., Battelle, B., Waloszek, D., 2005. Immunohistochemical localization of

neurotransmitters in the nervous system of larval *Limulus polyphemus* (Chelicerata,

Xiphosura): Evidence for a conserved protocerebral architecture in Euarthropoda.

Arthropod Struct. Dev. 34, 327–342. https://doi.org/10.1016/j.asd.2005.01.006

Hergenröder, R., Barth, F.G., 1983. Vibratory signals and spider behavior: How do the

sensory inputs from the eight legs interact in orientation? J. Comp. Physiol. A 152, 361–

371. https://doi.org/10.1007/BF00606241

Hildebrand, J.G., Shepherd, G.M., 1997. Mechanisms of olfactory discrimination: Converging

evidence for common principles across phyla. Annu. Rev. Neurosci. 20, 595–631.

https://doi.org/10.1146/annurev.neuro.20.1.595

Hill, D.E., 1975. The structure of the central nervous system of jumping spiders of the genus

Phidippus (Araneae: Salticidae). M.Sc. thesis; Oregon State University.

Homann, H., 1952. Die Nebenaugen der Araneen. Zweite Mitteilung. Zoologische Jahrbücher

Abteilung für Anatomie und Ontogenie der Tiere 72, 345–364.

Homann, H., 1950. Die Nebenaugen der Araneen. Zoologische Jahrbücher Abteilung für

Anatomie und Ontogenie der Tiere 71, 56–144.

Jakob, E.M., Long, S.M., Harland, D.P., Jackson, R.R., Carey, A., Searles, M.E., Porter, A.H.,
Canavesi, C., Rolland, J.P., 2018. Lateral eyes direct principal eyes as jumping spiders
track objects. Curr. Biol. 28, R1092–R1093. https://doi.org/10.1016/j.cub.2018.07.065

Kenyon, F.C., 1896. The meaning and structure of the so-called "mushroom bodies" of the
hexapod brain. Am. Nat. 643–650.

Klagges, B.R.E., Heimbeck, G., Godenschwege, T.A., Hofbauer, A., Pflugfelder, G.O.,
Reifegerste, R., Reisch, D., Schaupp, M., Buchner, S., Buchner, E., 1996. Invertebrate
synapsins: a single gene codes for several isoforms in *Drosophila*. J. Neurosci. 16, 3145–
3165.

Land, M.F., 1985a. The morphology and optics of spider eyes, in: Barth, F.G. (Ed.),
Neurobiology of Arachnids. Springer, Berlin, pp. 53–76. https://doi.org/10.1007/978-3-
642-70348-5_4

Land, M.F., 1985b. Fields of view of the eyes of primitive jumping spiders. J. Exp. Biol. 119,
381–384.

Land, M.F., 1972. Mechanisms of orientation and pattern recognition by jumping spiders
(Salticidae), in: Wehner, R. (Ed.), Information Processing in the Visual Systems of
Anthropods. Springer Berlin Heidelberg, pp. 231–247.

Land, M.F., 1971. Orientation by jumping spiders in the absence of visual feedback. J. Exp.
Biol. 54, 119–139.

Land, M.F., Barth, F.G., 1992. The quality of vision in the ctenid spider *Cupiennius salei*. J.
Exp. Biol. 164, 227–242.

Lehmann, T., Melzer, R.R., 2018. Also looking like *Limulus*? - retinula axons and visual
neuropils of Amblypygi (whip spiders). Front. Zool. 15, 52.
https://doi.org/10.1186/s12983-018-0293-6

Lehmann, T., Melzer, R.R., 2013. Looking like *Limulus*? - Retinula axons and visual neuropils
of the median and lateral eyes of scorpions. Front. Zool. 10, 40.
https://doi.org/10.1186/1742-9994-10-40

Lehmann, T., Melzer, R.R., Hörnig, M.K., Michalik, P., Sombke, A., Harzsch, S., 2016.
Arachnida (excluding scorpiones), in: Schmidt-Rhaesa, A., Harzsch, S., Purschke, G.
(Eds.), Structure and Evolution of Invertebrate Nervous Systems. Oxford University
Press, Oxford, pp. 453–477.

Loesel, R., Wolf, H., Kenning, M., Harzsch, S., Sombke, A., 2013. Architectural principles and
evolution of the arthropod central nervous system, in: Minelli, A. (Ed.), Arthropod

Biology and Evolution. Springer-Verlag, Berlin Heidelberg, pp. 299–342.

https://doi.org/10.1007/978-3-642-36160-9

Long, S.M., 2016. Spider Brain Morphology & Behavior. PhD thesis; University of

Massachusetts, Amherst.

Maddison, W.P., Hedin, M.C., 2003. Jumping spider phylogeny (Araneae: Salticidae).

Invertebr. Syst. 17, 529. https://doi.org/10.1071/is02044

Morehouse, N.I., Buschbeck, E.K., Zurek, D.B., Steck, M., Porter, M.L., 2017. Molecular

evolution of spider vision: New opportunities, familiar players. Biol. Bull. 233, 21–38.

https://doi.org/10.1086/693977

Nagata, T., Arikawa, K., Kinoshita, M., 2019. Photoreceptor projection from a four-tiered

retina to four distinct regions of the first optic ganglion in a jumping spider. J. Comp.

Neurol. 1–14. https://doi.org/10.1002/cne.24584

Nässel, D.R., 1999. Histamine in the brain of insects: a review. Microsc. Res. Tech. 44, 121–

136. https://doi.org/10.1002/(SICI)1097-0029(19990115/01)44:2/3<121::AID-

JEMT6>3.0.CO;2-F

O'Connor, M., Garm, A., Nilsson, D.E., 2009. Structure and optics of the eyes of the box

jellyfish *Chiropsella bronzie*. J. Comp. Physiol. A 195, 557–569.

https://doi.org/10.1007/s00359-009-0431-x

Orlando, E., Schmid, A., 2011. Colour blindness of the movement-detecting system of the

spider *Cupiennius salei*. J. Exp. Biol. 214, 546–550. https://doi.org/10.1242/jeb.051672

Ott, S.R., 2008. Confocal microscopy in large insect brains: Zinc-formaldehyde fixation

improves synapsin immunostaining and preservation of morphology in whole-mounts.

J. Neurosci. Methods 172, 220–230. https://doi.org/10.1016/j.jneumeth.2008.04.031

Paulus, H., 1979. Eye structure and the monophyly of the Arthropoda, in: AP, G. (Ed.),

Arthropod Phylogeny. Vam Nostrand Reinhold Company, New York., pp. 299-383.

Richter, S., Loesel, R., Purschke, G., Schmidt-Rhaesa, A., Scholtz, G., Stach, T., Vogt, L.,

Wanninger, A., Brenneis, G., Döring, C., Faller, S., Fritsch, M., Grobe, P., Heuer, C.M.,

Kaul, S., Møller, O.S., Müller, C.H., Rieger, V., Rothe, B.H., Stegner, M.E., Harzsch, S.,

2010. Invertebrate neurophylogeny: suggested terms and definitions for a

neuroanatomical glossary. Front. Zool. 7, 29. https://doi.org/10.1186/1742-9994-7-29

Saint Rémy, G., 1887. Contribution à l'étude du cerveau chez les Arthropodes trachéates.

Arch. Zool. Exp. Générale.

Schmid, A., 1998. Different functions of different eye types in the spider *Cupiennius salei*. J.

Exp. Biol. 201, 221–5.

Schmid, A., Duncker, M., 1993. Histamine immunoreactivity in the central nervous system of

the spider *Cupiennius salei*. Cell Tissue Res. 273, 533–545.

https://doi.org/10.1007/BF00333707

Schulze, E., Graupner, H., 1960. Anleitung zum mikroskopisch-technischen Arbeiten in

Biologie und Medizin. Akademische Verlagsgesellschaft.

Sombke, A., Harzsch, S., 2015. Immunolocalization of histamine in the optic neuropils of

Scutigera coleoptrata (Myriapoda: Chilopoda) reveals the basal organization of visual

systems in Mandibulata. Neurosci. Lett. 594, 111–6.

https://doi.org/10.1016/j.neulet.2015.03.029

Sombke, A., Harzsch, S., Hansson, B.S., 2011. Organization of deutocerebral neuropils and

olfactory behavior in the centipede *Scutigera coleoptrata* (Linnaeus, 1758) (Myriapoda:

Chilopoda). Chem. Senses 36, 43–61. https://doi.org/10.1093/chemse/bjq096

Sombke, A., Lipke, E., Michalik, P., Uhl, G., Harzsch, S., 2015. Potential and limitations of x-

ray micro-computed tomography in arthropod neuroanatomy: a methodological and

comparative survey. J. Comp. Neurol. 523, 1281–1295.

https://doi.org/10.1002/cne.23741

Spano, L., Long, S.M., Jakob, E.M., 2012. Secondary eyes mediate the response to looming

objects in jumping spiders (*Phidippus audax*, Salticidae). Biol. Lett. 8, 949–951.

https://doi.org/10.1098/rsbl.2012.0716

Steinhoff, P.O.M., Liedtke, J., Sombke, A., Schneider, J.M., Uhl, G., 2018. Early environmental

conditions affect the volume of higher-order brain centers in a jumping spider. J. Zool.

304, 182–192. https://doi.org/10.1111/jzo.12512

Steinhoff, P.O.M., Sombke, A., Liedtke, J., Schneider, J.M., Harzsch, S., Uhl, G., 2017. The

synganglion of the jumping spider *Marpissa muscosa* (Arachnida: Salticidae): Insights

from histology, immunohistochemistry and microCT analysis. Arthropod Struct. Dev.

46, 156–170. https://doi.org/10.1016/j.asd.2016.11.003

Strausfeld, N.J., 2012. Arthropod Brains: Evolution, Functional Elegance, and Historical

Significance, Cambridge: Harvard UP. Belknap Press of Harvard University Press,

Cambridge.

Strausfeld, N.J., Barth, F.G., 1993. Two visual systems in one brain: Neuropils serving the

secondary eyes of the spider *Cupiennius salei*. J. Comp. Neurol. 328, 43–62.

https://doi.org/10.1002/cne.903280105

Strausfeld, N.J., Weltzien, P., Barth, F.G., 1993. Two visual systems in one brain: Neuropils

serving the principal eyes of the spider *Cupiennius salei*. J. Comp. Neurol. 328, 63–75.

https://doi.org/10.1002/cne.903280105

Sullivan, J.M., Benton, J.L., Sandeman, D.C., Beltz, B.S., 2007. Adult neurogenesis: A common

strategy across diverse species. J. Comp. Neurol. 500, 574–584.

https://doi.org/10.1002/cne.21187

Weltzien, P., Barth, F.G., 1991. Volumetric measurements do not demonstrate that the

spider brain "central body" has a special role in web building. J. Morphol. 208, 91–98.

https://doi.org/10.1002/jmor.1052080104

Widmer, A., Höger, U., Meisner, S., French, A.S., Torkkeli, P.H., 2005. Spider peripheral

mechanosensory neurons are directly innervated and modulated by octopaminergic

efferents. J. Neurosci. 25, 1588–1598. https://doi.org/10.1523/jneurosci.4505-04.2005

Yamashita, S., Tateda, H., 1978. Spectral sensitivities of the anterior median eyes of the orb

web spiders, *Argiope bruennichii* and *A. amoena*. J. Exp. Biol. 74, 47–57.

Yamashita, S., Tateda, H., 1976. Spectral sensitivities of jumping spider eyes. J. Comp.

Physiol. A 105, 29–41. https://doi.org/10.1007/BF01380051

Zurek, D.B., Nelson, X.J., 2012a. Hyperacute motion detection by the lateral eyes of jumping

spiders. Vision Res. 66, 26–30. https://doi.org/10.1016/j.visres.2012.06.011

Zurek, D.B., Nelson, X.J., 2012b. Saccadic tracking of targets mediated by the anterior-lateral

eyes of jumping spiders. J. Comp. Physiol. A 198, 411–417.

https://doi.org/10.1007/s00359-012-0719-0

Zurek, D.B., Taylor, A.J., Evans, C.S., Nelson, X.J., 2010. The role of the anterior lateral eyes in

the vision-based behaviour of jumping spiders. J. Exp. Biol. 213, 2372–8.

https://doi.org/10.1242/jeb.042382

770
771
772

775 **Tables**

776 **Table 1** Primary and secondary antibodies

Labeling reagent	Dilution and specifications
Primary	

Monoclonal anti-synapsin antibody produced in mouse (RRID AB_528479)	1:1000, DSHB 3C11 Steinhoff et al. (2017), this study
Polyclonal anti α-tubulin antibody produced in rabbit (RRID AB_2637882)	1:1000 or 1:250, ThermoFisher PA5-58711
Monoclonal anti-acetylated α-tubulin antibody produced in mouse (RRID AB_477585)	1:1000, Sigma Aldrich T6793
Monoclonal anti-histamine antibody produced in rabbit (RRID AB_1541510)	1:1000, Progen 16043 Harzsch et al. (2005), Sombke and Harzsch (2015)
Secondary	
Polyclonal Cy3 anti-mouse IgG secondary anti-body produced in goat (RRID AB_2338680)	1:500, Jackson Immuno Research, 115–165-003
Polyclonal Alexa 488 anti-rabbit IgG secondary antibody produced in goat (RRID AB_2534074)	1:500, Invitrogen, A11006
Alexa Fluor 488 phalloidin (RRID AB_2315147)	1:50, Molecular Probes A-12379
Hoechst 33258	1:1000, Sigma 14530
YOYO 1 491	Molecular Probes Y-3601

778 **Table 2** Methods and antisera combinations

Methods	Antisera combinations/ staining	Coupled with	Specimens
Wholemount	anti-synapsin	Cy3 anti-mouse (goat)	5
Vibratome sections (50 µm; 80 µm)	anti-synapsin (mouse) anti- α-tubulin (rabbit)	Cy3 anti-mouse (goat) Alexa Fluor 488 anti-rabbit (goat) Nuclear labeling: Hoechst 33258	16
	anti-acetylated-α-tubulin (mouse)	Nuclear labeling: YOYO	6
	anti-histamine (rabbit)	Alexa Fluor 488 anti-rabbit (goat)	4
	anti-synapsin (mouse)	Cy3 anti-mouse (goat) Phalloidin	5
Paraffin sections (5 µm)	Azan-novum	/	2
microCT scans	1% idodine (in 99% ethanol)	/	4

780
781

Figures

Figure 1. The principal eye pathway in *Marpissa muscosa*. **a** Virtual horizontal section of the

dorsal protocerebrum obtained by microCT analysis shows the connections between the

AM1, AM2 and the AB. **b** Three-dimensional reconstruction based on a microCT image stack

shows the same connections as in a. **c-f** Maximum projection of image stacks (clsm), show

tubulin-immunoreactivity (grey) and somata (blue) in the protocerebrum of *Marpissa*

muscosa. **c** Horizontal section showing the AM2, the AB and connecting neurites (arrows). **d**

Sagittal section through the AM1, AM2 and the AB showing the connecting neurites between

AM2 and AB (arrow). **e** Sagittal section showing at least two lobes of the AM1, the thick tract

to the AM2 and the thinner connection to the AB (arrow). AL2/PL2 and MB are also visible. **f**

Horizontal section showing two lobes of the AM1, retinula cell axons terminating in those and

interneurons connecting AM1 with AM2. **Abbreviations: a** anterior, **AB** arcuate body,

AL2/PL2 second order visual neuropil of anterior lateral and posterior lateral eyes, **AM1** first

order visual neuropil of the anterior median eyes, **AM2** second order visual neuropil of the

anterior median eyes, **AMEr** retina of the anterior median eyes, **ChN** cheliceral neuropil, **d**

dorsal, **l** lateral, **MB** mushroom body, **PL1** first order visual neuropil of the posterior lateral

eyes, **PLEr** retina of the posterior lateral eyes, **PM1** first order visual neuropil of the posterior

median eyes.

Figure 2. Maximum projection of an image stack (clsm), showing tubulin-immunoreactivity (grey) and somata (blue) in the protocerebrum of *Marpissa muscosa*. The AL1 and PL1 are connected to both, the AL2/PL2 and the L2. The neurites forming these connections originate in SC1. The AL2/PL2 and the L2 are connected to the MBp via neurites, which originate in the SC2. The two MBp are connected with each other via the mushroom body bridge, which is formed by long neurites that project along the periphery of the MBp and seem to originate in the SC2. Some of these long neurites also connect the MBp with the MBh. Other neurites originate in SC3 and enter the protocerebrum medially. **Abbreviations: a** anterior, **AL1** first order visual neuropil of the anterior lateral eyes, **AL2/PL2** second order visual neuropil of ALE and PLE, **d** dorsal, **l** lateral, **L2** shared second order visual neuropil of the anterior lateral and posterior lateral eyes, **MBh** mushroom body haft, **MBp** mushroom body pedunculus, **SC1-3** soma cluster 1-3.

Figure 3. Paraffin sections through the anterior protocerebrum of *Marpissa muscosa.* **a**

Sagittal section showing the conspicuous tracts that connect the AL1 and PL1 with the L2. **b**

Horizontal section showing the convoluted and columnar structure of the AL1. Neurites that

connect the AL1 with the AB probably synapse in the AL1x. Long neurites connecting the PL1

with the AB are visible curving around L2. Other long neurites contributing to the tract to the

AB synapse in PM1. Arrows point to neurites contributing to the tract between first order

visual neuropils of the secondary eyes and the arcuate body. **c** Horizontal section showing

several neurites that connect the PL1 with the AL2/PL2, as well as the neurites contributing

to the tract between the PL1 and the AB (highlighted by arrows). **Abbreviations: a** anterior,

AB arcuate body, **AL1** first order visual neuropil of the anterior lateral eyes, **AL1x** neuropilar

subunit of the AL1, **AL2/PL2** second order visual neuropil of anterior lateral and posterior

lateral eyes, **ALEr** retina of the anterior lateral eyes, **AMEr** retina of the anterior median eyes,

d dorsal, **l** lateral, **L2** shared second order visual neuropil of the anterior lateral and posterior

lateral eyes, **PL1** first order visual neuropil of the posterior lateral eyes, **PLEr** retina of the

anterior lateral eyes, **PL1x** neuropilar subunit of the PL1, **PM1** first order visual neuropil of the

posterior median eyes**.**

Figure 4. Protocerebral neuropils depicted in different virtual horizontal sections obtained by

microCT analysis (**a, c, e**) and three-dimensional reconstructions based on microCT image

stacks (**b, d, f**). **a + b** The retina of the ALE is located very closely to its first order visual

neuropil, the AL1. The AL1 has a convoluted surface and is connected to the AL2/PL2 and the

L2. **c + d** The retina of the PLE almost comes in contact with its first order visual neuropil, the

PL1. The PL1 has a convoluted surface and is connected to the MB and the L2. **e + f** A

comparatively long optic nerve connects the retina of the PME with its first order visual

neuropil, the PM1. See supporting information for an interactive 3D-visualization of the

microCT reconstruction. **Abbreviations: a** anterior, **AL1** first order visual neuropil of the

anterior lateral eyes, **AL2/PL2** second order visual neuropil of anterior lateral and posterior

lateral eyes, **ALEr** retina of the anterior lateral eyes, **AMEr** retina of the anterior median eyes,

d dorsal, **l** lateral, **L2** shared second order visual neuropil of the anterior lateral and posterior

lateral eyes, **MBbr** mushroom body bridge, **MBh** mushroom body haft, **MBp** mushroom body

pedunculus, **PL1** first order visual neuropil of the posterior lateral eyes, **PLEr** retina of the

posterior lateral eyes, **PM1** first order visual neuropil of the posterior median eyes, **PMEr**

retina of the posterior median eyes.

Figure 5. Maximum projections of image stacks (clsm) showing tubulin-immunoreactivity
(grey), synapsin-immunoreactivity (magenta) and somata (blue) in the secondary eye visual
pathway of *Marpissa muscosa*. **a** Neurites from the PL1 pass between AL2/PL2 and L2 towards
the AB. **b** Neurites from AL1x join the tract from the PM1 that passes through between
AL2/PL2 and L2 (highlighted by arrows). **c** The PL1 is in close spatial relationship to the PLEr.
Parallel neurites connect the PL1 and PL1x with the Al2/PL2. **Abbreviations: a** anterior, **AB**
arcuate body, **AL1** first order visual neuropil of the anterior lateral eyes, **AL1x** neuropilar
subunit of the AL1, **AL2/PL2** second order visual neuropil of anterior lateral and posterior
lateral eyes, **AM2** second order visual neuropil of the anterior median eyes, **d** dorsal, **l** lateral,
L2 shared second order visual neuropil of the anterior lateral and posterior lateral eyes, **MBp**
mushroom body pedunculus, **PL1** first order visual neuropil of the posterior lateral eyes, **PL1x**

neuropilar subunit of the PL1, **PLEr** retina of the posterior lateral eyes, **PM1** first order visual

neuropil of the posterior median eyes.

Figure 6. Maximum projections of image stacks (clsm) showing tubulin-immunoreactivity (grey), synapsin-immunoreactivity (magenta) and somata (blue) in the secondary eye visual pathway of *Marpissa muscosa*. **a** The AL1 is connected to the AL2/PL2 and the L2 via two prominent neurite tracts. Some of the neurites contributing to these tracts originate in SC1. Neurites connecting the AL2/PL2 with the MBp originate in SC2. **b** The PL1 is connected to the AL2/PL2 and the L2 via two prominent neurite tracts. Some neurites that arise in SC2 pass between AL2/PL2 and MBp towards the L2. **c** The optic nerve of the PME (highlighted by arrow) terminates in the PM1. **c'** Higher magnification of C, showing the termination site of the PME optic nerve in the PM1. **d** Neurites from the PM1 pass between AL2/PL2 and L2. **Abbreviations: a** anterior, **AL1** first order visual neuropil of the anterior lateral eyes, **AL2/PL2**

second order visual neuropil of anterior lateral and posterior lateral eyes, **d** dorsal, **l** lateral, **L2** shared second order visual neuropil of the anterior lateral and posterior lateral eyes, **MBp** mushroom body pedunculus, **PL1** first order visual neuropil of the posterior lateral eyes, **PL1x** neuropilar subunit of the PL1, **PM1** first order visual neuropil of the posterior median eyes, **SC1-2** soma cluster 1-2.

Figure 7. Maximum projections of image stacks (clsm) showing tubulin-immunoreactivity
(grey), synapsin-immunoreactivity (magenta) and somata (blue) in the secondary eye visual
pathway of *Marpissa muscosa*. **a** Horizontal section through AL1, MBp and AL2/PL2. The AL1
connects to one of the two tightly adjoining clusters of AL2/PL2. Arrows point to primary
neurites connecting SC2 and AL2/PL2. Arrowheads point to neurites connecting the AL2/PL2
with the MBp. **b** Horizontal section through visual neuropils. The PM1 is situated between

AL1 and PL1, AL2/PL2 appear as a uniform array. The arrowhead points to neurites connecting PL1 with AB, neurites probably synapse in the PL1x. **c** A sagittal section through AL2/PL2 and MBp reveals two distinct clusters of AL2/PL2 and the chiasm formed by neurites that connect AL2/PL2 with MBp. **d** Tubulin-immunoreactivity shows long peripheral neurites contributing to the MBbr and connecting the MBp with the MBh. Arrows point to primary neurites connecting SC2 with the AL2/PL2. Arrowheads point to neurites connecting the MBp with the MBh. **Abbreviations: a** anterior, **AL1** first order visual neuropil of the anterior lateral eyes, **AL1x** neuropilar subunit of the AL1, **AL2/PL2** second order visual neuropil of anterior lateral and posterior lateral eyes, **d** dorsal, **l** lateral, **L2** shared second order visual neuropil of the anterior lateral and posterior lateral eyes, **MBbr** mushroom body bridge, **MBh** mushroom body haft, **MBp** mushroom body pedunculus, **PL1** first order visual neuropil of the posterior lateral eyes, **PL1x** neuropilar subunit of the PL1, **PM1** first order visual neuropil of the posterior median eyes, **SC1-3** soma cluster 1-3.

Figure 8. Maximum projections of image stacks (clsm), showing histamine-immunoreactivity in horizontal sections through the anteriormost protocerebrum of *Marpissa muscosa*. **a** Overview, shows strong histamine-immunoreactivity of retinula cell axon terminals in the AL1 and its substructure the AL1x, as well as in the PL1 and its substructure the PL1x. Retinula cell axons of the PME and their terminals in the PM1 exhibit strong histamine-immunoreactivity. **b** Magnification showing histamine-immunoreactive retinula cell axons and their terminals in the AL1x and PM1. Note the close spatial association of AL1x and PM1; AL1x is somewhat distorted due to sectioning. Arrows point to histamine-immunoreactive retinula cell axons and their terminals. **c** Magnification showing histamine-immunoreactive retinula cell axons and their terminals in the PL1 and PL1x. Arrows point to a retinula cell axon and a large terminal in the PL1x. **Abbreviations: a** anterior, **AL1** first order visual neuropil of the anterior lateral eyes, **AL1x** neuropilar subunit of the AL1, **d** dorsal, **l** lateral, **PL1** first order visual neuropil of the posterior lateral eyes, **PL1x** neuropilar subunit of the PL1, **PM1** first order visual neuropil of the posterior median eyes.

1059

Figure 9. The second order visual neuropil of anterior lateral and posterior lateral eyes in the
secondary eye visual pathway of *Marpissa muscosa.* **a** Three-dimensional volume rendering
of an anti-synapsin labeled whole-mount, view from posterodorsal. Note the two distinct
clusters of glomeruli that adjoin closely and coat the MBp anteriorly. **b** Single optical section
of the AL2/PL2 within an anti-synapsin labeled whole-mount shows two distinct clusters of
AL2/PL2 and a chiasm of neurites connecting them to the MBp. **c** Stack of optical sections

through the AL2/PL2 (maximum intensity projection), showing synapsin-immunoreactivity. The glomerular subunits within each individual AL2/PL2 may represent synaptic boutons. **d** Anti-synapsin (magenta) and phalloidin (green) double–labeling reveals glomerular substructures (boutons) of the AL2/PL2. **e** Three individual AL2/PL2 with glomerular substructures (boutons) labeled by anti-synapsin and phalloidin.

1099

Figure 10. Maximum projections of image stacks (clsm; **a,c-e**) and single optical section (clsm) showing synapsin-immunoreactivity (magenta), tubulin-immunoreactivity (grey) and somata (cyan) in the secondary eye visual system of *Cupiennius salei*. **a** In the principal eye visual pathway, the AM1 is connected to the AM2 via a thick tract. Connecting neurites between AM2 and the AB are also visible (arrow). **b** Within the secondary eye visual pathways, every eye is associated with its own first order visual neuropil (AL1, PM1 and PL2); the PM1 and PL1 appear to have a substructure (PM1x and PL1x). All three second order visual neuropils are sub-structured into glomeruli. **c** The AL2, PM2 and PL2 are all connected to the MB. **d** The MB of *C. salei* consists of a MBp, MBh and MBs. All three substructures form a continuous synapsin-rich neuropil. **e** Horizontal section showing the PM1 and the PM1x. **Abbreviations: a** anterior, **AB** arcuate body, **AL1** first order visual neuropil of the anterior lateral eyes, **AL2** second order visual neuropil of the anterior lateral eyes, **AM1** first order visual neuropil of the anterior median eyes, **AM2** second order visual neuropil of the anterior median eyes, **d** dorsal,

1113   **l** lateral, **MBh** mushroom body haft, **MBp** mushroom body pedunculus, **MBs** mushroom body

1114   shaft, **PL1** first order visual neuropil of the posterior lateral eyes, **PL1x** neuropilar subunit of

1115   the PL1, **PL2** second order visual neuropil of the posterior lateral eyes, **PM1** first order visual

1116   neuropil of the posterior median eyes, **PM1x** neuropilar subunit of the PM1, **PM2** second

1117   order visual neuropil of the posterior median eyes.

1118

1119

1120

1121

1122

1123

1124

1125

1126

1127

1128

1129

1130

1131

1132

1133

1134

1135

1136

1137

1138

1139

1140

1141

1142

1143

Figure 11. Schematic representations of the principal and secondary eye visual pathways in the brains of **a** *Marpissa muscosa* and **b** *Cupiennius salei.* Insets top left: 3D volume rendering of the prosomata of **a** *M. muscosa* and **b** *C. salei*, based on microCT analysis. **b** Left anterior median eye concealed by posterior median eye. **Abbreviations: AB** arcuate body, **AL1** first order visual neuropil of the anterior lateral eyes, **AL2** second order visual neuropil of the

anterior lateral eyes, **AL2/PL2** second order visual neuropil of anterior lateral and posterior

lateral eyes, **ALE** anterior lateral eye, **AME** anterior median eye, **L2** shared second order visual

neuropil of the anterior lateral and posterior lateral eyes, **MBbr** mushroom body bridge, **MBh**

mushroom body haft, **MBp** mushroom body pedunculus, **PL1** first order visual neuropil of the

posterior lateral eyes, **PL2** second order visual neuropil of the posterior lateral eyes, **PLE**

posterior lateral eye, **PME** posterior median eye, **PM1** first order visual neuropil of the

posterior median eyes, **PM2** second order visual neuropil of the posterior median eyes. **Filled**

black circles indicate somata with known positions, **dotted lines** represent primary neurites

and **arrows** indicate the assumed direction of information flow. **Asterisks** mark neuropilar

subregions of first order neuropils (AL1x and PL1x in *M. muscosa*, PM1x and PL1x in *C. salei*).