A cytomegalovirus immunevasin triggers integrated stress response-dependent

reorganization of the endoplasmic reticulum.

Running Title: HCMV UL148 reorganizes the ER.

Hongbo Zhang[a,1], Clarissa Read[d,e,1], Christopher C. Nguyen[a] , Mohammed N.A.

Siddiquey[a], Chaowei Shang[c], Cameron M. Hall[a], Jens von Einem[d], and Jeremy P. Kamil[a,b #]

[a]Department of Microbiology and Immunology,

[b]Center for Molecular and Tumor Virology

[c]Research Core Facility

LSU Health Sciences Center, Shreveport, Louisiana, USA

[d] Institute of Virology, Ulm University Medical Center,

[e]Central Facility for Electron Microscopy, Ulm University

Ulm, Germany

[1] these authors contributed equally to the study

Running Head: HCMV UL148 Remodels the ER

#Address correspondence to Jeremy P. Kamil, jkamil@lsuhsc.edu

ABSTRACT.   Human cytomegalovirus (HCMV) encodes an ER-resident glycoprotein, UL148, which activates the unfolded protein response (UPR) but is fully dispensable for viral replication in cultured cells. Hence, its previously ascribed roles in immune evasion and modulation of viral cell tropism are hypothesized to cause ER stress. Here, we show that UL148 is necessary and sufficient to drive the formation of large ER-derived structures that occupy up to 7% of the infected cell cytosol. The structures are found to be sites where UL148 coalesces with cellular proteins involved in ER quality control, such as Hrd1 and EDEM1. Transmission electron microscopy analyses reveal the structures to be comprised of tortuous, densely packed and apparently collapsed ER membranes that connect to distended cisternae. During induced ectopic expression of UL148-GFP fusion protein, punctate signals traffic to accumulate at prominent structures that exhibit poor recovery of fluorescence after photobleaching. Small molecule blockade of the integrated stress response (ISR) prevents the formation of puncta, leading to a uniform reticular fluorescent signal. Accordingly, ISR inhibition during HCMV infection abolishes the coalescence of UL148 and Hrd1 into discrete structures, which argues that UL148 requires the ISR to cause ER reorganization. Given that UL148 stabilizes immature forms of a receptor binding subunit for a viral envelope glycoprotein complex important for HCMV infectivity, our results imply that stress-dependent ER remodeling contributes to viral cell tropism.

IMPORTANCE.

Perturbations to ER morphology occur during infection with various intracellular pathogens and in certain genetic disorders. We identify that an HCMV gene product, UL148, profoundly reorganizes the ER during infection, and is sufficient to do so when expressed on its own. Our results reveal that UL148-dependent reorganization of the ER is a

prominent feature of HCMV infected cells. Moreover, we find that this example of virally

induced organelle remodeling requires the integrated stress response (ISR), a stress

adaptation pathway that contributes to a number of disease states. Since ER

reorganization accompanies the roles of UL148 in HCMV cell tropism and intracellular

retention of the immune cell co-stimulatory ligand CD58, our results may have implications

for understanding the mechanisms involved. Furthermore, our findings provide a basis to

utilize UL148 as a tool to investigate organelle responses to stress and to identify novel

drugs targeting the ISR.

INTRODUCTION.

UL148 is a human cytomegalovirus (HCMV) ER-resident glycoprotein that plays

roles in evasion of cell-mediated immunity and shows intriguing effects on cell tropism.

During infection of epithelial cells, viruses disrupted for *UL148* replicate to produce roughly

100-fold enhanced levels of infectious progeny virions compared to wildtype (1). These

effects correlate with reduced expression of glycoprotein O (gO), a subunit of a

heterotrimeric viral glycoprotein H (gH) / glycoprotein L (gL) complex (gH/gL/gO) that is

required for the infectivity of cell-free virions (2-4). The presence of gO in the context of the

heterotrimer endows the virus with the capacity to utilize the platelet derived growth factor

receptor α (PDGFRα) as an entry receptor (5-7). Accordingly, UL148 has been found to

stabilize immature forms of gO prior their assembly into gH/gL/gO heterotrimers (1, 8).

Despite that UL148 does not stably associate with gO, the data suggest an interaction with

gH (1).

UL148 also physically associates with CD58 (LFA-3), a co-stimulatory ligand for

natural killer cells and T-lymphocytes, preventing its presentation at cell surface (9).

Although the mechanisms by which UL148 stabilizes gO and retains CD58 within the ER

remain unknown, UL148 strongly contributes to activation of the unfolded protein response

(UPR) during infection, and is sufficient to activate the UPR when ectopically expressed in

non-infected cells (10). UL148 co-purifies from infected cells with SEL1L, an adaptor

subunit of ER-based E3 ubiquitin ligase Hrd1 that plays important roles in ER-associated

degradation (ERAD) of terminally misfolded glycoproteins (8). This suggests a physical

interaction with ERAD machinery, which may be germane to the mechanism by which

UL148 activates the UPR.

Here, we show that the expression of UL148 is necessary and sufficient to induce

unusual ER structures at which large quantities of ER factors involved in glycoprotein

quality control accumulate. In electron microscopy analyses, we find that the UL148

induced ER-structures are comprised of densely packed, tortuous ER membranes that form

connections with tubules of highly distended cisternal space. Furthermore, additional data

from inhibitor studies strongly argue that ER remodeling triggered by UL148 requires the

integrated stress response. Overall, our results reveal a striking ER perturbation induced

by HCMV, and which is entirely controlled by a single viral gene product. These findings

may have important implications for understanding how UL148 regulates viral cell tropism

and/or contributes to viral evasion of cell-mediated immunity.

RESULTS.

UL148 causes reorganization of ER quality control proteins into unusual globular structures.

The HCMV ER-resident glycoprotein UL148 was previously observed to co-localize with the ER marker calnexin during infection (1). Nonetheless, calnexin staining did not show the uniform reticular pattern characteristic for the ER marker. We later noticed that cells infected with a *UL148*-null virus showed uniform calnexin staining (see below). To formally determine whether UL148 influences calnexin localization, we compared fibroblasts at four days post-infection with either wildtype strain TB40/E virus (TB_WT) or a *UL148*-null mutant (TB_148$_{STOP}$)(8, 10), each derived from an infectious bacterial artificial chromosome (BAC) clone of HCMV strain TB40/E (11) (**FIG 1**). In cells infected with wildtype virus, calnexin antibodies stained unusual globular structures at the cell periphery, as expected (1) (**FIG 1A, 1C, SI FIG S1A**). However, in cells infected with the *UL148*-null virus, calnexin staining was uniform throughout the cytosol (**FIG 1B-C**), as would be expected for an ER marker in uninfected cells. The staining pattern for Hrd1, another ER marker, likewise indicated accumulation at unusual globular structures during wildtype HCMV infection, but not during infection with *UL148*-null mutant viruses (**FIG 1, S1A**). Because calnexin and Hrd1 staining showed uniform distribution in *UL148*-null virus infected cells, and because *UL148* is fully dispensable for efficient viral replication in fibroblasts (1), these results suggest that redistribution of these two ER markers depends on UL148.

As expected (12, 13), antibodies specific for glycoprotein H (gH), a viral envelope glycoprotein, stained a juxtanuclear compartment, termed the cytoplasmic virion assembly compartment (cVAC), which does not involve the ER (**FIG 1**). Similarly contrasting staining

patterns for Hrd1 and/or calnexin were observed in cells infected with wildtype versus

UL148-null mutants of clinical HCMV strains Merlin and TR (**FIG S1**). Reciprocally, we

restored a functional *UL148* at its native locus in the context of a BAC clone of HCMV strain

AD169, which spontaneously lost most of the gene in the course of extensive genetic

rearrangements that occurred during serial passage in cultured cells. In cells infected with

AD169 repaired for UL148, but not the *UL148*-null parental virus, the Hrd1 staining pattern

showed prominent globular structures (**FIG S1**).

Because similar differences in Hrd1 staining are observed between wildtype and

UL148-null infections of THP-1 macrophages, ARPE-19 epithelial cells and fibroblasts (**FIG**

1, FIG S1), we concluded that UL148-dependent reorganization of ER markers occurs in

multiple cell types. The calnexin staining pattern seen with four different primary clinical

isolates from patient throat swabs likewise indicated punctate globular structures. These

observations, taken together with results from formal comparisons of wildtype versus

UL148-null mutants of four different BAC-cloned HCMV strains (**FIG 1, FIG S1**), argue that

UL148 profoundly affects the ER during natural infection.

We measured the three-dimensional volume of UL148-dependent Hrd1 structures

from sixteen cells fixed at 96 h post infection (hpi) with strain TB40/E. On average, the

structures occupied 303.3 µm^3 (SEM: +/- 24.8) out of a total cell volume of 7657 µm^3

(SEM:+/- 651.0), which was estimated using a phalloidin-fluorophore conjugate to detect

the actin cytoskeleton (**FIG 1D-E**). Subtracting the volume of nuclei, as indicated by 4′,6-

diamidino-2-phenylindole [DAPI] (average nucleus: 1360 µm^3, SEM:+/-129.4), we calculate

that, on average, the structures occupy 5.2% of the cytosolic volume (SEM: +/-0.61%,

range: 2.0% – 7.0%). Based on these findings, taken together with results comparing

additional wildtype (WT) versus *UL148*-null mutant HCMV strains (**FIG S1**), we conclude

that the UL148-dependent ER structures are a prominent feature of HCMV-infected cells.

UL148 substantially contributes to activation of the unfolded protein response (UPR)

during HCMV infection (10), and co-purifies from infected cells with SEL1L (8), an adaptor

subunit for the E3 ubiquitin ligase Hrd1, which plays crucial roles in ER-associated

degradation (ERAD) of terminally misfolded glycoprotein substrates (14). Hence, the

accumulation of Hrd1 and calnexin at unusual structures during wildtype but not *UL148*-null

infection may suggest that structures form in response to defects in ER quality control

(ERQC) caused by UL148. Under conditions of proteasome inhibition, overexpression of

certain misfolded glycoproteins causes cellular factors involved in ERQC, such as calnexin,

to compartmentalize from the rest of the ER, while other ER markers such as BiP (Grp78)

or protein disulfide isomerase (PDI) remain largely unaltered (15-17).

To gain further insights into the nature of these peculiar ER structures, we set out to

develop a more comprehensive understanding of their protein composition by comparing

the localization of selected ER markers at 96 hpi during infection with either TB_148^{HA} (1),

a recombinant HCMV strain TB40/E virus that expresses UL148 fused at its C-terminus to

the nonapeptide epitope tag YPYDVPDYA from influenza A hemagglutinin (HA), or

TB_159^{HA}, a *UL148*-null comparator in which the UL148 homolog from rhesus

cytomegalovirus, *Rh159*, likewise fused to a C-terminal HA tag, is expressed in lieu of

UL148.

We considered the TB_159^{HA} virus to be an appropriate control for the following

reasons. Firstly, Rh159 and UL148 exhibit ~30% identity at the amino acid level and both

glycoproteins localize to the ER and block cell surface presentation immune cell activating

ligands; UL148 retains CD58, a ligand for CD2, while Rh159 retains NKG2D ligands of the

MIC- and ULBP families (1, 9, 18). Secondly, Rh159 is expressed from TB_159^{HA} at

comparable levels and with similar kinetics observed for UL148 from TB_148^{HA}, and the

two viruses replicate indistinguishably in fibroblasts (**FIG 2**). Moreover, as was the case in

ectopic expression (10), Rh159 does not appear to activate the UPR to the same extent

that UL148 does, as evidenced by lower levels of phospho-eIF2α and ATF4 during

TB_159[HA] infection (**FIG 2A**).

ERQC markers failed to coalesce into unusual structures during infection with

TB_159[HA] (**FIG 3**). The structures likewise failed to occur in rhesus fibroblasts during

infection with a recombinant rhesus cytomegalovirus (RhCMV) that we engineered to

express an HA-tag fused at the C-terminus of Rh159 (**FIG S1**). The latter argues against

the possibility that Rh159 somehow requires the context of rhesus cells to redistribute ER

markers in a manner analogous to what is seen for UL148 during HCMV infection. Taken

together, the data suggest that UL148 and Rh159 authentically differ in their effects on the

secretory pathway.

In cells infected with TB_148[HA], HA antibody staining indicated localization of UL148

to globular structures, as expected (1). Antibody signals from indirect confocal

immunofluorescence detection of cellular ER resident proteins involved in ERQC, including

calnexin, Hrd1, SEL1L, Herp, valosin containing protein (VCP, p97), and EDEM1, strongly

co-localized with signals from HA-tagged UL148 (**FIG 3-4**). These results indicate that a

number of ERQC factors coalesce with UL148 to form prominent globular structures in

infected cells. In contrast, we observed uniform ER staining patterns for PDI and

calreticulin (CALR), indicating that these ER markers do not localize to the UL148

structures (**FIG 3Q, 3S**). Intriguingly, antibody signals detecting reticulon 3 and ribophorin

1, which are markers for smooth ER and rough ER, respectively, each appreciably co-

localized with UL148 (HA) signal at the induced structures (**FIG 3K, 3M)**, which indicates

that the structures may involve both rough and smooth ER. LC3B failed to co-localize with

the structures (**FIG S2**), as might be expected given that the virus inhibits macroautophagy

at late times during infection (19, 20). Nevertheless, our results suggested that the induced

structures are enriched for a related mammalian ATG8 ortholog, GABARAP (**FIG 3O**).

In cells infected with TB_159^{HA}, all of the ER markers we examined showed uniform,

reticular staining, as did the anti-HA signal detecting Rh159 (**FIG 3B, D, F, H, J, L, N, R, T,**

V). EDEM1, in addition to showing reticular staining, also labeled puncta associated with

the nucleus (**FIG 3J, 3I**), which may represent enriched levels of the protein at the rough

ER membranes associated with the nuclear envelope, although we cannot exclude the

possibility of spurious intranuclear staining. GABARAP antibodies stained the cVAC in

TB_159^{HA} infected cells, but also showed a much weaker diffuse signal throughout the

cytosol. Even though we took steps to block viral Fc receptors, which can cause rabbit

antibodies to non-specifically label the cVAC (21), it is plausible that GABARAP antibody

signal from the cVAC reflects incomplete blocking of viral Fc receptors. For both viruses,

antibodies specific for the TGN marker syntaxin-6 (STX6) and gH, as expected, stained the

juxtanuclear cVAC structure (12, 13, 22), a post-ER compartment that excludes UL148 (1)

(**FIG S2**).

To quantify the degree of co-localization with UL148, we calculated Pearson's

correlation coefficients from a minimum of thirty TB_148^{HA} infected cells per staining

condition (fixation at 4 dpi), comparing the signal overlap for each ER marker to the HA

signal from UL148. The correlation coefficients (*r*) for staining patterns from antibodies

specific for EDEM1, Hrd1, reticulon-3 (RTN3), SEL1L, the ER retention motif KDEL, and

calnexin ranged from 0.8 to 0.72. These results suggest that these proteins, and lumenal

ER resident proteins such as BiP (GRP78) and GRP94 that contain a C-terminal KDEL

motif (EKDEL), extensively co-localize with UL148 at the induced structures (**FIG 4**).

Meanwhile, GABARAP, VCP, ribophorin-1 (RPN1), and Herp showed *r* values in the range

of 0.54 to 0.51, indicating a moderate degree of co-localization with UL148. However,

CALR and PDI, which like BiP, are lumenal ER residents, and the TGN marker syntaxin-6,

gave *r* values of close to zero, indicating that these markers do not appreciably co-localize

with UL148, as is consistent with our immunofluorescence data suggesting that these

markers negligibly associate with the unusual structures. From these results, we conclude

that the UL148-dependent ER structures are enriched for cellular markers involved in

glycoprotein quality control. In this regard, the structures resemble the "ER quality control

(ERQC) compartments" described by G. Lederkremer and colleagues (15, 16, 23). We

further conclude that Rh159 cannot substitute for UL148 to redistribute ERQC markers.

VCP and Hrd1 are recruited to incipient UL148 ER structures prior to calnexin.

To determine whether there might be differences in the kinetics of recruitment of

ERQC markers during formation of the ER structures, we examined a series of time points

from one to four days post-infection (dpi) with wildtype HCMV strain TB40/E carrying HA-

tagged UL148, staining for three different ERQC markers, calnexin, Hrd1, and VCP,

alongside UL148 (HA). At 1 dpi the signal from UL148 was only faintly detected, as

expected (1), while each of the ER markers was readily detected, providing a readout of

their staining patterns prior to being substantially perturbed by UL148 (**FIG S3**). At 2 dpi,

we detected robust anti-HA signal, indicating the presence of UL148. At this time point,

UL148 exhibited intense signal at small globular puncta, which we interpret to represent

incipient UL148 structures, as well as more diffuse staining of a reticular structure

consistent with undisturbed ER. Calnexin did not appreciably co-localize with the UL148

puncta until at least 3 dpi, and the structures were not readily visualized by calnexin

staining until 4 dpi (**FIG S3**). In contrast, signals from Hrd1 and VCP staining were sufficient

to mark the UL148 puncta by 2 dpi, which suggests that these markers may co-localize with

UL148 at earlier points during the genesis of the ER structures. Notably, the appearance of

Hrd1 at the structures prior to calnexin is consistent with our previous results showing that

UL148 co-purifies from infected cells with SEL1L, an adaptor subunit for Hrd1 (8). These

results imply that basal elements of the ERAD machinery, exemplified by Hrd1 and VCP,

might be recruited to UL148-induced ER structures prior to calnexin.

Visualization of UL148-induced ER structures by electron microscopy.

To discern the ultrastructural appearance of the UL148 structures and to formally ascertain

their relationship to the ER, we carried out transmission electron microscopy (TEM)

imaging of wildtype (TB_WT) and *UL148*-null (TB_148$_{STOP}$) infected fibroblasts at 5 dpi. In

high pressure frozen and freeze-substituted infected fibroblasts, cells with a high density of

viral nucleocapsids within the nucleus were selected for analysis, as this feature indicates

late time points during infection when UL148 is abundantly expressed. The TEM results

revealed prominent globular and oblong structures in the cytoplasm of wildtype virus

infected fibroblasts, but not *UL148*-null infected controls (**FIG 5-6**). The structures stand out

for their high electron-density, which may reflect the abundance of ERQC proteins together

with UL148 in these structures. Under higher magnification, these areas are characterized

by accumulations of densely packed ruffled membranes and membranous material, which

appear to either be collapsed ER, or ER tubules in very close association with each other.

The membrane accumulations were associated with smooth and partially rough ER

structures of seemingly enlarged cisternal space. Further, tomographic reconstruction of

scanning transmission electron microscopy (STEM) data suggest that densely packed ER

cisternae within the structures are interconnected and continuous in three-dimensional

space (**FIG 7B-C, Movie S1**).

UL148 accumulates in a detergent-insoluble form during infection.

Disease-associated variants of certain cellular proteins, such as the A103E mutant

of the calcium channel Orai1, and the E342K "Z" variant of alpha-1 antitrypsin (Z A1AT),

localize to anomalous ER structures reminiscent of those we observe to depend on UL148

(17, 24-27). A103E Orai1 and Z A1AT are found to accumulate in detergent-insoluble

forms within the ER or within ER membranes, respectively, which indicates aggregation or

polymerization of these proteins, and suggests a mechanism for the formation of ER

structures by the proteins (17, 24, 27, 28). We therefore set out to determine whether

differences in solubility might correlate with the differing potentials UL148 and Rh159 to

cause ER reorganization and to activate the UPR (10). To address this question, we

infected fibroblasts with TB_148[HA] or TB_159[HA] at MOI 1 and at various times post-

infection prepared cell lysates in radioimmunoprecipitation assay (RIPA) buffer. After

subjecting the lysates to high-speed centrifugation, we examined the relative levels of

UL148 and Rh159 in the detergent-soluble supernatant and detergent-insoluble pellet

fractions.

A substantial portion of UL148 was detected from the detergent-insoluble fractions at

all time points tested (**FIG 8**). Furthermore, the percentage of UL148 detected within the

insoluble fraction increased over time. In contrast, Rh159 was found only in the detergent-

soluble fraction. From these results, we conclude that UL148 accumulates in a detergent-

insoluble form during infection, and that Rh159 does not do so. Because the anti-HA

immunoreactive band detected in both the soluble and insoluble fractions from TB_148[HA]

infected cells showed a relative mobility of ~35 kD, which matched that expected for the

mature, endoH-sensitive glycoprotein (1), these findings argue that UL148 may form

aggregates or polymers within the ER.

UL148 is sufficient to compartmentalize the ER.

Our data thus far demonstrate that UL148 is necessary during infection to cause

redistribution of cellular ER markers for glycoprotein quality control processes, such that

substantial portion of the ERQC machinery appears to become sequestered away from the

rest of the organelle into novel membranous structures. Because a number of viral proteins

that remodel the ER during infection are sufficient to alter ER morphology when ectopically

expressed [e.g. (29),(30), reviewed in (31)], we wondered whether UL148 expression would

be sufficient to induce ER structures akin to those seen during HCMV infection. Making

use of "tet-on" ARPE-19 epithelial cells that inducibly express either UL148 or Rh159, each

carrying a C-terminal HA-tag (32), we induced transgene expression for 48 h and

subsequently stained for various cellular markers for the ER, for ERQC as well as for the

ATG8 family proteins LC3B and GABARAP, which play roles in macroautophagy and

related processes.

In cells expressing UL148 (i148[HA]), we observed that calnexin, Hrd1, EDEM1, and

VCP co-localized with UL148 at prominent globular structures reminiscent of those

observed during infection (**FIG S4**). The respective rough and smooth ER markers

ribophorin 1 (RPN1) and reticulon 3 (RTN3), as well as the ATG8 proteins LC3B and

GABARAP likewise co-localized to the UL148-induced structures. However, PDI and CALR

each failed to substantially co-localize with UL148 at the structures (**FIG S4**), or at best

showed only limited co-localization, consistent with our results from infected cells (**FIG 3-4**).

Signals from antibodies specific for the KDEL (EKDEL) motif important for ER retrieval of

luminal ER residents, such as BiP and GRP94, likewise showed only limited co-localization

with UL148. The PDI and KDEL results suggest that large regions of the ER are not

involved in the UL148 structures.

In cells expressing the UL148 homolog Rh159 (i159[HA]), we detected uniform

cytoplasmic distribution of ER markers, as well as of HA-tagged Rh159, similar to what we

observed during infection with the recombinant HCMV TB_159^{HA} (**FIG 3**). Furthermore, in

this setting the staining patterns for the ATG8 family proteins LC3B and GABARAP failed to

indicate any notable structures. Consistent with our previous study showing that UL148, but

not Rh159, is sufficient to activate the UPR (10), we observed accumulation of ATF4 and

phosphorylated eIF2α following doxycycline (dox) induced expression of UL148 but not

Rh159 (**FIG 9A**). Importantly, the intensity of HA signals detecting UL148 and Rh159

indicated that both ER resident immunevasins accumulate at roughly comparable levels

following dox induction. This argues against the possibility that gross protein expression

differences accounts for the differing effects of UL148 and Rh159 on UPR activation and on

the staining patterns for ER markers.

Pearson's correlation coefficient values were calculated to quantify the extent of

colocalization between UL148 and various cellular markers: EDEM1, Hrd1, VCP, calnexin

(CNX), RTN3, SEL1L, LC3B, and PDI (**FIG 9B**). The results quantitatively buttress the

confocal immunofluorescence results from these cells (**FIG S4**). We therefore conclude

that no other HCMV gene products are required for UL148 to cause ERQC factors to

segregate into discrete compartments. Hence, we conclude from these findings that UL148

is sufficient to cause large-scale reorganization of ERQC markers. Moreover, these results

suggest that reorganization of ER markers into discrete structures may be related to the

propensity of UL148 to induce ER stress.

To determine whether UL148-dependent ER remodeling could be visualized in real-

time using live cell imaging, we constructed new "tet-on" lentiviral vectors that inducibly

express either UL148 or Rh159 fused at their predicted C-terminal cytoplasmic tails to the

enhanced green fluorescent protein (GFP) from *Aequorea victoria* (33). Following lentiviral

transduction, we isolated puromycin-resistant ARPE-19 and subsequently subjected to

fluorescence-activated cell sorting (FACS) to enrich for cells that expressed GFP signal

following dox treatment. The resulting cell populations, which inducibly express either

UL148-GFP or Rh159-GFP, were designated i148^{GFP} and i159^{GFP}, respectively.

In live cell imaging studies, we observed that the GFP signal in dox-induced i148^{GFP}

cells first appeared in a reticular, largely uniform pattern, which was readily visible at 5 h

post-induction. However, by 9 h post-induction punctate signals began to appear. These

puncta were observed to traffic to sites of large-scale accumulation, where large

fluorescent structures progressively increased in size up until the termination of the

experiment at 19 h (**FIG S5, Movie S2**). In contrast, i159^{GFP} cells exhibited a largely

diffuse, reticular GFP signal at all time points monitored subsequent to transgene induction

(**FIG S5, Movie S3**). The live cell imaging results thus recapitulated the differences in the

HA-staining patterns observed between dox-induced i148^{HA} and i159^{HA} cells following

fixation, as well as those between HCMV expressing HA-tagged UL148 versus *UL148*-null

comparator viruses (**FIG 1, 3, Fig S1**).

UL148-GFP structures exhibit poor recovery of fluorescence after photobleaching.

To determine whether the protein contents of the UL148 induced structures exhibits

decreased mobility compared to non-perturbed ER regions, we carried out fluorescence

recovery after photobleaching (FRAP) studies. We photobleached regions of GFP signal in

i148^{GFP} or i159^{GFP} ARPE-19 cells that had been induced for transgene expression for 24 h,

and then monitored recovery of fluorescence over a 5 min time period. In Rh159-GFP

expressing cells (i159^{GFP}), fluorescence nearly recovered to pre-bleach levels within 3 min,

and by 5 min fully recovered (**FIG S6**). In contrast, when we photobleached a prominent

UL148-GFP structure, the GFP signal failed to appreciably recover fluorescence during the

same 5 min time period. Nonetheless, regions of reticular GFP signal from an i148^{GFP} cell,

which presumably represent ER regions not involved in an anomalous structure, recovered

fluorescent signal with kinetics similar to those observed during Rh159-GFP expression.

These results indicate that bleached UL148-GFP within the structures cannot rapidly be

exchanged with UL148-GFP from other portions of the organelle, which suggests that the

induced UL148 structures do not exchange their contents as efficiently as unperturbed ER.

Ectopically-expressed UL148 is degraded by proteasomes and not by autophagy.

The UL148 ER structures were found to be enriched with proteins such as Hrd1,

SEL1L, EDEM, and VCP (**FIG 3-4**, **9B, FIG S4**), which are posited to play key roles in

ERAD, a process during which misfolded glycoproteins are recognized, processed, and

dislocated across the ER membrane for degradation at cytosolic proteasomes. However,

we also detected elements of the machinery for autophagy in close association with the

UL148 structures. Namely, the mammalian ATG8 ortholog GABARAP co-localized with

UL148 during infection (**FIG 3-4**), and both GABARAP and another ATG8 ortholog, LC3B,

co-localized with UL148 at the ER structures during ectopic expression of UL148 (**FIG 9B,**

FIG S4). Therefore, we wished to determine whether UL148 is degraded by the

proteasome, as would be consistent with conventional ERAD, or by the lysosome, which

would suggest a role for autophagy-related processes, such as selective autophagy of the

ER (34), in dispensing with UL148, and presumably, in resolving the ER perturbations.

As a first step, we conducted a live-cell imaging "washout" experiment in which

i148^{GFP} cells were induced for transgene expression for 24 h, after which the growth

medium containing the transgene inducing agent (dox) was replaced with medium lacking

dox. During a 22 h imaging period following dox-washout, we observed the structures to

become progressively smaller as the reticular and punctate GFP signals gradually abated

(**SI FIG S7, Movie S7**). Because the results suggested that UL148-GFP structures could

be resolved over time, we induced UL148-GFP in tet-on ARPE-19 cells (i148^{gfp}) for 24 h,

washed out the inducing agent (dox), and then applied either epoxomicin (20 µM), an

irreversible inhibitor of the proteasome (35), or folimycin (115 nM), a proton pump inhibitor

that blocks autophagosome maturation and impedes lysosome-dependent degradation (36,

37). Of the two inhibitors, only epoxomicin stabilized UL148 (**SI FIG S7**). Although

folimycin treatment had no obvious effect on UL148-GFP levels, the drug markedly

increased the levels of LC3B-II, as would be expected with the lysosomal proton pump

inhibitor. From these results, we conclude that ectopically expressed UL148 is primarily

degraded by proteasome-dependent pathway.

UL148 requires the integrated stress response to cause ER reorganization.

We previously reported that UL148 triggers the UPR during ectopic expression, and

that UL148 contributes to activation of the PKR-like ER kinase (PERK) and inositol

requiring enzyme 1 (IRE1) during infection (10). The literature suggests that formation of

ERQC compartments requires PERK (38). PERK responds to ER stress by activating the

integrated stress response (ISR). In particular, PERK phosphorylates eIF2α at Ser51, and

the accumulation of phosphorylated eIF2α (eIF2α-P) globally attenuates mRNA translation

whilst stimulating the translation of a subset of mRNAs, such as those encoding ATF4 and

CHOP, which play roles in cellular adaptation to stress (39). Stress-regulated translation of

such mRNAs involves small upstream open reading frames (uORFs) in their 5' untranslated

regions that ordinarily suppress translation under basal conditions (40, 41).

To examine whether ER remodeling in response to UL148 requires the ISR, we

turned to a well characterized small molecule inhibitor of stress-regulated translation, ISRIB

(42-46). ISRIB is thought to act as a "molecular staple" that holds the guanine nucleotide

exchange factor eIF2B in an active decameric configuration (44-47), such that eIF2B will

continue to generate ternary complex (eIF2•GTP•Met-tRNAi) necessary for new cycles of

translational initiation, despite the presence of eIF2α-P. Because PERK is the kinase that

phosphorylates eIF2α in response to ER stress (48, 49), we also tested for effects of the

PERK inhibitor GSK2606414 (50). Having confirmed that ISRIB and GSK260641 do not

negatively impact UL148 expression during dox-induction of UL148-GFP **(FIG 10)**, we

treated i148^{GFP} cells with 200 nM ISRIB, 1.1 µM GSK260641, or DMSO vehicle control, and

carried out live cell imaging during induction of UL148-GFP expression.

ISRIB and GSK6060414 virtually abolished the formation of UL148 puncta through

13 h post induction (dox addition), a time point at which the DMSO control condition

showed abundant large structures (**FIG 10C-E, SI Movies S8-S10**). Based on our findings

(**FIG 1, 3, 5, 7, 9B, S4**), we interpret the formation and subsequent large-scale aggregation

of UL148-GFP puncta to faithfully indicate ER reorganization in response to UL148.

Therefore, these results suggested to us that pharmacological inhibition of either the ISR or

PERK prevent UL148 from remodeling the ER, which implies that ER remodeling in

response to ER stress requires stress-regulated translation.

Since our results with ISRIB and GSK6060414 indicated that the ISR is required for

ER remodeling to occur during ectopic expression of UL148, we next asked whether these

pharmacological agents would prevent UL148-dependent ER remodeling in the

physiologically authentic context of HCMV infection. Our live cell imaging studies from

ISRIB and GSK6060414 treatments indicated that UL148-GFP puncta could begin to

appear by ~19 h post treatment (**FIG 10, SI Movies S9-S10**), which suggested that the

pharmacological activity of the drugs might diminish over time. Thus to ensure that

blockade of PERK and the ISR would be sustained over the 4 d period that it takes for

structures to fully form during HCMV infection, we replenished drug treatments every 24 h,

replacing the spent media with media containing freshly reconstituted ISRIB, GSK6060414,

or DMSO vehicle control.

At 4 dpi, ER structures were readily visible in DMSO control treatment condition but

virtually abolished in the ISRIB and GSK2606414 conditions (**FIG 11A**). To quantify these

effects, we obtained Z-stacks from a minimum of 30 cells per treatment condition and used

Imaris 3D image analysis software to calculate the percentage of Hrd1 signal that

coalesced into discrete structures during infection. The results indicated highly significant

differences between the DMSO carrier alone setting compared to treatments with either

GSK2606414 or ISRIB (**FIG 11B**). In the DMSO control condition, on average, 10% of the

Hrd1 signal was found to be associated with the UL148 structures (arithmetic mean; range

5.8% – 17.6%), as indicated by structures delimited by HA signal detecting UL148. In the

presence of ISRIB or GSK2606414, however, these values were reduced to 2.8% (range:

0.69% – 7.1%) and 3.6% (range: 0.8%-6.5%), respectively. Reassuringly, roughly

equivalent levels of UL148 were detected in Western blot analyses of protein extracts from

drug treated and DMSO control conditions. Therefore, differences in UL148 expression are

unlikely to explain the observed failure of UL148 and Hrd1 to coalesce into ER structures

during inhibition of either the ISR or PERK. Meanwhile, ISRIB and GSK2606414

treatments both led to reduced levels of ATF4 accumulation, while the PERK inhibitor alone

was able to reduce the levels eIF2α-P, as indicated by a phospho-specific antibody.

Therefore, each of the drugs caused the expected effects on the PERK, eIF2α-P, and

ATF4 axis (**FIG 11C-D**).

A modest reduction in the levels of the viral envelope glycoprotein B (gB) was

observed during treatments with ISRIB or GSK2606414. This may suggest that the ISR is

required for optimal expression of the certain viral envelope glycoproteins. Even though

UL148 promotes ATF4 expression, likely via PERK activation (10), viruses that lack *UL148*

cause high levels of ATF4 accumulation at late times during infection, which is when viral

late gene products, such a gB, are expressed at their highest levels (10). Hence, the

activity of PERK and/or of the ISR may be required for optimal expression of viral envelope

glycoproteins. Regardless, the PERK inhibitor and ISRIB each prevented the appearance

of discrete UL148 ER structures while failing to substantially affect UL148 expression.

Therefore, the effects of the two inhibitors on UL148-dependent ER reorganization were

consistent with those seen during ectopic expression of the protein. Taken together, our

findings argue that the ISR is required for UL148 to cause remodeling of the organelle.

DISCUSSION

The ER structures that we have identified are noteworthy in several regards. Firstly,

this example of virally-induced ER remodeling is wholly dependent on a single viral gene

product, UL148. Of course, there are other examples of individual viral gene products that

are necessary to grossly perturb ER morphology during infection which are also found to be

sufficient to induce such perturbations [e.g., (29, 30, 51, 52)]. Nonetheless, the ER

structures caused by UL148 are— to the best of our knowledge, a hitherto undescribed

ultrastructural characteristic of the HCMV infected cell, which is surprising given their size,

scale, and prominence. Indeed, the average volume of the structures at 96 h postinfection

amounts to roughly 60% of the size of the nucleus of an uninfected human fibroblast [~500

μm^3, (53)]. Although UL148 expression is accompanied by UPR activation, ER

reorganization occurs without killing the host cell, even in the setting of ectopic expression

when viral functions that inhibit programmed cell death are absent. Nor does UL148 affect

the yield of infectious virus during replication in fibroblasts, since *UL148*-null mutant viruses

replicate indistinguishably from wildtype parental virus in this cell type (1, 8). As with all

enveloped viruses, HCMV requires the host cell secretory pathway to fold and process

enormous quantities of viral envelope glycoproteins that endow progeny virions with the

molecular machinery for infectivity. It is intriguing to consider how the infected cell tolerates

spatial reorganization of ER membranes and of the associated glycoprotein quality control

machinery without impacting the production of infectious progeny virions.

Furthermore, certain aspects of ER reorganization invoked by UL148 may be novel

to cell biology. In particular, the densely packed collapsed ER in close association with

varicosities that we observe to depend on UL148 during infection appear to differ from the

ER structures induced by other ER perturbagens. For instance, a hereditary form of

childhood cirrhosis caused by the *Z* allele of the alpha-1-antitrypsin (Z-A1AT) gene

(*SERPINA*) is characterized by polymerization of Z-A1AT within the ER (26), leading to its

accumulation in membrane-delimited inclusions (26, 27, 54). However, Z-A1AT inclusions

are not observed to associate with regions of collapsed ER, nor does Z-A1AT suffice to

activate the UPR, even though its expression sensitizes cells to other triggers of ER stress

(27, 55).

HCMV encodes at least one other ER-resident immunevasin that activates the UPR,

US11 (56). UL148 binds the NK-cell and T-cell costimulatory ligand CD58 to prevent its

transport to the cell surface while US11 targets the heavy chain of class I major

histocompatibility complex (MHC) for ER-associated degradation (57, 58). UL148 causes

CD58 to accumulate as an immature glycoform, presumably within the ER, but does not

lead to any obvious decrease in its overall abundance (9). Albeit that intracellular forms of

CD58 are in our hands refractory to detection by standard indirect immunofluorescence

protocols (data not shown), it seems reasonable to hypothesize that UL148 sequesters

CD58 into the unusual ER structures that it induces. Moreover, our detergent solubility

results suggest that UL148 may form aggregates or polymers in the course of preventing

CD58 presentation at the cell surface (FIG 8). It is worth noting that UL148, an

immunevasin targeting CD58, induces dramatic morphologic rearrangements of the ER,

while another HCMV immunevasin, US11, targeting MHC class I for degradation evidently

does not do so, even though both trigger ER stress. Nonetheless, it seems unlikely that

retention of CD58 would be required for ER reorganization. In fact, the capacity of UL148

to retain CD58 may be functionally separable from its peculiar effects on the morphology

and organization of the ER.

Despite that the relationship between the mechanism for CD58 retention and the

formation of UL148-dependent ER structures remains unresolved, our findings may

suggest a mechanism to explain the influence of UL148 on the tropism of the virus for

epithelial cells. These effects, exemplified by a ~100-fold replication advantage of *UL148*-

null viruses during infection of epithelial cells, correlate with decreased expression of

glycoprotein O (gO), a viral envelope glycoprotein, both in virions and in infected cells (1).

We previously reported that gO behaves as a constitutive ERAD substrate during infection

and that immature, newly synthesized forms of gO show enhanced stability in the presence

of UL148 (8). Our findings herein show that UL148 causes large-scale sequestration of

cellular factors important for ERAD, such as the ER mannosidase EDEM1 and the E3

ubiquitin ligase SEL1L/Hrd1, into large membranous structures. Moreover, the reticular

ER, as indicated by the staining patterns of antibodies specific for calreticulin, PDI, and the

KDEL motif, appears to remain largely intact following UL148-induced ER reorganization

(FIG 3, 10). Since these observations indicate that UL148 depletes ERAD factors from the

ER during the formation of the unusual structures, regions of ER not drawn into the

structures might be expected to offer a more permissive folding environment for

polypeptides, such as gO, which either fold slowly or inefficiently assemble into multiprotein

complexes.

Based on these observations, one might hypothesize that UL148 alters ER

proteostasis by physically dislocating (or sequestering) key elements of the "mannose

removal time clock" system that marks poorly folding glycoproteins for destruction via

ERAD (59, 60). Because gO is both the defining subunit of the heterotrimeric gH/gL/gO

envelope glycoprotein complex that governs HCMV entry and cell tropism (5-7, 61, 62), and

a constitutive ERAD substrate (1, 8), ER reorganization might be required for the effects of

UL148 on HCMV cell tropism. Going forward, it will be crucial to determine whether

classical ERAD substrates, such as the null Hong Kong variant of alpha-1-antitrypsin (63,

64) or ribophorin-332 (65, 66), are stabilized during UL148 expression, as would be

predicted if UL148 shifts ER proteostasis to negatively modulate ERAD.

In addition to having found that factors involved in proteasomal ERAD, such as Hrd1

and EDEM1, are enriched at the UL148 structures, we observed that GABARAP, a

mammalian ortholog of yeast ATG8, localizes to the UL148 ER structures during infection,

and that another ATG8 ortholog, LC3B, and GABARAP both associate with the structures

during ectopic expression of UL148. This may suggest roles for autophagy-related

pathways in UL148-dependent reorganization of the organelle. Since our results indicate

that UL148 is degraded by a proteasome-dependent pathway and not by lysosomes (**FIG**

S7), it seems unlikely that selective autophagy of the ER is directly involved in recycling or

degrading these ER structures. Nonetheless, ATG8 family proteins may contribute to

formation of the globular ER structures, since we observed trafficking of UL148-GFP

puncta to form large aggregates during live cell imaging (**FIG 12, 15, Movies S2, S8**). The

literature indicates that misfolded proteins traffic in a microtubule (MT)-dependent manner

to form pericentriolar structures termed aggresomes, which in the case of ERAD substrates

such as the ∆F508 mutant of the cystic fibrosis transmembrane conductance regulator

(CFTR), contain deglycosylated protein that presumably has already undergone dislocation

from the ER (67, 68). ATG8 family proteins such as LC3B and GABARAP not only play

roles in degradation of substrates via macroautophagy, but also bind MTs and recruit

machinery for MT-dependent transport of cargos (69-71). Therefore, GABARAP and LC3B

may be important for the recruitment of cellular machinery that transports perturbed ER

cargoes to sites of large-scale accumulation.

Although our results argue that the ISR is required for ER reorganization during

UL148 expression, precisely how UL148 triggers ER stress remains unknown. UL148 has

been found to co-purify from cells with SEL1L, a component of the ERAD machinery.

Thus, it seems plausible that UL148 may inhibit the Hrd1/SEL1L complex, which would

cause the buildup of unfolded proteins and thus trigger the UPR. However, inhibition of

ERAD in and of itself seems unlikely to account for the formation of ER structures. Another

non-mutually exclusive possibility is that UL148 multimerizes or aggregates in a manner

that constricts the ER lumen. For instance, the assembly of UL148 molecules on opposite

sides of the ER lumen might constrict the organelle in a manner consistent with the

collapsed regions of ER observed in our EM results (**FIG 6, 8**).

Additional work will be needed to decipher the molecular mechanisms by which

UL148 causes reorganization of the ER, and to determine its physiological relevance in the

setting of viral biology. Nonetheless, we have shown that UL148, when fused to a

fluorescent protein (FP), suffices both to trigger and to indicate the presence of a functional

ISR (**FIG 10-11, S5**). Hence, UL148-FP fusions may prove useful in high throughput

chemical-genetic screens to identify novel small molecule inhibitors of the ISR as well as to

identify cellular genes involved in stress-dependent remodeling of the ER. Interestingly, the

two pharmacological agents that block UL148-dependent ER remodeling, ISRIB and

GSK2606414, are known to prevent the formation of stress granules (SGs) (43, 72). SGs

are comprised of condensed aggregates of protein and RNA which occur due to stalled

mRNA translation in the context of disease states, such as amyotrophic lateral sclerosis

(73), as well as during treatments with toxic agents such as arsenite (43, 72).   It is

fascinating to consider that mechanistic parallels exist between the formation of SGs and

the UL148-dependent reorganization of the ER. Moreover, given the broad importance

macromolecular aggregation in pathological conditions such as neurodegenerative

diseases (74), certain of which also involve defects in ER proteostasis and aberrant

activation of the UPR, UL148 may hold promise as tool to discover new agents to

ameliorate disease.

MATERIALS AND METHODS.

Cells and virus.

hTERT-immortalized human foreskin fibroblasts (8), derived from ATCC HFF-1 cells

(SCRC-1041) were maintained in Dulbecco's modified Eagle's medium supplemented with

5%-10% newborn calf serum (Millipore Sigma) and antibiotics (complete DMEM) exactly as

described previously (8). i148^{HA} and i159^{HA} ARPE-19 epithelial cells (10), which upon

treatment with 100 ng/mL doxycycline, express HA-tagged UL148 or Rh159, respectively,

were likewise maintained in complete DMEM. For live cell imaging studies, i148^{HA} and

i159^{HA} ARPE-19 were maintained in Opti-MEM (Thermo Fisher) supplemented with 3%

certified tet-approved fetal bovine serum (FBS), 20 μg/mL gentamicin sulfate, 1 μg/mL

puromycin HCl and 10 μg/mL ciprofloxacin HCl. Telomerase-immortalized rhesus

fibroblasts (75) were a kind gift of Peter A. Barry, and were maintained in complete DMEM.

The acute monocytic leukemia cell line THP-1 (TIB-202) was obtained from ATCC

(Manassas, VA) and maintained as suspension cultures in RPMI 1640 medium

supplemented with 10% FBS (Millipore Sigma) and antibiotics (10 μg/mL ciprofloxacin and

25 μg/mL gentamicin). THP-1 were differentiated into adherent macrophages by incubating

for 48 h in the presence of 100 nM 2-O-tetradecanoylphorbol 13-acetate (Millipore Sigma),

and subsequently infected with the indicated viruses at an MOI of 5 TCID$_{50}$/cell.

Infectious bacterial artificial chromosome (BAC) clones of the following HCMV

strains were used for this study: TB40/E (also known as TB40-BAC4 or TB_WT) (11) as

well as its derivatives TB_148^{HA} and TB_148$_{STOP}$, which were used in our previous studies

(1, 8, 10); TR (TR*gfp*) (76, 77); Merlin repaired for *UL128* and *RL13* harboring *tetO*

sequences upstream of *UL131* (pAL1393) (78); AD169 repaired for *UL131* (AD_r131)(1,

79) and AD_r131_148^{HA}, a derivative of AD_r131 that carries a full length UL148 (from

strain TB40/E) tagged with an HA-epitope at the original *UL148* locus (8). An infectious

BAC clone of rhesus CMV (RhCMV) strain 68-1 (80) and a derivative that expresses an

HA-tagged Rh159 (details below) were also used for certain experiments.

The methods used for reconstitution of HCMV from purified BAC DNA, cultivation of virus

and preparation of stocks, including ultracentrifugation through sorbitol cushions and

determination of infectious titers by tissue culture infectious dose 50 ($TCID_{50}$) assay have

been described elsewhere (8, 10).

Construction of recombinant viruses and new plasmids for the study.

New recombinant viruses for this study were derived from BAC-cloned HCMV and

RhCMVs using *en passant* mutagenesis in GS1783 *E. coli,* a strain K12 derivative that

expresses the homing endonuclease I-SceI upon treatment with L-(+)-arabinose (81, 82).

Recombinant BACs were confirmed by Sanger DNA sequencing of the modified regions,

which was performed by Genewiz, Inc. (Piscataway, NJ) (not shown). Oligonucleotide

primers for construction of recombinant viruses and plasmids were custom synthesized by

Integrated DNA Technologies (Coralville, IA); sequences are provided in **Table S1**. Type II

restriction endonucleases, T4 DNA ligase, and Gibson assembly reagents (NEB HiFi

Assembly Master Mix) for generation of recombinant DNAs and for routine molecular

cloning and subcloning procedures were obtained from New England Biolabs (Ipswitch,

MA). KOD Hot Start DNA polymerase (EMD Millipore) was used for all PCR reactions.

To construct TB_159^HA, a strain TB40/E derivative in which the UL148 open reading

frame is replaced by Rh159 we carried out the following steps. Primers Rh159_Fw and

Rh159_HA_Rv were used to amplify and HA-tag the *Rh159* ORF from

pcDNA3.1+_Rh159_IRES_GFP (a gift from Klaus Früh, OHSU). The PCR product was

inserted into EcoRV-digested pEF1α using Gibson assembly. A PCR product containing

an ISce-I excisable kanamycin cassette was amplified from TB_148^{HA}_*ISce-Kan* integrate

BAC (1) using primers PpuISceKanGibs_Fw and PpuISceKanGibs_Rv and Gibson-

assembled into PpuMI-digested pEF1α_Rh159 plasmid to yield plasmid

pEF1α_Rh159_ISceKan. Primers Rh159_Fw_recomb and Rh159_Rv_recomb were used

to generate a PCR product from template plasmid pEF1α_Rh159_ISceKan. The PCR

product was electroporated into GS1783 E. coli harboring the TB_148^{HA} BAC, and Kan^r

colonies harboring TB_Δ148_Rh159_ISceKan integrate BACs were isolated on Luria-

Bertiani agar plates. Bacterial colonies representing kanamycin resistant integrates were

obtained and were subsequently resolved to 'scarlessly' remove the positive selection

marker by standard *en passant* protocols (81, 82) to yield TB_Δ148_Rh159^{HA}, which we

abbreviate herein as TB_159^{HA}. TB_159^{HA} was sequence-confirmed using primers

TB_159HAseq_F and TB_159HAseq_R. Similar strategies were used to modify the

RhCMV 68-1 BAC (80) to incorporate sequences encoding an HA-epitope immediately

before the stop codon of *Rh159*. An *en passant* strategy in which a shuttle plasmid

carrying *I-SceI-Kan^r* disrupted by in-frame nonsense codons was used insert premature

nonsense codons in the *UL148* CDS in the context of HCMV strains Merlin (pAL1393) and

TR (TR*gfp*), and was applied as described previously for generating TB_148$_{STOP}$, the

UL148$_{STOP}$ mutant of strain TB40/E (8).

To construct "tet-on" lentivirus vector plasmids containing UL148 fused to GFP, we

used primers Gibs_eGFP_Rv and 148_eGFP_Fw to amplify the GFP gene from a dsDNA

gBlock, EGFP-P2A-3XHA, synthesized by Integrated DNA Technologies (Coralville, IA),

which was a gift of Matthew D. Woolard (LSU Health Sciences Center, Shreveport, LA). In

a separate PCR, the *UL148* gene was amplified from plasmid pcDNA3.1-UL148^{HA} (10)

using primers Gib_148_Fw and 148_noStopRv. The two products were assembled, using

Gibson Assembly, together with EcoRV opened pcDNA3.1(+) (Invitrogen) to produce

pcDNA3.1-UL148-GFP.   After confirming the absence of spurious mutations by Sanger

DNA sequencing (Genewiz, not shown), the *UL148-GFP* cassette from pcDNA3.1-UL148-

GFP was released by EcoRI and NotI digestion and ligated into pOUPc turboRFP-link

plasmid which had been linearized using the same restriction sites. pOUPc turboRFP-link

is a derivative of lentiviral vector pOUPc (10) that was constructed by reinsertion of

turboRFP (tRFP). Briefly, tRFP was amplified from the original pOUPc-turboRFP using

primer pair tRFP linker Fw and tRFP linker Rv and then reinserted by Gibson Assembly

with pOUPc turboRFP that had been double-digested using EcoRI and Not I in order to

linearize the vector and remove the *turboRFP* (RFP) sequence.

To express Rh159 fused to eGFP, primers Gibs_eGFP_Rv and 159_eGFP_Fw were

used to amplify the GFP gene from EGFP-P2A-3XHA, and in a separate PCR reaction,

primers Gibs_159_Fw and 159_noStop_Rv Rh159 were used to amplify *Rh159* from

pcDNA3.1-Rh159[HA] (10). The latter two PCR products were assembled together with

EcoRV linearized pcDNA3.1(+) using Gibson Assembly, resulting in plasmid pcDNA3.1-

Rh159. The latter plasmid was used as template in a PCR reaction with primers

Gibs_Age_159_Fw and Gibs_Mlu_GFP_Rv. The resulting PCR product was Gibson

assembled into the lentiviral vector pOUPc (10) which had been double-digested with Mlu I

and Age I.

Drug treatments.

The PERK inhibitor, GSK2606414 (50), doxycycline hyclate, and folimycin were obtained

from MilliporeSigma (Burlington, MA). ISRIB (42-46) was obtained from MilliporeSigma or

APExBio (Boston, MA) and epoxomicin was procured Selleck Chemicals (Houston, TX).

ISRIB was dissolved in DMSO to make a 10,000× stock solution (2 mM) and used at a final

working concentration of 200 nM. GSK2606414 was dissolved in DMSO to make a

10,000× stock solution (11 mM) and used at a final working concentration of 1.1 µM.

Folimycin was prepared as a 1000× (115.46 µM) stock solution in DMSO and used at 115

nM final. Epoxomycin was prepared as a 100× (2 mM) stock solution in DMSO and used at

a final concentration of 20 µM.

Confocal microscopy and live-cell imaging. Confocal indirect immunofluorescence

microscopy imaging on fixed cells was carried out using Leica TCS SP5 Spectral Confocal

Microscope (Leica Microsystems, Heidelberg, Germany) using a Leica HCX PL APO CS

63x/1.4-0.6NA objective under oil immersion, except for the image shown in Figure 1C,

which was captured on a Nikon SIM-E and A1R confocal microscopy system (Nikon

Instruments, Melville, NY) using a Nikon SR Apo TIRF 100x/1.49NA objective under oil

immersion. Images for different fluorophore channels were acquired using sequential

scanning. Direct immunofluorescence live cell imaging data were collected using the Nikon

SIM-E microscope using a Nikon Apo 60x/1.40NA DIC objective.   The 3D projection shown

in FIG 1 was generated by NIS-Elements AR Analysis 4.60.00 (64-bit) software (Nikon)

from Z-stacks captured on the Nikon SIM-E / A1R microscope using a Nikon SR Apo TIRF

100x/1.49NA objective. For FIG 11, 3D projections were generated by Imaris x64 9.3.0

software (Bitplane, Inc.) in maximum intensity projection (MIP) mode.

For fixed cell imaging experiments other than those shown in Fig S1D, cells were

seeded on 12 mm circular, No. 1 thickness microscope cover glass (200121; Azer Scientific,

Morgantown, PA). At the indicated times post-treatment and or post-infection, cells were

washed with phosphate buffered saline (PBS) [137 mM NaCl, 2.7 mM KCl, 10 mM Na_2HPO_4,

and 1.8 mM KH_2PO_4, pH 7.4]; fixed for 15 min at room temperature in PBS containing 4%

(wt/vol) paraformaldehyde (Fisher Scientific, Waltham, MA), washed in PBS, permeabilized

for 10 min using 0.1% Triton X-100 (in PBS), subsequently washed again in PBS, and then

blocked for 45 min at 37 °C in PBS containing 5% (vol/vol) normal goat serum (Rockland

Immunochemicals, Limerick, PA). Cells were then washed three times in PBS followed by

incubation in 1% Human Fc Block (BD Biosciences, San Jose, CA) in PBS, for additional 45

min at 37 °C. Cells were then incubated in the presence of primary antibodies for 1 h at 37

°C or 4 °C overnight, and then washed three times with PBS containing 0.1% Tween-20

(PBST) for 5 min per wash. Alexa Fluor-labeled goat polyclonal secondary antibodies (all

from ThermoFisher Invitrogen, Waltham, MA, see **Table S2**) were used for secondary

detection. The slides were then mounted using Prolong Gold anti-fade reagent containing

DAPI (ThermoFisher) and placed under a Leica TCS SP5 confocal microscope for image

acquisition using a Leica 63× oil immersion objective lens (Leica Microsystems, Buffalo

Grove, IL).

For indirect immunofluorescence staining results of primary clinical isolates (SI Fig

S1D), four clinical HCMV isolates obtained by routine testing of throat swabs from patients

of the Ulm University Medical Center were provided by the diagnostic laboratory of the

Institute of Virology in Ulm. Sample material was applied to human fibroblasts and incubated

for several days until HCMV-positive cells could be detected. Infected cells were then seeded

together with uninfected fibroblasts, incubated for up to 5 days until plaques were formed

and processed for indirect immunofluorescence staining. ERQC compartments were

detected by staining for calnexin (CNX, E10; Santa Cruz Biotechnology, mouse, 1:50

dilution), the cVAC was detected by staining for HCMV pUL71 (rabbit, 1:500). Secondary

antibody used for detection of CNX was Alexa Fluor 555 labeled goat anti-mouse IgG

(1:1000) and for pUL71 detection, Alexa Fluor 488 conjugated goat anti-rabbit IgG (1:1000).

Confocal images were acquired using the 63× objective of a Zeiss Observer Z1 fluorescence

microscope equipped with Apotome and Zen software 2.3 (Carl Zeiss Microscopy GmbH,

Jena, Germany).

FRAP. FRAP studies were carried out on using live ARPE-19 cells from the i148[GFP] and

i159[GFP] populations as follows. Cells were seeded as above for live-cell imaging, induced

for transgene expression using dox (100 ng) for 24 h, and then placed on an incubated

sample stage of a Nikon A1R SIM-E imaging system equipped with an SR Apo TIRF

100x/1.49NA objective (Nikon). Three rectangular regions in image fields were defined for

measurement of (i) background signal, and of two regions with comparable initial GFP

signals, (ii) one designated as a control region (no photobleaching) and another for (iii)

photobleaching and recovery of signal after photobleaching. Photobleaching of selected

regions was carried out for 20 s using 405 nm laser light from a LU-N4 laser fiber (Nikon)

[power at source: 15 mW, power at objective: 8 mW]. Images and signal intensity

measurements were captured every 2 s at a rate of 1 frame per s (488 nm excitation, FITC

channel) starting immediately before photobleaching (t=1), and from t=20 s to t=320 s after.

Electron microscopy. Procedures to prepare samples for transmission electron microscopy

(TEM) included high-pressure freezing (HPF), freeze substitution, and Epon embedding,

which were carried out as described previously (83). Briefly, human fibroblasts were seeded

in µ-Slides (Ibidi GmbH, Martinsried, Germany) containing carbon-coated sapphire discs

(Engineering Office M. Wohlwend GmbH) 1 day prior to infection at 80 to 90% confluence.

Cells were infected with virus overnight at MOI 1. Medium containing viral inocula was

replaced with fresh medium the next day. Infected cells on sapphire discs were fixed by using

HPF with a Compact 01 high-pressure freezer (Engineering Office M. Wohlwend GmbH,

Sennwald, Switzerland) at 5 dpi. Thereafter, cells on sapphire discs were processed by

freeze-substitution and subsequently embedded in Epon (Fluka, Buchs, Switzerland).

Ultrathin sections of the Epon-embedded cells were cut with an ultramicrotome (Ultracut

UCT; Leica Microsystems, Wetzlar, Germany) and placed on Formvar-coated single-slot

grids (Plano GmbH, Wetzlar, Germany). Grids were examined in a Jeol JEM-1400 (Jeol Ltd.,

Tokyo, Japan) transmission electron microscope equipped with a charge-coupled-device

(CCD) camera at an acceleration voltage of 120 kV. Fixation and embedding of infected

cells for scanning transmission electron microscopy (STEM) tomography was the same as

described for TEM. Additional sample preparation steps were conducted as described

previously (84, 85). Tomogram acquisition was conducted on a STEM Jeol JEM-2100F with

an acceleration voltage of 200 kV. Tilt series were acquired from 600 nm thin sections from

+72° to -72° with a 1.5° increment using the bright field detector. Image series were

reconstructed to tomograms by weighted back projection with the IMOD software package

(86). 3D visualization of the membrane structures was performed using Avizo lite software

(Visualization Science Group, Burlington, MA, USA) by threshold segmentation.

Western blotting. Western blotting procedures, including primary and secondary

antibodies used for detection of HA tag, UL148, ATF4, eIF2α, P-eIF2α (Ser51), the HCMV

viral nuclear antigen IE1, and the conditions used for detection of Ser51 phosphorylated

eIF2α, were carried out as described previously (8, 10). Additional antibodies used in this

study were mouse anti-GAPDH (Cat. No. 60004-1, Proteintech, Rosemont, IL,), mouse

anti-gB clone 27-180 (87) (a generous gift of William J. Britt), and anti-GFP (D5.1) XP®

Rabbit mAb #2956 (Cell Signaling Technology, Danvers, MA).

Solubility analyses. 2.0×10^5 human fibroblasts were infected at an MOI of 1 TCID$_{50}$ per

cell with TB_148^{HA} or TB_159^{HA}. The following day, cells were washed twice with PBS to

remove viral inoculum and replenished with DMEM containing 5% newborn calf serum. At

the indicated times post-infection, cells were washed once with PBS and lysed by direct

addition of 100 μL RIPA buffer [25 mM HEPES (pH 7.5), 400 mM NaCl, 0.1% SDS, 0.5%

sodium deoxycholate, 1% NP-40, supplemented with 1$\times$ protease inhibitor cocktail

(ApexBio)]. Lysates were collected and rotated at 4°C for 1 h. Insoluble material was

pelleted by centrifugation (21,000 $\times$ g, 35 min). Supernatants containing soluble material

were transferred to a fresh microfuge tube, and 33 μL of 4$\times$ Laemmli sample buffer [200

mM Tris (pH 6.8), 8% SDS, 40% glycerol, 0.08% bromophenol blue] was added to bring

final volume to 133 μL. The pellet was disrupted in 133 μL of 1X Laemmli buffer prepared

by diluting 4$\times$ Laemmli buffer in RIPA buffer. Samples were reduced by addition of beta-

mercaptoethanol (5% final, v/v) and boiled at 90°C for 10 min. 40 μL of each sample was

resolved by SDS-PAGE (12% polyacrylamide gel), transferred overnight to nitrocellulose

membrane, and immunoblotted with antibodies against HA epitope or HCMV gB.

Quantitation of secondary antibody fluorescence signals were performed using an Odyssey

CLx scanner (Li-Cor, Inc., Lincoln, NE) in auto-scan mode. For each time-point, the signals

from RIPA-soluble and insoluble bands were summed to yield total signal, and the ratio of

insoluble band signal to total signal were also reported as percent insoluble HA over total

HA signal.

Statistical analyses. Statistical analyses were carried out using GraphPad Prism 8.1.0 for

MacOS (GraphPad, Inc., San Diego, CA).

ACKNOWLEDGEMENTS.

This project was supported by NIH grants R01-AI116851 (to J.P.K.) and P30-

GM110703. Its contents are solely the responsibility of the authors and do not necessarily

represent the official views of the NIAID or the NIGMS.

We are especially grateful to Erik L. Snapp (HHMI, Janelia Laboratories) for helpful

discussions and advice on interpretation of EM and confocal image data. We also thank

the following individuals for generously sharing reagents: Dong Yu (Washington University,

St. Louis, MO, currently GSK Vaccines), Thomas E. Shenk (Princeton University,

Princeton, NJ), Richard J. Stanton (Cardiff University, United Kingdom), William J. Britt

(University of Alabama, Birmingham), W. L. William Chang and Peter A. Barry (both of the

University of California, Davis; Davis, CA), Christian Sinzger (University Medical Center

Ulm, Ulm, Germany), Klaus Früh (Oregon Health Sciences University, Beaverton, OR), and

Gregory A. Smith (Northwestern University, Chicago, IL).

830 AUTHOR CONTRIBUTIONS.

831   Performed experiments: HZ, CR, CCN, MNAS, JvE. Data analysis and interpretation: HZ,

832   CR, CCN, JvE, CS, JPK. Designed experiments: HZ, CR, CCN, JvE, and JPK. Contributed

833   new reagents: HZ, CH, CCN, JPK. Electron microscopy and STEM tomography: CR and

834   JvE. Obtained funding: JPK. Wrote the manuscript: JPK with comments from JvE and CR.

837
838
FIG 1. UL148 reorganizes ER markers into anomalous structures during HCMV infection.
Fibroblasts infected at MOI 1 TCID$_{50}$/cell with either **(A)** wildtype HCMV strain TB40/E (TB_WT) or
(B) an isogenic *UL148*-null mutant (TB_148$_{STOP}$), were fixed at 96 h postinfection (hpi) and imaged
by confocal microscopy after staining with antibodies specific for calnexin or Hrd1 (magenta) and
glycoprotein H (gH, green), as indicated. DAPI (blue) signal was used to counterstain nuclei. **(C)**
Percentage of fibroblasts that contain Hrd1 structures at 96 hpi; fifty gH-positive cells for each
condition were scored for the presence or absence of UL148 structures (also see SI Fig S1A). **(D)**
3D confocal image projection of a TB_WT infected fibroblast stained at 96 hpi for Hrd1 and gH. **(E)**
Volumetric measurements were made from sixteen infected cells, stained with phalloidin-AlexaFluor
594 conjugate to estimate total cell volume, with Hrd1 antibody to estimate the volume of UL148-
dependent ER structures, and with DAPI to estimate nuclear volume.

FIG 2. Characterization of TB_148^(HA) and TB_159^(HA) viruses. (A) Human fibroblasts were infected at MOI 1 with the indicated recombinant HCMVs. Cell lysate samples harvested at the indicated times postinfection (h postinfection, hpi) were analyzed by Western blot to detect HA-tagged UL148 or Rh159, the 72 kD viral nuclear antigen IE1-72 (IE1), and GAPDH. **(B)** Single-cycle viral replication kinetic curves from MOI 1 infected fibroblasts were plotted by determining the titer in tissue culture infectious dose 50 ($TCID_{50}$) from supernatants collected at the indicated times postinfection.

FIG 3. UL148 localizes to unusual ER compartments that are enriched for glycoprotein quality control markers. Fibroblasts infected at MOI 1 with either TB_148HA (panels: **A, C, E, G, I, K, M, O, Q, S, U**) or TB_ 159HA (panels: **B, D, F, H, J, L, N, P, R, T, V**) were fixed at 96 h postinfection, and imaged by confocal microscopy after co-staining with antibodies specific for HA (UL148 / Rh159, magenta in merge) and the indicated cellular markers (green in merge). DAPI (blue) counterstaining is shown in merged image.

871
872
873 **FIG 4. Quantification of co-localization between UL148 and cellular markers.** A Pearson's
874 correlation coefficient (*r*) was calculated using NIH ImageJ software to estimate the degree of co-
875 localization between UL148 (HA signal) and each of the indicated cellular markers. A minimum of 30
876 cells were analyzed per marker. The arithmetic mean for each co-localization analysis result is shown
877 as a blue line, and data points for individual cells analyzed are plotted as circles.
878

879
880
FIG 5. TEM of ER structures in wild-type HCMV infected cells. Human fibroblasts infected with
wildtype HCMV (TB_WT) were fixed by high-pressure freezing and freeze substitution at day 5
postinfection and imaged using TEM. Panels (**A**) and (**B**) show cell overview at left. For each cell,
the boxed region is shown at higher magnification. Scale bars; left panels: 10 µm, right panels: 2
µm. rER: rough ER; Nu: nucleus; cVAC: cytoplasmic viral assembly compartment. Solid
arrowheads indicate the UL148-dependent ER structures of interest.

887
888
FIG 6. TEM of *UL148*-null HCMV infected cells. Human fibroblasts infected with a *UL148*-null
mutant (TB_148$_{STOP}$) were fixed by high-pressure freezing and freeze substitution at day 5
postinfection and imaged using TEM. At the upper left an overview panel of a representative cell is
shown with panels (**A**), (**B**), and (**C**) each boxed. The boxed regions are expanded at higher
magnification at right (**A**) and below (**B, C**). Scale bars for upper left overview panel: 10 μm, for
zoomed panels: 2 μm. rER: rough ER; Nu: nucleus; cVAC: cytoplasmic viral assembly
compartment.
896

897
898
FIG 7. STEM tomography of UL148-dependent ER structures in HCMV infected cells. Human
fibroblasts infected with wildtype HCMV (TB_WT) were fixed by high-pressure freezing and freeze
substitution at day 5 postinfection and tomograms were recorded by STEM. (**A**) Shows a virtual
section through the tomogram of a virus-induced membranous structure; asterisks denote ER
structures of distended lumenal space. White arrowheads indicate sites at which membranes
originating from distended ER cisternae continue into areas of involuted collapsed ER. Scale bar: 1
μm. (**B**) Shows the same virtual section as in (A) tilted and with a 3D visualization of the
membranous network (green) of the entire tomogram. The same distended ER cisternae as in (A)
are marked by asterisks. (**C**) Shows a cross section through (B) to visualize the membrane profile
of the membranous structures. The region delimited by a white box is shown in a higher
magnification on the right. Finer detail of the enlarged ER cisternae and the connections between
them are readily visible; asterisks indicate the same distended ER cisternae as in panels (A) and
(B). Also, see SI Movie S1. Scale bar: 1 μm.

912
913
FIG 8. Solubility analysis of UL148 and Rh159. (**A**) Human fibroblasts were infected at MOI 1
TCID$_{50}$/cell with HCMV strain TB40/E derived viruses TB_148^{HA}, which expresses UL148 fused at
its C-terminus to an HA-epitope tag, or TB_159^{HA}, which lacks *UL148* and instead expresses
rhesus CMV Rh159 carrying a C-terminal HA tag. At the indicated times postinfection (hpi),
infected cells were collected in radioimmunoprecipitation (RIPA) lysis buffer, centrifuged at 21,000 $\times$
g for 30 min, after which supernatant (S) and pellet (P) fractions were boiled in gel loading buffer
containing 2% sodium dodecyl sulfate (SDS). Equivalent portions of supernatant and pellet were
resolved by SDS-polyacrylamide gel electrophoresis, transferred to a nitrocellulose membrane for
detection of protein species immunoreactive to antibodies against HA epitope (HA) and HCMV
glycoprotein B (gB) and for total protein signal using Ponceau S reagent (Ponceau). (**B-C**). Signal
intensity of fluorophore-conjugated secondary antibodies in anti-HA Western blots were measured
from three independent biological replicates of the experiment shown in (A); error bars indicate
standard deviation. (**B**) The fluorescent signal for each infection time point condition (y-axis is in
arbitrary units, in hundred thousands). (**C**) The amount of signal found in the insoluble (pellet)
fraction relative to the total signal (pellet plus supernatant) for each infection time point are plotted
as percentage values.
930
931

FIG 9. ISR activation accompanies redistribution of ER markers during ectopic expression of UL148. (A) Lysates of 'tet-on' ARPE-19 cells expressing either UL148 or Rh159 fused to an HA tag, i148HA and i159HA, respectively, were collected at the indicated times post doxycycline (dox) induction and analyzed by Western blot for the expression of the indicated proteins and for the abundance of eIF2α phosphorylated at Ser51 (P-eIF2α) using a phospho-specific antibody. **(B)** Pearson's correlation coefficient (*r*) values were calculated using NIH ImageJ software estimate the degree of co-localization between UL148 (HA signal) and the indicated cellular markers. A minimum of 30 cells were analyzed per marker. Arithmetic means for each co-localization analysis result are shown as blue lines, and data points for individual cells analyzed are plotted as circles.

946
947
FIG 10. Inhibition of the integrated stress response impedes UL148-mediated ER remodeling.
(A-B) "Tet-on" ARPE-19 epithelial cells that inducibly express UL148 fused to green florescent
protein (gfp), i148^{gfp}, were induced for transgene expression using 100 ng/mL doxycycline (dox) in
the presence of DMSO carrier, ISRIB (200 nM), or GSK2606414 (1.1 µM) and monitored by anti-
UL148 Western blot for expression of UL148-GFP over a series of time points (h post treatment
with dox, hpt). **(C-E)** Live cell imaging of UL148-GFP expression patterns in the presence of ISRIB
(200 nM), GSK2606414 (1.1 µM), or DMSO vehicle (0.01%). Main scale bar represents 50 µm. For
inset panels at upper left of each image, which are magnified 2.4× relative to the main image, the
scale bar represents 10 µm. Also see SI Movies S8-S10.
957

958
959
FIG 11. Inhibition of the ISR prevents the coalescence of Hrd1 and UL148 into discrete
structures during infection. (**A**) Representative 3D maximum intensity projections of confocal
imaging Z-stacks obtained from cells infected at MOI 1 for 96 h with HCMV strain TB40/E carrying
HA-tagged *UL148* (TB_148^HA) and maintained in the presence of ISRIB (200 nM), GSK2606414 (1.1
µM), or DMSO carrier alone (0.1% vol/vol). In merged images HA signal is shown in magenta, Hrd1
in green, and DAPI counterstaining in blue. (**B**) The percent of Hrd1 antibody signal involved in
discrete structures at 96 hpi were calculated for a minimum of 30 cells per condition using Imaris x64
9.3.0 software. Statistical significance was determined using a one-way ANOVA followed by Tukey's
post-test; **** represents a P-value of <0.0001. The arithmetic mean for co-localization analysis
results are shown as blue lines with data points for individual cells analyzed plotted as circles, squares
or triangles, as indicated. (**C-D**) Western blot analyses of fibroblasts infected at MOI 1 with TB_148^HA
and maintained in the presence of ISRIB (200 nM), GSK2606414 (1.1 µM), or DMSO carrier alone
(0.01%); hpi: h post infection (hpi). Note: a phospho-specific antibody was used for detection of eIF2α
phosphorylated at Ser51 (P-eIF2α).

FIG S1: **Additional confocal microscopy results from infected cells.** (**A**) Lower magnification overview of results from FIG 1A-B; staining of gH and Hrd1 human fibroblasts 96 h postinfection (hpi) with wildtype HCMV strain TB40/E (WT) or *UL148*-null derivative TB_148$_{STOP}$. (**B**) gH and Hrd1 in human fibroblasts 96 hpi with either wildtype strain Merlin recovered from BAC pAL1393 (ME_WT) or a *UL148*-null derivative of the same virus (ME_148$_{STOP}$). (**C**): gH and Hrd1 at 96 hpi of human fibroblasts with wildtype HCMV strain TRgfp recovered from BAC-cloned TRgfp (TR_WT) or a *UL148*-null derivative of the same virus (TR_148$_{STOP}$). (**D**) Staining of the viral tegument protein UL71 and calnexin in human fibroblasts 5 days postinfection with primary clinical HCMV isolates obtained from patient throat swabs. (**E**) HA and Hrd1 in THP-1 macrophages fixed 96 hpi with HCMV strain TB40/E derivative viruses TB_148^{HA} or TB_159^{HA}. (**F**) HA and Hrd1 in ARPE-19 epithelial cells infected for 96 h with TB_148^{HA} or TB_159^{HA}. (**G**) Staining of HA and Hrd1 in ARPE-19 infected for 96 h with HCMV strain AD169 repaired for *UL131* and to which an HA-tagged *UL148* from TB40/E was restored to the native *UL148* locus (AD^{r131}_148^{HA}) versus parental AD169 repaired for *UL131* (AD^{r131}). Notes: (i) *UL131* is required for efficient infection of epithelial cells; (ii) for ADr131 we did not employ a viral marker (e.g., HA) to identify infected cells, so the appearance of syncytia and the characteristic kidney-bean shaped nucleus was used to indicate infected cells and a slightly lower magnification was used to best show these features. (**H**) HA and Hrd1 in telomerase-immortalized rhesus fibroblasts infected with BAC-derived rhesus CMV (RhCMV) strain 68-1 carrying an HA-tag at the C-terminus of Rh159.

1003
1004
1005 **FIG S2: LC3B, syntaxin-6, and gH staining in HCMV infected cells.** Human fibroblasts were
1006 infected with the indicated viruses for 96 h and then fixed, permeabilized, and stained using
1007 antibodies specific for the indicated proteins. Indirect immunofluorescence images were captured
1008 using a 63X objective on a Leica SP5 confocal microscope. gH: glycoprotein H, STX6: syntaxin-6.
1009

FIG S3: Cellular proteins involved in glycoprotein quality control are recruited with differing kinetics to UL148 ER structures. Fibroblasts infected with TB_148[HA] at 1 MOI were fixed at the indicated time points (days post infection, dpi) and imaged by confocal microscopy after staining with antibodies specific for HA (UL148, magenta), CNX (green, **A**) , Hrd1 (green, **B**) or VCP(green, **C**), DAPI (blue).

1018
1019 **FIG S4. UL148 is sufficient to remodel the ER.** Expression of HA-tagged UL148 or Rh159 was
1020 induced in "tet-on" ARPE-19 epithelial cells, i148HA and i159HA, respectively. Cells were fixed at 48
1021 h postinduction for indirect immunofluorescence staining for the indicated cellular markers (green)
1022 together with HA (magenta). Scale bar: 10 μm.

1023
1024
FIG S5. Live-cell Imaging of UL148-GFP and Rh159-GFP during induced ectopic expression.
'Tet-on' ARPE-19 epithelial cells that inducibly express either UL148 or Rh159 fused to green
florescent protein (gfp), i148^{gfp} (**A**) and i159^{gfp} (**B**), respectively, were induced for transgene
expression using 100 ng/mL doxycycline (dox) and imaged using live-cell microscopy. Images from
the selected time points (h post treatment with dox, hpt) are shown. The main scale bar represents
50 µm. For inset panels at upper left of each image, which are magnified 2.4× relative to the main
image, the scale bar represents 10 µm. Also see SI Movies S2-S3.
1032
1033
1034
1035

1036
FIG S6. FRAP. ARPE-19 cells that inducibly express Rh159-GFP, i159^{gfp} (**A**) or UL148-GFP, i148^{gfp}
(**B-C**) were doxycycline induced for 24 h (100 ng/mL doxycycline (dox). Separate regions of GFP
signal in selected cells were photobleached (405 nm laser) or left unbleached, while a third region
lacking GFP signal was chosen as a background reference were measured before and during
fluorescence recovery after photobleaching (FRAP). Note: background signal was not plotted
because values were resolvable from the x-axis. GFP signal intensity is plotted over a time period
(seconds, s) starting with an exposure at t=0 (immediately before photobleaching), and including
measurements taken every 2 s after photobleaching (0-20 s) until termination of the measurement
series at t=322 s (300 s of FRAP). (**D-F**) Images from the selected time points (s, seconds)
immediately before and after bleaching, and during fluorescence recovery period. Also see SI
Movies S4-S6.

A

h post washout 1 | 5 | 9

h post washout 13 | 17 | 21

50 μm

B

DMSO | epoxomicin
hpt 0 6 12 18 24 48 | 0 6 12 18 24 48

75 — GFP
75 — UL148
kD 20 — LC3B
37 — β-actin

C

DMSO | folimycin
hpt 6 12 18 24 48 | 6 12 18 24 48

75 — GFP
75 — UL148
kD 20 — LC3B
37 — β-actin

1048
1049
FIG S7: UL148-GFP is stabilized by inhibition of proteasomal but not lysosomal degradation.
"Tet-on" ARPE-19 cells that inducibly express UL148-GFP (i148[gfp]) were induced for 24 h by the
addition of 100 ng/mL doxycycline (dox), after which the dox inducing agent was washed out and
medium containing either the proteasome inhibitor epoxomicin (20 μM) or the proton pump inhibitor
folimycin (115 nM) was added and samples were harvested for Western blot analysis at the
indicated times post treatment (h post treatment, hpt) with folimycin or epoxomicin. DMSO was
added at 0.1% to control for folimycin, or 1% to control for epoxomicin. Also see SI Movie S7.
1057

TABLE S1. Oligonucleotide primers and synthetic DNAs.

Name	Sequence (5' to 3')
148_eGFP_Fw	ACCTGGTGTCGCGGCGTCGGATGGTGAGCAAGGGGGAGGAGCTTTTTACTGGCGTTGTTC
Gibs_eGFP_Rv	CGGCCGCCACTGTGCTGGATTACTTATACAGCTCGTCCATACCCAG
Gib_148_Fw	TGTGGTGGAATTCTGCAGATTGCCACCATGTTGCGCTTGCTG
148_noStopRv	CCGACGCCGCGACACCAGGTAGGTTATC
159_eGFP_Fw	TCTAAAACGCAGTCGTGAAGCTCATATGGTGAGCAAGGGGGAGGA
159_noStop_Rv	ATGAGCTTCACGACTGCGTTTTAGAAGTG
Gibs_159_Fw	TGTGGTGGAATTCTGCAGATTGCCACCATGGCCTACAACAGC
Rh159_Fw	TGTGGTGGAATTCTGCAGATACCATGGCCTACAACAGCTTCCTTCTCAGCTGCCTTACCATCGCAC TTTTACTG
Rh159_HA_Rv	CGGCCGCCACTGTGCTGGATCTATTAGGCGTAGTCTGGGACGTCGTATGGGTAATGAGCTTCACG ACTGCGTTTTAGAAGTG
PpuISceKanGibs_Fw	CATTCACCTGAACCCATAGGTCCTGCCTGGCAGAGAATCTTAGGGATAACAGGGTAATCG
PpuISceKanGibs _Rv	AGATTCTCTGCCAGGCAGGACCTATGGGTTCAGGTGAATGGCCAGTGTTACAACCAATTA
Rh159_Fw_recomb	GGGCGGTGGCCGGCACGCCGCATTTCCTAACCCGCGCAGCATGGCCTACAACAGCTTCCT
Rh159_Rv_recomb	CGCGAACGACGTGTGACGAGGACGTGGTTTCCGCAAGCCTTTAGGCGTAGTCTGGGACG
TB_159HAseq_F	AGTGTTTATGAGTCGGGCGG
TB_159HAseq_R	GTGGCGACGTGGATTTCTTG
EGFP-P2A-3XHA	GTCGACTGGATCCGGTACCGAATTCGCCACCATGGTGAGCAAGGGGGAGGAGCTTTTTACTGGC GTTGTTCCCATCCTGGTTGAGTTGGATGGGGATGTCAACGGCCACAAGTTCAGCGTTTCTGGAGA AGGAGAAGGCGACGCAACCTATGGGAAGCTCACTTTGAAATTCATATGCACCACCGGTAAACTCC CAGTCCCCTGGCCTACCCTGGTCACTACCTTGACATATGGAGTTCAGTGCTTTTCTCGATACCCCG ACCACATGAAGCAGCACGATTTTTTTCAAATCTGCAATGCCCGAGGGCTATGTACAAGAGAGGACCA TTTTCTTCAAAGACGACGGCAATTACAAGACCAGGGCCGAGGTCAAATTTGAAGGGGACACTCTTG TGAATCGGATCGAACTTAAAGGGATCGATTTCAAAGAGGACGGAAACATACTGGGGCATAAGCTG GAATACAACTACAATAGCCATAATGTTTACATTATGGCAGACAAGCAGAAAAACGGTATTAAGGTCA ACTTCAAGATACGGCCATAACATTGAAGATGGGTCAGTCCAGCTTGCTGACCATTATCAGCAGAATA CACCCATTGGCGATGGTCCCGTGCTTTTGCCTGACAACCACTACCTTTCTACCCAAAGCGCATTGT CAAAAGATCCTAACGAGAAGAGAGATCATATGGTCCTTTTGGAGTTCGTAACCGCAGCTGGTATAA CTCTGGGTATGGACGAGCTGTATAAGGGCTCAGGGGCTACAAATTTTAGTCTCCTTAAACAAGCCG GAGATGTCGAAGAGAACCCTGGGCCTTACCCATATGACGTTCCTGACTATGCCGGTTACCCCTAT GATGTGCCAGATTATGCCGGATCTTATCCTTACGATGTACCTGACTATGCAGGTGGGAGTGGCAT GAATTACGTGGGGCAGC
Gibs_Age_159_Fw	CCGTACCACTTCCTACCCTCGTAAACCGGTGCCACCATGGCCTACAACAG
Gibs_Mlu_GFP_Rv	ACCCGGCGCGGAGGCCAGATCTTAACGCGTTTACTTATACAGCTCGTCCATACC
tRFP linker Fw	CCACTTCCTACCCTCGTAAACCGGTGAATTCGCCACCATGAGCGAGCTGAT
tRFP linker Rv	GCGCGGAGGCCAGATCTTAACGCGTGCGGCCGCTCGAGAATTCATCTGTGCCCCAGTTTGC

1071
1072

TABLE S2. Antibodies Used in Immunofluorescence Microscopy

Primary Antibody	Host species	Isotype	Clone	Vendor	Catalog No.	Dilution used
Calnexin	Mouse	mAb IgG1	37	BD Biosciences	610523	1:50
Calnexin	Rabbit	mAb IgG	C5C9	Cell Signaling Tech.	2679	1:50
Syntaxin 6	Rabbit	mAb IgG	C34B2	Cell Signaling Tech.	2869	1:50
SYVN1 (Hrd1)	Rabbit	mAb IgG	D3O2A	Cell Signaling Tech.	14773	1:500
SEL1L	Rabbit	pAb	n/a	Sigma	S3699	1:50
KDEL	Mouse	IgG2a	10C3	Enzo Life Sci., Inc.	ADI-SPA-827-D	1:50
PDI	Rabbit	mAb IgG	C81H6	Cell Signaling Tech.	3501	1:50
Herp (HerpUD1)	Rabbit	pAb IgG	n/a	Proteintech	10813-1-AP	1:50
VCP	Mouse	IgG2a	5	Invitrogen / Life Technologies	MA3-004	1:50
Calreticulin	Rabbit	mAb IgG	D3E6	Cell Signaling Tech.	12238	1:200
Ribophorin 1 (RPN1)	Rabbit	pAb IgG		ThermoFisher / Invitrogen	PA5-65662	1:50
EDEM-1	Mouse	mAb IgM	D-1	Santa Cruz	sc-377394	1:50
GABARAP	Rabbit	mAb	E1J4E	Cell Signaling Tech.	13733	1:200
LC3B	Rabbit	mAb	D11	Cell Signaling Tech.	3868	1:200
HA epitope	Chicken	IgY		Bethyl	A190-106A	1:200
HCMV glycoprotein H (gH)	Mouse	IgG2b	14-4b	Gift of William J. Britt, M.D.	n/a	1: 50
Reticulon-3 (RTN3)	Mouse	IgG2a		ThermoFisher / Invitrogen	MA5-15538	1: 200
HA epitope	Rabbit	pAb		Bethyl	A190-108A	1:1000
Secondary Antibody	**Host species**	**Isotype**		**Vendor**	**Catalog No.**	**Dilution used**
Alexa Fluor 488 anti-Mouse IgG (H+L), cross adsorbed	Goat	IgG		ThermoFisher / Invitrogen	A11001	1:1000
Alexa Fluor 488 anti-Rabbit IgG (H+L) Cross-Adsorbed	Goat	IgG		ThermoFisher / Invitrogen	A11008	1:1000
Alexa Fluor 594 anti-Rabbit IgG (H+L) Cross-Adsorbed	Goat	IgG		ThermoFisher / Invitrogen	A11012	1:1000
Alexa Fluor 594 anti-Mouse IgG (H+L) Cross-Adsorbed	Goat	IgG		ThermoFisher / Invitrogen	A11005	1:1000
Alexa Fluor 647 anti-Chicken IgY (H+L)	Goat	IgG		ThermoFisher / Invitrogen	A11039	1:1000

1073
1074
1075

1076
1077
1078 **SUPPLEMENTARY VIDEO FILES.**
1079
Movie S1: STEM tomography of UL148 ER structures in wildtype HCMV infected fibroblasts at 5
dpi.
Movie S2: i148^{gfp} cells from 2-19 h post dox induction.
Movie S3: i159^{gfp} cells from 2-19 h post dox induction.
Movie S4: FRAP of UL148-GFP signal within ER structure.
Movie S5: FRAP of UL148-GFP signal outside ER structure.
Movie S6: FRAP of Rh159-GFP signal.
Movie S7: i148^{gfp} cells from 30 min to 22 h 07 min post washout of doxycycline (dox).
Movie S8: i148^{gfp} cells in the presence of 0.01% DMSO, from 2-19 h post dox induction.
Movie S9: i148^{gfp} cells in the presence of 1.1 µM GSK2606414, from 2-19 h post dox induction.
Movie S10: i148^{gfp} cells in the presence of 200 nM ISRIB, from 2-19 h post dox induction.
1091
1092

1093

1094

REFERENCES.

1. Li G, Nguyen CC, Ryckman BJ, Britt WJ, Kamil JP. 2015. A viral regulator of glycoprotein complexes contributes to human cytomegalovirus cell tropism. Proc Natl Acad Sci U S A 112:4471-6.
2. Wille PT, Knoche AJ, Nelson JA, Jarvis MA, Johnson DC. 2010. A human cytomegalovirus gO-null mutant fails to incorporate gH/gL into the virion envelope and is unable to enter fibroblasts and epithelial and endothelial cells. J Virol 84:2585-96.
3. Jiang XJ, Adler B, Sampaio KL, Digel M, Jahn G, Ettischer N, Stierhof YD, Scrivano L, Koszinowski U, Mach M, Sinzger C. 2008. UL74 of human cytomegalovirus contributes to virus release by promoting secondary envelopment of virions. J Virol 82:2802-12.
4. Zhou M, Lanchy JM, Ryckman BJ. 2015. Human Cytomegalovirus gH/gL/gO Promotes the Fusion Step of Entry into All Cell Types, whereas gH/gL/UL128-131 Broadens Virus Tropism through a Distinct Mechanism. J Virol 89:8999-9009.
5. Wu Y, Prager A, Boos S, Resch M, Brizic I, Mach M, Wildner S, Scrivano L, Adler B. 2017. Human cytomegalovirus glycoprotein complex gH/gL/gO uses PDGFR-alpha as a key for entry. PLoS Pathog 13:e1006281.
6. Kabanova A, Marcandalli J, Zhou T, Bianchi S, Baxa U, Tsybovsky Y, Lilleri D, Silacci-Fregni C, Foglierini M, Fernandez-Rodriguez BM, Druz A, Zhang B, Geiger R, Pagani M, Sallusto F, Kwong PD, Corti D, Lanzavecchia A, Perez L. 2016. Platelet-derived growth factor-alpha receptor is the cellular receptor for human cytomegalovirus gHgLgO trimer. Nat Microbiol 1:16082.
7. Wu K, Oberstein A, Wang W, Shenk T. 2018. Role of PDGF receptor-alpha during human cytomegalovirus entry into fibroblasts. Proc Natl Acad Sci U S A doi:10.1073/pnas.1806305115.
8. Nguyen CC, Siddiquey MNA, Zhang H, Li G, Kamil JP. 2018. Human Cytomegalovirus Tropism Modulator UL148 Interacts with SEL1L, a Cellular Factor That Governs Endoplasmic Reticulum-Associated Degradation of the Viral Envelope Glycoprotein gO. J Virol 92:e00688-18.
9. Wang ECY, Pjechova M, Nightingale K, Vlahava VM, Patel M, Ruckova E, Forbes SK, Nobre L, Antrobus R, Roberts D, Fielding CA, Seirafian S, Davies J, Murrell I, Lau B, Wilkie GS, Suarez NM, Stanton RJ, Vojtesek B, Davison A, Lehner PJ, Weekes MP, Wilkinson GWG, Tomasec P. 2018. Suppression of costimulation by human cytomegalovirus promotes evasion of cellular immune defenses. Proc Natl Acad Sci U S A 115:4998-5003.
10. Siddiquey MNA, Zhang H, Nguyen CC, Domma AJ, Kamil JP. 2018. The Human Cytomegalovirus Endoplasmic Reticulum-Resident Glycoprotein UL148 Activates the Unfolded Protein Response. J Virol 92:e00896-18.
11. Sinzger C, Hahn G, Digel M, Katona R, Sampaio KL, Messerle M, Hengel H, Koszinowski U, Brune W, Adler B. 2008. Cloning and sequencing of a highly productive, endotheliotropic virus strain derived from human cytomegalovirus TB40/E. J Gen Virol 89:359-68.
12. Das S, Vasanji A, Pellett PE. 2007. Three-dimensional structure of the human cytomegalovirus cytoplasmic virion assembly complex includes a reoriented secretory apparatus. J Virol 81:11861-9.
13. Sanchez V, Greis KD, Sztul E, Britt WJ. 2000. Accumulation of virion tegument and envelope proteins in a stable cytoplasmic compartment during human cytomegalovirus replication: characterization of a potential site of virus assembly. J Virol 74:975-86.

14. Lilley BN, Ploegh HL. 2005. Multiprotein complexes that link dislocation, ubiquitination, and
extraction of misfolded proteins from the endoplasmic reticulum membrane. Proc Natl Acad
Sci U S A 102:14296-301.
15. Kamhi-Nesher S, Shenkman M, Tolchinsky S, Fromm SV, Ehrlich R, Lederkremer GZ. 2001. A
novel quality control compartment derived from the endoplasmic reticulum. Mol Biol Cell
12:1711-23.
16. Leitman J, Shenkman M, Gofman Y, Shtern NO, Ben-Tal N, Hendershot LM, Lederkremer GZ.
2014. Herp coordinates compartmentalization and recruitment of HRD1 and misfolded
proteins for ERAD. Mol Biol Cell 25:1050-60.
17. Houck SA, Ren HY, Madden VJ, Bonner JN, Conlin MP, Janovick JA, Conn PM, Cyr DM. 2014.
Quality control autophagy degrades soluble ERAD-resistant conformers of the misfolded
membrane protein GnRHR. Mol Cell 54:166-179.
18. Sturgill ER, Malouli D, Hansen SG, Burwitz BJ, Seo S, Schneider CL, Womack JL, Verweij MC,
Ventura AB, Bhusari A, Jeffries KM, Legasse AW, Axthelm MK, Hudson AW, Sacha JB, Picker
LJ, Fruh K. 2016. Natural Killer Cell Evasion Is Essential for Infection by Rhesus
Cytomegalovirus. PLoS Pathog 12:e1005868.
19. Chaumorcel M, Souquere S, Pierron G, Codogno P, Esclatine A. 2008. Human
cytomegalovirus controls a new autophagy-dependent cellular antiviral defense
mechanism. Autophagy 4:46-53.
20. Chaumorcel M, Lussignol M, Mouna L, Cavignac Y, Fahie K, Cotte-Laffitte J, Geballe A, Brune
W, Beau I, Codogno P, Esclatine A. 2012. The human cytomegalovirus protein TRS1 inhibits
autophagy via its interaction with Beclin 1. J Virol 86:2571-84.
21. Alwine JC. 2012. The human cytomegalovirus assembly compartment: a masterpiece of viral
manipulation of cellular processes that facilitates assembly and egress. PLoS Pathog
8:e1002878.
22. Das S, Pellett PE. 2011. Spatial relationships between markers for secretory and endosomal
machinery in human cytomegalovirus-infected cells versus those in uninfected cells. J Virol
85:5864-79.
23. Leitman J, Ron E, Ogen-Shtern N, Lederkremer GZ. 2013. Compartmentalization of
endoplasmic reticulum quality control and ER-associated degradation factors. DNA Cell Biol
32:2-7.
24. Cox DW, Billingsley GD, Callahan JW. 1986. Aggregation of plasma Z type alpha 1-antitrypsin
suggests basic defect for the deficiency. FEBS Lett 205:255-60.
25. Le A, Ferrell GA, Dishon DS, Le QQ, Sifers RN. 1992. Soluble aggregates of the human PiZ
alpha 1-antitrypsin variant are degraded within the endoplasmic reticulum by a mechanism
sensitive to inhibitors of protein synthesis. J Biol Chem 267:1072-80.
26. Lomas DA, Evans DL, Finch JT, Carrell RW. 1992. The mechanism of Z alpha 1-antitrypsin
accumulation in the liver. Nature 357:605-7.
27. Ordonez A, Snapp EL, Tan L, Miranda E, Marciniak SJ, Lomas DA. 2013. Endoplasmic
reticulum polymers impair luminal protein mobility and sensitize to cellular stress in alpha1-
antitrypsin deficiency. Hepatology 57:2049-60.
28. Ekeowa UI, Freeke J, Miranda E, Gooptu B, Bush MF, Perez J, Teckman J, Robinson CV,
Lomas DA. 2010. Defining the mechanism of polymerization in the serpinopathies. Proc Natl
Acad Sci U S A 107:17146-51.

29. Suhy DA, Giddings TH, Jr., Kirkegaard K. 2000. Remodeling the endoplasmic reticulum by
poliovirus infection and by individual viral proteins: an autophagy-like origin for virus-
induced vesicles. J Virol 74:8953-65.
30. Windsor M, Hawes P, Monaghan P, Snapp E, Salas ML, Rodriguez JM, Wileman T. 2012.
Mechanism of collapse of endoplasmic reticulum cisternae during African swine fever virus
infection. Traffic 13:30-42.
31. Miller S, Krijnse-Locker J. 2008. Modification of intracellular membrane structures for virus
replication. Nat Rev Microbiol 6:363-74.
32. Siddiquey MNA, Zhang H, Nguyen CC, Domma AJ, Kamil JP. 2018. The human
cytomegalovirus ER resident glycoprotein UL148 activates the unfolded protein response. J
Virol doi:10.1128/JVI.00896-18.
33. Cormack BP, Valdivia RH, Falkow S. 1996. FACS-optimized mutants of the green fluorescent
protein (GFP). Gene 173:33-8.
34. Khaminets A, Heinrich T, Mari M, Grumati P, Huebner AK, Akutsu M, Liebmann L, Stolz A,
Nietzsche S, Koch N, Mauthe M, Katona I, Qualmann B, Weis J, Reggiori F, Kurth I, Hubner
CA, Dikic I. 2015. Regulation of endoplasmic reticulum turnover by selective autophagy.
Nature 522:354-8.
35. Meng L, Mohan R, Kwok BH, Elofsson M, Sin N, Crews CM. 1999. Epoxomicin, a potent and
selective proteasome inhibitor, exhibits in vivo antiinflammatory activity. Proc Natl Acad Sci
U S A 96:10403-8.
36. Muroi M, Shiragami N, Nagao K, Yamasaki M, Takatsuki A. 1993. Folimycin (concanamycin
A), a specific inhibitor of V-ATPase, blocks intracellular translocation of the glycoprotein of
vesicular stomatitis virus before arrival to the Golgi apparatus. Cell Struct Funct 18:139-49.
37. Muroi M, Takasu A, Yamasaki M, Takatsuki A. 1993. Folimycin (concanamycin A), an
inhibitor of V-type H(+)-ATPase, blocks cell-surface expression of virus-envelope
glycoproteins. Biochem Biophys Res Commun 193:999-1005.
38. Kondratyev M, Avezov E, Shenkman M, Groisman B, Lederkremer GZ. 2007. PERK-
dependent compartmentalization of ERAD and unfolded protein response machineries
during ER stress. Exp Cell Res 313:3395-407.
39. Harding HP, Novoa I, Zhang Y, Zeng H, Wek R, Schapira M, Ron D. 2000. Regulated
translation initiation controls stress-induced gene expression in mammalian cells. Mol Cell
6:1099-108.
40. Palam LR, Baird TD, Wek RC. 2011. Phosphorylation of eIF2 facilitates ribosomal bypass of
an inhibitory upstream ORF to enhance CHOP translation. J Biol Chem 286:10939-49.
41. Vattem KM, Wek RC. 2004. Reinitiation involving upstream ORFs regulates ATF4 mRNA
translation in mammalian cells. Proc Natl Acad Sci U S A 101:11269-74.
42. Sidrauski C, Acosta-Alvear D, Khoutorsky A, Vedantham P, Hearn BR, Li H, Gamache K,
Gallagher CM, Ang KK, Wilson C, Okreglak V, Ashkenazi A, Hann B, Nader K, Arkin MR,
Renslo AR, Sonenberg N, Walter P. 2013. Pharmacological brake-release of mRNA
translation enhances cognitive memory. Elife 2:e00498.
43. Sidrauski C, McGeachy AM, Ingolia NT, Walter P. 2015. The small molecule ISRIB reverses
the effects of eIF2alpha phosphorylation on translation and stress granule assembly. Elife 4.
44. Zyryanova AF, Weis F, Faille A, Alard AA, Crespillo-Casado A, Sekine Y, Harding HP, Allen F,
Parts L, Fromont C, Fischer PM, Warren AJ, Ron D. 2018. Binding of ISRIB reveals a
regulatory site in the nucleotide exchange factor eIF2B. Science 359:1533-1536.

45.   Sidrauski C, Tsai JC, Kampmann M, Hearn BR, Vedantham P, Jaishankar P, Sokabe M,
Mendez AS, Newton BW, Tang EL, Verschueren E, Johnson JR, Krogan NJ, Fraser CS,
Weissman JS, Renslo AR, Walter P. 2015. Pharmacological dimerization and activation of the
exchange factor eIF2B antagonizes the integrated stress response. Elife 4:e07314.
46.   Tsai JC, Miller-Vedam LE, Anand AA, Jaishankar P, Nguyen HC, Renslo AR, Frost A, Walter P.
2018. Structure of the nucleotide exchange factor eIF2B reveals mechanism of memory-
enhancing molecule. Science 359.
47.   Gordiyenko Y, Schmidt C, Jennings MD, Matak-Vinkovic D, Pavitt GD, Robinson CV. 2014.
eIF2B is a decameric guanine nucleotide exchange factor with a gamma2epsilon2 tetrameric
core. Nat Commun 5:3902.
48.   Harding HP, Zhang Y, Ron D. 1999. Protein translation and folding are coupled by an
endoplasmic-reticulum-resident kinase. Nature 397:271-4.
49.   Harding HP, Zhang Y, Bertolotti A, Zeng H, Ron D. 2000. Perk is essential for translational
regulation and cell survival during the unfolded protein response. Mol Cell 5:897-904.
50.   Axten JM, Medina JR, Feng Y, Shu A, Romeril SP, Grant SW, Li WH, Heerding DA, Minthorn E,
Mencken T, Atkins C, Liu Q, Rabindran S, Kumar R, Hong X, Goetz A, Stanley T, Taylor JD,
Sigethy SD, Tomberlin GH, Hassell AM, Kahler KM, Shewchuk LM, Gampe RT. 2012.
Discovery of 7-methyl-5-(1-{[3-(trifluoromethyl)phenyl]acetyl}-2,3-dihydro-1H-indol-5-yl)-
7H-p yrrolo[2,3-d]pyrimidin-4-amine (GSK2606414), a potent and selective first-in-class
inhibitor of protein kinase R (PKR)-like endoplasmic reticulum kinase (PERK). J Med Chem
55:7193-207.
51.   Tenorio R, Fernandez de Castro I, Knowlton JJ, Zamora PF, Lee CH, Mainou BA, Dermody TS,
Risco C. 2018. Reovirus sigmaNS and muNS Proteins Remodel the Endoplasmic Reticulum to
Build Replication Neo-Organelles. MBio 9.
52.   Miller S, Kastner S, Krijnse-Locker J, Buhler S, Bartenschlager R. 2007. The non-structural
protein 4A of dengue virus is an integral membrane protein inducing membrane alterations
in a 2K-regulated manner. J Biol Chem 282:8873-82.
53.   Lamond AI, Sleeman JE. 2003. Nuclear substructure and dynamics. Curr Biol 13:R825-8.
54.   Granell S, Baldini G, Mohammad S, Nicolin V, Narducci P, Storrie B, Baldini G. 2008.
Sequestration of mutated alpha1-antitrypsin into inclusion bodies is a cell-protective
mechanism to maintain endoplasmic reticulum function. Mol Biol Cell 19:572-86.
55.   Lawless MW, Greene CM, Mulgrew A, Taggart CC, O'Neill SJ, McElvaney NG. 2004.
Activation of endoplasmic reticulum-specific stress responses associated with the
conformational disease Z alpha 1-antitrypsin deficiency. J Immunol 172:5722-6.
56.   Tirosh B, Iwakoshi NN, Lilley BN, Lee AH, Glimcher LH, Ploegh HL. 2005. Human
cytomegalovirus protein US11 provokes an unfolded protein response that may facilitate
the degradation of class I major histocompatibility complex products. J Virol 79:2768-79.
57.   Wiertz EJ, Jones TR, Sun L, Bogyo M, Geuze HJ, Ploegh HL. 1996. The human
cytomegalovirus US11 gene product dislocates MHC class I heavy chains from the
endoplasmic reticulum to the cytosol. Cell 84:769-79.
58.   van den Boomen DJ, Timms RT, Grice GL, Stagg HR, Skodt K, Dougan G, Nathan JA, Lehner
PJ. 2014. TMEM129 is a Derlin-1 associated ERAD E3 ligase essential for virus-induced
degradation of MHC-I. Proc Natl Acad Sci U S A 111:11425-30.
59.   Caramelo JJ, Parodi AJ. 2008. Getting in and out from calnexin/calreticulin cycles. J Biol
Chem 283:10221-5.

60.   Hammond C, Braakman I, Helenius A. 1994. Role of N-linked oligosaccharide recognition,
1276   glucose trimming, and calnexin in glycoprotein folding and quality control. Proc Natl Acad
1277   Sci U S A 91:913-7.
1278   61.   Zhou M, Yu Q, Wechsler A, Ryckman BJ. 2013. Comparative analysis of gO isoforms reveals
1279   that strains of human cytomegalovirus differ in the ratio of gH/gL/gO and gH/gL/UL128-131
1280   in the virion envelope. J Virol 87:9680-90.
1281   62.   Zhang L, Zhou M, Stanton R, Kamil J, Ryckman BJ. 2018. Expression Levels of Glycoprotein O
1282   (gO) Vary between Strains of Human Cytomegalovirus, Influencing the Assembly of gH/gL
1283   Complexes and Virion Infectivity. J Virol 92.
1284   63.   Sifers RN, Brashears-Macatee S, Kidd VJ, Muensch H, Woo SL. 1988. A frameshift mutation
1285   results in a truncated alpha 1-antitrypsin that is retained within the rough endoplasmic
1286   reticulum. J Biol Chem 263:7330-5.
1287   64.   Choudhury P, Liu Y, Bick RJ, Sifers RN. 1997. Intracellular association between UDP-
1288   glucose:glycoprotein glucosyltransferase and an incompletely folded variant of alpha1-
1289   antitrypsin. J Biol Chem 272:13446-51.
1290   65.   Tsao YS, Ivessa NE, Adesnik M, Sabatini DD, Kreibich G. 1992. Carboxy terminally truncated
1291   forms of ribophorin I are degraded in pre-Golgi compartments by a calcium-dependent
1292   process. J Cell Biol 116:57-67.
1293   66.   de Virgilio M, Weninger H, Ivessa NE. 1998. Ubiquitination is required for the retro-
1294   translocation of a short-lived luminal endoplasmic reticulum glycoprotein to the cytosol for
1295   degradation by the proteasome. J Biol Chem 273:9734-43.
1296   67.   Johnston JA, Ward CL, Kopito RR. 1998. Aggresomes: a cellular response to misfolded
1297   proteins. J Cell Biol 143:1883-98.
1298   68.   Johnston JA, Illing ME, Kopito RR. 2002. Cytoplasmic dynein/dynactin mediates the
1299   assembly of aggresomes. Cell Motil Cytoskeleton 53:26-38.
1300   69.   Pankiv S, Alemu EA, Brech A, Bruun JA, Lamark T, Overvatn A, Bjorkoy G, Johansen T. 2010.
1301   FYCO1 is a Rab7 effector that binds to LC3 and PI3P to mediate microtubule plus end-
1302   directed vesicle transport. J Cell Biol 188:253-69.
1303   70.   Hummel T, Krukkert K, Roos J, Davis G, Klambt C. 2000. Drosophila Futsch/22C10 is a
1304   MAP1B-like protein required for dendritic and axonal development. Neuron 26:357-70.
1305   71.   Wang H, Bedford FK, Brandon NJ, Moss SJ, Olsen RW. 1999. GABA(A)-receptor-associated
1306   protein links GABA(A) receptors and the cytoskeleton. Nature 397:69-72.
1307   72.   Zhang K, Daigle JG, Cunningham KM, Coyne AN, Ruan K, Grima JC, Bowen KE, Wadhwa H,
1308   Yang P, Rigo F, Taylor JP, Gitler AD, Rothstein JD, Lloyd TE. 2018. Stress Granule Assembly
1309   Disrupts Nucleocytoplasmic Transport. Cell 173:958-971 e17.
1310   73.   Li YR, King OD, Shorter J, Gitler AD. 2013. Stress granules as crucibles of ALS pathogenesis. J
1311   Cell Biol 201:361-72.
1312   74.   Hetz C, Saxena S. 2017. ER stress and the unfolded protein response in neurodegeneration.
1313   Nat Rev Neurol 13:477-491.
1314   75.   Kirchoff V, Wong S, St JS, Pari GS. 2002. Generation of a life-expanded rhesus monkey
1315   fibroblast cell line for the growth of rhesus rhadinovirus (RRV). Arch Virol 147:321-33.
1316   76.   Smith IL, Taskintuna I, Rahhal FM, Powell HC, Ai E, Mueller AJ, Spector SA, Freeman WR.
1317   1998. Clinical failure of CMV retinitis with intravitreal cidofovir is associated with antiviral
1318   resistance. Arch Ophthalmol 116:178-85.

77. Murphy E, Yu D, Grimwood J, Schmutz J, Dickson M, Jarvis MA, Hahn G, Nelson JA, Myers
RM, Shenk TE. 2003. Coding potential of laboratory and clinical strains of human
cytomegalovirus. Proc Natl Acad Sci U S A 100:14976-81.
78. Stanton RJ, Baluchova K, Dargan DJ, Cunningham C, Sheehy O, Seirafian S, McSharry BP,
Neale ML, Davies JA, Tomasec P, Davison AJ, Wilkinson GW. 2010. Reconstruction of the
complete human cytomegalovirus genome in a BAC reveals RL13 to be a potent inhibitor of
replication. J Clin Invest 120:3191-208.
79. Hobom U, Brune W, Messerle M, Hahn G, Koszinowski UH. 2000. Fast screening procedures
for random transposon libraries of cloned herpesvirus genomes: mutational analysis of
human cytomegalovirus envelope glycoprotein genes. J Virol 74:7720-9.
80. Chang WL, Barry PA. 2003. Cloning of the full-length rhesus cytomegalovirus genome as an
infectious and self-excisable bacterial artificial chromosome for analysis of viral
pathogenesis. J Virol 77:5073-83.
81. Tischer BK, Smith GA, Osterrieder N. 2010. En passant mutagenesis: a two step markerless
red recombination system. Methods Mol Biol 634:421-30.
82. Tischer BK, von Einem J, Kaufer B, Osterrieder N. 2006. Two-step red-mediated
recombination for versatile high-efficiency markerless DNA manipulation in Escherichia coli.
Biotechniques 40:191-7.
83. Dietz AN, Villinger C, Becker S, Frick M, von Einem J. 2018. A Tyrosine-Based Trafficking
Motif of the Tegument Protein pUL71 Is Crucial for Human Cytomegalovirus Secondary
Envelopment. J Virol 92.
84. Hohn K, Sailer M, Wang L, Lorenz M, Schneider ME, Walther P. 2011. Preparation of
cryofixed cells for improved 3D ultrastructure with scanning transmission electron
tomography. Histochem Cell Biol 135:1-9.
85. Villinger C, Schauflinger M, Gregorius H, Kranz C, Hohn K, Nafeey S, Walther P. 2014. Three-
dimensional imaging of adherent cells using FIB/SEM and STEM. Methods Mol Biol
1117:617-38.
86. Kremer JR, Mastronarde DN, McIntosh JR. 1996. Computer visualization of three-
dimensional image data using IMOD. J Struct Biol 116:71-6.
87. Britt WJ, Jarvis MA, Drummond DD, Mach M. 2005. Antigenic domain 1 is required for
oligomerization of human cytomegalovirus glycoprotein B. J Virol 79:4066-79.
1350

1351