A cytomegalovirus immunevasin triggers integrated stress response-dependent

reorganization of the endoplasmic reticulum.

Running Title: HCMV UL148 reorganizes the ER.

Hongbo Zhang[a,1], Clarissa Read[d,e,1], Christopher C. Nguyen[a,1], Mohammed N.A.

Siddiquey[a], Chaowei Shang[c], Cameron M. Hall[a], Jens von Einem[d], and Jeremy P.

Kamil[a,b #]

[a]Department of Microbiology and Immunology,

[b]Center for Molecular and Tumor Virology

[c]Research Core Facility

LSU Health Sciences Center, Shreveport, Louisiana, USA

[d] Institute of Virology, Ulm University Medical Center,

[e]Central Facility for Electron Microscopy, Ulm University

Ulm, Germany

[1] these authors contributed equally to the study

Running Head: HCMV UL148 Remodels the ER

#Address correspondence to Jeremy P. Kamil, jkamil@lsuhsc.edu

ABSTRACT. Human cytomegalovirus (HCMV) encodes an ER-resident glycoprotein,

UL148, which activates the unfolded protein response (UPR) but is fully dispensable for

viral replication in cultured cells. Hence, previously ascribed roles of UL148 in immune

evasion and modulation of HCMV cell tropism are hypothesized to cause ER stress.

Here, we show that UL148 is necessary and sufficient to drive the formation of large

ER-derived structures that occupy up to 7% of the infected cell cytosol, and which

represent sites where UL148 coalesces together with ER quality control proteins such

as the E3 ubiquitin ligase Hrd1 and the ER mannosidase EDEM1. Ultrastructural

analyses of the structures reveal tortuous, densely packed segments of collapsed ER

which connect to distended cisternae. Notably, we observe that UL148 accumulates in

a detergent-insoluble form during infection while a homologous rhesus cytomegalovirus

immunevasin that fails to cause ER reorganization remains soluble. During induced

ectopic expression of UL148-GFP fusion protein, punctate signals are observed that

traffic to accumulate at prominent structures that exhibit poor recovery of fluorescence

after photobleaching. Small molecule blockade of the integrated stress response (ISR)

prevents puncta formation, leading to a uniform reticular fluorescent signal.

Accordingly, ISR inhibition during HCMV infection abolishes the coalescence of UL148

and Hrd1 into discrete structures. Given that UL148 stabilizes within the ER the

receptor binding subunit of an alternative viral envelope glycoprotein complex that is

required for infectivity, our results imply stress-dependent ER remodeling contributes to

viral cell tropism.

IMPORTANCE.

Perturbations to ER morphology occur during infection with various intracellular

pathogens and in certain genetic disorders. We identify that an HCMV gene product,

UL148, profoundly reorganizes the ER during infection, and is sufficient to do so when

expressed on its own. Although several viruses reorganize the ER, the viral proteins

necessary for such effects often play essential roles in viral replication. Although UL148

is fully dispensable for HCMV replication in cultured cells, UL148 intercepts the immune

cell co-stimulatory ligand CD58 before it reaches the cell surface and influences the

composition of viral envelope glycoprotein complexes required for cell entry. We find

that UL148 requires integrated stress response (ISR) to cause ER reorganization. Our

results reveal UL148-dependent reorganization of the ER as a prominent feature of

HCMV infected cells and provide a basis for the use of UL148 to screen for novel ISR

inhibitors.

INTRODUCTION.

UL148 is a human cytomegalovirus (HCMV) ER-resident glycoprotein that plays

roles in evasion of cell-mediated immunity and shows intriguing effects on cell tropism.

During infection of epithelial cells, viruses disrupted for *UL148* replicate to produce

roughly 100-fold enhanced levels of infectious progeny virions compared to wildtype (1).

These effects correlate with reduced expression of glycoprotein O (gO), a subunit of a

heterotrimeric viral glycoprotein H (gH) / glycoprotein L (gL) complex (gH/gL/gO) on the

virion envelope that is required for the infectivity of cell-free virions (2-4), and which

endows the virus with the capacity to utilize the platelet derived growth factor receptor α

(PDGFRα) as an entry receptor (5-7). Accordingly, UL148 has been found to stabilize

immature forms of gO prior their assembly into gH/gL/gO heterotrimers (1, 8). Despite

that UL148 does not stably associate with gO, the data suggest that it may interact with

gH (1).

UL148 also physically associates with CD58 (LFA-3), a co-stimulatory ligand for

natural killer cells and T-lymphocytes, preventing its presentation at cell surface (9).

Although the mechanisms by which UL148 stabilizes gO and retains CD58 within the

ER remain unknown, UL148 strongly contributes to activation of the unfolded protein

response (UPR) during infection, and is sufficient to activate the UPR when ectopically

expressed in non-infected cells (10). UL148 co-purifies from infected cells with SEL1L,

an adaptor subunit of ER-based E3 ubiquitin ligase Hrd1 that that plays important roles

in ER-associated degradation (ERAD) of terminally misfolded glycoproteins (8). Hence,

the potential association of UL148 with the ERAD machinery may suggest a mechanism

for its activation of the UPR.

Here, we show that expression of UL148 is necessary and sufficient to induce

unusual ER structures at which large quantities of ER factors involved in glycoprotein

quality control accumulate. In electron microscopy analyses, we find that the UL148

induced ER-structures are comprised of densely packed, tortuous ER membranes that

form connections with tubules of highly distended cisternal space. Our results may

have implications for understanding the mechanisms by which UL148 regulates viral cell

tropism and contributes to viral evasion of cell-mediated immunity. Further, our findings

suggest that ER remodeling events triggered by UL148 are indicative of an adaptive

response of eukaryotic cells to proteotoxic stress involving the secretory pathway.

RESULTS.

UL148 causes reorganization of ER quality control proteins into unusual globular structures.

The HCMV ER-resident glycoprotein UL148 was previously observed to co-localize with the ER marker calnexin during infection (1). Nonetheless, the calnexin staining did not show the uniform reticular staining pattern characteristic for the ER marker. However, we later noticed that cells infected with a *UL148*-null virus but not an isogenic wildtype control showed uniform calnexin staining (see below). To formally determine whether UL148 influences calnexin localization, we compared fibroblasts at four days post-infection with either wildtype (TB_WT) or *UL148*-null mutant (TB_148STOP) viruses derived from HCMV strain TB40/E (**FIG 1**). In cells infected with wildtype virus, calnexin antibodies stained unusual globular structures at the cell periphery, as expected (1) (**FIG 1A, 1C, SI FIG S1A**). However, in cells infected with the *UL148*-null virus, calnexin staining was uniform throughout the cytosol (**FIG 1B-C**), as would be expected for an ER marker in uninfected cells. The staining pattern for Hrd1, another ER marker, likewise indicated accumulation at unusual globular structures during wildtype HCMV infection, but not during infection with *UL148*-null mutant viruses (**FIG 1, SI FIG S1A**). Because calnexin and Hrd1 staining showed uniform distribution in *UL148*-null virus infected cells, and because *UL148* is fully dispensable for efficient viral replication in fibroblasts (1), these results suggest that redistribution of these ER markers during HCMV infection depends on UL148.

As expected (11, 12), antibodies specific for the HCMV envelope glycoprotein,

glycoprotein H (gH), stained a juxtanuclear compartment discrete from the ER, termed

the cytoplasmic virion assembly compartment (cVAC) (**FIG 1**). Localization of gH to the

cVAC was readily apparent in both wildtype and *UL148*-null virus infected cells. Similar

staining patterns for Hrd1 and/or calnexin were observed in cells infected with wildtype

or *UL148*-null mutants of clinical HCMV strains Merlin and TR, and in four primary

clinical isolates grown directly from throat swabs of infected patients (**SI, FIG S1B-D**).

Furthermore, similar differences in Hrd1 staining were observed between WT and

UL148-null viruses during infection of THP-1 macrophages and ARPE-19 epithelial cells

(**SI, FIG S1E**). Thus, the UL148-dependent effects on localization of calnexin and Hrd1

are not an anomaly particular to strain TB40/E. Nor does UL148-dependent

redistribution of ER markers occur only during infection of fibroblasts. These results

suggest that UL148-dependent redistribution of calnexin and Hrd1 occurs many if not all

cell types during wildtype HCMV infection.

We measured the three-dimensional volume of UL148-dependent Hrd1

structures from sixteen infected cells. We found that on average the structures

occupied 303.3 μm^3 (SEM: +/- 24.8) out of a total cell volume of 7657 μm^3 (SEM:+/-

651.0), as estimated by using a phalloidin-fluorophore conjugate to visualize the actin

cytoskeleton (**FIG 1E**). Subtracting the volume of nuclei, as indicated by 4′,6-diamidino-

2-phenylindole [DAPI] (average nucleus: 1360 μm^3, SEM:+/-129.4), we calculate that,

on average, the structures occupy 5.2% of the cytosolic volume (SEM: +/-0.61%, range:

2.0% – 7.0%). Based on these findings, taken together with results comparing

additional WT versus *UL148*-null mutant HCMV strains (**FIG S1**), we conclude that the

UL148-dependent ER structures are a prominent feature of HCMV-infected cells.

UL148 substantially contributes to activation of the unfolded protein response

(UPR) during HCMV infection (10), and co-purifies from infected cells with SEL1L (8),

an adaptor subunit for the E3 ubiquitin ligase Hrd1, which plays crucial roles in ER-

associated degradation (ERAD) of terminally misfolded glycoprotein substrates (13).

Hence, the accumulation of Hrd1 and calnexin at unusual structures during wildtype but

not *UL148*-null infection may suggest that the formation of these ER structures occurs in

response to defects in ER quality control (ERQC) caused by UL148. Under conditions

of proteasome inhibition, overexpression of certain misfolded glycoproteins causes

cellular factors involved in ERQC, such as calnexin to compartmentalize from the rest of

the ER, while other ER markers such as BiP (Grp78) or protein disulfide isomerase

remain largely unaltered (PDI) (14-16).

To gain further insights into the nature of these peculiar ER structures, we set out

to develop a more comprehensive understanding of their protein composition by

comparing the localization of selected ER markers in cells fixed at 96 h post-infection

(hpi) either with TB_148^{HA} (1), a derivative of HCMV strain TB40/E that expresses at the

C-terminus of UL148 the nonapeptide epitope YPYDVPDYA from influenza A

hemagglutinin (HA), or with TB_159^{HA}, an isogenic control virus in which we replaced

the *UL148* coding sequence with the UL148 homolog from rhesus cytomegalovirus,

Rh159, likewise fused at its C-terminus to an HA-tag. We considered the TB_159^{HA}

virus to be an appropriate control for the following reasons. Firstly, Rh159 and UL148

exhibit ~30% identity at the amino acid level and both glycoproteins localize to the ER

and block cell surface presentation immune cell activating ligands; UL148 retains CD58,

a ligand for CD2, while Rh159 retains NKG2D ligands of the MIC- and ULBP families (1,

9, 17). Secondly, Rh159 is expressed from TB_159^{HA} at comparable levels and with

similar kinetics observed for UL148 from TB_148^{HA}, and the two viruses replicate

indistinguishably in fibroblasts (**FIG 2**). Moreover, Rh159 does not appear to activate

the UPR (10), and ERQC markers fail to coalesce into unusual structures during

infection with TB_159^{HA} (**FIG 3**). Notably, ER structures fail to occur in rhesus

fibroblasts during infection with a recombinant rhesus cytomegalovirus (RhCMV)

engineered to express an HA-tag fused at the C-terminus of Rh159 (**FIG S1F**). This

argues against the possibility that Rh159 might somehow require the context of rhesus

cells to redistribute ER markers in a manner analogous to what is seen for UL148

during HCMV infection.

In cells infected with TB_148^{HA}, HA antibody staining indicated localization of

UL148 to globular structures, as expected (1). Antibody signals during indirect confocal

immunofluorescence detection of cellular ER resident proteins involved in ERQC,

including calnexin, Hrd1, SEL1L, Herp, valosin containing protein (VCP, p97), and

EDEM1, co-localized to a strong degree with signals from HA antibody detection of HA-

tagged UL148 (**FIG 3-4**). These results indicate that a number of ERQC factors

coalesce with UL148 to form prominent globular structures in infected cells. In contrast,

uniform ER staining patterns were observed with antibodies specific for PDI and

calreticulin (CALR), indicating that these ER markers do not co-localize to the UL148

structures (**FIGS 3Q, 3S**). Intriguingly, ER markers for smooth and rough ER,

ribophorin 1 and reticulon 3, both co-localized with UL148 (HA) signal at the structures

(**FIGS 3K, 3M**). This result suggests that rough and smooth ER may participate in the

ER structures. LC3B failed to co-localize with the structures (**FIG S2**), as expected

since macroautophagy is blocked at late times during HCMV infection (18, 19).

However, our results indicate that a related ATG8 family protein, GABARAP, co-

localizes with UL148 at the induced structures (**FIG 3O**).

In cells infected with TB_159^{HA}, all of the ER markers we examined showed

uniform, reticular staining, as did the anti-HA signal detecting Rh159 (**FIG 3B, D, F, H,**

J, L, N, P, R, T). EDEM1, in addition to showing reticular staining, also labeled puncta

associated with the nucleus (**FIG 3J**), which may represent enriched levels of the

protein at the rough ER membranes associated with the nuclear envelope, although we

cannot exclude the possibility of spurious intranuclear staining. Nonetheless, the

punctate nuclear signal for EDEM1 was much less pronounced in the TB_148^{HA}

infected cells (**FIG 3I**). GABARAP antibodies stained the cVAC in TB_159^{HA} infected

cells, but also showed a much weaker reticular signal that may indicate ER staining.

Even though we took steps to block viral Fc receptors, which can cause rabbit

antibodies to non-specifically label the cVAC (20), it is plausible that GABARAP

antibody signal from the cVAC reflects incomplete blocking of viral Fc receptors. For

both viruses, antibodies specific for syntaxin-6 (STX6) and gH, as expected, stained the

juxtanuclear cVAC structure (11, 12, 21), which was previously found to be altogether

distinct from the UL148 staining pattern (1)(**FIG S2**).

To quantify the degree of co-localization with UL148, we calculated Pearson's

correlation coefficients from a minimum of thirty TB_148^{HA} infected cells per staining

condition, comparing the signal overlap for each ER marker to the HA signal from

UL148. The correlation coefficients (*r*) for EDEM1, Hrd1, reticulon-3 (RTN3), SEL1L

and calnexin ranged from 0.8 to 0.72, which suggests that these proteins strongly co-

localize with UL148 by 4 days post-infection (dpi) (**FIG 4**). Meanwhile, GABARAP,

VCP, ribophorin-1 (RPN1), and Herp showed *r* values in the range of 0.54 to 0.51,

indicating moderate co-localization with UL148. As expected, the ER markers CALR

and PDI, and the TGN marker syntaxin-6, gave *r* values of close to zero, confirming that

these markers do not co-localize with UL148. From these results, we conclude that the

UL148-dependent ER structures are enriched for cellular markers associated with

glycoprotein quality control. In this regard, the UL148 ER structures resemble the "ER

quality control (ERQC) compartments" described by G. Lederkremer and colleagues

(14, 15, 22). Furthermore, the data indicate that Rh159 cannot substitute for UL148 to

induce these structures during HCMV infection of human fibroblasts, nor during RhCMV

infection of rhesus fibroblasts (**SI FIG S1**).

VCP and Hrd1 are recruited to incipient UL148 ER structures prior to calnexin.

To determine whether there might be differences in the kinetics of recruitment of

ERQC markers during formation of the ER structures, we examined a series of time

points from one to four days post-infection (dpi) with wild-type HCMV carrying HA-

tagged UL148, staining for three different ERQC markers, calnexin, Hrd1, and VCP,

alongside UL148 (HA), and acquired results using indirect immunofluorescence

confocal microscopy. At 1 dpi the signal from UL148 was only weakly detected, as

expected (1), while each of the ER markers was readily detected, providing a read-out

of their staining patterns prior to any substantial perturbation by UL148 (**FIG 5**). At 2

dpi, we detected robust anti-HA signal, indicating the presence of UL148. At this time

point, UL148 exhibited intense signal at small globular puncta, which we interpret to

represent incipient UL148 structures, as well as more diffuse staining of a reticular

structure consistent with ER. Calnexin (CNX) did not appreciably co-localize with the

UL148 puncta until at least 3 dpi, and the structures were not readily visualized from the

calnexin channel until 4 dpi (**FIG 5A**). In contrast, signals from Hrd1 and VCP staining

were sufficient to mark the UL148 puncta by 2 dpi, which suggests that these markers

may co-localize with UL148 at an earlier point during the genesis of the ER structures.

Notably, the appearance of Hrd1 at the structures prior to calnexin is consistent with our

previous results showing that UL148 co-purifies from infected cells with SEL1L, an

adaptor subunit for Hrd1 (8). These results imply that distal elements of the ERAD

machinery, exemplified by Hrd1 and VCP, may be recruited to UL148-induced ER

structures before calnexin.

Visualization of UL148-induced ER structures by electron microscopy.

To discern the ultrastructural appearance of the UL148 structures and to formally

ascertain their relationship to the ER, we carried out transmission electron microscopy

(TEM) imaging of wildtype (TB_WT) and *UL148*-null (TB_148$_{STOP}$) infected fibroblasts at

5 dpi. In high pressure frozen and freeze-substituted infected fibroblasts, cells with a

high density of viral nucleocapsids within the nucleus were selected for analysis, as this

feature indicates late time points during infection when UL148 is abundantly expressed.

The TEM results revealed prominent globular and oblong structures in the cytoplasm of

WT but not of *UL148*-null infected fibroblasts (**FIGS 6-7**). The structures stand out in

electron micrographs for their high electron-density, which may reflect the abundance of

ERQC proteins together with UL148 in these structures. Under higher magnification,

these areas are characterized by accumulations of densely packed ruffled membranes

and membranous material, which appear to be collapsed ER. Further, these membrane

accumulations were associated with smooth and partially rough ER structures with

seemingly enlarged cisternal space. Further, tomographic reconstruction of scanning

transmission electron microscopy (STEM) data suggest that densely packed ER

cisternae within the structures are interconnected and continuous in three-dimensional

space (**FIG 8B-C, SI Movie S1**).

A substantial fraction of UL148 is expressed in a detergent-insoluble form during

infection.

Certain disease-associated mutant alleles of cellular proteins, such as the A103E

mutant of the calcium channel Orai1 and the Z-allele of alpha-1 antitrypsin (Z A1AT),

localize to anomalous ER structures reminiscent of those we observed here to depend

on UL148 (16, 23-26). Furthermore, A103E Orai1 and Z A1AT are found to accumulate

in detergent-insoluble forms within the ER, which may represent aggregates or

polymers (16, 23, 26), and suggests a mechanism for the formation of ER structures by

the proteins. We therefore set out to determine whether differences in detergent

solubility between UL148 and Rh159 correlate with their differing potentials to

redistribute the ERQC machinery and to cause ER stress (10). To address this

question, we infected fibroblasts with TB_148[HA] or TB_159[HA] at MOI 1 and at various

times post-infection lysed the infected cells in radioimmunoprecipitation assay (RIPA)

buffer. After subjecting lysates to high-speed centrifugation, we examined the relative

levels of UL148 and Rh159 in the detergent-soluble supernatant and detergent-

insoluble pellet fractions.

A substantial fraction of UL148 was detected in the detergent-insoluble fractions

at all three time points tested (**FIG 9**). However, virtually all of the Rh159 signal was

detected in the detergent-soluble fraction. Only when anti-HA antibody signals were

shown at high exposure could faint levels of Rh159 be detected in the detergent-

insoluble fraction (**FIG 9A**, lower HA panel). From these results we concluded that

when compared to Rh159, UL148 is substantially less soluble in a detergent-based lysis

buffer, which suggests that UL148 may aggregate or assemble into higher-order

structures.

UL148 is sufficient to compartmentalize the ER.

Our data thus far demonstrate that UL148 is necessary during infection to cause

redistribution of cellular ER markers for glycoprotein quality control processes, such that

substantial portion of the ERQC machinery appears to become sequestered away from

the rest of the organelle into novel membranous structures. Because a number of viral

proteins that remodel the ER during infection are sufficient to alter ER morphology when

ectopically expressed [e.g. (27),(28), reviewed in (29)], we wondered whether UL148

expression is sufficient to induce ER structures akin to those seen during HCMV

infection. Making use of existing "tet-on" ARPE-19 epithelial cells that inducibly express

of either UL148 or Rh159, each carrying a C-terminal HA-tag (30), we induced

transgene expression for 48 h and subsequently stained for various cellular markers for

the ER, for ERQC as well as for LC3B and GABARAP, which are ATG8 family proteins

that mark autophagosome membranes.

In cells expressing UL148 (i148^{HA}), we observed that calnexin, Hrd1, EDEM1,

and VCP co-localize with UL148 at prominent globular structures reminiscent of those

observed during infection (**FIG 10**). The respective rough and smooth ER markers

ribophorin 1 (RPN1) and reticulon 3 (RTN3), as well as the ATG8 proteins LC3B and

GABARAP likewise co-localized to the UL148-induced structures. However, PDI and

CALR each failed to co-localize to the structures (**FIG 10**), consistent with our results

from infected cells (**FIG 3-4**).

In cells expressing the UL148 homolog Rh159 (i159^{HA}), we detected uniform

cytoplasmic distribution of ER markers, as well as of HA-tagged Rh159, similar to what

we observed during infection with the recombinant HCMV TB_159^{HA} (**FIG 3**). In this

setting, the staining patterns for the ATG8 family proteins LC3B and GABARAP failed to

indicate any notable structures. Consistent with our previous study showing that UL148,

but not Rh159, is sufficient to activate the UPR (10), we observed accumulation of ATF4

and phosphorylated eIF2α during induction of UL148 but not Rh159 (**FIG 11A**).

Importantly, HA signals indicated that UL148 and Rh159 accumulated at roughly

comparable levels following dox induction, which argues against the possibility that

gross differences in induced transgene expression accounts for the differences we

observed with regard to UPR activation and effects on staining patterns for ER markers.

We plotted the Pearson's correlation coefficient to estimate the extent of colocalization

between UL148 and various cellular markers: EDEM1, Hrd1, VCP, CNX, RTN3, SEL1L,

LC3B, and PDI (**FIG 11B**). The results quantitatively buttress the observations shown in

FIG 10.   From these results we concluded that no other HCMV gene products are

required for UL148 to cause ERQC factors to segregate into discrete compartments,

and hence, that UL148 is sufficient to cause large-scale reorganization of ERQC

markers. Further, these findings suggested to us that reorganization of ER markers into

discrete structures may be related to the propensity of UL148 to induce ER stress.

To determine whether UL148-dependent ER remodeling could be visualized in

real-time using live cell imaging, we constructed "tet-on" lentiviral vectors that inducibly

express either UL148 or Rh159 fused at their predicted C-terminal cytoplasmic tails to

the enhanced green fluorescent protein (GFP) from *Aequorea victoria* (31). Following

lentiviral transduction, puromycin-resistant ARPE-19 were isolated and subsequently

subjected to fluorescence-activated cell sorting (FACS) to enrich for cells that

expressed GFP signal following dox treatment. The resulting cell populations, which

inducibly express either UL148-GFP or Rh159-GFP, were designated i148^{GFP} and

i159^{GFP}, respectively.

In live cell imaging, we observed that the GFP signal in dox-induced i148^{GFP} cells

first appeared in a reticular, largely uniform pattern, which was readily visible at 5 h

post-induction. However, by 9 h post-induction punctate structures formed. These

puncta progressively grew larger up until the termination of the experiment at 19 h (**FIG**

12A, SI Movie 2). In contrast, i159^{GFP} cells exhibited a largely diffuse, reticular GFP

signal at all post-dox time points monitored (**FIG 12B, SI Movie 3**). The live cell

imaging results therefore recapitulated the differences in the HA-staining pattern that we

observed between dox-induced i148^{HA} and i159^{HA} cells following fixation, as well as

those between HA signals during HCMV infection with TB_148^{HA} and TB_159^{HA} (**FIGS**

1, 3, 10, SI Fig S1).

UL148-GFP structures exhibit poor recovery of fluorescence after

photobleaching.

To determine whether the protein contents of the UL148 induced structures

would exhibit decreased mobility compared to non-perturbed ER, we carried out

fluorescence recovery after photobleaching studies. We photobleached regions of GFP

signal in i148^{GFP} or i159^{GFP} ARPE-19 cells that had been induced for transgene

expression for 24 h and monitored recovery of signal over a 5 min period. In the case of

Rh159-GFP expressing cells (i159^{GFP}), fluorescence almost fully recovered within 3 min

(**FIG 13**). In contrast, when we photobleached a prominent UL148-GFP structure, the

GFP signal failed to appreciably recover fluorescence over the 5 min period for which

we obtained measurements. Nonetheless, an area of reticular GFP signal in an i148^{GFP}

cell, which presumably represented an ER region not involved in an anomalous ER

structure, recovered fluorescent signal with kinetics similar to those observed during

Rh159-GFP expression. These results suggested to us that UL148 structures do not

exchange their contents as efficiently as unperturbed ER, or that at least the UL148-

GFP within the structures is not able to be replaced rapidly with UL148-GFP protein

from other parts of the cell.

Ectopically-expressed UL148 is degraded by proteasomes and not by autophagy.

Although the UL148 ER structures were found to be enriched with proteins such

as Hrd1, SEL1L, EDEM, and VCP (**FIG 3-4**, **10-11B**) that are posited to play key roles in

ERAD, a process during which misfolded glycoproteins are recognized, processed, and

dislocated across the ER membrane for degradation at cytosolic proteasomes, we had

also obtained evidence suggesting that elements of the machinery for autophagy also

associated closely with the UL148 structures. Namely, the mammalian ATG8 ortholog

GABARAP co-localized with UL148 during infection (**FIG 3-4**), and both GABARAP and

another ATG8 ortholog, LC3B, both localized with the ER structures induced during

ectopic expression of UL148 (**FIG 10, 11B**). Therefore, we wished to determine

whether UL148 would degraded by the proteasome, as would be consistent with

conventional ERAD, or by the lysosome, which would suggest a functional role for

autophagy-related processes, such as selective autophagy of the ER (32), in dispensing

with UL148.

As a first step, we conducted a live-cell imaging "washout" experiment in which

i148^{GFP} cells were induced for transgene expression for 24 h, after which the growth

medium containing inducing agent (dox) was removed, the cells were washed, and then

the medium was lacking the inducing agent was applied to the cells. During a 22 h

imaging period following dox-washout, we observed the structures to become

progressively smaller as the reticular and punctate GFP signals abated (**FIG 14A, SI**

Movie S7). Because the results suggested that UL148-GFP structures could be

resolved over time, we induced UL148-GFP in tet-on ARPE-19 cells (i148^{gfp}) for 24 h,

washed out the inducing agent (dox) and then applied either epoxomicin (20 μM), an

irreversible inhibitor of the proteasome (33), or folimycin (115 nM), a proton pump

inhibitor that blocks autophagosome maturation and impedes the lysosome-dependent

protein degradation (34, 35). Of the two inhibitors, only epoxomicin markedly stabilized

UL148, as evidenced by Western blot signal for both GFP and UL148 (**FIG 14B-C**).

Folimycin had no obvious effect on UL148-GFP levels, even though the drug led to

increased levels of LC3B-II, as expected. From these results, we conclude that

ectopically expressed UL148 is primarily degraded by a proteasome-dependent

mechanism.

UL148 requires the integrated stress response to cause ER reorganization.

We previously defined that UL148 triggers the UPR during ectopic expression,

and that UL148 contributes to activation of the PKR-like ER kinase (PERK) and inositol

requiring enzyme 1 (IRE1) during infection (10). Although we failed to find any effect of

small interfering RNA (siRNA) or drug treatments targeting IRE1 on UL148-dependent

ER remodeling (not shown), we did observe that siRNA directed at PERK caused

UL148 to fail to form puncta or to co-localize with Hrd1 (not shown). Furthermore, the

literature suggests that formation of ERQC compartments requires PERK (36). PERK

responds to ER stress by activating the integrated stress response (ISR). PERK

activates the ISR by phosphorylating eIF2α at Ser51 (eIF2α-P); in turn, the

accumulation of eIF2α-P causes global attenuation of mRNA translation whilst

paradoxically stimulating translation of select mRNAs, such as those encoding ATF4

and CHOP, which play roles in adaptation to stress (37). Stress-regulated translation of

such mRNAs involves small upstream open reading frames, termed uORFs, in their 5'

untranslated regions that ordinarily suppress translation under basal conditions (38, 39).

To examine whether ER remodeling in response to UL148 requires the ISR, we

turned to a well characterized small molecule inhibitor of stress-regulated translation,

ISRIB (40-44). ISRIB is thought to function by acting as a "molecular staple" that holds

the guanine nucleotide exchange factor eIF2B in an active decameric configuration (42-

45), such that eIF2B will continue to generate ternary complex (eIF2•GTP•Met-tRNAi)

necessary for new cycles of translational initiation, despite the presence of phospho-

eIF2α (Ser51). Because PERK is the kinase that phosphorylates eIF2α in response to

ER stress (46, 47), we also tested for effects of the PERK inhibitor GSK2606414 (48).

Having confirmed that ISRIB and GSK260641 failed to negatively impact UL148

expression in dox-induced i148[HA] cells, when the drugs were applied at 200 nM or 1.1

µM, respectively **(FIG 15A-B)**, we treated i148[GFP] cells with 200 nM ISRIB, 1.1 µM

GSK260641, or DMSO vehicle control, and carried out live cell imaging during dox

induction of UL148-GFP.

Strikingly, ISRIB and GSK6060414 virtually abolished the formation of UL148

puncta through 13 h post induction (dox addition), a time point at which the DMSO

control condition showed abundant large structures (**FIG 15C-E, SI Movies 8-10**).

Based on our previous findings (**FIGs 1-3, 6-7, 10**), we interpret the formation and

subsequent large-scale aggregation of UL148-GFP puncta as a faithfully indicator of ER

remodeling in response to UL148. Therefore, these results suggested to us that

pharmacological treatments that inhibit the ISR and PERK prevent UL148 induced

remodeling of the ER, which further implies that ER remodeling in response to ER

stress requires stress-regulated translation.

Since our results with ISRIB and GSK6060414 during ectopic expression of

UL148 suggested that the ISR is required for UL148 mediated ER-remodeling, we next

asked whether these pharmacological agents would prevent UL148-dependent ER

remodeling in the authentic context of the HCMV infected cell. In live cell imaging

studies with ISRIB and GSK6060414, UL148-GFP puncta began to appear by ~17 h

post treatment (**FIG 15D-E, SI Movies 9-10**), suggesting that the inhibitory effects of the

drugs diminish over time. To ensure that the inhibitory activity would be maintained

over a four-day time course study in HCMV infected cells, we replaced the media at

each 24 h interval following infection to replenish freshly reconstituted ISRIB,

GSK6060414, or DMSO vehicle control.

Formation of ER structures was virtually ablated at 4 dpi during infection with

either ISRIB or GSK2606414 (**FIG 16A**). We used Imaris 3D imaging analysis of at

least 30 cells per condition to calculate the percentage of Hrd1 signals that coalesced

into discrete structures during infection, and the results corroborated statistically

significant differences between the DMSO carrier alone setting compared to treatments

with either GSK2606414 or ISRIB (**FIG 16B**). Furthermore, Western blot results

indicated roughly similar levels of UL148 in drug treated and DMSO control conditions,

and that ISRIB and GSK2606414 treatments both caused reduced levels of ATF4, while

PERK inhibitor alone was able to reduce the levels phospho-eIF2α (Ser51), detected by

a phospho-specific antibody, as expected (**FIG 16C-D**). A modest reduction, however,

was observed for the levels of viral envelope glycoprotein B (gB) during ISRIB and

GSK2606414 treatments, which suggests that the ISR may be required for optimal

expression of the certain viral envelope glycoproteins. Indeed, even viruses lacking

UL148 cause high levels of ATF4 accumulation at late times during infection, a time

when viral late genes, such a gB, are expressed at high levels (10). Regardless,

because the PERK inhibitor and ISRIB each prevented the appearance of discrete

UL148 ER structures while failing to substantially affect UL148 expression, these results

were consistent with those from our ectopic expression studies. Taken together, our

findings argue that the ISR is required for redistribution of ERQC markers in response to

UL148.

DISCUSSION

The ER structures we have identified herein are notable in several regards. Firstly, this example of virally-induced ER remodeling is wholly dependent on a single viral gene product, UL148. Of course, there are other examples of individual viral gene products that are necessary to grossly perturb ER morphology during infection and sufficient to induce such perturbations during ectopic expression [e.g., (27, 28, 49, 50)] Nonetheless, to the best of our knowledge the ER structures caused by UL148 are a hitherto undescribed ultrastructural characteristic of the HCMV infected cell, which is surprising given their size, scale, and prominence. Indeed, the average volume of the structures at 96 h postinfection amounts to roughly 60% of the size of the nucleus of an uninfected human fibroblast [~500 μm^3, (51)]. Although UL148 expression is accompanied by UPR activation, ER reorganization occurs without killing the host cell, even in the setting of ectopic expression when viral functions that inhibit programmed cell death are absent. Nor do we see any effects of UL148 on the yield of infectious virus during replication in fibroblasts, since *UL148*-null mutant viruses replicate indistinguishably from wildtype parental virus in cultured cells (1, 8). As with all enveloped viruses, HCMV requires the host cell secretory pathway to fold and process enormous quantities of viral envelope glycoproteins that endow progeny virions with the molecular machinery for infectivity. It is intriguing to consider how the infected cell tolerates spatial reorganization of ER membranes and the associated glycoprotein quality control machinery without any overtly impact on the production of infectious progeny virions.

Furthermore, certain aspects of ER reorganization invoked by UL148 may be

novel to cell biology. In particular, the densely packed collapsed ER in close

association with varicosities that we observe to depend on UL148 during infection

appear to differ from the ER structures caused by other ER perturbagens. For instance,

a hereditary form of childhood cirrhosis caused by the *Z* allele of the alpha-1-antitrypsin

(Z-A1AT) gene (*SERPINA*) is characterized by polymerization of Z-A1AT within the ER

(25), leading to its accumulation in membrane-delimited inclusions (25, 26, 52).

However, Z-A1AT inclusions are not observed to associate with regions of collapsed

ER, nor does Z-A1AT suffice to activate the UPR, even though its expression does

sensitize cells to other triggers of ER stress (26, 53).

HCMV encodes at least one other ER-resident immunevasin implicated to

activate the UPR, US11 (54). While UL148 binds the NK-cell and T-cell costimulatory

ligand CD58 to prevent its transport to the cell surface, US11 targets the heavy chain of

class I major histocompatibility complex (MHC) for ER-associated degradation (55, 56).

While UL148 retains CD58 in the ER and causes it to accumulate in presumably

immature glycoform that migrates with enhanced mobility in SDS-PAGE, there is no

obvious decrease in the overall abundance of CD58 (9). Even though intracellular

forms of CD58 in our hands are refractory to detection by indirect immunofluorescence

(not shown), we hypothesize that UL148 sequesters CD58 into the unusual ER

structures. Further, our detergent solubility results are consistent with the notion that

UL148 may form aggregates or polymers in the course of preventing CD58 presentation

at the cell surface (FIG 9). It is intriguing UL148, an immunevasin targeting CD58,

induces dramatic morphologic rearrangements of the ER, while another HCMV

immunevasin, US11, targeting MHC class I for degradation fails to do so, even though

both trigger ER stress. Although we cannot yet exclude the possibility, it is plausible

that retention of CD58 is required for ER reorganization.

While relationship between the mechanism for CD58 retention and the formation

of UL148-dependent ER structures remains unresolved, our findings suggest a

mechanism to explain the influence of UL148 on HCMV cell tropism. Because we find

that UL148 causes sequestration of the ER mannosidase EDEM1 and the basal E3

ubiquitin ligase SEL1L/Hrd1 into large membranous structures, regions of rough ER not

drawn into the structures might be expected to offer a more permissive folding

environment for poorly folding polypeptides that otherwise would be more efficiently

targeted for ERAD. Indeed, our results also indicate that the reticular ER, as indicated

by the staining pattern for calreticulin, PDI, and KDEL antibodies, remains largely intact

following UL148-induced ER reorganization of ER quality control markers into discrete

structures. Based on these observations, we hypothesize that UL148-mediated

sequestration of the glycoprotein quality control machinery alters the proteostatic

environment of the ER by physically dislocating key elements of the "mannose removal

time clock" (57, 58) that marks substrates for ERAD from marking said substrates for

destruction. Since glycoprotein O (gO) is at once a constitutive ERAD substrate (1, 8)

and a core component of a viral envelope glycoprotein complex that governs HCMV

entry and cell tropism (5-7, 59, 60), the hypothesis does not seem unreasonable on its

face. Obviously, however, it will be crucial to determine whether any generic ERAD

substrate are stabilized during UL148 expression, as would be predicted by this model.

Although many of the ERQC markers that co-localize with UL148 at the ER

structures, such as Hrd1 and EDEM1, are understood to play roles in pathways that

control proteasome dependent ERAD of terminally-misfolded glycoproteins, the

observation that GABARAP localizes to the structures during infection, and that both

LC3 and GABARAP do so during ectopic expression may suggest roles for autophagy-

related pathways in ER reorganization. However, our results indicate that UL148 is

degraded by a proteasome-dependent pathway and not by lysosomes. Therefore, it

seems unlikely that selective autophagy of the ER is involved in recycling or degrading

these ER structures.   Nonetheless, our live cell imaging results suggest that the UL148

ER structures originate as puncta that traffic to accumulate into larger structures, which

may suggest a relevance for co-localization with ATG8 family proteins.   Previous

studies have shown that misfolded proteins traffic in a microtubule (MT)-dependent

manner to form pericentriolar structures termed aggresomes (61, 62). ATG8 family

proteins such as LC3B not only play roles in degradation of substrates via

macroautophagy, but also bind MTs and recruit machinery for MT-dependent transport

of cargos (63-65). Therefore, it seems likely that the occurrence of GABARAP and

LC3B at the UL148 ER structures indicates recruitment of cellular machinery that

transports perturbed regions of ER to sites of large-scale accumulation.

Despite that our results indicate that the ISR is required for ER reorganization

during UL148 expression, precisely how UL148 triggers ER stress remains unknown.

UL148 has been found to co-purify from cells with SEL1L, a component of the ERAD

machinery. Thus, it seems plausible that UL148 may inhibit the Hrd1/SEL1L complex,

which would be predicted to cause the buildup of unfolded protein and thus trigger the

UPR. However, inhibition of ERAD alone seems unlikely to account for the formation of

ER structures. Another non-mutually exclusive possibility is that UL148 multimerizes in

a manner that constricts the ER lumen. Although the mechanism remains highly

speculative, the assembly UL148 molecules on opposite sides of the ER lumen might

constrict the organelle in a manner consistent with our EM results (FIG 6, 8).

MATERIALS AND METHODS.

Cells and Virus.

hTERT-immortalized human foreskin fibroblasts (8), derived from ATCC HFF-1 cells
(SCRC-1041) were maintained in Dulbecco's modified Eagle's medium supplemented
with 5%-10% newborn calf serum (Millipore Sigma) and antibiotics (complete DMEM)
exactly as described previously (8). i148^{HA} and i159^{HA} ARPE-19 epithelial cells (10),
which upon treatment with 100 ng/mL doxycycline, express HA-tagged UL148 or
Rh159, respectively, were likewise maintained in complete DMEM. For live cell imaging
studies, i148^{HA} and i159^{HA} ARPE-19 were maintained in Opti-MEM (Thermo Fisher)
supplemented with 3% certified tet-approves fetal bovine serum (FBS), 20 µg/mL
gentamicin sulfate, 1 µg/mL puromycin HCl and 10 µg/mL ciprofloxacin HCl.
Telomerase-immortalized rhesus fibroblasts (66) were a kind gift of Peter A. Barry, and
were maintained in complete DMEM. The acute monocytic leukemia cell line THP-1
(TIB-202) was obtained from ATCC (Manassas, VA) and were maintained in RPMI 1640
medium supplemented with 10% FBS (Millipore Sigma) and antibiotics (10 µg/mL
ciprofloxacin and 25 µg/mL gentamicin). Prior to infection with HCMV THP-1 were
differentiated into adherent macrophages by treatment with the phorbol ester, 2-O-
tetradecanoylphorbol 13-acetate (Millipore Sigma).

Infectious bacterial artificial chromosome (BAC) clones of HCMV strain TB40/E (also
known as TB40-BAC4 or TB_WT)(67), of strain TR (TR*gfp*)(68, 69), of strain Merlin

repaired for *UL128* and *RL13* harboring *tetO* sequences upstream of *UL131* (pAL1393)

(70), and of rhesus CMV strain 68-1 (71) were used in this study.

Construction of recombinant viruses and recombinant DNAs.

The TB40-BAC4 derivatives TB_148^HA and TB_148~STOP~ were used in our previous

studies and are described therein (1, 8, 10). New recombinant HCMVs for this study

were derived from BAC-cloned HCMV and RhCMVs using *en passant* mutagenesis in

GS1783 *E. coli,* a strain K12 derivative that expresses the homing endonuclease I-SceI

upon treatment with L-(+)-arabinose (72, 73). Recombinant BACs were confirmed by

Sanger DNA sequencing of the modified regions by Genewiz (Piscataway, NJ) (not

shown). Oligonucleotide primers for construction of recombinant viruses and plasmids

were custom synthesized by Integrated DNA Technologies (Coralville, IA); sequences

are provided in **Table S1**. Restriction enzymes, T4 DNA ligase, and Gibson assembly

reagents (NEB HiFi Assembly Master Mix) were obtained from New England Biolabs

(Ipswitch, MA).

To construct TB_159^HA, a strain TB40/E derivative in which the UL148 open

reading frame is replaced by Rh159 we carried out the following steps. Primers

Rh159_Fw and Rh159_HA_Rv were used to amplify and HA-tag the *Rh159* ORF from

pcDNA3.1+_Rh159_IRES_GFP (a gift from Klaus Früh, OHSU). The PCR product was

inserted into EcoRV-digested pEF1α using Gibson assembly. A PCR product

containing an ISce-I excisable kanamycin cassette was amplified from TB_148^HA_*ISce-*

Kan integrate BAC (1) using primers PpuISceKanGibs_Fw and PpuISceKanGibs _Rv

and Gibson-assembled into PpuMI-digested pEF1α_Rh159 plasmid to yield plasmid

pEF1α_Rh159_ISceKan. Primers Rh159_Fw_recomb and Rh159_Rv_recomb were

used to generate a PCR product from template plasmid pEF1α_Rh159_ISceKan. The

PCR product was electroporated into GS1783 E. coli harboring the TB_148^{HA} BAC, and

Kan^r colonies harboring TB_Δ148_Rh159_ISceKan integrate BACs were isolated on

Luria-Bertiani agar plates. Bacterial colonies representing Kan resistant integrates were

obtained, and integrates were subsequently resolved using standard *en passant*

protocols to yield TB_Δ148_Rh159^{HA}, which we abbreviate herein as TB_159^{HA}.

TB_159^{HA} was sequence-confirmed using primers TB_Rh159HA_seq_Fw and REV

TB_Rh159HA_seq. Similar strategies were used to modify the RhCMV 68-1 BAC (71)

so as to insert sequences encoding the nonapeptide HA-epitope such that Rh159 would

be expressed from the virus carrying a C-terminal HA tag. An *en passant* strategy in

which a shuttle plasmid carrying *I-SceI-Kan^r* disrupted by in-frame nonsense codons

was used insert premature nonsense codons in the *UL148* CDS in the context of HCMV

strains Merlin (pAL1393) and TR (TR*gfp*), and was applied as described previously for

generating TB_148$_{STOP}$, the UL148$_{STOP}$ mutant of strain TB40/E (8).

To construct "tet-on" lentivirus vector plasmids containing UL148 fused to GFP,

we used primers Gibs_eGFP_Rv and 148_eGFP_Fw to amplify the GFP gene from a

dsDNA gBlock, EGFP-P2A-3XHA, synthesized by Integrated DNA Technologies, gift of

Matthew D. Woolard, LSUHSC-S.   In a separate PCR, the *UL148* gene was amplified

from plasmid pcDNA3.1-UL148^{HA} (10) using primers Gib_148_Fw

148_noStopRv. The two products were assembled together with EcoRV opened

pcDNA3.1(+) (Invitrogen) to produce pcDNA3.1-UL148-GFP.   After confirming the

absence of spurious mutations by Sanger DNA sequencing (Genewiz), the *UL148-GFP*

cassette from pcDNA3.1-UL148-GFP was released by EcoRI and NotI digestion and

ligated into pOUPc turboRFP-link plasmid which had been linearized using the same

restriction sites. pOUPc turboRFP-link is a derivative of lentiviral vector pOUPc (10)

that was constructed by reinsertion of turboRFP (tRFP). Briefly, tRFP was amplified

from the original pOUPc-turboRFP using primer pair tRFP linker Fw and tRFP linker Rv

and then reinserted by Gibson Assembly with pOUPc turboRFP that had been double-

digested using EcoRI and Not I in order to linearize the vector and remove the

turboRFP (RFP) sequence. To express Rh159 fused to GFP, primers Gibs_eGFP_Rv

and 159_eGFP_Fw were used to amplify the GFP gene from EGFP-P2A-3XHA, and in

a separate PCR reaction, primers Gibs_159_Fw and 159_noStop_Rv Rh159 were

used to amplify *Rh159* from pcDNA3.1-Rh159[HA] (10). The latter two PCR products were

assembled together with EcoRV linearized pcDNA3.1(+) using Gibson Assembly,

resulting in plasmid pcDNA3.1-Rh159. The latter plasmid was used as template in a

PCR reaction with primers Gibs_Age_159_Fw and Gibs_Mlu_GFP_Rv. The resulting

PCR product was Gibson assembled into the lentiviral vector pOUPc (10) which had

been double-digested with Mlu I and Age I.

Drug treatments.

The PERK inhibitor, GSK2606414 (48), doxycycline hyclate, and folimycin were

obtained from MilliporeSigma (Burlington, MA). ISRIB (40-44) was obtained from

MilliporeSigma or APExBio (Boston, MA) and epoxomicin was procured Selleck

Chemicals (Houston, TX).   ISRIB was dissolved in DMSO and used at a final

concentration of 200 nM in live cell imaging and HCMV infection experiments but was

used at 2 μM in Western blotting experiments with i148[HA] ARPE-19 cells to address

whether the drug interfered with UL148 expression. GSK2606414 was also dissolved in

DMSO and used at a final working concentration of 1.1 μM. Folimycin was prepared as

a 1000× (115.46 μM) stock solution in DMSO and used at 115 nM final. Epoxomycin

was prepared as a 100× (2 mM) stock solution in DMSO and used at a final

concentration of 20 μM.

Confocal Microscopy and Live Cell Imaging. Confocal indirect immunofluorescence

microscopy imaging on fixed cells was carried out using Leica TCS SP5 Spectral

Confocal Microscope (Leica Microsystems, Heidelberg, Germany) using a Leica HCX

PL APO CS 63x/1.4-0.6NA objective under oil immersion, except for the image shown

in Figure 1C, which was captured on a Nikon SIM-E and A1R confocal microscopy

system (Nikon Instruments, Melville, NY) using a Nikon SR Apo TIRF 100x/1.49NA

objective under oil immersion. Images for different fluorophore channels were acquired

using sequential scanning. Direct immunofluorescence live cell imaging data were

collected using the Nikon SIM-E microscope using a Nikon Apo 60x/1.40NA DIC

objective.

The 3D projections shown in FIGs 1D and 16A were generated from Z-stacks

captured on the Nikon SIM-E / A1R microscope using a Nikon SR Apo TIRF

100x/1.49NA objective. For FIG 1D, the 3D projection was generated using NIS-

Elements AR Analysis 4.60.00 (64-bit) software (Nikon). For FIG 16A, 3D projections

were generated by Imaris x64 9.3.0 software (Bitplane, Inc.) in maximum intensity

projection (MIP) mode.

For fixed cell imaging experiments other than those shown in Fig S1 panels E, F,

G, and H, cells were seeded on 12 mm circular, No. 1 thickness microscope cover glass

(200121; Azer Scientific, Morgantown, PA). At the indicated times post-treatment and or

post-infection, cells were washed with phosphate buffered saline (PBS) [137 mM NaCl,

2.7 mM KCl, 10 mM Na_2HPO_4, and 1.8 mM KH_2PO_4, pH 7.4]; fixed for 15 min at room

temperature in PBS containing 4% (wt/vol) paraformaldehyde (Fisher Scientific, Waltham,

MA), washed in PBS, permeabilized for 10 min using 0.1% Triton X-100 (in PBS),

subsequently washed again in PBS, and then blocked for 45 min at 37 °C in PBS

containing 5% (vol/vol) normal goat serum (Rockland Immunochemicals, Limerick, PA).

Cells were then washed three times in PBS followed by incubation in 1% Human Fc Block

(BD Biosciences, San Jose, CA) in PBS, for additional 45 min at 37 °C. Cells were then

incubated in the presence of primary antibodies for 1 h at 37 °C or 4 °C overnight, and

then washed three times with PBS containing 0.1% Tween-20 (PBST) for 5 min per wash.

Alexa Fluor-labeled goat polyclonal secondary antibodies (all from ThermoFisher

Invitrogen, Waltham, MA, see **Table S2**) were used for secondary detection. The slides

were then mounted using Prolong Gold anti-fade reagent containing DAPI

(ThermoFisher) and placed under a Leica TCS SP5 confocal microscope for image

acquisition using a Leica 63× oil immersion objective lens (Leica Microsystems, Buffalo

Grove, IL).

For indirect immunofluorescence staining results of primary clinical isolates (SI Fig

S1 panels E-H), four clinical HCMV isolates obtained by routine testing of throat swabs

from patients of the Ulm University Medical Center were provided by the diagnostic

laboratory of the Institute of Virology in Ulm. Sample material was applied to human

700 fibroblasts and incubated for several days until HCMV-positive cells could be detected.

701 Infected cells were then seeded together with uninfected fibroblasts, incubated for up to

702 5 days until plaques were formed and processed for indirect immunofluorescence

703 staining. ERQC compartments were detected by staining for calnexin (CNX, E10; Santa

704 Cruz Biotechnology, mouse, 1:50 dilution), the cVAC was detected by staining for HCMV

705 pUL71 (rabbit, 1:500). Secondary antibody used for detection of CNX was Alexa Fluor

706 555 labeled goat anti-mouse IgG (1:1000) and for pUL71 detection, Alexa Fluor 488

707 conjugated goat anti-rabbit IgG (1:1000). Confocal images were acquired using the 63×

708 objective of a Zeiss Observer Z1 fluorescence microscope equipped with Apotome and

709 Zen software 2.3 (Carl Zeiss Microscopy GmbH, Jena, Germany).

711 **FRAP.** FRAP studies were carried out on using live ARPE-19 cells from the i148^{GFP}

712 and i159^{GFP} populations as follows. Cells were seeded as above for live-cell imaging,

713 induced for transgene expression using dox (100 ng) for 24 h, and then placed on an

714 incubated sample stage of a Nikon A1R SIM-E imaging system equipped with an SR

715 Apo TIRF 100x/1.49NA objective (Nikon). Three rectangular regions in image fields

716 were defined for measurement of (i) background signal, and of two regions with

717 comparable initial GFP signals, (ii) one designated as a control region (no

718 photobleaching) and another for (iii) photobleaching and recovery of signal after

719 photobleaching. Photobleaching of selected regions was carried out for 20 s using 405

720 nm laser light from a LU-N4 laser fiber (Nikon) [power at source: 15 mW, at objective: 8

721 mW]. Images and signal intensity measurements were captured every 2 s at a rate of 1

frame per s (488 nm excitation, FITC channel) starting immediately before

photobleaching (t=1), and from t=20 s to t=320 s after.

Electron Microscopy.   Procedures to prepare samples for transmission electron

microscopy (TEM) included high-pressure freezing (HPF), freeze substitution, and Epon

embedding, which were carried out as described previously (74).   Briefly, human

fibroblasts were seeded in μ-Slides (Ibidi GmbH, Martinsried, Germany) containing

carbon-coated sapphire discs (Engineering Office M. Wohlwend GmbH) 1 day prior to

infection at 80 to 90% confluence. Cells were infected with virus overnight at MOI 1.

Medium containing viral inocula was replaced with fresh medium the next day. Infected

cells on sapphire discs were fixed by using HPF with a Compact 01 high-pressure freezer

(Engineering Office M. Wohlwend GmbH, Sennwald, Switzerland) at 5 dpi.   Thereafter,

cells on sapphire discs were processed by freeze-substitution and subsequently

embedded in Epon (Fluka, Buchs, Switzerland). Ultrathin sections of the Epon-embedded

cells were cut with an ultramicrotome (Ultracut UCT; Leica Microsystems, Wetzlar,

Germany) and placed on Formvar-coated single-slot grids (Plano GmbH, Wetzlar,

Germany). Grids were examined in a Jeol JEM-1400 (Jeol Ltd., Tokyo, Japan)

transmission electron microscope equipped with a charge-coupled-device (CCD) camera

at an acceleration voltage of 120 kV.   Fixation and embedding of infected cells for

scanning transmission electron microscopy (STEM) tomography was the same as

described for TEM. Additional sample preparation steps were conducted as described

previously (75, 76). Tomogram acquisition was conducted on a STEM Jeol JEM-2100F

with an acceleration voltage of 200 kV. Tilt series were acquired from 600 nm thin sections

from +72° to -72° with a 1.5° increment using the bright field detector. Image series were

reconstructed to tomograms by weighted back projection with the IMOD software

package (77). Three-dimensional visualization of the membrane structures was

performed using Avizo lite software (Visualization Science Group, Burlington, MA, USA)

by threshold segmentation.

Western blotting. Western blotting procedures, including primary and secondary

antibodies used for detection of HA tag, ATF4, eIF2α, P-eIF2α (Ser51), the HCMV viral

nuclear antigen IE1, and the conditions used for detection of Ser51 phosphorylated

eIF2α, were carried out as described previously (8, 10). Additional antibodies used in

this study were mouse anti-GAPDH (Cat. No. 60004-1, Proteintech, Rosemont, IL,),

mouse anti-gB clone 27-180 (78) (a generous gift of William J. Britt), and anti-GFP

(D5.1) XP® Rabbit mAb #2956 (Cell Signaling Technology, Danvers, MA).

Statistical analyses. Statistical analyses were carried out using GraphPad Prism 8.1.0

for MacOS (GraphPad, San Diego, CA).

ACKNOWLEDGEMENTS.

This project was supported by NIH grants R01-AI116851 (to J.P.K.) and P30-

GM110703. Its contents are solely the responsibility of the authors and do not

necessarily represent the official views of the NIAID.

We are grateful to Erik L. Snapp (HHMI, Janelia Laboratories) for helpful

discussions concerning EM and confocal image data. We also thank Jacob E. Corn

 FIGURE LEGENDS:

FIG 1. UL148 reorganizes ER markers into anomalous structures during HCMV

infection.

Fibroblasts infected at 1 MOI with either **(A)** wildtype HCMV strain TB40/E (TB_WT) or

(B) an isogenic *UL148*-null mutant (TB_148STOP), were fixed at 96 h postinfection (hpi)

and imaged by confocal microscopy after staining with antibodies specific for CNX or Hrd1

(magenta) and glycoprotein H (gH, green), as indicated. DAPI (blue) signal was used to

counterstain nuclei. **(C)** Percentage of fibroblasts that contain Hrd1 structures at 96 hpi,

Fifty gH-positive cells for each condition were scored for presence or absence of UL148

structures (also see SI Fig S1A). (**D**) Three-dimensional confocal image projection of a

TB_WT infected fibroblast stained at 96 hpi for Hrd1 and gH. **(E)** Volumetric

measurements were made from sixteen infected cells, stained with phalloidin-AlexaFluor

594 conjugate to estimate total cell volume, with Hrd1 antibody to estimate the volume of

UL148-dependent ER structures, and with DAPI to estimate nuclear volume.

FIG 2. Characterization of TB_148^{HA} and TB_159^{HA} viruses. **(A)** Human fibroblasts

were infected at MOI 1 with the indicated recombinant HCMVs and whole cell lysate

samples harvested at the indicated times post-infection (h post-infection: hpi) were

analyzed by Western blot to detect HA-tagged UL148 or Rh159, the 72 kD viral nuclear

antigen IE1-72 (IE1), and GAPDH. **(B)** Single-cycle viral replication kinetic curves from

MOI 1 infected fibroblasts were plotted by determining the titer in tissue culture infectious

dose 50 (TCID$_{50}$) from supernatants collected at the indicated times postinfection.

FIG 3. UL148 localizes to unusual ER compartments enriched for glycoprotein

quality control markers. Fibroblasts infected at MOI 1 with either TB_148^HA (panels: **A,**

C, E, G, I, K, M, O, Q, S) or TB_ 159^HA (panels: **B, D, F, H, J, L, N, P, R, T**) were fixed

at 96 hpi, and imaged by confocal microscopy after co-staining with antibodies specific

for HA (UL148/Rh159, magenta in merge) and the indicated cellular markers (green in

merge). DAPI (blue) counterstaining is shown in merged image.

FIG 4. Quantification of Co-localization between UL148 and Cellular Markers. A

Pearson's correlation coefficient (r) was calculated using NIH ImageJ software to estimate

the degree of co-localization between UL148 (HA signal) and each of the indicated

cellular markers. A minimum of 30 cells were analyzed per marker. The arithmetic mean

for each co-localization analysis result is shown as a blue line, and data points for

individual cells analyzed are plotted as circles.

FIG 5. Cellular proteins involved in glycoprotein quality control are recruited with

differing kinetics to UL148 ER structures. Fibroblasts infected with TB_148^HA at 1 MOI

were fixed at the indicated time points (days post infection, dpi) and imaged by confocal

microscopy after staining with antibodies specific for HA (UL148, magenta), CNX (green,

A) , Hrd1 (green, **B**) or VCP(green, **C**), DAPI (blue).

FIG 6. TEM of ER structures in wild-type HCMV infected cells. Human fibroblasts

infected with wildtype HCMV (TB_WT) were fixed by high-pressure freezing and freeze

substitution at day 5 postinfection and imaged using TEM. Panels (**A**) and (**B**) show cell

overview at left. For each cell, the boxed region is shown at higher magnification.

Scale bars; left panels: 10 µm, right panels: 2 µm. rER: rough ER; Nu: nucleus; cVAC:

cytoplasmic viral assembly compartment. Solid arrowheads indicate the UL148-

dependent ER structures of interest.

FIG 7. TEM of *UL148*-null HCMV infected cells. Human fibroblasts infected with a

UL148-null mutant (TB_148STOP) were fixed by high-pressure freezing and freeze

substitution at day 5 postinfection and imaged using TEM. At the upper left an overview

panel of a representative cell is shown with panels **A, B, and C** each boxed. The boxed

regions are expanded at higher magnification at right (**A**) and below (**B, C**). Scale bars

for upper left overview panel: 10 µm, for zoomed panels: 2 µm. rER: rough ER; Nu:

nucleus; cVAC: cytoplasmic viral assembly compartment.

FIG 8. STEM tomography of UL148-dependent ER structures in HCMV infected

cells. Human fibroblasts infected with wildtype HCMV (TB_WT) were fixed by high-

pressure freezing and freeze substitution at day 5 postinfection and tomograms were

recorded by STEM. (**A**) Shows a virtual section through the tomogram of a virus-

induced membranous structure; asterisks denote ER structures of distended lumenal

space. White arrowheads indicate sites at which membranes originating from distended

ER cisternae continue into areas of involuted collapsed ER. Scale bar: 1 µm. (**B**)

Shows the same virtual section as in (A) tilted and with a 3D visualization of the

membranous network (green) of the entire tomogram. The same distended ER

cisternae as in (A) are marked by asterisks. (**C**) Shows a cross section through (B) to

visualize the membrane profile of the membranous structures. The region delimited by

a white box is shown in a higher magnification on the right. Finer detail of the enlarged

ER cisternae and the connections between them are readily visible; asterisks indicate

the same distended ER cisternae as in (A) and (B). Also, see SI Movie S1. Scale bar: 1

μm.

FIG 9. Solubility Analysis of UL148 and Rh159. (A) Human fibroblasts were infected at

MOI 1 $TCID_{50}$/cell with HCMV strain TB40/E derived viruses TB_148^{HA}, which

expresses UL148 fused at its C-terminus to an HA-epitope tag, or TB_159^{HA}, which

lacks *UL148* and instead expresses rhesus CMV Rh159 carrying a C-terminal HA tag.

At the indicated times postinfection (hpi), infected cells were collected

radioimmunoprecipitation (RIPA) lysis buffer. Lysates were then centrifuged at

21,000$\times$g for 30 min, after which supernatant (sup) and pellet (pel) fractions were boiled

in gel loading loading buffer containing 2% sodium dodecyl sulfate (SDS). Equivalent

portions of supernatant and pellet were then resolved by SDS-polyacrylamide gel

electrophoresis, and transferred to a nitrocellulose membrane for detection of protein

species immunoreactive to antibodies against HA epitope (HA) and HCMV glycoprotein

B (gB) and for total protein signal using Ponceau S reagent (Ponc.). (B-C). Signal

intensity of fluorescent conjugated secondary antibodies following anti-HA Western blot

was measured and plotted for detergent solubility analysis for experiment shown in (A)

or an independent biological replicate in which a 1% NP-40 lysis buffer was used

instead of RIPA buffer. Results are shown both as the arbitrary signal (in thousands) or

as a ratio of the signal from the pellet: soluble fractions, as indicated.

FIG 10. UL148 is sufficient to remodel the ER. HA-tagged UL148 or Rh159 were

doxycycline (dox) induced in the context of a "tet-on" lentiviral vector system in stably

transduced ARPE-19 epithelial cell populations, i148^{HA} and i159^{HA}, respectively. Cells

were fixed at 48 h postinduction for indirect immunofluorescence staining for the

indicated cellular markers (green) together with HA (magenta). Scale bar: 10 μm.

FIG 11. ISR activation accompanies redistribution of ER markers during ectopic

expression of UL148. (A) Lysates of tet-on ARPE-19 cells expressing either UL148 or

Rh159 fused to an HA tag, i148HA and i159HA, respectively, were collected at the

indicated times post doxycycline (dox) induction and analyzed by Western blot for the

expression of the indicated proteins and for the abundance of eIF2α phosphorylated at

Ser51 (P-eIF2α) using a phospho-specific antibody. **(B)** Pearson's correlation

coefficient (*r*) values were calculated using NIH ImageJ software estimate the degree of

co-localization between UL148 (HA signal) and the indicated cellular markers. A

minimum of 30 cells were analyzed per marker. Arithmetic means for each co-

localization analysis result are shown as blue lines, and data points for individual cells

analyzed are plotted as circles.

FIG 12. Live Cell Imaging of UL148-GFP and Rh159-GFP during Induced Ectopic

Expression. "Tet-on" ARPE-19 epithelial cells that inducibly express either UL148 or

Rh159 fused to green florescent protein (gfp), i148^{gfp} (**A**) and i159^{gfp} (**B**), respectively,

were induced for transgene expression using 100 ng/mL doxycycline (dox) and imaged

using live-cell microscopy. Images from the selected time points (h post treatment with

dox, hpt) are shown. Main scale bar represents 50 μm. For inset panels at upper left of

each image, which are magnified 2.4× relative to the main image, the scale bar

represents 10 μm. Also see SI Movies S2-S3.

FIG 13. FRAP analysis. ARPE-19 cells that inducibly express Rh159-GFP, i159^{gfp} (**A**)

or UL148-GFP, i148^{gfp} (**B-C**) were doxycycline induced for 24 h (100 ng/mL doxycycline

(dox). Separate regions of GFP signal in selected cells were photobleached (405 nm

laser) or left unbleached, while a third region lacking GFP signal was chosen as a

background reference were measured before and during fluorescence recovery after

photobleaching (FRAP). Note: background signal was not plotted because values were

resolvable from the x-axis. GFP signal intensity is plotted over a time period (seconds,

s) starting with an exposure at t=0 (immediately before photobleaching), and including

measurements taken every 2 s after photobleaching (0-20 s) until termination of the

measurement series at t=322 s (300 s of FRAP). (**D-F**) Images from the selected time

points (s, seconds) immediately before and after bleaching, and during fluorescence

recovery period. Also see SI Movies S4-S6.

FIG 14: UL148-GFP is stabilized by inhibition of proteasomal but not lysosomal

degradation. "Tet-on" ARPE-19 cells that inducibly express UL148-GFP (i148^{gfp}) were

induced for 24 h by the addition of 100 ng/mL doxycycline (dox), after which the dox

inducing agent was washed out and medium containing either the proteasome inhibitor

epoxomicin (20 μM) or the proton pump inhibitor folimycin (115 nM) was added and

samples were harvested for Western blot analysis at the indicated times post treatment

(h post treatment, hpt) with folimycin or epoximicin. Also see SI Movie S7.

FIG 15. Inhibition of the integrated stress response impedes UL148-mediated ER

remodeling. "Tet-on" ARPE-19 epithelial cells that inducibly express either UL148 or

Rh159 fused to green florescent protein (gfp), i148^{gfp} (**A**) and i159^{gfp} (**B**), respectively,

were induced for transgene expression using 100 ng/mL doxycycline (dox) in the

presence of absence of either ISRIB (200 nM) or GSK2606414 (1.1 μM) and monitored

by Western blot for expression of UL148-GFP over a series of time points (h post

treatment with dox, hpt). (**C-E**): Live-cell imaging of UL148-GFP and Rh159-GFP

expression patterns in the presence of ISRIB (200 nM), GSK2606414 (1.1 μM), or

DMSO vehicle alone. Main scale bar represents 50 μm. For inset panels at upper left of

each image, which are magnified 2.4× relative to the main image, the scale bar

represents 10 μm. Also see SI Movies S8-S10.

FIG 16. Inhibition of the ISR prevents the coalescence of Hrd1 and UL148 into

discrete structures during infection. (A) Representative 3D maximum intensity

projections of confocal imaging Z-stacks obtained from cells infected at MOI 1 for 96 h

with HCMV strain TB40/E carrying an HA-tagged *UL148*, TB_148^{HA}, and maintained in

the presence of ISRIB (200 nM), GSK2606414 (1.1 µM), or DMSO carrier alone (0.1%

vol/vol). In merged images HA signal is shown in magenta, Hrd1 in green, and DAPI

counterstaining in blue. (B) The percent of Hrd1 antibody signals that were involved in

discrete structures at 96 hpi were calculated for a minimum of 30 cells per condition were

determined using Imaris x64 9.3.0 software. Statistical significance was determined using

a one-way ANOVA followed by Tukey's post-test; **** represents a P-value of <0.0001.

The arithmetic mean for each co-localization analysis result is shown as a blue line, and

data points for individual cells analyzed are plotted as circles. (C-D) Western blot

analyses of fibroblasts infected at MOI 1 with TB_148^{HA} and maintained in the presence

of ISRIB (200 nM), GSK2606414 (1.1 µM), or DMSO carrier alone (0.1%); hpi: h post

infection (hpi). Note: a phospho-specific antibody was used for detection of eIF2α

phosphorylated at Ser51 (P-eIF2α).

952
953
954 **SI FIGURES and MOVIES.**

955
Fig S1: (A) Lower magnification overview of results from Fig 1A-B; confocal indirect

immunofluorescence staining of HCMV glycoprotein H (gH) and Hrd1 at 96 h

postinfection of human fibroblasts with wildtype HCMV strain TB40/E (WT) versus

UL148-null derivative TB_148$_{STOP}$. (B) Confocal indirect immunofluorescence staining

of gH and Hrd1 at 96 h postinfection of human fibroblasts with wildtype Merlin recovered

from BAC-cloned Merlin pAL1393 (ME_WT) or a *UL148*-null derivative of the same

virus (ME_148$_{STOP}$.) (C): Confocal indirect immunofluorescence staining of gH and

Hrd1 at 96 h postinfection of human fibroblasts with wildtype HCMV strain TRgfp

recovered from BAC-cloned TRgfp (TR_WT) or a *UL148*-null derivative of the same

virus (TR_148$_{STOP}$). (D) Confocal indirect immunofluorescence staining of the viral

tegument protein UL71 and calnexin at 5 days postinfection of human fibroblasts with

primary clinical HCMV isolates obtained from patient throat swabs. (E) Confocal

indirect immunofluorescence staining against HA and Hrd1 in THP-1 macrophages fixed

at 96 h postinfection with HCMV strain TB40/E derivative viruses TB_148^{HA} or

TB_159^{HA}; note: THP-1 monocytes were differentiated using phorbol ester treatment.

(**F**) Confocal indirect immunofluorescence staining of HA and Hrd1 in ARPE-19

epithelial cells infected for 96 h with either TB_148^{HA} or TB_159^{HA}. (**G**) Confocal indirect

immunofluorescence staining of HA and Hrd1 in ARPE-19 epithelial cells infected for 96

h with HCMV strain AD169 repaired for *UL131* and to which an HA-tagged UL148 CDS

from TB40/E was restored to the native *UL148* locus (AD^{r131}_148^{HA}) or with a parental

AD169 virus repaired for *UL131* to which *UL148* was not restored (AD^{r131}). Note: (i) a

functional *UL131* is required for efficient infection of epithelial cells; (ii) the difference in

scale bar between ADr131 and ADr131_148^{HA} is because with ADr131 we did not have

a viral marker (e.g., HA) to identify infected cells, so the appearance of syncytia and the

characteristic kidney-bean shaped nucleus was used to indicate infected cells, hence

the use of a slightly lower magnification. (**H**) Confocal indirect immunofluorescence

staining of telomerase-immortalized rhesus fibroblasts infected with BAC derived rhesus

CMV (RhCMV) strain 68-1 carrying an HA-tag at the C-terminus of Rh159.

FIG S2: Confocal microscopy of LC3B, syntaxin-6 and glycoprotein H staining

during HCMV infection. Human fibroblasts were infected with the indicated viruses for

96 h and then fixed, permeabilized, and stained using antibodies specific for the

indicated proteins. Indirect immunofluorescence images were captured using a 63X

objective on a Leica SP5 confocal microscope. gH: glycoprotein H, STX6: syntaxin-6.

993
MOVIE S1: TEM tomography of anomalous ER structures in wildtype HCMV infected

fibroblasts at day 5 postinfection.

MOVIES S2 – S7: Live cell imaging of GFP signal following dox induction of

transgene expression in i148^{gfp} and i159^{gfp} ARPE-19 epithelial cells.

Movie S2: i148^{gfp} cells from 2-19 h post dox induction.

Movie S3: i159^{gfp} cells from 2-19 h post dox induction.

1002

1003 **Movie S4**: **FRAP of UL148-GFP structure (ER within structure)**

1004 **Movie S5: FRAP of UL148 GFP structure (ER outside structure)**

1005 **Movie S6: FRAP of Rh159-GFP.**

1006

1007 **Movie S7: Imaging of UL148-GFP following washout of doxycycline.** (to 21 h 37

1008 min post washout).

1009

Movie S8: i148^{gfp} cells in the presence of 0.1% DMSO, from 2-19 h post dox

induction.

Movie S9: i148^{gfp} cells in the presence of 1.1 μM GSK2606414, from 2-19 h post

dox induction.

Movie S10: i148^{gfp} cells in the presence of 200 nM ISRIB, from 2-19 h post dox

induction.

1016

1017
1018

1019

1020

1021

1022

1.   Li G, Nguyen CC, Ryckman BJ, Britt WJ, Kamil JP. 2015. A viral regulator of glycoprotein complexes contributes to human cytomegalovirus cell tropism. Proc Natl Acad Sci U S A 112:4471-6.

2.   Wille PT, Knoche AJ, Nelson JA, Jarvis MA, Johnson DC. 2010. A human cytomegalovirus gO-null mutant fails to incorporate gH/gL into the virion envelope and is unable to enter fibroblasts and epithelial and endothelial cells. J Virol 84:2585-96.

3.   Jiang XJ, Adler B, Sampaio KL, Digel M, Jahn G, Ettischer N, Stierhof YD, Scrivano L, Koszinowski U, Mach M, Sinzger C. 2008. UL74 of human cytomegalovirus contributes to virus release by promoting secondary envelopment of virions. J Virol 82:2802-12.

4.   Zhou M, Lanchy JM, Ryckman BJ. 2015. Human Cytomegalovirus gH/gL/gO Promotes the Fusion Step of Entry into All Cell Types, whereas gH/gL/UL128-131 Broadens Virus Tropism through a Distinct Mechanism. J Virol 89:8999-9009.

5.   Wu Y, Prager A, Boos S, Resch M, Brizic I, Mach M, Wildner S, Scrivano L, Adler B. 2017. Human cytomegalovirus glycoprotein complex gH/gL/gO uses PDGFR-alpha as a key for entry. PLoS Pathog 13:e1006281.

6.   Kabanova A, Marcandalli J, Zhou T, Bianchi S, Baxa U, Tsybovsky Y, Lilleri D, Silacci-Fregni C, Foglierini M, Fernandez-Rodriguez BM, Druz A, Zhang B, Geiger R, Pagani M, Sallusto F, Kwong PD, Corti D, Lanzavecchia A, Perez L. 2016. Platelet-derived growth factor-alpha receptor is the cellular receptor for human cytomegalovirus gHgLgO trimer. Nat Microbiol 1:16082.

7.   Wu K, Oberstein A, Wang W, Shenk T. 2018. Role of PDGF receptor-alpha during human cytomegalovirus entry into fibroblasts. Proc Natl Acad Sci U S A doi:10.1073/pnas.1806305115.

8.   Nguyen CC, Siddiquey MNA, Zhang H, Li G, Kamil JP. 2018. Human Cytomegalovirus Tropism Modulator UL148 Interacts with SEL1L, a Cellular Factor That Governs Endoplasmic Reticulum-Associated Degradation of the Viral Envelope Glycoprotein gO. J Virol 92:e00688-18.

9.   Wang ECY, Pjechova M, Nightingale K, Vlahava VM, Patel M, Ruckova E, Forbes SK, Nobre L, Antrobus R, Roberts D, Fielding CA, Seirafian S, Davies J, Murrell I, Lau B, Wilkie GS, Suarez NM, Stanton RJ, Vojtesek B, Davison A, Lehner PJ, Weekes MP, Wilkinson GWG, Tomasec P. 2018. Suppression of costimulation by human cytomegalovirus promotes evasion of cellular immune defenses. Proc Natl Acad Sci U S A 115:4998-5003.

10.   Siddiquey MNA, Zhang H, Nguyen CC, Domma AJ, Kamil JP. 2018. The Human Cytomegalovirus Endoplasmic Reticulum-Resident Glycoprotein UL148 Activates the Unfolded Protein Response. J Virol 92:e00896-18.

11.   Das S, Vasanji A, Pellett PE. 2007. Three-dimensional structure of the human cytomegalovirus cytoplasmic virion assembly complex includes a reoriented secretory apparatus. J Virol 81:11861-9.

1061   12.   Sanchez V, Greis KD, Sztul E, Britt WJ. 2000. Accumulation of virion tegument and
1062   envelope proteins in a stable cytoplasmic compartment during human cytomegalovirus
1063   replication: characterization of a potential site of virus assembly. J Virol 74:975-86.
1064   13.   Lilley BN, Ploegh HL. 2005. Multiprotein complexes that link dislocation, ubiquitination,
1065   and extraction of misfolded proteins from the endoplasmic reticulum membrane. Proc
1066   Natl Acad Sci U S A 102:14296-301.
1067   14.   Kamhi-Nesher S, Shenkman M, Tolchinsky S, Fromm SV, Ehrlich R, Lederkremer GZ.
1068   2001. A novel quality control compartment derived from the endoplasmic reticulum.
1069   Mol Biol Cell 12:1711-23.
1070   15.   Leitman J, Shenkman M, Gofman Y, Shtern NO, Ben-Tal N, Hendershot LM, Lederkremer
1071   GZ. 2014. Herp coordinates compartmentalization and recruitment of HRD1 and
1072   misfolded proteins for ERAD. Mol Biol Cell 25:1050-60.
1073   16.   Houck SA, Ren HY, Madden VJ, Bonner JN, Conlin MP, Janovick JA, Conn PM, Cyr DM.
1074   2014. Quality control autophagy degrades soluble ERAD-resistant conformers of the
1075   misfolded membrane protein GnRHR. Mol Cell 54:166-179.
1076   17.   Sturgill ER, Malouli D, Hansen SG, Burwitz BJ, Seo S, Schneider CL, Womack JL, Verweij
1077   MC, Ventura AB, Bhusari A, Jeffries KM, Legasse AW, Axthelm MK, Hudson AW, Sacha
1078   JB, Picker LJ, Fruh K. 2016. Natural Killer Cell Evasion Is Essential for Infection by Rhesus
1079   Cytomegalovirus. PLoS Pathog 12:e1005868.
1080   18.   Chaumorcel M, Souquere S, Pierron G, Codogno P, Esclatine A. 2008. Human
1081   cytomegalovirus controls a new autophagy-dependent cellular antiviral defense
1082   mechanism. Autophagy 4:46-53.
1083   19.   Chaumorcel M, Lussignol M, Mouna L, Cavignac Y, Fahie K, Cotte-Laffitte J, Geballe A,
1084   Brune W, Beau I, Codogno P, Esclatine A. 2012. The human cytomegalovirus protein
1085   TRS1 inhibits autophagy via its interaction with Beclin 1. J Virol 86:2571-84.
1086   20.   Alwine JC. 2012. The human cytomegalovirus assembly compartment: a masterpiece of
1087   viral manipulation of cellular processes that facilitates assembly and egress. PLoS Pathog
1088   8:e1002878.
1089   21.   Das S, Pellett PE. 2011. Spatial relationships between markers for secretory and
1090   endosomal machinery in human cytomegalovirus-infected cells versus those in
1091   uninfected cells. J Virol 85:5864-79.
1092   22.   Leitman J, Ron E, Ogen-Shtern N, Lederkremer GZ. 2013. Compartmentalization of
1093   endoplasmic reticulum quality control and ER-associated degradation factors. DNA Cell
1094   Biol 32:2-7.
1095   23.   Cox DW, Billingsley GD, Callahan JW. 1986. Aggregation of plasma Z type alpha 1-
1096   antitrypsin suggests basic defect for the deficiency. FEBS Lett 205:255-60.
1097   24.   Le A, Ferrell GA, Dishon DS, Le QQ, Sifers RN. 1992. Soluble aggregates of the human PiZ
1098   alpha 1-antitrypsin variant are degraded within the endoplasmic reticulum by a
1099   mechanism sensitive to inhibitors of protein synthesis. J Biol Chem 267:1072-80.
1100   25.   Lomas DA, Evans DL, Finch JT, Carrell RW. 1992. The mechanism of Z alpha 1-antitrypsin
1101   accumulation in the liver. Nature 357:605-7.
1102   26.   Ordonez A, Snapp EL, Tan L, Miranda E, Marciniak SJ, Lomas DA. 2013. Endoplasmic
1103   reticulum polymers impair luminal protein mobility and sensitize to cellular stress in
1104   alpha1-antitrypsin deficiency. Hepatology 57:2049-60.

1105   27.   Suhy DA, Giddings TH, Jr., Kirkegaard K. 2000. Remodeling the endoplasmic reticulum by
1106   poliovirus infection and by individual viral proteins: an autophagy-like origin for virus-
1107   induced vesicles. J Virol 74:8953-65.
1108   28.   Windsor M, Hawes P, Monaghan P, Snapp E, Salas ML, Rodriguez JM, Wileman T. 2012.
1109   Mechanism of collapse of endoplasmic reticulum cisternae during African swine fever
1110   virus infection. Traffic 13:30-42.
1111   29.   Miller S, Krijnse-Locker J. 2008. Modification of intracellular membrane structures for
1112   virus replication. Nat Rev Microbiol 6:363-74.
1113   30.   Siddiquey MNA, Zhang H, Nguyen CC, Domma AJ, Kamil JP. 2018. The human
1114   cytomegalovirus ER resident glycoprotein UL148 activates the unfolded protein
1115   response. J Virol doi:10.1128/JVI.00896-18.
1116   31.   Cormack BP, Valdivia RH, Falkow S. 1996. FACS-optimized mutants of the green
1117   fluorescent protein (GFP). Gene 173:33-8.
1118   32.   Khaminets A, Heinrich T, Mari M, Grumati P, Huebner AK, Akutsu M, Liebmann L, Stolz
1119   A, Nietzsche S, Koch N, Mauthe M, Katona I, Qualmann B, Weis J, Reggiori F, Kurth I,
1120   Hubner CA, Dikic I. 2015. Regulation of endoplasmic reticulum turnover by selective
1121   autophagy. Nature 522:354-8.
1122   33.   Meng L, Mohan R, Kwok BH, Elofsson M, Sin N, Crews CM. 1999. Epoxomicin, a potent
1123   and selective proteasome inhibitor, exhibits in vivo antiinflammatory activity. Proc Natl
1124   Acad Sci U S A 96:10403-8.
1125   34.   Muroi M, Shiragami N, Nagao K, Yamasaki M, Takatsuki A. 1993. Folimycin
1126   (concanamycin A), a specific inhibitor of V-ATPase, blocks intracellular translocation of
1127   the glycoprotein of vesicular stomatitis virus before arrival to the Golgi apparatus. Cell
1128   Struct Funct 18:139-49.
1129   35.   Muroi M, Takasu A, Yamasaki M, Takatsuki A. 1993. Folimycin (concanamycin A), an
1130   inhibitor of V-type H(+)-ATPase, blocks cell-surface expression of virus-envelope
1131   glycoproteins. Biochem Biophys Res Commun 193:999-1005.
1132   36.   Kondratyev M, Avezov E, Shenkman M, Groisman B, Lederkremer GZ. 2007. PERK-
1133   dependent compartmentalization of ERAD and unfolded protein response machineries
1134   during ER stress. Exp Cell Res 313:3395-407.
1135   37.   Harding HP, Novoa I, Zhang Y, Zeng H, Wek R, Schapira M, Ron D. 2000. Regulated
1136   translation initiation controls stress-induced gene expression in mammalian cells. Mol
1137   Cell 6:1099-108.
1138   38.   Palam LR, Baird TD, Wek RC. 2011. Phosphorylation of eIF2 facilitates ribosomal bypass
1139   of an inhibitory upstream ORF to enhance CHOP translation. J Biol Chem 286:10939-49.
1140   39.   Vattem KM, Wek RC. 2004. Reinitiation involving upstream ORFs regulates ATF4 mRNA
1141   translation in mammalian cells. Proc Natl Acad Sci U S A 101:11269-74.
1142   40.   Sidrauski C, Acosta-Alvear D, Khoutorsky A, Vedantham P, Hearn BR, Li H, Gamache K,
1143   Gallagher CM, Ang KK, Wilson C, Okreglak V, Ashkenazi A, Hann B, Nader K, Arkin MR,
1144   Renslo AR, Sonenberg N, Walter P. 2013. Pharmacological brake-release of mRNA
1145   translation enhances cognitive memory. Elife 2:e00498.
1146   41.   Sidrauski C, McGeachy AM, Ingolia NT, Walter P. 2015. The small molecule ISRIB
1147   reverses the effects of eIF2alpha phosphorylation on translation and stress granule
1148   assembly. Elife 4.

1149   42.   Zyryanova AF, Weis F, Faille A, Alard AA, Crespillo-Casado A, Sekine Y, Harding HP, Allen
1150   F, Parts L, Fromont C, Fischer PM, Warren AJ, Ron D. 2018. Binding of ISRIB reveals a
1151   regulatory site in the nucleotide exchange factor eIF2B. Science 359:1533-1536.

1152   43.   Sidrauski C, Tsai JC, Kampmann M, Hearn BR, Vedantham P, Jaishankar P, Sokabe M,
1153   Mendez AS, Newton BW, Tang EL, Verschueren E, Johnson JR, Krogan NJ, Fraser CS,
1154   Weissman JS, Renslo AR, Walter P. 2015. Pharmacological dimerization and activation of
1155   the exchange factor eIF2B antagonizes the integrated stress response. Elife 4:e07314.

1156   44.   Tsai JC, Miller-Vedam LE, Anand AA, Jaishankar P, Nguyen HC, Renslo AR, Frost A, Walter
1157   P. 2018. Structure of the nucleotide exchange factor eIF2B reveals mechanism of
1158   memory-enhancing molecule. Science 359.

1159   45.   Gordiyenko Y, Schmidt C, Jennings MD, Matak-Vinkovic D, Pavitt GD, Robinson CV. 2014.
1160   eIF2B is a decameric guanine nucleotide exchange factor with a gamma2epsilon2
1161   tetrameric core. Nat Commun 5:3902.

1162   46.   Harding HP, Zhang Y, Ron D. 1999. Protein translation and folding are coupled by an
1163   endoplasmic-reticulum-resident kinase. Nature 397:271-4.

1164   47.   Harding HP, Zhang Y, Bertolotti A, Zeng H, Ron D. 2000. Perk is essential for translational
1165   regulation and cell survival during the unfolded protein response. Mol Cell 5:897-904.

1166   48.   Axten JM, Medina JR, Feng Y, Shu A, Romeril SP, Grant SW, Li WH, Heerding DA,
1167   Minthorn E, Mencken T, Atkins C, Liu Q, Rabindran S, Kumar R, Hong X, Goetz A, Stanley
1168   T, Taylor JD, Sigethy SD, Tomberlin GH, Hassell AM, Kahler KM, Shewchuk LM, Gampe
1169   RT. 2012. Discovery of 7-methyl-5-(1-{[3-(trifluoromethyl)phenyl]acetyl}-2,3-dihydro-1H-
1170   indol-5-yl)-7H-p yrrolo[2,3-d]pyrimidin-4-amine (GSK2606414), a potent and selective
1171   first-in-class inhibitor of protein kinase R (PKR)-like endoplasmic reticulum kinase
1172   (PERK). J Med Chem 55:7193-207.

1173   49.   Tenorio R, Fernandez de Castro I, Knowlton JJ, Zamora PF, Lee CH, Mainou BA, Dermody
1174   TS, Risco C. 2018. Reovirus sigmaNS and muNS Proteins Remodel the Endoplasmic
1175   Reticulum to Build Replication Neo-Organelles. MBio 9.

1176   50.   Miller S, Kastner S, Krijnse-Locker J, Buhler S, Bartenschlager R. 2007. The non-structural
1177   protein 4A of dengue virus is an integral membrane protein inducing membrane
1178   alterations in a 2K-regulated manner. J Biol Chem 282:8873-82.

1179   51.   Lamond AI, Sleeman JE. 2003. Nuclear substructure and dynamics. Curr Biol 13:R825-8.

1180   52.   Granell S, Baldini G, Mohammad S, Nicolin V, Narducci P, Storrie B, Baldini G. 2008.
1181   Sequestration of mutated alpha1-antitrypsin into inclusion bodies is a cell-protective
1182   mechanism to maintain endoplasmic reticulum function. Mol Biol Cell 19:572-86.

1183   53.   Lawless MW, Greene CM, Mulgrew A, Taggart CC, O'Neill SJ, McElvaney NG. 2004.
1184   Activation of endoplasmic reticulum-specific stress responses associated with the
1185   conformational disease Z alpha 1-antitrypsin deficiency. J Immunol 172:5722-6.

1186   54.   Tirosh B, Iwakoshi NN, Lilley BN, Lee AH, Glimcher LH, Ploegh HL. 2005. Human
1187   cytomegalovirus protein US11 provokes an unfolded protein response that may
1188   facilitate the degradation of class I major histocompatibility complex products. J Virol
1189   79:2768-79.

1190   55.   Wiertz EJ, Jones TR, Sun L, Bogyo M, Geuze HJ, Ploegh HL. 1996. The human
1191   cytomegalovirus US11 gene product dislocates MHC class I heavy chains from the
1192   endoplasmic reticulum to the cytosol. Cell 84:769-79.

56. van den Boomen DJ, Timms RT, Grice GL, Stagg HR, Skodt K, Dougan G, Nathan JA,
Lehner PJ. 2014. TMEM129 is a Derlin-1 associated ERAD E3 ligase essential for virus-
induced degradation of MHC-I. Proc Natl Acad Sci U S A 111:11425-30.
57. Caramelo JJ, Parodi AJ. 2008. Getting in and out from calnexin/calreticulin cycles. J Biol
Chem 283:10221-5.
58. Hammond C, Braakman I, Helenius A. 1994. Role of N-linked oligosaccharide
recognition, glucose trimming, and calnexin in glycoprotein folding and quality control.
Proc Natl Acad Sci U S A 91:913-7.
59. Zhou M, Yu Q, Wechsler A, Ryckman BJ. 2013. Comparative analysis of gO isoforms
reveals that strains of human cytomegalovirus differ in the ratio of gH/gL/gO and
gH/gL/UL128-131 in the virion envelope. J Virol 87:9680-90.
60. Zhang L, Zhou M, Stanton R, Kamil J, Ryckman BJ. 2018. Expression Levels of
Glycoprotein O (gO) Vary between Strains of Human Cytomegalovirus, Influencing the
Assembly of gH/gL Complexes and Virion Infectivity. J Virol 92.
61. Johnston JA, Ward CL, Kopito RR. 1998. Aggresomes: a cellular response to misfolded
proteins. J Cell Biol 143:1883-98.
62. Johnston JA, Illing ME, Kopito RR. 2002. Cytoplasmic dynein/dynactin mediates the
assembly of aggresomes. Cell Motil Cytoskeleton 53:26-38.
63. Pankiv S, Alemu EA, Brech A, Bruun JA, Lamark T, Overvatn A, Bjorkoy G, Johansen T.
2010. FYCO1 is a Rab7 effector that binds to LC3 and PI3P to mediate microtubule plus
end-directed vesicle transport. J Cell Biol 188:253-69.
64. Hummel T, Krukkert K, Roos J, Davis G, Klambt C. 2000. Drosophila Futsch/22C10 is a
MAP1B-like protein required for dendritic and axonal development. Neuron 26:357-70.
65. Wang H, Bedford FK, Brandon NJ, Moss SJ, Olsen RW. 1999. GABA(A)-receptor-
associated protein links GABA(A) receptors and the cytoskeleton. Nature 397:69-72.
66. Kirchoff V, Wong S, St JS, Pari GS. 2002. Generation of a life-expanded rhesus monkey
fibroblast cell line for the growth of rhesus rhadinovirus (RRV). Arch Virol 147:321-33.
67. Sinzger C, Hahn G, Digel M, Katona R, Sampaio KL, Messerle M, Hengel H, Koszinowski U,
Brune W, Adler B. 2008. Cloning and sequencing of a highly productive, endotheliotropic
virus strain derived from human cytomegalovirus TB40/E. J Gen Virol 89:359-68.
68. Smith IL, Taskintuna I, Rahhal FM, Powell HC, Ai E, Mueller AJ, Spector SA, Freeman WR.
1998. Clinical failure of CMV retinitis with intravitreal cidofovir is associated with
antiviral resistance. Arch Ophthalmol 116:178-85.
69. Murphy E, Yu D, Grimwood J, Schmutz J, Dickson M, Jarvis MA, Hahn G, Nelson JA,
Myers RM, Shenk TE. 2003. Coding potential of laboratory and clinical strains of human
cytomegalovirus. Proc Natl Acad Sci U S A 100:14976-81.
70. Stanton RJ, Baluchova K, Dargan DJ, Cunningham C, Sheehy O, Seirafian S, McSharry BP,
Neale ML, Davies JA, Tomasec P, Davison AJ, Wilkinson GW. 2010. Reconstruction of the
complete human cytomegalovirus genome in a BAC reveals RL13 to be a potent inhibitor
of replication. J Clin Invest 120:3191-208.
71. Chang WL, Barry PA. 2003. Cloning of the full-length rhesus cytomegalovirus genome as
an infectious and self-excisable bacterial artificial chromosome for analysis of viral
pathogenesis. J Virol 77:5073-83.

72. Tischer BK, Smith GA, Osterrieder N. 2010. En passant mutagenesis: a two step
markerless red recombination system. Methods Mol Biol 634:421-30.
73. Tischer BK, von Einem J, Kaufer B, Osterrieder N. 2006. Two-step red-mediated
recombination for versatile high-efficiency markerless DNA manipulation in Escherichia
coli. Biotechniques 40:191-7.
74. Dietz AN, Villinger C, Becker S, Frick M, von Einem J. 2018. A Tyrosine-Based Trafficking
Motif of the Tegument Protein pUL71 Is Crucial for Human Cytomegalovirus Secondary
Envelopment. J Virol 92.
75. Hohn K, Sailer M, Wang L, Lorenz M, Schneider ME, Walther P. 2011. Preparation of
cryofixed cells for improved 3D ultrastructure with scanning transmission electron
tomography. Histochem Cell Biol 135:1-9.
76. Villinger C, Schauflinger M, Gregorius H, Kranz C, Hohn K, Nafeey S, Walther P. 2014.
Three-dimensional imaging of adherent cells using FIB/SEM and STEM. Methods Mol
Biol 1117:617-38.
77. Kremer JR, Mastronarde DN, McIntosh JR. 1996. Computer visualization of three-
dimensional image data using IMOD. J Struct Biol 116:71-6.
78. Britt WJ, Jarvis MA, Drummond DD, Mach M. 2005. Antigenic domain 1 is required for
oligomerization of human cytomegalovirus glycoprotein B. J Virol 79:4066-79.
1254

1255

FIG 1. UL148 reorganizes ER markers into anomalous structures during HCMV infection.
Fibroblasts infected at 1 MOI with either **(A)** wildtype HCMV strain TB40/E (TB_WT) or **(B)** an isogenic *UL148*-null mutant (TB_148_STOP_), were fixed at 96 h postinfection (hpi) and imaged by confocal microscopy after staining with antibodies specific for CNX or Hrd1 (magenta) and glycoprotein H (gH, green), as indicated. DAPI (blue) signal was used to counterstain nuclei. **(C)** Percentage of fibroblasts that contain Hrd1 structures at 96 hpi, Fifty gH-positive cells for each condition were scored for presence or absence of UL148 structures (also see SI Fig S1A). (**D**) Three-dimensional confocal image projection of a TB_WT infected fibroblast stained at 96 hpi for Hrd1 and gH. **(E)** Volumetric measurements were made from sixteen infected cells, stained with phalloidin-AlexaFluor 594 conjugate to estimate total cell volume, with Hrd1 antibody to estimate the volume of UL148-dependent ER structures, and with DAPI to estimate nuclear volume.

FIG 2. Characterization of TB_148^HA and TB_159^HA viruses. **(A)** Human fibroblasts were infected at MOI 1 with the indicated recombinant HCMVs and whole cell lysate samples harvested at the indicated times post-infection (h post-infection: hpi) were analyzed by Western blot to detect HA-tagged UL148 or Rh159, the 72 kD viral nuclear antigen IE1-72 (IE1), and GAPDH. **(B)** Single-cycle viral replication kinetic curves from MOI 1 infected fibroblasts were plotted by determining the titer in tissue culture infectious dose 50 (TCID$_{50}$) from supernatants collected at the indicated times postinfection.

FIG 3. UL148 localizes to unusual ER compartments enriched for glycoprotein quality control markers. Fibroblasts infected at MOI 1 with either TB_148[HA] (panels: **A, C, E, G, I, K, M, O, Q, S**) or TB_ 159[HA] (panels: **B, D, F, H, J, L, N, P, R, T**) were fixed at 96 hpi, and imaged by confocal microscopy after co-staining with antibodies specific for HA (UL148/Rh159, magenta in merge) and the indicated cellular markers (green in merge). DAPI (blue) counterstaining is shown in merged image.

FIG 4. Quantification of Co-localization between UL148 and Cellular Markers. A Pearson's correlation coefficient (*r*) was calculated using NIH ImageJ software to estimate the degree of co-localization between UL148 (HA signal) and each of the indicated cellular markers. A minimum of 30 cells were analyzed per marker. The arithmetic mean for each co-localization analysis result is shown as a blue line, and data points for individual cells analyzed are plotted as circles.

FIG 5. Cellular proteins involved in glycoprotein quality control are recruited with differing kinetics to UL148 ER structures. Fibroblasts infected with TB_148[HA] at 1 MOI were fixed at the indicated time points (days post infection, dpi) and imaged by confocal microscopy after staining with antibodies specific for HA (UL148, magenta), CNX (green, **A**) , Hrd1 (green, **B**) or VCP(green, **C**), DAPI (blue).

FIG 6. TEM of ER structures in wild-type HCMV infected cells. Human fibroblasts infected with wildtype HCMV (TB_WT) were fixed by high-pressure freezing and freeze substitution at day 5 postinfection and imaged using TEM. Panels (**A**) and (**B**) show cell overview at left. For each cell, the boxed region is shown at higher magnification. Scale bars; left panels: 10 μm, right panels: 2 μm. rER: rough ER; Nu: nucleus; cVAC: cytoplasmic viral assembly compartment. Solid arrowheads indicate the UL148-dependent ER structures of interest.

FIG 7. TEM of *UL148*-null HCMV infected cells. Human fibroblasts infected with a *UL148*-null mutant (TB_148$_{STOP}$) were fixed by high-pressure freezing and freeze substitution at day 5 postinfection and imaged using TEM. At the upper left an overview panel of a representative cell is shown with panels **A, B, and C** each boxed. The boxed regions are expanded at higher magnification at right (**A**) and below (**B, C**). Scale bars for upper left overview panel: 10 μm, for zoomed panels: 2 μm. rER: rough ER; Nu: nucleus; cVAC: cytoplasmic viral assembly compartment.

FIG 8. STEM tomography of UL148-dependent ER structures in HCMV infected cells. Human fibroblasts infected with wildtype HCMV (TB_WT) were fixed by high-pressure freezing and freeze substitution at day 5 postinfection and tomograms were recorded by STEM. (**A**) Shows a virtual section through the tomogram of a virus-induced membranous structure; asterisks denote ER structures of distended lumenal space. White arrowheads indicate sites at which membranes originating from distended ER cisternae continue into areas of involuted collapsed ER. Scale bar: 1 µm. (**B**) Shows the same virtual section as in (A) tilted and with a 3D visualization of the membranous network (green) of the entire tomogram. The same distended ER cisternae as in (A) are marked by asterisks. (**C**) Shows a cross section through (B) to visualize the membrane profile of the membranous structures. The region delimited by a white box is shown in a higher magnification on the right. Finer detail of the enlarged ER cisternae and the connections between them are readily visible; asterisks indicate the same distended ER cisternae as in (A) and (B). Also, see SI Movie S1. Scale bar: 1 µm.

FIG 9. Solubility Analysis of UL148 and Rh159. (A) Human fibroblasts were infected at MOI 1 TCID$_{50}$/cell with HCMV strain TB40/E derived viruses TB_148^{HA}, which expresses UL148 fused at its C-terminus to an HA-epitope tag, or TB_159^{HA}, which lacks *UL148* and instead expresses rhesus CMV Rh159 carrying a C-terminal HA tag. At the indicated times postinfection (hpi), infected cells were collected radioimmunoprecipitation (RIPA) lysis buffer. Lysates were centrifuged at 21,000×g for 30 min, after which supernatant (sup) and pellet (pel) fractions were boiled in gel loading loading buffer containing 2% sodium dodecyl sulfate (SDS). Equivalent portions of supernatant and pellet were then resolved by SDS-polyacrylamide gel electrophoresis, and transferred to a nitrocellulose membrane for detection of protein species immunoreactive to antibodies against HA epitope (HA) and HCMV glycoprotein B (gB) and for total protein signal using Ponceau S reagent (Ponc.). (B-C). Signal intensity of fluorescent conjugated secondary antibodies following anti-HA Western blot was measured and plotted for detergent solubility analysis for experiment shown in (A) or an independent biological replicate in which a 1% NP-40 lysis buffer was used instead of RIPA buffer. Results are shown both as the arbitrary signal (in thousands) or as a ratio of the signal from the pellet: soluble fractions, as indicated.

FIG 10. UL148 is sufficient to remodel the ER. HA-tagged UL148 or Rh159 were doxycycline (dox) induced in the context of a "tet-on" lentiviral vector system in stably transduced ARPE-19 epithelial cell populations, i148^HA and i159^HA, respectively. Cells were fixed at 48 h postinduction for indirect immunofluorescence staining for the indicated cellular markers (green) together with HA (magenta). Scale bar: 10 μm.

FIG 11. ISR activation accompanies redistribution of ER markers during ectopic expression of UL148. (A) Lysates of tet-on ARPE-19 cells expressing either UL148 or Rh159 fused to an HA tag, i148HA and i159HA, respectively, were collected at the indicated times post doxycycline (dox) induction and analyzed by Western blot for the expression of the indicated proteins and for the abundance of eIF2α phosphorylated at Ser51 (P-eIF2α) using a phospho-specific antibody. (**B**) Pearson's correlation coefficient (*r*) values were calculated using NIH ImageJ software estimate the degree of co-localization between UL148 (HA signal) and the indicated cellular markers. A minimum of 30 cells were analyzed per marker. Arithmetic means for each co-localization analysis result are shown as blue lines, and data points for individual cells analyzed are plotted as circles.

FIG 12. Live Cell Imaging of UL148-GFP and Rh159-GFP during Induced Ectopic Expression. "Tet-on" ARPE-19 epithelial cells that inducibly express either UL148 or Rh159 fused to green florescent protein (gfp), i148^{gfp} (**A**) and i159^{gfp} (**B**), respectively, were induced for transgene expression using 100 ng/mL doxycycline (dox) and imaged using live-cell microscopy. Images from the selected time points (h post treatment with dox, hpt) are shown. Main scale bar represents 50 µm. For inset panels at upper left of each image, which are magnified 2.4× relative to the main image, the scale bar represents 10 µm. Also see SI Movies S2-S3.

FIG 13. FRAP analysis. ARPE-19 cells that inducibly express Rh159-GFP, i159^{gfp} (**A**) or UL148-GFP, i148^{gfp} (**B-C**) were doxycycline induced for 24 h (100 ng/mL doxycycline (dox). Separate regions of GFP signal in selected cells were photobleached (405 nm laser) or left unbleached, while a third region lacking GFP signal was chosen as a background reference were measured before and during fluorescence recovery after photobleaching (FRAP). Note: background signal was not plotted because values were resolvable from the x-axis. GFP signal intensity is plotted over a time period (seconds, s) starting with an exposure at t=0 (immediately before photobleaching), and including measurements taken every 2 s after photobleaching (0-20 s) until termination of the measurement series at t=322 s (300 s of FRAP). (**D-F**) Images from the selected time points (s, seconds) immediately before and after bleaching, and during fluorescence recovery period. Also see SI Movies S4-S6.

FIG 14: UL148-GFP is stabilized by inhibition of proteasomal but not lysosomal degradation. "Tet-on" ARPE-19 cells that inducibly express UL148-GFP (i148gfp) were induced for 24 h by the addition of 100 ng/mL doxycycline (dox), after which the dox inducing agent was washed out and medium containing either the proteasome inhibitor epoxomicin (20 µM) or the proton pump inhibitor folimycin (115 nM) was added and samples were harvested for Western blot analysis at the indicated times post treatment (h post treatment, hpt) with folimycin or epoximicin. Also see SI Movie S7.

FIG 15. Inhibition of the integrated stress response impedes UL148-mediated ER remodeling. "Tet-on" ARPE-19 epithelial cells that inducibly express either UL148 or Rh159 fused to green florescent protein (gfp), i148^{gfp} (**A**) and i159^{gfp} (**B**), respectively, were induced for transgene expression using 100 ng/mL doxycycline (dox) in the presence of absence of either ISRIB (200 nM) or GSK2606414 (1.1 µM) and monitored by Western blot for expression of UL148-GFP over a series of time points (h post treatment with dox, hpt). (**C-E**): Live-cell imaging of UL148-GFP and Rh159-GFP expression patterns in the presence of ISRIB (200 nM), GSK2606414 (1.1 µM), or DMSO vehicle alone. Main scale bar represents 50 µm. For inset panels at upper left of each image, which are magnified 2.4× relative to the main image, the scale bar represents 10 µm. Also see SI Movies S8-S10.

FIG 16. Inhibition of the ISR prevents the coalescence of Hrd1 and UL148 into discrete structures during infection. (A) Representative 3D maximum intensity projections of confocal imaging Z-stacks obtained from cells infected at MOI 1 for 96 h with HCMV strain TB40/E carrying an HA-tagged *UL148*, TB_148[HA], and maintained in the presence of ISRIB (200 nM), GSK2606414 (1.1 µM), or DMSO carrier alone (0.1% vol/vol). In merged images HA signal is shown in magenta, Hrd1 in green, and DAPI counterstaining in blue. (B) The percent of Hrd1 antibody signals that were involved in discrete structures at 96 hpi were calculated for a minimum of 30 cells per condition were determined using Imaris x64 9.3.0 software. Statistical significance was determined using a one-way ANOVA followed by Tukey's post-test; **** represents a P-value of <0.0001. The arithmetic mean for each co-localization analysis result is shown as a blue line, and data points for individual cells analyzed are plotted as circles. (C-D) Western blot analyses of fibroblasts infected at MOI 1 with TB_148[HA] and maintained in the presence of ISRIB (200 nM), GSK2606414 (1.1 µM), or DMSO carrier alone (0.1%); hpi: h post infection (hpi). Note: a phospho-specific antibody was used for detection of eIF2α phosphorylated at Ser51 (P-eIF2α).

Fig S1: (A) Lower magnification overview of results from Fig 1A-B; confocal indirect immunofluorescence staining of HCMV glycoprotein H (gH) and Hrd1 at 96 h postinfection of human fibroblasts with wildtype HCMV strain TB40/E (WT) versus *UL148*-null derivative TB_148$_{STOP}$. (B) Confocal indirect immunofluorescence staining of gH and Hrd1 at 96 h postinfection of human fibroblasts with wildtype Merlin recovered from BAC-cloned Merlin pAL1393 (ME_WT) or a *UL148*-null derivative of the same virus (ME_148$_{STOP}$.) (C): Confocal indirect immunofluorescence staining of gH and Hrd1 at 96 h postinfection of human fibroblasts with wildtype HCMV strain TRgfp recovered from BAC-cloned TRgfp (TR_WT) or a *UL148*-null derivative of the same virus (TR_148$_{STOP}$). (D) Confocal indirect immunofluorescence staining of the viral tegument protein UL71 and calnexin at 5 days postinfection of human fibroblasts with primary clinical HCMV isolates obtained from patient throat swabs. (E) Confocal indirect immunofluorescence staining against HA and Hrd1 in THP-1 macrophages fixed at 96 h postinfection with HCMV strain TB40/E derivative viruses TB_148^{HA} or TB_159^{HA}; note: THP-1 monocytes were differentiated using phorbol ester treatment. (**F**) Confocal indirect immunofluorescence staining of HA and Hrd1 in ARPE-19 epithelial cells infected for 96 h with either TB_148^{HA} or TB_159^{HA}. (**G**) Confocal indirect immunofluorescence staining of HA and Hrd1 in ARPE-19 epithelial cells infected for 96 h with HCMV strain AD169 repaired for *UL131* and to which an HA-tagged UL148 CDS from TB40/E was restored to the native *UL148* locus (AD^{r131}_148^{HA}) or with a parental AD169 virus repaired for *UL131* to which *UL148* was not restored (AD^{r131}). Notes: (i) a functional *UL131* is required for efficient infection of epithelial cells; (ii) the difference in scale bar between ADr131 and ADr131_148^{HA} is because with ADr131 we did not have a viral marker (e.g., HA) to identify infected cells, so the appearance of syncytia and the characteristic kidney-bean shaped nucleus was used to indicate infected cells, hence the use of a slightly lower magnification. (**H**) Confocal indirect immunofluorescence staining of telomerase-immortalized rhesus fibroblasts infected with BAC derived rhesus CMV (RhCMV) strain 68-1 carrying an HA-tag at the C-terminus of Rh159.

FIG S2: Confocal microscopy of LC3B, syntaxin-6 and glycoprotein H staining during HCMV infection. Human fibroblasts were infected with the indicated viruses for 96 h and then fixed, permeabilized, and stained using antibodies specific for the indicated proteins. Indirect immunofluorescence images were captured using a 63X objective on a Leica SP5 confocal microscope. gH: glycoprotein H, STX6: syntaxin-6.

TABLE S1. Oligonucleotide primers and synthetic DNAs used in this study

Name	Sequence (5' to 3')
148_eGFP_Fw	ACCTGGTGTCGCGGCGTCGGATGGTGAGCAAGGGGGAGGAGCTTTTTACTGGCGTTGTTC
Gibs_eGFP_Rv	CGGCCGCCACTGTGCTGGATTACTTATACAGCTCGTCCATACCCAG
Gib_148_Fw	TGTGGTGGAATTCTGCAGATTGCCACCATGTTGCGCTTGCTG
148_noStopRv	CCGACGCCGCGACACCAGGTAGGTTATC
159_eGFP_Fw	TCTAAAACGCAGTCGTGAAGCTCATATGGTGAGCAAGGGGGAGGA
159_noStop_Rv	ATGAGCTTCACGACTGCGTTTTAGAAGTG
Gibs_159_Fw	TGTGGTGGAATTCTGCAGATTGCCACCATGGCCTACAACAGC
Rh159_Fw	TGTGGTGGAATTCTGCAGATACCATGGCCTACAACAGCTTCCTTCTCAGCTGCCTTACCATCGCACTTTTACTG
Rh159_HA_Rv	CGGCCGCCACTGTGCTGGATCTATTAGGCGTAGTCTGGGACGTCGTATGGGTAATGAGCTTCACGACTGCGTTTTAGAAGTG
PpuISceKanGibs_Fw	CATTCACCTGAACCCATAGGTCCTGCCTGGCAGAGAATCTTAGGGATAACAGGGTAATCG
PpuISceKanGibs _Rv	AGATTCTCTGCCAGGCAGGACCTATGGGTTCAGGTGAATGGCCAGTGTTACAACCAATTA
Rh159_Fw_recomb	GGGCGGTGGCCGGCACGCCGCATTTCCTAACCCGCGCAGCATGGCCTACAACAGCTTCCT
Rh159_Rv_recomb	CGCGAACGACGTGTGACGAGGACGTGGTTTCCGCAAGCCTTTAGGCGTAGTCTGGGACG
TB_Rh159HA_seq_Fw	AGTGTTTATGAGTCGGGCGG
REV TB_Rh159HA_seq	GTGGCGACGTGGATTTCTTG
EGFP-P2A-3XHA	GTCGACTGGATCCGGTACCGAATTCGCCACCATGGTGAGCAAGGGGGAGGAGCTTTTTACTGGCGTTGTTCCCATCCTGGTTGAGTTGGATGGGGATGTCAACGGCCACAAGTTCAGCGTTTCTGGAGAAGGAGAAGGCGACGCAACCTATGGGAAGCTCACTTTGAAATTCATATGCACCACCGGTAAACTCCCAGTCCCCTGGCCTACCCTGGTCACTACCTTGACATATGGAGTTCAGTGCTTTTCTCGATACCCCGACCACATGAAGCAGCACGATTTTTTCAAATCTGCAATGCCCGAGGGCTATGTACAAGAGAGGACCATTTTCTTCAAAGACGACGGCAATTACAAGACCAGGGCCGAGGTCAAATTTGAAGGGGACACTCTTGTGAATCGGATCGAACTTAAAGGGATCGATTTCAAAGAGGACGGAAACATACTGGGGCATAAGCTGGAATACAACTACAATAGCCATAATGTTTACATTATGGCAGACAAGCAGAAAAACGGTATTAAGGTCAACTTCAAGATACGGCATAACATTGAAGATGGGTCAGTCCAGCTTGCTGACCATTATCAGCAGAATACACCCATTGGCGATGGTCCCGTGCTTTTGCCTGACAACCACTACCTTTCTACCCAAAGCGCATTGTCAAAAGATCCTAACGAGAAGAGAGATCATATGGTCCTTTTGGAGTTCGTAACCGCAGCTGGTATAACTCTGGGTATGGACGAGCTGTATAAGGGCTCAGGGGCTACAAATTTTAGTCTCCTTAAACAAGCCGGAGATGTCGAAGAGAACCCTGGGCCTTACCCATATGACGTTCCTGACTATGCCGGTTACCCCTATGATGTGCCAGATTATGCCGGATCTTATCCTTACGATGTACCTGACTATGCAGGTGGGAGTGGCATGAATTACGTGGGGCAGC
Gibs_Age_159_Fw	CCGTACCACTTCCTACCCTCGTAAACCGGTGCCACCATGGCCTACAACAG
Gibs_Mlu_GFP_Rv	ACCCGGCGCGGAGGCCAGATCTTAACGCGTTTACTTATACAGCTCGTCCATACC
tRFP linker Fw	CCACTTCCTACCCTCGTAAACCGGTGAATTCGCCACCATGAGCGAGCTGAT
tRFP linker Rv	GCGCGGAGGCCAGATCTTAACGCGTGCGGCCGCTCGAGAATTCATCTGTGCCCCAGTTTGC

TABLE S2. Antibodies Used in Immunofluorescence Microscopy

Primary Antibody	Host species	Isotype	Clone	Vendor	Catalog No.	Dilution used
Calnexin	Mouse	mAb IgG1	37	BD Biosciences	610523	1:50
Calnexin	Rabbit	mAb IgG	C5C9	Cell Signaling Tech.	2679	1:50
Syntaxin 6	Rabbit	mAb IgG	C34B2	Cell Signaling Tech.	2869	1:50
SYVN1 (Hrd1)	Rabbit	mAb IgG	D3O2A	Cell Signaling Tech.	14773	1:500
SEL1L	Rabbit	pAb	n/a	Sigma	S3699	1:50
KDEL	Mouse	IgG2a	10C3	Enzo Life Sci., Inc.	ADI-SPA-827-D	1:50
PDI	Rabbit	mAb IgG	C81H6	Cell Signaling Tech.	3501	1:50
Herp (HerpUD1)	Rabbit	pAb IgG	n/a	Proteintech	10813-1-AP	1:50
VCP	Mouse	IgG2a	5	Invitrogen / Life Technologies	MA3-004	1:50
Calreticulin	Rabbit	mAb IgG	D3E6	Cell Signaling Tech.	12238	1:200
Ribophorin 1 (RPN1)	Rabbit	pAb IgG		ThermoFisher / Invitrogen	PA5-65662	1:50
EDEM-1	Mouse	mAb IgM	D-1	Santa Cruz	sc-377394	1;50
GABARAP	Rabbit	mAb	E1J4E	Cell Signaling Tech.	13733	1:200
LC3B	Rabbit	mAb	D11	Cell Signaling Tech.	3868	1:200
HA epitope	Chicken	IgY		Bethyl	A190-106A	1:200
HCMV glycoprotein H (gH)	Mouse	IgG2b	14-4b	Gift of William J. Britt, M.D.	n/a	1: 50
Reticulon-3 (RTN3)	Mouse	IgG2a		ThermoFisher / Invitrogen	MA5-15538	1: 200
HA epitope	Rabbit	pAb		Bethyl	A190-108A	1:1000
Secondary Antibody	Host species	Isotype		Vendor	Catalog No.	Dilution used
Alexa Fluor 488 anti-Mouse IgG (H+L), cross adsorbed	Goat	IgG		ThermoFisher / Invitrogen	A11001	1:1000
Alexa Fluor 488 anti-Rabbit IgG (H+L) Cross-Adsorbed	Goat	IgG		ThermoFisher / Invitrogen	A11008	1:1000
Alexa Fluor 594 anti-Rabbit IgG (H+L) Cross-Adsorbed	Goat	IgG		ThermoFisher / Invitrogen	A11012	1:1000
Alexa Fluor 594 anti-Mouse IgG (H+L) Cross-Adsorbed	Goat	IgG		ThermoFisher / Invitrogen	A11005	1:1000
Alexa Fluor 647 anti-Chicken IgY (H+L)	Goat	IgG		ThermoFisher / Invitrogen	A11039	1:1000