A cytomegalovirus immunevasin triggers integrated stress response-dependent

reorganization of the endoplasmic reticulum.

Running Title: HCMV UL148 reorganizes the ER.

Hongbo Zhang[a,1], Clarissa Read[d,e,1], Christopher C. Nguyen[a,1], Mohammed N.A.

Siddiquey[a], Chaowei Shang[c], Cameron M. Hall[a], Jens von Einem[d], and Jeremy P.

Kamil[a,b #]

[a]Department of Microbiology and Immunology,

[b]Center for Molecular and Tumor Virology

[c]Research Core Facility

LSU Health Sciences Center, Shreveport, Louisiana, USA

[d] Institute of Virology, Ulm University Medical Center,

[e]Central Facility for Electron Microscopy, Ulm University

Ulm, Germany

[1] these authors contributed equally to the study

Running Head: HCMV UL148 Remodels the ER

#Address correspondence to Jeremy P. Kamil, jkamil@lsuhsc.edu

ABSTRACT. Human cytomegalovirus (HCMV) encodes an ER-resident glycoprotein,

UL148, which activates the unfolded protein response (UPR) but is fully dispensable for

viral replication in cultured cells. Hence, its previously ascribed roles in immune

evasion and modulation of viral cell tropism are hypothesized to cause ER stress. Here,

we show that UL148 is necessary and sufficient to drive the formation of large ER-

derived structures that occupy up to 7% of the infected cell cytosol. The structures are

found to be sites where UL148 coalesces together with cellular proteins involved in ER

quality control, such as Hrd1 and EDEM1. Ultrastructural analyses of the structures

reveal tortuous, densely packed segments of collapsed ER which connect to distended

cisternae. Notably, UL148 accumulates in a detergent-insoluble form during infection

while a homologous rhesus cytomegalovirus immunevasin that fails to cause ER

reorganization remains soluble. During induced ectopic expression of a UL148-GFP

fusion protein, punctate signals traffic to accumulate at prominent structures that exhibit

poor recovery of fluorescence after photobleaching. Small molecule blockade of the

integrated stress response (ISR) prevents the formation of puncta, leading to a uniform

reticular fluorescent signal. Accordingly, ISR inhibition during HCMV infection abolishes

the coalescence of UL148 and Hrd1 into discrete structures. Given that UL148

stabilizes immature forms of a receptor binding subunit for a viral envelope glycoprotein

complex of pivotal importance for HCMV infectivity, which is otherwise particularly

susceptible to ER associated degradation, our results imply that stress-dependent ER

remodeling contributes to viral cell tropism.

IMPORTANCE.

Perturbations to ER morphology occur during infection with various intracellular

pathogens and in certain genetic disorders. We identify that an HCMV gene product,

UL148, profoundly reorganizes the ER during infection, and is sufficient to do so when

expressed on its own. Our results reveal that UL148-dependent reorganization of the

ER is a prominent feature of HCMV infected cells. Moreover, we find that this example

of virally induced organelle remodeling requires the integrated stress response (ISR), a

stress adaptation pathway that contributes to a number of disease states. Since ER

reorganization accompanies the roles of UL148 in HCMV cell tropism and intracellular

retention of the immune cell co-stimulatory ligand CD58, our results may have

implications for understanding the mechanisms involved. Furthermore, our findings

provide a basis to utilize UL148 as a tool to investigate organelle responses to stress

and to identify novel drugs targeting the ISR.

INTRODUCTION.

UL148 is a human cytomegalovirus (HCMV) ER-resident glycoprotein that plays

roles in evasion of cell-mediated immunity and shows intriguing effects on cell tropism.

During infection of epithelial cells, viruses disrupted for *UL148* replicate to produce

roughly 100-fold enhanced levels of infectious progeny virions compared to wildtype (1).

These effects correlate with reduced expression of glycoprotein O (gO), a subunit of a

heterotrimeric viral glycoprotein H (gH) / glycoprotein L (gL) complex (gH/gL/gO) on the

virion envelope that is required for the infectivity of cell-free virions (2-4), and which

endows the virus with the capacity to utilize the platelet derived growth factor receptor α

(PDGFRα) as an entry receptor (5-7). Accordingly, UL148 has been found to stabilize

immature forms of gO prior their assembly into gH/gL/gO heterotrimers (1, 8). Despite

that UL148 does not stably associate with gO, the data suggest that it may interact with

gH (1).

UL148 also physically associates with CD58 (LFA-3), a co-stimulatory ligand for

natural killer cells and T-lymphocytes, preventing its presentation at cell surface (9).

Although the mechanisms by which UL148 stabilizes gO and retains CD58 within the

ER remain unknown, UL148 strongly contributes to activation of the unfolded protein

response (UPR) during infection, and is sufficient to activate the UPR when ectopically

expressed in non-infected cells (10). UL148 co-purifies from infected cells with SEL1L,

an adaptor subunit of ER-based E3 ubiquitin ligase Hrd1 that plays important roles in

ER-associated degradation (ERAD) of terminally misfolded glycoproteins (8). This

suggests a physical interaction with ERAD machinery, which may be germane to the

mechanism by which UL148 activates the UPR.

Here, we show that expression of UL148 is necessary and sufficient to induce

unusual ER structures at which large quantities of ER factors involved in glycoprotein

quality control accumulate. In electron microscopy analyses, we find that the UL148

induced ER-structures are comprised of densely packed, tortuous ER membranes that

form connections with tubules of highly distended cisternal space. Our results may

have implications for understanding the mechanisms by which UL148 regulates viral cell

tropism and contributes to viral evasion of cell-mediated immunity. Further, our findings

suggest that ER remodeling events triggered by UL148 are indicative of an adaptive

response of eukaryotic cells to proteotoxic stress involving the secretory pathway.

RESULTS.

UL148 causes reorganization of ER quality control proteins into unusual globular structures.

The HCMV ER-resident glycoprotein UL148 was previously observed to co-localize with the ER marker calnexin during infection (1). Nonetheless, calnexin staining did not show the uniform reticular pattern characteristic for the ER marker. We later noticed that cells infected with a *UL148*-null virus showed uniform calnexin staining (see below). To formally determine whether UL148 influences calnexin localization, we compared fibroblasts at four days post-infection with either wildtype (TB_WT) or *UL148*-null mutant (TB_148$_{STOP}$) viruses derived from an infectious bacterial artificial chromosome (BAC) clone of HCMV strain TB40/E (**FIG 1**). In cells infected with wildtype virus, calnexin antibodies stained unusual globular structures at the cell periphery, as expected (1) (**FIG 1A, 1C, SI FIG S1A**). However, in cells infected with the *UL148*-null virus, calnexin staining was uniform throughout the cytosol (**FIG 1B-C**), as would be expected for an ER marker in uninfected cells. The staining pattern for Hrd1, another ER marker, likewise indicated accumulation at unusual globular structures during wildtype HCMV infection, but not during infection with *UL148*-null mutant viruses (**FIG 1, SI FIG S1A**). Because calnexin and Hrd1 staining showed uniform distribution in *UL148*-null virus infected cells, and because *UL148* is fully dispensable for efficient viral replication in fibroblasts (1), these results suggest that redistribution of these ER markers depends on UL148.

As expected (11, 12), antibodies specific for the HCMV envelope glycoprotein,

glycoprotein H (gH), stained a juxtanuclear compartment, termed the cytoplasmic virion

assembly compartment (cVAC), which does not involve the ER (**FIG 1**). Similarly

contrasting staining patterns for Hrd1 and/or calnexin were observed in cells infected

with wildtype versus *UL148*-null mutants of clinical HCMV strains Merlin and TR (**FIG**

S1). Reciprocally, we restored a functional *UL148* at its native locus in the context of a

BAC clone of HCMV strain AD169, which spontaneously lost most of the gene in the

course of extensive genetic rearrangements and deletions that accumulated during

serial passage in tissue culture. In cells infected with AD169 repaired for UL148, but

not the *UL148*-null parental virus, the Hrd1 staining pattern showed obvious aggregation

into globular structures (FIG S1).

As similar differences in Hrd1 staining are observed between wildtype and

UL148-null infections of THP-1 macrophages, ARPE-19 epithelial cells and fibroblasts

(**FIG 1, FIG S1**), UL148-dependent reorganization of ER markers occurs in multiple cell

types. The calnexin staining pattern seen with four different primary clinical isolates

from patient throat swabs likewise indicated punctate globular structures. These

observations, taken together with the results from formal comparisons of wildtype

versus *UL148*-null mutants of four different BAC-cloned HCMV strains (**FIG 1, FIG S1**),

argue that UL148 profoundly affects the ER during natural infection.

We measured the three-dimensional volume of UL148-dependent Hrd1

structures from sixteen cells fixed at 96 hpi with strain TB40/E. On average, the

structures occupied 303.3 μm^3 (SEM: +/- 24.8) out of a total cell volume of 7657 μm^3

(SEM:+/- 651.0), as measured using a phalloidin-fluorophore conjugate to detect the

actin cytoskeleton (**FIG 1D-E**). Subtracting the volume of nuclei, as indicated by 4′,6-

diamidino-2-phenylindole [DAPI] (average nucleus: 1360 µm^3, SEM:+/-129.4), we

calculate that, on average, the structures occupy 5.2% of the cytosolic volume (SEM: +/-

0.61%, range: 2.0% – 7.0%). Based on these findings, taken together with results

comparing additional WT versus *UL148*-null mutant HCMV strains (**FIG S1**), we

conclude that the UL148-dependent ER structures are a prominent feature of HCMV-

infected cells.

UL148 substantially contributes to activation of the unfolded protein response

(UPR) during HCMV infection (10), and co-purifies from infected cells with SEL1L (8),

an adaptor subunit for the E3 ubiquitin ligase Hrd1, which plays crucial roles in ER-

associated degradation (ERAD) of terminally misfolded glycoprotein substrates (13).

Hence, the accumulation of Hrd1 and calnexin at unusual structures during wildtype but

not *UL148*-null infection may suggest that structures form in response to defects in ER

quality control (ERQC) caused by UL148. Under conditions of proteasome inhibition,

overexpression of certain misfolded glycoproteins causes cellular factors involved in

ERQC, such as calnexin, to compartmentalize from the rest of the ER, while other ER

markers such as BiP (Grp78) or protein disulfide isomerase (PDI) remain largely

unaltered (14-16).

To gain further insights into the nature of these peculiar ER structures, we set out

to develop a more comprehensive understanding of their protein composition by

comparing the localization of selected ER markers in cells fixed at 96 h post-infection

(hpi) either with TB_148^{HA} (1), a derivative of HCMV strain TB40/E that expresses at the

C-terminus of UL148 the nonapeptide epitope YPYDVPDYA from influenza A

hemagglutinin (HA), or with TB_159[HA], an isogenic control virus in which we replaced

the *UL148* coding sequence with the UL148 homolog from rhesus cytomegalovirus,

Rh159, likewise fused at its C-terminus to an HA-tag. We considered the TB_159[HA]

virus to be an appropriate control for the following reasons. Firstly, Rh159 and UL148

exhibit ~30% identity at the amino acid level and both glycoproteins localize to the ER

and block cell surface presentation immune cell activating ligands; UL148 retains CD58,

a ligand for CD2, while Rh159 retains NKG2D ligands of the MIC- and ULBP families (1,

9, 17). Secondly, Rh159 is expressed from TB_159[HA] at comparable levels and with

similar kinetics observed for UL148 from TB_148[HA], and the two viruses replicate

indistinguishably in fibroblasts (**FIG 2**). Moreover, Rh159 does not appear to activate

the UPR (10), and ERQC markers fail to coalesce into unusual structures during

infection with TB_159[HA] (**FIG 3**). Notably, ER structures likewise fail to occur in rhesus

fibroblasts during infection with a recombinant rhesus cytomegalovirus (RhCMV)

engineered to express an HA-tag fused at the C-terminus of Rh159 (**FIG S1**). This

argues against the possibility that Rh159 requires the context of rhesus cells to

redistribute ER markers in a manner analogous to what is seen for UL148 during HCMV

infection, and instead suggests that UL148 and Rh159 authentically differ in their effects

on the secretory pathway.

In cells infected with TB_148[HA], HA antibody staining indicated localization of

UL148 to globular structures, as expected (1). Antibody signals during indirect confocal

immunofluorescence detection of cellular ER resident proteins involved in ERQC,

including calnexin, Hrd1, SEL1L, Herp, valosin containing protein (VCP, p97), and

EDEM1, co-localized to a strong degree with signals from HA-tagged UL148 (**FIG 3-4**).

These results indicate that a number of ERQC factors coalesce with UL148 to form

prominent globular structures in infected cells. In contrast, uniform ER staining patterns

were observed with antibodies specific for PDI and calreticulin (CALR), indicating that

these ER markers do not localize to the UL148 structures (**FIGS 3Q, 3S**). Intriguingly,

antibody signals detecting reticulon 3 and ribophorin 1, which are markers for smooth

ER and rough ER, respectively, each appreciably co-localized with UL148 (HA) signal at

the induced structures (**FIGS 3K, 3M)**, which indicates that the structures may involve

both rough and smooth ER. LC3B failed to co-localize with the structures (**FIG S2**), as

would be expected given that the virus inhibits macroautophagy at late times during

infection (18, 19). Nevertheless, we obtained evidence suggesting the induced

structures are enriched for a related mammalian ATG8 ortholog, GABARAP (**FIG 3O**).

In cells infected with TB_159^{HA}, all of the ER markers we examined showed

uniform, reticular staining, as did the anti-HA signal detecting Rh159 (**FIG 3B, D, F, H,**

J, L, N, P, R, T). EDEM1, in addition to showing reticular staining, also labeled puncta

associated with the nucleus (**FIG 3J**), which may represent enriched levels of the

protein at the rough ER membranes associated with the nuclear envelope, although we

cannot exclude the possibility of spurious intranuclear staining. Nonetheless, the

punctate nuclear signal for EDEM1 was much less pronounced in the TB_148^{HA}

infected cells (**FIG 3I**). GABARAP antibodies stained the cVAC in TB_159^{HA} infected

cells, but also showed a much weaker reticular signal that may indicate ER staining.

Even though we took steps to block viral Fc receptors, which can cause rabbit

antibodies to non-specifically label the cVAC (20), it is plausible that GABARAP

antibody signal from the cVAC reflects incomplete blocking of viral Fc receptors. For

both viruses, antibodies specific for syntaxin-6 (STX6) and gH, as expected, stained the

juxtanuclear cVAC structure (11, 12, 21), which was previously found to be altogether

distinct from the UL148 staining pattern (1) (**FIG S2**).

To quantify the degree of co-localization with UL148, we calculated Pearson's

correlation coefficients from a minimum of thirty TB_148[HA] infected cells per staining

condition, comparing the signal overlap for each ER marker to the HA signal from

UL148. The correlation coefficients (r) for EDEM1, Hrd1, reticulon-3 (RTN3), SEL1L

and calnexin ranged from 0.8 to 0.72, which suggests that these proteins extensively

co-localize with UL148 at 4 days post-infection (dpi) (**FIG 4**). Meanwhile, GABARAP,

VCP, ribophorin-1 (RPN1), and Herp showed r values in the range of 0.54 to 0.51,

indicating a moderate degree of co-localization with UL148.   However, the ER markers

CALR and PDI, and the TGN marker syntaxin-6, gave r values of close to zero, which

confirms that these markers either do not co-localize with UL148 or show only a

negligible degree of association with UL148 at the induced structures.

From these results, we conclude that the UL148-dependent ER structures are

enriched for cellular markers associated with glycoprotein quality control. In this regard,

the structures resemble the "ER quality control (ERQC) compartments" described by G.

Lederkremer and colleagues (14, 15, 22). We further conclude that Rh159 cannot

substitute for UL148 to redistribute ERQC markers.

VCP and Hrd1 are recruited to incipient UL148 ER structures prior to calnexin.

To determine whether there might be differences in the kinetics of recruitment of

ERQC markers during formation of the ER structures, we examined a series of time

points from one to four days post-infection (dpi) with wild-type HCMV carrying HA-

tagged UL148, staining for three different ERQC markers, calnexin, Hrd1, and VCP,

alongside UL148 (HA). At 1 dpi the signal from UL148 was only faintly detected, as

expected (1), while each of the ER markers was readily detected, providing a readout of

their staining patterns prior to being substantially perturbed by UL148 (**FIG 5**). At 2 dpi,

we detected robust anti-HA signal, indicating the presence of UL148. At this time point,

UL148 exhibited intense signal at small globular puncta, which we interpret to represent

incipient UL148 structures, as well as more diffuse staining of a reticular structure

consistent with ER. Calnexin did not appreciably co-localize with the UL148 puncta until

at least 3 dpi, and the structures were not readily visualized by calnexin staining until 4

dpi (**FIG 5A**). In contrast, signals from Hrd1 and VCP staining were sufficient to mark

the UL148 puncta by 2 dpi, which suggests that these markers co-localize with UL148

at earlier points during the genesis of the ER structures. Notably, the appearance of

Hrd1 at the structures prior to calnexin is consistent with our previous results showing

that UL148 co-purifies from infected cells with SEL1L, an adaptor subunit for Hrd1 (8).

These results imply that basal elements of the ERAD machinery, exemplified by Hrd1

and VCP, may be recruited to UL148-induced ER structures prior to calnexin.

Visualization of UL148-induced ER structures by electron microscopy.

To discern the ultrastructural appearance of the UL148 structures and to formally

ascertain their relationship to the ER, we carried out transmission electron microscopy

(TEM) imaging of wildtype (TB_WT) and *UL148*-null (TB_148$_{STOP}$) infected fibroblasts at

5 dpi. In high pressure frozen and freeze-substituted infected fibroblasts, cells with a

high density of viral nucleocapsids within the nucleus were selected for analysis, as this

feature indicates late time points during infection when UL148 is abundantly expressed.

The TEM results revealed prominent globular and oblong structures in the cytoplasm of

wildtype virus infected fibroblasts, but not *UL148*-null infected controls (**FIGS 6-7**). The

structures stand out for their high electron-density, which may reflect the abundance of

ERQC proteins together with UL148 in these structures. Under higher magnification,

these areas are characterized by accumulations of densely packed ruffled membranes

and membranous material, which appear to be collapsed ER. Further, these membrane

accumulations were associated with smooth and partially rough ER structures with

seemingly enlarged cisternal space. Further, tomographic reconstruction of scanning

transmission electron microscopy (STEM) data suggest that densely packed ER

cisternae within the structures are interconnected and continuous in three-dimensional

space (**FIG 8B-C, SI Movie S1**).

UL148 accumulates in a detergent-insoluble form during infection.

Disease-associated variants of certain cellular proteins, such as the A103E

mutant of the calcium channel Orai1, and the E342K "Z" variant of alpha-1 antitrypsin (Z

A1AT), localize to anomalous ER structures reminiscent of those we observe to depend

on UL148 (16, 23-26). A103E Orai1 and Z A1AT are found to accumulate in detergent-

insoluble forms within the ER or within ER membranes, respectively, which indicates

aggregation or polymerization of these proteins, and suggests a mechanism for the

formation of ER structures by the proteins (16, 23, 26, 27). We therefore set out to

determine whether differences in solubility might correlate with the differing potentials

UL148 and Rh159 to cause ER reorganization and to activate the UPR (10). To

address this question, we infected fibroblasts with TB_148[HA] or TB_159[HA] at MOI 1 and

at various times post-infection prepared cell lysates in radioimmunoprecipitation assay

(RIPA) buffer. After subjecting the lysates to high-speed centrifugation, we examined

the relative levels of UL148 and Rh159 in the detergent-soluble supernatant and

detergent-insoluble pellet fractions.

A substantial portion of UL148 was detected from the detergent-insoluble

fractions at all time points tested (**FIG 9**). Furthermore, the percentage of UL148

detected within the insoluble fraction increased over time. In contrast, Rh159 was found

only in the detergent-soluble fraction. From these results, we conclude that UL148

accumulates in a detergent insoluble form during infection, and that Rh159 does not do

so. Because the relative mobility of the anti-HA immunoreactive band detected in

soluble and insoluble fractions obtained from TB_148[HA] infected cells showed a relative

mobility of ~35 kD, which is indicative of the mature, endoH-sensitive glycoprotein (1),

these findings suggest that UL148 may aggregate or polymerize within the ER.

UL148 is sufficient to compartmentalize the ER.

Our data thus far demonstrate that UL148 is necessary during infection to cause

redistribution of cellular ER markers for glycoprotein quality control processes, such that

substantial portion of the ERQC machinery appears to become sequestered away from

the rest of the organelle into novel membranous structures. Because a number of viral

proteins that remodel the ER during infection are sufficient to alter ER morphology when

ectopically expressed [e.g. (28),(29), reviewed in (30)], we wondered whether UL148

expression is sufficient to induce ER structures akin to those seen during HCMV

infection. Making use of existing "tet-on" ARPE-19 epithelial cells that inducibly express

of either UL148 or Rh159, each carrying a C-terminal HA-tag (31), we induced

transgene expression for 48 h and subsequently stained for various cellular markers for

the ER, for ERQC as well as for LC3B and GABARAP, which are ATG8 family proteins

that mark autophagosome membranes.

In cells expressing UL148 (i148^{HA}), we observed that calnexin, Hrd1, EDEM1,

and VCP co-localize with UL148 at prominent globular structures reminiscent of those

observed during infection (**FIG 10**). The respective rough and smooth ER markers

ribophorin 1 (RPN1) and reticulon 3 (RTN3), as well as the ATG8 proteins LC3B and

GABARAP likewise co-localized to the UL148-induced structures. However, PDI and

CALR each failed to substantially co-localize with UL148 at the structures (**FIG 10**), or at

best showed only limited co-localization, consistent with our results from infected cells

(**FIG 3-4**). Signals from antibodies specific for a KDEL motif important for ER retrieval

of luminal ER residents, such as BiP, likewise showed only limited co-localization with

UL148. These results suggest that large portions of the ER are not involved in the

UL148 structures.

In cells expressing the UL148 homolog Rh159 (i159^{HA}), we detected uniform

cytoplasmic distribution of ER markers, as well as of HA-tagged Rh159, similar to what

we observed during infection with the recombinant HCMV TB_159^{HA} (**FIG 3**).

Furthermore, In this setting the staining patterns for the ATG8 family proteins LC3B and

GABARAP failed to indicate any notable structures. Consistent with our previous study

showing that UL148, but not Rh159, is sufficient to activate the UPR (10), we observed

accumulation of ATF4 and phosphorylated eIF2α during induction of UL148 but not

Rh159 (**FIG 11A**). Importantly, the intensity of HA signals detecting UL148 and Rh159

indicated that both ER resident immunevasins accumulate at roughly comparable levels

following dox induction. This argues against the possibility that gross differences in

induced transgene expression might account for their differing effects on UPR activation

and on the staining patterns for ER markers.

Pearson's correlation coefficient values were calculated to quantify the extent of

colocalization between UL148 and various cellular markers: EDEM1, Hrd1, VCP,

calnexin (CNX), RTN3, SEL1L, LC3B, and PDI (**FIG 11B**). The results quantitatively

buttress the immunofluorescent staining data shown in FIG 10.   From these results, we

conclude that no other HCMV gene products are required for UL148 to cause ERQC

factors to segregate into discrete compartments. Hence, UL148 is sufficient to cause

large-scale reorganization of ERQC markers. Moreover, these findings suggest that

reorganization of ER markers into discrete structures may be related to the propensity

of UL148 to induce ER stress.

To determine whether UL148-dependent ER remodeling could be visualized in

real-time using live cell imaging, we constructed "tet-on" lentiviral vectors that inducibly

express either UL148 or Rh159 fused at their predicted C-terminal cytoplasmic tails to

the enhanced green fluorescent protein (GFP) from *Aequorea victoria* (32)*.* Following

lentiviral transduction, puromycin-resistant ARPE-19 were isolated and subsequently

subjected to fluorescence-activated cell sorting (FACS) to enrich for cells that

expressed GFP signal following dox treatment. The resulting cell populations, which

inducibly express either UL148-GFP or Rh159-GFP, were designated i148^{GFP} and

i159^{GFP}, respectively.

In live cell imaging studies, we observed that the GFP signal in dox-induced

i148^{GFP} cells first appeared in a reticular, largely uniform pattern, which was readily

visible at 5 h post-induction. However, by 9 h post-induction punctate signals began to

appear. These puncta were observed to traffic to sites of large scale accumulation,

where they formed larger fluorescent structures that progressively increased in size up

until the termination of the experiment at 19 h (**FIG 12A, SI Movie 2**). In contrast,

i159^{GFP} cells exhibited a largely diffuse, reticular GFP signal at all post-dox time points

monitored (**FIG 12B, SI Movie 3**). The live cell imaging results therefore recapitulated

the differences in the HA-staining pattern that we observed between dox-induced i148^{HA}

and i159^{HA} cells following fixation, as well as those between HA signals during HCMV

infection with TB_148^{HA} and TB_159^{HA} (**FIGS 1, 3, 10, SI Fig S1**).

UL148-GFP structures exhibit poor recovery of fluorescence after

photobleaching.

To determine whether the protein contents of the UL148 induced structures

would exhibit decreased mobility compared to non-perturbed ER, we carried out

fluorescence recovery after photobleaching studies. We photobleached regions of GFP

signal in i148^{GFP} or i159^{GFP} ARPE-19 cells that had been induced for transgene

expression for 24 h and monitored recovery of fluorescence over a 5 min period. In

Rh159-GFP expressing cells (i159^{GFP}), fluorescence almost fully recovered within 3 min,

and by 5 min fully recovered to pre-bleach levels (**FIG 13D**). In contrast, when we

photobleached a prominent UL148-GFP structure, the GFP signal failed to appreciably

recover fluorescence over the 5 min period (**FIG 13E**). Nonetheless, an area of reticular

GFP signal in an i148[GFP] cell that presumably represents a region of the ER not

involved in an anomalous structure, recovered fluorescent signal with kinetics similar to

those observed during Rh159-GFP expression. These results suggested to us that

UL148 structures do not exchange their contents as efficiently as unperturbed ER, or

that at least the UL148-GFP within the structures is not able to be replaced rapidly with

UL148-GFP fusion protein from other portions of the organelle.

Ectopically-expressed UL148 is degraded by proteasomes and not by autophagy.

The UL148 ER structures were found to be enriched with proteins such as Hrd1,

SEL1L, EDEM, and VCP (**FIG 3-4**, **10-11B**), which are posited to play key roles in

ERAD, a process during which misfolded glycoproteins are recognized, processed, and

dislocated across the ER membrane for degradation at cytosolic proteasomes.

However, we also detected elements of the machinery for autophagy in close

association with the UL148 structures. Namely, the mammalian ATG8 ortholog

GABARAP co-localized with UL148 during infection (**FIG 3-4**), and both GABARAP and

another ATG8 ortholog, LC3B, were found to co-localized with UL148 at the ER

structures during ectopic expression of UL148 (**FIG 10, 11B**). Therefore, we wished to

determine whether UL148 would degraded by the proteasome, as would be consistent

with conventional ERAD, or by the lysosome, which would suggest a role for autophagy-

related processes, such as selective autophagy of the ER (33), in dispensing with

UL148, and presumably, in resolving the ER perturbations.

As a first step, we conducted a live-cell imaging "washout" experiment in which

i148[GFP] cells were induced for transgene expression for 24 h, after which the growth

medium containing the transgene inducing agent (dox) was replaced with medium

lacking dox. During a 22 h imaging period following dox-washout, we observed the

structures to become progressively smaller as the reticular and punctate GFP signals

gradually abated (**FIG 14A, SI Movie S7**). Because the results suggested that UL148-

GFP structures could be resolved over time, we induced UL148-GFP in tet-on ARPE-19

cells (i148[gfp]) for 24 h, washed out the inducing agent (dox), and then applied either

epoxomicin (20 μM), an irreversible inhibitor of the proteasome (34), or folimycin (115

nM), a proton pump inhibitor that blocks autophagosome maturation and impedes the

lysosome-dependent protein degradation (35, 36). Of the two inhibitors, only

epoxomicin markedly stabilized UL148, as evidenced by the sustained Western blot

signal for both GFP and UL148 in this condition relative to either DMSO carrier or

folimycin treatments (**FIG 14B-C**). Despite that folimycin treatment had no obvious

effect on UL148-GFP levels, the drug markedly increased the levels of LC3B-II, as

would be expected with a lysosomal proton pump inhibitor. From these results, we

conclude that ectopically expressed UL148 is primarily degraded via a proteasome-

dependent mechanism.

UL148 requires the integrated stress response to cause ER reorganization.

We previously defined that UL148 triggers the UPR during ectopic expression,

and that UL148 contributes to activation of the PKR-like ER kinase (PERK) and inositol

requiring enzyme 1 (IRE1) during infection (10). Although we failed to find any effect of

small interfering RNA (siRNA) or drug treatments targeting IRE1 on UL148-dependent

ER remodeling (not shown), we did observe that siRNA directed at PERK caused

UL148 to fail to form puncta or to co-localize with Hrd1 (not shown). Furthermore, the

literature suggests that formation of ERQC compartments requires PERK (37). PERK

responds to ER stress by activating the integrated stress response (ISR). In particular,

PERK phosphorylates eIF2α at Ser51 (eIF2α-P), and the accumulation of eIF2α-P

causes global attenuation of mRNA translation whilst paradoxically stimulating

translation of select mRNAs, such as those encoding ATF4 and CHOP, which play roles

in adaptation to stress (38). Stress-regulated translation of such mRNAs involves small

upstream open reading frames (uORFs) in their 5' untranslated regions that ordinarily

suppress translation under basal conditions (39, 40).

To examine whether ER remodeling in response to UL148 requires the ISR, we

turned to a well characterized small molecule inhibitor of stress-regulated translation,

ISRIB (41-45). ISRIB is thought to function by acting as a "molecular staple" that holds

the guanine nucleotide exchange factor eIF2B in an active decameric configuration (43-

46), such that eIF2B will continue to generate ternary complex (eIF2•GTP•Met-tRNAi)

necessary for new cycles of translational initiation, despite the presence of phospho-

eIF2α (Ser51). Because PERK is the kinase that phosphorylates eIF2α in response to

ER stress (47, 48), we also tested for effects of the PERK inhibitor GSK2606414 (49).

Having confirmed that ISRIB and GSK260641 do not negatively impact UL148

expression during dox-induction of UL148-GFP **(FIG 15A-B)**, we treated i148[GFP] cells

with 200 nM ISRIB, 1.1 µM GSK260641, or DMSO vehicle control, and carried out live

cell imaging during dox induction.

Strikingly, ISRIB and GSK6060414 virtually abolished the formation of UL148

puncta through 13 h post induction (dox addition), a time point at which the DMSO

control condition showed abundant large structures (**FIG 15C-E, SI Movies 8-10**).

Based on our previous findings (**FIGs 1-3, 6-7, 10)**, we interpret the formation and

subsequent large-scale aggregation of UL148-GFP puncta to faithfully indicate of ER

reorganization in response to UL148. Therefore, these results suggested to us that

pharmacological treatments that inhibit the ISR and PERK prevent UL148 induced

remodeling of the ER, which further implies that ER remodeling in response to ER

stress requires stress-regulated translation.

Since our results with ISRIB and GSK6060414 suggest that the ISR is required

for UL148 mediated ER remodeling during ectopic expression of UL148, we next asked

whether these pharmacological agents would prevent UL148-dependent ER remodeling

in the physiologically authentic context of HCMV infection. However, our live cell

imaging studies with ISRIB and GSK6060414 indicated that following a pronounced

delay, UL148-GFP puncta nonetheless began to appear by ~19 h post treatment (**FIG**

15D-E, SI Movies 9-10), which suggested that the pharmacological activity of the drugs

might diminish over time. Therefore, in order to ensure that blockade of PERK and the

ISR would be sustained over a four-day period that it takes for structures to fully form

during HCMV infection, we replenished the drugs every 24 h by replacing the spent

media with media containing freshly reconstituted ISRIB, GSK6060414, or DMSO

vehicle control.

At 4 days postinfection, ER structures were virtually ablated in the ISRIB and

GSK2606414 treatment conditions (**FIG 16A**). To quantify these effects, we obtained Z-

stacks from a minimum of 30 cells per treatment condition and used Imaris 3D image

analysis software to calculate the percentage of Hrd1 signal that coalesced into discrete

structures during infection. The results indicated highly significant differences between

the DMSO carrier alone setting compared to treatments with either GSK2606414 or

ISRIB (**FIG 16B**). Reassuringly, roughly equivalent levels of UL148 were detected in

Western blot analyses of protein extracts from drug treated and DMSO control

conditions, which argues against the possibility that differences in UL148 expression

might explain the loss of structures involving UL148 and Hrd1. Furthermore, ISRIB and

GSK2606414 treatments both caused reduced levels of ATF4, while PERK inhibitor

alone was able to reduce the levels phospho-eIF2α (Ser51), detected by a phospho-

specific antibody. Therefore, each of the drugs caused their expected effects on the

PERK / eIF2α / ATF4 axis (**FIG 16C-D**).

Notably, a modest reduction in the levels of the viral envelope glycoprotein B

(gB) was observed during treatments with ISRIB or GSK2606414. This suggests that

the ISR may be required for optimal expression of the certain viral envelope

glycoproteins. Even viruses lacking *UL148* cause high levels of ATF4 accumulation at

late times during infection, which is when viral late gene products, such a gB, are

expressed at their highest levels (10). Hence, the activity of PERK, and/or ISR

regulated translation may be required for optimal expression of viral envelope

glycoproteins. Regardless, the PERK inhibitor and ISRIB each prevented the

appearance of discrete UL148 ER structures while failing to substantially affect UL148

expression. Therefore, the effects of the two inhibitors on UL148-dependent ER

reorganization were fully consistent with those seen during ectopic expression of the

immunevasin. Taken together, our findings argue that the ISR is required for UL148 to

cause remodeling of the organelle.

DISCUSSION

The ER structures that that we have identified are noteworthy in several regards.

Firstly, this example of virally-induced ER remodeling is wholly dependent on a single

viral gene product, UL148. Of course, there are other examples of individual viral gene

products that are necessary to grossly perturb ER morphology during infection which

are also found to be sufficient to induce such perturbations [e.g., (28, 29, 50, 51)]

Nonetheless, to the best of our knowledge the ER structures caused by UL148 are a

hitherto undescribed ultrastructural characteristic of the HCMV infected cell, which is

surprising given their size, scale, and prominence. Indeed, the average volume of the

structures at 96 h postinfection amounts to roughly 60% of the size of the nucleus of an

uninfected human fibroblast [~500 µm^3, (52)]. Although UL148 expression is

accompanied by UPR activation, ER reorganization occurs without killing the host cell,

even in the setting of ectopic expression when viral functions that inhibit programmed

cell death are absent. Nor does UL148 affect the yield of infectious virus during

replication in fibroblasts, since *UL148*-null mutant viruses replicate indistinguishably

from wildtype parental virus in this cell type (1, 8). As with all enveloped viruses, HCMV

requires the host cell secretory pathway to fold and process enormous quantities of viral

envelope glycoproteins that endow progeny virions with the molecular machinery for

infectivity. It is intriguing to consider how the infected cell tolerates spatial

reorganization of ER membranes and the associated glycoprotein quality control

machinery without impacting the production of infectious progeny virions.

Furthermore, certain aspects of ER reorganization invoked by UL148 may be

novel to cell biology. In particular, the densely packed collapsed ER in close

association with varicosities that we observe to depend on UL148 during infection

appear to differ from the ER structures induced by other ER perturbagens. For

instance, a hereditary form of childhood cirrhosis caused by the *Z* allele of the alpha-1-

antitrypsin (Z-A1AT) gene (*SERPINA*) is characterized by polymerization of Z-A1AT

within the ER (25), leading to its accumulation in membrane-delimited inclusions (25,

26, 53). However, Z-A1AT inclusions are not observed to associate with regions of

collapsed ER, nor does Z-A1AT suffice to activate the UPR, even though its expression

sensitizes cells to other triggers of ER stress (26, 54).

HCMV encodes at least one other ER-resident immunevasin that activates the

UPR, US11 (55). UL148 binds the NK-cell and T-cell costimulatory ligand CD58 to

prevent its transport to the cell surface while US11 targets the heavy chain of class I

major histocompatibility complex (MHC) for ER-associated degradation (56, 57). UL148

causes CD58 to accumulate as an immature glycoform, presumably within the ER, but

does not lead to any obvious decrease in its overall abundance (9). Albeit that

intracellular forms of CD58 appear to be refractory to detection by standard indirect

immunofluorescence protocols (not shown), it seems reasonable to hypothesize that

UL148 sequesters CD58 into the unusual ER structures that it induces. Moreover, our

detergent solubility results suggest that UL148 may form aggregates or polymers in the

course of preventing CD58 presentation at the cell surface (FIG 9). It is worth noting

that UL148, an immunevasin targeting CD58, induces dramatic morphologic

rearrangements of the ER, while another HCMV immunevasin, US11, targeting MHC

class I for degradation evidently does not do so, even though both trigger ER stress.

Nonetheless, it seems unlikely that retention of CD58 would be required for ER

reorganization. In fact, the capacity of UL148 to retain CD58 may be functionally

separable from its peculiar effects on the morphology and organization of the ER.

Despite that the relationship between the mechanism for CD58 retention and the

formation of UL148-dependent ER structures remains unresolved, our findings may

suggest a mechanism to explain the influence of UL148 the tropism of the virus for

epithelial cells. These effects, which are exemplified by a ~100-fold replication

advantage of *UL148*-null viruses in epithelial cells, correlate with decreased expression

of glycoprotein O (gO) (1). We previously reported that gO behaves as a constitutive

ERAD substrate during infection and that immature, newly synthesized forms of gO

show enhanced stability in the presence of UL148 (8), and we now show that UL148

causes large-scale sequestration of the proteasomal ERAD factors, such as the ER

mannosidase EDEM1 and the E3 ubiquitin ligase SEL1L/Hrd1, into large membranous

structures. Collectively, these observations may suggest that regions of rough ER not

drawn into the structures might offer a more permissive folding environment for

polypeptides that are slow to fold or assemble into multiprotein complexes.

Indeed, our results indicate that the reticular ER, as indicated by the staining

pattern for calreticulin, PDI, and KDEL antibodies, remains largely intact following

UL148-induced ER reorganization of ER quality control markers into discrete structures.

Based on these findings, we hypothesize that UL148 alters ER proteostasis by

physically dislocating key elements of the "mannose removal time clock" system that

marks poorly folding glycoproteins for destruction via ERAD (58, 59). Because gO is

both the defining subunit of the heterotrimeric gH/gL/gO envelope glycoprotein complex

that governs HCMV entry and cell tropism (5-7, 60, 61), and a constitutive ERAD

substrate (1, 8), the hypothesis that ER reorganization is required for the effects of

UL148 on HCMV cell tropism does not seem unreasonable on its face. Going forward,

it will be crucial to determine whether classical ERAD substrates, such as the null Hong

Kong variant of alpha-1-antitrypsin (62, 63) or ribophorin-332 (64, 65), are stabilized

during UL148 expression, as would be predicted if UL148 shifts ER proteostasis to

negatively modulate ERAD.

In addition to having found that factors involved in proteasomal ERAD, such as

Hrd1 and EDEM1, are enriched at the UL148 structures, we also observed that

GABARAP, a mammalian ortholog of yeast ATG8, localizes to the UL148 ER structures

during infection, and that another ATG8 ortholog, LC3B, and GABARAP both associate

with the structures during ectopic expression of UL148. This may suggest roles for

autophagy-related pathways in UL148-dependent reorganization of the organelle.

Since our results indicate that UL148 is degraded by a proteasome-dependent pathway

and not by lysosomes, it seems unlikely that selective autophagy of the ER is directly

involved in recycling or degrading these ER structures. Nonetheless, ATG8 family

proteins may contribute to formation of large structures, vis-a-vis the trafficking of

UL148-GFP puncta observed during our live cell imaging studies. Previous work

indicates that misfolded proteins traffic in a microtubule (MT)-dependent manner to form

pericentriolar structures termed aggresomes (66, 67). ATG8 family proteins such as

LC3B and GABARAP not only play roles in degradation of substrates via

macroautophagy, but also bind MTs and recruit machinery for MT-dependent transport

of cargos (68-70). Therefore, GABARAP and LC3B may be important for the

recruitment of cellular machinery that transports perturbed ER cargoes to sites of large-

scale accumulation.

Although that our results argue that the ISR is required for ER reorganization

during UL148 expression, precisely how UL148 triggers ER stress remains unknown.

UL148 has been found to co-purify from cells with SEL1L, a component of the ERAD

machinery. Thus, it seems plausible that UL148 may inhibit the Hrd1/SEL1L complex,

which would cause the buildup of unfolded proteins and thus trigger the UPR. However,

inhibition of ERAD in and of itself seems unlikely to account for the formation of ER

structures. Another non-mutually exclusive possibility is that UL148 multimerizes or

aggregates in a manner that constricts the ER lumen. For instance, the assembly of

UL148 molecules on opposite sides of the ER lumen might constrict the organelle in a

manner consistent with the collapsed regions of ER observed in our EM results (FIGS 6,

8).

Despite that additional work will certainly be needed to decipher the molecular

mechanisms by which UL148 causes reorganization of the ER, and to determine its

physiological relevance to viral biology. Nonetheless, we have shown that UL148, when

fused to a fluorescent protein, suffices to trigger and robustly indicate the ISR (FIGS 15-

16), which suggests that UL148 could be used in screens to identify novel inhibitors of

the ISR, as well as in genetic screens to identify cellular genes involved in stress-

dependent remodeling of the ER. Interestingly, the two pharmacological agents that we

have identified to block UL148-dependent ER remodeling, ISRIB and GSK2606414, are

also known to disperse stress granules (SGs) and to prevent their formation in the first

place. SGs are comprised of condensed aggregates of protein and RNA that occur due

to stalled mRNA translation in disease states, such as amyotrophic lateral sclerosis

(71), and during treatments with toxic agents such as arsenite (42, 72).   It is fascinating

to consider that mechanistic parallels exist between the formation of SGs and the

UL148-dependent reorganization of the ER. Moreover, given the broad importance

macromolecular aggregation in pathological conditions such as neurodegenerative

diseases (73), certain of which also involve defects in ER proteostasis and aberrant

activation of the UPR, UL148 may hold promise as tool to discover new agents to

ameliorate disease.

MATERIALS AND METHODS.

Cells and Virus.

hTERT-immortalized human foreskin fibroblasts (8), derived from ATCC HFF-1 cells

(SCRC-1041) were maintained in Dulbecco's modified Eagle's medium supplemented

with 5%-10% newborn calf serum (Millipore Sigma) and antibiotics (complete DMEM)

exactly as described previously (8). i148^{HA} and i159^{HA} ARPE-19 epithelial cells (10),

which upon treatment with 100 ng/mL doxycycline, express HA-tagged UL148 or

Rh159, respectively, were likewise maintained in complete DMEM. For live cell imaging

studies, i148^{HA} and i159^{HA} ARPE-19 were maintained in Opti-MEM (Thermo Fisher)

supplemented with 3% certified tet-approves fetal bovine serum (FBS), 20 µg/mL

gentamicin sulfate, 1 µg/mL puromycin HCl and 10 µg/mL ciprofloxacin HCl.

Telomerase-immortalized rhesus fibroblasts (74) were a kind gift of Peter A. Barry, and

were maintained in complete DMEM. The acute monocytic leukemia cell line THP-1

(TIB-202) was obtained from ATCC (Manassas, VA) and were maintained in RPMI 1640

medium supplemented with 10% FBS (Millipore Sigma) and antibiotics (10 µg/mL

ciprofloxacin and 25 µg/mL gentamicin). THP-1 were differentiated into adherent

macrophages by incubating for 48 h in the presence of 100 nM 2-O-

tetradecanoylphorbol 13-acetate (Millipore Sigma), and subsequently infected with the

indicated viruses at an MOI of 5 TCID$_{50}$/cell.

Infectious bacterial artificial chromosome (BAC) clones of the following HCMV

strains were used for this study: TB40/E (also known as TB40-BAC4 or TB_WT) (75)

as well as its derivatives TB_148^{HA} and TB_148$_{STOP}$, which were used in our previous

studies (1, 8, 10); TR (TR*gfp*)(76, 77); Merlin repaired for *UL128* and *RL13* harboring

tetO sequences upstream of *UL131* (pAL1393) (78); AD169 repaired for *UL131*

(AD_r131)(1, 79) and AD_r131_148[HA], a derivative of AD_r131 that carries a full length

UL148 (from strain TB40/E) tagged with an HA-epitope at the original *UL148* locus (8).

An infectious BAC clone of rhesus CMV (RhCMV) strain 68-1 (80) and a derivative that

expresses an HA-tagged Rh159 (details below) were also used for certain experiments.

The methods used for reconstitution of HCMV from purified BAC DNA, cultivation

of virus and preparation of stocks, including ultracentrifugation through sorbitol cushions

and determination of infectious titers by tissue culture infectious dose 50 (TCID$_{50}$) assay

have been described elsewhere (8, 10).

Construction of recombinant viruses and recombinant DNAs.

New recombinant viruses for this study were derived from BAC-cloned HCMV

and RhCMVs using *en passant* mutagenesis in GS1783 *E. coli,* a strain K12 derivative

that expresses the homing endonuclease I-SceI upon treatment with L-(+)-arabinose

(81, 82). Recombinant BACs were confirmed by Sanger DNA sequencing of the

modified regions, which was performed by Genewiz, Inc. (Piscataway, NJ) (not shown).

Oligonucleotide primers for construction of recombinant viruses and plasmids were

custom synthesized by Integrated DNA Technologies (Coralville, IA); sequences are

provided in **Table S1**. Type II restriction endonucleases, T4 DNA ligase, and Gibson

assembly reagents (NEB HiFi Assembly Master Mix) for generation of recombinant

DNAs and for routine molecular cloning and subcloning procedures were obtained from

New England Biolabs (Ipswitch, MA). KOD Hot Start DNA polymerase (EMD Millipore)

was used for all PCR reactions.

To construct TB_159^{HA}, a strain TB40/E derivative in which the UL148 open

reading frame is replaced by Rh159 we carried out the following steps. Primers

Rh159_Fw and Rh159_HA_Rv were used to amplify and HA-tag the *Rh159* ORF from

pcDNA3.1+_Rh159_IRES_GFP (a gift from Klaus Früh, OHSU). The PCR product was

inserted into EcoRV-digested pEF1α using Gibson assembly. A PCR product

containing an ISce-I excisable kanamycin cassette was amplified from TB_148^{HA}_*ISce-*

Kan integrate BAC (1) using primers PpuISceKanGibs_Fw and PpuISceKanGibs _Rv

and Gibson-assembled into PpuMI-digested pEF1α_Rh159 plasmid to yield plasmid

pEF1α_Rh159_ISceKan. Primers Rh159_Fw_recomb and Rh159_Rv_recomb were

used to generate a PCR product from template plasmid pEF1α_Rh159_ISceKan. The

PCR product was electroporated into GS1783 E. coli harboring the TB_148^{HA} BAC, and

Kan^r colonies harboring TB_Δ148_Rh159_ISceKan integrate BACs were isolated on

Luria-Bertiani agar plates. Bacterial colonies representing Kan resistant integrates were

obtained, and integrates were subsequently resolved using standard *en passant*

protocols to yield TB_Δ148_Rh159^{HA}, which we abbreviate herein as TB_159^{HA}.

TB_159^{HA} was sequence-confirmed using primers TB_Rh159HA_seq_Fw and REV

TB_Rh159HA_seq. Similar strategies were used to modify the RhCMV 68-1 BAC (80)

so as to insert sequences encoding the nonapeptide HA-epitope such that Rh159 would

be expressed from the virus carrying a C-terminal HA tag. An *en passant* strategy in

which a shuttle plasmid carrying *I-SceI-Kan^r* disrupted by in-frame nonsense codons

was used insert premature nonsense codons in the *UL148* CDS in the context of HCMV

strains Merlin (pAL1393) and TR (TR*gfp*), and was applied as described previously for

generating TB_148$_{STOP}$, the UL148$_{STOP}$ mutant of strain TB40/E (8).

To construct "tet-on" lentivirus vector plasmids containing UL148 fused to GFP,

we used primers Gibs_eGFP_Rv and 148_eGFP_Fw to amplify the GFP gene from a

dsDNA gBlock, EGFP-P2A-3XHA, synthesized by Integrated DNA Technologies, gift of

Matthew D. Woolard, LSUHSC-S.   In a separate PCR, the *UL148* gene was amplified

from plasmid pcDNA3.1-UL148^{HA} (10) using primers Gib_148_Fw

148_noStopRv. The two products were assembled together with EcoRV opened

pcDNA3.1(+) (Invitrogen) to produce pcDNA3.1-UL148-GFP.   After confirming the

absence of spurious mutations by Sanger DNA sequencing (Genewiz), the *UL148-GFP*

cassette from pcDNA3.1-UL148-GFP was released by EcoRI and NotI digestion and

ligated into pOUPc turboRFP-link plasmid which had been linearized using the same

restriction sites. pOUPc turboRFP-link is a derivative of lentiviral vector pOUPc (10)

that was constructed by reinsertion of turboRFP (tRFP). Briefly, tRFP was amplified

from the original pOUPc-turboRFP using primer pair tRFP linker Fw and tRFP linker Rv

and then reinserted by Gibson Assembly with pOUPc turboRFP that had been double-

digested using EcoRI and Not I in order to linearize the vector and remove the

turboRFP (RFP) sequence. To express Rh159 fused to eGFP, primers Gibs_eGFP_Rv

and 159_eGFP_Fw were used to amplify the GFP gene from EGFP-P2A-3XHA, and in

a separate PCR reaction, primers Gibs_159_Fw and 159_noStop_Rv Rh159 were

used to amplify *Rh159* from pcDNA3.1-Rh159^{HA} (10). The latter two PCR products were

assembled together with EcoRV linearized pcDNA3.1(+) using Gibson Assembly,

resulting in plasmid pcDNA3.1-Rh159. The latter plasmid was used as template in a

PCR reaction with primers Gibs_Age_159_Fw and Gibs_Mlu_GFP_Rv. The resulting

PCR product was Gibson assembled into the lentiviral vector pOUPc (10) which had

been double-digested with Mlu I and Age I.

Drug treatments.

The PERK inhibitor, GSK2606414 (49), doxycycline hyclate, and folimycin were

obtained from MilliporeSigma (Burlington, MA). ISRIB (41-45) was obtained from

MilliporeSigma or APExBio (Boston, MA) and epoxomicin was procured Selleck

Chemicals (Houston, TX).   ISRIB was dissolved in DMSO to make a 10,000× stock

solution (2 mM) and used at a final working concentration of 200 nM. GSK2606414 was

dissolved in DMSO to make a 10,000× stock solution (11 mM) and used at a final

working concentration of 1.1 µM. Folimycin was prepared as a 1000× (115.46 µM)

stock solution in DMSO and used at 115 nM final. Epoxomycin was prepared as a 100×

(2 mM) stock solution in DMSO and used at a final concentration of 20 µM.

Confocal Microscopy and Live Cell Imaging. Confocal indirect immunofluorescence

microscopy imaging on fixed cells was carried out using Leica TCS SP5 Spectral

Confocal Microscope (Leica Microsystems, Heidelberg, Germany) using a Leica HCX

PL APO CS 63x/1.4-0.6NA objective under oil immersion, except for the image shown

in Figure 1C, which was captured on a Nikon SIM-E and A1R confocal microscopy

system (Nikon Instruments, Melville, NY) using a Nikon SR Apo TIRF 100x/1.49NA

objective under oil immersion. Images for different fluorophore channels were acquired

using sequential scanning. Direct immunofluorescence live cell imaging data were

collected using the Nikon SIM-E microscope using a Nikon Apo 60x/1.40NA DIC

objective.

The 3D projections shown in FIGs 1D and 16A were generated from Z-stacks

captured on the Nikon SIM-E / A1R microscope using a Nikon SR Apo TIRF

100x/1.49NA objective. For FIG 1D, the 3D projection was generated using NIS-

Elements AR Analysis 4.60.00 (64-bit) software (Nikon). For FIG 16A, 3D projections

were generated by Imaris x64 9.3.0 software (Bitplane, Inc.) in maximum intensity

projection (MIP) mode.

For fixed cell imaging experiments other than those shown in Fig S1 panels E, F,

G, and H, cells were seeded on 12 mm circular, No. 1 thickness microscope cover glass

(200121; Azer Scientific, Morgantown, PA). At the indicated times post-treatment and or

post-infection, cells were washed with phosphate buffered saline (PBS) [137 mM NaCl,

2.7 mM KCl, 10 mM Na_2HPO_4, and 1.8 mM KH_2PO_4, pH 7.4]; fixed for 15 min at room

temperature in PBS containing 4% (wt/vol) paraformaldehyde (Fisher Scientific, Waltham,

MA), washed in PBS, permeabilized for 10 min using 0.1% Triton X-100 (in PBS),

subsequently washed again in PBS, and then blocked for 45 min at 37 °C in PBS

containing 5% (vol/vol) normal goat serum (Rockland Immunochemicals, Limerick, PA).

Cells were then washed three times in PBS followed by incubation in 1% Human Fc Block

(BD Biosciences, San Jose, CA) in PBS, for additional 45 min at 37 °C. Cells were then

incubated in the presence of primary antibodies for 1 h at 37 °C or 4 °C overnight, and

then washed three times with PBS containing 0.1% Tween-20 (PBST) for 5 min per wash.

Alexa Fluor-labeled goat polyclonal secondary antibodies (all from ThermoFisher

Invitrogen, Waltham, MA, see **Table S2**) were used for secondary detection. The slides

were then mounted using Prolong Gold anti-fade reagent containing DAPI

(ThermoFisher) and placed under a Leica TCS SP5 confocal microscope for image

acquisition using a Leica 63× oil immersion objective lens (Leica Microsystems, Buffalo

Grove, IL).

For indirect immunofluorescence staining results of primary clinical isolates (SI Fig

S1 panels E-H), four clinical HCMV isolates obtained by routine testing of throat swabs

from patients of the Ulm University Medical Center were provided by the diagnostic

laboratory of the Institute of Virology in Ulm. Sample material was applied to human

fibroblasts and incubated for several days until HCMV-positive cells could be detected.

Infected cells were then seeded together with uninfected fibroblasts, incubated for up to

5 days until plaques were formed and processed for indirect immunofluorescence

staining. ERQC compartments were detected by staining for calnexin (CNX, E10; Santa

Cruz Biotechnology, mouse, 1:50 dilution), the cVAC was detected by staining for HCMV

pUL71 (rabbit, 1:500). Secondary antibody used for detection of CNX was Alexa Fluor

555 labeled goat anti-mouse IgG (1:1000) and for pUL71 detection, Alexa Fluor 488

conjugated goat anti-rabbit IgG (1:1000). Confocal images were acquired using the 63×

objective of a Zeiss Observer Z1 fluorescence microscope equipped with Apotome and

Zen software 2.3 (Carl Zeiss Microscopy GmbH, Jena, Germany).

FRAP. FRAP studies were carried out on using live ARPE-19 cells from the i148[GFP]

and i159[GFP] populations as follows. Cells were seeded as above for live-cell imaging,

induced for transgene expression using dox (100 ng) for 24 h, and then placed on an

incubated sample stage of a Nikon A1R SIM-E imaging system equipped with an SR

Apo TIRF 100x/1.49NA objective (Nikon). Three rectangular regions in image fields

were defined for measurement of (i) background signal, and of two regions with

comparable initial GFP signals, (ii) one designated as a control region (no

photobleaching) and another for (iii) photobleaching and recovery of signal after

photobleaching. Photobleaching of selected regions was carried out for 20 s using 405

nm laser light from a LU-N4 laser fiber (Nikon) [power at source: 15 mW, at objective: 8

mW]. Images and signal intensity measurements were captured every 2 s at a rate of 1

frame per s (488 nm excitation, FITC channel) starting immediately before

photobleaching (t=1), and from t=20 s to t=320 s after.

Electron Microscopy. Procedures to prepare samples for transmission electron

microscopy (TEM) included high-pressure freezing (HPF), freeze substitution, and Epon

embedding, which were carried out as described previously (83). Briefly, human

fibroblasts were seeded in µ-Slides (Ibidi GmbH, Martinsried, Germany) containing

carbon-coated sapphire discs (Engineering Office M. Wohlwend GmbH) 1 day prior to

infection at 80 to 90% confluence. Cells were infected with virus overnight at MOI 1.

Medium containing viral inocula was replaced with fresh medium the next day. Infected

cells on sapphire discs were fixed by using HPF with a Compact 01 high-pressure freezer

(Engineering Office M. Wohlwend GmbH, Sennwald, Switzerland) at 5 dpi. Thereafter,

cells on sapphire discs were processed by freeze-substitution and subsequently

embedded in Epon (Fluka, Buchs, Switzerland). Ultrathin sections of the Epon-embedded

cells were cut with an ultramicrotome (Ultracut UCT; Leica Microsystems, Wetzlar,

Germany) and placed on Formvar-coated single-slot grids (Plano GmbH, Wetzlar,

Germany). Grids were examined in a Jeol JEM-1400 (Jeol Ltd., Tokyo, Japan)

transmission electron microscope equipped with a charge-coupled-device (CCD) camera

at an acceleration voltage of 120 kV. Fixation and embedding of infected cells for

scanning transmission electron microscopy (STEM) tomography was the same as

described for TEM. Additional sample preparation steps were conducted as described

previously (84, 85). Tomogram acquisition was conducted on a STEM Jeol JEM-2100F

with an acceleration voltage of 200 kV. Tilt series were acquired from 600 nm thin

sections from +72° to -72° with a 1.5° increment using the bright field detector. Image

series were reconstructed to tomograms by weighted back projection with the IMOD

software package (86). Three-dimensional visualization of the membrane structures was

performed using Avizo lite software (Visualization Science Group, Burlington, MA, USA)

by threshold segmentation.

Western blotting. Western blotting procedures, including primary and secondary

antibodies used for detection of HA tag, ATF4, eIF2α, P-eIF2α (Ser51), the HCMV viral

nuclear antigen IE1, and the conditions used for detection of Ser51 phosphorylated

eIF2α, were carried out as described previously (8, 10). Additional antibodies used in

this study were mouse anti-GAPDH (Cat. No. 60004-1, Proteintech, Rosemont, IL,),

mouse anti-gB clone 27-180 (87) (a generous gift of William J. Britt), and anti-GFP

(D5.1) XP® Rabbit mAb #2956 (Cell Signaling Technology, Danvers, MA).

Solubility analyses. 2.0×10^5 human fibroblasts were infected at an MOI of 1 TCID$_{50}$

per cell with TB_148[HA] or TB_159[HA]. The following day, cells were washed twice with

PBS to remove viral inoculum and replenished with DMEM containing 5% newborn calf

serum. At the indicated times post-infection, cells were washed once with PBS and

lysed by direct addition of 100 μL RIPA buffer [25 mM HEPES (pH 7.5), 400 mM NaCl,

0.1% SDS, 0.5% sodium deoxycholate, 1% NP-40, supplemented with $1\times$ protease

inhibitor cocktail (ApexBio)]. Lysates were collected and rotated at 4°C for 1 h.

Insoluble material was pelleted by centrifugation (21,000 × g, 35 min). Supernatants

containing soluble material were transferred to a fresh microfuge tube, and 33 μL of 4×

Laemmli sample buffer [200 mM Tris (pH 6.8), 8% SDS, 40% glycerol, 0.08%

bromophenol blue] was added to bring final volume to 133 μL. The pellet was disrupted

in 133 μL of 1X Laemmli buffer prepared by diluting 4× Laemmli buffer in RIPA buffer.

Samples were reduced by addition of beta-mercaptoethanol (5% final, v/v) and boiled at

90°C for 10 min. 40 μL of each sample was resolved by SDS-PAGE (12%

polyacrylamide gel), transferred overnight to nitrocellulose membrane, and

immunoblotted with antibodies against HA epitope or HCMV gB. Quantitation of

secondary antibody fluorescence signals were performed using an Odyssey CLx

scanner (Li-Cor, Inc., Lincoln, NE) in auto-scan mode. For each time-point, the signals

from RIPA-soluble and insoluble bands were summed to yield total signal, and the ratio

of insoluble band signal to total signal were also reported as percent insoluble HA over

total HA signal.

Statistical analyses. Statistical analyses were carried out using GraphPad Prism 8.1.0

for MacOS (GraphPad, Inc., San Diego, CA).

ACKNOWLEDGEMENTS.

This project was supported by NIH grants R01-AI116851 (to J.P.K.) and P30-

GM110703. Its contents are solely the responsibility of the authors and do not

necessarily represent the official views of the NIAID or the NIGMS.

We are especially grateful to Erik L. Snapp (HHMI, Janelia Laboratories) for

helpful discussions concerning interpretation of EM and confocal image data. We also

thank Jacob E. Corn and Jin Rui "Amos" Liang (both of ETH Zurich, Switzerland), Dong

Yu (Washington University, St. Louis, MO, currently GSK Vaccines), Thomas E. Shenk

(Princeton University, Princeton, NJ), Richard J. Stanton (Cardiff University, United

Kingdom), William J. Britt (University of Alabama, Birmingham), W. L. William Chang

and Peter A. Barry (both of the University of California, Davis; Davis, CA), Christian

Sinzger (University Medical Center Ulm, Ulm, Germany), Klaus Früh (Oregon Health

Sciences University, Beaverton, OR), and Gregory A. Smith (Northwestern University,

Chicago, IL) for generously sharing reagents.

AUTHOR CONTRIBUTIONS.

Performed experiments: HZ, CR, CCN, MNAS, JvE. Data analysis and interpretation:

HZ, CR, CCN, JvE, CS, JPK. Wrote the manuscript: JPK with comments from JvE and

CR.   Conceived of the study: JPK.   Designed experiments: HZ, JvE, CR, CCN.

Contributed new reagents: HZ, CH, CCN, JPK.   Electron microscopy: CR and JvE.

Protein solubility analysis: CCN. Obtained funding: JPK.

HZ performed all confocal immunofluorescence experiments with the exception of the SI

Figure on primary clinical isolates from patient throat swabs (JvE), viral replication kinetics

experiments, and also constructed novel reagents, including lentivirus vector plasmids for

isolating i148-GFP and i159-GFP ARPE-19 cell populations. HZ also carried out FRAP

studies, and performed all of the live cell imaging experiments, and all Western blots in

the study, except those otherwise specified below. CR carried out the TEM and STEM

tomography studies. CCN constructed the TB_159^{HA} virus and conceived of and

performed the protein solubility analyses comparing UL148 and Rh159. MS carried out

Western blotting studies on the effects of folimycin versus epoxomicin on UL148

degradation. CH constructed the RhCMV 68-1 BAC carrying an HA-tagged Rh159. CS

assisted with confocal imaging and with Imaris and Nikon software analyses of the 3D

images.

 FIGURE LEGENDS:

FIG 1. UL148 reorganizes ER markers into anomalous structures during HCMV

infection.

Fibroblasts infected at 1 MOI with either **(A)** wildtype HCMV strain TB40/E (TB_WT) or

(B) an isogenic *UL148*-null mutant (TB_148$_{STOP}$), were fixed at 96 h postinfection (hpi)

and imaged by confocal microscopy after staining with antibodies specific for calnexin or

Hrd1 (magenta) and glycoprotein H (gH, green), as indicated. DAPI (blue) signal was

used to counterstain nuclei. **(C)** Percentage of fibroblasts that contain Hrd1 structures at

96 hpi; fifty gH-positive cells for each condition were scored for the presence or absence

of UL148 structures (also see SI Fig S1A). (**D**) Three-dimensional confocal image

projection of a TB_WT infected fibroblast stained at 96 hpi for Hrd1 and gH. **(E)**

Volumetric measurements were made from sixteen infected cells, stained with phalloidin-

AlexaFluor 594 conjugate to estimate total cell volume, with Hrd1 antibody to estimate

the volume of UL148-dependent ER structures, and with DAPI to estimate nuclear

volume.

FIG 2. Characterization of TB_148^{HA} and TB_159^{HA} viruses. (A) Human fibroblasts

were infected at MOI 1 with the indicated recombinant HCMVs and whole cell lysate

samples harvested at the indicated times post-infection (h post-infection: hpi) were

analyzed by Western blot to detect HA-tagged UL148 or Rh159, the 72 kD viral nuclear

antigen IE1-72 (IE1), and GAPDH. **(B)** Single-cycle viral replication kinetic curves from

MOI 1 infected fibroblasts were plotted by determining the titer in tissue culture infectious

dose 50 ($TCID_{50}$) from supernatants collected at the indicated times postinfection.

FIG 3. UL148 localizes to unusual ER compartments enriched for glycoprotein

quality control markers. Fibroblasts infected at MOI 1 with either TB_148^{HA} (panels: **A,**

C, E, G, I, K, M, O, Q, S) or TB_ 159^{HA} (panels: **B, D, F, H, J, L, N, P, R, T**) were fixed

at 96 hpi, and imaged by confocal microscopy after co-staining with antibodies specific

for HA (UL148/Rh159, magenta in merge) and the indicated cellular markers (green in

merge). DAPI (blue) counterstaining is shown in merged image.

FIG 4. Quantification of Co-localization between UL148 and Cellular Markers. A

Pearson's correlation coefficient (*r*) was calculated using NIH ImageJ software to estimate

the degree of co-localization between UL148 (HA signal) and each of the indicated

cellular markers. A minimum of 30 cells were analyzed per marker. The arithmetic mean

for each co-localization analysis result is shown as a blue line, and data points for

individual cells analyzed are plotted as circles.

FIG 5. Cellular proteins involved in glycoprotein quality control are recruited with

differing kinetics to UL148 ER structures. Fibroblasts infected with TB_148^{HA} at 1 MOI

were fixed at the indicated time points (days post infection, dpi) and imaged by confocal

microscopy after staining with antibodies specific for HA (UL148, magenta), CNX (green,

A) , Hrd1 (green, **B**) or VCP(green, **C**), DAPI (blue).

FIG 6. TEM of ER structures in wild-type HCMV infected cells. Human fibroblasts

infected with wildtype HCMV (TB_WT) were fixed by high-pressure freezing and freeze

substitution at day 5 postinfection and imaged using TEM. Panels (**A**) and (**B**) show cell

overview at left. For each cell, the boxed region is shown at higher magnification.

Scale bars; left panels: 10 µm, right panels: 2 µm. rER: rough ER; Nu: nucleus; cVAC:

cytoplasmic viral assembly compartment. Solid arrowheads indicate the UL148-

dependent ER structures of interest.

FIG 7. TEM of *UL148*-null HCMV infected cells. Human fibroblasts infected with a

UL148-null mutant (TB_148$_{STOP}$) were fixed by high-pressure freezing and freeze

substitution at day 5 postinfection and imaged using TEM. At the upper left an overview

panel of a representative cell is shown with panels **A, B, and C** each boxed. The boxed

regions are expanded at higher magnification at right (**A**) and below (**B, C**). Scale bars

for upper left overview panel: 10 µm, for zoomed panels: 2 µm. rER: rough ER; Nu:

nucleus; cVAC: cytoplasmic viral assembly compartment.

FIG 8. STEM tomography of UL148-dependent ER structures in HCMV infected

cells. Human fibroblasts infected with wildtype HCMV (TB_WT) were fixed by high-

pressure freezing and freeze substitution at day 5 postinfection and tomograms were

recorded by STEM. (**A**) Shows a virtual section through the tomogram of a virus-

induced membranous structure; asterisks denote ER structures of distended lumenal

space. White arrowheads indicate sites at which membranes originating from distended

ER cisternae continue into areas of involuted collapsed ER. Scale bar: 1 µm. (**B**)

Shows the same virtual section as in (A) tilted and with a 3D visualization of the

membranous network (green) of the entire tomogram. The same distended ER

cisternae as in (A) are marked by asterisks. (**C**) Shows a cross section through (B) to

visualize the membrane profile of the membranous structures. The region delimited by

a white box is shown in a higher magnification on the right. Finer detail of the enlarged

ER cisternae and the connections between them are readily visible; asterisks indicate

the same distended ER cisternae as in (A) and (B). Also, see SI Movie S1. Scale bar: 1

µm.

FIG 9. Solubility Analysis of UL148 and Rh159. (**A**) Human fibroblasts were infected at

MOI 1 $TCID_{50}$/cell with HCMV strain TB40/E derived viruses TB_148^{HA}, which

expresses UL148 fused at its C-terminus to an HA-epitope tag, or TB_159^{HA}, which

lacks *UL148* and instead expresses rhesus CMV Rh159 carrying a C-terminal HA tag.

At the indicated times postinfection (hpi), infected cells were collected

radioimmunoprecipitation (RIPA) lysis buffer, centrifuged at 21,000$\times$g for 30 min, after

which supernatant (sup) and pellet (pel) fractions were boiled in gel loading buffer

containing 2% sodium dodecyl sulfate (SDS). Equivalent portions of supernatant and

pellet were resolved by SDS-polyacrylamide gel electrophoresis, transferred to a

nitrocellulose membrane for detection of protein species immunoreactive to antibodies

against HA epitope (HA) and HCMV glycoprotein B (gB) and for total protein signal

using Ponceau S reagent (Ponceau). (**B-C**). Signal intensity of fluorophore-conjugated

secondary antibodies in anti-HA Western blots were measured from three independent

biological replicates of the experiment shown in (A); error bars indicate standard

deviation. (**B**) The fluorescent signal for each infection time point condition (Y-axis

indicates arbitrary units, in hundred thousands). (**C**) The amount of signal found in the

insoluble (pellet) fraction relative to the total signal (pellet plus supernatant) for each

infection time point are plotted as percentage values.

FIG 10. UL148 is sufficient to remodel the ER. HA-tagged UL148 or Rh159 were

doxycycline (dox) induced in the context of a "tet-on" lentiviral vector system in stably

transduced ARPE-19 epithelial cell populations, i148^{HA} and i159^{HA}, respectively. Cells

were fixed at 48 h postinduction for indirect immunofluorescence staining for the

indicated cellular markers (green) together with HA (magenta). Scale bar: 10 µm.

FIG 11. ISR activation accompanies redistribution of ER markers during ectopic

expression of UL148. (A) Lysates of tet-on ARPE-19 cells expressing either UL148 or

Rh159 fused to an HA tag, i148HA and i159HA, respectively, were collected at the

indicated times post doxycycline (dox) induction and analyzed by Western blot for the

expression of the indicated proteins and for the abundance of eIF2α phosphorylated at

Ser51 (P-eIF2α) using a phospho-specific antibody. (**B**) Pearson's correlation

coefficient (*r*) values were calculated using NIH ImageJ software estimate the degree of

co-localization between UL148 (HA signal) and the indicated cellular markers. A

minimum of 30 cells were analyzed per marker. Arithmetic means for each co-

localization analysis result are shown as blue lines, and data points for individual cells

analyzed are plotted as circles.

FIG 12. Live Cell Imaging of UL148-GFP and Rh159-GFP during Induced Ectopic Expression. "Tet-on" ARPE-19 epithelial cells that inducibly express either UL148 or Rh159 fused to green florescent protein (gfp), i148^{gfp} (**A**) and i159^{gfp} (**B**), respectively, were induced for transgene expression using 100 ng/mL doxycycline (dox) and imaged using live-cell microscopy. Images from the selected time points (h post treatment with dox, hpt) are shown. The main scale bar represents 50 μm. For inset panels at upper left of each image, which are magnified 2.4× relative to the main image, the scale bar represents 10 μm. Also see SI Movies S2-S3.

FIG 13. FRAP analysis. ARPE-19 cells that inducibly express Rh159-GFP, i159^{gfp} (**A**) or UL148-GFP, i148^{gfp} (**B-C**) were doxycycline induced for 24 h (100 ng/mL doxycycline (dox). Separate regions of GFP signal in selected cells were photobleached (405 nm laser) or left unbleached, while a third region lacking GFP signal was chosen as a background reference were measured before and during fluorescence recovery after photobleaching (FRAP). Note: background signal was not plotted because values were resolvable from the x-axis. GFP signal intensity is plotted over a time period (seconds, s) starting with an exposure at t=0 (immediately before photobleaching), and including measurements taken every 2 s after photobleaching (0-20 s) until termination of the measurement series at t=322 s (300 s of FRAP). (**D-F**) Images from the selected time points (s, seconds) immediately before and after bleaching, and during fluorescence recovery period. Also see SI Movies S4-S6.

1008

FIG 14: UL148-GFP is stabilized by inhibition of proteasomal but not lysosomal

degradation. "Tet-on" ARPE-19 cells that inducibly express UL148-GFP (i148^{gfp}) were

induced for 24 h by the addition of 100 ng/mL doxycycline (dox), after which the dox

inducing agent was washed out and medium containing either the proteasome inhibitor

epoxomicin (20 µM) or the proton pump inhibitor folimycin (115 nM) was added and

samples were harvested for Western blot analysis at the indicated times post treatment

(h post treatment, hpt) with folimycin or epoxomicin. Also see SI Movie S7.

1016

1017

FIG 15. Inhibition of the integrated stress response impedes UL148-mediated ER

remodeling. "Tet-on" ARPE-19 epithelial cells that inducibly express either UL148 or

Rh159 fused to green florescent protein (gfp), i148^{gfp} (**A**) and i159^{gfp} (**B**), respectively,

were induced for transgene expression using 100 ng/mL doxycycline (dox) in the

presence of absence of either ISRIB (200 nM) or GSK2606414 (1.1 µM) and monitored

by Western blot for expression of UL148-GFP over a series of time points (h post

treatment with dox, hpt). (**C-E**): Live-cell imaging of UL148-GFP and Rh159-GFP

expression patterns in the presence of ISRIB (200 nM), GSK2606414 (1.1 µM), or

DMSO vehicle alone. Main scale bar represents 50 µm. For inset panels at upper left of

each image, which are magnified 2.4× relative to the main image, the scale bar

represents 10 µm. Also see SI Movies S8-S10.

1029

1030

FIG 16. Inhibition of the ISR prevents the coalescence of Hrd1 and UL148 into discrete structures during infection. (**A**) Representative 3D maximum intensity projections of confocal imaging Z-stacks obtained from cells infected at MOI 1 for 96 h with HCMV strain TB40/E carrying an HA-tagged *UL148*, TB_148^{HA}, and maintained in the presence of ISRIB (200 nM), GSK2606414 (1.1 µM), or DMSO carrier alone (0.1% vol/vol). In merged images HA signal is shown in magenta, Hrd1 in green, and DAPI counterstaining in blue. (**B**) The percent of Hrd1 antibody signal involved in discrete structures at 96 hpi were calculated for a minimum of 30 cells per condition were determined using Imaris x64 9.3.0 software. Statistical significance was determined using a one-way ANOVA followed by Tukey's post-test; **** represents a P-value of <0.0001. The arithmetic mean for co-localization analysis results are shown as blue lines with data points for individual cells analyzed plotted as circles. (**C-D**) Western blot analyses of fibroblasts infected at MOI 1 with TB_148^{HA} and maintained in the presence of ISRIB (200 nM), GSK2606414 (1.1 µM), or DMSO carrier alone (0.1%); hpi: h post infection (hpi). Note: a phospho-specific antibody was used for detection of eIF2α phosphorylated at Ser51 (P-eIF2α).

SI FIGURES and MOVIES.

FIG S1: **Additional indirect immunofluorescent confocal microscopy results from infected cells.** (**A**) Lower magnification overview of results from FIGS 1A-B; staining of HCMV glycoprotein H (gH) and Hrd1 at 96 h postinfection (hpi) of human fibroblasts with wildtype HCMV strain TB40/E (WT) versus *UL148*-null derivative TB_148$_{STOP}$. (**B**)

Staining of gH and Hrd1 at 96 hpi of human fibroblasts with wildtype Merlin recovered

from BAC-cloned Merlin pAL1393 (ME_WT) or a *UL148*-null derivative of the same

virus (ME_148$_{STOP}$.) (**C**): Staining of gH and Hrd1 at 96 hpi of human fibroblasts with

wildtype HCMV strain TRgfp recovered from BAC-cloned TRgfp (TR_WT) or a *UL148*-

null derivative of the same virus (TR_148$_{STOP}$). (**D**) Staining of the viral tegument protein

UL71 and calnexin at 5 days postinfection of human fibroblasts with primary clinical

HCMV isolates obtained from patient throat swabs. (**E**) Staining against HA and Hrd1 in

THP-1 macrophages fixed at 96 hpi with HCMV strain TB40/E derivative viruses

TB_148^{HA} or TB_159^{HA}; note: THP-1 monocytes were differentiated using phorbol ester

treatment. (**F**) Staining of HA and Hrd1 in ARPE-19 epithelial cells infected for 96 h with

either TB_148^{HA} or TB_159^{HA}. (**G**) Staining of HA and Hrd1 in ARPE-19 epithelial cells

infected for 96 h with HCMV strain AD169 repaired for *UL131* and to which an HA-

tagged UL148 CDS from TB40/E was restored to the native *UL148* locus (AD^{r131}_148^{HA})

or with a parental AD169 virus repaired for *UL131* to which *UL148* was not restored

(AD^{r131}). Notes: (i) a functional *UL131* is required for efficient infection of epithelial cells;

(ii) the difference in scale bar is because with ADr131 we did not employ a viral marker

(e.g., HA) to identify infected cells, so the appearance of syncytia and the characteristic

kidney-bean shaped nucleus was used to indicate infected cells and a slightly lower

magnification was used to best show these features. (**H**) Staining for HA and Hrd1 in

telomerase-immortalized rhesus fibroblasts infected with BAC-derived rhesus CMV

(RhCMV) strain 68-1 carrying an HA-tag at the C-terminus of Rh159.

1075

1076 **FIG S2: Confocal microscopy of LC3B, syntaxin-6 and glycoprotein H staining**

1077 **during HCMV infection.** Human fibroblasts were infected with the indicated viruses for

1078 96 h and then fixed, permeabilized, and stained using antibodies specific for the

1079 indicated proteins. Indirect immunofluorescence images were captured using a 63X

1080 objective on a Leica SP5 confocal microscope. gH: glycoprotein H, STX6: syntaxin-6.

1081

1082 **Movie S1: STEM tomography of anomalous ER structures in wildtype HCMV**

1083 **infected fibroblasts at day 5 postinfection.**

1084

Movies S2 – S7: Live cell imaging of GFP signal following dox induction of

transgene expression in i148^{gfp} and i159^{gfp} ARPE-19 epithelial cells:

Movie S2: i148^{gfp} cells from 2-19 h post dox induction.

Movie S3: i159^{gfp} cells from 2-19 h post dox induction.

Movie S4: **FRAP of UL148-GFP structure (ER within structure)**

Movie S5: FRAP of UL148 GFP structure (ER outside structure)

Movie S6: FRAP of Rh159-GFP.

1092

Movie S7: Imaging of UL148-GFP following washout of doxycycline. (to 21 h 37

min post washout).

Movie S8: i148^{gfp} cells in the presence of 0.1% DMSO, from 2-19 h post dox

induction.

Movie S9: i148^{gfp} cells in the presence of 1.1 µM GSK2606414, from 2-19 h post

dox induction.

Movie S10: i148^{gfp} cells in the presence of 200 nM ISRIB, from 2-19 h post dox

induction.

1101

1102

LITERATURE CITED

1. Li G, Nguyen CC, Ryckman BJ, Britt WJ, Kamil JP. 2015. A viral regulator of glycoprotein complexes contributes to human cytomegalovirus cell tropism. Proc Natl Acad Sci U S A 112:4471-6.

2. Wille PT, Knoche AJ, Nelson JA, Jarvis MA, Johnson DC. 2010. A human cytomegalovirus gO-null mutant fails to incorporate gH/gL into the virion envelope and is unable to enter fibroblasts and epithelial and endothelial cells. J Virol 84:2585-96.

3. Jiang XJ, Adler B, Sampaio KL, Digel M, Jahn G, Ettischer N, Stierhof YD, Scrivano L, Koszinowski U, Mach M, Sinzger C. 2008. UL74 of human cytomegalovirus contributes to virus release by promoting secondary envelopment of virions. J Virol 82:2802-12.

4. Zhou M, Lanchy JM, Ryckman BJ. 2015. Human Cytomegalovirus gH/gL/gO Promotes the Fusion Step of Entry into All Cell Types, whereas gH/gL/UL128-131 Broadens Virus Tropism through a Distinct Mechanism. J Virol 89:8999-9009.

5. Wu Y, Prager A, Boos S, Resch M, Brizic I, Mach M, Wildner S, Scrivano L, Adler B. 2017. Human cytomegalovirus glycoprotein complex gH/gL/gO uses PDGFR-alpha as a key for entry. PLoS Pathog 13:e1006281.

6. Kabanova A, Marcandalli J, Zhou T, Bianchi S, Baxa U, Tsybovsky Y, Lilleri D, Silacci-Fregni C, Foglierini M, Fernandez-Rodriguez BM, Druz A, Zhang B, Geiger R, Pagani M, Sallusto F, Kwong PD, Corti D, Lanzavecchia A, Perez L. 2016. Platelet-derived growth factor-alpha receptor is the cellular receptor for human cytomegalovirus gHgLgO trimer. Nat Microbiol 1:16082.

7. Wu K, Oberstein A, Wang W, Shenk T. 2018. Role of PDGF receptor-alpha during human cytomegalovirus entry into fibroblasts. Proc Natl Acad Sci U S A doi:10.1073/pnas.1806305115.

8. Nguyen CC, Siddiquey MNA, Zhang H, Li G, Kamil JP. 2018. Human Cytomegalovirus Tropism Modulator UL148 Interacts with SEL1L, a Cellular Factor That Governs Endoplasmic Reticulum-Associated Degradation of the Viral Envelope Glycoprotein gO. J Virol 92:e00688-18.

9. Wang ECY, Pjechova M, Nightingale K, Vlahava VM, Patel M, Ruckova E, Forbes SK, Nobre L, Antrobus R, Roberts D, Fielding CA, Seirafian S, Davies J, Murrell I, Lau B, Wilkie GS, Suarez NM, Stanton RJ, Vojtesek B, Davison A, Lehner PJ, Weekes MP, Wilkinson GWG, Tomasec P. 2018. Suppression of costimulation by human cytomegalovirus promotes evasion of cellular immune defenses. Proc Natl Acad Sci U S A 115:4998-5003.

10. Siddiquey MNA, Zhang H, Nguyen CC, Domma AJ, Kamil JP. 2018. The Human Cytomegalovirus Endoplasmic Reticulum-Resident Glycoprotein UL148 Activates the Unfolded Protein Response. J Virol 92:e00896-18.

11. Das S, Vasanji A, Pellett PE. 2007. Three-dimensional structure of the human cytomegalovirus cytoplasmic virion assembly complex includes a reoriented secretory apparatus. J Virol 81:11861-9.

12. Sanchez V, Greis KD, Sztul E, Britt WJ. 2000. Accumulation of virion tegument and envelope proteins in a stable cytoplasmic compartment during human cytomegalovirus replication: characterization of a potential site of virus assembly. J Virol 74:975-86.

1146   13.   Lilley BN, Ploegh HL. 2005. Multiprotein complexes that link dislocation, ubiquitination,
1147   and extraction of misfolded proteins from the endoplasmic reticulum membrane. Proc
1148   Natl Acad Sci U S A 102:14296-301.
1149   14.   Kamhi-Nesher S, Shenkman M, Tolchinsky S, Fromm SV, Ehrlich R, Lederkremer GZ.
1150   2001. A novel quality control compartment derived from the endoplasmic reticulum.
1151   Mol Biol Cell 12:1711-23.
1152   15.   Leitman J, Shenkman M, Gofman Y, Shtern NO, Ben-Tal N, Hendershot LM, Lederkremer
1153   GZ. 2014. Herp coordinates compartmentalization and recruitment of HRD1 and
1154   misfolded proteins for ERAD. Mol Biol Cell 25:1050-60.
1155   16.   Houck SA, Ren HY, Madden VJ, Bonner JN, Conlin MP, Janovick JA, Conn PM, Cyr DM.
1156   2014. Quality control autophagy degrades soluble ERAD-resistant conformers of the
1157   misfolded membrane protein GnRHR. Mol Cell 54:166-179.
1158   17.   Sturgill ER, Malouli D, Hansen SG, Burwitz BJ, Seo S, Schneider CL, Womack JL, Verweij
1159   MC, Ventura AB, Bhusari A, Jeffries KM, Legasse AW, Axthelm MK, Hudson AW, Sacha
1160   JB, Picker LJ, Fruh K. 2016. Natural Killer Cell Evasion Is Essential for Infection by Rhesus
1161   Cytomegalovirus. PLoS Pathog 12:e1005868.
1162   18.   Chaumorcel M, Souquere S, Pierron G, Codogno P, Esclatine A. 2008. Human
1163   cytomegalovirus controls a new autophagy-dependent cellular antiviral defense
1164   mechanism. Autophagy 4:46-53.
1165   19.   Chaumorcel M, Lussignol M, Mouna L, Cavignac Y, Fahie K, Cotte-Laffitte J, Geballe A,
1166   Brune W, Beau I, Codogno P, Esclatine A. 2012. The human cytomegalovirus protein
1167   TRS1 inhibits autophagy via its interaction with Beclin 1. J Virol 86:2571-84.
1168   20.   Alwine JC. 2012. The human cytomegalovirus assembly compartment: a masterpiece of
1169   viral manipulation of cellular processes that facilitates assembly and egress. PLoS Pathog
1170   8:e1002878.
1171   21.   Das S, Pellett PE. 2011. Spatial relationships between markers for secretory and
1172   endosomal machinery in human cytomegalovirus-infected cells versus those in
1173   uninfected cells. J Virol 85:5864-79.
1174   22.   Leitman J, Ron E, Ogen-Shtern N, Lederkremer GZ. 2013. Compartmentalization of
1175   endoplasmic reticulum quality control and ER-associated degradation factors. DNA Cell
1176   Biol 32:2-7.
1177   23.   Cox DW, Billingsley GD, Callahan JW. 1986. Aggregation of plasma Z type alpha 1-
1178   antitrypsin suggests basic defect for the deficiency. FEBS Lett 205:255-60.
1179   24.   Le A, Ferrell GA, Dishon DS, Le QQ, Sifers RN. 1992. Soluble aggregates of the human PiZ
1180   alpha 1-antitrypsin variant are degraded within the endoplasmic reticulum by a
1181   mechanism sensitive to inhibitors of protein synthesis. J Biol Chem 267:1072-80.
1182   25.   Lomas DA, Evans DL, Finch JT, Carrell RW. 1992. The mechanism of Z alpha 1-antitrypsin
1183   accumulation in the liver. Nature 357:605-7.
1184   26.   Ordonez A, Snapp EL, Tan L, Miranda E, Marciniak SJ, Lomas DA. 2013. Endoplasmic
1185   reticulum polymers impair luminal protein mobility and sensitize to cellular stress in
1186   alpha1-antitrypsin deficiency. Hepatology 57:2049-60.
1187   27.   Ekeowa UI, Freeke J, Miranda E, Gooptu B, Bush MF, Perez J, Teckman J, Robinson CV,
1188   Lomas DA. 2010. Defining the mechanism of polymerization in the serpinopathies. Proc
1189   Natl Acad Sci U S A 107:17146-51.

28. Suhy DA, Giddings TH, Jr., Kirkegaard K. 2000. Remodeling the endoplasmic reticulum by
poliovirus infection and by individual viral proteins: an autophagy-like origin for virus-
induced vesicles. J Virol 74:8953-65.
29. Windsor M, Hawes P, Monaghan P, Snapp E, Salas ML, Rodriguez JM, Wileman T. 2012.
Mechanism of collapse of endoplasmic reticulum cisternae during African swine fever
virus infection. Traffic 13:30-42.
30. Miller S, Krijnse-Locker J. 2008. Modification of intracellular membrane structures for
virus replication. Nat Rev Microbiol 6:363-74.
31. Siddiquey MNA, Zhang H, Nguyen CC, Domma AJ, Kamil JP. 2018. The human
cytomegalovirus ER resident glycoprotein UL148 activates the unfolded protein
response. J Virol doi:10.1128/JVI.00896-18.
32. Cormack BP, Valdivia RH, Falkow S. 1996. FACS-optimized mutants of the green
fluorescent protein (GFP). Gene 173:33-8.
33. Khaminets A, Heinrich T, Mari M, Grumati P, Huebner AK, Akutsu M, Liebmann L, Stolz
A, Nietzsche S, Koch N, Mauthe M, Katona I, Qualmann B, Weis J, Reggiori F, Kurth I,
Hubner CA, Dikic I. 2015. Regulation of endoplasmic reticulum turnover by selective
autophagy. Nature 522:354-8.
34. Meng L, Mohan R, Kwok BH, Elofsson M, Sin N, Crews CM. 1999. Epoxomicin, a potent
and selective proteasome inhibitor, exhibits in vivo antiinflammatory activity. Proc Natl
Acad Sci U S A 96:10403-8.
35. Muroi M, Shiragami N, Nagao K, Yamasaki M, Takatsuki A. 1993. Folimycin
(concanamycin A), a specific inhibitor of V-ATPase, blocks intracellular translocation of
the glycoprotein of vesicular stomatitis virus before arrival to the Golgi apparatus. Cell
Struct Funct 18:139-49.
36. Muroi M, Takasu A, Yamasaki M, Takatsuki A. 1993. Folimycin (concanamycin A), an
inhibitor of V-type H(+)-ATPase, blocks cell-surface expression of virus-envelope
glycoproteins. Biochem Biophys Res Commun 193:999-1005.
37. Kondratyev M, Avezov E, Shenkman M, Groisman B, Lederkremer GZ. 2007. PERK-
dependent compartmentalization of ERAD and unfolded protein response machineries
during ER stress. Exp Cell Res 313:3395-407.
38. Harding HP, Novoa I, Zhang Y, Zeng H, Wek R, Schapira M, Ron D. 2000. Regulated
translation initiation controls stress-induced gene expression in mammalian cells. Mol
Cell 6:1099-108.
39. Palam LR, Baird TD, Wek RC. 2011. Phosphorylation of eIF2 facilitates ribosomal bypass
of an inhibitory upstream ORF to enhance CHOP translation. J Biol Chem 286:10939-49.
40. Vattem KM, Wek RC. 2004. Reinitiation involving upstream ORFs regulates ATF4 mRNA
translation in mammalian cells. Proc Natl Acad Sci U S A 101:11269-74.
41. Sidrauski C, Acosta-Alvear D, Khoutorsky A, Vedantham P, Hearn BR, Li H, Gamache K,
Gallagher CM, Ang KK, Wilson C, Okreglak V, Ashkenazi A, Hann B, Nader K, Arkin MR,
Renslo AR, Sonenberg N, Walter P. 2013. Pharmacological brake-release of mRNA
translation enhances cognitive memory. Elife 2:e00498.
42. Sidrauski C, McGeachy AM, Ingolia NT, Walter P. 2015. The small molecule ISRIB
reverses the effects of eIF2alpha phosphorylation on translation and stress granule
assembly. Elife 4.

43.   Zyryanova AF, Weis F, Faille A, Alard AA, Crespillo-Casado A, Sekine Y, Harding HP, Allen
F, Parts L, Fromont C, Fischer PM, Warren AJ, Ron D. 2018. Binding of ISRIB reveals a
regulatory site in the nucleotide exchange factor eIF2B. Science 359:1533-1536.
44.   Sidrauski C, Tsai JC, Kampmann M, Hearn BR, Vedantham P, Jaishankar P, Sokabe M,
Mendez AS, Newton BW, Tang EL, Verschueren E, Johnson JR, Krogan NJ, Fraser CS,
Weissman JS, Renslo AR, Walter P. 2015. Pharmacological dimerization and activation of
the exchange factor eIF2B antagonizes the integrated stress response. Elife 4:e07314.
45.   Tsai JC, Miller-Vedam LE, Anand AA, Jaishankar P, Nguyen HC, Renslo AR, Frost A, Walter
P. 2018. Structure of the nucleotide exchange factor eIF2B reveals mechanism of
memory-enhancing molecule. Science 359.
46.   Gordiyenko Y, Schmidt C, Jennings MD, Matak-Vinkovic D, Pavitt GD, Robinson CV. 2014.
eIF2B is a decameric guanine nucleotide exchange factor with a gamma2epsilon2
tetrameric core. Nat Commun 5:3902.
47.   Harding HP, Zhang Y, Ron D. 1999. Protein translation and folding are coupled by an
endoplasmic-reticulum-resident kinase. Nature 397:271-4.
48.   Harding HP, Zhang Y, Bertolotti A, Zeng H, Ron D. 2000. Perk is essential for translational
regulation and cell survival during the unfolded protein response. Mol Cell 5:897-904.
49.   Axten JM, Medina JR, Feng Y, Shu A, Romeril SP, Grant SW, Li WH, Heerding DA,
Minthorn E, Mencken T, Atkins C, Liu Q, Rabindran S, Kumar R, Hong X, Goetz A, Stanley
T, Taylor JD, Sigethy SD, Tomberlin GH, Hassell AM, Kahler KM, Shewchuk LM, Gampe
RT. 2012. Discovery of 7-methyl-5-(1-{[3-(trifluoromethyl)phenyl]acetyl}-2,3-dihydro-1H-
indol-5-yl)-7H-p yrrolo[2,3-d]pyrimidin-4-amine (GSK2606414), a potent and selective
first-in-class inhibitor of protein kinase R (PKR)-like endoplasmic reticulum kinase
(PERK). J Med Chem 55:7193-207.
50.   Tenorio R, Fernandez de Castro I, Knowlton JJ, Zamora PF, Lee CH, Mainou BA, Dermody
TS, Risco C. 2018. Reovirus sigmaNS and muNS Proteins Remodel the Endoplasmic
Reticulum to Build Replication Neo-Organelles. MBio 9.
51.   Miller S, Kastner S, Krijnse-Locker J, Buhler S, Bartenschlager R. 2007. The non-structural
protein 4A of dengue virus is an integral membrane protein inducing membrane
alterations in a 2K-regulated manner. J Biol Chem 282:8873-82.
52.   Lamond AI, Sleeman JE. 2003. Nuclear substructure and dynamics. Curr Biol 13:R825-8.
53.   Granell S, Baldini G, Mohammad S, Nicolin V, Narducci P, Storrie B, Baldini G. 2008.
Sequestration of mutated alpha1-antitrypsin into inclusion bodies is a cell-protective
mechanism to maintain endoplasmic reticulum function. Mol Biol Cell 19:572-86.
54.   Lawless MW, Greene CM, Mulgrew A, Taggart CC, O'Neill SJ, McElvaney NG. 2004.
Activation of endoplasmic reticulum-specific stress responses associated with the
conformational disease Z alpha 1-antitrypsin deficiency. J Immunol 172:5722-6.
55.   Tirosh B, Iwakoshi NN, Lilley BN, Lee AH, Glimcher LH, Ploegh HL. 2005. Human
cytomegalovirus protein US11 provokes an unfolded protein response that may
facilitate the degradation of class I major histocompatibility complex products. J Virol
79:2768-79.
56.   Wiertz EJ, Jones TR, Sun L, Bogyo M, Geuze HJ, Ploegh HL. 1996. The human
cytomegalovirus US11 gene product dislocates MHC class I heavy chains from the
endoplasmic reticulum to the cytosol. Cell 84:769-79.

57. van den Boomen DJ, Timms RT, Grice GL, Stagg HR, Skodt K, Dougan G, Nathan JA,
Lehner PJ. 2014. TMEM129 is a Derlin-1 associated ERAD E3 ligase essential for virus-
induced degradation of MHC-I. Proc Natl Acad Sci U S A 111:11425-30.
58. Caramelo JJ, Parodi AJ. 2008. Getting in and out from calnexin/calreticulin cycles. J Biol
Chem 283:10221-5.
59. Hammond C, Braakman I, Helenius A. 1994. Role of N-linked oligosaccharide
recognition, glucose trimming, and calnexin in glycoprotein folding and quality control.
Proc Natl Acad Sci U S A 91:913-7.
60. Zhou M, Yu Q, Wechsler A, Ryckman BJ. 2013. Comparative analysis of gO isoforms
reveals that strains of human cytomegalovirus differ in the ratio of gH/gL/gO and
gH/gL/UL128-131 in the virion envelope. J Virol 87:9680-90.
61. Zhang L, Zhou M, Stanton R, Kamil J, Ryckman BJ. 2018. Expression Levels of
Glycoprotein O (gO) Vary between Strains of Human Cytomegalovirus, Influencing the
Assembly of gH/gL Complexes and Virion Infectivity. J Virol 92.
62. Sifers RN, Brashears-Macatee S, Kidd VJ, Muensch H, Woo SL. 1988. A frameshift
mutation results in a truncated alpha 1-antitrypsin that is retained within the rough
endoplasmic reticulum. J Biol Chem 263:7330-5.
63. Choudhury P, Liu Y, Bick RJ, Sifers RN. 1997. Intracellular association between UDP-
glucose:glycoprotein glucosyltransferase and an incompletely folded variant of alpha1-
antitrypsin. J Biol Chem 272:13446-51.
64. Tsao YS, Ivessa NE, Adesnik M, Sabatini DD, Kreibich G. 1992. Carboxy terminally
truncated forms of ribophorin I are degraded in pre-Golgi compartments by a calcium-
dependent process. J Cell Biol 116:57-67.
65. de Virgilio M, Weninger H, Ivessa NE. 1998. Ubiquitination is required for the retro-
translocation of a short-lived luminal endoplasmic reticulum glycoprotein to the cytosol
for degradation by the proteasome. J Biol Chem 273:9734-43.
66. Johnston JA, Ward CL, Kopito RR. 1998. Aggresomes: a cellular response to misfolded
proteins. J Cell Biol 143:1883-98.
67. Johnston JA, Illing ME, Kopito RR. 2002. Cytoplasmic dynein/dynactin mediates the
assembly of aggresomes. Cell Motil Cytoskeleton 53:26-38.
68. Pankiv S, Alemu EA, Brech A, Bruun JA, Lamark T, Overvatn A, Bjorkoy G, Johansen T.
2010. FYCO1 is a Rab7 effector that binds to LC3 and PI3P to mediate microtubule plus
end-directed vesicle transport. J Cell Biol 188:253-69.
69. Hummel T, Krukkert K, Roos J, Davis G, Klambt C. 2000. Drosophila Futsch/22C10 is a
MAP1B-like protein required for dendritic and axonal development. Neuron 26:357-70.
70. Wang H, Bedford FK, Brandon NJ, Moss SJ, Olsen RW. 1999. GABA(A)-receptor-
associated protein links GABA(A) receptors and the cytoskeleton. Nature 397:69-72.
71. Li YR, King OD, Shorter J, Gitler AD. 2013. Stress granules as crucibles of ALS
pathogenesis. J Cell Biol 201:361-72.
72. Zhang K, Daigle JG, Cunningham KM, Coyne AN, Ruan K, Grima JC, Bowen KE, Wadhwa
H, Yang P, Rigo F, Taylor JP, Gitler AD, Rothstein JD, Lloyd TE. 2018. Stress Granule
Assembly Disrupts Nucleocytoplasmic Transport. Cell 173:958-971 e17.
73. Hetz C, Saxena S. 2017. ER stress and the unfolded protein response in
neurodegeneration. Nat Rev Neurol 13:477-491.

74.   Kirchoff V, Wong S, St JS, Pari GS. 2002. Generation of a life-expanded rhesus monkey
fibroblast cell line for the growth of rhesus rhadinovirus (RRV). Arch Virol 147:321-33.

75.   Sinzger C, Hahn G, Digel M, Katona R, Sampaio KL, Messerle M, Hengel H, Koszinowski U,
Brune W, Adler B. 2008. Cloning and sequencing of a highly productive, endotheliotropic
virus strain derived from human cytomegalovirus TB40/E. J Gen Virol 89:359-68.

76.   Smith IL, Taskintuna I, Rahhal FM, Powell HC, Ai E, Mueller AJ, Spector SA, Freeman WR.
1998. Clinical failure of CMV retinitis with intravitreal cidofovir is associated with
antiviral resistance. Arch Ophthalmol 116:178-85.

77.   Murphy E, Yu D, Grimwood J, Schmutz J, Dickson M, Jarvis MA, Hahn G, Nelson JA,
Myers RM, Shenk TE. 2003. Coding potential of laboratory and clinical strains of human
cytomegalovirus. Proc Natl Acad Sci U S A 100:14976-81.

78.   Stanton RJ, Baluchova K, Dargan DJ, Cunningham C, Sheehy O, Seirafian S, McSharry BP,
Neale ML, Davies JA, Tomasec P, Davison AJ, Wilkinson GW. 2010. Reconstruction of the
complete human cytomegalovirus genome in a BAC reveals RL13 to be a potent inhibitor
of replication. J Clin Invest 120:3191-208.

79.   Hobom U, Brune W, Messerle M, Hahn G, Koszinowski UH. 2000. Fast screening
procedures for random transposon libraries of cloned herpesvirus genomes: mutational
analysis of human cytomegalovirus envelope glycoprotein genes. J Virol 74:7720-9.

80.   Chang WL, Barry PA. 2003. Cloning of the full-length rhesus cytomegalovirus genome as
an infectious and self-excisable bacterial artificial chromosome for analysis of viral
pathogenesis. J Virol 77:5073-83.

81.   Tischer BK, Smith GA, Osterrieder N. 2010. En passant mutagenesis: a two step
markerless red recombination system. Methods Mol Biol 634:421-30.

82.   Tischer BK, von Einem J, Kaufer B, Osterrieder N. 2006. Two-step red-mediated
recombination for versatile high-efficiency markerless DNA manipulation in Escherichia
coli. Biotechniques 40:191-7.

83.   Dietz AN, Villinger C, Becker S, Frick M, von Einem J. 2018. A Tyrosine-Based Trafficking
Motif of the Tegument Protein pUL71 Is Crucial for Human Cytomegalovirus Secondary
Envelopment. J Virol 92.

84.   Hohn K, Sailer M, Wang L, Lorenz M, Schneider ME, Walther P. 2011. Preparation of
cryofixed cells for improved 3D ultrastructure with scanning transmission electron
tomography. Histochem Cell Biol 135:1-9.

85.   Villinger C, Schauflinger M, Gregorius H, Kranz C, Hohn K, Nafeey S, Walther P. 2014.
Three-dimensional imaging of adherent cells using FIB/SEM and STEM. Methods Mol
Biol 1117:617-38.

86.   Kremer JR, Mastronarde DN, McIntosh JR. 1996. Computer visualization of three-
dimensional image data using IMOD. J Struct Biol 116:71-6.

87.   Britt WJ, Jarvis MA, Drummond DD, Mach M. 2005. Antigenic domain 1 is required for
oligomerization of human cytomegalovirus glycoprotein B. J Virol 79:4066-79.

1361

1362

FIG 1. UL148 reorganizes ER markers into anomalous structures during HCMV infection.
Fibroblasts infected at 1 MOI with either **(A)** wildtype HCMV strain TB40/E (TB_WT) or **(B)** an isogenic *UL148*-null mutant (TB_148$_{STOP}$), were fixed at 96 h postinfection (hpi) and imaged by confocal microscopy after staining with antibodies specific for calnexin or Hrd1 (magenta) and glycoprotein H (gH, green), as indicated. DAPI (blue) signal was used to counterstain nuclei. **(C)** Percentage of fibroblasts that contain Hrd1 structures at 96 hpi; fifty gH-positive cells for each condition were scored for the presence or absence of UL148 structures (also see SI Fig S1A). (**D**) Three-dimensional confocal image projection of a TB_WT infected fibroblast stained at 96 hpi for Hrd1 and gH. **(E)** Volumetric measurements were made from sixteen infected cells, stained with phalloidin-AlexaFluor 594 conjugate to estimate total cell volume, with Hrd1 antibody to estimate the volume of UL148-dependent ER structures, and with DAPI to estimate nuclear volume.

FIG 2. Characterization of TB_148^{HA} and TB_159^{HA} viruses. **(A)** Human fibroblasts were infected at MOI 1 with the indicated recombinant HCMVs and whole cell lysate samples harvested at the indicated times post-infection (h post-infection: hpi) were analyzed by Western blot to detect HA-tagged UL148 or Rh159, the 72 kD viral nuclear antigen IE1-72 (IE1), and GAPDH. **(B)** Single-cycle viral replication kinetic curves from MOI 1 infected fibroblasts were plotted by determining the titer in tissue culture infectious dose 50 (TCID$_{50}$) from supernatants collected at the indicated times postinfection.

FIG 3. UL148 localizes to unusual ER compartments enriched for glycoprotein quality control markers. Fibroblasts infected at MOI 1 with either TB_148[HA] (panels: **A, C, E, G, I, K, M, O, Q, S**) or TB_159[HA] (panels: **B, D, F, H, J, L, N, P, R, T**) were fixed at 96 hpi, and imaged by confocal microscopy after co-staining with antibodies specific for HA (UL148/Rh159, magenta in merge) and the indicated cellular markers (green in merge). DAPI (blue) counterstaining is shown in merged image.

FIG 4. Quantification of Co-localization between UL148 and Cellular Markers. A Pearson's correlation coefficient (*r*) was calculated using NIH ImageJ software to estimate the degree of co-localization between UL148 (HA signal) and each of the indicated cellular markers. A minimum of 30 cells were analyzed per marker. The arithmetic mean for each co-localization analysis result is shown as a blue line, and data points for individual cells analyzed are plotted as circles.

FIG 5. Cellular proteins involved in glycoprotein quality control are recruited with differing kinetics to UL148 ER structures. Fibroblasts infected with TB_148[HA] at 1 MOI were fixed at the indicated time points (days post infection, dpi) and imaged by confocal microscopy after staining with antibodies specific for HA (UL148, magenta), CNX (green, **A**), Hrd1 (green, **B**) or VCP(green, **C**), DAPI (blue).

FIG 6. TEM of ER structures in wildtype HCMV infected cells. Human fibroblasts infected with wildtype HCMV (TB_WT) were fixed by high-pressure freezing and freeze substitution at day 5 postinfection and imaged using TEM. Panels (**A**) and (**B**) show cell overview at left. For each cell, the boxed region is shown at higher magnification. Scale bars; left panels: 10 µm, right panels: 2 µm. rER: rough ER; Nu: nucleus; cVAC: cytoplasmic viral assembly compartment. Solid arrowheads indicate the UL148-dependent ER structures of interest.

FIG 7. TEM of *UL148*-null HCMV infected cells. Human fibroblasts infected with a *UL148*-null mutant (TB_148$_{STOP}$) were fixed by high-pressure freezing and freeze substitution at day 5 postinfection and imaged using TEM. At the upper left an overview panel of a representative cell is shown with panels **A, B, and C** each boxed. The boxed regions are expanded at higher magnification at right (**A**) and below (**B, C**). Scale bars for upper left overview panel: 10 µm, for zoomed panels: 2 µm. rER: rough ER; Nu: nucleus; cVAC: cytoplasmic viral assembly compartment.

FIG 8. STEM tomography of UL148-dependent ER structures in HCMV infected cells. Human fibroblasts infected with wildtype HCMV (TB_WT) were fixed by high-pressure freezing and freeze substitution at day 5 postinfection and tomograms were recorded by STEM. (**A**) Shows a virtual section through the tomogram of a virus-induced membranous structure; asterisks denote ER structures of distended lumenal space. White arrowheads indicate sites at which membranes originating from distended ER cisternae continue into areas of involuted collapsed ER. Scale bar: 1 μm. (**B**) Shows the same virtual section as in (A) tilted and with a 3D visualization of the membranous network (green) of the entire tomogram. The same distended ER cisternae as in (A) are marked by asterisks. (**C**) Shows a cross section through (B) to visualize the membrane profile of the membranous structures. The region delimited by a white box is shown in a higher magnification on the right. Finer detail of the enlarged ER cisternae and the connections between them are readily visible; asterisks indicate the same distended ER cisternae as in (A) and (B). Also, see SI Movie S1. Scale bar: 1 μm.

FIG 9. Solubility Analysis of UL148 and Rh159. (**A**) Human fibroblasts were infected at MOI 1 TCID$_{50}$/cell with HCMV strain TB40/E derived viruses TB_148^{HA}, which expresses UL148 fused at its C-terminus to an HA-epitope tag, or TB_159^{HA}, which lacks *UL148* and instead expresses rhesus CMV Rh159 carrying a C-terminal HA tag. At the indicated times postinfection (hpi), infected cells were collected radioimmunoprecipitation (RIPA) lysis buffer, centrifuged at 21,000×g for 30 min, after which supernatant (sup) and pellet (pel) fractions were boiled in gel loading buffer containing 2% sodium dodecyl sulfate (SDS). Equivalent portions of supernatant and pellet were resolved by SDS-polyacrylamide gel electrophoresis, transferred to a nitrocellulose membrane for detection of protein species immunoreactive to antibodies against HA epitope (HA) and HCMV glycoprotein B (gB) and for total protein signal using Ponceau S reagent (Ponceau). (**B-C**). Signal intensity of fluorophore-conjugated secondary antibodies in anti-HA Western blots were measured from three independent biological replicates of the experiment shown in (A); error bars indicate standard deviation. (**B**) The fluorescent signal for each infection time point condition (Y-axis indicates arbitrary units, in hundred thousands). (**C**) The amount of signal found in the insoluble (pellet) fraction relative to the total signal (pellet plus supernatant) for each infection time point are plotted as percentage values.

FIG 10. UL148 is sufficient to remodel the ER. HA-tagged UL148 or Rh159 were doxycycline (dox) induced in the context of a "tet-on" lentiviral vector system in stably transduced ARPE-19 epithelial cell populations, i148[HA] and i159[HA], respectively. Cells were fixed at 48 h postinduction for indirect immunofluorescence staining for the indicated cellular markers (green) together with HA (magenta). Scale bar: 10 μm.

FIG 11. ISR activation accompanies redistribution of ER markers during ectopic expression of UL148. (**A**) Lysates of tet-on ARPE-19 cells expressing either UL148 or Rh159 fused to an HA tag, i148HA and i159HA, respectively, were collected at the indicated times post doxycycline (dox) induction and analyzed by Western blot for the expression of the indicated proteins and for the abundance of eIF2α phosphorylated at Ser51 (P-eIF2α) using a phospho-specific antibody. (**B**) Pearson's correlation coefficient (*r*) values were calculated using NIH ImageJ software estimate the degree of co-localization between UL148 (HA signal) and the indicated cellular markers. A minimum of 30 cells were analyzed per marker. Arithmetic means for each co-localization analysis result are shown as blue lines, and data points for individual cells analyzed are plotted as circles.

FIG 12. Live Cell Imaging of UL148-GFP and Rh159-GFP during Induced Ectopic Expression. "Tet-on" ARPE-19 epithelial cells that inducibly express either UL148 or Rh159 fused to green florescent protein (gfp), i148gfp (**A**) and i159gfp (**B**), respectively, were induced for transgene expression using 100 ng/mL doxycycline (dox) and imaged using live-cell microscopy. Images from the selected time points (h post treatment with dox, hpt) are shown. The main scale bar represents 50 μm. For inset panels at upper left of each image, which are magnified 2.4× relative to the main image, the scale bar represents 10 μm. Also see SI Movies S2-S3.

FIG 13. FRAP analysis. ARPE-19 cells that inducibly express Rh159-GFP, i159^{gfp} (**A**) or UL148-GFP, i148^{gfp} (**B-C**) were doxycycline induced for 24 h (100 ng/mL doxycycline (dox). Separate regions of GFP signal in selected cells were photobleached (405 nm laser) or left unbleached, while a third region lacking GFP signal was chosen as a background reference were measured before and during fluorescence recovery after photobleaching (FRAP). Note: background signal was not plotted because values were resolvable from the x-axis. GFP signal intensity is plotted over a time period (seconds, s) starting with an exposure at t=0 (immediately before photobleaching), and including measurements taken every 2 s after photobleaching (0-20 s) until termination of the measurement series at t=322 s (300 s of FRAP). (**D-F**) Images from the selected time points (s, seconds) immediately before and after bleaching, and during fluorescence recovery period. Also see SI Movies S4-S6.

FIG 14: UL148-GFP is stabilized by inhibition of proteasomal but not lysosomal degradation. "Tet-on" ARPE-19 cells that inducibly express UL148-GFP (i148^{gfp}) were induced for 24 h by the addition of 100 ng/mL doxycycline (dox), after which the dox inducing agent was washed out and medium containing either the proteasome inhibitor epoxomicin (20 μM) or the proton pump inhibitor folimycin (115 nM) was added and samples were harvested for Western blot analysis at the indicated times post treatment (h post treatment, hpt) with folimycin or epoxomicin. Also see SI Movie S7.

FIG 15. Inhibition of the integrated stress response impedes UL148-mediated ER remodeling. "Tet-on" ARPE-19 epithelial cells that inducibly express either UL148 or Rh159 fused to green florescent protein (gfp), i148^gfp (**A**) and i159^gfp (**B**), respectively, were induced for transgene expression using 100 ng/mL doxycycline (dox) in the presence of absence of either ISRIB (200 nM) or GSK2606414 (1.1 μM) and monitored by Western blot for expression of UL148-GFP over a series of time points (h post treatment with dox, hpt). (**C-E**): Live-cell imaging of UL148-GFP and Rh159-GFP expression patterns in the presence of ISRIB (200 nM), GSK2606414 (1.1 μM), or DMSO vehicle alone. Main scale bar represents 50 μm. For inset panels at upper left of each image, which are magnified 2.4× relative to the main image, the scale bar represents 10 μm. Also see SI Movies S8-S10.

FIG 16. Inhibition of the ISR prevents the coalescence of Hrd1 and UL148 into discrete structures during infection. (**A**) Representative 3D maximum intensity projections of confocal imaging Z-stacks obtained from cells infected at MOI 1 for 96 h with HCMV strain TB40/E carrying an HA-tagged *UL148*, TB_148[HA], and maintained in the presence of ISRIB (200 nM), GSK2606414 (1.1 µM), or DMSO carrier alone (0.1% vol/vol). In merged images HA signal is shown in magenta, Hrd1 in green, and DAPI counterstaining in blue. (**B**) The percent of Hrd1 antibody signal involved in discrete structures at 96 hpi were calculated for a minimum of 30 cells per condition were determined using Imaris x64 9.3.0 software. Statistical significance was determined using a one-way ANOVA followed by Tukey's post-test; **** represents a P-value of <0.0001. The arithmetic mean for co-localization analysis results are shown as blue lines with data points for individual cells analyzed plotted as circles. (**C-D**) Western blot analyses of fibroblasts infected at MOI 1 with TB_148[HA] and maintained in the presence of ISRIB (200 nM), GSK2606414 (1.1 µM), or DMSO carrier alone (0.1%); hpi: h post infection (hpi). Note: a phospho-specific antibody was used for detection of eIF2α phosphorylated at Ser51 (P-eIF2α).

FIG S1: **Additional indirect immunofluorescent confocal microscopy results from infected cells.**
(**A**) Lower magnification overview of results from FIGS 1A-B; staining of HCMV glycoprotein H (gH) and Hrd1 at 96 h postinfection (hpi) of human fibroblasts with wildtype HCMV strain TB40/E (WT) versus *UL148*-null derivative TB_148$_{STOP}$. (**B**) Staining of gH and Hrd1 at 96 hpi of human fibroblasts with wildtype Merlin recovered from BAC-cloned Merlin pAL1393 (ME_WT) or a *UL148*-null derivative of the same virus (ME_148$_{STOP}$.) (**C**): Staining of gH and Hrd1 at 96 hpi of human fibroblasts with wildtype HCMV strain TRgfp recovered from BAC-cloned TRgfp (TR_WT) or a *UL148*-null derivative of the same virus (TR_148$_{STOP}$). (**D**) Staining of the viral tegument protein UL71 and calnexin at 5 days postinfection of human fibroblasts with primary clinical HCMV isolates obtained from patient throat swabs. (**E**) Staining against HA and Hrd1 in THP-1 macrophages fixed at 96 hpi with HCMV strain TB40/E derivative viruses TB_148^{HA} or TB_159^{HA}; note: THP-1 monocytes were differentiated using phorbol ester treatment. (**F**) Staining of HA and Hrd1 in ARPE-19 epithelial cells infected for 96 h with either TB_148^{HA} or TB_159^{HA}. (**G**) Staining of HA and Hrd1 in ARPE-19 epithelial cells infected for 96 h with HCMV strain AD169 repaired for *UL131* and to which an HA-tagged UL148 CDS from TB40/E was restored to the native *UL148* locus (AD^{r131}_148^{HA}) or with a parental AD169 virus repaired for *UL131* to which *UL148* was not restored (AD^{r131}). Notes: (i) a functional *UL131* is required for efficient infection of epithelial cells; (ii) the difference in scale bar is because with ADr131 we did not employ a viral marker (e.g., HA) to identify infected cells, so the appearance of syncytia and the characteristic kidney-bean shaped nucleus was used to indicate infected cells and a slightly lower magnification was used to best show these features. (**H**) Staining for HA and Hrd1 in telomerase-immortalized rhesus fibroblasts infected with BAC-derived rhesus CMV (RhCMV) strain 68-1 carrying an HA-tag at the C-terminus of Rh159.

FIG S2: Confocal microscopy of LC3B, syntaxin-6 and glycoprotein H staining during HCMV infection. Human fibroblasts were infected with the indicated viruses for 96 h and then fixed, permeabilized, and stained using antibodies specific for the indicated proteins. Indirect immunofluorescence images were captured using a 63X objective on a Leica SP5 confocal microscope. gH: glycoprotein H, STX6: syntaxin-6.

TABLE S1. Oligonucleotide primers and synthetic DNAs used in this study

Name	Sequence (5' to 3')
148_eGFP_Fw	ACCTGGTGTCGCGGCGTCGGATGGTGAGCAAGGGGGAGGAGCTTTTTACTGGCGTTGTTC
Gibs_eGFP_Rv	CGGCCGCCACTGTGCTGGATTACTTATACAGCTCGTCCATACCCAG
Gib_148_Fw	TGTGGTGGAATTCTGCAGATTGCCACCATGTTGCGCTTGCTG
148_noStopRv	CCGACGCCGCGACACCAGGTAGGTTATC
159_eGFP_Fw	TCTAAAACGCAGTCGTGAAGCTCATATGGTGAGCAAGGGGGAGGA
159_noStop_Rv	ATGAGCTTCACGACTGCGTTTTAGAAGTG
Gibs_159_Fw	TGTGGTGGAATTCTGCAGATTGCCACCATGGCCTACAACAGC
Rh159_Fw	TGTGGTGGAATTCTGCAGATACCATGGCCTACAACAGCTTCCTTCTCAGCTGCCTTACCATCGCACTTTTACTG
Rh159_HA_Rv	CGGCCGCCACTGTGCTGGATCTATTAGGCGTAGTCTGGGACGTCGTATGGGTAATGAGCTTCACGACTGCGTTTTAGAAGTG
PpuISceKanGibs_Fw	CATTCACCTGAACCCATAGGTCCTGCCTGGCAGAGAATCTTAGGGATAACAGGGTAATCG
PpuISceKanGibs _Rv	AGATTCTCTGCCAGGCAGGACCTATGGGTTCAGGTGAATGGCCAGTGTTACAACCAATTA
Rh159_Fw_recomb	GGGCGGTGGCCGGCACGCCGCATTTCCTAACCCGCGCAGCATGGCCTACAACAGCTTCCT
Rh159_Rv_recomb	CGCGAACGACGTGTGACGAGGACGTGGTTTCCGCAAGCCTTTAGGCGTAGTCTGGGACG
TB_Rh159HA_seq_Fw	AGTGTTTATGAGTCGGGCGG
REV TB_Rh159HA_seq	GTGGCGACGTGGATTTCTTG
EGFP-P2A-3XHA	GTCGACTGGATCCGGTACCGAATTCGCCACCATGGTGAGCAAGGGGGAGGAGCTTTTTACTGGCGTTGTTCCCATCCTGGTTGAGTTGGATGGGGATGTCAACGGCCACAAGTTCAGCGTTTCTGGAGAAGGAGAAGGCGACGCAACCTATGGGAAGCTCACTTTGAAATTCATATGCACCACCGGTAAACTCCCAGTCCCCTGGCCTACCCTGGTCACTACCTTGACATATGGAGTTCAGTGCTTTTCTCGATACCCCGACCACATGAAGCAGCACGATTTTTTCAAATCTGCAATGCCCGAGGGCTATGTACAAGAGAGGACCATTTTCTTCAAAGACGACGGCAATTACAAGACCAGGGCCGAGGTCAAATTTGAAGGGGACACTCTTGTGAATCGGATCGAACTTAAAGGGATCGATTTCAAAGAGGACGGAAACATACTGGGGCATAAGCTGGAATACAACTACAATAGCCATAATGTTTACATTATGGCAGACAAGCAGAAAAACGGTATTAAGGTCAACTTCAAGATACGGCATAACATTGAAGATGGGTCAGTCCAGCTTGCTGACCATTATCAGCAGAATACACCCATTGGCGATGGTCCCGTGCTTTTGCCTGACAACCACTACCTTTCTACCCAAAGCGCATTGTCAAAAGATCCTAACGAGAAGAGAGATCATATGGTCCTTTTGGAGTTCGTAACCGCAGCTGGTATAACTCTGGGTATGGACGAGCTGTATAAGGGCTCAGGGGCTACAAATTTTAGTCTCCTTAAACAAGCCGGAGATGTCGAAGAGAACCCTGGGCCTTACCCATATGACGTTCCTGACTATGCCGGTTACCCCTATGATGTGCCAGATTATGCCGGATCTTATCCTTACGATGTACCTGACTATGCAGGTGGGAGTGGCATGAATTACGTGGGGCAGC
Gibs_Age_159_Fw	CCGTACCACTTCCTACCCTCGTAAACCGGTGCCACCATGGCCTACAACAG
Gibs_Mlu_GFP_Rv	ACCCGGCGCGGAGGCCAGATCTTAACGCGTTTACTTATACAGCTCGTCCATACC
tRFP linker Fw	CCACTTCCTACCCTCGTAAACCGGTGAATTCGCCACCATGAGCGAGCTGAT
tRFP linker Rv	GCGCGGAGGCCAGATCTTAACGCGTGCGGCCGCTCGAGAATTCATCTGTGCCCCAGTTTGC

TABLE S2. Antibodies Used in Immunofluorescence Microscopy

Primary Antibody	Host species	Isotype	Clone	Vendor	Catalog No.	Dilution used
Calnexin	Mouse	mAb IgG1	37	BD Biosciences	610523	1:50
Calnexin	Rabbit	mAb IgG	C5C9	Cell Signaling Tech.	2679	1:50
Syntaxin 6	Rabbit	mAb IgG	C34B2	Cell Signaling Tech.	2869	1:50
SYVN1 (Hrd1)	Rabbit	mAb IgG	D3O2A	Cell Signaling Tech.	14773	1:500
SEL1L	Rabbit	pAb	n/a	Sigma	S3699	1:50
KDEL	Mouse	IgG2a	10C3	Enzo Life Sci., Inc.	ADI-SPA-827-D	1:50
PDI	Rabbit	mAb IgG	C81H6	Cell Signaling Tech.	3501	1:50
Herp (HerpUD1)	Rabbit	pAb IgG	n/a	Proteintech	10813-1-AP	1:50
VCP	Mouse	IgG2a	5	Invitrogen / Life Technologies	MA3-004	1:50
Calreticulin	Rabbit	mAb IgG	D3E6	Cell Signaling Tech.	12238	1:200
Ribophorin 1 (RPN1)	Rabbit	pAb IgG		ThermoFisher / Invitrogen	PA5-65662	1:50
EDEM-1	Mouse	mAb IgM	D-1	Santa Cruz	sc-377394	1;50
GABARAP	Rabbit	mAb	E1J4E	Cell Signaling Tech.	13733	1:200
LC3B	Rabbit	mAb	D11	Cell Signaling Tech.	3868	1:200
HA epitope	Chicken	IgY		Bethyl	A190-106A	1:200
HCMV glycoprotein H (gH)	Mouse	IgG2b	14-4b	Gift of William J. Britt, M.D.	n/a	1: 50
Reticulon-3 (RTN3)	Mouse	IgG2a		ThermoFisher / Invitrogen	MA5-15538	1: 200
HA epitope	Rabbit	pAb		Bethyl	A190-108A	1:1000
Secondary Antibody	**Host species**	**Isotype**		**Vendor**	**Catalog No.**	**Dilution used**
Alexa Fluor 488 anti-Mouse IgG (H+L), cross adsorbed	Goat	IgG		ThermoFisher / Invitrogen	A11001	1:1000
Alexa Fluor 488 anti-Rabbit IgG (H+L) Cross-Adsorbed	Goat	IgG		ThermoFisher / Invitrogen	A11008	1:1000
Alexa Fluor 594 anti-Rabbit IgG (H+L) Cross-Adsorbed	Goat	IgG		ThermoFisher / Invitrogen	A11012	1:1000
Alexa Fluor 594 anti-Mouse IgG (H+L) Cross-Adsorbed	Goat	IgG		ThermoFisher / Invitrogen	A11005	1:1000
Alexa Fluor 647 anti-Chicken IgY (H+L)	Goat	IgG		ThermoFisher / Invitrogen	A11039	1:1000