A cytomegalovirus immunevasin triggers integrated stress response-dependent

reorganization of the endoplasmic reticulum.

Running Title: HCMV UL148 reorganizes the ER.

Hongbo Zhang[a,1], Clarissa Read[d,e,1], Christopher C. Nguyen[a,1], Mohammed N.A.

Siddiquey[a], Chaowei Shang[c], Cameron M. Hall[a], Jens von Einem[d], and Jeremy P.

Kamil[a,b #]

[a]Department of Microbiology and Immunology,

[b]Center for Molecular and Tumor Virology

[c]Research Core Facility

LSU Health Sciences Center, Shreveport, Louisiana, USA

[d] Institute of Virology, Ulm University Medical Center,

[e]Central Facility for Electron Microscopy, Ulm University

Ulm, Germany

[1] these authors contributed equally to the study

Running Head: HCMV UL148 Remodels the ER

#Address correspondence to Jeremy P. Kamil, jkamil@lsuhsc.edu

ABSTRACT. Human cytomegalovirus (HCMV) encodes an ER-resident glycoprotein,

UL148, which activates the unfolded protein response (UPR) but is fully dispensable for

viral replication in cultured cells. Hence, its previously ascribed roles in immune

evasion and modulation of viral cell tropism are hypothesized to cause ER stress. Here,

we show that UL148 is necessary and sufficient to drive the formation of large ER-

derived structures that occupy up to 7% of the infected cell cytosol. The structures are

found to be sites where UL148 coalesces with cellular proteins involved in ER quality

control, such as Hrd1 and EDEM1. Transmission electron microscopy analyses reveal

the structures to be comprised of tortuous, densely packed and apparently collapsed

ER membranes that connect to distended cisternae. During induced ectopic expression

of UL148-GFP fusion protein, punctate signals traffic to accumulate at prominent

structures that exhibit poor recovery of fluorescence after photobleaching. Small

molecule blockade of the integrated stress response (ISR) prevents the formation of

puncta, leading to a uniform reticular fluorescent signal. Accordingly, ISR inhibition

during HCMV infection abolishes the coalescence of UL148 and Hrd1 into discrete

structures, which argues that UL148 requires the ISR to cause ER reorganization.

Given that UL148 stabilizes immature forms of a receptor binding subunit for a viral

envelope glycoprotein complex important for HCMV infectivity, our results imply that

stress-dependent ER remodeling contributes to viral cell tropism.

IMPORTANCE.

Perturbations to ER morphology occur during infection with various intracellular

pathogens and in certain genetic disorders. We identify that an HCMV gene product,

UL148, profoundly reorganizes the ER during infection, and is sufficient to do so when

expressed on its own. Our results reveal that UL148-dependent reorganization of the

ER is a prominent feature of HCMV infected cells. Moreover, we find that this example

of virally induced organelle remodeling requires the integrated stress response (ISR), a

stress adaptation pathway that contributes to a number of disease states. Since ER

reorganization accompanies the roles of UL148 in HCMV cell tropism and intracellular

retention of the immune cell co-stimulatory ligand CD58, our results may have

implications for understanding the mechanisms involved. Furthermore, our findings

provide a basis to utilize UL148 as a tool to investigate organelle responses to stress

and to identify novel drugs targeting the ISR.

INTRODUCTION.

UL148 is a human cytomegalovirus (HCMV) ER-resident glycoprotein that plays roles in evasion of cell-mediated immunity and shows intriguing effects on cell tropism. During infection of epithelial cells, viruses disrupted for *UL148* replicate to produce roughly 100-fold enhanced levels of infectious progeny virions compared to wildtype (1). These effects correlate with reduced expression of glycoprotein O (gO), a subunit of a heterotrimeric viral glycoprotein H (gH) / glycoprotein L (gL) complex (gH/gL/gO) that is required for the infectivity of cell-free virions (2-4). The presence of gO in the context of the heterotrimer endows the virus with the capacity to utilize the platelet derived growth factor receptor α (PDGFRα) as an entry receptor (5-7). Accordingly, UL148 has been found to stabilize immature forms of gO prior their assembly into gH/gL/gO heterotrimers (1, 8). Despite that UL148 does not stably associate with gO, the data suggest an interaction with gH (1).

UL148 also physically associates with CD58 (LFA-3), a co-stimulatory ligand for natural killer cells and T-lymphocytes, preventing its presentation at cell surface (9). Although the mechanisms by which UL148 stabilizes gO and retains CD58 within the ER remain unknown, UL148 strongly contributes to activation of the unfolded protein response (UPR) during infection, and is sufficient to activate the UPR when ectopically expressed in non-infected cells (10). UL148 co-purifies from infected cells with SEL1L, an adaptor subunit of ER-based E3 ubiquitin ligase Hrd1 that plays important roles in ER-associated degradation (ERAD) of terminally misfolded glycoproteins (8). This suggests a physical interaction with ERAD machinery, which may be germane to the mechanism by which UL148 activates the UPR.

Here, we show that the expression of UL148 is necessary and sufficient to

induce unusual ER structures at which large quantities of ER factors involved in

glycoprotein quality control accumulate. In electron microscopy analyses, we find that

the UL148 induced ER-structures are comprised of densely packed, tortuous ER

membranes that form connections with tubules of highly distended cisternal space.

Furthermore, additional data from inhibitor studies strongly argue that ER remodeling

triggered by UL148 requires the integrated stress response. Overall, our results reveal

a striking ER perturbation induced by HCMV, and which is entirely controlled by a single

viral gene product. These findings may have important implications for understanding

how UL148 regulates viral cell tropism and/or contributes to viral evasion of cell-

mediated immunity.

RESULTS.

UL148 causes reorganization of ER quality control proteins into unusual globular structures.

The HCMV ER-resident glycoprotein UL148 was previously observed to co-localize with the ER marker calnexin during infection (1). Nonetheless, calnexin staining did not show the uniform reticular pattern characteristic for the ER marker. We later noticed that cells infected with a *UL148*-null virus showed uniform calnexin staining (see below). To formally determine whether UL148 influences calnexin localization, we compared fibroblasts at four days post-infection with either wildtype strain TB40/E virus (TB_WT) or a *UL148*-null mutant (TB_148$_{STOP}$)(8, 10), each derived from an infectious bacterial artificial chromosome (BAC) clone of HCMV strain TB40/E (11) (**FIG 1**). In cells infected with wildtype virus, calnexin antibodies stained unusual globular structures at the cell periphery, as expected (1) (**FIG 1A, 1C, SI FIG S1A**). However, in cells infected with the *UL148*-null virus, calnexin staining was uniform throughout the cytosol (**FIG 1B-C**), as would be expected for an ER marker in uninfected cells. The staining pattern for Hrd1, another ER marker, likewise indicated accumulation at unusual globular structures during wildtype HCMV infection, but not during infection with *UL148*-null mutant viruses (**FIG 1, SI FIG S1A**). Because calnexin and Hrd1 staining showed uniform distribution in *UL148*-null virus infected cells, and because *UL148* is fully dispensable for efficient viral replication in fibroblasts (1), these results suggest that redistribution of these two ER markers depends on UL148.

As expected (12, 13), antibodies specific for glycoprotein H (gH), a viral envelope

glycoprotein, stained a juxtanuclear compartment, termed the cytoplasmic virion

assembly compartment (cVAC), which does not involve the ER (**FIG 1**). Similarly

contrasting staining patterns for Hrd1 and/or calnexin were observed in cells infected

with wildtype versus *UL148*-null mutants of clinical HCMV strains Merlin and TR (**FIG**

S1). Reciprocally, we restored a functional *UL148* at its native locus in the context of a

BAC clone of HCMV strain AD169, which spontaneously lost most of the gene in the

course of extensive genetic rearrangements that occurred during serial passage in

cultured cells. In cells infected with AD169 repaired for UL148, but not the *UL148*-null

parental virus, the Hrd1 staining pattern showed prominent globular structures (**FIG S1**).

Because similar differences in Hrd1 staining are observed between wildtype and

UL148-null infections of THP-1 macrophages, ARPE-19 epithelial cells and fibroblasts

(**FIG 1, FIG S1**), we concluded that UL148-dependent reorganization of ER markers

occurs in multiple cell types. The calnexin staining pattern seen with four different

primary clinical isolates from patient throat swabs likewise indicated punctate globular

structures. These observations, taken together with results from formal comparisons of

wildtype versus *UL148*-null mutants of four different BAC-cloned HCMV strains (**FIG 1,**

FIG S1), argue that UL148 profoundly affects the ER during natural infection.

We measured the three-dimensional volume of UL148-dependent Hrd1

structures from sixteen cells fixed at 96 h post infection (hpi) with strain TB40/E. On

average, the structures occupied 303.3 µm^3 (SEM: +/- 24.8) out of a total cell volume of

7657 µm^3 (SEM:+/- 651.0), which was estimated using a phalloidin-fluorophore

conjugate to detect the actin cytoskeleton (**FIG 1D-E**). Subtracting the volume of nuclei,

as indicated by 4′,6-diamidino-2-phenylindole [DAPI] (average nucleus: 1360 μm^3,

SEM:+/-129.4), we calculate that, on average, the structures occupy 5.2% of the

cytosolic volume (SEM: +/-0.61%, range: 2.0% – 7.0%). Based on these findings, taken

together with results comparing additional wildtype (WT) versus *UL148*-null mutant

HCMV strains (**FIG S1**), we conclude that the UL148-dependent ER structures are a

prominent feature of HCMV-infected cells.

UL148 substantially contributes to activation of the unfolded protein response

(UPR) during HCMV infection (10), and co-purifies from infected cells with SEL1L (8),

an adaptor subunit for the E3 ubiquitin ligase Hrd1, which plays crucial roles in ER-

associated degradation (ERAD) of terminally misfolded glycoprotein substrates (14).

Hence, the accumulation of Hrd1 and calnexin at unusual structures during wildtype but

not *UL148*-null infection may suggest that structures form in response to defects in ER

quality control (ERQC) caused by UL148. Under conditions of proteasome inhibition,

overexpression of certain misfolded glycoproteins causes cellular factors involved in

ERQC, such as calnexin, to compartmentalize from the rest of the ER, while other ER

markers such as BiP (Grp78) or protein disulfide isomerase (PDI) remain largely

unaltered (15-17).

To gain further insights into the nature of these peculiar ER structures, we set out

to develop a more comprehensive understanding of their protein composition by

comparing the localization of selected ER markers at 96 hpi during infection with either

TB_148^{HA} (1), a recombinant HCMV strain TB40/E virus that expresses UL148 fused at

its C-terminus to the nonapeptide epitope tag YPYDVPDYA from influenza A

hemagglutinin (HA), or TB_159^{HA}, a *UL148*-null comparator in which the UL148

homolog from rhesus cytomegalovirus, *Rh159*, likewise fused to a C-terminal HA tag, is

expressed in lieu of UL148.

We considered the TB_159^{HA} virus to be an appropriate control for the following

reasons. Firstly, Rh159 and UL148 exhibit ~30% identity at the amino acid level and

both glycoproteins localize to the ER and block cell surface presentation immune cell

activating ligands; UL148 retains CD58, a ligand for CD2, while Rh159 retains NKG2D

ligands of the MIC- and ULBP families (1, 9, 18). Secondly, Rh159 is expressed from

TB_159^{HA} at comparable levels and with similar kinetics observed for UL148 from

TB_148^{HA}, and the two viruses replicate indistinguishably in fibroblasts (**FIG 2**).

Moreover, as was the case in ectopic expression (10), Rh159 does not appear to

activate the UPR to the same extent that UL148 does, as evidenced by lower levels of

phospho-eIF2α and ATF4 during TB_159^{HA} infection (**FIG 2A**).

ERQC markers failed to coalesce into unusual structures during infection with

TB_159^{HA} (**FIG 3**). The structures likewise failed to occur in rhesus fibroblasts during

infection with a recombinant rhesus cytomegalovirus (RhCMV) that we engineered to

express an HA-tag fused at the C-terminus of Rh159 (**FIG S1**). The latter argues

against the possibility that Rh159 somehow requires the context of rhesus cells to

redistribute ER markers in a manner analogous to what is seen for UL148 during HCMV

infection. Taken together, the data suggest that UL148 and Rh159 authentically differ in

their effects on the secretory pathway.

In cells infected with TB_148^{HA}, HA antibody staining indicated localization of

UL148 to globular structures, as expected (1). Antibody signals from indirect confocal

immunofluorescence detection of cellular ER resident proteins involved in ERQC,

including calnexin, Hrd1, SEL1L, Herp, valosin containing protein (VCP, p97), and

EDEM1, strongly co-localized with signals from HA-tagged UL148 (**FIG 3-4**). These

results indicate that a number of ERQC factors coalesce with UL148 to form prominent

globular structures in infected cells. In contrast, we observed uniform ER staining

patterns for PDI and calreticulin (CALR), indicating that these ER markers do not

localize to the UL148 structures (**FIGS 3Q, 3S**). Intriguingly, antibody signals detecting

reticulon 3 and ribophorin 1, which are markers for smooth ER and rough ER,

respectively, each appreciably co-localized with UL148 (HA) signal at the induced

structures (**FIGS 3K, 3M)**, which indicates that the structures may involve both rough

and smooth ER. LC3B failed to co-localize with the structures (**FIG S2**), as might be

expected given that the virus inhibits macroautophagy at late times during infection (19,

20). Nevertheless, our results suggested that the induced structures are enriched for a

related mammalian ATG8 ortholog, GABARAP (**FIG 3O**).

In cells infected with TB_159^{HA}, all of the ER markers we examined showed

uniform, reticular staining, as did the anti-HA signal detecting Rh159 (**FIG 3B, D, F, H,**

J, L, N, P, R, T). EDEM1, in addition to showing reticular staining, also labeled puncta

associated with the nucleus (**FIG 3J, 3I**), which may represent enriched levels of the

protein at the rough ER membranes associated with the nuclear envelope, although we

cannot exclude the possibility of spurious intranuclear staining. GABARAP antibodies

stained the cVAC in TB_159^{HA} infected cells, but also showed a much weaker diffuse

signal throughout the cytosol. Even though we took steps to block viral Fc receptors,

which can cause rabbit antibodies to non-specifically label the cVAC (21), it is plausible

that GABARAP antibody signal from the cVAC reflects incomplete blocking of viral Fc

receptors. For both viruses, antibodies specific for the TGN marker syntaxin-6 (STX6)

and gH, as expected, stained the juxtanuclear cVAC structure (12, 13, 22), a post-ER

compartment that excludes UL148 (1) (**FIG S2**).

To quantify the degree of co-localization with UL148, we calculated Pearson's

correlation coefficients from a minimum of thirty TB_148[HA] infected cells per staining

condition (fixation at 4 dpi), comparing the signal overlap for each ER marker to the HA

signal from UL148. The correlation coefficients (r) for staining patterns from antibodies

specific for EDEM1, Hrd1, reticulon-3 (RTN3), SEL1L, the ER retention motif KDEL, and

calnexin ranged from 0.8 to 0.72. These results suggest that these proteins, and

lumenal ER resident proteins such as BiP (GRP78) and GRP94 that contain a C-

terminal KDEL motif (EKDEL), extensively co-localize with UL148 at the induced

structures (**FIG 4**). Meanwhile, GABARAP, VCP, ribophorin-1 (RPN1), and Herp

showed r values in the range of 0.54 to 0.51, indicating a moderate degree of co-

localization with UL148. However, CALR and PDI, which like BiP, are lumenal ER

residents, and the TGN marker syntaxin-6, gave r values of close to zero, indicating that

these markers do not appreciably co-localize with UL148, as is consistent with our

immunofluorescence data suggesting that these markers negligibly associate with the

unusual structures. From these results, we conclude that the UL148-dependent ER

structures are enriched for cellular markers involved in glycoprotein quality control. In

this regard, the structures resemble the "ER quality control (ERQC) compartments"

described by G. Lederkremer and colleagues (15, 16, 23). We further conclude that

Rh159 cannot substitute for UL148 to redistribute ERQC markers.

233 **VCP and Hrd1 are recruited to incipient UL148 ER structures prior to calnexin.**

234 To determine whether there might be differences in the kinetics of recruitment of

235 ERQC markers during formation of the ER structures, we examined a series of time

236 points from one to four days post-infection (dpi) with wildtype HCMV strain TB40/E

237 carrying HA-tagged UL148, staining for three different ERQC markers, calnexin, Hrd1,

238 and VCP, alongside UL148 (HA). At 1 dpi the signal from UL148 was only faintly

239 detected, as expected (1), while each of the ER markers was readily detected, providing

240 a readout of their staining patterns prior to being substantially perturbed by UL148 (**FIG**

241 **S3**). At 2 dpi, we detected robust anti-HA signal, indicating the presence of UL148. At

242 this time point, UL148 exhibited intense signal at small globular puncta, which we

243 interpret to represent incipient UL148 structures, as well as more diffuse staining of a

244 reticular structure consistent with undisturbed ER. Calnexin did not appreciably co-

245 localize with the UL148 puncta until at least 3 dpi, and the structures were not readily

246 visualized by calnexin staining until 4 dpi (**FIG S3**). In contrast, signals from Hrd1 and

247 VCP staining were sufficient to mark the UL148 puncta by 2 dpi, which suggests that

248 these markers may co-localize with UL148 at earlier points during the genesis of the ER

249 structures. Notably, the appearance of Hrd1 at the structures prior to calnexin is

250 consistent with our previous results showing that UL148 co-purifies from infected cells

251 with SEL1L, an adaptor subunit for Hrd1 (8). These results imply that basal elements of

252 the ERAD machinery, exemplified by Hrd1 and VCP, might be recruited to UL148-

253 induced ER structures prior to calnexin.

254

255 **Visualization of UL148-induced ER structures by electron microscopy.**

To discern the ultrastructural appearance of the UL148 structures and to formally

ascertain their relationship to the ER, we carried out transmission electron microscopy

(TEM) imaging of wildtype (TB_WT) and *UL148*-null (TB_148$_{STOP}$) infected fibroblasts at

5 dpi. In high pressure frozen and freeze-substituted infected fibroblasts, cells with a

high density of viral nucleocapsids within the nucleus were selected for analysis, as this

feature indicates late time points during infection when UL148 is abundantly expressed.

The TEM results revealed prominent globular and oblong structures in the cytoplasm of

wildtype virus infected fibroblasts, but not *UL148*-null infected controls (**FIG 5-6**). The

structures stand out for their high electron-density, which may reflect the abundance of

ERQC proteins together with UL148 in these structures. Under higher magnification,

these areas are characterized by accumulations of densely packed ruffled membranes

and membranous material, which appear to either be collapsed ER, or ER tubules in

very close association with each other. The membrane accumulations were associated

with smooth and partially rough ER structures of seemingly enlarged cisternal space.

Further, tomographic reconstruction of scanning transmission electron microscopy

(STEM) data suggest that densely packed ER cisternae within the structures are

interconnected and continuous in three-dimensional space (**FIG 7B-C, SI Movie S1**).

UL148 accumulates in a detergent-insoluble form during infection.

Disease-associated variants of certain cellular proteins, such as the A103E

mutant of the calcium channel Orai1, and the E342K "Z" variant of alpha-1 antitrypsin (Z

A1AT), localize to anomalous ER structures reminiscent of those we observe to depend

on UL148 (17, 24-27). A103E Orai1 and Z A1AT are found to accumulate in detergent-

insoluble forms within the ER or within ER membranes, respectively, which indicates

aggregation or polymerization of these proteins, and suggests a mechanism for the

formation of ER structures by the proteins (17, 24, 27, 28). We therefore set out to

determine whether differences in solubility might correlate with the differing potentials

UL148 and Rh159 to cause ER reorganization and to activate the UPR (10). To

address this question, we infected fibroblasts with TB_148[HA] or TB_159[HA] at MOI 1 and

at various times post-infection prepared cell lysates in radioimmunoprecipitation assay

(RIPA) buffer. After subjecting the lysates to high-speed centrifugation, we examined

the relative levels of UL148 and Rh159 in the detergent-soluble supernatant and

detergent-insoluble pellet fractions.

A substantial portion of UL148 was detected from the detergent-insoluble

fractions at all time points tested (**FIG 8**). Furthermore, the percentage of UL148

detected within the insoluble fraction increased over time. In contrast, Rh159 was found

only in the detergent-soluble fraction. From these results, we conclude that UL148

accumulates in a detergent-insoluble form during infection, and that Rh159 does not do

so. Because the anti-HA immunoreactive band detected in both the soluble and

insoluble fractions from TB_148[HA] infected cells showed a relative mobility of ~35 kD,

which matched that expected for the mature, endoH-sensitive glycoprotein (1), these

findings argue that UL148 may form aggregates or polymers within the ER.

UL148 is sufficient to compartmentalize the ER.

Our data thus far demonstrate that UL148 is necessary during infection to cause

redistribution of cellular ER markers for glycoprotein quality control processes, such that

substantial portion of the ERQC machinery appears to become sequestered away from

the rest of the organelle into novel membranous structures. Because a number of viral

proteins that remodel the ER during infection are sufficient to alter ER morphology when

ectopically expressed [e.g. (29),(30), reviewed in (31)], we wondered whether UL148

expression would be sufficient to induce ER structures akin to those seen during HCMV

infection. Making use of "tet-on" ARPE-19 epithelial cells that inducibly express either

UL148 or Rh159, each carrying a C-terminal HA-tag (32), we induced transgene

expression for 48 h and subsequently stained for various cellular markers for the ER, for

ERQC as well as for the ATG8 family proteins LC3B and GABARAP, which play roles in

macroautophagy and related processes.

In cells expressing UL148 (i148^{HA}), we observed that calnexin, Hrd1, EDEM1,

and VCP co-localized with UL148 at prominent globular structures reminiscent of those

observed during infection (**SI FIG 4**). The respective rough and smooth ER markers

ribophorin 1 (RPN1) and reticulon 3 (RTN3), as well as the ATG8 proteins LC3B and

GABARAP likewise co-localized to the UL148-induced structures. However, PDI and

CALR each failed to substantially co-localize with UL148 at the structures (**SI FIG 4**), or

at best showed only limited co-localization, consistent with our results from infected

cells (**FIG 3-4**). Signals from antibodies specific for the KDEL (EKDEL) motif important

for ER retrieval of luminal ER residents, such as BiP and GRP94, likewise showed only

limited co-localization with UL148. The PDI and KDEL results suggest that large

regions of the ER are not involved in the UL148 structures.

In cells expressing the UL148 homolog Rh159 (i159^{HA}), we detected uniform

cytoplasmic distribution of ER markers, as well as of HA-tagged Rh159, similar to what

we observed during infection with the recombinant HCMV TB_159^HA (**FIG 3**).

Furthermore, in this setting the staining patterns for the ATG8 family proteins LC3B and

GABARAP failed to indicate any notable structures. Consistent with our previous study

showing that UL148, but not Rh159, is sufficient to activate the UPR (10), we observed

accumulation of ATF4 and phosphorylated eIF2α following doxycycline (dox) induced

expression of UL148 but not Rh159 (**FIG 9A**). Importantly, the intensity of HA signals

detecting UL148 and Rh159 indicated that both ER resident immunevasins accumulate

at roughly comparable levels following dox induction. This argues against the possibility

that gross protein expression differences accounts for the differing effects of UL148 and

Rh159 on UPR activation and on the staining patterns for ER markers.

Pearson's correlation coefficient values were calculated to quantify the extent of

colocalization between UL148 and various cellular markers: EDEM1, Hrd1, VCP,

calnexin (CNX), RTN3, SEL1L, LC3B, and PDI (**FIG 9B**). The results quantitatively

buttress the confocal immunofluorescence results from these cells (**FIG S4**). We

therefore conclude that no other HCMV gene products are required for UL148 to cause

ERQC factors to segregate into discrete compartments. Hence, we conclude from

these findings that UL148 is sufficient to cause large-scale reorganization of ERQC

markers. Moreover, these results suggest that reorganization of ER markers into

discrete structures may be related to the propensity of UL148 to induce ER stress.

To determine whether UL148-dependent ER remodeling could be visualized in

real-time using live cell imaging, we constructed new "tet-on" lentiviral vectors that

inducibly express either UL148 or Rh159 fused at their predicted C-terminal cytoplasmic

tails to the enhanced green fluorescent protein (GFP) from *Aequorea victoria* (33)*.*

Following lentiviral transduction, we isolated puromycin-resistant ARPE-19 and

subsequently subjected to fluorescence-activated cell sorting (FACS) to enrich for cells

that expressed GFP signal following dox treatment. The resulting cell populations,

which inducibly express either UL148-GFP or Rh159-GFP, were designated i148^{GFP}

and i159^{GFP}, respectively.

In live cell imaging studies, we observed that the GFP signal in dox-induced

i148^{GFP} cells first appeared in a reticular, largely uniform pattern, which was readily

visible at 5 h post-induction. However, by 9 h post-induction punctate signals began to

appear. These puncta were observed to traffic to sites of large-scale accumulation,

where large fluorescent structures progressively increased in size up until the

termination of the experiment at 19 h (**FIG S5, SI Movie 2**). In contrast, i159^{GFP} cells

exhibited a largely diffuse, reticular GFP signal at all time points monitored subsequent

to transgene induction (**FIG S5, SI Movie 3**). The live cell imaging results thus

recapitulated the differences in the HA-staining patterns observed between dox-induced

i148^{HA} and i159^{HA} cells following fixation, as well as those between HCMV expressing

HA-tagged UL148 versus *UL148*-null comparator viruses (**FIGS 1, 3, SI Fig S1**).

UL148-GFP structures exhibit poor recovery of fluorescence after

photobleaching.

To determine whether the protein contents of the UL148 induced structures

exhibits decreased mobility compared to non-perturbed ER regions, we carried out

fluorescence recovery after photobleaching (FRAP) studies. We photobleached regions

of GFP signal in i148^{GFP} or i159^{GFP} ARPE-19 cells that had been induced for transgene

expression for 24 h, and then monitored recovery of fluorescence over a 5 min time

period. In Rh159-GFP expressing cells (i159^{GFP}), fluorescence nearly recovered to pre-

bleach levels within 3 min, and by 5 min fully recovered (**SI FIG S6**). In contrast, when

we photobleached a prominent UL148-GFP structure, the GFP signal failed to

appreciably recover fluorescence during the same 5 min time period. Nonetheless,

regions of reticular GFP signal from an i148^{GFP} cell, which presumably represent ER

regions not involved in an anomalous structure, recovered fluorescent signal with

kinetics similar to those observed during Rh159-GFP expression. These results

indicate that bleached UL148-GFP within the structures cannot rapidly be exchanged

with UL148-GFP from other portions of the organelle, which suggests that the induced

UL148 structures do not exchange their contents as efficiently as unperturbed ER.

Ectopically-expressed UL148 is degraded by proteasomes and not by autophagy.

The UL148 ER structures were found to be enriched with proteins such as Hrd1,

SEL1L, EDEM, and VCP (**FIG 3-4**, **9B, SI FIG S4**), which are posited to play key roles

in ERAD, a process during which misfolded glycoproteins are recognized, processed,

and dislocated across the ER membrane for degradation at cytosolic proteasomes.

However, we also detected elements of the machinery for autophagy in close

association with the UL148 structures. Namely, the mammalian ATG8 ortholog

GABARAP co-localized with UL148 during infection (**FIG 3-4**), and both GABARAP and

another ATG8 ortholog, LC3B, co-localized with UL148 at the ER structures during

ectopic expression of UL148 (**FIG 9B, SI FIG S4**). Therefore, we wished to determine

whether UL148 is degraded by the proteasome, as would be consistent with

conventional ERAD, or by the lysosome, which would suggest a role for autophagy-

related processes, such as selective autophagy of the ER (34), in dispensing with

UL148, and presumably, in resolving the ER perturbations.

As a first step, we conducted a live-cell imaging "washout" experiment in which

i148^{GFP} cells were induced for transgene expression for 24 h, after which the growth

medium containing the transgene inducing agent (dox) was replaced with medium

lacking dox. During a 22 h imaging period following dox-washout, we observed the

structures to become progressively smaller as the reticular and punctate GFP signals

gradually abated (**SI FIG S7, SI Movie S7**). Because the results suggested that UL148-

GFP structures could be resolved over time, we induced UL148-GFP in tet-on ARPE-19

cells (i148^{gfp}) for 24 h, washed out the inducing agent (dox), and then applied either

epoxomicin (20 μM), an irreversible inhibitor of the proteasome (35), or folimycin (115

nM), a proton pump inhibitor that blocks autophagosome maturation and impedes

lysosome-dependent degradation (36, 37). Of the two inhibitors, only epoxomicin

stabilized UL148 (**SI FIG S7**). Although folimycin treatment had no obvious effect on

UL148-GFP levels, the drug markedly increased the levels of LC3B-II, as would be

expected with the lysosomal proton pump inhibitor. From these results, we conclude

that ectopically expressed UL148 is primarily degraded by proteasome-dependent

pathway.

UL148 requires the integrated stress response to cause ER reorganization.

We previously reported that UL148 triggers the UPR during ectopic expression,

and that UL148 contributes to activation of the PKR-like ER kinase (PERK) and inositol

requiring enzyme 1 (IRE1) during infection (10). The literature suggests that formation

of ERQC compartments requires PERK (38). PERK responds to ER stress by

activating the integrated stress response (ISR). In particular, PERK phosphorylates

eIF2α at Ser51, and the accumulation of phosphorylated eIF2α (eIF2α-P) globally

attenuates mRNA translation whilst stimulating the translation of a subset of mRNAs,

such as those encoding ATF4 and CHOP, which play roles in cellular adaptation to

stress (39). Stress-regulated translation of such mRNAs involves small upstream open

reading frames (uORFs) in their 5' untranslated regions that ordinarily suppress

translation under basal conditions (40, 41).

To examine whether ER remodeling in response to UL148 requires the ISR, we

turned to a well characterized small molecule inhibitor of stress-regulated translation,

ISRIB (42-46). ISRIB is thought to act as a "molecular staple" that holds the guanine

nucleotide exchange factor eIF2B in an active decameric configuration (44-47), such

that eIF2B will continue to generate ternary complex (eIF2•GTP•Met-tRNAi) necessary

for new cycles of translational initiation, despite the presence of eIF2α-P. Because

PERK is the kinase that phosphorylates eIF2α in response to ER stress (48, 49), we

also tested for effects of the PERK inhibitor GSK2606414 (50). Having confirmed that

ISRIB and GSK260641 do not negatively impact UL148 expression during dox-induction

of UL148-GFP **(FIG 10)**, we treated i148^{GFP} cells with 200 nM ISRIB, 1.1 µM

GSK260641, or DMSO vehicle control, and carried out live cell imaging during induction

of UL148-GFP expression.

ISRIB and GSK6060414 virtually abolished the formation of UL148 puncta

through 13 h post induction (dox addition), a time point at which the DMSO control

condition showed abundant large structures (**FIG 10C-E, SI Movies 8-10**). Based on

our findings (**FIG 1, 3, 5, 7, 9B, S4)**, we interpret the formation and subsequent large-

scale aggregation of UL148-GFP puncta to faithfully indicate ER reorganization in

response to UL148. Therefore, these results suggested to us that pharmacological

inhibition of either the ISR or PERK prevent UL148 from remodeling the ER, which

implies that ER remodeling in response to ER stress requires stress-regulated

translation.

Since our results with ISRIB and GSK6060414 indicated that the ISR is required

for ER remodeling to occur during ectopic expression of UL148, we next asked whether

these pharmacological agents would prevent UL148-dependent ER remodeling in the

physiologically authentic context of HCMV infection. Our live cell imaging studies from

ISRIB and GSK6060414 treatments indicated that UL148-GFP puncta could begin to

appear by ~19 h post treatment (**FIG 10, SI Movies 9-10**), which suggested that the

pharmacological activity of the drugs might diminish over time. Thus to ensure that

blockade of PERK and the ISR would be sustained over the 4 d period that it takes for

structures to fully form during HCMV infection, we replenished drug treatments every 24

h, replacing the spent media with media containing freshly reconstituted ISRIB,

GSK6060414, or DMSO vehicle control.

At 4 dpi, ER structures were readily visible in DMSO control treatment condition

but virtually abolished in the ISRIB and GSK2606414 conditions (**FIG 11A**). To quantify

these effects, we obtained Z-stacks from a minimum of 30 cells per treatment condition

and used Imaris 3D image analysis software to calculate the percentage of Hrd1 signal

that coalesced into discrete structures during infection. The results indicated highly

significant differences between the DMSO carrier alone setting compared to treatments

with either GSK2606414 or ISRIB (**FIG 11B**). In the DMSO control condition, on

average, 10% of the Hrd1 signal was found to be associated with the UL148 structures

(arithmetic mean; range 5.8% – 17.6%), as indicated by structures delimited by HA

signal detecting UL148. In the presence of ISRIB or GSK2606414, however, these

values were reduced to 2.8% (range: 0.69% – 7.1%) and 3.6% (range: 0.8%-6.5%),

respectively. Reassuringly, roughly equivalent levels of UL148 were detected in

Western blot analyses of protein extracts from drug treated and DMSO control

conditions. Therefore, differences in UL148 expression are unlikely to explain the

observed failure of UL148 and Hrd1 to coalesce into ER structures during inhibition of

either the ISR or PERK. Meanwhile, ISRIB and GSK2606414 treatments both led to

reduced levels of ATF4 accumulation, while the PERK inhibitor alone was able to

reduce the levels eIF2α-P, as indicated by a phospho-specific antibody. Therefore,

each of the drugs caused the expected effects on the PERK, eIF2α-P, and ATF4 axis

(**FIG 11C-D**).

A modest reduction in the levels of the viral envelope glycoprotein B (gB) was

observed during treatments with ISRIB or GSK2606414. This may suggest that the ISR

is required for optimal expression of the certain viral envelope glycoproteins. Even

though UL148 promotes ATF4 expression, likely via PERK activation (10), viruses that

lack *UL148* cause high levels of ATF4 accumulation at late times during infection, which

is when viral late gene products, such a gB, are expressed at their highest levels (10).

Hence, the activity of PERK and/or of the ISR may be required for optimal expression of

viral envelope glycoproteins. Regardless, the PERK inhibitor and ISRIB each prevented

the appearance of discrete UL148 ER structures while failing to substantially affect

UL148 expression. Therefore, the effects of the two inhibitors on UL148-dependent ER

reorganization were consistent with those seen during ectopic expression of the protein.

Taken together, our findings argue that the ISR is required for UL148 to cause

remodeling of the organelle.

DISCUSSION

The ER structures that we have identified are noteworthy in several regards.

Firstly, this example of virally-induced ER remodeling is wholly dependent on a single

viral gene product, UL148. Of course, there are other examples of individual viral gene

products that are necessary to grossly perturb ER morphology during infection which

are also found to be sufficient to induce such perturbations [e.g., (29, 30, 51, 52)].

Nonetheless, the ER structures caused by UL148 are— to the best of our knowledge, a

hitherto undescribed ultrastructural characteristic of the HCMV infected cell, which is

surprising given their size, scale, and prominence. Indeed, the average volume of the

structures at 96 h postinfection amounts to roughly 60% of the size of the nucleus of an

uninfected human fibroblast [~500 μm^3, (53)]. Although UL148 expression is

accompanied by UPR activation, ER reorganization occurs without killing the host cell,

even in the setting of ectopic expression when viral functions that inhibit programmed

cell death are absent. Nor does UL148 affect the yield of infectious virus during

replication in fibroblasts, since *UL148*-null mutant viruses replicate indistinguishably

from wildtype parental virus in this cell type (1, 8). As with all enveloped viruses, HCMV

requires the host cell secretory pathway to fold and process enormous quantities of viral

envelope glycoproteins that endow progeny virions with the molecular machinery for

infectivity. It is intriguing to consider how the infected cell tolerates spatial

reorganization of ER membranes and of the associated glycoprotein quality control

machinery without impacting the production of infectious progeny virions.

Furthermore, certain aspects of ER reorganization invoked by UL148 may be

novel to cell biology. In particular, the densely packed collapsed ER in close

association with varicosities that we observe to depend on UL148 during infection

appear to differ from the ER structures induced by other ER perturbagens. For

instance, a hereditary form of childhood cirrhosis caused by the *Z* allele of the alpha-1-

antitrypsin (Z-A1AT) gene (*SERPINA*) is characterized by polymerization of Z-A1AT

within the ER (26), leading to its accumulation in membrane-delimited inclusions (26,

27, 54). However, Z-A1AT inclusions are not observed to associate with regions of

collapsed ER, nor does Z-A1AT suffice to activate the UPR, even though its expression

sensitizes cells to other triggers of ER stress (27, 55).

HCMV encodes at least one other ER-resident immunevasin that activates the

UPR, US11 (56). UL148 binds the NK-cell and T-cell costimulatory ligand CD58 to

prevent its transport to the cell surface while US11 targets the heavy chain of class I

major histocompatibility complex (MHC) for ER-associated degradation (57, 58). UL148

causes CD58 to accumulate as an immature glycoform, presumably within the ER, but

does not lead to any obvious decrease in its overall abundance (9). Albeit that

intracellular forms of CD58 are in our hands refractory to detection by standard indirect

immunofluorescence protocols (data not shown), it seems reasonable to hypothesize

that UL148 sequesters CD58 into the unusual ER structures that it induces. Moreover,

our detergent solubility results suggest that UL148 may form aggregates or polymers in

the course of preventing CD58 presentation at the cell surface (FIG 8). It is worth

noting that UL148, an immunevasin targeting CD58, induces dramatic morphologic

rearrangements of the ER, while another HCMV immunevasin, US11, targeting MHC

class I for degradation evidently does not do so, even though both trigger ER stress.

Nonetheless, it seems unlikely that retention of CD58 would be required for ER

reorganization. In fact, the capacity of UL148 to retain CD58 may be functionally

separable from its peculiar effects on the morphology and organization of the ER.

Despite that the relationship between the mechanism for CD58 retention and the

formation of UL148-dependent ER structures remains unresolved, our findings may

suggest a mechanism to explain the influence of UL148 on the tropism of the virus for

epithelial cells. These effects, exemplified by a ~100-fold replication advantage of

UL148-null viruses during infection of epithelial cells, correlate with decreased

expression of glycoprotein O (gO), a viral envelope glycoprotein, both in virions and in

infected cells (1). We previously reported that gO behaves as a constitutive ERAD

substrate during infection and that immature, newly synthesized forms of gO show

enhanced stability in the presence of UL148 (8). Our findings herein show that UL148

causes large-scale sequestration of cellular factors important for ERAD, such as the ER

mannosidase EDEM1 and the E3 ubiquitin ligase SEL1L/Hrd1, into large membranous

structures. Moreover, the reticular ER, as indicated by the staining patterns of

antibodies specific for calreticulin, PDI, and the KDEL motif, appears to remain largely

intact following UL148-induced ER reorganization (FIG 3, 10). Since these

observations indicate that UL148 depletes ERAD factors from the ER during the

formation of the unusual structures, regions of ER not drawn into the structures might

be expected to offer a more permissive folding environment for polypeptides, such as

gO, which either fold slowly or inefficiently assemble into multiprotein complexes.

Based on these observations, one might hypothesize that UL148 alters ER

proteostasis by physically dislocating (or sequestering) key elements of the "mannose

removal time clock" system that marks poorly folding glycoproteins for destruction via

ERAD (59, 60). Because gO is both the defining subunit of the heterotrimeric gH/gL/gO

envelope glycoprotein complex that governs HCMV entry and cell tropism (5-7, 61, 62),

and a constitutive ERAD substrate (1, 8), ER reorganization might be required for the

effects of UL148 on HCMV cell tropism. Going forward, it will be crucial to determine

whether classical ERAD substrates, such as the null Hong Kong variant of alpha-1-

antitrypsin (63, 64) or ribophorin-332 (65, 66), are stabilized during UL148 expression,

as would be predicted if UL148 shifts ER proteostasis to negatively modulate ERAD.

In addition to having found that factors involved in proteasomal ERAD, such as

Hrd1 and EDEM1, are enriched at the UL148 structures, we observed that GABARAP,

a mammalian ortholog of yeast ATG8, localizes to the UL148 ER structures during

infection, and that another ATG8 ortholog, LC3B, and GABARAP both associate with

the structures during ectopic expression of UL148. This may suggest roles for

autophagy-related pathways in UL148-dependent reorganization of the organelle.

Since our results indicate that UL148 is degraded by a proteasome-dependent pathway

and not by lysosomes (**FIG S7**), it seems unlikely that selective autophagy of the ER is

directly involved in recycling or degrading these ER structures. Nonetheless, ATG8

family proteins may contribute to formation of the globular ER structures, since we

observed trafficking of UL148-GFP puncta to form large aggregates during live cell

imaging (**FIG 12, 15, Movies S2, S8**). The literature indicates that misfolded proteins

traffic in a microtubule (MT)-dependent manner to form pericentriolar structures termed

aggresomes, which in the case of ERAD substrates such as the ΔF508 mutant of the

cystic fibrosis transmembrane conductance regulator (CFTR), contain deglycosylated

protein that presumably has already undergone dislocation from the ER (67, 68). ATG8

family proteins such as LC3B and GABARAP not only play roles in degradation of

substrates via macroautophagy, but also bind MTs and recruit machinery for MT-

dependent transport of cargos (69-71). Therefore, GABARAP and LC3B may be

important for the recruitment of cellular machinery that transports perturbed ER cargoes

to sites of large-scale accumulation.

Although our results argue that the ISR is required for ER reorganization during

UL148 expression, precisely how UL148 triggers ER stress remains unknown. UL148

has been found to co-purify from cells with SEL1L, a component of the ERAD

machinery. Thus, it seems plausible that UL148 may inhibit the Hrd1/SEL1L complex,

which would cause the buildup of unfolded proteins and thus trigger the UPR. However,

inhibition of ERAD in and of itself seems unlikely to account for the formation of ER

structures. Another non-mutually exclusive possibility is that UL148 multimerizes or

aggregates in a manner that constricts the ER lumen. For instance, the assembly of

UL148 molecules on opposite sides of the ER lumen might constrict the organelle in a

manner consistent with the collapsed regions of ER observed in our EM results (**FIG 6,**

8).

Additional work will be needed to decipher the molecular mechanisms by which

UL148 causes reorganization of the ER, and to determine its physiological relevance in

the setting of viral biology. Nonetheless, we have shown that UL148, when fused to a

fluorescent protein (FP), suffices both to trigger and to indicate the presence of a

functional ISR (**FIG 10-11, S5**). Hence, UL148-FP fusions may prove useful in high

throughput chemical-genetic screens to identify novel small molecule inhibitors of the

ISR as well as to identify cellular genes involved in stress-dependent remodeling of the

ER. Interestingly, the two pharmacological agents that block UL148-dependent ER

remodeling, ISRIB and GSK2606414, are known to prevent the formation of stress

granules (SGs) (43, 72). SGs are comprised of condensed aggregates of protein and

RNA which occur due to stalled mRNA translation in the context of disease states, such

as amyotrophic lateral sclerosis (73), as well as during treatments with toxic agents

such as arsenite (43, 72).   It is fascinating to consider that mechanistic parallels exist

between the formation of SGs and the UL148-dependent reorganization of the ER.

Moreover, given the broad importance macromolecular aggregation in pathological

conditions such as neurodegenerative diseases (74), certain of which also involve

defects in ER proteostasis and aberrant activation of the UPR, UL148 may hold promise

as tool to discover new agents to ameliorate disease.

MATERIALS AND METHODS.

Cells and virus.

hTERT-immortalized human foreskin fibroblasts (8), derived from ATCC HFF-1 cells

(SCRC-1041) were maintained in Dulbecco's modified Eagle's medium supplemented

with 5%-10% newborn calf serum (Millipore Sigma) and antibiotics (complete DMEM)

exactly as described previously (8). i148^{HA} and i159^{HA} ARPE-19 epithelial cells (10),

which upon treatment with 100 ng/mL doxycycline, express HA-tagged UL148 or

Rh159, respectively, were likewise maintained in complete DMEM. For live cell imaging

studies, i148^{HA} and i159^{HA} ARPE-19 were maintained in Opti-MEM (Thermo Fisher)

supplemented with 3% certified tet-approved fetal bovine serum (FBS), 20 µg/mL

gentamicin sulfate, 1 µg/mL puromycin HCl and 10 µg/mL ciprofloxacin HCl.

Telomerase-immortalized rhesus fibroblasts (75) were a kind gift of Peter A. Barry, and

were maintained in complete DMEM. The acute monocytic leukemia cell line THP-1

(TIB-202) was obtained from ATCC (Manassas, VA) and maintained as suspension

cultures in RPMI 1640 medium supplemented with 10% FBS (Millipore Sigma) and

antibiotics (10 µg/mL ciprofloxacin and 25 µg/mL gentamicin). THP-1 were

differentiated into adherent macrophages by incubating for 48 h in the presence of 100

nM 2-O-tetradecanoylphorbol 13-acetate (Millipore Sigma), and subsequently infected

with the indicated viruses at an MOI of 5 TCID$_{50}$/cell.

Infectious bacterial artificial chromosome (BAC) clones of the following HCMV

strains were used for this study: TB40/E (also known as TB40-BAC4 or TB_WT) (11)

as well as its derivatives TB_148^{HA} and TB_148$_{STOP}$, which were used in our previous

studies (1, 8, 10); TR (TR*gfp*) (76, 77); Merlin repaired for *UL128* and *RL13* harboring

tetO sequences upstream of *UL131* (pAL1393) (78); AD169 repaired for *UL131*

(AD_r131)(1, 79) and AD_r131_148[HA], a derivative of AD_r131 that carries a full length

UL148 (from strain TB40/E) tagged with an HA-epitope at the original *UL148* locus (8).

An infectious BAC clone of rhesus CMV (RhCMV) strain 68-1 (80) and a derivative that

expresses an HA-tagged Rh159 (details below) were also used for certain experiments.

The methods used for reconstitution of HCMV from purified BAC DNA, cultivation of

virus and preparation of stocks, including ultracentrifugation through sorbitol cushions

and determination of infectious titers by tissue culture infectious dose 50 ($TCID_{50}$) assay

have been described elsewhere (8, 10).

Construction of recombinant viruses and new plasmids for the study.

New recombinant viruses for this study were derived from BAC-cloned HCMV

and RhCMVs using *en passant* mutagenesis in GS1783 *E. coli,* a strain K12 derivative

that expresses the homing endonuclease I-SceI upon treatment with L-(+)-arabinose

(81, 82). Recombinant BACs were confirmed by Sanger DNA sequencing of the

modified regions, which was performed by Genewiz, Inc. (Piscataway, NJ) (not shown).

Oligonucleotide primers for construction of recombinant viruses and plasmids were

custom synthesized by Integrated DNA Technologies (Coralville, IA); sequences are

provided in **Table S1**. Type II restriction endonucleases, T4 DNA ligase, and Gibson

assembly reagents (NEB HiFi Assembly Master Mix) for generation of recombinant

DNAs and for routine molecular cloning and subcloning procedures were obtained from

New England Biolabs (Ipswitch, MA). KOD Hot Start DNA polymerase (EMD Millipore)

was used for all PCR reactions.

To construct TB_159^{HA}, a strain TB40/E derivative in which the UL148 open

reading frame is replaced by Rh159 we carried out the following steps. Primers

Rh159_Fw and Rh159_HA_Rv were used to amplify and HA-tag the *Rh159* ORF from

pcDNA3.1+_Rh159_IRES_GFP (a gift from Klaus Früh, OHSU). The PCR product was

inserted into EcoRV-digested pEF1α using Gibson assembly. A PCR product

containing an ISce-I excisable kanamycin cassette was amplified from TB_148^{HA}_*ISce-*

Kan integrate BAC (1) using primers PpuISceKanGibs_Fw and PpuISceKanGibs _Rv

and Gibson-assembled into PpuMI-digested pEF1α_Rh159 plasmid to yield plasmid

pEF1α_Rh159_ISceKan. Primers Rh159_Fw_recomb and Rh159_Rv_recomb were

used to generate a PCR product from template plasmid pEF1α_Rh159_ISceKan. The

PCR product was electroporated into GS1783 E. coli harboring the TB_148^{HA} BAC, and

Kan^{r} colonies harboring TB_Δ148_Rh159_ISceKan integrate BACs were isolated on

Luria-Bertiani agar plates. Bacterial colonies representing kanamycin resistant

integrates were obtained and were subsequently resolved to 'scarlessly' remove the

positive selection marker by standard *en passant* protocols (81, 82) to yield

TB_Δ148_Rh159^{HA}, which we abbreviate herein as TB_159^{HA}. TB_159^{HA} was

sequence-confirmed using primers TB_159HAseq_F and TB_159HAseq_R. Similar

strategies were used to modify the RhCMV 68-1 BAC (80) to incorporate sequences

encoding an HA-epitope immediately before the stop codon of *Rh159*. An *en passant*

strategy in which a shuttle plasmid carrying *I-SceI-Kan^{r}* disrupted by in-frame nonsense

codons was used insert premature nonsense codons in the *UL148* CDS in the context

of HCMV strains Merlin (pAL1393) and TR (TR*gfp*), and was applied as described

previously for generating TB_148$_{STOP}$, the UL148$_{STOP}$ mutant of strain TB40/E (8).

To construct "tet-on" lentivirus vector plasmids containing UL148 fused to GFP,

we used primers Gibs_eGFP_Rv and 148_eGFP_Fw to amplify the GFP gene from a

dsDNA gBlock, EGFP-P2A-3XHA, synthesized by Integrated DNA Technologies

(Coralville, IA), which was a gift of Matthew D. Woolard (LSU Health Sciences Center,

Shreveport, LA).   In a separate PCR, the *UL148* gene was amplified from plasmid

pcDNA3.1-UL148^{HA} (10) using primers Gib_148_Fw and 148_noStopRv. The two

products were assembled, using Gibson Assembly, together with EcoRV opened

pcDNA3.1(+) (Invitrogen) to produce pcDNA3.1-UL148-GFP.   After confirming the

absence of spurious mutations by Sanger DNA sequencing (Genewiz, not shown), the

UL148-GFP cassette from pcDNA3.1-UL148-GFP was released by EcoRI and NotI

digestion and ligated into pOUPc turboRFP-link plasmid which had been linearized

using the same restriction sites. pOUPc turboRFP-link is a derivative of lentiviral vector

pOUPc (10) that was constructed by reinsertion of turboRFP (tRFP). Briefly, tRFP was

amplified from the original pOUPc-turboRFP using primer pair tRFP linker Fw and tRFP

linker Rv and then reinserted by Gibson Assembly with pOUPc turboRFP that had been

double-digested using EcoRI and Not I in order to linearize the vector and remove the

turboRFP (RFP) sequence.

To express Rh159 fused to eGFP, primers Gibs_eGFP_Rv and 159_eGFP_Fw

were used to amplify the GFP gene from EGFP-P2A-3XHA, and in a separate PCR

reaction, primers Gibs_159_Fw and 159_noStop_Rv Rh159 were used to amplify

Rh159 from pcDNA3.1-Rh159^{HA} (10). The latter two PCR products were assembled

together with EcoRV linearized pcDNA3.1(+) using Gibson Assembly, resulting in

plasmid pcDNA3.1-Rh159. The latter plasmid was used as template in a PCR reaction

with primers Gibs_Age_159_Fw and Gibs_Mlu_GFP_Rv. The resulting PCR product

was Gibson assembled into the lentiviral vector pOUPc (10) which had been double-

digested with Mlu I and Age I.

Drug treatments.

The PERK inhibitor, GSK2606414 (50), doxycycline hyclate, and folimycin were

obtained from MilliporeSigma (Burlington, MA). ISRIB (42-46) was obtained from

MilliporeSigma or APExBio (Boston, MA) and epoxomicin was procured Selleck

Chemicals (Houston, TX).   ISRIB was dissolved in DMSO to make a 10,000× stock

solution (2 mM) and used at a final working concentration of 200 nM. GSK2606414 was

dissolved in DMSO to make a 10,000× stock solution (11 mM) and used at a final

working concentration of 1.1 µM. Folimycin was prepared as a 1000× (115.46 µM)

stock solution in DMSO and used at 115 nM final. Epoxomycin was prepared as a 100×

(2 mM) stock solution in DMSO and used at a final concentration of 20 µM.

Confocal microscopy and live-cell imaging. Confocal indirect immunofluorescence

microscopy imaging on fixed cells was carried out using Leica TCS SP5 Spectral

Confocal Microscope (Leica Microsystems, Heidelberg, Germany) using a Leica HCX

PL APO CS 63x/1.4-0.6NA objective under oil immersion, except for the image shown

in Figure 1C, which was captured on a Nikon SIM-E and A1R confocal microscopy

system (Nikon Instruments, Melville, NY) using a Nikon SR Apo TIRF 100x/1.49NA

objective under oil immersion. Images for different fluorophore channels were acquired

using sequential scanning. Direct immunofluorescence live cell imaging data were

collected using the Nikon SIM-E microscope using a Nikon Apo 60x/1.40NA DIC

objective.   The 3D projection shown in FIG 1 was generated by NIS-Elements AR

Analysis 4.60.00 (64-bit) software (Nikon) from Z-stacks captured on the Nikon SIM-E /

A1R microscope using a Nikon SR Apo TIRF 100x/1.49NA objective. For FIG 11, 3D

projections were generated by Imaris x64 9.3.0 software (Bitplane, Inc.) in maximum

intensity projection (MIP) mode.

For fixed cell imaging experiments other than those shown in Fig S1D, cells were

seeded on 12 mm circular, No. 1 thickness microscope cover glass (200121; Azer

Scientific, Morgantown, PA). At the indicated times post-treatment and or post-infection,

cells were washed with phosphate buffered saline (PBS) [137 mM NaCl, 2.7 mM KCl, 10

mM Na_2HPO_4, and 1.8 mM KH_2PO_4, pH 7.4]; fixed for 15 min at room temperature in PBS

containing 4% (wt/vol) paraformaldehyde (Fisher Scientific, Waltham, MA), washed in

PBS, permeabilized for 10 min using 0.1% Triton X-100 (in PBS), subsequently washed

again in PBS, and then blocked for 45 min at 37 °C in PBS containing 5% (vol/vol) normal

goat serum (Rockland Immunochemicals, Limerick, PA). Cells were then washed three

times in PBS followed by incubation in 1% Human Fc Block (BD Biosciences, San Jose,

CA) in PBS, for additional 45 min at 37 °C. Cells were then incubated in the presence of

primary antibodies for 1 h at 37 °C or 4 °C overnight, and then washed three times with

PBS containing 0.1% Tween-20 (PBST) for 5 min per wash. Alexa Fluor-labeled goat

polyclonal secondary antibodies (all from ThermoFisher Invitrogen, Waltham, MA, see

Table S2) were used for secondary detection.   The slides were then mounted using

Prolong Gold anti-fade reagent containing DAPI (ThermoFisher) and placed under a

Leica TCS SP5 confocal microscope for image acquisition using a Leica 63× oil

immersion objective lens (Leica Microsystems, Buffalo Grove, IL).

For indirect immunofluorescence staining results of primary clinical isolates (SI Fig

S1D), four clinical HCMV isolates obtained by routine testing of throat swabs from patients

of the Ulm University Medical Center were provided by the diagnostic laboratory of the

Institute of Virology in Ulm. Sample material was applied to human fibroblasts and

incubated for several days until HCMV-positive cells could be detected. Infected cells

were then seeded together with uninfected fibroblasts, incubated for up to 5 days until

plaques were formed and processed for indirect immunofluorescence staining. ERQC

compartments were detected by staining for calnexin (CNX, E10; Santa Cruz

Biotechnology, mouse, 1:50 dilution), the cVAC was detected by staining for HCMV

pUL71 (rabbit, 1:500). Secondary antibody used for detection of CNX was Alexa Fluor

555 labeled goat anti-mouse IgG (1:1000) and for pUL71 detection, Alexa Fluor 488

conjugated goat anti-rabbit IgG (1:1000). Confocal images were acquired using the 63×

objective of a Zeiss Observer Z1 fluorescence microscope equipped with Apotome and

Zen software 2.3 (Carl Zeiss Microscopy GmbH, Jena, Germany).

FRAP. FRAP studies were carried out on using live ARPE-19 cells from the i148^{GFP}

and i159^{GFP} populations as follows. Cells were seeded as above for live-cell imaging,

induced for transgene expression using dox (100 ng) for 24 h, and then placed on an

incubated sample stage of a Nikon A1R SIM-E imaging system equipped with an SR

Apo TIRF 100x/1.49NA objective (Nikon). Three rectangular regions in image fields

were defined for measurement of (i) background signal, and of two regions with

comparable initial GFP signals, (ii) one designated as a control region (no

photobleaching) and another for (iii) photobleaching and recovery of signal after

photobleaching. Photobleaching of selected regions was carried out for 20 s using 405

nm laser light from a LU-N4 laser fiber (Nikon) [power at source: 15 mW, power at

objective: 8 mW]. Images and signal intensity measurements were captured every 2 s

at a rate of 1 frame per s (488 nm excitation, FITC channel) starting immediately before

photobleaching (t=1), and from t=20 s to t=320 s after.

Electron microscopy.   Procedures to prepare samples for transmission electron

microscopy (TEM) included high-pressure freezing (HPF), freeze substitution, and Epon

embedding, which were carried out as described previously (83).   Briefly, human

fibroblasts were seeded in µ-Slides (Ibidi GmbH, Martinsried, Germany) containing

carbon-coated sapphire discs (Engineering Office M. Wohlwend GmbH) 1 day prior to

infection at 80 to 90% confluence. Cells were infected with virus overnight at MOI 1.

Medium containing viral inocula was replaced with fresh medium the next day. Infected

cells on sapphire discs were fixed by using HPF with a Compact 01 high-pressure freezer

(Engineering Office M. Wohlwend GmbH, Sennwald, Switzerland) at 5 dpi. Thereafter,

cells on sapphire discs were processed by freeze-substitution and subsequently

embedded in Epon (Fluka, Buchs, Switzerland). Ultrathin sections of the Epon-embedded

cells were cut with an ultramicrotome (Ultracut UCT; Leica Microsystems, Wetzlar,

Germany) and placed on Formvar-coated single-slot grids (Plano GmbH, Wetzlar,

Germany). Grids were examined in a Jeol JEM-1400 (Jeol Ltd., Tokyo, Japan)

transmission electron microscope equipped with a charge-coupled-device (CCD) camera

at an acceleration voltage of 120 kV. Fixation and embedding of infected cells for

scanning transmission electron microscopy (STEM) tomography was the same as

described for TEM. Additional sample preparation steps were conducted as described

previously (84, 85). Tomogram acquisition was conducted on a STEM Jeol JEM-2100F

with an acceleration voltage of 200 kV. Tilt series were acquired from 600 nm thin

sections from +72° to -72° with a 1.5° increment using the bright field detector. Image

series were reconstructed to tomograms by weighted back projection with the IMOD

software package (86). 3D visualization of the membrane structures was performed using

Avizo lite software (Visualization Science Group, Burlington, MA, USA) by threshold

segmentation.

Western blotting. Western blotting procedures, including primary and secondary

antibodies used for detection of HA tag, UL148, ATF4, eIF2α, P-eIF2α (Ser51), the

HCMV viral nuclear antigen IE1, and the conditions used for detection of Ser51

phosphorylated eIF2α, were carried out as described previously (8, 10). Additional

antibodies used in this study were mouse anti-GAPDH (Cat. No. 60004-1, Proteintech,

Rosemont, IL,), mouse anti-gB clone 27-180 (87) (a generous gift of William J. Britt),

and anti-GFP (D5.1) XP® Rabbit mAb #2956 (Cell Signaling Technology, Danvers, MA).

Solubility analyses. 2.0×10^5 human fibroblasts were infected at an MOI of 1 TCID$_{50}$

per cell with TB_148^{HA} or TB_159^{HA}. The following day, cells were washed twice with

PBS to remove viral inoculum and replenished with DMEM containing 5% newborn calf

serum. At the indicated times post-infection, cells were washed once with PBS and

lysed by direct addition of 100 μL RIPA buffer [25 mM HEPES (pH 7.5), 400 mM NaCl,

0.1% SDS, 0.5% sodium deoxycholate, 1% NP-40, supplemented with 1$\times$ protease

inhibitor cocktail (ApexBio)]. Lysates were collected and rotated at 4°C for 1 h.

Insoluble material was pelleted by centrifugation (21,000 $\times$ g, 35 min). Supernatants

containing soluble material were transferred to a fresh microfuge tube, and 33 μL of 4$\times$

Laemmli sample buffer [200 mM Tris (pH 6.8), 8% SDS, 40% glycerol, 0.08%

bromophenol blue] was added to bring final volume to 133 μL. The pellet was disrupted

in 133 μL of 1X Laemmli buffer prepared by diluting 4$\times$ Laemmli buffer in RIPA buffer.

Samples were reduced by addition of beta-mercaptoethanol (5% final, v/v) and boiled at

90°C for 10 min. 40 μL of each sample was resolved by SDS-PAGE (12%

polyacrylamide gel), transferred overnight to nitrocellulose membrane, and

immunoblotted with antibodies against HA epitope or HCMV gB. Quantitation of

secondary antibody fluorescence signals were performed using an Odyssey CLx

scanner (Li-Cor, Inc., Lincoln, NE) in auto-scan mode. For each time-point, the signals

from RIPA-soluble and insoluble bands were summed to yield total signal, and the ratio

of insoluble band signal to total signal were also reported as percent insoluble HA over

total HA signal.

Statistical analyses. Statistical analyses were carried out using GraphPad Prism 8.1.0

for MacOS (GraphPad, Inc., San Diego, CA).

ACKNOWLEDGEMENTS.

This project was supported by NIH grants R01-AI116851 (to J.P.K.) and P30-

GM110703. Its contents are solely the responsibility of the authors and do not

necessarily represent the official views of the NIAID or the NIGMS.

We are especially grateful to Erik L. Snapp (HHMI, Janelia Laboratories) for

helpful discussions and detailed advice on interpretation of EM and confocal image

data. We also thank Dong Yu (Washington University, St. Louis, MO, currently GSK

Vaccines), Thomas E. Shenk (Princeton University, Princeton, NJ), Richard J. Stanton

(Cardiff University, United Kingdom), William J. Britt (University of Alabama,

Birmingham), W. L. William Chang and Peter A. Barry (both of the University of

California, Davis; Davis, CA), Christian Sinzger (University Medical Center Ulm, Ulm,

Germany), Klaus Früh (Oregon Health Sciences University, Beaverton, OR), and

Gregory A. Smith (Northwestern University, Chicago, IL) for generously sharing

reagents.

AUTHOR CONTRIBUTIONS.

Performed experiments: HZ, CR, CCN, MNAS, JvE. Data analysis and interpretation:

HZ, CR, CCN, JvE, CS, JPK. Designed experiments: HZ, CR, CCN, JvE, and JPK.

Contributed new reagents: HZ, CH, CCN, JPK. Electron microscopy and STEM

tomography: CR and JvE. Obtained funding: JPK. Wrote the manuscript: JPK with

comments from JvE and CR.

869
870
FIG 1. UL148 reorganizes ER markers into anomalous structures during HCMV infection.
Fibroblasts infected at 1 MOI with either **(A)** wildtype HCMV strain TB40/E (TB_WT) or **(B)** an
isogenic *UL148*-null mutant (TB_148_{STOP}), were fixed at 96 h postinfection (hpi) and imaged by
confocal microscopy after staining with antibodies specific for calnexin or Hrd1 (magenta) and
glycoprotein H (gH, green), as indicated. DAPI (blue) signal was used to counterstain nuclei. **(C)**
Percentage of fibroblasts that contain Hrd1 structures at 96 hpi; fifty gH-positive cells for each
condition were scored for the presence or absence of UL148 structures (also see SI Fig S1A).
(**D**) 3D confocal image projection of a TB_WT infected fibroblast stained at 96 hpi for Hrd1 and
gH. **(E)** Volumetric measurements were made from sixteen infected cells, stained with phalloidin-
AlexaFluor 594 conjugate to estimate total cell volume, with Hrd1 antibody to estimate the volume
of UL148-dependent ER structures, and with DAPI to estimate nuclear volume.

887
888
FIG 2. Characterization of TB_148^HA and TB_159^HA viruses. **(A)** Human fibroblasts were
infected at MOI 1 with the indicated recombinant HCMVs. Cell lysate samples harvested at the
indicated times postinfection (h postinfection, hpi) were analyzed by Western blot to detect HA-
tagged UL148 or Rh159, the 72 kD viral nuclear antigen IE1-72 (IE1), and GAPDH. **(B)** Single-
cycle viral replication kinetic curves from MOI 1 infected fibroblasts were plotted by determining
the titer in tissue culture infectious dose 50 (TCID$_{50}$) from supernatants collected at the indicated
times postinfection.

896
FIG 3. UL148 localizes to unusual ER compartments that are enriched for glycoprotein
897
quality control markers. Fibroblasts infected at MOI 1 with either TB_148[HA] (panels: **A, C, E,**
898
G, I, K, M, O, Q, S) or TB_ 159[HA] (panels: **B, D, F, H, J, L, N, P, R, T**) were fixed at 96 h
899
postinfection, and imaged by confocal microscopy after co-staining with antibodies specific for HA
900
(UL148 / Rh159, magenta in merge) and the indicated cellular markers (green in merge). DAPI
901
(blue) counterstaining is shown in merged image.
902

903
904
905

906
907 **FIG 4. Quantification of co-localization between UL148 and cellular markers.** A Pearson's
908 correlation coefficient (*r*) was calculated using NIH ImageJ software to estimate the degree of co-
909 localization between UL148 (HA signal) and each of the indicated cellular markers. A minimum
910 of 30 cells were analyzed per marker. The arithmetic mean for each co-localization analysis result
911 is shown as a blue line, and data points for individual cells analyzed are plotted as circles.
912

913
914
FIG 5. TEM of ER structures in wild-type HCMV infected cells. Human fibroblasts infected
with wildtype HCMV (TB_WT) were fixed by high-pressure freezing and freeze substitution at
day 5 postinfection and imaged using TEM. Panels (**A**) and (**B**) show cell overview at left. For
each cell, the boxed region is shown at higher magnification. Scale bars; left panels: 10 μm,
right panels: 2 μm. rER: rough ER; Nu: nucleus; cVAC: cytoplasmic viral assembly
compartment. Solid arrowheads indicate the UL148-dependent ER structures of interest.

FIG 6. TEM of *UL148*-null HCMV infected cells. Human fibroblasts infected with a *UL148*-
null mutant (TB_148$_{STOP}$) were fixed by high-pressure freezing and freeze substitution at day 5
postinfection and imaged using TEM. At the upper left an overview panel of a representative
cell is shown with panels (**A**), **(B)**, and **(C)** each boxed. The boxed regions are expanded at
higher magnification at right (**A**) and below (**B, C**). Scale bars for upper left overview panel: 10
µm, for zoomed panels: 2 µm. rER: rough ER; Nu: nucleus; cVAC: cytoplasmic viral assembly
compartment.
930

931
932
FIG 7. STEM tomography of UL148-dependent ER structures in HCMV infected cells.
Human fibroblasts infected with wildtype HCMV (TB_WT) were fixed by high-pressure freezing
and freeze substitution at day 5 postinfection and tomograms were recorded by STEM. (**A**)
Shows a virtual section through the tomogram of a virus-induced membranous structure;
asterisks denote ER structures of distended lumenal space. White arrowheads indicate sites at
which membranes originating from distended ER cisternae continue into areas of involuted
collapsed ER. Scale bar: 1 µm. (**B**) Shows the same virtual section as in (A) tilted and with a
3D visualization of the membranous network (green) of the entire tomogram. The same
distended ER cisternae as in (A) are marked by asterisks. (**C**) Shows a cross section through
(B) to visualize the membrane profile of the membranous structures. The region delimited by a
white box is shown in a higher magnification on the right. Finer detail of the enlarged ER
cisternae and the connections between them are readily visible; asterisks indicate the same
distended ER cisternae as in panels (A) and (B). Also, see SI Movie S1. Scale bar: 1 µm.

946
947
FIG 8. Solubility analysis of UL148 and Rh159. (**A**) Human fibroblasts were infected at MOI
1 $TCID_{50}$/cell with HCMV strain TB40/E derived viruses TB_148[HA], which expresses UL148
fused at its C-terminus to an HA-epitope tag, or TB_159[HA], which lacks *UL148* and instead
expresses rhesus CMV Rh159 carrying a C-terminal HA tag. At the indicated times
postinfection (hpi), infected cells were collected in radioimmunoprecipitation (RIPA) lysis buffer,
centrifuged at 21,000 × *g* for 30 min, after which supernatant (S) and pellet (P) fractions were
boiled in gel loading buffer containing 2% sodium dodecyl sulfate (SDS). Equivalent portions of
supernatant and pellet were resolved by SDS-polyacrylamide gel electrophoresis, transferred to
a nitrocellulose membrane for detection of protein species immunoreactive to antibodies against
HA epitope (HA) and HCMV glycoprotein B (gB) and for total protein signal using Ponceau S
reagent (Ponceau). (**B-C**). Signal intensity of fluorophore-conjugated secondary antibodies in
anti-HA Western blots were measured from three independent biological replicates of the
experiment shown in (A); error bars indicate standard deviation. (**B**) The fluorescent signal for
each infection time point condition (y-axis is in arbitrary units, in hundred thousands). (**C**) The
amount of signal found in the insoluble (pellet) fraction relative to the total signal (pellet plus
supernatant) for each infection time point are plotted as percentage values.
964
965

966
967
FIG 9. ISR activation accompanies redistribution of ER markers during ectopic
expression of UL148. (A) Lysates of tet-on ARPE-19 cells expressing either UL148 or Rh159
fused to an HA tag, i148^{HA} and i159^{HA}, respectively, were collected at the indicated times post
doxycycline (dox) induction and analyzed by Western blot for the expression of the indicated
proteins and for the abundance of eIF2α phosphorylated at Ser51 (P-eIF2α) using a phospho-
specific antibody. (**B**) Pearson's correlation coefficient (*r*) values were calculated using NIH
ImageJ software estimate the degree of co-localization between UL148 (HA signal) and the
indicated cellular markers. A minimum of 30 cells were analyzed per marker. Arithmetic means
for each co-localization analysis result are shown as blue lines, and data points for individual
cells analyzed are plotted as circles.
978
979
980

981
982
FIG 10. Inhibition of the integrated stress response impedes UL148-mediated ER
remodeling. (A-B) "Tet-on" ARPE-19 epithelial cells that inducibly express UL148 fused to
green florescent protein (gfp), i148^{gfp}, were induced for transgene expression using 100 ng/mL
doxycycline (dox) in the presence of DMSO carrier, ISRIB (200 nM), or GSK2606414 (1.1 µM)
and monitored by anti-UL148 Western blot for expression of UL148-GFP over a series of time
points (h post treatment with dox, hpt). (**C-E**) Live-cell imaging of UL148-GFP expression
patterns in the presence of ISRIB (200 nM), GSK2606414 (1.1 µM), or DMSO vehicle (0.01%).
Main scale bar represents 50 µm. For inset panels at upper left of each image, which are
magnified 2.4× relative to the main image, the scale bar represents 10 µm. Also see SI Movies
S8-S10.
993

994
995
996
FIG 11. Inhibition of the ISR prevents the coalescence of Hrd1 and UL148 into discrete
structures during infection. (**A**) Representative 3D maximum intensity projections of confocal
imaging Z-stacks obtained from cells infected at MOI 1 for 96 h with HCMV strain TB40/E carrying
an HA-tagged *UL148*, TB_148[HA], and maintained in the presence of ISRIB (200 nM),
GSK2606414 (1.1 µM), or DMSO carrier alone (0.1% vol/vol). In merged images HA signal is
shown in magenta, Hrd1 in green, and DAPI counterstaining in blue. (**B**) The percent of Hrd1
antibody signal involved in discrete structures at 96 hpi were calculated for a minimum of 30 cells
per condition using Imaris x64 9.3.0 software. Statistical significance was determined using a
one-way ANOVA followed by Tukey's post-test; **** represents a P-value of <0.0001. The
arithmetic mean for co-localization analysis results are shown as blue lines with data points for
individual cells analyzed plotted as circles, squares or triangles, as indicated. (**C-D**) Western blot
analyses of fibroblasts infected at MOI 1 with TB_148[HA] and maintained in the presence of ISRIB
(200 nM), GSK2606414 (1.1 µM), or DMSO carrier alone (0.01%); hpi: h post infection (hpi). Note:
a phospho-specific antibody was used for detection of eIF2α phosphorylated at Ser51 (P-eIF2α).

FIG S1: **Additional confocal microscopy results from infected cells.** (**A**) Lower magnification view of results from FIG 1; staining of gH and Hrd1 at 96 hpi of human fibroblasts with wildtype HCMV strain TB40/E (WT) versus *UL148*-null derivative TB_148$_{STOP}$. (**B**) gH and Hrd1 at 96 hpi of human fibroblasts with wildtype HCMV strain Merlin (ME_WT) or a *UL148*-null derivative of the same virus (ME_148$_{STOP}$). (**C**): gH and Hrd1 at 96 hpi of human fibroblasts with wildtype HCMV strain TRgfp recovered from BAC-cloned TRgfp (TR_WT) or a *UL148*-null derivative of the same virus (TR_148$_{STOP}$). (**D**) Staining of the viral tegument protein UL71 and calnexin of human fibroblasts infected with primary clinical HCMV isolates obtained from patient throat swabs (at 5 dpi). (**E**) HA and Hrd1 in THP-1 macrophages (differentiated using phorbol ester treatment) fixed at 96 hpi with HCMV strain TB40/E derivative viruses TB_148^{HA} or TB_159^{HA}. (**F**) HA and Hrd1 in ARPE-19 epithelial cells infected for 96 h with either TB_148^{HA} or TB_159^{HA}. (**G**) HA and Hrd1 in ARPE-19 infected for 96 h with HCMV strain AD169 repaired for *UL131* and to which an HA-tagged *UL148* from TB40/E was restored to the native *UL148* locus (AD^{r131}_148^{HA}) vs parental AD169 repaired for *UL131* (AD^{r131}). For ADr131 we did not employ a viral marker (e.g., HA) to identify infected cells, so the appearance of syncytia and the characteristic kidney-bean shaped nucleus was used to indicate infected cells, and a slightly lower magnification was used to best show these features. (**H**) HA and Hrd1 in telomerase-immortalized rhesus fibroblasts infected with RhCMV strain 68-1 carrying an HA-tag at the C-terminus of Rh159.

1039
1040
1041 **FIG S2: LC3B, syntaxin-6, and gH staining in HCMV infected cells.** Human fibroblasts
1042 were infected with the indicated viruses for 96 h and then fixed, permeabilized, and stained
1043 using antibodies specific for the indicated proteins. Indirect immunofluorescence images were
1044 captured using a 63X objective on a Leica SP5 confocal microscope. gH: glycoprotein H, STX6:
1045 syntaxin-6.

FIG S3: Cellular proteins involved in glycoprotein quality control are recruited with differing kinetics to UL148 ER structures. Fibroblasts infected with TB_148^{HA} at 1 MOI were fixed at the indicated time points (days post infection, dpi) and imaged by confocal microscopy after staining with antibodies specific for HA (UL148, magenta), CNX (green, **A**) , Hrd1 (green, **B**) or VCP(green, **C**), DAPI (blue).

1061
FIG S4. UL148 is sufficient to remodel the ER. Expression of HA-tagged UL148 or Rh159
was induced in "tet-on" ARPE-19 epithelial cells, i148HA and i159HA, respectively. Cells were
fixed at 48 h postinduction for indirect immunofluorescence staining for the indicated cellular
markers (green) together with HA (magenta). Scale bar: 10 μm.

FIG S5. Live-cell Imaging of UL148-GFP and Rh159-GFP during induced ectopic expression. "Tet-on" ARPE-19 epithelial cells that inducibly express either UL148 or Rh159 fused to green florescent protein (gfp), i148^{gfp} (**A**) and i159^{gfp} (**B**), respectively, were induced for transgene expression using 100 ng/mL doxycycline (dox) and imaged using live-cell microscopy. Images from the selected time points (h post treatment with dox, hpt) are shown. The main scale bar represents 50 µm. For inset panels at upper left of each image, which are magnified 2.4× relative to the main image, the scale bar represents 10 µm. Also see SI Movies S2-S3.

1080
1081
FIG S6. FRAP. ARPE-19 cells that inducibly express Rh159-GFP, i159^{gfp} (**A**) or UL148-GFP,
i148^{gfp} (**B-C**) were doxycycline induced for 24 h (100 ng/mL doxycycline (dox). Separate regions
of GFP signal in selected cells were photobleached (405 nm laser) or left unbleached, while a
third region lacking GFP signal was chosen as a background reference were measured before
and during fluorescence recovery after photobleaching (FRAP). Note: background signal was
not plotted because values were resolvable from the x-axis. GFP signal intensity is plotted over
a time period (seconds, s) starting with an exposure at t=0 (immediately before photobleaching),
and including measurements taken every 2 s after photobleaching (0-20 s) until termination of
the measurement series at t=322 s (300 s of FRAP). (**D-F**) Images from the selected time points
(s, seconds) immediately before and after bleaching, and during fluorescence recovery period.
Also see SI Movies S4-S6.

FIG S7: UL148-GFP is stabilized by inhibition of proteasomal but not lysosomal degradation. "Tet-on" ARPE-19 cells that inducibly express UL148-GFP (i148^{gfp}) were induced for 24 h by the addition of 100 ng/mL doxycycline (dox), after which the dox inducing agent was washed out and medium containing either the proteasome inhibitor epoxomicin (20 µM) or the proton pump inhibitor folimycin (115 nM) was added and samples were harvested for Western blot analysis at the indicated times post treatment (h post treatment, hpt) with folimycin or epoxomicin. DMSO was added at 0.1% to control for folimycin, or 1% to control for epoxomicin. Also see SI Movie S7.

TABLE S1. Oligonucleotide primers and synthetic DNAs.

Name	Sequence (5' to 3')
148_eGFP_Fw	ACCTGGTGTCGCGGCGTCGGATGGTGAGCAAGGGGGAGGAGCTTTTTACTGGCGTTGTTC
Gibs_eGFP_Rv	CGGCCGCCACTGTGCTGGATTACTTATACAGCTCGTCCATACCCAG
Gib_148_Fw	TGTGGTGGAATTCTGCAGATTGCCACCATGTTGCGCTTGCTG
148_noStopRv	CCGACGCCGCGACACCAGGTAGGTTATC
159_eGFP_Fw	TCTAAAACGCAGTCGTGAAGCTCATATGGTGAGCAAGGGGGAGGA
159_noStop_Rv	ATGAGCTTCACGACTGCGTTTTAGAAGTG
Gibs_159_Fw	TGTGGTGGAATTCTGCAGATTGCCACCATGGCCTACAACAGC
Rh159_Fw	TGTGGTGGAATTCTGCAGATACCATGGCCTACAACAGCTTCCTTCTCAGCTGCCTTACCATCGCACTTTTACTG
Rh159_HA_Rv	CGGCCGCCACTGTGCTGGATCTATTAGGCGTAGTCTGGGACGTCGTATGGGTAATGAGCTTCACGACTGCGTTTTAGAAGTG
PpuISceKanGibs_Fw	CATTCACCTGAACCCATAGGTCCTGCCTGGCAGAGAATCTTAGGGATAACAGGGTAATCG
PpuISceKanGibs _Rv	AGATTCTCTGCCAGGCAGGACCTATGGGTTCAGGTGAATGGCCAGTGTTACAACCAATTA
Rh159_Fw_recomb	GGGCGGTGGCCGGCACGCCGCATTTCCTAACCCGCGCAGCATGGCCTACAACAGCTTCCT
Rh159_Rv_recomb	CGCGAACGACGTGTGACGAGGACGTGGTTTCCGCAAGCCTTTAGGCGTAGTCTGGGACG
TB_159HAseq_F	AGTGTTTATGAGTCGGGCGG
TB_159HAseq_R	GTGGCGACGTGGATTTCTTG
EGFP-P2A-3XHA	GTCGACTGGATCCGGTACCGAATTCGCCACCATGGTGAGCAAGGGGGAGGAGCTTTTTACTGGCGTTGTTCCCATCCTGGTTGAGTTGGATGGGGATGTCAACGGCCACAAGTTCAGCGTTTCTGGAGAAGGAGAAGGCGACGCAACCTATGGGAAGCTCACTTTGAAATTCATATGCACCACCGGTAAACTCCCAGTCCCCTGGCCTACCCTGGTCACTACCTTGACATATGGAGTTCAGTGCTTTTCTCGATACCCCGACCACATGAAGCAGCACGATTTTTTCAAATCTGCAATGCCCGAGGGCTATGTACAAGAGAGGACCATTTTCTTCAAAGACGACGGCAATTACAAGACCAGGGCCGAGGTCAAATTTGAAGGGGACACTCTTGTGAATCGGATCGAACTTAAAGGGATCGATTTCAAAGAGGACGGAAACATACTGGGGCATAAGCTGGAATACAACTACAATAGCCATAATGTTTACATTATGGCAGACAAGCAGAAAAACGGTATTAAGGTCAACTTCAAGATACGGCATAACATTGAAGATGGGGTCAGTCCAGCTTGCTGACCATTATCAGCAGAATACACCCATTGGCGATGGTCCCGTGCTTTTGCCTGACAACCACTACCTTTCTACCCAAAGCGCATTGTCAAAAGATCCTAACGAGAAGAGAGATCATATGGTCCTTTTGGAGTTCGTAACCGCAGCTGGTATAACTCTGGGTATGGACGAGCTGTATAAGGGCTCAGGGGCTACAAATTTTAGTCTCCTTAAACAAGCCGGAGATGTCGAAGAGAACCCTGGGCCTTACCCATATGACGTTCCTGACTATGCCGGTTACCCCTATGATGTGCCAGATTATGCCGGATCTTATCCTTACGATGTACCTGACTATGCAGGTGGGAGTGGCATGAATTACGTGGGGCAGC
Gibs_Age_159_Fw	CCGTACCACTTCCTACCCTCGTAAACCGGTGCCACCATGGCCTACAACAG
Gibs_Mlu_GFP_Rv	ACCCGGCGCGGAGGCCAGATCTTAACGCGTTTACTTATACAGCTCGTCCATACC
tRFP linker Fw	CCACTTCCTACCCTCGTAAACCGGTGAATTCGCCACCATGAGCGAGCTGAT
tRFP linker Rv	GCGCGGAGGCCAGATCTTAACGCGTGCGGCCGCTCGAGAATTCATCTGTGCCCCAGTTTGC

TABLE S2. Antibodies Used in Immunofluorescence Microscopy

Primary Antibody	Host species	Isotype	Clone	Vendor	Catalog No.	Dilution used
Calnexin	Mouse	mAb IgG1	37	BD Biosciences	610523	1:50
Calnexin	Rabbit	mAb IgG	C5C9	Cell Signaling Tech.	2679	1:50
Syntaxin 6	Rabbit	mAb IgG	C34B2	Cell Signaling Tech.	2869	1:50
SYVN1 (Hrd1)	Rabbit	mAb IgG	D3O2A	Cell Signaling Tech.	14773	1:500
SEL1L	Rabbit	pAb	n/a	Sigma	S3699	1:50
KDEL	Mouse	IgG2a	10C3	Enzo Life Sci., Inc.	ADI-SPA-827-D	1:50
PDI	Rabbit	mAb IgG	C81H6	Cell Signaling Tech.	3501	1:50
Herp (HerpUD1)	Rabbit	pAb IgG	n/a	Proteintech	10813-1-AP	1:50
VCP	Mouse	IgG2a	5	Invitrogen / Life Technologies	MA3-004	1:50
Calreticulin	Rabbit	mAb IgG	D3E6	Cell Signaling Tech.	12238	1:200
Ribophorin 1 (RPN1)	Rabbit	pAb IgG		ThermoFisher / Invitrogen	PA5-65662	1:50
EDEM-1	Mouse	mAb IgM	D-1	Santa Cruz	sc-377394	1:50
GABARAP	Rabbit	mAb	E1J4E	Cell Signaling Tech.	13733	1:200
LC3B	Rabbit	mAb	D11	Cell Signaling Tech.	3868	1:200
HA epitope	Chicken	IgY		Bethyl	A190-106A	1:200
HCMV glycoprotein H (gH)	Mouse	IgG2b	14-4b	Gift of William J. Britt, M.D.	n/a	1: 50
Reticulon-3 (RTN3)	Mouse	IgG2a		ThermoFisher / Invitrogen	MA5-15538	1: 200
HA epitope	Rabbit	pAb		Bethyl	A190-108A	1:1000
Secondary Antibody	Host species	Isotype		Vendor	Catalog No.	Dilution used
Alexa Fluor 488 anti-Mouse IgG (H+L), cross adsorbed	Goat	IgG		ThermoFisher / Invitrogen	A11001	1:1000
Alexa Fluor 488 anti-Rabbit IgG (H+L) Cross-Adsorbed	Goat	IgG		ThermoFisher / Invitrogen	A11008	1:1000
Alexa Fluor 594 anti-Rabbit IgG (H+L) Cross-Adsorbed	Goat	IgG		ThermoFisher / Invitrogen	A11012	1:1000
Alexa Fluor 594 anti-Mouse IgG (H+L) Cross-Adsorbed	Goat	IgG		ThermoFisher / Invitrogen	A11005	1:1000
Alexa Fluor 647 anti-Chicken IgY (H+L)	Goat	IgG		ThermoFisher / Invitrogen	A11039	1:1000

1116
1117
1118

1119
1120
1121   **SUPPLEMENTARY VIDEO FILES.**
1122
1123   **Movie S1: STEM tomography of UL148 ER structures in wildtype HCMV infected**
1124   **fibroblasts at 5 dpi.**
1125
1126   **Movies S2 – S3: Live cell imaging of GFP signal following dox induction of transgene**
1127   **expression in i148^{gfp} and i159^{gfp} ARPE-19 epithelial cells.**
1128   **Movie S2: i148^{gfp} cells from 2-19 h post dox induction.**
1129   **Movie S3: i159^{gfp} cells from 2-19 h post dox induction.**
1130
1131   **Movies S4-S6: FRAP.**
1132   **Movie S4**: **FRAP of UL148-GFP structure (ER within structure).**
1133   **Movie S5: FRAP of UL148 GFP structure (ER outside structure).**
1134   **Movie S6: FRAP of Rh159-GFP.**
1135
1136   **Movie S7: Imaging of UL148-GFP following washout of doxycycline (dox):** live cell imaging
1137   capturing 30 min to 22 h 07 min post washout of dox**.**
1138
1139   **Movies S8-S10: ISR blockade.**
1140   **Movie S8: i148^{gfp} cells in the presence of 0.01% DMSO, from 2-19 h post dox induction.**
1141   **Movie S9: i148^{gfp} cells in the presence of 1.1 µM GSK2606414, from 2-19 h post dox**
1142   **induction.**
1143   **Movie S10: i148^{gfp} cells in the presence of 200 nM ISRIB, from 2-19 h post dox induction.**
1144
1145
1146
1147

REFERENCES.

1148

1149

1.   Li G, Nguyen CC, Ryckman BJ, Britt WJ, Kamil JP. 2015. A viral regulator of glycoprotein
complexes contributes to human cytomegalovirus cell tropism. Proc Natl Acad Sci U S A
112:4471-6.
2.   Wille PT, Knoche AJ, Nelson JA, Jarvis MA, Johnson DC. 2010. A human cytomegalovirus
gO-null mutant fails to incorporate gH/gL into the virion envelope and is unable to enter
fibroblasts and epithelial and endothelial cells. J Virol 84:2585-96.
3.   Jiang XJ, Adler B, Sampaio KL, Digel M, Jahn G, Ettischer N, Stierhof YD, Scrivano L,
Koszinowski U, Mach M, Sinzger C. 2008. UL74 of human cytomegalovirus contributes to
virus release by promoting secondary envelopment of virions. J Virol 82:2802-12.
4.   Zhou M, Lanchy JM, Ryckman BJ. 2015. Human Cytomegalovirus gH/gL/gO Promotes the
Fusion Step of Entry into All Cell Types, whereas gH/gL/UL128-131 Broadens Virus
Tropism through a Distinct Mechanism. J Virol 89:8999-9009.
5.   Wu Y, Prager A, Boos S, Resch M, Brizic I, Mach M, Wildner S, Scrivano L, Adler B. 2017.
Human cytomegalovirus glycoprotein complex gH/gL/gO uses PDGFR-alpha as a key for
entry. PLoS Pathog 13:e1006281.
6.   Kabanova A, Marcandalli J, Zhou T, Bianchi S, Baxa U, Tsybovsky Y, Lilleri D, Silacci-Fregni
C, Foglierini M, Fernandez-Rodriguez BM, Druz A, Zhang B, Geiger R, Pagani M, Sallusto
F, Kwong PD, Corti D, Lanzavecchia A, Perez L. 2016. Platelet-derived growth factor-
alpha receptor is the cellular receptor for human cytomegalovirus gHgLgO trimer. Nat
Microbiol 1:16082.
7.   Wu K, Oberstein A, Wang W, Shenk T. 2018. Role of PDGF receptor-alpha during human
cytomegalovirus entry into fibroblasts. Proc Natl Acad Sci U S A
doi:10.1073/pnas.1806305115.
8.   Nguyen CC, Siddiquey MNA, Zhang H, Li G, Kamil JP. 2018. Human Cytomegalovirus
Tropism Modulator UL148 Interacts with SEL1L, a Cellular Factor That Governs
Endoplasmic Reticulum-Associated Degradation of the Viral Envelope Glycoprotein gO. J
Virol 92:e00688-18.
9.   Wang ECY, Pjechova M, Nightingale K, Vlahava VM, Patel M, Ruckova E, Forbes SK,
Nobre L, Antrobus R, Roberts D, Fielding CA, Seirafian S, Davies J, Murrell I, Lau B, Wilkie
GS, Suarez NM, Stanton RJ, Vojtesek B, Davison A, Lehner PJ, Weekes MP, Wilkinson
GWG, Tomasec P. 2018. Suppression of costimulation by human cytomegalovirus
promotes evasion of cellular immune defenses. Proc Natl Acad Sci U S A 115:4998-5003.
10.   Siddiquey MNA, Zhang H, Nguyen CC, Domma AJ, Kamil JP. 2018. The Human
Cytomegalovirus Endoplasmic Reticulum-Resident Glycoprotein UL148 Activates the
Unfolded Protein Response. J Virol 92:e00896-18.
11.   Sinzger C, Hahn G, Digel M, Katona R, Sampaio KL, Messerle M, Hengel H, Koszinowski U,
Brune W, Adler B. 2008. Cloning and sequencing of a highly productive, endotheliotropic
virus strain derived from human cytomegalovirus TB40/E. J Gen Virol 89:359-68.
12.   Das S, Vasanji A, Pellett PE. 2007. Three-dimensional structure of the human
cytomegalovirus cytoplasmic virion assembly complex includes a reoriented secretory
apparatus. J Virol 81:11861-9.

13.   Sanchez V, Greis KD, Sztul E, Britt WJ. 2000. Accumulation of virion tegument and
envelope proteins in a stable cytoplasmic compartment during human cytomegalovirus
replication: characterization of a potential site of virus assembly. J Virol 74:975-86.
14.   Lilley BN, Ploegh HL. 2005. Multiprotein complexes that link dislocation, ubiquitination,
and extraction of misfolded proteins from the endoplasmic reticulum membrane. Proc
Natl Acad Sci U S A 102:14296-301.
15.   Kamhi-Nesher S, Shenkman M, Tolchinsky S, Fromm SV, Ehrlich R, Lederkremer GZ.
2001. A novel quality control compartment derived from the endoplasmic reticulum.
Mol Biol Cell 12:1711-23.
16.   Leitman J, Shenkman M, Gofman Y, Shtern NO, Ben-Tal N, Hendershot LM, Lederkremer
GZ. 2014. Herp coordinates compartmentalization and recruitment of HRD1 and
misfolded proteins for ERAD. Mol Biol Cell 25:1050-60.
17.   Houck SA, Ren HY, Madden VJ, Bonner JN, Conlin MP, Janovick JA, Conn PM, Cyr DM.
2014. Quality control autophagy degrades soluble ERAD-resistant conformers of the
misfolded membrane protein GnRHR. Mol Cell 54:166-179.
18.   Sturgill ER, Malouli D, Hansen SG, Burwitz BJ, Seo S, Schneider CL, Womack JL, Verweij
MC, Ventura AB, Bhusari A, Jeffries KM, Legasse AW, Axthelm MK, Hudson AW, Sacha
JB, Picker LJ, Fruh K. 2016. Natural Killer Cell Evasion Is Essential for Infection by Rhesus
Cytomegalovirus. PLoS Pathog 12:e1005868.
19.   Chaumorcel M, Souquere S, Pierron G, Codogno P, Esclatine A. 2008. Human
cytomegalovirus controls a new autophagy-dependent cellular antiviral defense
mechanism. Autophagy 4:46-53.
20.   Chaumorcel M, Lussignol M, Mouna L, Cavignac Y, Fahie K, Cotte-Laffitte J, Geballe A,
Brune W, Beau I, Codogno P, Esclatine A. 2012. The human cytomegalovirus protein
TRS1 inhibits autophagy via its interaction with Beclin 1. J Virol 86:2571-84.
21.   Alwine JC. 2012. The human cytomegalovirus assembly compartment: a masterpiece of
viral manipulation of cellular processes that facilitates assembly and egress. PLoS Pathog
8:e1002878.
22.   Das S, Pellett PE. 2011. Spatial relationships between markers for secretory and
endosomal machinery in human cytomegalovirus-infected cells versus those in
uninfected cells. J Virol 85:5864-79.
23.   Leitman J, Ron E, Ogen-Shtern N, Lederkremer GZ. 2013. Compartmentalization of
endoplasmic reticulum quality control and ER-associated degradation factors. DNA Cell
Biol 32:2-7.
24.   Cox DW, Billingsley GD, Callahan JW. 1986. Aggregation of plasma Z type alpha 1-
antitrypsin suggests basic defect for the deficiency. FEBS Lett 205:255-60.
25.   Le A, Ferrell GA, Dishon DS, Le QQ, Sifers RN. 1992. Soluble aggregates of the human PiZ
alpha 1-antitrypsin variant are degraded within the endoplasmic reticulum by a
mechanism sensitive to inhibitors of protein synthesis. J Biol Chem 267:1072-80.
26.   Lomas DA, Evans DL, Finch JT, Carrell RW. 1992. The mechanism of Z alpha 1-antitrypsin
accumulation in the liver. Nature 357:605-7.
27.   Ordonez A, Snapp EL, Tan L, Miranda E, Marciniak SJ, Lomas DA. 2013. Endoplasmic
reticulum polymers impair luminal protein mobility and sensitize to cellular stress in
alpha1-antitrypsin deficiency. Hepatology 57:2049-60.

28.   Ekeowa UI, Freeke J, Miranda E, Gooptu B, Bush MF, Perez J, Teckman J, Robinson CV,
Lomas DA. 2010. Defining the mechanism of polymerization in the serpinopathies. Proc
Natl Acad Sci U S A 107:17146-51.
29.   Suhy DA, Giddings TH, Jr., Kirkegaard K. 2000. Remodeling the endoplasmic reticulum by
poliovirus infection and by individual viral proteins: an autophagy-like origin for virus-
induced vesicles. J Virol 74:8953-65.
30.   Windsor M, Hawes P, Monaghan P, Snapp E, Salas ML, Rodriguez JM, Wileman T. 2012.
Mechanism of collapse of endoplasmic reticulum cisternae during African swine fever
virus infection. Traffic 13:30-42.
31.   Miller S, Krijnse-Locker J. 2008. Modification of intracellular membrane structures for
virus replication. Nat Rev Microbiol 6:363-74.
32.   Siddiquey MNA, Zhang H, Nguyen CC, Domma AJ, Kamil JP. 2018. The human
cytomegalovirus ER resident glycoprotein UL148 activates the unfolded protein
response. J Virol doi:10.1128/JVI.00896-18.
33.   Cormack BP, Valdivia RH, Falkow S. 1996. FACS-optimized mutants of the green
fluorescent protein (GFP). Gene 173:33-8.
34.   Khaminets A, Heinrich T, Mari M, Grumati P, Huebner AK, Akutsu M, Liebmann L, Stolz
A, Nietzsche S, Koch N, Mauthe M, Katona I, Qualmann B, Weis J, Reggiori F, Kurth I,
Hubner CA, Dikic I. 2015. Regulation of endoplasmic reticulum turnover by selective
autophagy. Nature 522:354-8.
35.   Meng L, Mohan R, Kwok BH, Elofsson M, Sin N, Crews CM. 1999. Epoxomicin, a potent
and selective proteasome inhibitor, exhibits in vivo antiinflammatory activity. Proc Natl
Acad Sci U S A 96:10403-8.
36.   Muroi M, Shiragami N, Nagao K, Yamasaki M, Takatsuki A. 1993. Folimycin
(concanamycin A), a specific inhibitor of V-ATPase, blocks intracellular translocation of
the glycoprotein of vesicular stomatitis virus before arrival to the Golgi apparatus. Cell
Struct Funct 18:139-49.
37.   Muroi M, Takasu A, Yamasaki M, Takatsuki A. 1993. Folimycin (concanamycin A), an
inhibitor of V-type H(+)-ATPase, blocks cell-surface expression of virus-envelope
glycoproteins. Biochem Biophys Res Commun 193:999-1005.
38.   Kondratyev M, Avezov E, Shenkman M, Groisman B, Lederkremer GZ. 2007. PERK-
dependent compartmentalization of ERAD and unfolded protein response machineries
during ER stress. Exp Cell Res 313:3395-407.
39.   Harding HP, Novoa I, Zhang Y, Zeng H, Wek R, Schapira M, Ron D. 2000. Regulated
translation initiation controls stress-induced gene expression in mammalian cells. Mol
Cell 6:1099-108.
40.   Palam LR, Baird TD, Wek RC. 2011. Phosphorylation of eIF2 facilitates ribosomal bypass
of an inhibitory upstream ORF to enhance CHOP translation. J Biol Chem 286:10939-49.
41.   Vattem KM, Wek RC. 2004. Reinitiation involving upstream ORFs regulates ATF4 mRNA
translation in mammalian cells. Proc Natl Acad Sci U S A 101:11269-74.
42.   Sidrauski C, Acosta-Alvear D, Khoutorsky A, Vedantham P, Hearn BR, Li H, Gamache K,
Gallagher CM, Ang KK, Wilson C, Okreglak V, Ashkenazi A, Hann B, Nader K, Arkin MR,
Renslo AR, Sonenberg N, Walter P. 2013. Pharmacological brake-release of mRNA
translation enhances cognitive memory. Elife 2:e00498.

43.   Sidrauski C, McGeachy AM, Ingolia NT, Walter P. 2015. The small molecule ISRIB
1280   reverses the effects of eIF2alpha phosphorylation on translation and stress granule
1281   assembly. Elife 4.
1282   44.   Zyryanova AF, Weis F, Faille A, Alard AA, Crespillo-Casado A, Sekine Y, Harding HP, Allen
1283   F, Parts L, Fromont C, Fischer PM, Warren AJ, Ron D. 2018. Binding of ISRIB reveals a
1284   regulatory site in the nucleotide exchange factor eIF2B. Science 359:1533-1536.
1285   45.   Sidrauski C, Tsai JC, Kampmann M, Hearn BR, Vedantham P, Jaishankar P, Sokabe M,
1286   Mendez AS, Newton BW, Tang EL, Verschueren E, Johnson JR, Krogan NJ, Fraser CS,
1287   Weissman JS, Renslo AR, Walter P. 2015. Pharmacological dimerization and activation of
1288   the exchange factor eIF2B antagonizes the integrated stress response. Elife 4:e07314.
1289   46.   Tsai JC, Miller-Vedam LE, Anand AA, Jaishankar P, Nguyen HC, Renslo AR, Frost A, Walter
1290   P. 2018. Structure of the nucleotide exchange factor eIF2B reveals mechanism of
1291   memory-enhancing molecule. Science 359.
1292   47.   Gordiyenko Y, Schmidt C, Jennings MD, Matak-Vinkovic D, Pavitt GD, Robinson CV. 2014.
1293   eIF2B is a decameric guanine nucleotide exchange factor with a gamma2epsilon2
1294   tetrameric core. Nat Commun 5:3902.
1295   48.   Harding HP, Zhang Y, Ron D. 1999. Protein translation and folding are coupled by an
1296   endoplasmic-reticulum-resident kinase. Nature 397:271-4.
1297   49.   Harding HP, Zhang Y, Bertolotti A, Zeng H, Ron D. 2000. Perk is essential for translational
1298   regulation and cell survival during the unfolded protein response. Mol Cell 5:897-904.
1299   50.   Axten JM, Medina JR, Feng Y, Shu A, Romeril SP, Grant SW, Li WH, Heerding DA,
1300   Minthorn E, Mencken T, Atkins C, Liu Q, Rabindran S, Kumar R, Hong X, Goetz A, Stanley
1301   T, Taylor JD, Sigethy SD, Tomberlin GH, Hassell AM, Kahler KM, Shewchuk LM, Gampe
1302   RT. 2012. Discovery of 7-methyl-5-(1-{[3-(trifluoromethyl)phenyl]acetyl}-2,3-dihydro-1H-
1303   indol-5-yl)-7H-p yrrolo[2,3-d]pyrimidin-4-amine (GSK2606414), a potent and selective
1304   first-in-class inhibitor of protein kinase R (PKR)-like endoplasmic reticulum kinase
1305   (PERK). J Med Chem 55:7193-207.
1306   51.   Tenorio R, Fernandez de Castro I, Knowlton JJ, Zamora PF, Lee CH, Mainou BA, Dermody
1307   TS, Risco C. 2018. Reovirus sigmaNS and muNS Proteins Remodel the Endoplasmic
1308   Reticulum to Build Replication Neo-Organelles. MBio 9.
1309   52.   Miller S, Kastner S, Krijnse-Locker J, Buhler S, Bartenschlager R. 2007. The non-structural
1310   protein 4A of dengue virus is an integral membrane protein inducing membrane
1311   alterations in a 2K-regulated manner. J Biol Chem 282:8873-82.
1312   53.   Lamond AI, Sleeman JE. 2003. Nuclear substructure and dynamics. Curr Biol 13:R825-8.
1313   54.   Granell S, Baldini G, Mohammad S, Nicolin V, Narducci P, Storrie B, Baldini G. 2008.
1314   Sequestration of mutated alpha1-antitrypsin into inclusion bodies is a cell-protective
1315   mechanism to maintain endoplasmic reticulum function. Mol Biol Cell 19:572-86.
1316   55.   Lawless MW, Greene CM, Mulgrew A, Taggart CC, O'Neill SJ, McElvaney NG. 2004.
1317   Activation of endoplasmic reticulum-specific stress responses associated with the
1318   conformational disease Z alpha 1-antitrypsin deficiency. J Immunol 172:5722-6.
1319   56.   Tirosh B, Iwakoshi NN, Lilley BN, Lee AH, Glimcher LH, Ploegh HL. 2005. Human
1320   cytomegalovirus protein US11 provokes an unfolded protein response that may
1321   facilitate the degradation of class I major histocompatibility complex products. J Virol
1322   79:2768-79.

57.   Wiertz EJ, Jones TR, Sun L, Bogyo M, Geuze HJ, Ploegh HL. 1996. The human
cytomegalovirus US11 gene product dislocates MHC class I heavy chains from the
endoplasmic reticulum to the cytosol. Cell 84:769-79.

58.   van den Boomen DJ, Timms RT, Grice GL, Stagg HR, Skodt K, Dougan G, Nathan JA,
Lehner PJ. 2014. TMEM129 is a Derlin-1 associated ERAD E3 ligase essential for virus-
induced degradation of MHC-I. Proc Natl Acad Sci U S A 111:11425-30.

59.   Caramelo JJ, Parodi AJ. 2008. Getting in and out from calnexin/calreticulin cycles. J Biol
Chem 283:10221-5.

60.   Hammond C, Braakman I, Helenius A. 1994. Role of N-linked oligosaccharide
recognition, glucose trimming, and calnexin in glycoprotein folding and quality control.
Proc Natl Acad Sci U S A 91:913-7.

61.   Zhou M, Yu Q, Wechsler A, Ryckman BJ. 2013. Comparative analysis of gO isoforms
reveals that strains of human cytomegalovirus differ in the ratio of gH/gL/gO and
gH/gL/UL128-131 in the virion envelope. J Virol 87:9680-90.

62.   Zhang L, Zhou M, Stanton R, Kamil J, Ryckman BJ. 2018. Expression Levels of
Glycoprotein O (gO) Vary between Strains of Human Cytomegalovirus, Influencing the
Assembly of gH/gL Complexes and Virion Infectivity. J Virol 92.

63.   Sifers RN, Brashears-Macatee S, Kidd VJ, Muensch H, Woo SL. 1988. A frameshift
mutation results in a truncated alpha 1-antitrypsin that is retained within the rough
endoplasmic reticulum. J Biol Chem 263:7330-5.

64.   Choudhury P, Liu Y, Bick RJ, Sifers RN. 1997. Intracellular association between UDP-
glucose:glycoprotein glucosyltransferase and an incompletely folded variant of alpha1-
antitrypsin. J Biol Chem 272:13446-51.

65.   Tsao YS, Ivessa NE, Adesnik M, Sabatini DD, Kreibich G. 1992. Carboxy terminally
truncated forms of ribophorin I are degraded in pre-Golgi compartments by a calcium-
dependent process. J Cell Biol 116:57-67.

66.   de Virgilio M, Weninger H, Ivessa NE. 1998. Ubiquitination is required for the retro-
translocation of a short-lived luminal endoplasmic reticulum glycoprotein to the cytosol
for degradation by the proteasome. J Biol Chem 273:9734-43.

67.   Johnston JA, Ward CL, Kopito RR. 1998. Aggresomes: a cellular response to misfolded
proteins. J Cell Biol 143:1883-98.

68.   Johnston JA, Illing ME, Kopito RR. 2002. Cytoplasmic dynein/dynactin mediates the
assembly of aggresomes. Cell Motil Cytoskeleton 53:26-38.

69.   Pankiv S, Alemu EA, Brech A, Bruun JA, Lamark T, Overvatn A, Bjorkoy G, Johansen T.
2010. FYCO1 is a Rab7 effector that binds to LC3 and PI3P to mediate microtubule plus
end-directed vesicle transport. J Cell Biol 188:253-69.

70.   Hummel T, Krukkert K, Roos J, Davis G, Klambt C. 2000. Drosophila Futsch/22C10 is a
MAP1B-like protein required for dendritic and axonal development. Neuron 26:357-70.

71.   Wang H, Bedford FK, Brandon NJ, Moss SJ, Olsen RW. 1999. GABA(A)-receptor-
associated protein links GABA(A) receptors and the cytoskeleton. Nature 397:69-72.

72.   Zhang K, Daigle JG, Cunningham KM, Coyne AN, Ruan K, Grima JC, Bowen KE, Wadhwa
H, Yang P, Rigo F, Taylor JP, Gitler AD, Rothstein JD, Lloyd TE. 2018. Stress Granule
Assembly Disrupts Nucleocytoplasmic Transport. Cell 173:958-971 e17.

73. Li YR, King OD, Shorter J, Gitler AD. 2013. Stress granules as crucibles of ALS
pathogenesis. J Cell Biol 201:361-72.
74. Hetz C, Saxena S. 2017. ER stress and the unfolded protein response in
neurodegeneration. Nat Rev Neurol 13:477-491.
75. Kirchoff V, Wong S, St JS, Pari GS. 2002. Generation of a life-expanded rhesus monkey
fibroblast cell line for the growth of rhesus rhadinovirus (RRV). Arch Virol 147:321-33.
76. Smith IL, Taskintuna I, Rahhal FM, Powell HC, Ai E, Mueller AJ, Spector SA, Freeman WR.
1998. Clinical failure of CMV retinitis with intravitreal cidofovir is associated with
antiviral resistance. Arch Ophthalmol 116:178-85.
77. Murphy E, Yu D, Grimwood J, Schmutz J, Dickson M, Jarvis MA, Hahn G, Nelson JA,
Myers RM, Shenk TE. 2003. Coding potential of laboratory and clinical strains of human
cytomegalovirus. Proc Natl Acad Sci U S A 100:14976-81.
78. Stanton RJ, Baluchova K, Dargan DJ, Cunningham C, Sheehy O, Seirafian S, McSharry BP,
Neale ML, Davies JA, Tomasec P, Davison AJ, Wilkinson GW. 2010. Reconstruction of the
complete human cytomegalovirus genome in a BAC reveals RL13 to be a potent inhibitor
of replication. J Clin Invest 120:3191-208.
79. Hobom U, Brune W, Messerle M, Hahn G, Koszinowski UH. 2000. Fast screening
procedures for random transposon libraries of cloned herpesvirus genomes: mutational
analysis of human cytomegalovirus envelope glycoprotein genes. J Virol 74:7720-9.
80. Chang WL, Barry PA. 2003. Cloning of the full-length rhesus cytomegalovirus genome as
an infectious and self-excisable bacterial artificial chromosome for analysis of viral
pathogenesis. J Virol 77:5073-83.
81. Tischer BK, Smith GA, Osterrieder N. 2010. En passant mutagenesis: a two step
markerless red recombination system. Methods Mol Biol 634:421-30.
82. Tischer BK, von Einem J, Kaufer B, Osterrieder N. 2006. Two-step red-mediated
recombination for versatile high-efficiency markerless DNA manipulation in Escherichia
coli. Biotechniques 40:191-7.
83. Dietz AN, Villinger C, Becker S, Frick M, von Einem J. 2018. A Tyrosine-Based Trafficking
Motif of the Tegument Protein pUL71 Is Crucial for Human Cytomegalovirus Secondary
Envelopment. J Virol 92.
84. Hohn K, Sailer M, Wang L, Lorenz M, Schneider ME, Walther P. 2011. Preparation of
cryofixed cells for improved 3D ultrastructure with scanning transmission electron
tomography. Histochem Cell Biol 135:1-9.
85. Villinger C, Schauflinger M, Gregorius H, Kranz C, Hohn K, Nafeey S, Walther P. 2014.
Three-dimensional imaging of adherent cells using FIB/SEM and STEM. Methods Mol
Biol 1117:617-38.
86. Kremer JR, Mastronarde DN, McIntosh JR. 1996. Computer visualization of three-
dimensional image data using IMOD. J Struct Biol 116:71-6.
87. Britt WJ, Jarvis MA, Drummond DD, Mach M. 2005. Antigenic domain 1 is required for
oligomerization of human cytomegalovirus glycoprotein B. J Virol 79:4066-79.
1406
1407