1   **cRacle: R Tools for Estimating Climate from Vegetation[1]**

3   Robert S. Harbert[2,3,4], Alex A. Baryiames[2]

5 [2]Department of Biology, Stonehill College, 320 Washington St., North Easton, MA 02357

6 [3]American Museum of Natural History, 79[th] St & Central Park West, New York, NY 10024

8 Email Address: RH: rharbert@stonehill.edu

10 Number of words:

16 *Acknowledgements –*

17 The authors would like to thank Stonehill College for their support of undergraduate research

18 that made this study possible.

20   ABSTRACT

Premise of the study – The Climate Reconstruction Analysis using Coexistence Likelihood

Estimation (CRACLE) method for the estimation of climate from vegetation is a robust set of

modeling tools for estimating climate and paleoclimate that makes use of large repositories of

biodiversity data and open-access R software.

Methods — Here we implement a new R library for the estimation of climate from

vegetation.The 'cRacle' library implements functions for data access, aggregation, and models to

estimate climate from plant community composition. 'cRacle' is modular and features many

best-practice features.

Results – Performance tests using modern vegetation survey data from North and South America

shows that CRACLE outperforms alternative methods. CRACLE estimates of mean annual

temperature (MAT) are usually within 1°C of the actual when optimal model parameters are

used. Generalized Boosted Regression (GBR) model correction is also shown here to improve on

CRACLE models by reducing bias.

Discussion – CRACLE provides accurate estimates of climate from modern plant communities.

Non-parametric CRACLE modeling coupled to GBR model correction produces the most accurate

results to date. The 'cRacle' R library streamlines the estimation of climate from plant

community data, and will make this modeling more accessible to a wider range of users.

Key words: CRACLE, paleoclimate, R, ENM, GBIF, fossil, climate.

INTRODUCTION

'cRacle' is an open-source R software package that performs the Climate Reconstruction

Analysis using Coexistence Likelihood Estimation (CRACLE) model for the estimation of climate

from biological community data (Harbert & Nixon, 2015). CRACLE performs an empirical

likelihood estimation of climatic parameters given a set of taxa (species, genera, or families) and

their modern distributions via point-locality data from the Global Biodiversity Information Facility

(GBIF) or similar primary biodiversity databases (e.g., iNaturalist, iDigBio, BIEN).

The primary application of CRACLE and related methods is in the reconstruction of

paleoclimate from fossil floras. All current applications of CRACLE make assumptions about

climatic niche stability through time because the rely on the Nearest Living Relative (NLR)

approach to infer elements of fossil taxon niche dimensions by observing the niche space

occupied by a closely related extant relative (Mosbrugger and Utescher, 1997; Utescher, 2014;

Harris *et al*. 2014). The NLR assumptions will be most valid when the fossil taxa can be placed as

members of extant species and least accurate when fossils are members of extinct lineages. For

this reason, NLR and CRACLE will produce the most reliable reconstructions of relatively recent

(e.g., Pleistocene and Pliocene) paleoclimate, but may produce reasonable results for much

older fossil assemblages where the plant fossil taxa can be placed reliably in extant groups with

well sampled modern distributions (e.g., Harris *et al.* 2014).

CRACLE modeling was recently applied to the estimation of paleoclimate using Late

Quaternary (<50ka – *kilo anuum*) plant macrofossils from packrat (*Neotoma* spp.) middens in

Western North America (Harbert, Nixon 2018). Analysis of the packrat midden paleoclimate

estimates revealed a history of rapid climate change during and just after the Late Pleistocene

deglaciation, followed by Holocene warm and dry periods of 1-2°C above the modern (1950-

2000) averages (Harbert and Nixon, 2018)

Community assemblages from excavated permafrost at five Pliocene sites in the

Canadian Arctic Archipelago analyzed with CRACLE (Fletcher *et al.*, 2017). This work revealed

that the Canadian arctic was up to 22°C warmer than today during the Early Pliocene (~3.6Ma –

mega anuum) and supported many cool-temperate plant taxa. However, Fletcher *et al.* (2017)

carefully compared results derived from CRACLE estimates using species only identifications vs.

generic level identifications and suggest that care must be taken when interpreting CRACLE

results when generic level identifications are used to avoid being misled by model imprecision.

Modified bioclimatic envelope method that are closely related to CRACLE were recently

used to reconstruct the Paleocene-Eocene Thermal Maximum (~55Ma) and the Eocene Thermal

Maximum (53.5Ma) by way of plant macrofossils from the Green River Basin in southwestern

Wyoming and plant microfossils found in Arctic marine sediment cores (Hyland et al., 2018;

Willard et al., 2019).

CRACLE models have been applied to non-plant communities as well. Pliocene Arctic

climate estimates based on fossil beetle communities using CRACLE, revealed notable biases in

beetle based climate reconstructions relative to plant proxy and stable isotope methods

(Fletcher *et al.*, 2018).

Despite CRACLE existing for the first four years as only a single R script published as a

supplement to the original paper (Harbert and Nixon, 2015) the method has proven to be a

useful tool to the community (presently 16 Google Scholar citations), yielding informative results

for several fossil systems. Here, we present the 'cRacle' R package. 'cRacle' is designed to

streamline the process of CRACLE modeling, data acquisition and pre-processing of primary

biodiversity data (from GBIF and other related databases), and CRACLE result visualization. The

'cRacle' R package is built from modular functions that perform segments of the modeling

protocol that allow for easy to read R scripts as well as future modular developments of related

methods.

For comparison with the CRACLE methods, we have also implemented the related

Thompson's Mutual Climatic Range (MCR; Thompson et al., 2012). We include both the

weighted and unweighted MCR methods as representatives of the many available MCR or

Coexistence Approach (CA) range-intersect methods (Mosbrugger and Utescher, 1997; Sinka and

Atkinson, 1999; Utescher et al., 2014) for the estimation of climate from vegetation (ECV) as part

of the 'cRacle' R package. The 'cRacle' package is distributed under the open MIT License and all

code is available on GitHub (https://www.github.com/rsh249/cRacle).

                                                     METHODS

Full documentation of 'cRacle' R code is available with the package code

(https://github.com/rsh249/cRacle). Below is an outline of the major functionality of the 'cRacle'

package and explanation of many of the core functions available.

'cRacle' Package Functions – <u>Data Acquisition</u> – Downloading primary biodiversity data

from GBIF (www.gbif.org), iNaturalist (www.inaturalist.org), and BISON (www.bison.usgs.gov) are

supported in 'cRacle' through the functions *gbif_get(), get_bison()*, and *inat()*. Each of these

functions accepts arguments for a single taxon (genus, species, or family) to query and a

maximum number of records to return. Data pre-processing, cleaning for outliers, spatial bias

reduction, and climate data extraction for a set of georeferenced occurrence data is done

through the *extraction()* function. Data downloading for multiple taxa and preprocessing

functionality can be done in one step using the *getextr()* function, which is implemented with

parallel computing capability to reduce user wait time.

<u>Likelihood Modeling</u> – The primary modeling that CRACLE requires is the calculation of

probability density functions using both parametric (Gaussian Normal) and non-parametric

(Kernel Density) estimates. These calculations can be performed with 'cRacle' directly from the

output of *extraction()* using the function(s) *dens_obj()* and *densform()* for multiple and single

taxa respectively. Note that *dens_obj()* is simply a wrapper function for *densform()* designed to

make multi-species likelihood model building simpler. Both parametric and non-parametric

likelihood functions are always calculated and stored in the object produced by *densform()* and

dens_obj(). This way all calculations are done in tandem for a dataset and results for both

methods can be examined at the end of the CRACLE process.

The likelihood modeling carried out in the *densform()* function includes several options

that can be manipulated by the user. The Kernel Density Estimation (KDE) employed by CRACLE

can now be implemented with several kernel estimators including the standard Gaussian,

Epanechnikov, cosine, optcosine, biweight, rectangular, and triangular kernels based on the R

'stats' package function *density()* (R Core Team, 2018). The kernel bandwidth can be optimized

using standard criteria via the argument 'bw' in the *densform()* and *dens_obj()* functions.

Likelihoods can be calculated based on a standard or conditional probability via the 'manip'

argument where conditional probabilities take into account the background distribution of each

climate parameter using a set of randomly sampled points from within a normalized distance

from each primary occurrence location or from a user defined set of climate data. Finally, the

resulting likelihood distributions can be trimmed to empirical ranges or confidence intervals to

avoid unintended extrapolation using the 'clip' argument.

CRACLE Joint Likelihood – After likelihood functions are estimated the next key CRACLE

step is to calculate first the joint likelihood and then the maximum of the joint likelihood

distribution. 'cRacle' implements the function *and_fun()* for calculating the joint likelihood and

or_fun() for creating unions between taxa for merging likelihood features between groups (i.e.,

between species in a clade). The output of *and_fun()* is a set of joint likelihood features for all

parameters, and this can be summarized by the function *get_optim()* which finds the optimal

climate values for each parameter. The output of *get_optim()* provides the user with CRACLE

results.

Example Pseudocode – The general CRACLE modeling flow will use the following

functions in order: *getextr() -> dens_obj()-> and_fun()-> get_optim();* to produce climate

estimates from a list of taxa. This code would: download data and extract climate data, estimate

likelihood distributions, calculate the joint likelihood, and find the optimum of the joint likelihood

for each climate variable analyzed.

Visualizing Likelihood and Joint Likelihood Functions – 'cRacle' provides a set of functions

for visualizing the likelihood functions as standard probability density distributions. These

functions are *densplot()* for plotting single distribution objects (i.e., for one taxon/climate

parameter pair), *addplot()* for adding to an existing *densplot()* figure, and *multiplot()* for plotting

distributions for multiple taxa given a single climate parameter. For example, a user may plot the

distributions for all taxa in their study for Mean Annual Temperature using *multiplot()* and the

output from *dens_obj()*, and then add the joint likelihood function using *addplot()* and the model

object output from *and_fun().*

Issue Tracking and Versioning – The repository at https://github.com/rsh249/cRacle.git

will maintain the current development version of 'cRacle'. Future major releases will be

submitted to CRAN. Issues with running 'cRacle' code are accepted on the Git repository's

'Issues' feature at https://github.com/rsh249/cRacle/issues or by email with the corresponding

author of this publication.

Data access – Primary biodiversity data, occurrence coordinates for your target taxa, and

climate data are required for CRACLE modeling. We recommend primary biodiversity data

downloaded from the Global Biodiversity Information Facility (GBIF) and support data access

through 'cRacle' functions. Climate data must be in the form of a raster object readable by the R

'raster' package (Hijmans, 2018). Climate data from the WorldClim project (Hijmans et al., 2005;

Fick and Hijmans, 2017) are recommended, but users may also use raster data downloaded from

other sources such as CHELSA (Karger et al., 2017), and ENVIREM (Title and Bemmels, 2018).

Estimation Climate from Vegetation – (ECV) – As originally described CRACLE consists of

two methods for likelihood estimation: parametric and non-parametric probability functions.

'cRacle' implements these two approaches in tandem for each step of the likelihood analysis,

and has implemented options about whether to calculate the likelihoods from standard

probability or conditional probability. For comparison we have implemented calculations for the

Mutual Climatic Range (MCR) method for ECV described by Thompson *et al.* (2012), though the

unweighted MCR described is analogous to the Coexistence Approach (CA) described elsewhere

(Mosbrugger and Utescher, 1997)

CRACLE (Harbert and Nixon, 2015) defines both a parametric and non-parametric

probability density function for each taxon – climate parameter pair. CRACLE estimates

likelihood for a set of taxa ($t_{1:i}$). The CRACLE Parametric –Normal – probabilities are estimated as:

$$f(x|t) = \frac{1}{\sqrt{2\sigma^2\pi}} e^{\frac{(x-\mu)^2}{2\sigma^2}}$$

179   Whereas the CRACLE Non-Parametric – Kernel Density Estimation (KDE) – probabilities are

180   estimated as:

$$181 \qquad f(x|t) = \frac{1}{n}\sum_{n=1}^{i} K_h(\bar{x} - x_i)$$

182   Where K is a Kernel function of properties XXXX, and 'h' is the kernel bandwidth, a smoothing

183   parameter with value > 0. We recommend using either the "optcosine" or "gaussian" kernels

184   and a bandwidth calculated using Silverman's Rule of Thumb (Silverman, 1986). 'cRacle' allows

185   for the calculation of probability density functions conditioned by the probability density of a

186   random background sample for each climate parameter defined as:

$$187 \qquad f(c) = \frac{1}{n}\sum_{n=1}^{i} K_h(\bar{c} - c_i)$$

188   Where 'c' is the set of background climate given by a sample of points from within the study

189   area. The conditional probability for a taxon (t) is defined as:

$$190 \qquad f(t|x) = \frac{f(x|t)}{f(c)}$$

192   For the estimation of climate from vegetation CRACLE modeling calculates the joint likelihood for

193   multiple taxon probability functions as:

$$194 \qquad L(x|t_{1:i}) = \sum_{n=1}^{i} \ln[f(x|t_n)]$$

195   Where the maximum value of $L(x|\ t_{1:i})$ corresponds to the value of the climate parameter 'x' that

196   is most likely to lead to the coexistence of that set of taxa ($t_{1:i}$).

198 Note that 'cRacle' also implements the calculation of a weighted mean by variance that can be

199 substituted for the parametric CRACLE functions. Weighted means ($\bar{x}$) is calculated given a set of

200 taxon means ($x_{1:i}$) and standard deviations ($\sigma_{1:i}$):

201
$$\bar{x} = \frac{\sum_{i=1}^{n}(x_i\,\sigma_i^{-2})}{\sum_{i=1}^{n}\sigma_i^{-2}}$$

Modern Validation – To test the performance of CRACLE that can be expected for

paleoclimate modeling we set two experimental analyses to test aspects of the CRACLE modeling

process.

First, we sample iNaturalist data (www.inaturalist.org) by a grid search of North America

for plant species coexisting in 0.1x0.1 degree bounded search areas using the 'rinat' R package

(Barve and Hart, 2017). Then we build CRACLE estimates for Mean Annual Temperature using

the WorldClim 2.0 dataset at 2.5 arcminutes resolution (Fick and Hijmans, 2017) for parametric

and non-parametric CRACLE as well as MCR and MCR un (unweighted) methods. Primary

biodiversity data were obtained from GBIF (www.gbif.org) using the 'cRacle' *get_dist_all()*

function and the GBIF RESTful JSON based API. Model overfitting for the iNaturalist dataset was

tested by spatially thinning the raw GBIF data using the 'cRacle' *extraction()* function for factors

of 2,4,6,8, and 10 times the dimensions of the 2.5 arcminute climate raster cells. The 'cRacle'

thinning procedure is an imperfect, but efficient, spatial thinning method that is consistent with

best-practices in Ecological Niche Modeling literature (Aiello-Lammens *et al.* 2015). CRACLE and

MCR model output for each thinning level were summarized and reported for general guidance.

The R code for the iNaturalist grid search test is available at

https://github.com/rsh249/cracle_examples/inat_grid_search.

Second, the Botanical Information and Ecology Network (BIEN 4.0) database was queried

for published vegetation plot surveys from North and South America using the 'rbien' R package

(Maitner *et al.* 2017). For plot surveys with >5 plant taxa with distribution data available from

GBIF we estimated climate parameters for all 19 bioclimatic variables from the WorldClim 2.0

data (Fick and Hijmans, 2017) using parametric CRACLE methods and spatial thinning to a factor

of 2 times the 2.5 arcminute climate grid. Error in CRACLE are non-random (Harbert and Nixon,

2015), therefore non-linear regression modeling may be able to account for and correct

common CRACLE errors. CRACLE results for 70,391 plot surveys were then partitioned 50:50 into

training and testing sets by random sampling with replacement and Generalized Boosted

Regression (GBR) models were fit to the training data for each of the 19 climate parameters

using the R 'gbm' package (Greenwell *et al.* 2019). The GBR models made adjusted climate

estimates using the CRACLE results in the test set to determine whether error patterns in

CRACLE can be predicted by GBR and therefore may be corrected for in independent analyses.

The R code for the iNaturalist grid search is available:

https://github.com/rsh249/cracle_examples/cracle_bien.

236   RESULTS

Modern Validation – iNaturalist - The grid search of iNaturalist in North America yielded

climate estimations by CRACLE (both parametric and non-parametric) and the Mutual Climatic

Range (MCR) unweighted and weighted approaches for 285 sites at spatial thinning factors of 0,

2, 4, 6, 8, and 10 times the 2.5 arcminute climate raster. A total of 1,204,925 unique records

were accessed from GBIF for this analysis for a total of 1541 species. The sampled localities

ranged in mean annual temperature (MAT) from 0.4-26.1°C. The top performing (lowest error

rates) estimates were the CRACLE non-parametric ('kde' in Figure 1) and the MCR weighted

('mcr.w' in Figure 1) methods. Mean errors for CRACLE were ~1°C and median errors were

generally less than the means. Spatial thinning did not change mean or median error rates for

any of the tested methods (Fig. 1B,C).

Figure 1. iNaturalist Grid Search Climate Estimation Results. A) Geographic search area (shading) with successful climate estimation locations marked (black squares). B) Mean Anomaly Rates, and C) Median Anomaly Rates for CRACLE ('kde' non-parametric, and 'means' parametric) and

MCR ('uw' unweighted and 'w' weighted) results for factor levels of 0 ('raw'), 2, 4, 6, 8, and 10

times the 2.5 arcminute climate raster.

BIEN – Non-parametric CRACLE estimates of all 19 WorldClim Bioclim variables

(www.worldclim.org/bioclim) from the WorldClim 2.0 dataset (Fick and Hijmans, 2017) were

generated for 70,391 vegetation plot surveys accessed through the BIEN 4.0 database using a

total of 2,560,261 georeferenced specimen records from GBIF.

Mean absolute anomaly rates for temperature parameters were between 1-2°C with

Pearson's Correlation coefficients greater than 0.9 in the best scenarios (Table 1), but up to 5°C

and correlation coefficients <0.7 for estimates of mean temperature during the wettest and

driest quarters of the year (BIO8 and BIO9). The mean absolute anomaly rates for mean annual

precipitation (MAP – BIO12) was 179mm with a correlation of 0.79 (Table 1).

Generalized Boosted Regression (GBR) models were used to correct CRACLE estimates

using independent random samples without replacement of 50% of the data for training and

testing datasets. GBR correction improved CRACLE performance in the test set by 20-50% (Table

1, 2; Figure 2). CRACLE+GBM corrected anomalies are smaller and more symmetrically

distributed around '0' than the non-corrected CRACLE results (Figure 2D), and CRACLE+GBM

estimates are more strongly correlated with the WorldClim data than the raw CRACLE results

(Table 1, 2).

Figure 2. CRACLE estimates and GBR error correction demonstration for BIEN vegetation plot
surveys and representative climate parameters. BIEN vegetation plot surveys from 70,391 unique

272 localities (A) were analyzed with CRACLE (B) to estimate the 19 Bioclim variables (showing BIO1,

273 5, 6, 12, 17, and 18 here as top performing examples) from the WorldClim2.0 dataset (Fick and

274 Hijmans, 2017). Generalized Boosted Regression error correction modeling was trained and

275 tested (C) on independent subsets of plot data. GBR correction yielded overall reduced error

276 rates and less biased estimates in many cases (D).

277   DISCUSSION

The modern validation analyses conducted here should provide a good baseline for

expected CRACLE performance going forward. More thorough testing in future studies is

certainly welcome, but most modeling choices are should be made by the user to reflect any

unique properties of their study system. The results we provide here indicate that the best

estimates that are generated from the 'cRacle' software are the CRACLE non-parametric

estimates with spatially thinned data, these yielding mean errors of approximately 0.5°C less

than the Mutual Climatic Range (MCR) method, an analogous method to the widely used

Coexistence Approach (Mosbrugger and Utescher, 1997). Furthermore, CRACLE results

presented here compare favorably, though indirectly, to recent applications of the Weighted

Average – Partial Least Squares (WA-PLS) modeling that is common for analysis of fossil pollen

samples that uses modern analog communities to build proxy models (Montade *et al.*, 2019). A

direct comparison of CRACLE and WA-PLS methods is necessary future work.

Through our analysis of BIEN hosted vegetation plot surveys we show that CRACLE can

produce accurate climate estimates for a variety of both temperature and precipitation

parameters (Figure 2), though some parameters are better predicted than others (Table 1).

Notably, BIO8 (Mean Temperature of the Warmest Quarter), BIO9 (Mean Temperature of the

Coldest Quarter), BIO13 (Precipitation of the Wettest Month), and BIO15 (Precipitation

Seasonality – Coeff. Var.) are relatively poorly predicted by CRACLE, likely due to less direct

impact of those variables on plant distributions. Whereas estimates of BIO1 (MAT), BIO5 (Max

Temperature), BIO6 (Min Temperature), and BIO18 (Precipitation of the Driest Quarter) yield the

highest correlation with known values (Table 1). We also show that CRACLE model correction via

Generalized Boosted Regression (GBR) can lower error rates from the CRACLE baseline (Figure 2;

Table 2). We expect that further regression model fitting on expanded datasets may yield further

refinement, however, the GBR model correction is implemented in 'cRacle' *get_optim()* as an

option for users to correct non-parameteric CRACLE estimates for the WorldClim 2.0 Bioclim

variables only.

Resources and Tutorials – We are actively developing resources and tutorials in support

of the 'cRacle' R package. Demonstration code and short projects will be maintained on a

continuing basis at https://github.com/rsh249/cracle_examples. We aim to provide a series of

web tutorials for various CRACLE modeling tasks here to guide beginner users through the

process. Tutorials, and issue tracking are distributed through the main 'cRacle' repository:

https://github.com/rsh249/cRacle.git.

Conclusion – 'cRacle' is a new and actively maintained resource for climate estimation

from biological community composition. These estimates are particularly relevant to the study of

fossil systems where often the best indication of past climate is the community of plant and

animal fossils. We show that users of 'cRacle' should expect estimates that are accurate (e.g.,

within 1°C for MAT) when applying best-practices.

316   AUTHOR CONTRIBUTIONS

AAB conceived of and implemented the iNaturalist grid search and CRACLE analysis and provided

valuable testing, benchmarking, and feedback data on the 'cRacle' R library. RSH designed the

BIEN vegetation plots CRACLE study and implemented the Generalized Boosted Regression

model correction analysis. RSH designed and wrote the 'cRacle' R package code.

LITERATURE CITED

Aiello-Lammens, M.E., R.A. Boria, A. Radosavljevic, B. Vilela, and R.P. Anderson. 2015. spThin: an

R package for spatial thinning of species occurrence records for use in ecological niche models.

Ecography, *38*(5): 541-545.

Barve, V., and E. Hart. 2017. 'rinat': Access iNaturalist Data Through APIs. R package version

0.1.5. https://cran.r-project.org/package=rinat

Fick, S.E. and R.J. Hijmans. 2017. WorldClim 2: new 1-km spatial resolution climate surfaces for

global land areas. *International journal of climatology* 37(12):4302-4315.

Fletcher, T., R. Feng, A.M. Telka, J.V. Matthews Jr, and A. Ballantyne. 2017. Floral Dissimilarity

and the Influence of Climate in the Pliocene High Arctic: Biotic and Abiotic Influences on Five

Sites on the Canadian Arctic Archipelago. *Frontiers in Ecology and Evolution* 5:19.

Fletcher, T.L., A.Z. Csank, and A.P. Ballantyne. 2018. Identifying bias in cold season temperature

reconstructions by beetle mutual climatic range methods in the Pliocene Canadian High Arctic.

Palaeogeography, Palaeoclimatology, Palaeoecology 514: 672-676.

Greenwell, B., B. Boehmke, J. Cunningham, and GBM Developers. 2019. 'gbm': Generalized

Boosted Regression Models. R package version 2.1.5 https://cran.r-project.org/package=gbm

Harbert, R.S. and K.C. Nixon. 2015. Climate reconstruction analysis using coexistence likelihood

estimation (CRACLE): A method for the estimation of climate using vegetation. *American Journal*

of Botany 102(8): 1277-1289.

HARBERT, R.S. and NIXON, K.C., 2018. Quantitative Late Quaternary Climate Reconstruction from

Plant Macrofossil Communities in Western North America. *Open Quaternary* 4(1):8.

Harris, A.J., Papeş, M., GAO, Y.D. and Watson, L., 2014. Estimating paleoenvironments using

ecological niche models of nearest living relatives: A case study of Eocene Aesculus L. *Journal of*

Systematics and Evolution, *52*(1): 16-34.

Hijmans, R.J., S.E. Cameron, J.L. Parra, P.G. Jones, A. Jarvis. 2005. Very high resolution

interpolated climate surfaces for global land areas. *International Journal of Climatology: A*

Journal of the Royal Meteorological Society 25(15):1965-1978.

Hijmans, R.J. (2018). raster: Geographic Data Analysis and Modeling. R package version 2.8-4.

https://CRAN.R-project.org/package=raster

Hyland, E.G., K.W. Huntington, N.D. Sheldon, and T. Reichgelt. 2018. Temperature seasonality in

the North American continental interior during the Early Eocene Climatic Optimum. *Climate of*

the Past 14(10): 1391-1404.

Karger, D.N., O. Conrad, J. Böhner, T. Kawohl, H. Kreft, R.W. Soria-Auza, N.E. Zimmermann, H.P.

Linder, and M. Kessler. 2017. Climatologies at high resolution for the earth's land surface areas.

Scientific Data 4:170122.

Maitner, B.S., B. Boyle, N. Casler, R. Condit, J. Donoghue, S.M. Durán, D. Guaderrama, C.E.

Hinchliff, P.M. Jørgensen, N.J. Kraft, and B. McGill. 2018. The bien r package: A tool to access the

Botanical Information and Ecology Network (BIEN) database. *Methods in Ecology and Evolution*

9(2):373-379

Montade, V., O. Peyron, C. Favier, J.P Francois, and S.G. Haberle. 2019. A pollen–climate

calibration from western Patagonia for palaeoclimatic reconstructions. *Journal of Quaternary*

Science 34(1): 76-86.

Mosbrugger, V. and T. Utescher. 1997. The coexistence approach—a method for quantitative

reconstructions of Tertiary terrestrial palaeoclimate data using plant fossils. *Palaeogeography,*

Palaeoclimatology, Palaeoecology 134(1-4):61-86.

R Core Team (2018). R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Silverman, BW. 1986. Density estimation for statistics and data analysis. Routledge. DOI:

https://doi.org/10.1201/9781315140919

Thompson, R.S., K.H. Anderson, R.T. Pelltier, L.E. Strickland, P.J. Bartlein, and S.L. Shafer. 2012.

Quantitative estimation of climatic parameters from vegetation data in North America by the

mutual climatic range technique. *Quaternary Science Reviews* 51:18-39.

Title, P.O. and J.B. Bemmels. 2018. ENVIREM: an expanded set of bioclimatic and topographic

variables increases flexibility and improves performance of ecological niche modeling. *Ecography*

41(2):291-307.

Utescher, T., A.A. Bruch, B. Erdei, L. François, D. Ivanov, F.M.B. Jacques, A.K. Kern, V.

Mosbrugger, and R.A. Spicer. 2014. The Coexistence Approach—theoretical background and

practical considerations of using plant fossils for climate quantification. *Palaeogeography,*

Palaeoclimatology, Palaeoecology, 410: 58-73.

Willard, D.A., T.H. Donders, T. Reichgelt, D.R. Greenwood, F. Sangiorgi, F. Peterse, K.G. Nierop, j.

Frieling, S. Schouten, and A. Sluijs. 2019. Arctic vegetation, temperature, and hydrology during

Early Eocene transient global warming events. *Global and Planetary Change* 178:139-152.

Zaborac-Reed, S.J. and E.B. Leopold. 2016. Determining the paleoclimate and elevation of the

late Eocene Florissant flora: support from the coexistence approach. *Canadian Journal of Earth*

Sciences 53(6): 565-573.

TABLES

Table 1: CRACLE performance statistics for 19 Bioclim parameters estimated for BIEN vegetation plot data (n=35,195 plots)								
Climate_Parameter	Mean Anomaly	Mean Absolute Anomaly	Median Absolute Anomaly	RMSE	NRMSE	Pearson's *r*	Spearman's *r*	Units
wc2.0_bio_2.5m_01	0.34	1.40	1.09	1.86	0.05	0.95	0.96	°C
wc2.0_bio_2.5m_02	-0.31	1.01	0.74	1.39	0.07	0.82	0.79	°C
wc2.0_bio_2.5m_03	-0.25	2.51	1.80	3.44	0.04	0.87	0.87	°C
wc2.0_bio_2.5m_04	-23.93	64.64	42.35	92.68	0.06	0.84	0.85	°C
wc2.0_bio_2.5m_05	-0.19	1.71	1.19	2.32	0.07	0.92	0.95	°C
wc2.0_bio_2.5m_06	0.64	2.13	1.57	2.93	0.05	0.91	0.92	°C
wc2.0_bio_2.5m_07	-0.94	2.42	1.97	3.19	0.07	0.79	0.83	°C
wc2.0_bio_2.5m_08	1.00	5.60	3.27	7.81	0.18	0.69	0.63	°C
wc2.0_bio_2.5m_09	-1.00	5.64	2.43	8.80	0.16	0.60	0.63	°C
wc2.0_bio_2.5m_10	0.03	1.30	0.97	1.74	0.06	0.94	0.96	°C
wc2.0_bio_2.5m_11	0.31	1.89	1.36	2.66	0.05	0.92	0.93	°C
wc2.0_bio_2.5m_12	26.20	179.17	111.69	275.09	0.06	0.79	0.83	mm/year
wc2.0_bio_2.5m_13	7.32	25.04	11.31	47.92	0.07	0.70	0.75	mm/month
wc2.0_bio_2.5m_14	9.89	15.49	9.72	21.79	0.10	0.77	0.76	mm/month
wc2.0_bio_2.5m_15	-7.11	11.66	8.30	16.12	0.15	0.69	0.71	--
wc2.0_bio_2.5m_16	21.84	68.96	31.48	135.63	0.08	0.69	0.76	mm/3*month
wc2.0_bio_2.5m_17	26.80	45.61	27.49	64.92	0.08	0.80	0.76	mm/3*month
wc2.0_bio_2.5m_18	-6.20	34.76	23.16	52.69	0.05	0.90	0.91	mm/3*month
wc2.0_bio_2.5m_19	34.97	99.35	73.22	142.96	0.10	0.57	0.69	mm/3*month

Table 2: GBR Corrected CRACLE performance statistics for 19 Bioclim parameters estimated for BIEN vegetation plot data (n=35,195 plots)								
Climate_Parameter	Mean Anomaly	Mean Absolute Anomaly	Median Absolute Anomaly	RMSE	NRMSE	Pearson's *r*	Spearman's *r*	Units
wc2.0_bio_2.5m_01	-0.06	1.11	0.85	1.51	0.04	0.96	0.96	°C
wc2.0_bio_2.5m_02	0.00	0.84	0.60	1.18	0.06	0.86	0.81	°C
wc2.0_bio_2.5m_03	-0.01	2.08	1.48	2.89	0.04	0.91	0.89	°C
wc2.0_bio_2.5m_04	1.03	51.65	36.94	72.73	0.05	0.90	0.87	°C*100
wc2.0_bio_2.5m_05	-0.11	1.18	0.83	1.64	0.05	0.96	0.95	°C
wc2.0_bio_2.5m_06	-0.16	1.75	1.18	2.49	0.04	0.93	0.93	°C
wc2.0_bio_2.5m_07	-0.03	1.85	1.33	2.57	0.06	0.85	0.86	°C
wc2.0_bio_2.5m_08	-0.38	4.43	3.00	6.17	0.15	0.74	0.70	°C
wc2.0_bio_2.5m_09	0.41	4.86	2.38	7.66	0.14	0.71	0.69	°C
wc2.0_bio_2.5m_10	-0.06	0.97	0.73	1.32	0.05	0.97	0.96	°C
wc2.0_bio_2.5m_11	-0.15	1.61	1.14	2.32	0.04	0.94	0.94	°C
wc2.0_bio_2.5m_12	12.84	147.83	95.59	225.75	0.05	0.86	0.86	mm/year
wc2.0_bio_2.5m_13	1.52	19.52	10.52	35.44	0.05	0.78	0.82	mm/month

wc2.0_bio_2.5m_14	0.04	10.83	7.50	15.50	0.08	0.85	0.86	mm/month
wc2.0_bio_2.5m_15	2.11	7.56	4.77	11.17	0.11	0.75	0.73	--
wc2.0_bio_2.5m_16	3.25	51.70	28.72	97.37	0.06	0.79	0.83	mm/3*month
wc2.0_bio_2.5m_17	-0.80	32.85	22.45	46.69	0.07	0.87	0.87	mm/3*month
wc2.0_bio_2.5m_18	2.48	28.98	21.01	42.83	0.04	0.93	0.92	mm/3*month
wc2.0_bio_2.5m_19	11.46	70.76	46.68	114.25	0.08	0.71	0.76	mm/3*month

 Figure Legends

Figure 1. iNaturalist Grid Search Climate Estimation Results. A) Geographic search area (shading)

with successful climate estimation locations marked (black squares). B) Mean Anomaly Rates,

and C) Median Anomaly Rates for CRACLE ('kde' non-parametric, and 'means' parametric) and

MCR ('uw' unweighted and 'w' weighted) results for factor levels of 0 ('raw'), 2, 4, 6, 8, and 10

times the 2.5 arcminute climate raster.

Figure 2. CRACLE estimates and GBR error correction demonstration for BIEN vegetation plot

surveys and representative climate parameters. BIEN vegetation plot surveys from 70,391 unique

localities (A) were analyzed with CRACLE (B) to estimate the 19 Bioclim variables (showing BIO1,

5, 6, 12, 17, and 18 here as top performing examples) from the WorldClim2.0 dataset (Fick and

Hijmans, 2017). Generalized Boosted Regression error correction modeling was trained and

tested (C) on independent subsets of plot data. GBR correction yielded overall reduced error

rates and less biased estimates in many cases (D).