Microbial evolution reshapes soil carbon feedbacks to climate change

Elsa Abs[1,2]*, Scott R. Saleska[3], Regis Ferriere[2,3,4]

[1] Department of Ecology and Evolutionary Biology, University of California, Irvine, USA

[2] International Center for Interdisciplinary Global Environmental Studies (iGLOBES),

CNRS, Ecole Normale Supérieure, Paris Sciences & Lettres University,

University of Arizona, Tucson, AZ 85721, USA

[3] Department of Ecology & Evolutionary Biology, University of Arizona,

Tucson, AZ 85721, USA

[4] Institut de Biologie (IBENS), Ecole Normale Supérieure,

Paris Sciences & Lettres University, CNRS, INSERM, 75005 Paris, France

*Corresponding author, eabs@uci.edu

Abstract

Microbial decomposition of soil organic matter is a key component of the global carbon cycle. As Earth's climate changes, the response of microbes and microbial enzymes to rising temperatures will largely determine the soil carbon feedback to atmospheric CO2. However, while increasing attention focuses on physiological and ecological mechanisms of microbial responses, the role of evolutionary adaptation has been little studied. To address this gap, we developed an ecosystem-evolutionary model of a soil microbe-enzyme system under warming. Constraining the model with observations from five contrasting sites reveals evolutionary aggravation of soil carbon loss to be the most likely outcome; however, temperature-dependent increases in mortality could cause an evolutionary buffering effect instead. We generally predict a strong latitudinal pattern, from small evolutionary effects at low latitude to large evolutionary effects at high latitudes. Accounting for evolutionary mechanisms will likely be critical for improving projections of Earth system responses to climate change.

Microorganisms are key drivers of global biogeochemical cycles[1]. In terrestrial ecosystems,

soil microbes decompose organic matter, returning carbon to the atmosphere as carbon

dioxide (CO_2)[2]. *In vitro* and *in situ* experiments suggest that changes in microbial

decomposition with warming are an important feedback to climate[3–5]. Soil microbial

populations may respond to increasing temperature through physiological mechanisms such

as individual metabolic adjustment[6,7] and ecological mechanisms such as shifts in

population abundance or community composition[8,9]. Given the short generation time, large

population sizes and standing genetic variation of many microbial organisms, evolutionary

adaptive responses of microbial populations to warming are also likely[10,11]. However, how

microbial evolutionary adaptation may contribute to carbon-climate feedbacks is

unknown[12].

Key to microbial decomposition of soil organic matter is the production by microbes of

extracellular enzymes (exoenzymes), that diffuse locally in the soil and bind to soil organic

matter compounds[13]. Because the fitness cost of exoenzyme production[14] (reduced

allocation to growth, Fig. 1a) is paid by individual microbes whereas fitness benefits (larger

resource pool) are received by microbial collectives[15], we expect genetic variation in

exoenzyme production[16] to be under strong selection[15,17]. Our objective is to evaluate how

exoenzyme production responds to selection under environmental warming, and how the

evolutionary response of exoenzyme production impacts the response of soil organic

carbon stock (SOC).

To this end, we develop and analyze a novel ecosystem-evolutionary model, starting

with an ecosystem model of microbe-enzyme decomposition first proposed by Allison *et al.*

(2010)[18] (Fig. 1a), and then modified to take microbial evolutionary adaptation into

account. Amongst ecosystem models of soil microbial decomposition (reviewed in Abs and

Ferriere (2019)), Allison *et al.*[18]'s model is the simplest of the *CDMZ* type, where *C*

denotes a single pool of soil organic carbon, *D*, dissolved organic carbon, *M*, the biomass of

the microbial community, and *Z*, a single pool of exoenzymes. The focal microbial trait is

the fraction of assimilated carbon allocated to exoenzyme production[19–21], or 'exoenzyme

allocation fraction', whose mean value across microbial strains or species is denoted by φ.

The balance of assimilated carbon, $1 - \varphi$, is allocated to microbial growth. In the absence

of evolution, the enzyme allocation fraction, φ, represents the mean of an unchanging

distribution of enzyme allocation behaviors across a community of microbes. With natural

selection acting on the trait's genetic variation across the community, φ evolves as the

microbial community adapts to the environmental conditions, e.g. mean soil temperature.

Our ecosystem-evolutionary model predicts the evolutionarily stable value, φ^*, as a

function of temperature, and how the response of φ^* to warming affects the decomposition

rate and SOC stock (C at equilibrium).

By comparing the response of the SOC stock that our ecosystem-evolutionary model

predicts (EVO response) to the response predicted by the ecosystem *CDMZ* model in the

absence of evolution, assuming φ to be a fixed parameter (ECOS response), we can

evaluate the contribution of microbial evolutionary adaptation (EVO effect) to the direction

and magnitude of the SOC stock response to climate warming (Fig. 1b, c). To illustrate

how EVO effects may vary in real ecosystems, we use available data[22] on the

decomposition kinetic parameters in five sites of increasing latitude and decreasing mean

annual temperature. We evaluate ECOS and EVO responses for each site, and compare

them within and among sites. Our analysis identifies parameters and temperature

dependencies that critically influence the strength of evolutionary effects. We discuss how

these evolutionary effects relate to previous consideration of 'adaptation' in microbial

community responses to climate warming[18,23,24], and how evolutionary adaptation may

interact with ecological responses such as species sorting and community shifts[25–27]. We

conclude by highlighting how our results could inform future empirical work.

Model overview

We use the microbe-enzyme model of litter decomposition first introduced in ref. 18[18] to

describe the ecosystem dynamics of soil organic carbon (C), dissolved organic carbon (D),

microbial biomass (M), and extracellular enzyme abundance (Z), given litter input, leaching

rates, and soil temperature (Fig. 1a and Supplementary Figure 1). The effect of temperature

is mediated by enzymes kinetics, with exoenzymes driving the decomposition rate, and

intra-cellular enzymes involved in resource uptake and microbial biomass synthesis. The

model parameters (microbial life history parameters, thermal dependencies, enzyme

parameters, carbon inputs) were constrained by experimental and observational data[18,20,22]

(see Abs and Ferriere 2019 for a review). Within these parameter ranges, the model outputs

are consistent with target empirical values (target C of the order of 100 mg cm^{-3}, M about

2% of C, Z about 1% of M, and limiting D close to zero). Zhang et $al.$[28] provided some

more direct validation by successfully fitting a $CDMZ$ model to time series of field

measurements of soil respiration from a specific ecosystem (semiarid savannah subject to

episodic rainfall pulses).

In general, as temperature increases, the baseline ecosystem model without evolution

predicts[18] a decline in equilibrium SOC due to faster enzyme kinetics, a positive feedback

to warming. We call this the ECOS (for ECOSystem-only) response of the system. To

investigate the effect of evolutionary adaptation on decomposition, we include microbial

evolution in the ecosystem model. Our focus is on soil bacteria (as opposed to fungi),

which typically have large population size and short generation time. DOC uptaken by

individual cells is allocated to exoenzyme production (fraction φ) or microbial biomass,

with enzyme production efficiency denoted by γ_Z and microbial growth efficiency (MGE)

denoted by γ_M. We assume that microbes may vary individually in their exoenzyme

allocation fraction and that some of this variation has a genetic basis. We then derive the

selection gradient on the enzyme allocation fraction φ and compute the evolutionarily stable

value, φ^*. Changing temperature alters the selection gradient, hence φ^*. Knowing how φ^*

changes as temperature increases, we can evaluate how the ecosystem equilibrium changes

from both the direct effect of warming on enzyme kinetics, and the indirect effect mediated

by microbial evolutionary adaptation to warming (Fig. 1b, c).

At the molecular and cellular level, the effect of warming on microbial decomposition

is mediated by the temperature sensitivity of intra- and extra-cellular enzymatic

activity[22,29,30]. In our baseline 'kinetics-only' scenario of temperature-dependent

decomposition, we assume that microbial uptake parameters (maximum uptake rate and

half-saturation constant) and exoenzyme kinetics parameters (maximum decomposition rate

and half-saturation constant) increase with temperature[5,31] in a logistic manner. We

consider two additional scenarios for the influence of temperature on decomposition. In the

microbial mortality scenario, the microbial death rate also increases with temperature[32].

This could be due to a higher risk of predation or pathogenic infection at higher

temperatures, or faster microbial senescence due to higher protein turnover[32]. In the

microbial growth efficiency (MGE) scenario, MGE (the fraction of carbon allocated to

growth that actually contributes to microbial biomass, as opposed to being released as CO_2

via growth respiration) decreases with temperature[18,24], which could be due to higher

maintenance costs at higher temperature[33].

Results

Evolution of the enzyme allocation fraction trait. The adapted exoenzyme allocation

fraction, φ^*, corresponds to a maximum of microbial fitness relative to other trait values,

and therefore depends on the parameters defining microbial net growth rate: MGE,

maximum uptake rate, local competitive advantage to exoenzyme producers, or

'competition asymmetry', and mortality. From our model, φ^* at temperature T is derived

(methods) as:

(1) $$\varphi^*(T) = 1 - \frac{d_M}{\gamma_M\, v_{max}^U} - \frac{1}{c_0}$$

where γ_M denotes the MGE; v_{max}^U, the maximum uptake rate; d_M, the death rate, and c_0, the

degree of competition asymmetry. The latter quantifies the accessibility of the exoenzyme

'public good' to microbes; it measures the differential availability of enzymatically

produced dissolved organic carbon (DOC) to different microbial strains. Competition

asymmetry is shaped by diffusion of exoenzymes and DOC, and by microbial mobility, and

is thus likely influenced by soil physical properties, such as texture or moisture. For

simplicity, we assume that competition asymmetry is independent of temperature.

At a fixed temperature, the model predicts that microbes invest less in exoenzymes

when exposed to hostile conditions, which may translate into high mortality (d_M), low

MGE (γ_M), low maximal uptake rate (v_{max}^U), and/or low competitive advantage to

producers (c_0). Conversely, microbes are selected to allocate more resource into producing

exoenzymes when more favourable growth conditions allow them to obtain a better 'return

on investment'[34].

According to equation (1), the adaptive response of exoenzyme allocation to warming

depends on how the MGE, maximum uptake, and mortality vary with temperature. We

investigate how parameters influence the direction and sensitivity of the adaptive enzyme

allocation fraction to temperature by calculating the derivative of φ^* with respect to T, for

each scenario of temperature dependence. In the baseline 'kinetics-only' scenario (in which

v^U_{max} is the only variable in (1) that depends on temperature) the response of φ^* to

temperature variation is given by

(2) $\dfrac{d\varphi^*}{dT} = \dfrac{E^U_v \, d_M}{\gamma_M R (T+273)^2 \, v^U_{max}(T)}$

According to equation (2), increasing temperature creates more favorable growth

conditions for microbes through higher resource uptake capacity (higher v^U_{max}), thus

microbes evolve higher investment in exoenzyme production (positive $d\varphi^*/dT$)

(Supplementary Figure 5a), resulting in lower equilibrium SOC (an evolutionary

enhancement of the positive feedback to warming seen in the baseline ECOS response).

Because φ^* is proportional to the product of the exponential of $1/T$ and to $1/T^2$, its

sensitivity to temperature is strongest across low temperatures (equation (2),

Supplementary Figure 5a). The adaptive response is generally predicted to be strongest

(and the enhancement of the positive climate feedback greatest) when warming improves

microbial growth potential (by increasing v^U_{max}) under initially hostile conditions (high d_M,

low γ_M, low v^U_{max} due to low intrinsic uptake rate v^U_0 and/or high activation energy E^U_v, low

initial temperature T_0) (equation (2)).

In the temperature-dependent mortality scenario, both mortality and uptake respond

exponentially to temperature. Microbes evolve a higher investment in exoenzyme

production if the change in resource uptake capacity remains greater than the change in

mortality rate; otherwise, microbes evolve a lower investment (Supplementary Note 3 and

Supplementary Figure 5b-c). In the MGE-temperature dependent scenario, the response of

φ^* to temperature variation is proportional to the inverse of an exponential function of

temperature (through the uptake rate, v^U_{max}) times a linear function of temperature[32]

(through MGE). As a consequence, at low initial temperature, the adaptive response of the

allocation strategy to warming is mainly driven by the thermal dependence of resource

uptake, and microbes adapt by increasing their resource investment in exoenzyme

production. For ecosystems that are initially warmer, the microbial adaptive response to

warming is mainly driven by the reduction of MGE, which leads to lower allocation to

exoenzymes (Supplementary Note 3 and Supplementary Figure 5d).

There are two cases where the evolutionary enhancement of positive feedbacks to

warming (due to higher enzyme kinetic rates) could be reversed, leading to lower enzyme

production and less feedback to warming: the case of temperature sensitive mortality, and

the case of temperature-dependent MGE in an initially warm ecosystem (Supplementary

Fig. 5c, d). In all other cases, evolutionary adaptation to warming leads to larger resource

allocation to exoenzyme production and a stronger positive feedback, with a greater

response in initially colder ecosystems.

Comparing the ECOS and the EVO responses. The evolutionary response of SOC

equilibrium to warming combines the non-evolutionary response of the ecosystem and the

evolutionary adaptation of enzyme production (Fig. 1c). In all three scenarios of

temperature dependence, the SOC non-evolutionary equilibrium generally decreases as

temperature or resource allocation to exoenzymes increases (Fig. 1b, Supplementary Figure

4, Supplementary Note 5). The negative effect of temperature results from the SOC

equilibrium being mostly sensitive to the maximum decomposition rate (v_{max}^D)

(Supplementary Note 2); all other temperature-dependent parameters have a lesser

influence on the SOC equilibrium across their range of variation with temperature. As the

maximum decomposition rate increases with warming, the SOC equilibrium decreases. The

SOC equilibrium is always lower in systems where microbes invest more resources in

exoenzymes, because all other parameters being fixed, a larger enzyme allocation fraction

entails that more exoenzymes are produced per unit time, which leads to more SOC

decomposed per unit time (Supplementary Note 1). We therefore predict that, without

evolution, the equilibrium soil carbon stock shrinks as the climate warms, in all scenarios

of temperature-dependence (Fig. 3). The strongest losses occur in initially cold systems due

to the non-linear response of the maximum decomposition rate (v_{max}^D) and therefore of

SOC equilibrium to temperature (Fig. 3, Supplementary Figure 4). We therefore expect

evolution to alter the non-evolutionary response according to the scenario, from

aggravating soil carbon loss in systems where microbes adapt to warming with higher

exoenzyme production, to buffering carbon loss from soils where microbes adaptively

respond to warming with a lower exoenzyme allocation fraction (Fig. 1c).

As expected, the simulated evolutionary response of SOC equilibrium to warming

(Figs. 2, 3) mirrors the adaptive response of the enzyme allocation fraction to warming in

all scenarios of temperature dependence (Supplementary Figure 5a-d). In the baseline

scenario, we showed that microbes always evolve a higher enzyme allocation fraction in

response to rising temperature, and we predict that evolution should amplify the non-

evolutionary soil carbon loss due to warming (Fig. 3a). The effect of evolution is strongest

in ecosystems characterized by conditions that are hostile to microbial growth (low initial

temperature, T_0; high mortality, d_M; low MGE, γ_M; low maximum uptake rate, v_0^U) (Fig. 2,

Fig. 3a, Supplementary Figure 6), where the adaptive response of the enzyme allocation

fraction (change in φ^*, equation (2), Supplementary Figure 5a) has been shown to be the

most pronounced. Strong evolutionary effects are robust to the other model parameters –

enzyme parameters (efficiency, production) and environmental parameters (litter input,

leaching) (Supplementary Figures 6 and 7, Supplementary Note 4), which have no

influence on the relationship between φ^* and temperature (equations (1) and (2)). This

relationship, however, cannot explain why stronger evolutionary effects are predicted when

the differential accessibility to resources between microbial strains is small (low c_0)

(equation (2), Fig. 2c-d). In this case, low competition asymmetry selects for microbes

allocating little to exoenzymes (low φ^*), shaping systems that are more sensitive to change

in enzyme allocation fraction due to the non-linear response of SOC equilibrium to φ

(Supplementary Note 1, Supplementary Figure 3).

In the temperature-dependent mortality scenario, the strength of the effect of

temperature on mortality is an important determinant of the adaptive response to warming.

When mortality is moderately sensitive to temperature, the positive response of φ^* to

warming is attenuated. As a result, the evolutionary aggravation of soil carbon loss is less

severe (Fig. 3b). When mortality is strongly sensitive to temperature, warming creates more

hostile conditions for microbial growth, therefore φ^* decreases with warming. Evolutionary

adaptation then buffers the loss of soil carbon (negative EVO effect, Fig. 3c).

In the temperature-dependent MGE scenario, the direction (positive or negative) of the

response of φ^* to warming is determined by the initial temperature T_0. The evolutionary

effect parallels the response of φ^*. At low T_0, φ^* increases strongly with temperature,

causing an aggravation of soil carbon loss (positive evolutionary effect). In contrast, at high

T_0, φ^* decreases strongly with warming, thus opposing the non-evolutionary response

(negative evolutionary effect) and promoting carbon sequestration instead (Fig. 3d). At

intermediate T_0, φ^* is barely sensitive to temperature, and the effect of evolutionary

adaptation to warming is negligible.

Our general analysis of the ecosystem *CDMZ* model has shown that non-evolutionary

response of equilibrium SOC to warming always involves soil carbon loss and may only

vary in amplitude. In contrast, the adaptive response of microbial enzyme allocation, which

is shaped by an interaction among the non-linear temperature dependences of microbial

traits, can drive negative as well as positive responses of equilibrium soil carbon to

warming. Adaptive evolution generally amplifies soil carbon loss. However, in ecosystems

where soil microbial mortality increases with temperature, this general evolutionary effect

can be reversed; adaptive evolution may then reduce soil carbon loss, or even cause carbon

sequestration.

Model predictions using empirical data from five sites. To illustrate how evolutionary

effects may vary in real ecosystems, we used available data[22] on the decomposition kinetic

parameters from five sites of increasing latitude and decreasing mean annual temperature

(Costa Rica, California, West Virginia, Maine, and Alaska, Fig. 4). We evaluated non-

evolutionary and evolutionary responses for each site under three levels of competition

asymmetry (as quantified by the local competitive advantage to producers, c_0) (Fig. 4a-h).

Under our baseline scenario, EVO effects correlate strongly with mean annual temperature,

even more so for low competition asymmetry (Fig. 4i). Stronger EVO effects occur in

colder sites, as found in the general analysis (Fig. 3a). In contrast, the non-evolutionary

response does not correlate with mean annual temperature (Fig. 4a). As a result, a

temperate site such as Maine exhibits a weak non-evolutionary response that can be
strongly amplified by evolution, whereas the warm Costa Rica site shows a strong non-
evolutionary response that is little affected by evolution.
These results are quantitatively attenuated but qualitatively unaffected when microbial
mortality increases moderately with temperature (Fig. 4b, f, j). With a stronger effect of
temperature on microbial mortality, all sites show the evolutionary buffering effect (Fig.
4k) found in the general analysis (Fig. 3c). The intensity of evolutionary buffering is
independent of the sites' mean annual temperature, whereas it varies significantly with
competition asymmetry (Fig. 4k). Under the temperature-dependent MGE scenario, non-
evolutionary and evolutionary responses are reduced in magnitude compared to the baseline
scenario (Fig. 4d, h), particularly in cold sites. However, in these sites, EVO effects are
enhanced dramatically (Fig. 4l). Thus, in a site as cold as Alaska, a significant evolutionary
loss of soil carbon is predicted, whereas the non-evolutionary-driven loss of soil carbon
would be negligible (Fig. 4l).
286
Discussion
As global warming increases environmental temperatures, our model predicts evolution of
the enzyme allocation fraction, with potentially large effects on the decomposition process
and SOC stock (Fig. 1). The size of evolutionary effects is most sensitive to MGE,
microbial mortality, activation energy of uptake maximal rate, competition asymmetry,
initial temperature (Fig. 2), and to the traits' temperature sensitivity (Fig. 3). Implications
of our findings for large geographic scales across terrestrial ecosystems, are revealed by
specifying the model for the five contrasting sites for which exoenzyme kinetics data are
available[22]. We find evolutionary aggravation of soil carbon loss to be the most likely
outcome, with a strong latitudinal pattern, from small evolutionary effects at low latitude to
large evolutionary effects at high latitudes (Fig. 4). Strong temperature-dependence of
microbial mortality, however, would dramatically change the evolutionary pattern, possibly
causing an attenuation of soil carbon loss or even carbon sequestration in response to
warming.
301

These modeling results are broadly consistent with empirical work showing that changes in

soil C stocks are driven by changes in microbial enzyme activity [35,36], and that these

changes in activity arise from the multiple effects of temperature on enzyme kinetics,

microbial pool size, and microbial allocation to exoenzymes[21,37]. Measured mass-specific

potential enzyme activity, used as a proxy for allocation to exoenzymes, was shown to

generally increase with warming, a pattern also predicted by our evolutionary model[21]. In

particular, Steinweg *et al.*[21] found that the rise in enzyme allocation was greatest for

moderate warming and less pronounced for strong warming, which matches our model

predictions under the scenario of temperature-dependent MGE. A large body of empirical

observations and controlled experiments at various time and spatial scales highlights a

general response of soil respiration and soil C stocks to warming, involving an ephemeral

increase of respiration, no significant change in SOC, and a decrease in microbial biomass.

Only the temperature-dependent mortality scenario with evolution could match these

patterns. It has been argued, however, that most experiments remain insufficient to rule out

model predictions that depart from these patterns[38].

317   The mechanism (physiological, ecological and/or genetic changes) by which

enzyme allocation varies in these experiments is unknown; future research should address

the role that evolutionary processes may have played in shaping these responses. It is

important, in both experimental and modeling studies, to distinguish among different

potential biological mechanisms. Next, we consider the distinct roles of physiological

acclimation (phenotypic plasticity within individuals) and ecological community assembly

(shifts in species composition in whole communities) versus evolutionary adaptation.

Acclimation responses, whereby individuals' physiological mechanisms buffer the kinetics

effect of warming and maintain homeostasis in key life-history traits such as growth and

maintenance, are sometimes referred to as 'adaptation' (e.g. of microbial carbon use

efficiency in response to warming)[18,23,24]. However, this is phenotypic plasticity at the

individual level, rather than genetic adaptation at the population level. In our model,

constant mortality (in the baseline and temperature-dependent MGE scenarios) and constant

MGE (in the baseline and temperature-dependent mortality scenarios) can be interpreted as

expressions of such phenotypic plasticity[18,32]. Our results thus show that plasticity, whereby

individual cells buffer growth efficiency and/or mortality against temperature variation,

does not necessarily impede or overwhelm evolutionary adaptation to climate warming. It

is precisely under the assumptions that MGE and mortality remain constant with respect to

temperature that the strongest adaptive change in exoenzyme allocation is predicted.

The enzyme allocation fraction itself might be plastic. Indeed, Steinweg *et al.[21]*

interpreted the positive effect of temperature on allocation to enzyme production as a cell-

level response to larger nutrient needs driven by higher maintenance costs. In our model,

warming causes larger rates of nutrient uptakes (higher v_{max}^U) and this can lead to an

adaptive increase of the enzyme allocation fraction (equation (1)). From the general

evolutionary theory of phenotypic plasticity[39], we expect the evolutionary optimal reaction

norm of enzyme allocation fraction to follow the same pattern, i.e. φ increasing with

warming, provided that the cost of plasticity is not too high and does not alter too markedly

the selection gradient on φ.

An interesting way of testing the role of evolutionary adaptation *vs.* plasticity would be

to monitor the effect of warming on experimentally evolving bacterial communities at

different levels of medium diffusivity or porosity[40]. Our model predicts that the diffusion of

resources (DOC), that is likely dependent on physical quality of the soil medium, has a

strong influence on the adaptive evolution of microbial exoenzyme production in response

to warming, but not on the ecosystem (non-evolutionary) response driven by enzyme

kinetics and physiological plasticity. Thus, comparing population-scale decomposition and

respiration across treatments that factorially cross temperature and medium quality may

contribute to disentangle the effects of adaptation *vs.* plasticity.

An even greater challenge is to tease apart evolutionary adaptation and ecological responses

such as species sorting or community shifts in species or functional group abundances, and

assess their relative importance in the response of microbial communities to environmental

change[25–27]. On the theoretical side, evolutionary and non-evolutionary trait-based models

make fundamentally different predictions[41]. Evolutionary models address the dynamics of

trait distributions driven in trait space by natural selection and genetic variation. In contrast,

ecological community models focus on the dynamics of a given set of species (with possible additions of potentially very different ones via dispersal[42]). In ecological community models, warming may result in a shift of community composition, but only within the set of initially assembled species. As a consequence, different (initial) assemblages may lead to different responses[43], and the stochasticity of community assembly may therefore contribute to variation in functional responses[44]. In contrast, adaptive evolution fueled by genetic variation (due to *de novo* mutations and/or mobile genetic elements[45]) drives a gradual exploration of the community trait space and convergence towards evolutionarily attractors ('fitness peaks'). This is expected to yield more predictable community responses, because evolving communities are less constrained by the (historically contingent) set of species and corresponding traits that initially assembled.

On the empirical side, there is growing experimental evidence for soil bacterial communities shifting, with functional consequences on decomposition, in response to climate[43]. On the other hand, evolution experiments on *Pseudomonas* bacteria showed that local adaptation was as important as community composition in shaping the community response to elevated temperature over the course of a two-month experiment[46]. Our model could be extended to disentangle the role of evolutionary adaptation vs. ecological shifts in response to warming. To represent the functional diversity of a microbial community exploiting a diversity of substrates, different SOC pools could be included[47], alongside specifying different types of enzymes and different microbial functional groups that produce them[20]. In our model parameterization, microbial functional groups would potentially differ in traits such as MGE (γ_M), enzyme cost (γ_Z) and uptake rate (v_{max}^U)[20], and assuming heritable variation, the evolution of these traits would drive the adaptive response of the microbial functional community to environmental change. In an evolution experiment using *Neurospora discreta* as a model system to assess adaptation of soil fungi to warming[48], Allison *et al.*[49] did find evidence for evolutionary responses in MGE and uptake rates. By letting multiple traits evolve, our model extended to multiple substrates could be used first to generate an evolutionarily stable community at a given temperature (following Sauterey *et al.*'s[50] approach for including evolutionary dynamics in models of

ocean planktonic communities), and then evaluate both ecological and evolutionary

responses to warming. Ideally, mechanistic models of the kind studied here, without and

with genetic variation in exoenzyme allocation and other adaptive traits, could be fitted to

experimental data and compared in an inferential model selection framework[51]. Microbial

systems offer much promise to achieve such a level of model-data integration. The trait-

based approach for microbial communities could facilitate model validation by leveraging

genomic and metagenomic data mapped to soil microbial function[52].

In sum, this work shows that evolutionary feedbacks, operating alongside

physiological and ecological responses, may profoundly alter ecosystem function, revealing

a critical need for evidence of the role of evolutionary change in microbial communities

facing environmental change. Rather surprisingly, studies that investigate how rapid

evolution in communities affect ecosystem function are dominated by studies of large

organisms, e.g. fish[53]. Rapid evolutionary adaptation of microbial populations to

environmental change, especially temperature gradients[54], is well documented in laboratory

systems, but there is limited evidence for the role of positive selection and rapid evolution

in the response of natural microbial communities to environmental change[25]. In the

laboratory, evolution experiments on single-strain bacterial cultures exposed to a

temperature gradient could be analysed by sequencing genes involved in the synthesis of

carbon-mineralization enzymes and reconstructing time series of allelic frequency spectra[55].

In such experiments, concurrent measurements of respiration, enzyme activity and

microbial biomass could be correlated to putative changes in allele frequencies. A few

controlled evolution studies in soil[56,57], including the resident community, exist for bacteria

and viruses, showing patterns of reciprocal (antagonistic) evolution of resistance and

infectivity over a matter of weeks. For more complex microbial communities, studies of the

human gut are paving the way, suggesting significant short-term evolution occurring in

natural microbial populations that are embedded in larger microbial communities[58]. Thus,

there is mounting evidence that the diversity and structural complexity of microbial

communities does not offset their evolutionary potential to respond to environmental

change.

Given the large population size and short generation time of many microorganisms,
microbial evolution is likely an essential component of ecosystem response to warming.
Microbial evolutionary adaptation to warming, and its impact on the decomposition of soil
organic matter, can radically change soil carbon dynamics. Overall, we expect evolutionary
effects to vary greatly among ecosystems that differ in biotic (microbial life history and
physiology) and abiotic (temperature, soil texture and moisture) characteristics. Empirical
data suggest that natural values of enzyme allocation fraction are low[30,34] and fall in the
range for which our model predicts large evolutionary responses of decomposition to
warming. Predicting geographic variation in evolutionary responses and effects across large
geographic scales calls for more empirical data on variation in ecosystem (competition)
traits, especially how the competitive advantage to exoenzyme producers varies with soil
physical properties, and in microbial physiological traits, especially microbial mortality.
Both the general analysis and the numerical application to specific sites underscore the
critical effect of the shapes of the traits' temperature dependencies. Our work highlights the
need empirically to probe the thermal dependence of enzymatic and microbial
physiological traits, especially mortality, across wide enough temperature ranges.
In spite of an increasing effort to document and understand the ecosystem impact of
microbial physiological and ecological responses to climate warming[18,24,59], no Earth
system model that seeks to represent the role of living organisms in climate feedbacks has
yet included evolutionary mechanisms of adaptation. Our model is a critical first step. Next
steps will involve accounting for soil carbon stabilization on a timescale longer than
respiration[60]; for vegetation types, the associated diversity of organic substrates in litter,
and corresponding diversity of microbial decomposers[20]; and for the interaction of
biogeochemical cycles and associated stoichiometric constraints[19,20,34,61,62]. As
demonstrated by successful Earth-scale modeling of phytoplankton abundance and
distribution in the global ocean[63], future models that take these new steps will trade off their
added structural complexity with the 'self-parameterization' that the processes of trait
heritable variation and natural selection drive in the models themselves. As this research
program unfolds, we expect projections of future climate and carbon cycle feedbacks, and

their uncertainty, to be significantly impacted by adaptive evolution, from local to global

scales.

Methods

Ecosystem model. Based on ref. 18[18] (Fig. 1a, Supplementary Fig. 1), the ecosystem model
has four state variables measured in unit mass of carbon: soil (non decomposed) organic
carbon (SOC), C; soil decomposed soluble organic carbon (DOC), D; microbial biomass,
M; and exoenzyme concentration, Z. Exoenzyme production drives the decomposition
process of C into D, which is the only source of carbon for microbes. The model accounts
for microbial production and death, exoenzyme decay, recycling of dead microbes and
degraded exoenzymes, C input from plant litter, and leaching of C and D.

Model equations. State variables C, D, M, Z obey equations (1a-d):

(1a) $\quad \dfrac{dC}{dt} = I - \dfrac{v_{max}^{D}\,C}{K_m^{D}+C} Z - e_C C$

(1b) $\quad \dfrac{dD}{dt} = \dfrac{v_{max}^{D}\,C}{K_m^{D}+C} Z + d_M M + d_Z Z - \dfrac{v_{max}^{U}\,D}{K_m^{U}+D} M - e_D D$

(1c) $\quad \dfrac{dM}{dt} = (1 - \varphi)\gamma_M \dfrac{v_{max}^{U}\,D}{K_m^{U}+D} M - d_M M$

(1d) $\quad \dfrac{dZ}{dt} = \varphi\, \gamma_Z \dfrac{v_{max}^{U}\,D}{K_m^{U}+D} M - d_Z Z$

In equation (1a), decomposition follows from Michaelis-Menten kinetics of Z binding
substrate C; there is a constant input, I, of soil organic (non decomposed) carbon from
aboveground litter, and a loss due to leaching at constant rate e_C. In equation (1b), D is
produced by decomposition and the recycling of dead microbial biomass and inactive
enzymes; D is consumed by microbial uptake, and lost by leaching at constant rate e_D. In
equation (1c), growth of microbial biomass M is driven by the rate of D uptake (a Monod
function of D) times the fraction of uptaken D turned into biomass, $(1 - \varphi)\,\gamma_M$, minus
microbial mortality at constant rate d_M. In equation (1d), enzyme variation is driven by the
rate of D uptake times the fraction allocated to enzyme production, φ, and production
efficiency, γ_Z, minus enzyme deactivation at constant rate, d_Z.

Ecosystem equilibria. The ecosystem model possesses either one globally stable

equilibrium, or three equilibria (one of which is always unstable) (Supplementary Fig. 2).

There are thresholds $\varphi_{\min}$ and $\varphi_{\max}$ such that the single globally stable equilibrium exists for

$\varphi < \varphi_{\min}$ or $\varphi > \varphi_{\max}$ and is given by $C = I/e_C$, $D = 0$, $M = 0$, $Z = 0$. Thus, at this equilibrium,

the microbial population is extinct and no decomposition occurs. For $\varphi_{\min} < \varphi < \varphi_{\max}$, the

microbial population can either go extinct (then the system stabilizes at the same

equilibrium as before) or persists at or around a non-trivial equilibrium, which can be

solved for analytically (equation (S1)). Note that $\varphi_{\min}$ and $\varphi_{\max}$ depend on all microbial and

model parameters (Supplementary Figure 3, Supplementary Note 1).

Effect of temperature on model parameters. Decomposition is predicted to respond to

warming[5] due to the temperature sensitivity of enzymatic activity[22,29,64]. Microbial

assimilation may also vary with temperature if the microbial membrane proteins involved

in nutrient uptake are sensitive to warming. Following ref. 18[18], we assume that exoenzyme

kinetics parameters (maximum decomposition rate v_{max}^D and half-saturation constant K_m^D)

and microbial uptake parameters (maximum uptake rate $v^U{}_{max}$ and half-saturation constant

K_m^U) follow Arrhenius relations with temperature. This defines our baseline 'kinetics-only'

scenario of temperature-dependent decomposition:

(2a) $v_{max}^D = v_0^D \; e^{-\frac{E_v^D}{R(T+273)}}$

(2b) $K_m^D = K_0^D \; e^{-\frac{E_K^D}{R(T+273)}}$

(2c) $v_{max}^U = v_0^U \; e^{-\frac{E_v^U}{R(T+273)}}$

(2d) $K_m^U = K_0^U \; e^{-\frac{E_K^U}{R(T+273)}}$

where T is temperature in Celsius, R is the ideal gas constant, and the E parameters denote

the corresponding activation energies.

We consider two additional scenarios for the influence of temperature on

decomposition. In the temperature-dependent microbial mortality scenario[32], the microbial

death rate increases with temperature. This could be due to a higher risk of predation or

pathogenic infection at higher temperatures, or faster microbial senescence due to higher

protein turnover[32]. In this scenario, the microbial death rate d_M depends on temperature

according to

(3) $d_M(T) = d_{M0}e^{-\frac{E_{dM}}{R(T+273)}}$

as in ref. 32[32].

In the temperature-dependent microbial growth efficiency (MGE) scenario, the MGE

decreases with temperature[18,24,32], possibly due to higher maintenance costs at higher

temperature[33]. This is modeled by making the microbial growth efficiency γ_M vary linearly

with temperature[18,22,47,65]:

(4) $\gamma_M(T) = \gamma_{M,ref} - m\ (T - T_{ref})$

with $T_{ref} = 20\ °C$.

How scenarios of temperature-dependence and parameter values influence the

response of equilibrium C to temperature is shown in Supplementary Figure 4 and

commented on in the Supplementary Note 4.

Evolutionary Dynamics. The enzyme allocation fraction φ is a 'public good' trait: as an

individual microbe produces exoenzymes, it experiences an energetic cost and obtains a

benefit – access to decomposed organic carbon – that depends on its own and other

microbes's production in the spatial neighborhood[66,67]. As a public good trait, φ is under

strong direct negative selection: 'cheaters' that produce less or no exoenzymes, and thus

avoid the cost while reaping the benefit of enzyme production by cooperative neighbors,

should be at a selective advantage. In a highly diffusive environment in which exoenzymes

are well mixed, φ would evolve to zero, leading to evolutionary suicide[68]. However, in a

more realistic spatially distributed environment with limited exoenzyme diffusion,

microbes with a given trait are more likely to interact with phenotypically similar microbes,

which puts more cooperative microbes at a competitive advantage over less cooperative

strains[66,69,70]. This generates indirect positive selection on trait φ.

We derived the trait value φ^* at which negative and positive selections balance, giving

the evolutionarily stable microbial strategy (Supplementary Note 3), as

(5) $\varphi^*(T) = 1 - \dfrac{d_M}{\gamma_M \, v_{max}^U} - \dfrac{1}{c_0}$

where c_0 measures the competitive advantage, due to spatially local interactions, of any

given strain over a slightly less cooperative strain, or 'competition asymmetry'

(Supplementary Note 3). The parameter c_0 is likely to depend on the diffusivity of D, which

may itself vary with soil properties such as texture or water content.

As temperature rises from T_0 to T, the direction and magnitude of the microbial

adaptive response is measured by $\Delta\varphi^* = \varphi^*(T) - \varphi^*(T_0)$, which depends on the scenario

of temperature dependence. The evolutionary response (EVO response) of SOC is given by

(6)   EVO response $= \Delta C_{EVO}(T_0, T) = C(T, \varphi^*(T)) - C(T_0, \varphi^*(T_0))$

where $C(T, \varphi)$ denotes ecosystem equilibrium C at temperature T, given enzyme allocation

fraction φ. The EVO response is to be compared with the response in the absence of

evolution (ECOS response):

(7)   ECOS response $= \Delta C_{ECOS}(T_0, T) = C(T, \varphi^*(T_0)) - C(T_0, \varphi^*(T_0))$

in which the enzyme allocation fraction is fixed at its T_0-adapted value, $\varphi^*(T_0)$ (Fig. 1c).

We measure the magnitude of the evolutionary effect (EVO effect) as the difference

between the EVO response averaged over the temperature range (T_0, T) and the ECOS

response averaged over the same temperature range, normalized by the ECOS response:

(8) $EVO\ effect = \dfrac{\left| \int_{T_0}^{T} \Delta C_{EVO}(T_0,T) - \int_{T_0}^{T} \Delta C_{ECOS}(T_0,T) \right|}{\left| \int_{T_0}^{T} \Delta C_{ECOS}(T_0,T) \right|}$

This evaluation allows us to compare EVO effects across systems that differ in the

magnitude of their ECOS response. In all simulations we use $T = T_0 + \Delta T$ where $\Delta T = 5$

°C. In general, the ECOS and EVO responses are monotonic, close-to-linear functions of T

over the considered temperature ranges $(T_0, T_0 + \Delta T)$, which makes all our comparative

analyses almost insensitive to our choice of ΔT.

Parameter default values and sensitivity analysis. Under the temperature-dependent

kinetics-only scenario, the ecosystem model (equations (1) and (2)) includes seven

microbial parameters (φ, γ_M, d_M, v_0^U, E_v^U, K_0^U, E_K^U), six enzyme parameters (γ_Z, d_Z, v_0^D, E_v^D,

K_0^D, E_K^D) and four environmental parameters (I, e_C, e_D, T). Our set of default parameter

values is derived from Allison *et al.* (2010)[18] (Supplementary Table 1). The enzyme

allocation fraction default value is 10% at 20 °C[34]. For the dependence of enzyme kinetics

parameters on temperature, $v_{max}^D(T)$ and $K_m^D(T)$, we selected the Arrhenius equations that

best fit data from California[22] (mean annual $T = 17$ °C) and match values at 20 °C (0.42 and

600, respectively)[18]. For the uptake kinetic parameters, we obtained $v_{max}^U(T)$ by selecting

v_0^U that best fits the Arrhenius equation in ref. 18[18] with $E_v^U = 35$ and we obtained $K_m^U(T)$

by selecting K_0^U and E_K^U that best fit the linear relation used in ref. 18[18]. To parametrize the

temperature-dependent mortality ($d_{M,ref}$, T_{ref}) and MGE ($\gamma_{M,ref}$, m, T_{ref}) models, we used

values from ref. 32[32] and tested two values of E_{dM} ($E_{dM} = 0$ is the enzyme only temperature-

dependent model). For greater realism, we used a higher value of the exoenzyme

deactivation rate (twice the value used in ref. 18[18]) and constrained the range of all

parameters in order to enhance stability and produce relative stock sizes that are consistent

with empirical data, so that at equilibrium M is about 1% of C, Z is about 1% of M and D is

limiting (hence close to 0 at equilibrium[71,72]),

We analysed the model sensitivity by varying parameters over two orders of magnitude

(as in ref. 18[18]) – except γ_M and γ_Z for which we used the whole range over which the non-

trivial ecosystem equilibrium is stable (Supplementary Table 2). To assess the significance

of our findings for real ecosystems, we focused on five sites for which empirical data[22]

could be used to constrain the model. The five sites contrast strongly in their initial

temperature, T_0, and decomposition kinetics, $v_{max}^D(T)$ and $K_m^D(T)$ for which we selected the

Arrhenius equations (2) that best fit the relations used in ref. 22[22] (Supplementary Table 3).

Data availability

No datasets were generated or analysed during the current study.

Code availability

The sensitivity analysis and figures that support the findings of this study were coded on

Mathematica. The code file has been deposited in "MicFigCode",

https://github.com/elsaabs/MicFigCode/tree/master.

References

1. Falkowski, P. G., Fenchel, T. & Delong, E. F. The microbial engines that drive Earth's biogeochemical cycles. *Science* **320**, 1034–1039 (2008).

2. Waksman, S. A. & Starkey, R. L. *The soil and the microbe*. (John Wiley And Sons; New York, 1931).

3. Frey, S. D., Lee, J., Melillo, J. M. & Six, J. The temperature response of soil microbial efficiency and its feedback to climate. *Nat. Clim. Chang.* **3**, 395 (2013).

4. Singh, B. K., Bardgett, R. D., Smith, P. & Reay, D. S. Microorganisms and climate change: terrestrial feedbacks and mitigation options. *Nat. Rev. Microbiol.* **8**, 779–790 (2010).

5. Davidson, E. A. & Janssens, I. A. Temperature sensitivity of soil carbon decomposition and feedbacks to climate change. *Nature* **440**, 165–173 (2006).

6. Bradford, M. A. *et al.* Thermal adaptation of soil microbial respiration to elevated temperature. *Ecol. Lett.* **11**, 1316–1327 (2008).

7. Tucker, C. L., Bell, J., Pendall, E. & Ogle, K. Does declining carbon-use efficiency explain thermal acclimation of soil respiration with warming? *Glob. Chang. Biol.* **19**, 252–263 (2013).

8. Creamer, C. A. *et al.* Microbial community structure mediates response of soil C decomposition to litter addition and warming. *Soil Biol. Biochem.* **80**, 175–188 (2015).

9. Wei, H. *et al.* Thermal acclimation of organic matter decomposition in an artificial forest soil is related to shifts in microbial community structure. *Soil Biol. Biochem.* **71**, 1–12 (2014).

629   10.   Padfield, D., Yvon-Durocher, G., Buckling, A., Jennings, S. & Yvon-Durocher, G.

630   Rapid evolution of metabolic traits explains thermal adaptation in phytoplankton.

631   *Ecol. Lett.* **19**, 133–142 (2016).

632   11.   Schaum, C.-E. *et al.* Adaptation of phytoplankton to a decade of experimental

633   warming linked to increased photosynthesis. *Nat Ecol Evol* **1**, 94 (2017).

634   12.   Monroe, J. G. *et al.* Ecoevolutionary Dynamics of Carbon Cycling in the

635   Anthropocene. *Trends Ecol. Evol.* **33**, 213–225 (2018).

636   13.   Ratledge, C. Biodegradation of oils, fats and fatty acids. in *Biochemistry of microbial*

637   *degradation* (ed. Ratledge, C.) 89–141 (Springer Netherlands, 1994).

638   14.   Harder, W. & Dijkhuizen, L. Physiological responses to nutrient limitation. *Annu. Rev.*

639   *Microbiol.* **37**, 1–23 (1983).

640   15.   Velicer, G. J. Social strife in the microbial world. *Trends Microbiol.* **11**, 330–337

641   (2003).

642   16.   Trivedi, P. *et al.* Microbial regulation of the soil carbon cycle: evidence from gene-

643   enzyme relationships. *ISME J.* **10**, 2593–2604 (2016).

644   17.   Rainey, P. B. & Rainey, K. Evolution of cooperation and conflict in experimental

645   bacterial populations. *Nature* **425**, 72–74 (2003).

646   18.   Allison, S. D., Wallenstein, M. D. & Bradford, M. A. Soil-carbon response to

647   warming dependent on microbial physiology. *Nat. Geosci.* **3**, 336 (2010).

648   19.   Sinsabaugh, R. L. & Moorhead, D. L. Resource allocation to extracellular enzyme

649   production: A model for nitrogen and phosphorus control of litter decomposition. *Soil*

650   *Biol. Biochem.* **26**, 1305–1311 (1994).

20.   Allison, S. D. A trait-based approach for modelling microbial litter decomposition.

Ecol. Lett. **15**, 1058–1070 (2012).

21.   Steinweg, J. M., Dukes, J. S., Paul, E. A. & Wallenstein, M. D. Microbial responses to

multi-factor climate change: effects on soil enzymes. *Front. Microbiol.* **4**, 146 (2013).

22.   German, D. P., Marcelo, K. R. B., Stone, M. M. & Allison, S. D. The Michaelis-

Menten kinetics of soil extracellular enzymes in response to temperature: a cross-

latitudinal study. *Glob. Chang. Biol.* **18**, 1468–1479 (2012).

23.   Allison, S. D. Modeling adaptation of carbon use efficiency in microbial communities.

Front. Microbiol. **5**, 571 (2014).

24.   Wieder, W. R., Bonan, G. B. & Allison, S. D. Global soil carbon projections are

improved by modelling microbial processes. *Nat. Clim. Chang.* **3**, 909 (2013).

25.   O'Brien, S., Hodgson, D. J. & Buckling, A. The interplay between microevolution and

community structure in microbial populations. *Curr. Opin. Biotechnol.* **24**, 821–825

(2013).

26.   Boon, E. *et al.* Interactions in the microbiome: communities of organisms and

communities of genes. *FEMS Microbiol. Rev.* **38**, 90–118 (2014).

27.   Strauss, S. Y. Ecological and evolutionary responses in complex communities:

implications for invasions and eco-evolutionary feedbacks. *Oikos* **123**, 257–266

(2014).

28.   Zhang, X. *et al.* Assessing five evolving microbial enzyme models against field

measurements from a semiarid savannah—What are the mechanisms of soil

respiration pulses? *Geophys. Res. Lett.* **41**, 6428–6434 (2014).

29.   Wallenstein, M. D., Mcmahon, S. K. & Schimel, J. P. Seasonal variation in enzyme

activities and temperature sensitivities in Arctic tundra soils. *Glob. Chang. Biol.* **15**,

675  1631–1639 (2009).

30. Burns, R. G. *et al.* Soil enzymes in a changing environment: Current knowledge and future directions. *Soil Biol. Biochem.* **58**, 216–234 (2013).

31. Hochachka, P. W. & Somero, G. N. *Biochemical Adaptation: Mechanism and Process in Physiological Evolution*. (Oxford University Press, 2002).

32. Hagerty, S. B. *et al.* Accelerated microbial turnover but constant growth efficiency with warming in soil. *Nat. Clim. Chang.* **4**, 903 (2014).

33. Sinsabaugh, R. L., Manzoni, S., Moorhead, D. L. & Richter, A. Carbon use efficiency of microbial communities: stoichiometry, methodology and modelling. *Ecol. Lett.* **16**, 930–939 (2013).

34. Schimel, J. P. & Weintraub, M. N. The implications of exoenzyme activity on microbial carbon and nitrogen limitation in soil: a theoretical model. *Soil Biol. Biochem.* **35**, 549–563 (2003).

35. Carreiro, M. M., Sinsabaugh, R. L., Repert, D. A. & Parkhurst, D. F. MICROBIAL ENZYME SHIFTS EXPLAIN LITTER DECAY RESPONSES TO SIMULATED NITROGEN DEPOSITION. *Ecology* **81**, 2359–2365 (2000).

36. Waldrop, M. P., Zak, D. R., Sinsabaugh, R. L., Gallo, M. & Lauber, C. NITROGEN DEPOSITION MODIFIES SOIL CARBON STORAGE THROUGH CHANGES IN MICROBIAL ENZYMATIC ACTIVITY. *Ecol. Appl.* **14**, 1172–1177 (2004).

37. Malik, A. A., Puissant, J., Goodall, T., Allison, S. D. & Griffiths, R. I. Soil microbial communities with greater investment in resource acquisition have lower growth yield. *Soil Biol. Biochem.* **132**, 36–39 (2019).

38. Sulman, B. N. *et al.* Multiple models and experiments underscore large uncertainty in

soil carbon dynamics. *Biogeochemistry* **141**, 109–123 (2018).

39. Scheiner, S. M. Genetics and evolution of phenotypic plasticity. *Annu. Rev. Ecol. Syst.*

24, 35–68 (1993).

40. Rebolleda-Gómez, M. & Travisano, M. The Cost of Being Big: Local Competition,

Importance of Dispersal, and Experimental Evolution of Reversal to Unicellularity.

Am. Nat. **192**, 731–744 (2018).

41. Stegen, J. C., Enquist, B. J. & Ferrière, R. Eco-evolutionary community dynamics:

covariation between diversity and invasibility across temperature gradients. *Am. Nat.*

180, E110–26 (2012).

42. Evans, S., Martiny, J. B. H. & Allison, S. D. Effects of dispersal and selection on

stochastic assembly in microbial communities. *ISME J.* **11**, 176–185 (2017).

43. Glassman, S. I. *et al.* Decomposition responses to climate depend on microbial

community composition. *Proc. Natl. Acad. Sci. U. S. A.* **115**, 11994–11999 (2018).

44. Albright, M. B. N., Chase, A. B. & Martiny, J. B. H. Experimental Evidence that

Stochasticity Contributes to Bacterial Composition and Functioning in a Decomposer

Community. *MBio* **10**, (2019).

45. Billiard, S., Collet, P., Ferrière, R., Méléard, S. & Tran, V. C. Stochastic dynamics for

adaptation and evolution of microorganisms. *arXiv [math.PR]* (2016).

46. Gómez, P. *et al.* Local adaptation of a bacterium is as important as its presence in

structuring a natural microbial community. *Nat. Commun.* **7**, 12453 (2016).

47. Wang, G., Post, W. M. & Mayes, M. A. Development of microbial-enzyme-mediated

decomposition model parameters through steady-state and dynamic analyses. *Ecol.*

Appl. **23**, 255–272 (2013).

48. Romero-Olivares, A. L., Taylor, J. W. & Treseder, K. K. Neurospora discreta as a

model to assess adaptation of soil fungi to warming. *BMC Evolutionary Biology* **15**,

(2015).

49. Allison, S. D., Romero-Olivares, A. L., Lu, L., Taylor, J. W. & Treseder, K. K.

Temperature acclimation and adaptation of enzyme physiology in Neurospora discreta.

Fungal Ecol. **35**, 78–86 (2018).

50. Sauterey, B., Ward, B. A., Follows, M. J., Bowler, C. & Claessen, D. When

everything is not everywhere but species evolve: an alternative method to model

adaptive properties of marine ecosystems. *J. Plankton Res.* **37**, 28–47 (2015).

51. Pontarp, M. *et al.* The Latitudinal Diversity Gradient: Novel Understanding through

Mechanistic Eco-evolutionary Models. *Trends Ecol. Evol.* **34**, 211–223 (2019).

52. Trivedi, P., Anderson, I. C. & Singh, B. K. Microbial modulators of soil carbon

storage: integrating genomic and metabolic knowledge for global prediction. *Trends

Microbiol.* **21**, 641–651 (2013).

53. Harmon, L. J. *et al.* Evolutionary diversification in stickleback affects ecosystem

functioning. *Nature* **458**, 1167–1170 (2009).

54. Bennett, A. F. & Lenski, R. E. An experimental test of evolutionary trade-offs during

temperature adaptation. *Proc. Natl. Acad. Sci. U. S. A.* **104 Suppl 1**, 8649–8654

(2007).

55. Good, B. H., McDonald, M. J., Barrick, J. E., Lenski, R. E. & Desai, M. M. The

dynamics of molecular evolution over 60,000 generations. *Nature* **551**, 45–50 (2017).

56. Gómez, P. & Buckling, A. Bacteria-phage antagonistic coevolution in soil. *Science*

332, 106–109 (2011).

57. Buckling, A. & Rainey, P. B. Antagonistic coevolution between a bacterium and a

bacteriophage. *Proc. Biol. Sci.* **269**, 931–936 (2002).

58. Garud, N. R., Good, B. H., Hallatschek, O. & Pollard, K. S. Evolutionary dynamics of

bacteria in the gut microbiome within and across hosts. *PLoS Biol.* **17**, e3000102

(2019).

59. Treseder, K. K. *et al.* Integrating microbial ecology into ecosystem models: challenges

and priorities. *Biogeochemistry* **109**, 7–18 (2012).

60. Tang, J. & Riley, W. J. Weaker soil carbon–climate feedbacks resulting from

microbial and abiotic interactions. *Nat. Clim. Chang.* **5**, 56 (2014).

61. Harte, J. & Kinzig, A. P. Mutualism and competition between plants and

decomposers: implications for nutrient allocation in ecosystems. *Am. Nat.* **141**, 829–

846 (1993).

62. Kaiser, C., Franklin, O., Richter, A. & Dieckmann, U. Social dynamics within

decomposer communities lead to nitrogen retention and organic matter build-up in

soils. *Nat. Commun.* **6**, 8960 (2015).

63. Follows, M. J., Dutkiewicz, S., Grant, S. & Chisholm, S. W. Emergent biogeography

of microbial communities in a model ocean. *Science* **315**, 1843–1846 (2007).

64. Stone, M. M. *et al.* Temperature sensitivity of soil enzyme kinetics under N-

fertilization in two temperate forests. *Glob. Chang. Biol.* **18**, 1173–1184 (2012).

65. Li, J., Wang, G., Allison, S. D., Mayes, M. A. & Luo, Y. Soil carbon sensitivity to

temperature and carbon use efficiency compared across microbial-ecosystem models

of varying complexity. *Biogeochemistry* **119**, 67–84 (2014).

66. Allison, S. D. Cheaters, diffusion and nutrients constrain decomposition by microbial

767  enzymes in spatially structured environments: Constraints on enzymatic

768  decomposition. *Ecol. Lett.* **8**, 626–635 (2005).

769 67. Nadell, C. D., Drescher, K. & Foster, K. R. Spatial structure, cooperation and

770  competition in biofilms. *Nat. Rev. Microbiol.* **14**, 589–600 (2016).

771 68. Ferriere, R. & Legendre, S. Eco-evolutionary feedbacks, adaptive dynamics and

772  evolutionary rescue theory. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **368**, 20120081

773  (2013).

774 69. Durrett, R. & Levin, S. The Importance of Being Discrete (and Spatial). *Theor. Popul.*

775  *Biol.* **46**, 363–394 (1994).

776 70. Chao, L. & Levin, B. R. Structured habitats and the evolution of anticompetitor toxins

777  in bacteria. *Proc. Natl. Acad. Sci. U. S. A.* **78**, 6324–6328 (1981).

778 71. Fierer, N., Strickland, M. S., Liptzin, D., Bradford, M. A. & Cleveland, C. C. Global

779  patterns in belowground communities. *Ecol. Lett.* **12**, 1238–1249 (2009).

780 72. Ladd, J. N., Amato, M., Zhou, L.-K. & Schultz, J. E. Differential effects of rotation,

781  plant residue and nitrogen fertilizer on microbial biomass and organic matter in an

782  Australian alfisol. *Soil Biol. Biochem.* **26**, 821–831 (1994).

End Notes

Acknowledgements We thank Rachel Gallery, Pierre-Henri Gouyon, Moira Hough,

Hélène Leman, Laura Meredith, and Mitch Pavao-Zuckerman for discussion. E.A. was

supported by fellowships from Ecole Doctorale Frontières du Vivant and MemoLife

Laboratory of Excellence (PIA-10-LBX-54). S.R.S. was supported by a grant from the

Genomic Science Program of the United States Department of Energy (DOE) (DE-

SC0016440) and the University of Arizona's Agnese Nelms Haury Program in

Environment and Social Justice. R.F. acknowledges support from FACE Partner University

Fund, CNRS Mission pour l'Interdisciplinarité, and PSL University (IRIS OCAV and PSL-

University of Arizona Mobility Program).

Author contributions R.F. conceived the study. All authors developed the model. E.A.

performed the analysis. E.A. and R.F. wrote the first version of the manuscript. All authors

finalized the paper.

Competing interests The authors declare no competing interests.

Additional information

Supplementary information is available for this paper at https://doi.org/xxx

Correspondence and requests for materials should be addressed to E.A. or R.F.

     Legends

     Figure 1. Effect of temperature and enzyme allocation fraction on SOC ecosystem

     equilibrium. a, Structure of the microbial-enzyme ecosystem model (see Methods for

     details): SOC stock is the balance of plant input, I, and loss by exoenzyme-mediated

     degradation to DOC (D), which in turn is allocated between (fraction φ) production of

     enzymes (Z) and (fraction $1-\varphi$) growth of microbial biomass (M). **b**, Effect of temperature

     and enzyme allocation fraction, φ, on SOC equilibrium, C, in the baseline scenario of

     temperature dependence. **c**, Response of SOC ecosystem equilibrium, C, to a 5°C increase

     in temperature (from 20 °C to 25 °C) as a function of enzyme allocation fraction, φ.

     Parameters are set to their default values (Supplementary Table 1), except $I = 5\ 10^{-3}$, $v_0^U =$

     10^5, $E_v^U = 38$ and $c_0 = 1.17$.

     Figure 2. Effect of microbial evolutionary adaptation on the SOC equilibrium

     response to + 5 °C warming (EVO effect). Temperature influences enzyme kinetics only

     (baseline scenario of temperature dependence). **a**, Influence of microbial biomass

     production efficiency, γ_M, and microbial mortality rate, d_M. **b**, Influence of microbial

     resource acquisition traits v_0^U and E_v^U. **c-d**, Influence of competition asymmetry, c_0, and

     initial temperature, T_0. In all figures, constant parameters are set to their default values

     (Supplementary Table 1) and I is set to $5\ 10^{-3}$. Points A1 and B1 indicate the default

     parameter values. Point A2 (respectively B2) exemplifies values of γ_M and d_M (resp. v_0^U and

     E_v^U) for which the EVO effect is strong. Panel **c** (resp. **d**) shows the influence of c_0 and T_0

     on the EVO effect at A2 (resp. B2).

     Figure 3. Ecosystem and ecosystem-evolutionary responses of SOC equilibrium to

     warming (up to + 5 °C) for three scenarios of temperature dependence. Ecosystem and

     ecosystem-evolutionary changes in SOC equilibrium C given by Eq. (3a) (without

     evolution, dashed curves) and Eq. (3b) (with evolution, plain curves) are plotted as a

     function of the increase in temperature. *Blue curves*, initial temperature $T_0 = 5°C$. *Black*

curves, $T_0 = T_{ref} = 20°C$. *Red curves*, $T_0 = 30°C$. *Insets*, Direction and magnitude of EVO

effect (%), from - 150 % to + 150 %, color code indicates T_0 as before. **a**, Baseline scenario

of temperature dependence (enzyme kinetics only). **b**, Temperature-dependent microbial

turnover, with $E_{dM} = 25 < E_v^U$. **c**, Temperature- dependent microbial turnover, with $E_{dM} =$

$55 > E_v^U$. **d**, Temperature-dependent MGE, with $m = 0.014$. Parameters values correspond

to point B2 in Fig. 2 ($I = 5\ 10^{-3}$, $v_0^U = 10^5$, $E_v^U = 38$, $c_0 = 1.17$); other parameters are set to

their default values (Table S1).

Figure 4. Ecosystem and ecosystem-evolutionary responses of SOC equilibrium to + 5

°C warming predicted for five sites. a-d, ECOS response. **e-h**, EVO response. **i-l**, EVO

effect. AK: Alaska, boreal forest, $T_0 = 0.1°C$. ME: Maine, temperate forest, $T_0 = 5°C$. WV:

West Virginia, temperate forest, $T_0 = 9°C$. CA: California, temperate grassland, $T_0 = 17°C$.

CR: Costa Rica, tropical rain forest, $T_0 = 26°C$. First row (**a, e, i**): baseline scenario of

temperature dependence. Second row (**b, f, j**): temperature-dependent microbial turnover

scenario with $E_{dM} = 25 < E_v^U$. Third row (**c, g, k**): temperature- dependent microbial

turnover scenario with $E_{dM} = 55 > E_v^U$. Fourth row (**d, h, l**): temperature-dependent MGE

scenario ($m = - 0.014$). The influence of competition asymmetry, c_0, is shown (low: $c_0 =$

1.17, intermediate: $c_0 = 1.34$, high: $c_0 = 1.5$). For clarity, vertical axis for ECOS and EVO

responses are truncated at - 65 mg C cm^{-3}. Actual values for AK with $c_0 = 1.17$ are ECOS

response = - 170 mg C cm^{-3} and EVO response = - 556 mg C cm^{-3}; actual value for WV

with $c_0 = 1.17$ is EVO response = - 92.8 mg C cm^{-3}. Parameter values correspond to point

B2 in Fig. 2 ($I = 5\ 10^{-3}$, $v_0^U = 10^5$, $E_v^U = 38$, $c_0 = 1.17$); other parameters are set to their

default values (Supplementary Table 1).

 Figure 1

a

b

c

 Figure 2

 Figure 3

 Figure 4