Modelling the population dynamics of *Plasmodium falciparum* gametocytes in

humans during malaria infection

Pengxing Cao[1, *], Katharine A. Collins[2], Sophie Zaloumis[3], Thanaporn Wattanakul[4,5], Joel

Tarning[4,5], Julie A. Simpson[3], James S. McCarthy[6], James M. McCaw[1,3,7, *]

1. School of Mathematics and Statistics, University of Melbourne, Melbourne, Australia.

2. Department of Medical Microbiology, Radboud University Medical Center, Nijmegen,

Netherland.

3. Centre for Epidemiology and Biostatistics, Melbourne School of Population and Global

Health, University of Melbourne, Melbourne, Australia

4. Mahidol-Oxford Tropical Medicine Research Unit, Faculty of Tropical Medicine,

Mahidol University, Bangkok, Thailand.

5. Centre for Tropical Medicine and Global Health, Nuffield Department of Medicine,

University of Oxford, Oxford, United Kingdom

6. QIMR Berghofer Medical Research Institute, Brisbane, Queensland, Australia.

7. Epidemiology, Peter Doherty Institute for Infection and Immunity, Parkville, Victoria,

Australia.

* Address correspondence to Pengxing Cao (pengxing.cao@unimelb.edu.au) or James M.

McCaw (jamesm@unimelb.edu.au).

Abstract

Every year over two hundred million people are infected with the malaria parasite. Renewed efforts to eliminate malaria has highlighted the potential to interrupt transmission from humans to mosquitoes which is mediated through the gametocytes. Reliable prediction of transmission requires an improved understanding of *in vivo* kinetics of gametocytes. Here we study the population dynamics of *Plasmodium falciparum* gametocytes in human hosts by establishing a framework which incorporates improved measurements of parasitaemia in humans, a novel mathematical model of gametocyte dynamics, and model validation using a Bayesian hierarchical inference method. We found that the novel mathematical model provides an excellent fit to the available clinical data from 17 volunteers infected with *P. falciparum*, and reliably predicts observed gametocyte levels. We estimated the *P. falciparum*'s sexual commitment rate and gametocyte sequestration time in humans to be 0.54% (95% credible interval: 0.30-1.00) per life cycle and 8.39 (6.54-10.59) days respectively. Furthermore, we used the data-calibrated model to predict the effects of those gametocyte dynamics parameters on human-to-mosquito transmissibility, providing a method to link within-human host kinetics of malaria infection to epidemiological-scale infection and transmission patterns.

Malaria is a mosquito-borne parasitic disease caused by protozoan parasites of the *Plasmodium*

genus. It is estimated to have caused approximately 219 million new cases and 435,000 deaths

(primarily due to *Plasmodium falciparum*) in 2017[1]. New tools will be required to achieve the

ambitious goal of malaria elimination. Among tools proposed are novel interventions to block

transmission of infection from human hosts to vector mosquitoes[2]. Falciparum malaria is

transmitted from the human host to the mosquito when terminally differentiated male and female

sexual stages of the parasite, called gametocytes, are taken up by the female *Anopheles* mosquito

during a blood meal[3,4]. The level of gametocytes in the blood, often referred to as

gametocytaemia, is highly associated with the probability of human-to-mosquito transmission[5,6].

Gametocyte levels below a certain threshold (i.e. < 1000 per mL blood) minimise the probability

for a mosquito to take up both a male and female gametocyte during a blood meal, necessary to

propagate infection[7]. An accurate understanding of the kinetics of gametocytes in human hosts is

essential to predict the probability of transmission. A mathematical model that accurately

captures the processes that give rise to observed gametocyte kinetics would be an important

predictive tool to facilitate the design of effective intervention strategies.

Nonetheless, there is significant uncertainty surrounding fundamental aspects of *Plasmodium*

falciparum gametocyte population dynamics in humans. Parameters such as how many

gametocytes are produced during each asexual life cycle, the period of time in which early

gametocytes disappear from the circulation before mature gametocytes appear (referred to as

sequestration), and the period in which gametocytes circulate are poorly quantified. These gaps

in understanding are due to a range of technical and logistic limitations. The first is the relatively

poor sensitivity of the standard diagnostic test, namely microscopic examination of blood films.

In vivo estimates of gametocyte kinetic parameters have been primarily based on historical data

from neurosyphilis patients whose illness was treated with so-called malariotherapy[8,9]. In these

studies, the limit of quantification was approximately 10^4 parasites/mL blood, at least two orders

of magnitude higher than that of current quantitative PCR (qPCR) assays[10]. This high limit of

quantification prevents an accurate estimation of the onset of the emergence of both asexual

parasites and mature gametocytes in peripheral blood. The second limitation is that the available

estimates of gametocyte population dynamics parameters based on *in vitro* cultures[11,12] may not

applicable to natural infection with *P. falciparum* gametocytes due to the *in vitro* conditions that

may not mimic what occurs in the human host[3].

Recent advances in experimental medicine using volunteer infection studies (VIS), otherwise

known as controlled human malaria infection (CHMI) studies, allow prospective study design

and data collection with the explicit aim of collecting *in vivo* data[13], in particular an improved

quantification of *P. falciparum* gametocyte kinetics by qPCR applied in a novel VIS[7].

Furthermore, the models and fitting methods used in the neurosyphilis patient studies have been

superseded for parameter estimation by increasingly sophisticated within-host models[14] and

improvements in computational algorithms for Bayesian statistical inference[15]. Therefore, there

is an emerging opportunity to improve our quantitative understanding of the population

dynamics of *P. falciparum* gametocytes in human hosts by combining the novel VIS data and

advanced modelling approaches.

In this paper, we developed a novel mathematical model of gametocyte dynamics, fitted the

model to the VIS data and estimated the gametocyte dynamics parameters using a Bayesian

hierarchical inference method (details are given in the Materials and Methods). We demonstrate

that the data-calibrated model can reliably predict the time course of gametocytaemia and thus

should form an essential part of modelling studies of malaria transmission.

Results

Model fitting and validation

The outcome variable used in model fitting was the total parasitaemia (total circulating asexual

parasites and gametocytes per mL blood) collected from a previously published VIS[7], with other

measurements from the same study, such as the asexual parasitaemia (circulating asexual

parasites per mL blood) and gametocytaemia (circulating female and male gametocytes per mL

blood), used to validate model predictions.

The results of fitting the mathematical model to total parasitaemia data for all 17 volunteers are

given in Figure 1 (upper panels) where 12 volunteers experienced recrudescence while the rest

did not. The median of posterior predictions and 95% prediction interval (PI) were computed

from 5000 model simulations based on 5000 samples from the posterior parameter distribution

(see Materials and Methods). The results show that the model describes the total parasitaemia

data very well (median of posterior predictions vs. observed data). Furthermore, the narrow 95%

PI indicates a low level of uncertainty in predicted total parasitaemia.

Figure 1: Results of model fitting for all 17 volunteers. Data are presented by circles. The
median of posterior predictions (solid line) and 95% prediction interval (PI, shaded
area) are generated by 5000 model simulations based on 5000 samples from the
posterior parameter distribution (see the Materials and Methods).

Having calibrated the model against total parasitaemia, the 5000 posterior parameter sets were

used to calculate the median of posterior predictions and 95% PI of the asexual parasitaemia and

gametocytaemia versus time profiles. Model predictions of the asexual parasitaemia and

gametocytaemia for all 17 volunteers are shown in Figure 2 and Figure 3 respectively (curves:

median prediction; shaded areas: 95% PI) and are compared to the asexual parasitaemia and

gametocytaemia data (circles). We emphasise that this was not a fitting exercise, rather an

independent validation of the calibrated model.

Time post-inoculation (days)

Figure 2: Comparison of model predictions and clinical data for the asexual parasitaemia

for all 17 volunteers. Data are presented by circles. The median of posterior

predictions (solid curve) and 95% PI (shaded area) are generated by 5000 model

simulations based on 5000 samples from the posterior parameter distribution (see the

Materials and Methods). Note that one asexual parasite/mL was the level of detection

and no data are available for Volunteer 101 and 106 to validate the model predictions.

Figure 3: Comparison of model predictions and clinical data for the gametocytaemia for all 17 volunteers. Data are presented by circles. The median of posterior predictions (solid curve) and 95% PI (shaded area) are generated by 5000 model simulations based on 5000 samples from the posterior parameter distribution (see the Materials and Methods). Note that one gametocyte/mL was the level of detection.

For the majority of asexual parasitaemia data there is an excellent predictive capability for the model (see Figure 2), in particular the accurate predictions for both the recrudescent case where a portion of asexual parasitaemia data diverge from the total parasitaemia measurement (e.g. Volunteer 103, 104, 105, 201, 203, 304 and 307) and the non-recrudescent case where the posterior-median prediction curve (solid red curve) lies below the limit of detection (one asexual parasite/mL) (e.g. Volunteer 202, 301 and 302). However, there are some discrepant observations. The model under-predicts (see Volunteer 204) or over-predicts (see Volunteer 303, 305 and 306) a portion of the asexual parasitaemia data. Furthermore, for some volunteers such

as 202, 301 and 302, the 95% PI (red shaded area) extends into the detectable region again after
the asexual parasitaemia reaches below the detection limit, indicating that there was a small
chance that the patients may have suffered a recrudescence during the observation period (of
course, they did not) or after the observation period (although this predication cannot be
evaluated because artemisinin combination therapy was given immediately after the period).
Despite the above discrepant observations for asexual parasitaemia, the model predictions of
gametocytaemia are very consistent with the data for all 17 volunteers, as shown in Figure 3.

Estimation of gametocyte population dynamics parameters

The model calibration process provided the joint posterior density for the model parameters,
which were used to estimate some key biological parameters governing the population dynamics
of *P. falciparum* gametocytes (see the Material and Methods for details). As shown in Table 1,
the sexual commitment rate – the percentage of asexual parasites entering sexual development
during each asexual life cycle – is estimated to be 0.54% (95% credible interval (CI): 0.30%,
1.00%). This *in vivo* estimate is much lower than 11% that was estimated from *in vitro* data[11].
The gametocyte sequestration time is the average time that immature gametocytes (stages I—IV)
cannot be observed in the peripheral circulation and was estimated to be 8.39 days (95% CI:
6.54, 10.59 days). The estimate for the circulating gametocyte lifespan is 63.5 days, with a broad
95% CI (12.7 to 1513.9 days) resulting from the long-tailed posterior distribution (see Figure S1,
Supplementary Information) and is much longer than the previous *in vitro* estimate of 16—32
days[12] (note that our lower bound of the 95% CI is lower than the *in vitro* estimated range). The
wide estimate for the circulating gametocyte lifespan, and in particular the high upper bound of
the 95% CI, is primarily due to the limited observation time in the VIS as explored in more detail
in the Discussion.

Table 1: Estimates of some key biological parameters and comparison with the literature.
The estimates of the biological parameters (middle column) are shown as the median and 95%
credible interval (CI) of the marginal posterior parameter distribution (see Figure S1 in the
Supplementary Information). Estimates from the literature (third column) are shown in the
format of either "mean estimate (95% confidence interval)" or "mean estimate [minimum –

maximum estimate]" or simply "a low estimate – a high estimate". Note some quoted estimates

are from studies of *P. falciparum* (*Pf*) from either *in vivo* or *in vitro* studies.

Biological parameters (unit)	Median estimate (95%CI)	Estimates in the literature
Sexual commitment rate (%/life cycle)	0.54 (0.30, 1.00)	11 (6.2, 15.8)[11] (*Pf; in vitro*) 0.64 [0.03 – 13.51][9] (*Pf; in vivo*)
Gametocyte sequestration time (days)	8.39 (6.54, 10.59)	7.4 [4 – 12][9] (*Pf; in vivo*)
Circulating gametocyte lifespan (days)	63.5 (12.7, 1513.9)	16 – 32[12] (*Pf; in vitro*) 6.4 [1.3 – 22.2][9] (*Pf; in vivo*)
Parasite multiplication factor (/life cycle)	21.8 (17.6, 26.9)	10 – 33[16] (*Pf; in vivo*) 16.4 (15.1, 17.8)[17] (*Pf; in vivo*)

As shown in Table 1, there are both similarities and differences in parameter estimates for *P.*

falciparum based on our analysis and the analysis of historical neurosyphilis patient data[9]. We

found that they exhibited very similar *in vivo* sexual commitment rate (median 0.54% vs. mean

0.64% with overlapping plausible ranges) and gametocyte sequestration time (median 8.39 days

vs. mean 7.4 days with overlapping plausible ranges). On the other hand, we found that the

circulating gametocyte lifespan for *P. falciparum* was much longer than that estimated from the

neurosyphilis patient data[9] (median 63.5 days vs. 6.4 days with very different ranges).

Finally, we provided an estimate for the parasite multiplication factor, an important parameter

that quantifies the *net* replication rate of asexual parasites and also influences the rate of

gametocyte generation. Our posterior-median estimate is 21.8 parasites per life cycle (95% CI:

17.6, 26.9), consistent with previous estimates which lie in the range 10—33[16], and a bit larger

than an updated estimate calculated from a pooled analysis of parasite counts from 177

volunteers infected with the same *P. falciparum* strain (16.4 parasites per life cycle)[17].

Predicting the impact of gametocyte kinetics on human-to-mosquito transmissibility

Having validated our mathematical model of asexual parasitaemia and gametocyte dynamics, we were able to predict how the gametocyte dynamics parameters influence the transmissibility of *P. falciparum* malaria from humans to mosquitoes in various epidemiological scenarios. In particular, we focused on the early phase of infection where the innate immune response is minimal and treatment has not been administered (in order to avoid complications that our mathematical model was not designed to capture). Two scenarios were considered:

- Predicting the potential infectiousness of newly hospitalised malaria patients for various values of sexual commitment rate and gametocyte sequestration time. In the model, gametocytaemia was assumed to be a surrogate of the potential infectiousness and newly hospitalised patients were assumed to exhibit a total parasitaemia of approximately 10^8 parasites/mL based on published clinical data from Cambodia and Thailand[18]. As illustrated in Figure 4A, we simulated the model (for different sexual commitment rates and gametocyte sequestration times) and looked at the critical gametocytaemia level (indicated by G_c) corresponding to the time when the total parasitaemia (wave-like black curves) first reached 10^8 parasites/mL (their associations are indicated by the dotted lines and arrows).

- Predicting the non-infectious period of malaria patients for various values of sexual commitment rate and gametocyte sequestration time. In the model, the non-infectious period was defined to be time from the inoculation of infected red blood cells to the time when the gametocytaemia reach 10^3 parasites/mL, which is a threshold allowing for human-to-mosquito transmission[7]. Note that this non-infectious period does not include the latent period due to the liver stage which should be considered if the starting time were taken as the inoculation by mosquito. As illustrated in Figure 4B, we simulated the model (for different sexual commitment rates and gametocyte sequestration times) and identified the critical time (indicated by t_c) when the gametocytaemia (blue curve) first reached 10^3

203   parasites/mL (their associations are indicated by the dotted lines and arrows).

Figure 4: Simulation of two scenarios predicting the dependence of human-to-mosquito transmissibility on the sexual commitment rate and gametocyte sequestration time. (A) illustration of the first scenario: predicting the critical gametocytaemia level (indicated by G_c) at the time when the total parasitaemia reach 10^8 parasites/mL. (B) illustration of the second scenario: predicting the non-infectious period (indicated by t_c), which is defined to be time from inoculation of infected red blood cells to the time when the gametocytaemia reach 10^3 parasites/mL (a threshold for human-to-mosquito transmission[7]). (C and D) Heatmaps showing the dependence of the critical gametocytaemia G_c and the non-infectious period t_c on the sexual commitment rate and gametocyte sequestration time. The black dots represent the benchmark value obtained by simulating the gametocyte population dynamics model using the median estimates of the posterior samples of the population mean parameters (see the

Materials and Methods for details). The red curves indicate the cases corresponding to

gametocytaemia of 10^3 parasites/mL.

Figure 4C shows that a higher sexual commitment rate or a lower gametocyte sequestration time

leads to a higher gametocytaemia (G_c) at the time of hospitalisation. The red curve in Figure 3C

indicates the level curve of 10^3 gametocytes/mL (i.e. the threshold for infectiousness as

mentioned above) dividing the heatmap into two regions. To the left, G_c is below 10^3

gametocytes/mL, suggesting clinical presentation precedes infectiousness, while to the right G_c is

above 10^3 gametocytes/mL and the converse applies. The G_c value obtained by model simulation

using the median estimates of the population mean parameters (indicated by the black dot) is

below 10^3 gametocytes/mL, suggesting that newly hospitalized malaria patients are less likely to

be infectious, and thus efforts to identify and treat infections in a timely manner may have a

substantial impact in terms of reduced transmission potential.

Figure 4D reinforces the result in Figure 4C using the non-infectious period (t_c). As the sexual

commitment rate increases or the gametocyte sequestration time decreases, t_c decreases.

However, for relatively large values of the sexual commitment rate (e.g. > 20%), we observed an

increase in t_c as the sexual commitment rate increases (see the top-right corner of Figure 4D).

This is because an increased fraction of sexual conversion can lead to both a decrease in the rate

of asexual parasite growth and an increase in the number of sexual parasites, and the impact of

the former more than counterbalances that of the latter.

Discussion

We have developed a novel mathematical model of gametocyte dynamics, that integrates both

multi-state asexual parasite's life cycle and the development of gametocytes for the first time.

Model parameters were estimated by fitting the model to data from 17 malaria-naïve volunteers

inoculated with *P. falciparum*-infected red blood cells (3D7 strain). Compared to previous

studies, our work is distinguished by three novel contributions: (1) the use of a prospectively

planned clinical trial to collect more accurate quantitative data of parasite levels measured by

qPCR; (2) the development of a novel population dynamics mathematical model which allows

for robust and biologically-informed extrapolation and hypothesis testing/scenario analysis; and

(3) the use of a Bayesian hierarchical inference method for model calibration and parameter

estimation.

For gametocyte kinetic parameters, we found that our *in vivo* estimate of the *P. falciparum*

sexual commitment rate was similar to that found in the neurosyphilis patient data[9] but was much

smaller than previous *in vitro* estimates (see Table 1). Novel VIS data using biomarkers specific

to early sexual parasites (e.g. PfGEXP5[19]) may help to validate and improve our *in vivo* estimate.

Furthermore, our *in vivo* estimate of the circulating gametocyte lifespan is substantially larger

than previous *in vitro* estimates and is also larger than that found in the neurosyphilis patient

data[9] (see Table 1). One way to improve our *in vivo* estimate would be to use new *P. falciparum*

data with gametocytaemia measurements over a longer period of time to capture the natural

decay of circulating gametocytes, which was not observed in the current VIS data (see Figure 3).

We also predicted the effects of altered gametocyte kinetic parameters on the transmissibility

from human to mosquito, focusing on two scenarios: the infectiousness of newly hospitalised

malaria patients (i.e. the gametocytaemia when asexual parasitaemia first reaches a hospitalised

level of 10^8 parasites/mL in the model); and the non-infectious period of malaria patients (i.e. the

time from the inoculation of infected red blood cells to the time when the gametocytaemia reach

a transmission threshold of 10^3 parasites/mL in the model). We explored how the sexual

commitment rate and gametocyte sequestration time influenced the gametocyte level and the

non-infectious period. We would like to emphasize that human-to-mosquito transmissibility is

determined by both the gametocytaemia level in humans and the relationship between

gametocytaemia and the probability of transmission per bite. A reliable prediction of the former

is essential but not a sole determinant of transmissibility. Therefore, it is also important to

improved our quantitative understanding of the latter, which could be complicated and also

determined by the densities of female and male gametocytes[5,6,20].

Our study has some limitations. The gametocyte population dynamics model, that has been

shown to have sufficient complexity to reproduce the clinical observations, is still a rather coarse

simplification of the actual biological processes. For example, the model does not assume an

adaptive sexual commitment rate[21], nor does it consider the mechanisms of sexual

commitment[22]. Furthermore, the model assumes a constant gametocyte death rate but does not

consider other non-constant formulations as have been previously proposed[8]. Another limitation

is that we assumed a fixed duration for the asexual life cycle of 42 hours, while previous work by
our group suggests that the life cycle may be altered by up to a few hours in response to
antimalarial drugs (e.g. artemisinin[23,24]), although it is not known if piperaquine (which was
administered in this VIS) has a similar effect.

In conclusion, we have developed a novel mathematical model of gametocyte population
dynamics, and demonstrated that it reliably predicts time series data of gametocytaemia. The
model provides a powerful predictive tool for informing the design of future volunteer infection
studies which aim to test transmission-blocking interventions. Furthermore, the within human
host transmission model can be incorporated into epidemiological-scale models to refine
predictions of the impacts of various antimalarial treatments and transmission interventions.

Materials and Methods

Study population and measurements

The data used in this modelling study are from a previously published VIS[7] where 17 malaria-
naïve volunteers were inoculated with *P. falciparum*-infected red blood cells (3D7 strain). The
study was approved by the QIMR Berghofer Human Research Ethics Committee and registered
with ClinicalTrials.gov (NCT02431637 and NCT02431650). The volunteers were treated with
480 mg piperaquine phosphate (PQP) on day 7 or 8 post-inoculation to attenuate asexual parasite
growth and a second dose of 960 mg PQP was given to any volunteer for treatment of
recrudescent asexual parasitaemia. All participants received a course of artemether/lumefantrine
and, if required, a single dose of primaquine (45 mg) to clear all parasites at the end of the study.
Parasitaemia in the volunteers were monitored at least daily since the inoculation.

The data contain the measurements of total parasitaemia (total circulating asexual parasites and
gametocytes per mL blood), asexual parasitaemia (circulating asexual parasites per mL blood),
gametocytaemia (circulating female and male gametocytes per mL blood) and plasma
concentration of PQP, at multiple time points after inoculation (see Figure 1—3 and Figure S2 in
the Supplementary Information). Further details about the VIS are given in[7]. We emphasize that
the data used in model fitting is the total parasitaemia (from the first measurement to the time

before any treatment other than PQP) and the other data (i.e. asexual parasitaemia and
gametocytaemia) are used to validate the model.

Gametocyte dynamics model

The model extends the published models of asexual parasite life cycle[18,25] by incorporating the
development of gametocytes. The model is comprised of three parts describing three populations
of parasites: asexual parasites (P), sexual parasites (P_G) and gametocytes (G). Note that
biologically speaking the phrase "sexual parasites" includes gametocytes, but in order to model
the sexual parasites which are indistinguishable from asexual parasites under a microscope, we
treat those sexual parasites and gametocytes as two different populations. A schematic diagram
of the development of those populations based on current knowledge[11,22] is shown in Figure 5.

Figure 5: Schematic diagram showing the model compartments and transitions. Square
compartments in the inner loop represent the asexual parasite population which
follows a life cycle of maturation and replication every a_L hours. Sexual commitment
occurs from age a_s and a fraction of asexual parasites become sexual (the bigger
square compartments in the outer loop) and eventually enter the development of stage

I—V gametocytes (the round compartments). The compartments with a dashed

boundary are sequestered to tissues and thus not measurable in a blood smear. The

notation of each compartment is consistent with those in model equations and is

explained in the main text.

Asexual parasites develop and replicate in the red blood cells (RBCs) till cell rupture at the end

of each life cycle and the released free parasites (merozoites) can initiate new life cycles if

successfully invading susceptible RBCs. At the time of inoculation (i.e. $t = 0$ hours in the

model), we define the inoculum size to be P_{init} and assume the age distribution of inoculated

parasites is Gaussian with mean μ and standard deviation σ. As time increments by one hour, the

asexual parasites of age a at time t (denoted as $P(a,t)$) follow the iterative equation:

$$P(a,t) = \begin{cases} P(a-1,t-1)e^{-\overline{k_d}-\delta_P}, & a = 2,3,\dots,a_L \\ r_P P(a_L,t-1)e^{-\overline{k_d}-\delta_P}, & a = 1 \end{cases} \qquad (1)$$

where $\overline{k_d}$ represents the average rate of asexual parasite killing by PQP and δ_P is the rate of

asexual parasite death due to processes other than PQP. $\overline{k_d}$ is approximated by the average of

$k_d(t-1)$ and $k_d(t)$ and $k_d(t) = k_{max}C(t)^\gamma/(C(t)^\gamma + EC_{50}{}^\gamma)$ where k_{max} is the maximum

killing rate and EC_{50} is the PQP concentration at which half maximum killing is achieved. $C(t)$

is the PQP concentration which is simulated by a pharmacokinetic model introduced below. a_L is

the length of each life cycle and r_P is the parasite replication rate indicating the average number

of newly infected RBCs attributable to the rupture of a single infected RBC at the end of its life

cycle. Sexual commitment is assumed to occur at the first age of the trophozoite stage (denoted

to be a_s) and a percentage (f) of asexual parasites leave asexual life cycle and start sexual

development in the next hour, which is modelled by

$$P(a_s+1,t) = (1-f)P(a_s,t-1)e^{-\overline{k_d}-\delta_P}, \qquad (2)$$

$$P_G(a_s+1,t) = fP(a_s,t-1)e^{-\overline{k_d}-\delta_P}. \qquad (3)$$

The first equation describes the proportion of parasites remaining in the asexual life cycle while

the second equation describes the proportion of parasites becoming sexual parasites (P_G).

According to Figure 5, the sexual parasites continue the rest of the life cycle and a part of the

next life cycle (note that they appear indistinguishable from asexual parasites under microscopy)

before becoming stage I gametocytes. The process is modelled by

$$346 \quad P_G(a,t) = \begin{cases} P_G(a-1, t-1)e^{-\delta_P}, & a = 2, 3, \dots, a_L \text{ except } a = a_s \text{ and } a = a_s + 1 \\ r_P P_G(a_L, t-1)e^{-\delta_P}, & a = 1 \end{cases} \quad (4)$$

347   Note that no $\overline{k_d}$ for P_G because PQP does not kill sexual parasites and gametocytes. The changes

348   of the sequestered stage I—IV gametocytes (G_1—G_4) are governed by difference equations

$$349 \quad G_1(t) = G_1(t-1)e^{-(m+\delta_G)} + \frac{P_G(a_s - 1, t-1)e^{-\delta_P}\left(1 - e^{-(m+\delta_G)}\right)}{m + \delta_G}, \quad (5)$$

$$350 \quad G_2(t) = G_2(t-1)e^{-(m+\delta_G)} + \frac{mG_1(t-1)\left(1 - e^{-(m+\delta_G)}\right)}{m + \delta_G}, \quad (6)$$

$$351 \quad G_3(t) = G_3(t-1)e^{-(m+\delta_G)} + \frac{mG_2(t-1)\left(1 - e^{-(m+\delta_G)}\right)}{m + \delta_G}, \quad (7)$$

$$352 \quad G_4(t) = G_4(t-1)e^{-(m+\delta_G)} + \frac{mG_3(t-1)\left(1 - e^{-(m+\delta_G)}\right)}{m + \delta_G}, \quad (8)$$

353   where m is the rate of gametocyte maturation and δ_G is the death rate of sequestered

354   gametocytes. Stage V gametocytes are circulating in bloodstream (and therefore can be measured

355   from the peripheral blood film) modelled by

$$356 \quad G_5(t) = G_5(t-1)e^{-\delta_{Gm}} + \frac{mG_4(t-1)\left(1 - e^{-\delta_{Gm}}\right)}{\delta_{Gm}}, \quad (9)$$

where δ_{Gm} is the death rate of mature circulating gametocytes.

The total parasitaemia in the model is given by $\sum_{a=1}^{a=a_s-1}[P(a,t) + P_G(a,t)] + G_5(t)$, which was

fitted to the VIS data. After model fitting, we simulated the asexual parasitaemia $\sum_{a=1}^{a=a_s-1} P(a,t)$

and gametocytaemia $G_5(t)$ and compared them with associated data for model validation. Table

2 presents all the model parameters and their units and descriptions.

Table 2: Details of the gametocyte dynamics model parameters. The parameter ranges of the

prior distributions are chosen based on previous estimates[25] and biological plausibility. We

assumed parasites younger than 25h are circulating and thus fix a_s to be 25h. For 3D7 strain, the

life cycle is approximately 39—45h (based on *in vitro* estimates[26] and *in vivo* estimates by

Wockner et al.[17]) and we fix a_L to be 42h. The boundaries of the uniform prior distribution (U)

also indicate the lower and upper bounds of the corresponding parameters.

Parameter	Unit	Description	Prior distribution

P_{init}	parasites/mL	inoculation size	U(0, 10)
μ	h	mean of the initial parasite age distribution	U(0, 35)
σ	h	SD of the initial parasite age distribution	U(0, 20)
r_P	(unitless)	parasite replication rate	U(0, 100)
k_{max}	h^{-1}	maximum rate of parasite killing by PQP	U(0, 1)
EC_{50}	ng/mL	half-maximum effective PQP concentration	U(1, 100)
γ	(unitless)	Hill coefficient for PQP	U(0, 20)
f	(unitless)	sexual commitment rate (in percentage)	U(0, 1)
δ_P	h^{-1}	death rate of asexual and sexual parasites	U(0, 0.2)
m	h^{-1}	maturation rate of gametocytes	U(0, 0.1)
δ_G	h^{-1}	death rate of sequestered gametocytes	U(0, 0.1)
δ_{Gm}	h^{-1}	death rate of circulating gametocytes	U(0, 0.1)
a_s	h	sequestration age of asexual parasites	fixed to be 25
a_L	h	length of life cycle of asexual parasites	fixed to be 42

368 SD: standard deviation; h: hour.

Pharmacokinetic model of piperaquine (PQP)

In the within-host model, the killing rate $k_d(t)$ is determined by PQP concentration $C(t)$ which

was simulated from a pharmacokinetic (PK) model introduced in this section. The PK model,

provided by Thanaporn Wattanakul and Joel Tarning (Mahidol-Oxford Tropical Medicine

Research Unit, Bangkok), is a three-compartment disposition model with two transit

compartments for absorption (see schematic diagram in Figure S3, Supplementary Information).

Based on Figure S3, the model is formulated to be a system of ordinary differential equations:

376
$$\frac{dD}{dt} = -k_T D, \tag{10}$$

377
$$\frac{dT_1}{dt} = k_T D - k_T T_1, \tag{11}$$

378
$$\frac{dT_2}{dt} = k_T T_1 - k_T T_2, \tag{12}$$

379
$$\frac{dC}{dt} = \frac{k_T T_2 + q_1 P_1 + q_2 P_2 - q_1 C - q_2 C - q_c C}{V_c}, \tag{13}$$

$$380 \quad \frac{dP_1}{dt} = \frac{q_1 C - q_1 P_1}{V_1} \, , \tag{14}$$

$$381 \quad \frac{dP_2}{dt} = \frac{q_2 C - q_2 P_2}{V_2} \, , \tag{15}$$

The model parameters were obtained by fitting the model output $C(t)$ to the PQP concentration

data from the VIS in MATLAB (version 2016b; The MathWorks, Natick, MA) using *lsqcurvefit*.

The MATLAB code (with detailed comments) are publicly available at

https://doi.org/10.26188/5cde4c26c8201. Note that for volunteers whose number of data points is

less than the number of parameters in the PK model (such that optimization fails), the parameter

values are given by the estimates from another VIS (provided by Thanaporn Wattanakul and Joel

Tarning). The best-fit curve and associated parameter values for all volunteers are provided in

Figure S2 and Table S1 in the Supplementary Information.

390   Fitting the model to parasitaemia data

We took a Bayesian hierarchical modelling approach[27] to fit the gametocyte dynamics model to

the data from all 17 volunteers. In detail, each volunteer has 12 model parameters (i.e. those in

Table 2 except a_s and a_L; also called the individual parameters) and lower and upper bounds of

the parameters are given in Table 2. If denoting the individual parameters to be θ_{ind} and lower

and upper bounds to be b_L and b_U respectively, the following transformations are used to convert

the bounded individual parameters to unbounded ones (denoted by φ_{ind}) in order to in order to

improve computational efficiency[28,29]:

$$398 \quad \varphi_{ind} = \ln\left(\frac{\theta_{ind} - b_L}{b_U - \theta_{ind}}\right), \tag{16}$$

399   φ_{ind} obeys a multivariate normal distribution $\mathcal{N}(\varphi_{pop}, \Omega_{pop})$ where

$$400 \quad \varphi_{pop} = \ln\left(\frac{\theta_{pop} - b_L}{b_U - \theta_{pop}}\right) \tag{17}$$

and θ_{pop} is a vector containing 12 population mean parameters (hyperparameters) corresponding

to the 12 gametocyte dynamics model parameters. Ω_{pop} is the covariance matrix. For more

efficient sampling process, $\varphi_{ind} \sim \mathcal{N}(\varphi_{pop}, \Omega_{pop})$ was reparameterised to a non-centred form

$\varphi_{ind} = \varphi_{pop} + \omega_{pop} L \eta$, where ω_{pop} is the diagonal standard deviation (SD) matrix whose

diagonal elements are the 12 population SD parameters (hyperparameters); L is the lower

Cholesky factor of the correlation matrix; η obeys standard multivariate normal distribution.

Note that $\Omega_{pop} = \omega_{pop}LL^T\omega_{pop}$ where LL^T is the correlation matrix. The prior distributions for

the 12 population mean parameters θ_{pop} are given by uniform distributions with bounds given in

Table 2. The prior distribution for the 12 population SD parameters is standard half-normal and

the prior distribution for the lower Cholesky factor of the correlation matrix L is given by

Cholesky LKJ correlation distribution with shape parameter of 2[29,30]. The distribution of the

observed parasitaemia measurements is assumed to be a lognormal distribution with mean given

by the model-simulated values and SD parameter with prior distribution of a half-Cauchy

distribution with a location parameter of zero and a scale parameter of 5. The distribution for the

observed parasitaemia measurements was used to calculate the likelihood function and the M3

method[31] was used to penalise the likelihood for data points below the limit of detection for the

total parasitaemia (10 parasites/mL[7]).

Model fitting was implemented in R (version 3.2.3)[32] and Stan (RStan 2.17.3)[29] using the

Hamiltonian Monte Carlo (HMC) optimized by the No-U-Turn Sampler (NUTS) to draw

samples from the joint posterior distribution of the parameters including the individual

parameters (12 parameters for each volunteers) and population mean parameters (12

hyperparameters). Five chains with different starting points (set by different random seeds) were

implemented and 1000 posterior samples retained from each chain after a burn-in of 1000

iterations (in total 5000 samples were drawn from the joint posterior distribution). The marginal

posterior and prior distributions of the population mean and SD parameters are shown in Figure

S4 and S5 in the Supplementary Information. The marginal posterior distributions of the

individual parameters for all 17 volunteers are shown in Figure S6—S17 (using violin plots) in

the Supplementary Information. For each volunteer, the 5000 sets of individual parameters are

used to simulate the gametocyte population dynamics model and generate 5000 simulated model

outputs (e.g. 5000 time series of total parasitaemia, asexual parasitaemia or gametocytaemia).

The posterior prediction and 95% prediction interval (PI) are given by the median and quantiles

of 2.5% and 97.5% of the 5000 model outputs at each time respectively (see Figure 1—3 for

example).

The estimates of some key biological parameters (Table 1) were calculated using the 5000

posterior draws of the 12 population mean parameters, i.e. median and 2.5%- and 97.5%-quantile

(95% credible interval). The sexual commitment rate was calculated directly by f_{pop} (the

population mean parameter for f). Sequestered and circulating gametocyte lifespan were calculated by $1/\delta_{G_{pop}}/24$ and $1/\delta_{Gm_{pop}}/24$ respectively (the factor of 24 converts hours into days). Gametocyte sequestration time was calculated by $4/m_{pop}/24$ where 4 indicates four sequestered state (stage I to IV) and 24 converts hours into days. Parasite multiplication factor is calculated by $r_{P_{pop}}\exp(-\delta_{P_{pop}}a_L)(1-f_{pop})$ where the term $\exp(-\delta_{P_{pop}}a_L)(1-f_{pop})$ gives the fraction of surviving asexual parasites after death and sexual conversion per life cycle.

The gametocyte dynamics model with parameters given by the median estimates of the population mean parameters was used to simulate the two scenarios predicting the dependence of human-to-mosquito transmissibility on the sexual commitment rate and gametocyte sequestration time (Figure 4).

Final analysis and visualization were performed in MATLAB. All computer codes (R, Stan, MATLAB), data and fitting results (CSV and MAT files) and an instruction document (note that reading the document first will make the code much easy to follow) are publicly available at https://doi.org/10.26188/5cde4c26c8201.

References

1. The World Health Organization. *World Malaria Report 2018*. (2018).

2. malERA: An updated research agenda for diagnostics, drugs, vaccines, and vector control in malaria elimination and eradication. *PLoS Med.* **14**, e1002455 (2017).

3. Bousema, T. & Drakeley, C. Epidemiology and infectivity of Plasmodium falciparum and Plasmodium vivax gametocytes in relation to malaria control and elimination. *Clin. Microbiol. Rev.* **24**, 377–410 (2011).

4. Josling, G. A. & Llinás, M. Sexual development in Plasmodium parasites: Knowing when it's time to commit. *Nat. Rev. Microbiol.* **13**, 573–587 (2015).

5. Churcher, T. S. *et al.* Predicting mosquito infection from Plasmodium falciparum gametocyte density and estimating the reservoir of infection. *Elife* **2**, e00626 (2013).

6. Bradley, J. *et al.* Predicting the likelihood and intensity of mosquito infection from sex specific Plasmodium falciparum gametocyte density. *Elife* **7**, e34463 (2018).

7. Collins, K. A. *et al.* A controlled human malaria infection model enabling evaluation of transmission-blocking interventions. *J. Clin. Invest.* **128**, 1551–1562 (2018).

466   8.   Diebner, H. H. *et al.* Modelling the transition of asexual blood stages of Plasmodium

467   falciparum to gametocytes. *J. Theor. Biol.* **202**, 113–127 (2000).

468   9.   Eichner, M. *et al.* Genesis, sequestration and survival of Plasmodium falciparum

469   gametocytes: Parameter estimates from fitting a model to malariatherapy data. *Trans. R.*

470   *Soc. Trop. Med. Hyg.* **95**, 497–501 (2001).

471   10.   Rockett, R. J. *et al.* A real-time, quantitative PCR method using hydrolysis probes for the

472   monitoring of Plasmodium falciparum load in experimentally infected human volunteers.

473   *Malar. J.* **10**, 48 (2011).

474   11.   Filarsky, M. *et al.* GDV1 induces sexual commitment of malaria parasites by antagonizing

475   HP1-dependent gene silencing. *Science.* **359**, 1259–1263 (2018).

476   12.   Gebru, T., Lalremruata, A., Kremsner, P. G., Mordmüller, B. & Held, J. Life-span of in

477   vitro differentiated Plasmodium falciparum gametocytes. *Malar. J.* **16**, 330 (2017).

478   13.   McCarthy, J. S. *et al.* A Pilot Randomised Trial of Induced Blood-Stage Plasmodium

479   falciparum Infections in Healthy Volunteers for Testing Efficacy of New Antimalarial

480   Drugs. *PLoS One* **6**, e21914 (2011).

481   14.   Khoury, D. S. *et al.* Within-host modeling of blood-stage malaria. *Immunol. Rev.* **285**,

482   168–193 (2018).

483   15.   Piray, P., Dezfouli, A., Heskes, T., Frank, M. J. & Daw, N. D. Hierarchical Bayesian

484   inference for concurrent model fitting and comparison for group studies. *bioRxiv* 393561

485   (2018). doi:10.1101/393561

486   16.   Wockner, L. F., Hoffmann, I., O'Rourke, P., McCarthy, J. S. & Marquart, L. Comparison

487   of statistical models to estimate parasite growth rate in the induced blood stage malaria

488   model. *Malar. J.* **16**, 352 (2017).

489   17.   Wockner, L. F. *et al.* In vivo growth rate of Plasmodium falciparum: analysis of parasite

490   growth data from malaria volunteer infection studies. *Submitted*

491   18.   Saralamba, S. *et al.* Intrahost modeling of artemisinin resistance in Plasmodium

492   falciparum. *Proc. Natl. Acad. Sci.* **108**, 397–402 (2010).

493   19.   Tibúrcio, M. *et al.* Specific expression and export of the Plasmodium falciparum

494   Gametocyte EXported Protein-5 marks the gametocyte ring stage. *Malar. J.* **14**, 334

495   (2015).

20.   Da, D. F. *et al.* Experimental study of the relationship between Plasmodium gametocyte

density and infection success in mosquitoes; implications for the evaluation of malaria

transmission-reducing interventions. *Exp. Parasitol.* **149**, 74–83 (2015).

21.   Schneider, P. *et al.* Adaptive plasticity in the gametocyte conversion rate of malaria

parasites. *PLOS Pathog.* **14**, e1007371 (2018).

22.   Bancells, C. *et al.* Revisiting the initial steps of sexual development in the malaria parasite

Plasmodium falciparum. *Nat. Microbiol.* **4**, 144–154 (2019).

23.   Dogovski, C. *et al.* Targeting the Cell Stress Response of Plasmodium falciparum to

Overcome Artemisinin Resistance. *PLOS Biol.* **13**, e1002132 (2015).

24.   Cao, P. *et al.* A mechanistic model quantifies artemisinin-induced parasite growth

retardation in blood-stage Plasmodium falciparum infection. *J. Theor. Biol.* **430**, 117–127

(2017).

25.   Zaloumis, S. *et al.* Assessing the utility of an anti-malarial pharmacokinetic-

pharmacodynamic model for aiding drug clinical development. *Malar. J.* **11**, 303 (2012).

26.   Duffy, S. & Avery, V. M. Plasmodium falciparum in vitro continuous culture conditions:

A comparison of parasite susceptibility and tolerance to anti-malarial drugs throughout the

asexual intra-erythrocytic life cycle. *Int. J. Parasitol. Drugs Drug Resist.* **7**, 295–302

(2017).

27.   Gelman, A. *et al. Bayesian Data Analysis, Third Edition (Texts in Statistical Science).*

(Taylor & Francis, 2013).

28.   Lesaffre, E., Rizopoulos, D. & Tsonaka, R. The logistic transform for bounded outcome

scores. *Biostatistics* **8**, 72–85 (2007).

29.   Stan Development Team. *Stan Modeling Language Users Guide and Reference Manual.*

(2017).

30.   Lewandowski, D., Kurowicka, D. & Joe, H. Generating random correlation matrices based

on vines and extended onion method. *J. Multivar. Anal.* **100**, 1989–2001 (2009).

31.   Ahn, J. E., Karlsson, M. O., Dunne, A. & Ludden, T. M. Likelihood based approaches to

handling data below the quantification limit using NONMEM VI. *J. Pharmacokinet.*

Pharmacodyn. **35**, 401–421 (2008).

32.   R Core Team. R: A language and environment for statistical computing. http://www.R-

project.org/. *R Foundation for Statistical Computing, Vienna, Austria* (2017).

Acknowledgments

We acknowledge useful conversations with David S. Khoury, Deborah Cromer and Miles P. Davenport (Kirby Institute, UNSW Australia, Sydney, Australia) and assistance in drafting the manuscript from Laura Cascales (QIMR Berghofer, Brisbane, Australia). We thank Jörg J. Möhrle, Head of Translation from Medicines for Malaria Venture for his support and permission to use the trial data. The work is supported by an Australian Research Council (ARC) Discovery project (DP170103076) and a National Health and Medical Research Centre (NHMRC) of Australia Project Grant (1025319) and supported in part by the Australian Centre for Research Excellence on Malaria Elimination, funded by the NHMRC (1134989). SGZ is funded by an ARC Discovery Early Career Researcher Award (170100785). JAS is supported by a NHMRC Senior Research Fellowship (1104975). This research was supported by use of the Nectar Research Cloud, a collaborative Australian research platform supported by the National Collaborative Research Infrastructure Strategy (NCRIS).

Author contributions

P.C., J.A.S., J.S.M. and J.M.M conceived the study; K.A.C. and J.S.M. provided the clinical data; P.C. analyzed the data with contributions from K.A.C., J.A.S., J.S.M. and J.M.M.; P.C. developed the mathematical model; S.Z. developed the statistical method; T.W. and J.T. provided the pharmacokinetic model for piperaquine; P.C. performed model fitting, simulation and analysis; P.C. wrote the first draft of the manuscript; All authors reviewed and commented on the manuscript.

Competing interests

We have no competing interests.