1   **Competition among host-specific lineages of *Poecilochirus***
2   ***carabi* mites influences the extent of co-adaptation with their**
3   ***Nicrophorus vespilloides* burying beetle hosts**
4
5

Authors: Syuan-Jyun Sun[1,2,*] and Rebecca M. Kilner[1]
Affiliations:
[1]Department of Zoology, University of Cambridge, Downing Street, Cambridge, CB2
3EJ, UK
[2]International Degree Program in Climate Change and Sustainable Development,
National Taiwan University, Taipei 10617, Taiwan
12
13 **Corresponding author:** Syuan-Jyun Sun; sjs243@ntu.edu.tw

14
15 **Keywords:** host-parasite interactions, population divergence, reciprocal transplant,
16 local adaptation, burying beetles, mites
17
18
19
20

Abstract:

Reciprocal selection between symbiotic organisms and their hosts can generate

variations in local adaptation between them. Symbionts often form species complexes

with lineages partially adapted to various hosts. However, it is unclear how

interactions among these lineages influences geographic variation in host-symbiont

local adaptation. We addressed this shortcoming with experiments on burying beetles

Nicrophorus vespilloides and their specialist phoretic mite *Poecilochirus carabi* in

two adjacent woodlands. Burying beetles transport these mites to vertebrate carrion

upon which they both reproduce. *P. carabi* appears to be a species complex, with

distinct lineages that specialise on breeding alongside different *Nicrophorus* species.

We found that in one wood (Gamlingay Woods), *N. vespilloides* carries a mixture of

mite lineages, with each lineage corresponding to one of the four *Nicrophorus* species

that inhabits this wood. However, two burying beetle species coexist in neighbouring

Waresley Woods and here *N. vespilloides* predominantly carries the mite lineage that

favours *N. vespilloides*. Mite lineage mixing alters the degree of local adaptation for

both *N. vespilloides* and the *P. carabi* mites, affecting reproductive success variably

across different woodlands. In Gamlingay, mite lineage mixing reduced *N.*

vespilloides reproductive success, while experimentally purifying mites lineage

enhanced it. The near pure lineage of *vespilloides* mites negligibly impacted Waresley

N. vespilloides. Mite reproductive success varied with host specificity: Gamlingay

mites had greatest reproductive success on Gamlingay beetles, and performed less

well with Waresley beetles. By contrast, Waresley mites had consistent reproductive

success, regardless of beetle's woodland of origin. We conclude that there is some

evidence that *N. vespilloides* and its specific mite lineage have co-adapted. However,

neither *N. vespilloides* nor its mite lineage adapted to breed alongside other mite

lineages. This, we suggest, causes variation between Waresley and Gaminglay Woods

in the extent of local adaptation between *N. vespilloides* beetles and their *P. carabi*

mites.

Introduction

Ever since Darwin (1859), evidence has been gathering that natural selection causes populations to adapt in different ways to their different local environments. More recent work suggests that adaptation can happen on very local scales, even when there is still gene flow between populations (e.g. Sun et al., 2020; J. N. Thompson, 2013). Nevertheless, the extent of local adaptation varies across landscapes, between populations of the same species (e.g. J. N. Thompson, 1999, 2005). This is especially true for adaptations that arise from co-evolutionary interactions between symbiotic species (by which we mean 'intimately co-existing' species), which are engaged in reciprocal selection. In specialist antagonistic interactions, or obligate mutualisms, each party exerts such strong selection on the other that there can be little differentiation between populations (e.g. et al., 2004; Johnson et al., 2010; Lively & Dybdahl, 2000). A key challenge is to understand the ecological factors that generate such geographical mosaics of co-evolution and co-adaptation (Thompson, 2013).

Environmental differences among populations are known to cause variation in the outcome of any species interaction. These might arise from variation in the availability of an essential resource (e.g. Johnson et al., 2010), or the presence of a common enemy species (e.g. Hopkins et al., 2017; Sun & Kilner, 2020), or differences in the abiotic environment (e.g.Kersch & Fonseca, 2005). Theoretical analyses suggest that this variation, in turn, can cause dramatic differences in the trajectory of local co-evolution (e.g. Nuismer et al., 2000). Therefore, determining ecological correlates of variation in the extent of reciprocal selection between the same two partner species can potentially explain why there is geographical variation in the pattern of local adaptation among populations (e.g. Garrido et al., 2012; Gorter et al., 2016; Johnson et al., 2010).

However, in reality, interactions between species rarely involve just two partner species. For example, one insect species commonly pollinates more than one plant species, while each plant species can be pollinated by more than one insect species (e.g. J. N. Thompson, 2013). Similarly, one cactus species can be in a protective mutualism with multiple species of ants (Ness et al., 2006), while multiple bumblebee species *Bombus* spp. commonly interact with multiple mite species (Haas et al., 2019). Likewise, host species richness and abundance are positively correlated with parasite species richness (e.g. Hechinger & Lafferty, 2005). Multispecies associations are likely to generate variation in the strength of reciprocal selection. For example, multiple infection of different parasite species or strains can differentially influence

the fitness among different host species interacting in the same community,

depending on the susceptibility and tolerance of each host species (Friesen et al.,

2017). Therefore understanding interactions among different strains or species of

symbionts is likely to help account for geographical variation in the pattern of local

co-adaptation between symbionts and their hosts.

Here we apply this rationale to understanding the extent of local co-adaptation

between burying beetles *Nicrophorus vespilloides* (Coleoptera: Silphidae) and the

species complex of *Poecilochirus carabi* mites with whom they interact

(Mesostigmata: Parasitidae). Both the burying beetle and the mite require small

vertebrate carrion to breed, but only the beetle is capable of flight. The mites travel

with the beetle between breeding opportunities whilst sexually immature

deutonymphs, attached as benign passengers to the adult beetle's thorax. When they

locate a dead body, the adult beetles convert the carrion into an edible nest for their

young, and then stay with their brood to defend and nourish them. Meanwhile, the

mites disembark, moult into the adult stage, mate, and produce offspring alongside the

beetle larvae (Schwarz & Müller, 1992). Breeding alongside mites can be costly for

beetles (De Gasperin et al., 2015; De Gasperin & Kilner, 2015; Nehring et al., 2019),

which select adaptations in the beetles to mitigate these costs. When their duties of

care are complete, adult beetles fly off in search of new reproductive opportunities,

carrying with them the next generation of mites. The timing of the mite's lifecycle is

therefore closely aligned with the duration of beetle parental care (Schwarz & Müller,

1992).

P. carabi exists as a species complex because it comprises distinct lineages of

mites that are each specialised to breed on different species of burying beetle

Nicrophorus spp (Nehring et al., 2017). Each burying beetle species differs slightly in

its duration of care. This has favoured divergent adaptation in the mite populations

associated with each species of burying beetle, which in turn has generated distinct

mite lineages (Canitz et al., 2021; Müller & Schwarz, 1990; Wilson, 1982).

Nevertheless, mite lineages can still interbreed and cannot be told apart

morphologically. The key way to distinguish them is behaviourally, through their

relative preference for different burying beetle species (Brown & Wilson, 1992;

Schwarz, 1996; Wilson, 1982). Previous work suggests that when sympatric

Nicrophorus species do not differ much in their duration of care, then each species of

burying beetles carries a mixture of the different mite lineages associated with each of

the sympatric beetles and the mites hybridise across lineages (Brown & Wilson, 1992).

However, in populations where *Nicrophorus* species differ considerably in their

duration of care (Brown & Wilson, 1992; Scott, 1998), especially when larger beetle

species have relatively longer larval development time than smaller beetle species,

each species is more likely to carry its own specialist mite lineage and each mite

lineage is more likely to be reproductively isolated (Brown & Wilson, 1992).

We focused on *N. vespilloides* and *P. carabi* inhabiting two neighbouring yet

geographically isolated populations (Gamlingay and Waresley Woods) in

Cambridgeshire, UK. Previous studies have shown that there are differences in the

Nicrophorus guild inhabiting each woodland (Sun et al., 2020). Both woods contain

the smallest UK burying beetle *N. vespilloides* and the larger burying beetle *N.*

humator. However, only Gamlingay Woods is additionally routinely inhabited by the

intermediate-sized *N. interruptus* and *N. investigator*. We tested the following

predictions that:

(1) *N. vespilloides* from Gamlingay Woods should be more likely to carry a

mixture of mite lineages, whereas *N. vespilloides* from Waresley Woods should be

more likely to carry the *N. vespilloides* lineage of mites, reflecting the different

species that routinely co-exist in each wood. To test this prediction, we conducted two

consecutive choice experiments for mites to choose among all four *Nicrophorus*

beetles in the first generation of *P. carabi*, and then bred these mites based on their

choice. Their offspring were tested again to evaluate the consistency in choice and the

extent to which each mite lineage was mixed in each woodland.

(2) *N. vespilloides* should have lower reproductive success when breeding

alongside a mixture of *P. carabi* lineages than when breeding with its specialist *P.*

carabi lineage, with whom the beetle is more likely to be co-adapted. We

experimentally manipulated the mite composition, either mixing all four lineages of *P.*

carabi or allowing the pure *N. vespilloides* lineage of *P. carabi* to breed alongside *N.*

vespilloides. At dispersal, we determined the reproductive success of both beetles and

mites.

(3) *N.vespilloides* and *P. carabi* should be divergently co-adapted in Gamlingay

and Waresley woods, if they differ in whether they carry pure versus mixed lineages

of mites. Specifically, we predicted that *N. vespilloides* and its specialist lineage of

mites should be locally adapted to each other. We tested this prediction using

reciprocal transplant experiments by exposing mites from either Gamlingay Woods or

Waresley Woods to *N. vespilloides* beetles from Gamlingay and Waresley Woods.

Just as before, we determined the reproductive success of both beetles and mites at
dispersal.
156
Methods
Study species
Gamlingay and Waresley Woods are fragments of the Wild Wood that covered
England until deforestation (c. 1000-4000 years ago), and are roughly 2.5km apart
(Sun et al., 2020). *N. vespilloides* are the most abundant burying beetles in each wood
(81.6% in Gamlingay Woods and 93.7% in Waresley Woods (Sun et al., 2020)), and
the *P. carabi* mite species complex is the most commonly found mite species
associating with *Nicrophorus* beetles (Schwarz et al., 1998).
165
Field observations
Surveys of burying beetle communities were conducted in Gamlingay (Latitude:
52.15555°; Longitude: −0.19286°) and Waresley (Latitude: 52.17487°; Longitude:
−0.17354°) Woods in Cambridgeshire, UK. From June to October in 2016-2017, five
traps at each site were baited with mouse carcasses and hung in the same location,
separated by at least 150 m. We checked the traps every 2-3 weeks and collected all
Nicrophorus spp. under permit. Traps were then refilled with fresh mice at each
collection. The number and sex of each beetle species were recorded at each location
for each trapping event. Each beetle's pronotum width was measured to the nearest
0.01 mm as a standardised measurement of body size (Jarrett et al., 2017). We
separated mites from their beetle hosts under anaesthetic by CO_2. The data presented
here are related to those from an earlier published study within this system that
focussed only on the correlation between body size and mite load in *N. vespilloides* in
2017 (Sun et al., 2019). In the current study, we have used a more extensive dataset to
thoroughly investigate the mite load on each *Nicrophorus* spp. during 2016-2017.
181
182 **Origin and maintenance of burying beetles and mites**
183 Both species were kept under laboratory conditions at $21 \pm 2°C$ and on a 16:8 light to
184 dark cycle.
185
Beetles were kept individually in plastic boxes (12cm x 8cm x 2cm) filled with moist
soil. Field-collected beetles were kept for at least two weeks before they were
subjected to experimentation to even out any differences in sexual maturity and

nutritional status. To breed beetles, *N. vespilloides* collected from the field sites were

paired on a mouse in a breeding box lined with damp soil. All breeding boxes were then

placed into cupboards to mimic underground environments. After eight days, we

collected the dispersing larvae and transferred them to eclosion boxes (10 x 10 x 2 cm,

25 compartments) filled with moist soil. At eclosion, each emerging beetle was moved

to a plastic container (12 x 8 x 2 cm) with moist soil. We fed beetles twice a week with

minced beef for 2-3 weeks until they were sexually mature.

Mites were maintained in distinct populations, according to their woodland of origin,

and kept apart from burying beetles except when breeding. To breed mites, each

month we transferred 15 mite deutonymphs chosen at random, and a pair of beetles

from the same population, to a new breeding box (17 x 12 x 6 cm with 2 cm of soil)

furnished with a fresh mouse carcass ($n = 10$ for each population). After breeding,

beetle parents and third-instar larvae were removed from the box. The mites remained

and were given another adult beetle, and thereafter supplied with minced beef twice a

week.

Prediction 1: Gamlingay N. vespilloides *carry a mixture of* P. carabi *lineages*

To investigate population differences in the number of mite lineages present in each

wood, we used consecutive choice experiments. *P. carabi* from either Gamlingay or

Waresley Woods were allowed to choose between one of the following field-collected

burying beetle species: *N. vespilloides*, *N. humator*, *N. interruptus*, and *N.*

investigator. Each species was represented by one individual, drawn at random from a

pool of field beetles (184 *N. vespilloides*, 98 *N. humator*, 129 *N. interruptus* and 100

N. investigator). Burying beetles in this pool were haphazardly chosen from four field

populations (Gamlingay Woods, Waresley Woods, Madingley Woods (Latitude:

52.22658°; Longitude: 0.04303°), and Thetford Forest (Latitude: 52.41286°;

Longitude: 0.75167°). This allowed us to remove any beetle population-level effect

on mite preferences, and to focus entirely on the effect of the beetle's species on

influencing mite preferences.

We used mites that had been bred for one generation in the lab so that the tested

mites had no prior experience which might influence their choice of beetle. Four

burying beetles (of the same sex, one from each species; $n = 43$ trials where beetles

were all males and 38 trials where beetles were all females) were introduced into a

plastic container (17 x 12 x 6 cm), around which they could move freely. At the same

time 50 mites were also introduced – either from the Gamlingay population or from

the Waresley population. It is unlikely that the mites hindered each other in attaching

to host beetles because wild-caught beetles can carry up to 200-300 mites individually,

even on *N. vespilloides*, the smallest of the UK species (Schwarz et al., 1998; Sun et

al., 2019). The container held moist soil to a depth of 2 cm and minced beef in *ad*

libitum quantities. All beetles and mites were starved for a day prior to the experiment

to increase the likelihood that they gathered together on the mince, to mimic the

natural context in which mites transfer between beetles hosts at feeding/ breeding

opportunities. The number of mites carried by each beetle species was recorded 24 h

later, and used to assess the mixing of different mite lineages in each population.

Mites that chose *N. vespilloides*, *N. humator*, *N. interruptus*, and *N. investigator* were

defined as P-ves, P-hum, P-int, and P-inv, respectively. We then bred these mites

separately on a fresh mouse carcass, with one mouse carcass for each lineage of mite

identified by the choice test. The offspring of these breedings were then tested again

for their burying beetle preferences, as a further test of extent to which the mite

lineages were mixed in each woodland. For this second choice test, we introduced 10

mites and one beetle from each of the four species of the same sex ($n = 92$ trials

where beetles were all males and 50 trials where beetles were all females) in a plastic

container, and counted the number of mites on each beetle after 24 h. We tested 10

mites in the second choice experiment because we harvested relatively few P-hum

and P-inv in the first choice experiment, and this meant there were fewer offspring

available for testing.

Prediction 2: A mixture of P. carabi lineages negatively affects N. vespilloides

These experiments were focused on *N. vespilloides* and *P. carabi* mites drawn from

Gamlingay Woods. We experimentally manipulated the composition of the mite

community (= a group of 10 deutonymphs) associated with each burying beetle,

generating two treatments: a) pure *N. vespilloides* lineage of *P. carabi* and b) a

mixture of all four lineages of *P. carabi*. In the mixture, we introduced 3 P-ves, 1

P-hum, 4 P-int, and 2 P-inv, based on the relative proportion of the four *P. carabi*

lineages from *Prediction 1*. The mites used were descendants of the second generation

of *P. carabi* from the experiment above, and lineages were determined from the

preferences they exhibited in the second generation of choice experiment. Ten mites

were introduced at beetle pairing, directly onto the carcass. Pairs of beetles were

sequentially assigned to one of the three mite treatments, introduced into a breeding

box (17 x 12 x 6 cm with 2 cm of soil) and given a 15-20 g (17.71 ± 0.16) mouse

carcass to breed upon. At larval dispersal, i.e. 8 days after pairing, we counted all

larvae and weighed the whole brood (to the nearest 0.1mg). We calculated the average

larval mass for each brood (total brood mass divided by the number of larvae). To

determine the reproductive success of mites, we used CO_2 to detach dispersing mite

deutonymphs from adult beetles, at the end of the breeding event.

Prediction 3: Populations have adapted divergently

To assess the extent of local adaptation, we adopted a fully factorial design of

experimental reciprocal mite infestation (e.g. Blanquart et al., 2013; Garrido et al.,

2012; Nuismer & Gandon, 2008). These experiments were carried out in three blocks.

Each beetle population (Gamlingay/Waresley) was infested with either 10 Gamlingay

mite deutonymphs or 10 Waresley mite deutonymphs. Mites were introduced directly

onto 15-20 g carcasses (16.97 ± 0.12 g), when beetles were paired. Pairs of beetles

were unrelated, to prevent inbreeding. We took the same measurements of beetle and

mite reproductive success as in the previous experiments, when larvae dispersed away

from the carcass 8 days after pairing.

Statistical analyses

We analysed the data using generalized linear mixed models (GLMM) with the glmer

function in the *lme4* package in R version 3.4.3 (R Development Core Team, 2014).

To obtain minimal adequate models, we applied a stepwise approach to exclude

non-significant variables and interactions (Crawley, 2007). We included block as a

random effect in all models, where appropriate. Tukey HSD tests were used for

post-hoc pairwise comparisons, as necessary, using the *lsmeans* package (Lenth,

2016). All data are provided in electronic supplementary material.

Prediction 1: Gamlingay N. vespilloides *carry a mixture of* P. carabi *lineages*

We analysed variation in the number of mites carried by each beetle species using a

Poisson model, with an offset of the log total number of mites allowed to make a

choice in each trial. We included mite population (Gamlingay/Waresley), beetle

species (*N. vespilloides*, *N. humator*, *N. interruptus*, *and N. investigator*), and their

interaction as explanatory variables. Sex and body size of beetles were included as

covariates. We also included as random factors the sampling year (2017 or 2018), and

trial ID. We analysed the data in a similar manner for the second choice experiment,

testing for consistency of beetle species preference among mites in the next

generation. Of the 16 possible outcomes this two generation experiment yielded (N =

4 species for mites to choose in Gen 1 x N = 4 species for mites to choose in Gen 2),

we analysed only those the mites that made the same choice of beetle species as their

parents (i.e. N = 4 outcomes).

Prediction 2: A mixture of P. carabi *lineages negatively affects* N. vespilloides

We used generalized linear models (GLM) to analyse three measures of beetle

reproductive success when exposed to different experimental combinations of mite

lineages: brood size (using a Poisson distribution), brood mass and average larval

mass (both using a Gaussian distribution). We included mite treatment as an

explanatory variable, and additionally included carcass mass as a covariate when

analysing brood size and brood mass. For analysis of average larval mass, we

included larval density (brood size divided by carcass mass) as a covariate. We also

analysed variation in mite reproductive success using a negative binomial GLM with

mite treatments and carcass mass as explanatory variables.

Prediction 3: Populations have adapted divergently

Local adaptation of beetles to mites

We analysed three measures of beetle reproductive success when exposed to different

mite populations using GLMMs: brood size, brood mass and average larval mass.

Beetle wood of origin (Gamlingay/Waresley), mite wood of origin

(Gamlingay/Waresley) and their interaction were included as explanatory variables.

We included carcass mass as a covariate for the analysis of brood size and brood mass,

whereas we included larval density (brood size divided by carcass mass) as a

covariate for the analysis of average larval mass. In all models, block was included as

a random factor.

Local adaptation of mites to beetles We analysed the number of dispersing mites

deutonymphs present at the end of each reproductive bout using a negative binomial

GLMM. Beetle wood of origin (Gamlingay/Waresley), mite wood of origin

(Gamlingay/Waresley), and their interaction were included as explanatory variables.

Carcass mass was included as a covariate, whereas block was included as a random

factor.

Results

Field observation

In total, 1465 *Nicrophorus* individuals were caught over the two sampling years (780 and 685 for Gamlingay and Waresley Woods, respectively), carrying a total of 17,249 *P. carabi* mite deutonymphs on four beetle species (Fig. 1). We found that the mite load on each *Nicrophorus* species varied differently between populations (beetle species x population interaction, $\chi^2 = 46.19$, d.f. = 3, $p < 0.001$; Fig. 1). Specifically, *N. vespilloides* from Gamlingay had an average lower number of mites compared to *N. interruptus* (*post-hoc* comparison: $z = -2.86$, $p = 0.022$), but carried similar number of mites compared to those of *N. humator* (*post-hoc* comparison: $z = -0.73$, $p = 0.886$) and *N. investigator* (*post-hoc* comparison: $z = -1.53$, $p = 0.418$). In Waresley Wood, however, *N. vespilloides* carried more mites than *N. humator* (*post-hoc* comparison: $z = 6.80$, $p < 0.001$), but we could detect no difference in the mite load carried by *N. vespilloides* and *N. interruptus* (*post-hoc* comparison: $z = 2.22$, $p = 0.118$), nor between *N. vespilloides* and *N. investigator* (*post-hoc* comparison: $z = 1.14$, $p = 0.663$). Moreover, comparing mite abundance on the beetle species, between woodlands, we found that Gamlingay *N. vespilloides* had lower number of mites than Waresley *N. vespilloides* (*post-hoc* comparison: $z = -4.18$, $p < 0.001$). In contrast, *N. humator* from Gamlingay had higher mite abundance than *N. humator* from Waresley (*post-hoc* comparison: $z = 4.45$, $p < 0.001$), and there was a tendency for Gamlingay *N. interruptus* to carry more mites than those from Waresley (*post-hoc* comparison: $z = 1.72$, $p = 0.086$). We could detect no significant difference between Gamlingay and Waresley in mite abundance on *N. investigator* (*post-hoc* comparison: $z = 0.90$, $p = 0.370$).

Prediction 1: Gamlingay *N. vespilloides* carry a mixture of *P. carabi* lineages

The first lab-bred generation of *P. carabi* differed significantly in their relative preference for the different *Nicrophorus* species, depending on whether they were derived originally from Gamlingay or Waresley Woods (mite population x beetle species interaction, $\chi^2 = 152.62$, d.f. = 3, $p < 0.001$; Fig. 2a; Table 1). *P. carabi* derived from Gamlingay Woods and Waresley Woods showed divergent preferences for the different *Nicrophorus* species (Gamlingay: $\chi^2 = 294.08$, d.f. = 3, $p < 0.001$; Waresley: $\chi^2 = 459.69$, d.f. = 3, $p < 0.001$). Gamlingay mites were similarly likely to favour *N. interruptus* and *N. vespilloides*. They favoured *N. investigator* less frequently and *N. humator* even less (Fig. 2a; Table 1). In contrast, Waresley *P.*

carabi showed a clear preference for *N. vespilloides*. Their next-favoured beetle

species was *N. interruptus*, followed by *N. investigator*, and then *N. humator* (Fig. 2a;

Table 1). We used *post-hoc* comparisons to compare the strength of the mite

preference for each *Nicrophorus* species between populations. We found that

Gamlingay mites showed stronger preference for *N. humator* ($z = 5.99$, $p < 0.001$) and

N. interruptus ($z = 5.45$, $p < 0.001$) than mites from Waresley. They were also slightly

inclined to favour *N. investigator*, but this pattern was not statistically significant ($z =$

1.93, $p = 0.054$). By contrast, Waresley mites showed a higher preference for *N.*

vespilloides than Gamlingay mites ($z = -9.04$, $p < 0.001$).

We bred mites that showed the same preference for burying beetle species, and

tested whether the preferences of the offspring matched those of their parents, to test

for indirect evidence that mites were segregating into genetic lineages, as inferred

from their host preference. The extent to which the beetle preferences aligned

between the generations varied by woodland (mite population x beetle species

interaction, $\chi^2 = 50.42$, d.f. $= 3$, $p < 0.001$). We found that Waresley P-ves mites

consistently had stronger preference for *N. vespilloides* than Gamlingay mites ($t =$

-2.17, $p = 0.030$; Fig. 2b). In contrast, both P-hum and P-int mites from Gamlingay

Woods showed stronger consistency in their choice of *N. humator* ($t = 5.48$, $p < 0.001$;

Fig. 2b) and *N. interruptus* ($t = 4.92$, $p < 0.001$; Fig. 2b), respectively, compared to

those from Waresley Woods. Gamlingay P-inv mites also shown a non-significant

tendency to prefer *N. investigator* compared to Waresley P-inv mites ($t = 1.78$, $p =$

0.075; Fig. 2b).

Prediction 2: A mixture of *P. carabi* lineages negatively affects *N. vespilloides*

We tested whether differential mixing of mite lineages in Gamlingay Woods caused

variations in beetle reproductive success. We created experimental mite communities,

manipulated to different degrees to contain mites with different beetle preferences.

We found that beetles breeding alongside mites that varied in their preference for

different beetle species produced smaller ($\chi^2 = 33.40$, d.f. $= 1$, $p < 0.001$; Fig. 3a) and

lighter broods ($\chi^2 = 5.39$, d.f. $= 1$, $p = 0.02$; Fig. 3b) than beetles breeding with a pure

population of *N. vespilloides*-specific mites. We found no effect of the experimental

mite community composition on average larval mass ($\chi^2 = 2.55$, d.f. $= 1$, $p = 0.110$;

Fig. 3c). The beetle preferences of the mites also explained variation in mite

reproductive success (Fig. 3d). Mites produced more offspring in a pure population of

N. vespilloides-specific *P. carabi* than when in a mixture of different *P. carabi*

'lineages' ($\chi^2 = 18.13$, d.f. = 1, $p < 0.001$).

Prediction 3: Populations have adapted divergently

1. *Local adaptation of beetles to mite populations*

Brood size

Beetles exposed to Gamlingay and Waresley mites produced a similar brood size (Fig.

4a; Table 2). In addition, beetles from Waresley Woods consistently produced larger

broods compared to those from Gamlingay Woods (Fig. 4a; Table 2).

Brood mass

We found that different mite populations affected burying beetle fitness in different

ways, depending on the burying beetle's woodland of origin (Table 2). In Gamlingay

Woods, we found that Gamlingay beetles breeding with Gamlingay mites produced

lighter broods compared to when they bred with Waresley mites (*post-hoc*

comparison, local mites v. foreign mites: $t = -2.28$, $p = 0.024$). Beetles from Waresley

Woods produced broods that did not differ significantly in mass whether they were

breeding alongside Waresley mites or Gamlingay mites (*post-hoc* comparison, local

mites v. foreign mites: $t = 0.76$, $p = 0.451$).

Average larval mass

We found that the woodland origin of the mites and of the beetles interacted in

affecting average larval mass (Fig. 4c; Table 2). When beetles bred alongside mites

from the same woodland population, their larvae were smaller than when they bred

alongside mites from the other woodland for Gamlingay beetles (*post-hoc* comparison,

local mites v. foreign mites: $t = -4.08$, $p < 0.001$) and for Waresley beetles (*post-hoc*

comparison, local mites v. foreign mites: $t = 2.35$, $p = 0.020$), but the extent of

reduction in average larval mass by breeding with local mites was greater in

Gamlingay beetles than in Waresley beetles (Table 2).

2. *Local adaptation of mites to beetle populations*

We found that the reproductive success of the mites depended on the woodland origin

of the beetles but not that of the mites. Gamlingay mites had greater reproductive

success when breeding alongside Gamlingay beetles rather than with Waresley

beetles (Fig. 4d; Table 2). Gamlingay and Waresley mites performed equally well on

Waresley beetles (Fig. 4d; Table 2).

Discussion

The mosaic theory of co-evolution (Thompson, 2005) suggests that populations differ

in the extent of co-adaptation between interacting species because the structure of

selection varies between populations; because the strength of reciprocal selection

varies between populations (Agrawal & Zhang, 2021); or because genetic variation

influences the capacity for co-adaptation in different populations (Whitham et al.,

2020); or some combination of all three of these factors. The novel contribution of

this study is to show how interactions among the lineages of one species are

themselves a source of selection, both within and between species (Sun et al., 2019)

(Table 3).

The differences we report here between Gamlingay and Waresley Woods can be

traced back to the contrast in the *Nicrophorus* guild that inhabits each wood (Sun et

al., 2020). In Gamlingay Woods, four *Nicrophorus* species co-exist. In support of

Prediction (1), we found that *N. vespilloides* carries a mixture of mite lineages from

all four *Nicrophorus* species. By contrast, in Waresley Woods, there are routinely

only two burying beetle species (Sun et al., 2020). Waresley derived mites displayed a

marked preference for associating with *N. vespilloides*, and this preference persisted

between generations. Therefore, we conclude that Waresley *N. vespilloides* carries a

near pure-bred lineage of P-ves mites. Perhaps *N. vespilloides* and *N. humator* differ

too much in the duration of parental care for mixed lineages to persist in Waresley

Woods (see Brown & Wilson 1992). This remains to be determined in future work.

Experimentally manipulating the lineages present in the mite community at each

breeding event revealed two ways in which the composition of the mite community

exerts selection. First, and unexpectedly, we found that mites themselves had lower

breeding success when breeding as a mixed community than when breeding as a pure

P-ves population. This suggests that the mite lineages impose selection on each other

and that each mite lineage is locally adapted to breed alongside its own kind, perhaps

through mechanisms that minimise local mate competition (Nehring & Müller, 2009).

Male mites in particular are known to be lethally competitive (Nehring & Müller,

2009) and it would be interesting to determine whether the incidence of death is

greater when the lineages are mixed.

Unexpectedly, we also found that *N. vespilloides* somehow modulates the
selection that the mixed mite lineages impose upon each other during reproduction.
Furthermore, the increase in mite reproductive success was more pronounced when
Gamlingay *N. vespilloides* were present than when Waresley *N. vespilloides* bred
alongside the mites, suggesting that the Gamlingay *P. carabi* mites have divergently
adapted to breed in association with Gamlingay *N. vespilloides*.

From the Gamlingay *N. vespilloides'* perspective, however, the increase in mite
reproductive success came at the cost of the beetle's reproductive success. Therefore
the second way in which the mixed community of mites is divergent from the pure
P-ves population lies in relative strength of selection it imposes on *N. vespilloides* – in
support of Prediction (2). Perhaps as a direct consequence of increased competition
among the mite lineages, or perhaps through other means, the mixed community more
negatively affected Gamlingay burying beetle reproductive success than did the pure
P-ves population of mites.

A key way in which mites reduce burying beetle reproductive success is by
consuming beetle eggs and newly hatched larvae (de Gasperin & Kilner 2015, Sun &
Kilner 2020 eLife). With greater numbers of mites breeding, potentially, in the mixed
mite communities (Nehring & Muller 2009), perhaps there was a greater incidence of
mite attack on beetle offspring. Consistent with this interpretation, we found that
Waresley *N. vespilloides* actually had higher reproductive success than Gamlingay *N.*
vespilloides when breeding alongside the mixed mite community. This could be
because Waresley females lay more eggs than Gamlingay females (Sun et al., 2020)
and so can better withstand the greater incidence of offspring attack by mites, as
evidenced by Waresley beetles consistently producing larger broods, irrespective of
mite presence and woodland of origin (Fig. 4a).

In summary, overall we find that *N. vespilloides* and the *N. vespilloides* mite
lineage have co-adapted to each other (Table 3). However, neither *N. vespilloides* nor
the *N. vespilloides* mite lineage is adapted to breed alongside other mite lineages.
Local adaptation probably involves some form of tolerance rather than any specific
defences against mite attacks on reproductive success (Svensson & Råberg, 2010).
Since Gamlingay *N. vespilloides* carries a mixture of mite lineages, the extent of local
adaptation between *N. vespilloides* beetles and their *P. carabi* mites is greater in
Waresley Woods than in Gamlingay Woods.

Animals are commonly hosts to diverse communities of symbionts. The general
conclusion we draw from our work is that variation in the extent of local adaptation

between symbionts and their hosts depends on two factors that have been relatively

overlooked: (1) the environmental effects that shape how these communities arise,

and (2) the nature of interactions among different strains of symbiont. Nevertheless,

each factor has been shown to be important in recent work on other species. For

example, in deep-sea hydrothermal vent snails, environmental conditions have been

found to have a greater effect than host genetic variation in determining symbiont

composition and adaptation to local geochemical conditions (Breusing et al., 2022).

Similarly, in pea aphids, cooperative relationships among symbiont strains have been

found to influence the nature of host-symbiont interactions and adaptations (Peng et

al., 2023). Whether these two factors are just as influential in influencing the extent of

local adaptation between other species of hosts and their symbionts remains to be

investigated.

Conflict of interest

The authors declare that there is no competing interest.

References

Agrawal, A. A., & Zhang, X. (2021). The evolution of coevolution in the study of
species interactions. *Evolution*, *75*(7), 1594–1606.
https://doi.org/10.1111/EVO.14293

Blanquart, F., Kaltz, O., Nuismer, S. L., & Gandon, S. (2013). A practical guide to
measuring local adaptation. *Ecology Letters*, *16*(9), 1195–1205.
https://doi.org/10.1111/ele.12150

Breusing, C., Genetti, M., Russell, S. L., Corbett-Detig, R. B., & Beinart, R. A. (2022).
Horizontal transmission enables flexible associations with locally adapted
symbiont strains in deep-sea hydrothermal vent symbioses. *Proceedings of the
National Academy of Sciences of the United States of America*, *119*(14).
https://doi.org/10.1073/PNAS.2115608119/-/DCSUPPLEMENTAL

Brown, J. M., & Wilson, D. S. (1992). Local specialization of phoretic mites on
sympatric carrion beetle hosts. *Ecology*, *73*(2), 463–478.
https://doi.org/10.2307/1940753

Canitz, J., Sikes, D. S., Knee, W., Baumann, J., Haftaro, P., Steinmetz, N., Nave, M.,
Eggert, A.-K., Hwang, W., & Nehring, V. (2021). Cryptic diversity within the
Poecilochirus carabi mite species complex phoretic on Nicrophorus burying

beetles: Phylogeny, biogeography, and host specificity. *Molecular Ecology*, *31*(2),

658–674. https://doi.org/10.1111/mec.16248

Crawley, M. J. (2007). *The R Book*. John Wiley & Sons, Ltd.

https://doi.org/10.1002/9780470515075

De Gasperin, O., Duarte, A., & Kilner, R. M. (2015). Interspecific interactions explain

variation in the duration of paternal care in the burying beetle. *Animal Behaviour*,

109, 199–207. https://doi.org/10.1016/j.anbehav.2015.08.014

De Gasperin, O., & Kilner, R. M. (2015). Friend or foe: inter-specific interactions and

conflicts of interest within the family. *Ecological Entomology*, *40*, 787–795.

https://doi.org/10.1111/een.12259

Forde, S. E., Thompson, J. N., & Bohannan, B. J. M. (2004). Adaptation varies through

space and time in a coevolving host–parasitoid interaction. *Nature*, *431*(7010),

841–844. https://doi.org/10.1038/nature02906

Friesen, O. C., Poulin, R., & Lagrue, C. (2017). Differential impacts of shared parasites

on fitness components among competing hosts. *Ecology and Evolution*, *7*(13),

4682–4693. https://doi.org/10.1002/ece3.3062

Garrido, E., Andraca-Gómez, G., & Fornoni, J. (2012). Local adaptation:

simultaneously considering herbivores and their host plants. *New Phytologist*,

193(2), 445–453. https://doi.org/10.1111/j.1469-8137.2011.03923.x

Gorter, F. A., Scanlan, P. D., & Buckling, A. (2016). Adaptation to abiotic conditions

drives local adaptation in bacteria and viruses coevolving in heterogeneous

environments. *Biology Letters*, *12*(2), 20150879.

https://doi.org/10.1098/rsbl.2015.0879

Haas, S. M., Cardinal, S., Beaulieu, F., & Forrest, J. R. K. (2019). Mite-y bees: bumble

bees (*Bombus* spp., Hymenoptera: Apidae) host a relatively homogeneous mite

(Acari) community, shaped by bee species identity but not by geographic

proximity. *Ecological Entomology*. https://doi.org/10.1111/een.12706

Hechinger, R. F., & Lafferty, K. D. (2005). Host diversity begets parasite diversity:

bird final hosts and trematodes in snail intermediate hosts. *Proceedings of the*

Royal Society B: Biological Sciences, *272*(1567), 1059–1066.

https://doi.org/10.1098/rspb.2005.3070

Hopkins, S. R., Wojdak, J. M., & Belden, L. K. (2017). Defensive symbionts mediate

host-parasite interactions at multiple scales. *Trends in Parasitology*, *33*(1), 53–64.

https://doi.org/10.1016/j.pt.2016.10.003

Jarrett, B. J. M., Schrader, M., Rebar, D., Houslay, T. M., & Kilner, R. M. (2017).

Cooperative interactions within the family enhance the capacity for evolutionary

change in body size. *Nature Ecology & Evolution*, *1*(7), 0178.

https://doi.org/10.1038/s41559-017-0178

Johnson, N. C., Wilson, G. W. T., Bowker, M., Wilson, J., & Miller, R. M. (2010).

Resource limitation is a driver of local adaptation in mycorrhizal symbioses.

Proceedings of the National Academy of Sciences of the United States of America,

107(5), 2093–2098. https://doi.org/10.1073/pnas.0906710107

Kersch, M. F., & Fonseca, C. R. (2005). Abiotic factors and the conditional outcome of

an ant–plant mutualism. *Ecology*, *86*(8), 2117–2126.

https://doi.org/10.1890/04-1916

Lenth, R. V. (2016). Least-squares means: the R package lsmeans. *Journal of*

Statistical Software, *69*(1), 1–33. https://doi.org/10.18637/jss.v069.i01

Lively, C. M., & Dybdahl, M. F. (2000). Parasite adaptation to locally common host

genotypes. *Nature*, *405*(6787), 679–681. https://doi.org/10.1038/35015069

Müller, J. K., & Schwarz, H. H. (1990). Differences in carrier-preference and evidence

of reproductive isolation between mites of *Poecilochirus carabi* (Acari,

Parasitidae) living phoretically on two sympatric *Necrophorus* species

(Coleoptera, Silphidae). *Zoologische Jahrbücher, Abteilung Für Systematik,*

Ökologie Und Geographie Der Tiere, *117*(1), 23–30.

https://www.cabdirect.org/cabdirect/abstract/19910504588

Nehring, V., & Müller, J. K. (2009). Social environment affects the life history tactic of

a phoretic mite. *Journal of Evolutionary Biology*, *22*(8), 1616–1623.

https://doi.org/10.1111/j.1420-9101.2009.01771.x

Nehring, V., Müller, J. K., & Steinmetz, N. (2017). Phoretic Poecilochirus mites

specialize on their burying beetle hosts. *Ecology and Evolution*, *7*(24),

10743–10751. https://doi.org/10.1002/ECE3.3591

Nehring, V., Teubner, H., & König, S. (2019). Dose-independent virulence in phoretic

mites that parasitize burying beetles. *International Journal for Parasitology*,

49(10), 759–767. https://doi.org/10.1016/J.IJPARA.2019.05.011

Ness, J. H., Morris, W. F., & Bronstein, J. L. (2006). Integrating quality and quantity of

mutualistic service to contrast ant species protecting *Ferocactus wislizeni*.

Ecology, *87*(4), 912–921.

Nuismer, S. L., & Gandon, S. (2008). Moving beyond common□garden and transplant

designs: insight into the causes of local adaptation in species interactions. *The*

American Naturalist, *171*(5), 658–668. https://doi.org/10.1086/587077

Nuismer, S. L., Thompson, J. N., & Gomulkiewicz, R. (2000). Coevolutionary clines

across selection mosaics. *Evolution; International Journal of Organic Evolution*,

54(4), 1102–1115.

Peng, L., Hoban, J., Joffe, J., Smith, A. H., Carpenter, M., Marcelis, T., Patel, V.,

Lynn-Bell, N., Oliver, K. M., & Russell, J. A. (2023). Cryptic community

structure and metabolic interactions among the heritable facultative symbionts of

the pea aphid. *Journal of Evolutionary Biology*.

https://doi.org/10.1111/JEB.14216

R Development Core Team. (2014). R: A language and environment for statistical

computing. In *R Foundation for Statistical Computing*.

Schwarz, H. H. (1996). Host range and behavioral preferences in German sibling

species of the Poecilochirus carabi complex (Acari: Mesostigmata: Parasitidae).

International Journal of Acarology, *22*(2), 135–140.

https://doi.org/10.1080/01647959608684087

Schwarz, H. H., & Müller, J. K. (1992). The dispersal behaviour of the phoretic mite

Poecilochirus carabi (Mesostigmata, Parasitidae): adaptation to the breeding

biology of its carrier *Necrophorus vespilloides* (Coleoptera, Silphidae). *Oecologia*,

89(4), 487–493. https://doi.org/10.1007/BF00317154

Schwarz, H. H., Starrach, M., & Koulianos, S. (1998). Host specificity and permanence

of associations between mesostigmatic mites (Acari: Anactinotrichida) and

burying beetles (Coleoptera: Silphidae: *Nicrophorus*). *Journal of Natural History*,

32(2), 159–172. https://doi.org/10.1080/00222939800770101

Scott, M. P. (1998). The ecology and behavior of burying beetles. *Annu. Rev. Entomol*,

43(4), 595–618. https://doi.org/10.1086/596276

Sun, S.-J., Catherall, A. M., Pascoal, S., Jarrett, B. J. M., Miller, S. E., Sheehan, M. J.,

& Kilner, R. M. (2020). Rapid local adaptation linked with phenotypic plasticity.

Evolution Letters, *4*(4), 345–359. https://doi.org/10.1002/evl3.176

Sun, S.-J., Horrocks, N. P. C., & Kilner, R. M. (2019). Conflict within species

determines the value of a mutualism between species. *Evolution Letters*, *3*(2),

185–197. https://doi.org/10.1002/evl3.109

Sun, S.-J., & Kilner, R. M. (2020). Temperature stress induces mites to help their

carrion beetle hosts by eliminating rival blowflies. *ELife*, *9*(e55649), e55649.

https://doi.org/10.7554/eLife.55649

Svensson, E. I., & Råberg, L. (2010). Resistance and tolerance in animal enemy-victim

coevolution. *Trends in Ecology & Evolution*, *25*(5), 267–274.

https://doi.org/10.1016/j.tree.2009.12.005

Thompson, J. N. (1999). Specific hypotheses on the geographic mosaic of coevolution.

The American Naturalist, *153*(s5), S1–S14. https://doi.org/10.1086/303208

Thompson, J. N. (2005). *The geographic mosaic of coevolution*. University of Chicago

Press.

Thompson, J. N. (2013). *Relentless evolution*. University of Chicago Press.

Whitham, T. G., Allan, G. J., Cooper, H. F., & Shuster, S. M. (2020). Intraspecific

genetic variation and species interactions contribute to community evolution.

Https://Doi.Org/10.1146/Annurev-Ecolsys-011720-123655, *51*, 587–612.

https://doi.org/10.1146/ANNUREV-ECOLSYS-011720-123655

Wilson, D. S. (1982). Genetic polymorphism for carrier preference in a phoretic mite.

Annals of the Entomological Society of America, *75*(3), 293–296.

https://doi.org/10.1093/aesa/75.3.293

Data accessibility

The datasets supporting this article are available as electronic supplementary material and also have been deposited at Github (https://github.com/syuanjyunsun/local_adaptation).

Authors' contributions

S.-J. S. collected the data and performed the analyses. All authors designed the study and contributed to writing the manuscript.

Funding

S.-J.S. was supported by the Taiwan Cambridge Scholarship from the Cambridge Commonwealth, European & International Trust, NTU New Faculty Founding Research Grant, National Science and Technology Council 2030 Cross-Generation Young Scholars Program (111-2628-B-002-050-), and the Yushan Fellow Program provided by the Ministry of Education. R.M.K. was supported by a European Research Council Consolidator grant 301785 BALDWINIAN_BEETLES and a Wolfson Merit Award from the Royal Society.

Acknowledgements

We thank the other members of the Kilner Group, especially S. Aspinall and C. Swannack for logistical support, and S. Issar for providing the field-caught beetles for the choice experiments.

Figures:

Figure 1. Number of *P. carabi* carried by each *Nicrophorus spp.* in Gamlingay and
Waresley Woods. *N. vespilloides*, *N. humator*, *N. interruptus*, and *N. investigator*
were represented as ves, hum, int, and inv, respectively. Means with standard error
are shown, whereas individual data were pointed with a jitter effect to prevent overlap.
(Note that *N. interruptus* and *N. investigator* are routinely found in Gamlingay Woods,
though not in Waresley Woods – see Sun et al. 2020 for further details)

687
688

Figure 2. Population differences of mite preferences between Gamlingay (G) and Waresley (W) Woods, tested over two generations. (a) Proportion of mites that were attracted to each *Nicrophorus* spp. in the first generation (ves = *N. vespilloides*, hum = *N. humator*, int = *N. interruptus*, inv = *N. investigator*) and (b) proportion of mites that were attracted to their parents' preferred *Nicrophorus* spp. P-ves, P-hum, P-int, and P-inv represent mites that chose *N. vespilloides*, *N. humator*, *N. interruptus*, and *N. investigator*, respectively. Means with standard error are shown. The dashed line at 25% represents the proportion of *P.carabi* associating simply by chance with one of the *Nicrophorus* species.

Figure 3. Reproductive success of beetles and mites from Gamlingay Woods, following experimental manipulations of the mite lineages on each carcass. Reproductive success of beetles was measured as (a) brood size, (b) brood mass and (c) average larval mass, whereas mite reproductive success was measured as (d) the number of deutonymphs dispersing with adult beetles. In the mite treatments, 'mix' means that beetles bred alongside 10 mites as a mixture of all four 'lineages' and 'ves' means that beetles bred alongside 10 mites from pure *N. vespilloides* lineage. Means with standard error are shown. Each point represents a different reproductive event.

Figure 4. Burying beetle and mite reproductive success in relation to woodland of
origin and mite/beetle treatments. Beetles / mites deriving originally from Gamlingay
Woods are shown in blue, whereas those deriving originally from Waresley Woods
are shown in red. The beetle / mite treatments refer to the origin of the partner species.
Reproductive success of beetles was measured as (a) brood size, (b) brood mass and
(c) average larval mass, whereas mite reproductive success was measured as (d) the
number of deutonymphs dispersing with adult beetles. Means with standard error are
shown. Each point represents a different reproductive event.

719
720

Table 1. Results of Tukey's post hoc comparisons for beetle species x population interaction in the first choice experiment.

Gamlingay

Groups compared	Estimate	SE	z-ratio	*P*-value
hum to int	-1.22	0.08	-15.54	**<.0001**
hum to inv	-0.29	0.09	-3.24	**0.007**
hum to ves	-1.10	0.09	-11.57	**<.0001**
int to inv	0.93	0.07	13.39	**<.0001**
int to ves	0.12	0.06	1.89	0.233
inv to ves	-0.81	0.08	-9.98	**<.0001**

Waresley

Groups compared	Estimate	SE	z-ratio	*P*-value
hum to int	-1.53	0.10	-15.56	**<.0001**
hum to inv	-0.75	0.11	-6.93	**<.0001**
hum to ves	-2.21	0.11	-20.57	**<.0001**
int to inv	0.79	0.08	10.48	**<.0001**
int to ves	-0.68	0.06	-11.36	**<.0001**
inv to ves	-1.47	0.08	-19.23	**<.0001**

722   Abbreviations: ves, *N. vespilloides*; hum, *N. humator*; int, *N. interruptus*; inv, *N. investigator*.

723   *p* values < 0.05 were highlighted in bold.

Table 2. Results from the final models analysing the fitness components of local adaptation between beetles and mites.

Dependent variable	Explanatory variables	X^2	d.f.	p value
Brood size	mite population	0.67	1	0.412
	beetle population	6.36	1	**0.012**
	carcass mass	24.82	1	**<0.001**
Brood mass	mite population	5.19	1	**0.023**
	beetle population	6.64	1	**0.010**
	carcass mass	14.32	1	**<0.001**
	mite population*beetle population	4.73	1	**0.030**
Average larval mass	mite population	16.69	1	**<0.001**
	beetle population	13.38	1	**<0.001**
	larval density	158.84	1	**<0.001**
	mite population*beetle population	21.17	1	**<0.001**
Number of mite offspring	mite population	2.28	1	0.131
	beetle population	7.75	1	**0.005**
	carcass mass	0.40	1	0.526

p values < 0.05 were highlighted in bold.

729   **Table 3.** Summary of experimental results investigating the extent of local co-adaptation between *N.*

730   *vespilloides* and *P. carabi* in Gamlingay *versus* Waresley Woods

Focal population	Have greatest reproductive success when breeding alongside:
Gamlingay *N. vespilloides*	P-ves mites > mixed lineage mites
Waresley *N. vespilloides*	P-ves mites = mixed lineage mites
Gamlingay *P. carabi*	Gamlingay *N. vespilloides* > Waresley *N. vespilloides*
Waresley *P. carabi*	Gamlingay *N. vespilloides* > Waresley *N. vespilloides*

732