Gateway mechanism confers ultrasensitivity on motor's function

Oshri Afanzar[1†], Diana Di Paolo[2], Miriam Eisenstein[3], Kohava Levi[1], Anne
Plochowietz[2], Achillefs N. Kapanidis[2], Richard Michael Berry[2] and Michael
Eisenbach[1*]

[1]Department of Biomolecular Sciences, The Weizmann Institute of Science,
7610001 Rehovot, Israel

[2]Biological Physics Research Group, Clarendon Laboratory, Department of
Physics, University of Oxford, Oxford, UK

[3]Department of Chemical Research Support, The Weizmann Institute of Science,
7610001 Rehovot, Israel

* To whom correspondence should be addressed. Email
m.eisenbach@weizmann.ac.il.

Summary

In bacteria such as *Escherichia coli*, information from the chemotaxis system to
the flagellar motor is transduced by a single protein, CheY. The motor responds to
CheY by switching its direction of rotation from the default direction to the other.
This response is ultrasensitive, but the mechanism underlying this ultrasensitive
switching is unknown. Here we resolved this mechanism. We show that the motor
employs three sequential steps of CheY binding to distinct sites at the motor, each
with a different outcome. Binding to the first site activates CheY at the motor.
Binding to the second site generates transient motor switching and enables binding
to the third site, which stabilizes the switched state. We demonstrate by
mathematical modeling that such a gateway mechanism provides ultrasensitivity.
The unique combination of tight regulation and flexibility provided by this
mechanism raises the possibility that similar mechanisms may be functional in
other systems as well.

---

† Present address: Department of Chemical and Systems Biology, Stanford University, Stanford,
CA 94305, USA

Keywords

Bacterial chemotaxis, bacterial flagellar motor, CheY, flagellar rotation, flagellar motor, flagellar switch.

Introduction

Chemotaxis of bacteria is achieved by modulating the direction of flagellar rotation. This is done by a switch — a large complex at the base of the flagellar motor, consisting of multiple copies of the proteins FliM, FliN and FliG (Figure 1A). In bacteria like *Escherichia coli*, the switch shifts the direction of rotation from the default, counterclockwise, to clockwise in response to binding the signaling protein CheY, which shuttles back and forth between the chemotaxis receptor complex and the flagellar motor. CheY can be activated by phosphorylation to generate clockwise rotation. This activation is regulated by CheA and CheZ as specific kinase and phosphatase, respectively (Figure 1B). A receptor-mediated attractant response (or removal of a repellent) inhibits CheA activity; stimulation by repellents (or attractant removal) enhances its activity (for reviews — [1-4]). Another covalent modification that activates CheY to generate clockwise rotation is lysine-acetylation [5-7]. The regulation of acetylation is known to involve Acs and CobB as acetyl-transferase and deacetylase, respectively (Figure 1B) [8,9]. It is not known whether other components are involved in regulation of CheY acetylation and what role acetylation plays in bacterial chemotaxis [10].

Earlier studies of the corresponding author's group revealed that phosphorylated CheY (CheY~P) mainly binds to the motor at the *N*-terminus of FliM (FliM$_N$) [11,12]. Subsequently, two additional sites with weaker binding of CheY were reported (termed hereafter 'low-affinity sites'), one at FliN, to which the binding is FliM$_N$-dependent [13], and one at FliM at other location than FliM$_N$ [14]. CheY binding to FliM$_N$ [11,12,15,16] and, apparently to FliN [13], is regulated by phosphorylation. NMR analysis in *Thermotoga maritima* identified the middle domain of FliM (FliM$_M$) as a low-affinity binding site for CheY [17], suggesting that similar binding may also occur in *E. coli*.

The motor response to CheY binding is ultrasensitive, meaning that the dependence of clockwise generation on CheY concentration is highly cooperative [18]. The mechanism underlying this ultrasensitivity is unknown and, as a matter of fact, it is quite intriguing in view of the observation that CheY binding to the motor is non-cooperative [19,20].

Each cell contains multiple flagellar motors. When a considerable fraction of flagella rotate clockwise, the cell preforms a vigor turning motion termed tumbling [21,22], as a result of which the subsequent linear swimming is in a randomly

new direction. It was proposed [23] and demonstrated [24] that anisotropic tumbling (in
which only some of the motors rotate clockwise for a very short time), does not
randomize the swimming direction and maintains directional persistence of
swimming. This behavior was proposed to markedly improve the performance of
collective migration [24], implying an evolutionary advantage. Maintenance of
directional persistence requires extremely short intervals of clockwise rotation. It is
unclear how an ultrasensitive motor can produce relatively long clockwise intervals
for tumbling and short intervals for directional persistence.

The questions raised above are essentially parts of a wider question — how
the motor switches from one direction of rotation to another. Indeed, in spite of
decades of studies of the motor's function, the molecular mechanism underlying
switching has remained obscure. Suspecting that one of the reasons for this
situation is the inability of earlier studies to distinguish between the functions of
each of the individual CheY-binding sites at the switch (due to the masking effect
of the high-affinity binding of CheY to $FliM_N$), we commenced a study aimed at
revealing the functions of each of these sites.

Results

CheY binding to the motor's low-affinity sites generates clockwise rotation

To determine whether CheY binding to the low-affinity sites is functional and
physiologically relevant, we investigated the binding and functional interaction of
CheY with motors in which FliM had been truncated to remove $FliM_N$ (termed
hereafter $FliM_{\Delta N}$), which contains the high-affinity CheY-binding site. To study
binding, we employed *in vivo* Förster resonance energy transfer (FRET). We
measured the interaction between overexpressed CheY-mCherry and $FliM_{\Delta N}$-
YPet-labeled motors, using a $\Delta cheZ$ background to ensure that CheY-mCherry
was mostly phosphorylated (Figure 1C for an explanatory scheme; Figure S1 for a
detailed representative analysis). Addition and subsequent removal of the
attractant serine caused reduction or enhancement, respectively, of the FRET
signal, implying the interaction of CheY-mCherry with the low-affinity sites (Figure
1D, blue). The amplitude of this response was similar to that of non-truncated FliM
(termed hereafter $FliM_{wt}$)-YPet motors (Figure 1D, red). Nevertheless, the
response was slower in $FliM_{\Delta N}$ cells, and the CheY-mCherry concentration had to
be an order of magnitude higher in the case of $FliM_{\Delta N}$ cells (~170 µM vs. ~15 µM
in $FliM_{\Delta N}$ and $FliM_{wt}$ cells, respectively). The response of CheY-mCherry under
conditions that do not allow its phosphorylation ($\Delta cheA$ background, i.e., the kinase
is missing and the phosphatase is present) was complex and smaller in amplitude
(Figure 1D, yellow). No response was observed in a negative control with

overexpressed mCherry (Figure 1D, purple). To determine whether this binding to the low-affinity sites is functional, we examined whether it can generate clockwise rotation of the motor. We tethered cells containing FliM$_{\Delta N}$-YPet motors and ~100 µM CheY in a $\Delta cheZ$ background (to ensure phosphorylation of CheY) to glass via their flagella (Figure 1E), and analyzed their direction of rotation with an automated home-made software. The cells responded to positive stimuli (attractant addition or repellent removal) with reduced clockwise rotation (Figure 1F). Repellent stimulation had hardly any effect in this background (Figure 1G). CheY acetylation, stimulated by the acetyl donor acetate, also generated clockwise rotation in FliM$_{\Delta N}$ cells (Figure 1H, blue). It is worth noting that the decrease in clockwise level in response to acetate removal was biphasic. Although acetate also functions as a repellent, the clockwise generation was the outcome of acetylation because the repellent benzoate, which acts as a repellent by the same mechanism as does acetate [25,26] but does not serve as an acetyl donor [6,8], did not generate clockwise rotation (Figure 1H, red; Figure S4 for sequential additions of acetate and other repellents). These results suggest that CheY binding to the low-affinity sites is functional in the sense of generating clockwise rotation, and that acetylation is apparently more potent than phosphorylation for this.

FliM$_N$ filters CheY from the cytoplasm and activates it

The finding that FliM$_N$ is not essential for clockwise generation raised the question of what its role is. FliM$_N$ promotes the interaction of CheY with the motor because, in its absence, CheY must be overexpressed for clockwise generation. Therefore, a reasonable possibility is that FliM$_N$ tethers CheY to the motor, thereby elevating the local concentration of CheY at the low-affinity sites [13,17]. According to this possibility, FliM$_N$ acts as a 'fishing line', increasing the rate of CheY association with the clockwise-generating low-affinity sites at the motor. However, this might not be the only function of FliM$_N$. It has been found that FliM$_N$-bound CheY adopts an intermediate conformation between the active and inactive states [27] and enhances the rate of CheY phosphorylation by small phosphodonors [28]. This begs the question of whether these observed features serve to increase CheY's potency to generate clockwise rotation when it is tethered to FliM$_N$. To examine this possibility, we studied the motor behavior in a FliM$_{\Delta N}$ strain that expresses CheY fused to FliM$_N$ (FliM$_N$–CheY) and compared it with the motor's behavior of the same strain that expresses wild-type CheY instead of FliM$_N$–CheY. The expression of both proteins was induced in parallel and under identical conditions to different levels by an IPTG-inducible promoter. We made this comparison in $\Delta cheZ$ and $\Delta cheA$ backgrounds, elevating and lowering CheY phosphorylation, respectively. In a $\Delta cheZ$ background, cells expressing FliM$_N$–CheY~P spent much more time in clockwise rotation than cells expressing CheY~P (Figure 2A; blue and red,

respectively). In Δ*cheA* background, cells expressing FliM$_N$–CheY could generate
low levels of clockwise rotation whereas cells expressing wild-type CheY could not
produce clockwise rotation (Figure 2A, B; cyan and purple, respectively). This
implies that CheY can acquire an active clockwise-generating conformation as a
result of binding to FliM$_N$. Acetylation, mediated by saturating concentrations of
acetate, greatly enhanced clockwise rotation (Figure 2A-B; green). The response
of the cells was sigmoidal with respect to acetate concentration (Figure 2C for
responses and Figure 2D for summarized response curve). The reason for this
sigmoidity is unclear. The cooperative deactivation of Acs by PatZ-mediated
acetylation [29] could be a reason this sigmoidity, provided that Acs acetylation
inhibits the conversion of acetate to AcAMP but not the subsequent conversion of
AcAMP to AcCoA. The response to acetate removal was monophasic at low
acetate concentrations but biphasic at high concentrations (Figure 2C; black lines).
This biphasic response to acetate removal in cells expressing FliM$_N$–CheY differed
from the cognate biphasic response in cells expressing non-modified CheY (Figure
1H) in the sense that here, contrary to CheY-expressing cells, the slow phase
preceded the rapid one. This difference implies that the biphasic response is not
due to acetate metabolism. The clockwise-generating effect of acetate was
acetylation specific as benzoate had no effect on the clockwise rotation (Figure
2E). Notably, at high FliM$_N$-CheY levels, both stimulus types, attractants and
repellents, caused reduced clockwise rotation. However, unlike the response to
benzoate and leucine, which started as a repellent response and then reversed,
the response to acetate was clockwise elevation from the very beginning (Figure
2F). The inverted repellent response observed herein at high FliM$_N$-CheY
concentration resembles the inverted attractant response reported for CheY-
containing gutted *E. coli* cells [30].

To substantiate the conclusion that FliM$_N$ activates CheY independently of
phosphorylation, we took advantage of CheY(D13K) being a constitutively active
non-phosphorylatable variant of CheY [31] and compared its dose-dependent
clockwise-generating activity with that of FliM$_N$-CheY(D13K) in Δ*cheA* background.
The FliM$_N$ fusion extremely increased the clockwise generation potency of
CheY(D13K) (Figure 2G), so much so that clockwise rotation could be generated
without induction, just by the leaky expression of the protein from the plasmid.
Leaky expression levels were roughly equivalent to endogenous CheY expression
levels, suggesting that, with respect to the CheY concentration required for
clockwise generation, the FliM$_N$ fusion to CheY(D13K) substantially compensated
for the FliM$_N$ deletion from the motor. This strongly supports the notion that FliM$_N$
functions to promote switching in a phosphorylation-independent manner.

Taken together, the results suggest that FliM$_N$ not only tethers CheY to the
motor but also activates it, and that phosphorylation and acetylation further

enhance this activation. However, high CheY~P-FliM$_N$ levels result in an inverted
response. Another function suggested for FliM$_N$ was to provide some of the
interface required for the interaction of CheY~P with FliN [13]. While our results do
not test the validity of this suggestion, it should be noted that this function is
phosphorylation dependent whereas the function of FliM$_N$, demonstrated herein,
appears to be phosphorylation independent.

Both CheY binding and clockwise generation are biphasic processes

Clockwise generation in the absence of FliM$_N$ indicates that one or more of the
other CheY-binding sites, i.e., of the low-affinity sites, is involved in CheY binding.
The way that CheY interacts with these sites is of special interest in view of the
fact that the motor's response to CheY is ultrasensitive [18] even though the binding
of CheY to the motor is non-cooperative [19,20]. Since these published binding
experiments essentially reflected binding to the high-affinity site at FliM$_N$, we
investigated whether CheY binding to the low-affinity sites is cooperative and, if
so, whether it can explain the ultrasensitive response of the motor. To this end, we
measured the amount of CheY bound to FliM$_{\Delta N}$ motors as a function of the
intracellular concentration of CheY. We determined the amount of bound CheY by
FRET photobleaching, measuring the increase in fluorescence intensity of the
donor, FliM$_{\Delta N}$-YPet, due to photobleaching of the acceptor, CheY~P-mCherry
(Figure 3A, B). This technique was used previously for FRET measurements of
chemotactic signaling at the population level [32]. We preformed the measurements
in vivo, in single cells, using a $\Delta cheZ$ background to promote CheY
phosphorylation (Figure 3C, black curve, for one representative experiment out of
three). We used parallel measurements with free mCherry to control for non-
specific binding (Figure 3C, purple curve). We found that, in all measurements, the
fitted binding curve could be described as the sum of two binding processes
(Figure 3D; black for the average fitted curve of three experiments; red and blue
for the binding processes derived from this black binding curve). It appears that a
saturation process was predominant at low CheY concentrations and a
cooperative-like process was dominant at high concentrations. It should be noted
that none of the measured FRET phases was due to FRET signals between free
FliM$_{\Delta N}$-YPet and CheY-mCherry molecules in the cytoplasm. This is for two
reasons. First, no CheY~P-mCherry binding to FliM$_{\Delta N}$-YPet was detected *in vitro*
at these CheY concentrations. And second, only spots of high intensity FliM$_{\Delta N}$-
YPet signal were measured. Likewise, the non-cooperative and cooperative
binding phases were not due to CheY~P-mCherry binding to FliM$_{\Delta N}$-YPet in the
cytoplasm and in the motor, respectively. This is because clockwise rotation could
be observed at CheY concentrations lower than those of the cooperative phase. If

only the latter contributed to CheY binding to the motor, clockwise rotation could not have been observed.

Two CheY-binding phases, as found above, may well give rise to a biphasic motor response to CheY, meaning that the outcome of each binding phase is a different clockwise-generation mechanism. If so, there is a good chance that the clockwise intervals generated by each mechanism would be distributed around a different interval length. Furthermore, because each of the binding phases occurs at a different concentration range of CheY (Figure 3D) and because the extent of clockwise rotation is dependent on the intracellular CheY concentration (Figure 2A), we expected that the clockwise distribution related to each mechanism would be dependent on the clockwise level. To put all these to the test, we measured the lengths of clockwise intervals in FliM$_{\Delta N}$ motors and studied their distributions at different clockwise levels.

We quantified switching by video analysis of tethered $\Delta cheA$ cells containing FliM$_{\Delta N}$ motors and expressing FliM$_N$-CheY in the presence of acetate (10 mM, pH 7.0), recorded at 75 frames/s. We defined clockwise intervals as intervals that have at least 4 consecutive frames (52 ms) of clockwise rotation at a rate higher than 1 Hz, which is the measured rotation rate of fluctuations of non-rotating cells distributed around 0±0.5 Hz (±SD). This 4-frame threshold excluded short events that might not be clockwise rotation mediated by CheY. This is according to a negative control consisting of a $\Delta cheY$ strain, in which the generation of clockwise rotation cannot be mediated by CheY. In this negative-control strain we studied 1 min recordings of 344 cells, and found ~100 clockwise intervals, all shorter than 50 ms. Figure 4A shows an example of a clockwise interval lasting 4 frames, the shortest duration that exceeded the threshold (see also supplementary Movie S1). To avoid artefacts that might have been caused by interactions between the cell body and the surface, we also excluded from the analysis intervals that were flanked by pause events (Figure 4B; pause events were defined as absolute rotation rate <1 Hz for more than 2 consecutive frames). We grouped clockwise intervals from cells that had similar average clockwise levels and produced from each group a distribution of interval lengths (Figure 4C). At low clockwise levels, the distribution appeared as a fast, monophasic exponential decay (Figure 4C, blue; note the logarithmic scale of the ordinate, in which exponential decays are seen as straight lines). The distribution became biphasic at intermediate clockwise levels (Figure 4C, gray), and slow monophasic at high clockwise levels (Figure 4C, orange). To calculate the contribution of each of the phases to clockwise rotation time, we fitted each of the distributions in Figure 4C with a biexponential expression, and extracted the average time of clockwise rotation of each process in each clockwise level (Figure 4D for demonstration of a calculation for a given clockwise level). Like the first phase of CheY binding to the

motor (Figure 3D), the first phase of clockwise generation steadily increased in
clockwise rotation time up to saturation, generating only very short intervals of
clockwise rotation (Figure 4E, blue). Like the second binding phase, the second
clockwise-generation phase emerged at intermediate clockwise levels, producing
long intervals of clockwise rotation (Figure 4E, red). The average reversal
frequency of these motors was roughly 8-fold higher than that of wild-type motors
(Figure 4F compared with peak reversal frequency of ~0.8 s^{-1} in wild-type cells
containing constitutively active CheY [33]), meaning overall shorter clockwise
intervals. This difference is probably a reflection of the lower affinity of $FliM_{\Delta N}$
motors for CheY relative to that of motors containing $FliM_{wt}$. These results suggest
that the two phases of CheY binding to the motor bring about two phases of
clockwise generation.

The measurements above (Figure 4C-F) were based on motors of cells in
which acetylation of CheY was promoted. We verified that similar distributions
were obtained when CheY was activated by phosphorylation, by D13K mutation,
and by $FliM_N$ fusion (Figure S5), meaning that the biphasic motor response was
not the product of acetylation dynamics. Interestingly, when clockwise rotation was
generated by $FliM_N$-CheY (in the absence of any other activation), only short
intervals were observed (Figure S5B). This observation taken together with the
limited clockwise-generation activity of this variant (Figure 2A, cyan) suggests that
the generation of the second phase depends on excessive CheY activation. Thus,
it appears that adequate CheY activation is required for the formation of a biphasic
response.

The first phase of clockwise generation enables directional persistence in
swimming

Since the first clockwise-generation phase was characterized by very brief
clockwise intervals (Figure 4E, blue), and since generation of such short intervals
only slightly deflects the swimming trajectory and maintains directional persistence
[23,24], we investigated whether the first clockwise-generation phase indeed
preserves directional persistence. We assessed the distribution of angular
deflection in swimming cells, with an expectation to find that the distribution peaks
at small turning angles, compatible with directional persistence. For this, we
measured the angular shift in 1 s trajectories of *$fliM_{\Delta N}$ $\Delta cheA$* cells expressing
$FliM_N$-CheY in the absence or presence of acetate (Figure S6; blue and red,
respectively). As a control for noise, we made similar measurements in cells that
do not express CheY and, therefore, do not tumble (Figure S6; green and purple
for the absence or presence of acetate, respectively). The experimental
distributions were significantly different from the control distributions (P = $6x10^{-9}$
and $4x10^{-26}$ for the absence and presence of acetate, respectively, according to a

two-sample Kolmogorov-Smirnov test). In the absence of CheY, turning angles can be related to noise. The same noise is expected to appear also in the experimental distributions. To de-noise the experimental distributions, we subtracted them from the control distributions (Figure 4G). We found that both in the absence or presence of acetate, the subtracted distribution peaked at 20^o (Figure 4G), which is the angular deflection expected to maintain directional persistence of swimming. It is worth noting that, under some assumptions, a peak distribution of 20^o turning angle was calculated to be the product of an average clockwise intervals of 50 ms [34]. This value is approximately the average clockwise interval length of the first phase of clockwise generation in our measurements (Figure 4E and Figure S5B). Thus, the first phase of clockwise generation appears to produce angular deflection while maintaining directional persistence. Somewhat larger proportion of high deflection angles were observed in the presence of acetate, suggesting that acetate is less supportive of directional persistence.

CheY binding to FliN is apparently responsible for the first phase of clockwise generation

The simplest explanation of biphasic binding that results in biphasic clockwise generation is CheY binding to two different sites on FliM$_{\Delta N}$ motors. Since, as mentioned above, the known CheY-binding sites in FliM$_{\Delta N}$ motors are the low-affinity sites at FliN and FliM$_M$, we determined whether or not this explanation is correct by studying the generation of clockwise rotation in a *fliN*(A93D) mutant, in which the interaction of FliM$_N$-CheY~P with FliN is greatly impaired [13]. We expected that if CheY indeed binds to two different motor sites, the distribution of interval lengths would be skewed by the FliN mutation. To this end, we first characterized how FliM$_N$-CheY affects clockwise generation in the mutant. We found that clockwise generation by FliM$_N$-CheY~P was highly impaired in the *fliN*(A93D) mutant (Figure 4H, blue), supporting the conclusion of Sarkar *et al.* [13] that FliN is involved in FliM$_N$-CheY~P binding. Pushing phosphorylation further by the repellent benzoate did not increase clockwise rotation (Figure 4H, red point). Unlike phosphorylation, FliM$_N$-CheY acetylation and, the more so, acetylation and phosphorylation together, led to significant clockwise rotation (Figure 4H, green and purple, respectively). To determine whether the *fliN*(A93D) mutation affected any of the switching phases, we calculated the distributions of clockwise interval length of the mutant under phosphorylating and acetylating conditions (Δ*cheZ* background in the presence of acetate). As in the case of non-mutated FliN, two phases of clockwise generation were observed (Figure 4I). However, the first clockwise-generation phase was saturated already at the lowest clockwise level (Figure 4J, blue) along with the emergence of the second phase (Figure 4J, red), and very short intervals of clockwise rotation were hardly observable (Figure 4I).

Similar observations were made when the cells lacked CheA (in which only
acetylation occurred; Figure 4K), suggesting that the generation of long intervals
were the result of CheY acetylation. The occurrence of long intervals could also be
demonstrated by the peak reversal frequency of these motors, which was roughly
3-fold lower than that of wild-type FliM$_{\Delta N}$ motors (Figure 4L; cf. Figure 4G). This
implies that short intervals of clockwise rotation, manifested as frequent reversals,
are associated with CheY binding to FliN, and long intervals, manifested as low
reversal frequency, is mostly dependent on CheY binding to another site.

346 CheY binding to FliM$_M$ is apparently responsible for the second phase of
347 clockwise generation

The suppression of short intervals of clockwise rotation when CheY interaction with
FliN was impaired strongly suggested that long intervals are produced by CheY
interaction with a motor site other than FliN. An obvious candidate site for this
binding is the other low-affinity binding site, FliM$_M$, shown to bind CheY in *T.*
maritima [17]. To examine the plausibility of CheY binding to FliM$_M$ in *E. coli*, we
employed *in vivo* crosslinking of cells expressing FliM$_{\Delta N}$-YPet, using a non-specific
crosslinker, glutaraldehyde. (We wish to point out that FRET experiments *in vivo*
cannot distinguish CheY binding to FliM from CheY binding to FliN due to the
physical proximity of FliM to FliN.) The advantage of using crosslinking is that,
beyond being carried out *in vivo*, it can detect weak interactions. We tracked the
crosslinking products by SDS-PAGE and scanning the gel for FliM$_{\Delta N}$-YPet
fluorescence. This enabled us to quantify the extent of complex formation. Among
many other crosslinking products, we obtained a product at the size of a complex
between FliM$_{\Delta N}$-YPet and CheY (Figure 5A, lanes 1–3). The formation of this
complex was dependent on CheY overexpression (Figure S7), implying that the
complex indeed contained CheY. These observations are consistent with the
possibility of CheY binding to FliM$_M$ *in vivo*, but they do not rule out a possibility of
CheY binding to another FliM domain (e.g., to the C terminus domain of FliM).
Various direct *in vitro* binding assays between purified CheY and FliM$_{\Delta N}$ were not
conclusive, probably due to the low affinity of FliM$_{\Delta N}$ for CheY.

To examine the likelihood that CheY binding to FliM$_M$ is functional, we
employed docking analysis between CheY and FliM$_M$ based on their binding
interface in *T. maritima* (Supplementary Methods for detailed description). We
especially paid attention to the β4α4 loop of CheY, known to be at the CheY-FliM$_M$
binding interface in *T. maritima* [17] and predicted in our earlier all-atom molecular
dynamics simulation to be activated by CheY acetylation at its clockwise-
generating site Lys-91 [35]. In the docking analysis, we produced a number of
models of CheY binding to FliM$_M$ in *T. maritima*, and translated those fitted well
with the NMR results of Dyer *et al.* [17] to *E. coli* proteins by superposing the active

and inactive forms of CheY upon the model of FliM$_M$. We substantiated the
superposition results by docking the *E. coli* proteins independently of the *T.*
maritima model. This docking demonstrated that while the active conformation of
CheY well docked onto FliM$_M$ (Figure 5B, up-left docking model), the inactive
conformation of the β4α4 loop of CheY clashed with FliM$_M$ (Figure 5B, down-right).
This is consistent with the proposition that acetylation generates clockwise rotation
by enhancing the binding of CheY to FliM$_M$, and that this enhancement is done by
modulation of the β4α4 conformation.

The analysis predicted two prominent latching interfaces of CheY with
FliM$_M$. One interface is electrostatic and is mostly contributed by arginine at
position 94 of FliM (Figure S8A). The other interface is a hydrophobic patch in FliM
(Figure S8B). To determine the functional relevance of the first interface, we
examined the rotation of motors of cells expressing FliM$_{\Delta N}$(R94S) or FliM$_{\Delta N}$(R94L)
in a Δ*cheZ* background and overexpressing FliM$_N$-CheY. These motors hardly
rotated clockwise even when benzoate was used to further stimulate CheY
phosphorylation. Very low levels of clockwise rotation were observed following the
promotion of CheY acetylation by acetate (Figure 5C). These results are in line
with the docking model's prediction that FliM(R94) is involved in CheY binding. To
determine the functional relevance of the second interface, we examined whether
replacement of a charged residue to a hydrophobic one within the hydrophobic
CheY-FliM$_M$ binding interface (E214W) would increase clockwise generation.
Indeed, clockwise rotation in the *fliM*$_{\Delta N}$(E214W) mutant in a Δ*cheZ* background
was observed at much lower FliM$_N$-CheY expression levels (Figure 5D). It appears
that the mutation affected the motor's sensitivity to CheY rather than affecting its
intrinsic clockwise level because the clockwise rotation levels of *fliM*$_{\Delta N}$(E214W) in
a Δ*cheA* background, where FliM$_N$-CheY is mostly inactive and does not contribute
much to clockwise generation, were low and comparable to cells containing non-
mutated FliM$_{\Delta N}$ (Figure S9; blue). Nevertheless, when FliM$_N$-CheY was activated
by acetate, the response of cells with FliM$_{\Delta N}$(E214W) motors exceeded that of
FliM$_{\Delta N}$ (Figure S9; red).

To determine whether the proposed interaction of CheY with FliM$_M$ affects
the second phase of clockwise generation, we examined the distributions of
clockwise interval lengths and reversal frequency in *fliM*$_{\Delta N}$(R94SL) and
fliM$_{\Delta N}$(E214W) mutants. Indeed, only the first phase of clockwise generation was
apparent in *fliM*$_{\Delta N}$(R94SL) mutants (Figure 5E, F). The peak reversal frequency of
the mutants was roughly similar to that of its wild-type FliM$_{\Delta N}$ parent but the
mutants did not reach high clockwise levels (Figure 5G). The distributions of
clockwise intervals produced by FliM$_{\Delta N}$(E214W) motors were approximately
monophasic at all clockwise levels (Figure 5I), with interval lengths very similar to

those exhibited by FliN(A93D) motors (Figure 5J *versus* Figure 4J). Consistently,
the peak reversal frequency of the FliM$_{\Delta N}$(E214W) motors was between the values
of wild-type FliM$_{\Delta N}$ motors and FliN(A93D) FliM$_{\Delta N}$ motors (Figure 5K *versus* Figure
4F, L). It appears that the E214W mutation increased the affinity of FliM$_M$ for CheY
to a level higher than that of FliN for CheY. Consequently, intervals initiated by
CheY binding to FliN were immediately extended by subsequent binding to FliM$_M$.
Furthermore, the inability of the *fliM*$_{\Delta N}$(R94SL) mutant to exhibit long clockwise
intervals suggests that CheY binding to FliM$_M$ is required for stable clockwise
rotation.

Another observation made in these experiments is that the rotation rate in
the clockwise intervals was lower in the *fliM*$_{\Delta N}$(R94SL) mutant (Figure 5H) relative
to the *fliM*$_{\Delta N}$(E214W) mutant (Figure 5L) and the wild-type FliM$_{\Delta N}$ parent (Figure
S5). Notably, the distribution of clockwise rotation rates as well as the distribution
of clockwise interval lengths of the *fliM*$_{\Delta N}$(R94SL) mutant (Figure 5H, E) expressing
FliM$_N$-CheY under phosphorylating conditions were similar to those of non-mutated
FliM$_{\Delta N}$ cells (Figure S5B). Low clockwise rotation rates in motors that are not stably
rotating clockwise is consistent with a conformational spread model [36], where low
clockwise rotation rate is suggestive of incomplete switching events (i.e., events in
which the conformation of the motor did not stabilize in a coherent clockwise
conformation). Taken together, these results suggest that the second phase of
switching is due to CheY interaction with FliM$_M$, and that the outcome of this
interaction is stabilization of the rotation in the clockwise direction.

438 CheY~P and CheY~Ac preferentially bind to FliN and FliM$_M$, respectively

The observation that clockwise generation in FliN(A93D) motors could only be
observed in the presence of acetate (Figure 4H) raised the possibility that the low-
affinity binding sites have different preferences for CheY~P and for CheY~Ac. To
examine this possibility, we first investigated whether CheY~P and CheY~Ac have
different *in vivo* affinities for the motor. To this end, we compared *in vivo* the dwell
time of single CheY molecules at FliM-YPet motors between phosphorylating
conditions, acetylating conditions, and conditions under which CheY is both
phosphorylated and acetylated. We electroporated CheY(I95V) molecules labeled
with a maleimide modification of the photo-stable organic dye Atto647 into FliM-
YPet expressing cells [37]. The experiments were carried out both in a $\Delta cheZ$
background to make CheY fully phosphorylated and in a $\Delta cheA$ background to
make CheY non-phosphorylated. The *cheY*(I95V) mutation was designed to
increase CheY affinity for FliM$_N$ [38], thus enhancing sampling of otherwise rare
binding events. Custom-written software tracked CheY(I95V)-Atto647 molecules
and estimated motor locations in each cell using FliM-YPet fluorescence images.
CheY(I95V)-Atto647 molecules dwelling within 75 nm of a motor for >30 ms were

interpreted as binding to the motor (Figure 6A; see Methods). Presence of acetate
increased the dwell time of CheY at the motor (Figure 6B and 6C for the survival
probabilities of CheY~P and CheY, respectively, to remain bound to the motor;
Movie S2). Notably, the very long dwell events in Movie S2 were only detected in
the presence of acetate. The extension of CheY's dwell time at the motor by
acetylation taken together with the generation of long clockwise intervals by
acetate even when CheY binding to FliN is impaired (Figure 4I) suggest that $FliM_M$
has a preference for CheY~Ac.

To examine whether, indeed, CheY~P and CheY~Ac preferentially bind to
FliN and $FliM_M$, respectively, we measured the concentration-dependent response
of $FliM_{\Delta N}$ motors to CheY~P, CheY~Ac, and CheY with both modifications
combined, expecting to observe concentration dependence that resembles the two
phases of CheY binding to the motor (Figure 3D). Unmodified CheY did not
produce clockwise rotation (pre-acetate stimulation in Figure 6D; purple curve in
Figure 6F). CheY acetylation achieved by acetate addition increased, after a short
delay, the fraction of time spent in clockwise rotation (Figure 6D). The extent of the
response was sigmoidally dependent on the CheY level (Figure 6F, red).
Increasing concentrations of CheY~P (CheY in a $\Delta cheZ$ background) enhanced
clockwise rotation up to saturation (Figure 6F, blue). Acetylation of CheY~P further
enhanced clockwise rotation (Figure 6E; Figure 6F, green). Notably, the clockwise
curve of CheY~P (Figure 6F, blue) resembled the first binding phase (Figure 3D,
blue), including the CheY~P concentration required for achieving saturation.
Likewise, the clockwise curve of CheY~Ac (Figure 6F, red) resembled the second
phase of binding (Figure 3D, red). These resemblances endorse the conclusion
that the first binding phase (blue) in Figure 3D is more associated with CheY~P
binding and the second phase (red) is mainly associated with CheY~Ac binding.
They also suggest that the motor hypersensitivity to CheY is the outcome, at least
in part, of the apparently cooperative binding of CheY~Ac and CheY~Ac~P to
$FliM_M$ (see Discussion).

Discussion

The molecular mechanism underlying the motor's switching from one direction of
rotation to the other, which is the essence of bacterial chemotaxis, is yet unknown.
Here we showed that clockwise generation is mediated by CheY interaction with
three different sites at the motor. We revealed the function of each of these sites
and the probable sequence of events at the motor during switching from
counterclockwise to clockwise rotation.

Functions of FliM$_N$

We found that the first site to which CheY binds, a high-affinity site at FliM$_N$, is not essential for clockwise generation (Figures 1D, 2A). It tethers CheY to the motor and activates it. We propose that this CheY activation at the motor is a preliminary step in the process of clockwise generation (Figure 7A), which may have evolved to filter out crosstalk with proteins having CheY-like topology in other two-component signaling pathways. Binding of such proteins to FliM$_N$ is expected to be futile.

Functions of the other CheY-binding sites in the switching mechanism

We showed that, in FliM$_{\Delta N}$ motors, CheY binding to the motor is biphasic (Figure 3D) as a result of binding to two different sites at FliN and FliM$_M$. We further demonstrated that CheY binding to these sites generates short and long clockwise intervals, respectively, reflected in a biphasic clockwise generation (Figures 4–6), and that FliN has preference for CheY~P and FliM$_M$ for CheY~Ac. We propose that CheY~P binding to FliN makes the first catalytic step of clockwise generation, producing mainly brief episodes of clockwise rotation (Figure 7B). CheY~Ac binding to FliM$_M$ makes the second catalytic step of clockwise generation, produces long clockwise intervals and, thereby, stabilizes the rotation in clockwise direction (Figure 7C). Because the conformation of FliM is different in the counterclockwise and clockwise states, it is reasonable that FliM$_M$ can only bind CheY when it is in its clockwise state, meaning that the generation of brief clockwise rotation by the preceding binding step to FliN enables binding to FliM$_M$ and generation of longer clockwise rotation. Indeed, when CheY interaction with FliM$_M$ was turned down, the motor produced only short intervals of clockwise rotation (Figure 5E-G) that were relatively slow in rate (Figure 5H), suggesting incomplete switching according to the conformational-spread model [36]. This observation was not captured in earlier attempts to characterize the switching process, probably because of the masking effect of FliM$_N$ in those experiments. The presence of FliM$_N$ is also the reason for earlier findings [9,39] that acetylation reduces the binding of CheY to FliM$_{wt}$ *in vitro*.

Physiological significance

This switching mechanism provides dynamics and flexibility to the system in two senses. First, at the intermediate stage, the system can either proceed to a stable rotation in the new direction or shift back to the original direction. And second, it provides the system flexibility with respect to the swimming patterns. It appears that the first CheY binding phase is associated with swimming patterns that enable slight reorientation while maintaining directional persistence (Figure 4G), and the second phase, which produces long clockwise intervals, results in tumbling events

that lead to sharp reorientation. In this way this switching mechanism increases
chemotactic performance. With these advantages, it would not be surprising if
similar mechanisms are found in the future in the output of other signaling systems
as well.

Motor's ultrasensitivity can emerge from a gateway mechanism

The response of wild-type motors to CheY is ultrasensitive [18]. Since CheY binding
to $FliM_{wt}$-containing motors is non-cooperative [19,20], meaning that CheY binding to
$FliM_N$ does not generate ultrasensitivity, it is conceivable that CheY binding to FliN
and $FliM_M$ does generate ultrasensitivity. The question is how. A reasonable
possibility is that CheY can only (or mainly) bind to $FliM_M$ when the motor is already
rotating clockwise. The finding that FliM subunits within the motor have different
orientations in the counterclockwise and clockwise states [40] is consistent with this
possibility, as one orientation may sterically block CheY binding. In such a
mechanism, which we term a gateway mechanism, clockwise rotation, which
results from CheY binding to FliN, exposes another CheY-binding site at $FliM_M$.
The binding of CheY to that site stabilizes clockwise rotation. This mechanism is
in line with the biphasic curve of CheY binding to the motor (Figure 3D). To
examine whether a gateway mechanism can confer ultrasensitivity on the motor's
response to CheY, we constructed a simplified kinetic model of the motor in which
CheY binds to $FliM_M$ only if FliN is already CheY-bound. The simplified model
consists of a single FliN-binding site and two $FliM_M$-binding sites (Figure 8A; the
number of binding sites is not expected to affect the conclusion). A requirement
from such a model is that the binding affinities of CheY for its binding sites at the
motor would affect the ultrasensitivity of the motor. Therefore, we examined CheY
binding to the motor as a function of the ratio between CheY affinity for FliN and
CheY affinity for $FliM_M$. When the dissociation constant for CheY binding to FliN
was much lower than for the binding to $FliM_M$, the modeled binding was
multiphasic, but not ultrasensitive (Figure 8B, blue). When the dissociation
constant for the binding of CheY for FliN was much higher than for the binding to
$FliM_M$, the binding was ultrasensitive (Figure 8B, orange). When the dissociation
constants were comparable, the binding was a simple saturation process (Figure
8B, gray; note the logarithmic scale). Consistently, the apparent Hill coefficients of
the curves shifted from positive cooperativity (Figure 8B, insert, orange), through
zero cooperativity (Figure 8 B, insert; grey), to negative cooperativity (Figure 8 B,
insert, blue). Thus, ultrasensitivity can arise from a gating mechanism.

Conclusion

The flagellar motor likely binds CheY at a non-catalytic site, $FliM_N$, with a resultant
activation of CheY, and at two different catalytic sites, FliN and $FliM_M$, with a

resultant generation of short and long intervals of clockwise rotation, respectively.
The generation of short and long intervals can support directional persistence and
randomization of direction in the tactic behavior of bacteria. Ultrasensitivity in the
motor response to CheY may be produced by a gateway mechanism of switching,
in which the extent of ultrasensitivity emerges from the difference in affinities of the
binding sites for CheY.

Methods

Strains and plasmids

To produce *fliM$_N$* truncation, *fliM-YPet* was cloned from the strain JPA945 [41] (kindly
provided by J. Armitage) with genomic flanks of 500 base pairs to the Pst1 sites of
pDS132 suicide plasmid [42]. FliM residues 1–16 were truncated from the resulting
plasmid by RF-cloning. The constructed plasmid was used to perform allelic
exchange with strain JPA945 to produce strain EW566 bearing *fliMΔ(1-16)-YPet*
genomic mutation. The mutation was verified by PCR sequencing. Plasmidic
mutations were produced by RF cloning. The genomic deletion mutations of *cheA*
and *cheZ* were produced by subjecting cells to P1 transduction with phage
containing the genetic background of strain JW1870 or JW1877, respectively. See
Tables S2-3 for detailed strains and plasmids list.

Growth conditions

Strains, diluted 1:100 from overnight cultures, were cultured to mid-late
exponential phase at 30°C in tryptone broth with appropriate antibiotics to maintain
the plasmids. For CheY expression, cells were grown to mid-late exponential
phase and induced by IPTG for 3-4 h. It appeared that dilution of overnight cultures,
which spent 1–2 days on the bench at room temperature, produced better
responses to acetate.

Analysis of the direction of flagellar rotation

594 The direction of flagellar rotation was determined by the tethering assay [43].
595 Analysis of the rotation was done automatically by an in-house custom-made
596 MATLAB program. For interval analysis, each subjected interval was considered
597 valid if it was flanked by two other valid intervals or by rotation in the opposite
598 direction. See Supplementary Methods for details and other considerations.

In vivo FRET

Cells were immobilized onto a polylysine-coated 8-well glass bottom (ibidi, Germany), washed 4 times with motility buffer and visualized using an inverted confocal microscope set (IX81, Olympus, Japan) with an UPLSAPO 60x0 NA:1.35 objective equipped and excitation lasers at 488 and 559 nm for YPet and mCherry, respectively. FRET was measured by acceptor photobleaching with high intensity 559 nm laser. FRET efficiency was estimated as ($I_{Post\ bleaching}$ − $I_{Prior\ bleaching}$) / $I_{Post\ bleaching}$. See Supplementary Methods for details.

Single molecule analysis

The locations of CheY(I95V)-Atto647 and FliM-YPet were automatically estimated by a custom-made MATLAB script. Motor locations were identified as peaks of fluorescence. The exact motor location was determined by identifying the maxima location of a 2D Gaussian, fitted to each motor spot. For the identification of motors, several images were acquired, normalized and averaged before Atto647 imaging. The location of CheY(I95V)-Atto647 was determined similarly to motor location in each frame. CheY(I95V)-Atto647 was considered bound to the motor when it was within 75 nm of a motor location for at least 30 ms (3 frames). Shorter co-localizations were excluded to avoid false positive measurements (e.g., molecules diffusing near the motor). The value of 30 ms was chosen as the half of the expected mean dwell time of CheY(I95V)-Atto647 with $FliM_N$ [~60 ms given a K_d = 3.9 µM for CheY(I95V)-$FliM_N$ interaction and an estimated k_{on} = $4x10^6$ $M^{-1}s^{-1}$ [20,38]. The mean-square displacement in 2 dimensions due to free diffusion is $<r^2>$ = $4Dt$. Setting $<r^2>$ = $(75\ nm)^2$ and t = 30 ms gives D = 0.047 $\mu m^2\ s^{-1}$, which is lower than the diffusion coefficient of CheY in the cell. Thus, the false-positive rate due to diffusing molecules is expected to be negligible. See Supplementary Methods for a detailed description of protein labeling and optical setup.

References

1. Berg, H. C. The rotary motor of bacterial flagella. *Annu. Rev. Biochem.* **72,** 19–54 (2003).
2. Eisenbach, M. in *Chemotaxis* (ed. Eisenbach, M.) 53–215 (Imperial College Press, 2004).
3. Terashima, H., Kojima, S. & Homma, M. Flagellar motility in bacteria structure and function of flagellar motor. *Int Rev Cell Mol Biol* **270,** 39–85 (2008).
4. Porter, S. L., Wadhams, G. H. & Armitage, J. P. Signal processing in complex chemotaxis pathways. *Nature reviews* **9,** 153–165 (2011).
5. Wolfe, A. J., Conley, M. P. & Berg, H. C. Acetyladenylate plays a role in controlling the direction of flagellar rotation. *Proc. Natl. Acad. Sci. U.S.A.* **85,** 6711–6715 (1988).

6.   Barak, R., Welch, M., Yanovsky, A., Oosawa, K. & Eisenbach, M.
Acetyladenylate or its derivative acetylates the chemotaxis protein CheY *in*
vitro and increases its activity at the flagellar switch. *Biochemistry (USA)*
31, 10099–10107 (1992).
7.   Barak, R., Abouhamad, W. N. & Eisenbach, M. Both acetate kinase and
acetyl Coenzyme A synthetase are involved in acetate-stimulated change
in the direction of flagellar rotation in *Escherichia coli*. *J. Bacteriol.* **180,**
985–988 (1998).
8.   Barak, R., Prasad, K., Shainskaya, A., Wolfe, A. J. & Eisenbach, M.
Acetylation of the chemotaxis response regulator CheY by acetyl-CoA
synthetase from *Escherichia coli*. *J. Mol. Biol.* **342,** 383–401 (2004).
9.   Li, R. *et al.* CobB regulates *Escherichia coli* chemotaxis by deacetylating
the response regulator CheY. *Mol. Microbiol.* **76,** 1162–1174 (2010).
10.   Barak, R. & Eisenbach, M. Acetylation of the response regulator, CheY, is
involved in bacterial chemotaxis. *Mol. Microbiol.* **40,** 731–743 (2001).
11.   Welch, M., Oosawa, K., Aizawa, S. I. & Eisenbach, M. Phosphorylation-
dependent binding of a signal molecule to the flagellar switch of bacteria.
Proc. Natl. Acad. Sci. U.S.A. **90,** 8787–8791 (1993).
12.   Bren, A. & Eisenbach, M. The N terminus of the flagellar switch protein,
FliM, is the binding domain for the chemotactic response regulator, CheY.
J. Mol. Biol. **278,** 507–514 (1998).
13.   Sarkar, M. K., Paul, K. & Blair, D. Chemotaxis signaling protein CheY binds
to the rotor protein FliN to control the direction of flagellar rotation in
Escherichia coli. *Proc. Natl. Acad. Sci. U.S.A.* **107,** 9370–9375 (2010).
14.   Mathews, M. A. A., Tang, H. L. & Blair, D. F. Domain analysis of the FliM
protein of *Escherichia coli*. *J. Bacteriol.* **180,** 5580–5590 (1998).
15.   Barak, R. & Eisenbach, M. Correlation between phosphorylation of the
chemotaxis protein CheY and its activity at the flagellar motor.
Biochemistry (USA) **31,** 1821–1826 (1992).
16.   McEvoy, M., Bren, A., Eisenbach, M. & Dahlquist, F. W. Identification of the
binding interfaces on CheY for two of its targets, the phosphatase CheZ
and the flagellar switch protein FliM. *J. Mol. Biol.* **289,** 1423–1433 (1999).
17.   Dyer, C. M., Vartanian, A. S., Zhou, H. & Dahlquist, F. W. A molecular
mechanism of bacterial flagellar motor switching. *J. Mol. Biol.* **388,** 71–84
(2009).
18.   Cluzel, P., Surette, M. & Leibler, S. An ultrasensitive bacterial motor
revealed by monitoring signaling proteins in single cells. *Science* **287,**
1652–1655 (2000).
19.   Sagi, Y., Khan, S. & Eisenbach, M. Binding of the chemotaxis response
regulator CheY to the isolated, intact switch complex of the bacterial
flagellar motor: Lack of cooperativity. *J. Biol. Chem.* **278,** 25867–25871
(2003).
20.   Sourjik, V. & Berg, H. C. Binding of the *Escherichia coli* response regulator
CheY to its target measured *in vivo* by fluorescence resonance energy
transfer. *Proc. Natl. Acad. Sci. U.S.A.* **99,** 12669–12674 (2002).

21. Berg, H. C. & Brown, D. A. Chemotaxis inEscherichia colianalysed by
three-dimensional tracking. *Nature* **239,** 500–504 (1972).

22. Turner, L., Ryu, W. S. & Berg, H. C. Real-time imaging of fluorescent
flagellar filaments. *J. Bacteriol.* **182,** 2793–2801 (2000).

23. Vladimirov, N., Lebiedz, D. & Sourjik, V. Predicted auxiliary navigation
mechanism of peritrichously flagellated chemotactic bacteria. *PLoS*
computational biology **6,** e1000717 (2010).

24. Saragosti, J. *et al.* Directional persistence of chemotactic bacteria in a
traveling concentration wave. *Proc. Natl. Acad. Sci. U.S.A.* **108,** 16235–
16240 (2011).

25. Kihara, M. & Macnab, R. M. Cytoplasmic pH mediates pH taxis and weak-
acid repellent taxis of bacteria. *J. Bacteriol.* **145,** 1209–1221 (1981).

26. Repaske, D. R. & Adler, J. Change in intracellular pH of *Escherichia coli*
mediates the chemotactic response to certain attractants and repellents. *J.*
Bacteriol. **145,** 1196–1208 (1981).

27. Dyer, C. M. & Dahlquist, F. W. Switched or not?: the structure of
unphosphorylated CheY bound to the N terminus of FliM. *J. Bacteriol.* **188,**
7354–7363 (2006).

28. Schuster, M., Silversmith, R. E. & Bourret, R. B. Conformational coupling in
the chemotaxis response regulator CheY. *Proc. Natl. Acad. Sci. U.S.A.* **98,**
6003–6008 (2001).

29. de Diego Puente, T. *et al.* The Protein Acetyltransferase PatZ from
Escherichia coli Is Regulated by Autoacetylation-induced Oligomerization.
J. Biol. Chem. **290,** 23077–23093 (2015).

30. Barak, R. & Eisenbach, M. Chemotactic-like response of *Escherichia coli*
cells lacking the known chemotaxis machinery but containing
overexpressed CheY. *Mol. Microbiol.* **31,** 1125–1137 (1999).

31. Bourret, R. B., Hess, J. F. & Simon, M. I. Conserved aspartate residues
and phosphorylation in signal transduction by the chemotaxis protein
CheY. *Proc. Natl. Acad. Sci. U.S.A.* **87,** 41–45 (1990).

32. Kentner, D. & Sourjik, V. Dynamic map of protein interactions in
theEscherichia colichemotaxis pathway. *Mol Syst Biol* **5,** 238 (2009).

33. Scharf, B. E., Fahrner, K. A., Turner, L. & Berg, H. C. Control of direction of
flagellar rotation in bacterial chemotaxis. *Proc. Natl. Acad. Sci. U.S.A.* **95,**
201–206 (1998).

34. Saragosti, J., Silberzan, P. & Buguin, A. Modeling E. coli tumbles by
rotational diffusion. Implications for chemotaxis. *PLoS ONE* **7,** e35412
(2012).

35. Fraiberg, M. *et al.* CheY's acetylation sites responsible for generating
clockwise flagellar rotation in *Escherichia coli*. *Mol. Microbiol.* **95,** 231–244
(2015).

36. Bai, F. *et al.* Conformational spread as a mechanism for cooperativity in the
bacterial flagellar switch. *Science* **327,** 685–689 (2010).

37. Di Paolo, D., Afanzar, O., Armitage, J. P. & Berry, R. M. Single-molecule
imaging of electroporated dye-labelled CheY in live Escherichia coli.

Philosophical Transactions of the Royal Society of London Series B -
Biological Sciences **371,** 20150492 (2016).

38. Schuster, M., Zhao, R., Bourret, R. B. & Collins, E. J. Correlated switch
binding and signaling in bacterial chemotaxis. *J. Biol. Chem.* **275,** 19752–
19758 (2000).

39. Liarzi, O. *et al.* Acetylation represses the binding of CheY to its target
proteins. *Mol. Microbiol.* **76,** 932–943 (2010).

40. Paul, K., Brunstetter, D., Titen, S. & Blair, D. F. A molecular mechanism of
direction switching in the flagellar motor of *Escherichia coli*. *Proc. Natl.*
Acad. Sci. U.S.A. **108,** 17171–17176 (2011).

41. Delalez, N. J. *et al.* Signal-dependent turnover of the bacterial flagellar
switch protein FliM. *Proc. Natl. Acad. Sci. U.S.A.* **107,** 11347–11351
(2010).

42. Philippe, N., Alcaraz, J.-P., Coursange, E., Geiselmann, J. & Schneider, D.
Improvement of pCVD442, a suicide plasmid for gene allele exchange in
bacteria. *Plasmid* **51,** 246–255 (2004).

43. Silverman, M. & Simon, M. Flagellar rotation and the mechanism of
bacterial motility. *Nature* **249,** 73–74 (1974).

44. Thomas, D. R., Francis, N. R., Xu, C. & DeRosier, D. J. The three-
dimensional structure of the flagellar rotor from a clockwise-locked mutant
ofSalmonella entericaserovar Typhimurium. *J. Bacteriol.* **188,** 7039–7048
(2006).

45. Goldbeter, A. & Koshland, D. E. An amplified sensitivity arising from
covalent modification in biological systems. *Proc. Natl. Acad. Sci. U.S.A.*
78, 6840–6844 (1981).

Acknowledgments

We are indebted to Dr. David Blair and Paige Wheatley in his group for numerous
extensive discussions, for critical reading of earlier versions of the manuscript, for
disclosing their unpublished data, and for preparing some of the mutants used in
this study. We thank Dr. James E. Ferrell for his contribution in the construction of
the mathematical model, Dr. Ady Vaknin and Vered Frank in his group for their
useful suggestions and technical assistance, Dr. S. Roy Caplan for helpful
discussions, Vladimir Kiss for assistance with operating the confocal microscope,
Dr. Judith Armitage for supplying bacterial strains, and Yana Gurevich for technical
assistance. This study was supported by grant no. 2013197 from the US-Israel
Binational Science Foundation and by grant no. 66/14 from the Israel Science
Foundation to M.E., by a UK BBSRC grant no. BB/H01795X/1 and an ERC Starter
grant no. ERC 261227 to A.N.K., and by a BBSRC grant to R.M.B. D.D.P. was
supported by a UK EPSRC DTC studentship. A.P. was supported by a UK EPSRC
DTA studentship and the German National Academic Foundation (Studienstiftung)

and the Phizackerley Senior Scholarship in Medical Sciences at Balliol College,
Oxford. O.A. was supported by an EMBO short term travel fellowship to Oxford.
771

Figures

Figure 1. CheY binds to low-affinity sites at the motor to produce clockwise
rotation. **A**, An electron density map of the switch complex of the flagellar motor of
Salmonella, a relative of *E. coli*, that demonstrates the organization within the
switch. The map was produced by [44] and downloaded from
http://dx.doi.org/10.1093/nar/gkv1126. **B**, CheY interactions. **C**, Experimental
scheme of the FRET experiment. When CheY-mCherry is in proximity to FliM-
YPet, there is energy transfer from the excited YPet fluorophore to mCherry, as a
result of which excitation of YPet results in red fluorescence of mCherry. The
attractant serine lowers the phosphorylation level of CheY with a consequent
dissociation from FliM. Therefore, following attractant addition, FRET is reduced
and excitation of YPet results in yellow emission typical of YPet fluorescence. **D**,
Representative FRET measurements of cells in response to an attractant stimulus

(0.1 mM serine). (See Figure S1 for the detailed analysis and Figure S2 for other
measurements.) FRET ratio is the ratio of mCherry to YPet fluorescence. CheY-
mCherry and mCherry concentrations were ~170 µM in the case of FliM$_{\Delta N}$ and ~15
µM in the case of wild-type FliM (Figure S3A for calibration of CheY concentration).
Data were smoothed with a mean window at the size of 10 data points. **E**,
Experimental scheme for measurement of tethered motors: Motors (red point) are
tethered to glass in a flow chamber *via* anti-flagellin antibody (red). When observed
from above, the direction of cell rotation indicates the direction of motor rotation.
F, The response of tethered *fliM*$_{\Delta N}$ Δ*cheZ* cells (strain EW635), induced for CheY
expression from a plasmid (200 µM IPTG), to positive chemotactic stimulation.
Lines and shaded regions of the average fraction of time spent in clockwise rotation
are the mean ± SEM. The arrow shows the time of stimulation. N is the number of
cells. MeAsp and leucine were used at 1 mM, benzoate (pH 7.0) at 50 mM. **G**, As
in F, but for negative chemotactic stimulation. **H**, Response of tethered *fliM*$_{\Delta N}$ cells
(strain sPW416), induced for CheY expression from a plasmid (800 µM IPTG), to
acetate and benzoate (both at 50 mM, pH 7.0). Lines and shaded regions of the
average fraction of time spent in clockwise rotation are the mean ± SEM. Down-
and up-arrows mark the introduction and the removal of stimuli, respectively.

804
Figure 2. FliM$_N$ is an activator of CheY. **A**, Dependence of clockwise rotation on
the levels of CheY~P and FliM$_N$-CheY~P. The figure shows the mean of clockwise
rotation in tethered *fliM$_{\Delta N}$ ΔcheZ* cells expressing CheY (red) or FliM$_N$-CheY (blue;
strains EW694 and EW697, respectively) and *fliM$_{\Delta N}$ ΔcheA* cells expressing CheY
(purple) or FliM$_N$-CheY (cyan; strains EW693 and EW696, respectively), with IPTG
as an inducer. For FliM$_N$-CheY~Ac (green), acetate (10 mM, pH 7.0) was
employed. Each data point is the average of all measurements at a given IPTG
concentration, weighted by the sample number of each experiment. For data, see
Table S1. **B**, Distributions of the data in A. **C**, Mean response of *fliM$_{\Delta N}$ ΔcheA* cells
expressing FliM$_N$-CheY (EW696) to introduction or removal of varying acetate
concentrations (pH 7.0). FliM$_N$-CheY expression was induced with 800 μM IPTG.
Black lines mark the apparent biphasic response to acetate removal. N is the
number of cells. **D**, Changes in clockwise rotation induced by acetate, calculated

from C as the difference in clockwise levels between time segments 0-20 s and
280-300 s (stimulus introduction; red) or between time segments 580-600 s and
300-320 s (stimulus removal; blue). Black line is a Hill fit. **E**, Comparison of the
mean response of cells to acetate and benzoate. Details as in C, but different cells
were examined. Acetate and benzoate at 10 mM (pH 7.0). **F**, Mean response of
fliM$_{\Delta N}$ cells expressing FliM$_N$-CheY (sPW417) to the introduction or removal of
acetate, benzoate (each at 50 mM, pH 7.0), leucine and MeAsp (each at 1 mM).
FliM$_N$-CheY expression was induced with 800 μM IPTG. **G**, Dependence of
clockwise rotation on the levels of CheY(D13K) and FliM$_N$-CheY(D13K). As in A-B
for tethered *fliM$_{\Delta N}$* Δ*cheA* cells expressing CheY(D13K) or FliM$_N$-CheY(D13K)
(strains EW737 and EW739, respectively).

Figure 3. CheY binding to the motor is biphasic. **A**, Experimental scheme: the ratio
of YPet fluorescence before and after mCherry photobleaching indicates the extent
of CheY binding by FliM$_{\Delta N}$. **B**, Demonstration of FRET acquisition, showing the
enhancement of FliM$_{\Delta N}$-YPet fluorescence due to mCherry photobleaching. **C**, One
representative experiment out of 3 (black points and curve). Purple data points and
curves show a control performed in parallel with free mCherry. FRET efficiency
was calculated as the ratio of yPET fluorescence before and after photobleaching
of mCherry. The data points are the mean ± SEM of 100 observations each. Curve
is a fit to a double Hill expression (n_{H1} = 2, n_{H2} = 4). **D**, FRET response curve as a
function of the CheY-mCherry concentration. Black curve is the average fit of three
different normalized experiments to a double-Hill expression; shaded region is ±
SEM. Red and blue curves were derived from the black curve. The conversion of
mCherry fluorescence intensity to CheY concentration was estimated according to
Figure S3B-C.

Figure 4. Clockwise generation is a biphasic process initiated by CheY binding to FliN. **A**, A demonstration of clockwise interval measurement. Red points, axes of rotation; black, the cell body. Clockwise rotation is apparent between ~50 and 90 ms. **B**, Demonstration of clockwise intervals considered valid and invalid (left and right, respectively). **C**, Distribution of clockwise interval lengths at different average clockwise levels in $\Delta cheA$ cells containing FliM$_{\Delta N}$ motors and expressing FliM$_N$-CheY (strain EW696), in the presence of acetate (10 mM, pH 7.0). Color gradient changes in 0.1 steps from 0 to 1 clockwise rotation. The black lines show the distributions associated with the first and second phases. **D**, Demonstration of a calculation of clockwise rotation time of each phase at each clockwise level. Distributions as in D were fitted with a biexponential decay expression. In each distribution and for each phase of exponential decay, the average interval length was calculated as the exponential pre-factor divided by the decay rate. **E**, Average interval length in each of the phases as a function of clockwise level for the distributions in C. **F**, Average reversal frequency as a function of clockwise level.

Same cells shown in C. **G**, Small deflection angles of swimming in cells that exhibit
mostly a single switching phase. The distributions shown herein are the angle
deflection distributions of free-swimming cells in Figure S6, after subtracting the
control distributions of cells that do not express CheY (in which deflection angles
represent noise). For clarity, the distributions were normalized according to the
maximal value of the distribution. The experimental cells were $\Delta cheA$ containing
FliM$_{\Delta N}$ motors and expressing FliM$_N$-CheY (strain EW696). The control cells were
$\Delta cheY$ (strain UU1631). The measurements were performed in the presence and
absence of acetate (10 mM, pH 7.0). **H**, Effects of FliM$_N$-CheY acetylation and
phosphorylation on clockwise rotation of tethered cells containing FliM$_{\Delta N}$
FliN(A93D) motors. Acetate and benzoate concentrations were 10 mM each (pH
7.0). Non-phosphorylated and phosphorylated CheY was produced by using
$\Delta cheA$ and $\Delta cheZ$ backgrounds (strains EW713 and EW714, respectively). Each
data point is the average of all measurements at a given IPTG concentration,
weighted by the sample number of each experiment. Distributions were drawn with
respect to experiments with same range of average clockwise rotation. For data
see Table S1. **I-L**, As in C, E, F, except that in I, J, L the cells used were $\Delta cheZ$ (I,
J, L) or $\Delta cheA$ (K) containing FliM$_{\Delta N}$ FliN(A93D) motors and expressing FliM$_N$-
CheY (strain EW714 or EW713, respectively).

Figure 5. The second switching phase is associated with CheY binding to FliM$_M$.
A, CheY crosslinks with FliM$_{ΔN}$-YPet. Cells overexpressing CheY or FliM$_N$-CheY
from a plasmid were crosslinked with glutaraldehyde. The cytoplasm of the cells
was resolved by SDS-PAGE. The gel was imaged for FliM$_{ΔN}$-YPet fluorescence.
The figure shows a gel where three different samples of CheY and FliM$_N$-CheY
were resolved. Full details as in Figure S7. **B**, Docking simulation between the
predicted binding interfaces of CheY and FliM$_M$ in *E. coli*. The solvent accessible
surface of FliM$_M$ is shown in green, with a binding depression area and its
surrounding ridge highlighted in yellow and brown, respectively. **Up-left**, The
predicted interaction of activated CheY with FliM$_M$ in *E. coli*. CheY is depicted as a
beige ribbon, with residues shown as ball and stick corresponding to the interacting
residues in *T. maritima*. Mg^{2+} ions are shown as magenta spheres. **Down-right**,
The active form of the β4α4 loop of CheY is predicted to promote CheY interaction
with FliM$_M$. The predicted CheY-FliM$_M$ complex in *E. coli* rotated by approximately
180° about the vertical axis aligned with the inactivated conformation of CheY

(orange coil). The region in the inactivated conformation that clashes with $FliM_M$
was previously predicted to acquire an active conformation in the case of
CheY(K91) acetylation [35]. **C**, Clockwise rotation of tethered *fliM(R94S)$_{\Delta N}$ $\Delta cheZ$*
cells or *fliM(R94L)$_{\Delta N}$ $\Delta cheZ$* cells (strain EW731 or EW733, respectively)
expressing $FliM_N$-CheY in the presence of acetate (10 mM, pH 7.0). $FliM_N$-CheY
expression was induced by 8 mM IPTG. For data see Table S1. **D**, Clockwise
rotation of tethered *fliM*(E214W)$_{\Delta N}$ $\Delta cheZ$ cells at various levels of IPTG for the
induction of $FliM_N$-CheY expression. Each data point is the average of all
measurements at similar IPTG concentration, weighted by the sample number of
each experiment. For data see Table S1. **E**, The distribution of clockwise intervals
for *fliM(R94SL)$_{\Delta N}$ $\Delta cheZ$* cells expressing $FliM_N$-CheY in the presence of acetate
(10 mM, pH 7). **F**, Average interval length in each of the phases as a function of
clockwise level for the distributions in E. **G**, Average reversal frequency as a
function of clockwise level. Same cells shown in E. **H**, A histogram of clockwise
rotation velocities at different clockwise levels. Same cells shown in E. **I-L**, Same
as E-H, except that cells of *fliM*(E214W)$_{\Delta N}$ $\Delta cheZ$ expressing $FliM_N$-CheY were
used in the absence of acetate.

912
Figure 6. CheY~Ac and CheY~P differ in their effective concentration and motor
binding kinetics. **A**, Demonstration of single-molecule tracking in a single cell. Left,
Imaged FliM-YPet fluorescence. White dots indicate the estimated motor locations.
White circles illustrate a 75 nm radius around these locations. Middle, Contour map
of the normalized sum of probabilities of the localization events of CheY(I95V)-
Atto647 (from blue to red is low to high; black is zero probability). White circles
show motor locations from the left panel. Right, Bright-field image of the cell. **B**,
Survival probabilities phosphorylated CheY(I95V)-Atto647 at the motor under
acetylating and non-acetylating conditions (presence or absence of acetate [50
mM, pH 7.0], green and blue, respectively). CheY was mostly phosphorylated due
to the Δ*cheZ* background (strain EW669). Dashed lines are confidence bounds of
the distribution at 95%. The number of cells recorded was 82 and 40 (without and
with acetate, respectively). The number of trajectories in which CheY was found to

interact with FliM was 2414 and 2660 (without and with acetate). **C**, As in B for a
Δ*cheA* background (strain EW668). The number of cells recorded was 63 and 56
(without and with acetate). The number of trajectories in which CheY was found to
interact with FliM was 1316 and 1904 (without and with acetate). **D**, Response of
tethered *fliM*$_{ΔN}$ Δ*cheA* cells (strain EW634) to acetate (50 mM, pH 7.0) at different
CheY concentrations. Lines and shaded regions are mean ± SEM. **E**, Response
of tethered *fliM*$_{ΔN}$ Δ*cheZ* cells (strain EW635) to acetate. Details as in D. **F**,
Contribution of acetylation and phosphorylation to switching to clockwise rotation.
The points are the mean clockwise rotation calculated from D, E at time segments
0-20 s and 160-180 s.

Figure 7. A model for clockwise generation by CheY~P and CheY~Ac. The model
suggests the most likely interactions of CheY at the motor. Yellow motor ring, FliN.
Red motor ring, FliM in a conformation that cannot bind CheY. Green motor ring,
FliM in a conformation that can bind CheY. Red background region is
counterclockwise rotation. Yellow background region is frequent switching. Green
shaded region is consecutive clockwise rotation. See text for details.

Figure 8. Kinetic model demonstrates varying levels of ultrasensitivity produced
by a gateway system. **A**, The kinetic model. Yellow ring represents CheY binding
site at FliN. The ring at the middle of the motor represents CheY binding sites in
$FliM_M$, when the motor is rotating counterclockwise (red) or clockwise (green). **B**,
Binding curve solution of the kinetic model for different K_3 values (binding to both
FliN and $FliM_M$). K_1 was set as 1. K_2 was set as 10^6 K_1. Different colors are solutions
for different K_3 values. Ligand levels were normalized to half-saturation point
(EC_{50}) of each curve. Note that normalized ligand concentrations are shown in log-
scale. The solution of the model was generated in Mathematica. ***Insert***, Calculated
apparent Hill-coefficient for the curves in the main panel. Apparent Hill-coefficient
was calculated as $\log_{10}(81)/\log_{10}(EC_{90}/EC_{10})$ [45].