Curious connections: white matter pathways supporting individual differences in epistemic and perceptual curiosity

Abbreviated title: Curious connections

Ashvanti Valji[1], Alisa Priemysheva[1], Carl J. Hodgetts[1], Alison G. Costigan[1],

Greg D. Parker[2,1], Kim S. Graham[1], Andrew D. Lawrence[1], Matthias J. Gruber[1]

[1]Cardiff University Brain Research Imaging Centre (CUBRIC), School of Psychology, Cardiff University, Cardiff, United Kingdom, CF24 4HQ.

[2]Experimental MRI Centre (EMRIC), School of Bioscience, Cardiff University, Cardiff, United Kingdom, CF10 3AX.

Corresponding Authors:

Ashvanti Valji, ValjiA@cardiff.ac.uk;

Matthias J. Gruber, GruberM@cardiff.ac.uk;

Number of pages: 33

Number of Tables: 0

Number of Figures: 5

Number of words for Abstract: 244

Number of words for Introduction: 636

Number of words for Discussion: 1484

Conflict of interests: The authors declare no competing financial interests

Acknowledgements: We would like to thank the funders of this research as well as Ofer Pasternak for providing the free-water corrected pipeline, John Evans and Peter Hobden for scanning support, Samuel Ridgeway and Bethany Coad for support in data collection, Sonya Foley and Marie-Lucie Read for support in data pre-processing.

1 **Abstract**

2 Across the lifespan, curiosity motivates us to learn, yet curiosity varies strikingly between

3 individuals. Such individual differences have been shown for two distinct dimensions of

4 curiosity: *epistemic curiosity* (EC), the desire to acquire knowledge about facts, and

5 *perceptual curiosity* (PC), the desire for sensory information. It is not known, however,

6 whether both dimensions of curiosity depend on different brain networks and whether inter-

7 individual differences in curiosity depend on variation in anatomical connectivity within these

8 networks. Here, we investigated the neuroanatomical connections underpinning individual

9 variation in trait curiosity. Fifty-one female participants underwent a two-shell diffusion MRI

10 sequence and completed questionnaires measuring EC and PC. Using deterministic

11 spherical deconvolution tractography we extracted microstructural metrics (fractional

12 anisotropy (FA) and mean diffusivity (MD)) from two key white matter tracts: the fornix

13 (implicated in novelty processing, exploration, information seeking and episodic memory)

14 and the inferior longitudinal fasciculus (ILF) (implicated in semantic learning and memory). In

15 line with our predictions, we found that EC – but not PC – correlated with ILF microstructure.

16 Fornix microstructure, in contrast, correlated with both EC and PC with posterior

17 hippocampal fornix fibres - associated with posterior hippocampal network connectivity -

18 linked to PC specifically. These findings suggest that differences in distinct dimensions of

19 curiosity map systematically onto specific white matter tracts underlying well characterized

20 brain networks. Furthermore, the results pave the way to study the anatomical substrates of

21 inter-individual differences in dimensions of trait curiosity that motivate the learning of

22 distinct forms of knowledge and skills.

Significance statement

Despite recent interest in curiosity states and the broad spectrum of variation in stable tendencies to experience or express curiosity, the biological correlates of trait curiosity are unknown. Here, we found that specific types of curiosity correlate with microstructure of specific white matter tracts in the brain - the inferior longitudinal fasciculus and the fornix. Our findings on the relationship between specific aspects of curiosity and anatomical connections underlying well characterized brain networks highlight the specificity of trait curiosity. Furthermore, our findings pave the way to further understand inter-individual differences in curiosity and which aspects of curiosity benefit language, memory and other cognitive processes cultivating a deeper knowledge and skill set.

Introduction

Curiosity is described as the desire for new information that motivates seeking out and acquiring knowledge (Loewenstein, 1994; Litman, 2005). The momentary experience of curiosity can be seen as a motivational state that facilitates knowledge acquisition (Silvia & Kashdan, 2009; Gottlieb and Oudeyer, 2018). Consistent with this idea, studies have shown that states of high curiosity enhance long-term memory (Kang et al., 2009; Gruber et al., 2014; McGillivray et al., 2015; Marvin and Shohamy, 2016; Stare et al., 2018; Galli et al., 2018). Furthermore, recent neuroimaging evidence suggests that state curiosity enhances memory via increased activation in the mesolimbic dopaminergic circuit including the hippocampus (Gruber et al., 2014; Kang et al., 2009). Notably, the positive effects of state curiosity on memory have been found to greatly vary between individuals in that individual variations observed in the midbrain and hippocampus activity predict the magnitude of memory enhancements (Gruber et al., 2014).

Over the last decades, between-person differences in curiosity as a personality trait (i.e. tendencies to experience and express curiosity) have been well characterized. Based on Berlyne's (1954) suggestion that different types of curiosity are aroused by opportunities for new knowledge or sensory stimulation, trait curiosity has been split into two broad facets: curiosity as engagement with semantic knowledge - *epistemic curiosity* (EC); or as engagement with sensory stimuli - *perceptual curiosity* (PC). Building on Loewenstein's (1994) model of aversive curiosity, Litman and colleagues further proposed that these two aspects of curiosity can be further separated into diversive/interest-based and specific/deprivation-based curiosity. Diversive/interest curiosity is linked to positive affect and is thought to energize and direct exploration with the ultimate goal of stimulating one's interest and reduce boredom. In contrast, specific/deprivation curiosity is accompanied by a negative, frustrated feeling of information deprivation and uncertainty, associated with a specific knowledge gap, that people are motivated to eliminate (Berlyne, 1966; Litman, 2005, 2008, Litman and Spielberger, 2003; Litman and Jimerson, 2004).

The neuroanatomical substrates underpinning individual differences in trait curiosity

are unknown. Studies investigating higher-order personality traits subsuming curiosity,

however, provide a fruitful starting point to investigate the neuroanatomical connections

underlying trait curiosity (DeYoung, 2014; Woo et al., 2014). For example, Privado et al.

(2017) found an association between 'openness to experience' and microstructure of the

inferior longitudinal fasciculus (ILF), a ventral, temporo-occipital association tract implicated

in semantic memory (Herbet et al., 2018; Hodgetts et al., 2015, 2017). Additionally, Cohen et

al. (2009) showed that individual differences in novelty seeking were associated with

microstructure of the fornix, a key pathway that connects the hippocampus - involved in

novelty detection, exploration, information seeking and episodic memory (O'Keefe and

Nadel, 1978; Kumaran and Maguire, 2009; Murray et al., 2017; Voss et al., 2017) - to the

thalamus, ventral striatum, amygdala and prefrontal cortex (Saunders and Aggleton, 2007;

Aggleton et al. 2015).

Here, we used multi-shell diffusion MRI and spherical deconvolution tractography to

investigate whether individual differences in ILF and fornix microstructure would be

associated with individual differences in trait curiosity. Given the importance of ILF to

semantic cognition (Hodgetts et al., 2017; Ripollés et al., 2017; Herbet et al., 2018), we

predicted an association between ILF microstructure and EC but not PC. In contrast, given

that hippocampal circuitry supports novelty detection, exploratory behaviour and information

seeking in many domains (O'Keefe and Nadel, 1978; Kumaran and Maguire, 2009; Murray

et al., 2017; Voss et al., 2017) we predicted an association between fornix microstructure

and *both* EC and PC. Further, given evidence of a posterior (fine-grained) to anterior (gist-

based) gradient of representational specialization along the long-axis of the hippocampus

(Ranganath and Ritchey, 2012; Poppenk et al. 2013; Strange et al., 2014; Murray et al.,

2017), we predicted that fornical fibres associated with posterior and anterior hippocampus

(Christiansen et al., 2017; Saunders and Aggleton, 2007) would be more strongly associated

with PC and EC, respectively.

Materials and Methods

Participants

Fifty-one healthy female adult undergraduate students, with a mean age of 20 years (SD ±

1, range = 19-24) participated. They provided written consent prior to participating in the

study, which was approved by the Cardiff University Research Ethics Committee, and

received a remuneration of approximately £25 for their participation.

Trait curiosity measures

Participants completed the *Epistemic Curiosity Scale (EC)* (Litman, 2008) and the

Perceptual Curiosity Scale (PC) (Collins et al., 2004), along with other self-report measures

not relevant to the current study. The EC scale consists of five interest EC items and five

deprivation EC items with participants answering on a scale ranging from 1 (*almost never*) to

4 (*almost always*). The interest EC items are associated with behaviours that stimulate

positive affect, or involve learning something completely new (e.g. "I enjoy learning about

subjects that are unfamiliar to me"). In contrast, deprivation EC items describe behaviours

that reduce negative feelings of information deprivation and uncertainty (e.g. "I can spend

hours on a single problem because I just can't rest without knowing the answer"). The PC

scale (Collins et al., 2004) comprised of twelve items (6 diversive PC items and 6 specific

PC items) and again participants respond on a scale that ranged from 1 (*almost never*) to 4

(*almost always*). The diversive PC items describe exploratory behaviours in which one seeks

out new places and a broad range of sensory stimulation (e.g. "I like to discover new places

to go"), whereas specific PC describes exploration of novel, specific and sensorially

stimulating stimuli (e.g. "When I hear a strange sound, I usually try to find out what caused

it"). The Cronbach's alpha coefficients for the scales were all >= .70 suggesting good

internal consistency.

Imaging acquisition

Imaging data were obtained at CUBRIC, Cardiff University on a 3 Tesla MRI scanner (Siemens Magnetom Prisma) with a 32-channel head coil. T1-weighted structural 3D images were acquired using an MPRAGE sequence (orientation = sagittal; TR = 2250ms; TE = 3.06ms; TI = 900ms; flip angle = 9°; FOV = 256mm²; slice thickness = 1mm; voxel size = 1mm³; number of slices = 224; bandwidth = 230Hz/pixel; total acquisition time = 7 minutes 36 seconds).

Diffusion weighted images were acquired using a multi-shell sequence (orientation = transversal/axial; TR = 9400ms; TE = 67.0ms; FOV = 256mm²; slice thickness = 2mm; voxel size = 2mm³; number of slices = 80). Diffusion gradients were applied in (i) 30 isotropic directions by using a diffusion-weighted factor b=1200sec/mm², (ii) in 60 isotropic directions by using a diffusion-weighted factor b=2400sec/mm², and (iii) a volume without diffusion gradients (b=0sec/mm²) (bandwidth = 1954Hz/pixel; total acquisition time = 15 minutes 51 seconds).

Diffusion MRI pre-processing

T1-weighted structural images were subjected to a 'brain-tissue only' mask using FSL's Brain Extraction Tool (RRID:SCR_002823; Smith, 2002). Using ExploreDTI (RRID:SCR_001643; v4.8.3; Leemans et al., 2009) each b-value image was then co-registered to the T1 structural image. Subsequently, all b-value images were corrected for head motion and eddy currents within ExploreDTI. Tensor fitting was conducted on the b-1200 data given the tensor model assumes hindered (Gaussian) diffusion, and at lower b-values more of the signal is due to hindered rather than restricted diffusion (Jones et al., 2013). To correct for voxel-wise partial volume artefacts arising from free water contamination, the two-compartment 'Free Water Elimination' (FWE) procedure was applied to the current b-1200 data – this improves reconstruction of white matter tracts near the

ventricles such as the fornix (Pasternak et al., 2009, 2014), yielding whole brain voxel-wise

free-water corrected FA and MD tissue maps. Following FWE, corrected diffusion tensor-

derived structural metrics were computed. Fractional anisotropy (FA), reflects the extent to

which diffusion within biological tissue is anisotropic (constrained along a single axis). MD

(10^{-3} mm^2 s^{-1}) reflects overall degree of diffusivity (Vettel et al., 2017). The resulting free

water corrected FA and MD maps were inputs for the tractography analysis.

Tractography

As higher b-values allow for better fibre orientation estimations (Vettel et al., 2017), we

performed tractography on the b-2400 data using damped Richardson-Lucy spherical

deconvolution (dRL-SD). Spherical deconvolution provides a direct estimate of the

underlying distribution of fibre orientations in the brain and when applied to tractography

leads to accurate reconstructions of the major white matter pathway, and an improved ability

to describe complex white matter anatomy (Dell'Acqua and Tournier, 2018). The algorithm

extracted peaks in the fibre orientation density function (fODF) at the centre of each voxel,

where streamlines along the orientation of the fODF peaks were reconstructed using a step

size of 0.5mm. Streamline tracts were terminated if the direction of the pathway changed

through an angle greater than 45° or if the fODF threshold fell below 0.05.

In ExploreDTI, manual tractography was carried out using AND, NOT, and SEED

ROI gates on colour-coded FA maps to extract specific white matter tracts. AND gates

(**Figure 1** - green) were used to extract fibres that passed through the gate, NOT gates

(**Figure 1** - red) were used to exclude any fibres that passed through the gate, and finally

SEED gates (**Figure 1** - blue) were used as a starting point to extract fibres that passed

through this gate and then to include only those fibres that then passed through any added

AND gates. Manual tractography was carried out on a minimum of 15 subjects in order to

calculate a tract model to perform automated tractography on all 51 data sets (Explore DTI;

Parker et al., 2013). This procedure enables the construction of white matter tracts in space

in which streamlines belonging to particular anatomical features of interest consistently

project to distinct sub-regions, allowing the reconstruction of streamline data by observing

their projected positions (Parker et al., 2013). After running the automated tractography

software each tract was visually inspected, and any erroneous fibres were pruned using

additional NOT gates. These tract masks from the b=2400 data were then intersected with

the b=1200 free-water corrected FA and MD maps to derive free-water corrected tract-

specific measures of FA and MD values for statistical analysis.

Inferior Longitudinal Fasciculus tractography. The ILF (**Figure 1B**) was reconstructed using

a two-ROI approach in each hemisphere (Wakana et al., 2007). In the mid-saggital slice of

the brain, the coronal crosshair was placed posterior to the corpus callosum. In the coronal

plane a SEED gate was drawn around the entire cortex of interest. Next in the coronal view,

the last slice where the temporal lobe was separate from the frontal lobe was identified and

one AND gate was drawn around the temporal lobe. Any stray fibres not consistent with the

ILF pathway were removed with NOT gates. FA and MD of the right and left ILF were

summed and averaged to provide a bilateral measure for the main analyses.

Fornix tractography. The fornix (**Figure 1A**) was traced in line with the landmarks described

in Catani and Thiebaut de Schotten (2008). In the mid saggital slice of the brain, the coronal

crosshair was placed at the anterior commissure and moved approximately 6 voxels

posterior in the brain. In the coronal plane, one AND gate was drawn around the fornix

bundle where the anterior pillars enter the body of the fornix. Finally, NOT gates were drawn

around any protruding areas that were not part of the fornix.

Anterior and posterior hippocampal fornix tractography. In addition, we employed a method

adapted from prior work to reconstruct the anterior and posterior hippocampal fornix fibres

(Christiansen et al., 2017). Both anterior and posterior hippocampal fornix reconstructions

required the AND and NOT gates that were applied during whole fornix tractography. Some
NOT gates were augmented to enable better extraction of the anterior and posterior
hippocampal streamlines of the fornix. A standard landmark for the anterior-posterior
hippocampal boundary was proposed to be a small bundle of grey matter that outlines the
most anterior extent of the parahippocampal gyrus that is called uncal apex or uncus
(Poppenk et al., 2013). This landmark was identified for each hemisphere separately when
carrying out manual tractography of the anterior and posterior hippocampal fornix. In order to
perform this, the uncus was first localised at its anterior part and traced to its posterior
boundary. The first coronal slice in which the uncus was not visible anymore was used as
the landmark in order distinguish between fibres that project into anterior (head of the
hippocampus) and posterior hippocampus (body and tail of the hippocampus) (**Figure 1C**).

After the left and right hemispheric landmarks were identified, one NOT gate on each
hemisphere was drawn around the hippocampus to set boundaries for posterior
hippocampal fornix tracts, removing fibres that pass through these NOT gates (**Figure 1D**).
After the posterior hippocampal fornix was identified, the same coordinates of the anterior-
posterior hippocampal boundary landmark for each hemisphere were used to replace the
NOT gates with AND gates for the left and right anterior hippocampal fornix reconstruction
(**Figure 1E**). The posterior, left, and right anterior hippocampal fornix were saved as
separate tracts to aid subsequent automated tractography (**Figure 2**). Note that diffusion
tensor metrics of the whole fornix and those averaged across anterior and posterior
hippocampal fornix segments were highly correlated (FA, $r(51) = 0.940$, $p < 0.001$; MD, $r(51)$
$= 0.942$, $p < 0.001$) indicating that the anterior and posterior hippocampal fornix
reconstructions matched the whole fornix reconstructions.

Figure 1. Automated tractography reconstructions of the fornix, its anterior and posterior hippocampal fornix fibres and the inferior longitudinal fasciculus (ILF). AND (green), NOT (red), and SEED (blue) ROI gates for each of the tracts are displayed on the sagittal midline plane. (A) Fornix tractography using AND and NOT gates. (B) Left ILF tractography using SEED, AND and NOT gates. (C) Location of AND and NOT gates for tractography of the anterior and posterior hippocampal fornix, respectively. (D) Posterior hippocampal fornix tractography using one additional NOT gate placed between the head and the body of the hippocampus to only include fornical fibres that connect with posterior hippocampus (i.e., hippocampal body and tail). (E) Anterior hippocampal fornix tractography using one additional AND gate placed between the head and body of the hippocampus (i.e., identical location as NOT gate in (D)) to include fibres that pass through this ROI gate to the anterior hippocampus.

A Posterior hippocampal fornix fibres

B Left anterior hippocampal fornix fibres

C Right anterior hippocampal fornix fibres

| Right | x = 75 | y = 54 | z = 46 |
| Left | x = 38 | y = 54 | z = 44 |

Posterior hippocampus

x = 39   y = 64   z = 40

Left anterior hippocampus

x = 72   y = 64   z = 42

Right anterior hippocampus

Figure 2. Automated tractography reconstructions of anterior and posterior hippocampal fornix fibres on coronal slices. Tractography of the fornix fibres projecting to the posterior hippocampus (**A**). Tractography of fornix fibres projecting to the left anterior hippocampus (**B**). Tractography of the fornix fibres projecting to the right anterior hippocampus (**C**).

Experimental Design and Statistical analyses

For the questionnaire data, in the event of missing responses (2 participants failed to give a response to one PC item), the mean value of the remaining items that were answered in the full scale was calculated which then replaced the missing item score. For each curiosity subscale (i.e., the two subscales of PC and EC), we calculated a total score for each participant. Participants' data with diffusion tensor metrics +/– 3SD beyond the group mean were considered as outliers and removed from respective analyses. This resulted in one participant's data being removed from all analyses involving ILF MD and a different participant's data being removed from analyses including bilaterally averaged ILF FA.

To test for associations between curiosity trait scores and microstructure of our selected anatomical tracts, we conducted directional *Pearson's* correlations using MATLAB

(RRID:SCR_001622). Since higher FA and lower MD is typically associated with higher

microstructural integrity (Vettel et al., 2017), we predicted a positive correlation between

levels of curiosity and FA and a negative correlation with MD.

To determine whether the *Pearson's* correlation coefficient *r* was statistically

significant, we performed non-parametric permutation tests that randomly permute the real

data between participants. Permutation tests were conducted separately for the two

microstructure metrics (i.e., FA and MD) and for the EC and PC scales. Importantly, we

corrected for multiple comparisons across the subscales within a curiosity scale (e.g.,

diversive- and specific PC). The steps were as follows: First, we performed *Pearson's*

correlations on the real data (i.e., correlations between the scores of the two curiosity

subscales and the microstructure measure (e.g., diversive PC with ILF MD and specific PC

with ILF MD)). Thereby, we obtained the empirical correlation coefficients reflecting the

relationship between the two curiosity subscales and a specific microstructure measure.

Second, within each curiosity subscale, we shuffled the curiosity scores across participants,

which resulted in pairs containing a curiosity score and a microstructure value that is

randomly assigned across participants. On these shuffled data, we then calculated surrogate

Pearson's coefficients for the two curiosity subscale scores and the microstructure metric,

and saved the maximum surrogate *Pearson's r* across the two correlations (i.e., subscale-

microstructure$_{max}$) (Groppe, Urbach and Kutas, 2011). Third, the second step was repeated

5000 times. Based on the 5000 permutations, we created a null distribution of all surrogate

subscale-microstructure$_{max}$ coefficient values and determined the alpha cut-off point

($p < 0.05$; one-sided; i.e., 4750th data point of the surrogate null distribution) in order to test

the statistical significance of the real *Pearson's* coefficients reflecting the relationship

between the two subscales and the microstructure measure. This approach allowed us to

correct for multiple comparisons across the two subscales within each curiosity scale. In

follow-up analyses for specific curiosity subscales (e.g., interest EC subscale), we also

performed follow-up permutation tests that corrected for multiple comparisons across both

hemispheres (e.g., left and right ILF MD). The 95% confidence intervals (CI) for each

correlation was derived using a bootstrapping method based on 1000 iterations.

Results

Epistemic curiosity – but not perceptual curiosity – correlates with ILF microstructure

ILF FA. We conducted a series of permutation tests that investigated the relationships

between curiosity trait scores and microstructure in *a-priori* selected anatomical tracts. For

each permutation test, we corrected for multiple comparisons for the two subscales

separately within the EC and PC scale. The first permutation test targeted ILF FA and EC.

We found that bilaterally averaged ILF FA did not significantly correlate with either subscale

of EC (deprivation EC, $r(50) = 0.143$, $p_{corr} = 0.243$, 95% CI [-0.105, 0.364]; interest EC, $r(50)$

$= 0.191$, $p_{corr} = 0.151$, 95% CI [-0.0734, 0.440]). A further permutation test was conducted on

bilaterally averaged ILF FA with the two subscales of PC, where again neither subscale

significantly correlated with bilateral ILF FA (specific PC, $r(50) = 0.109$, $p_{corr} = 0.329$, 95% CI

[-0.229, 0.427]; diversive PC, $r(50) = 0.207$; $p_{corr} = 0.122$, 95% CI [-0.109, 0.453]).

ILF MD. Targeting ILF MD, a permutation test revealed a significant negative correlation

between ILF MD and deprivation EC ($r(50) = -0.388$, $p_{corr} = 0.004$, 95% CI [-0.572, -0.124],

Figure 3A) and a significant negative correlation between ILF MD and interest EC ($r(50) =$

-0.289, $p_{corr} = 0.038$, 95% CI [-0.489, 0.007], **Figure 3B**). In contrast, bilateral ILF MD was

not significantly correlated with any subscale of PC (specific PC, $r(50) = -0.134$, $p_{corr} = 0.267$,

95% CI [-0.392, 0.157]); diversive PC ($r(50) = 0.020$, $p_{corr} = 0.710$, 95% CI [-0.260, 0.271]).

Figure 3. Inferior longitudinal fasciculus microstructure only shows relationship with epistemic curiosity. These results were obtained from non-parametric permutation tests that corrected for multiple comparisons across the two subscales within the Epistemic Curiosity scale (EC). A significant relationship was found between MD of the inferior longitudinal fasciculus (ILF) with deprivation- and interest EC (**A**, **B**, respectively). The line of best fit is shown on each scatter plot with 50 data points.

Neuropsychological evidence suggests that semantic knowledge is represented bilaterally in the anterior temporal lobes (ATL) but may show subtle inter-hemispheric (left > right) gradations for verbal stimuli (Rice et al., 2015). Therefore, we asked whether the significant correlation between bilateral ILF MD and both EC subscales were driven specifically by the left as opposed to the right ILF. Separate permutation tests were conducted for each subscale of EC with left ILF MD and right ILF MD as the two variables of interest (i.e., correcting for multiple comparisons across the two hemispheres). The first permutation test on deprivation EC found that both left and right ILF MD significantly correlated with deprivation EC (left ILF: $r(50)$ = -0.341, p_{corr} = 0.016, 95% CI [-0.566, -0.078]; right ILF: $r(50)$ = -0.358, p_{corr} = 0.012, 95% CI [-0.564, -0.106]). The second permutation test investigating whether interest EC correlates with left and right ILF MD indicated a numerical negative relationship for both tracts but neither reached significance with the adopted multiple comparisons correction (left ILF: $r(50)$ = -0.254, p_{corr} = 0.066, 95% CI [-0.491, 0.086]); right ILF: $r(50)$ = -0.267, p_{corr} = 0.051, 95% CI [-0.472, -0.056]).

Interest epistemic curiosity correlates with fornix microstructure

Fornix FA. Regarding fornix FA, permutation tests revealed a significant positive correlation between interest EC and fornix FA (r(51) = 0.281, p_{corr} = 0.039, 95% CI [-0.008, 0.491], **Figure 4**). In contrast, deprivation EC showed no significant correlation with fornix FA (r(51) = 0.155, p_{corr} = 0.214, 95% CI [-0.120, 0.422]). A second permutation test was conducted on fornix FA with the two subscales of PC, diversive and specific, but neither subscale significantly correlated with fornix FA (specific PC, r(51) = 0.111, p_{corr} = 0.328, 95% CI [-0.266, 0.4252]; diversive PC, r(51) = 0.064, p_{corr} = 0.466, 95% CI [-0.204, 0.351]).

Fornix MD. Despite the earlier findings of a significant positive correlation between interest EC and fornix FA, permutation tests revealed no significant negative correlation between fornix MD and interest EC (r(51) = -0.110, p_{corr} = 0.332, 95% CI [-0.372, 0.171]) or deprivation EC (r(51) = -0.029, p_{corr} = 0.574, 95% CI [-0.314, 0.296]). The second permutation test, investigating the association between fornix MD and the two subscales of PC, also showed that neither specific nor diversive PC significantly correlated with fornix MD (specific PC, r(51) = -0.250, p_{corr} = 0.070, 95% CI [-0.499, 0.054]; diversive PC, (r(51) = -0.159; p_{corr} = 0.214, 95% CI [-0.398, 0.113]).

Figure 4. Fornix microstructure shows relationship with interest epistemic curiosity. These results obtained from non-parametric permutation tests correcting for multiple

comparisons across subscales within the Epistemic Curiosity scales (EC). A significant relationship was found between fractional anisotropy (FA) of the whole fornix and interest EC. The line of best fit is shown on the scatterplot with 51 data points.

Specific perceptual curiosity shows relationship with posterior hippocampal fornix

Recent accounts postulate a posterior-anterior gradient of representational granularity along the long axis of the hippocampus, linked to a gradient in anatomical connectivity (Aggleton, 2012; Strange et al., 2014), from 'fine' perceptual detail to 'course' or gist-like representations (Poppenk et al., 2013; Robin and Moscovitch, 2017; Sheldon et al., 2019). This account suggests that a correlation might be evident between posterior hippocampal fornix and PC, and anterior hippocampal fornix and EC, respectively. To test this account, we explored the relationship between *specific* PC (i.e., associated with a detailed perceptual knowledge gap) and anterior/posterior hippocampal fornix MD, given that the correlation between specific PC and whole fornix MD did not reach significance in the preliminary analyses. Conversely, to pinpoint how EC is associated with the anterior/posterior hippocampal fornix FA, we focussed our analyses on *interest* EC for which we had found a positive correlation with whole fornix FA.

A first permutation test (corrected for multiple comparisons) targeted the three individual fornix segmentations (i.e., left anterior, right anterior, bilateral posterior hippocampal fornix). (Note the anterior hippocampal fornical fibres form the lateral fornix but posterior hippocampal fornical fibres form the medial fornix which cannot easily be separated into separate hemispheres). We found that specific PC significantly correlated with posterior hippocampal fornix MD (r(51) = -0.277, p_{corr} = 0.047, 95% CI [-0.528,0.056], **Figure 5**), but it did not correlate significantly with left or right anterior hippocampal fornix MD (left: (r(51 = -0.189, p_{corr} = 0.176, 95% CI [-0.451,0.062]); right: (r(51) = -0.028, p_{corr} =

371 0.610, 95% CI [-0.289,0.264]). This finding suggests that specific PC might mainly be

372 supported by fornical fibres that have connections to the posterior hippocampus.

Figure 5. Posterior hippocampal fornix microstructure shows relationship with

aspects of perceptual curiosity. Results obtained from non-parametric permutation tests

correcting for multiple comparisons across subscales within the Perceptual Curiosity scales

(PC). A significant relationship was found between MD of the posterior hippocampal fornix

(i.e., fornix fibres that project specifically into posterior hippocampus) and specific PC. The

line of best fit is shown on the scatterplot with 51 data points.

In contrast, although we found that interest EC significantly correlates with *whole*

fornix FA, the three distinct fornix segmentations did not reveal significant correlations with

interest EC (left anterior hippocampal fornix FA, r(51) = 0.269, p_{corr} = 0.065, 95% CI [-0.029,

0.521]; right anterior hippocampal fornix FA (r(51) = 0.080, p_{corr} = 0.479, 95% CI [-0.161,

0.307]); posterior hippocampal fornix FA, r(51) = 0.272, p_{corr} = 0.062, 95% CI [-0.009,

0.479]).

In summary, we found that two individual subscales that tap into epistemic and

perceptual curiosity traits showed significant correlations with fornix microstructure. In

particular, we found that whole fornix FA significantly correlated with interest EC whereas

posterior hippocampal fornix microstructure significantly correlated with specific PC.

Discussion

Curiosity motivates us to seek out information and it facilitates knowledge acquisition (Loewenstein, 1994; Litman, 2005; Silvia and Kashdan, 2009; Gottlieb and Oudeyer, 2018). While a fledgling line of research has shown that curiosity states - the momentary experience of curiosity - enhance hippocampus-dependent memory (for a review, see Gruber et al., 2019), there is also a broad spectrum of variation in *stable* tendencies to experience or express curiosity. Here, we found that ILF microstructure correlated with both interest and deprivation EC traits, but not with PC traits. Additionally, fornix microstructure was associated with interest - but not deprivation - EC, and specific - but not diverse - PC. In particular, while microstructure of the whole fornix correlated with interest EC, specific PC correlated with the posterior hippocampal fornix. These findings support the notion that curiosity is a multifaceted motivational construct and that distinct aspects of curiosity map onto specific white matter tracts underlying well-characterized brain networks.

Epistemic curiosity and ILF

The ILF, which connects ventral aspects of ATL, occipito-temporal, and occipital cortex (Herbet et al., 2018), appears critical for bidirectional interactions between an ATL-based bilateral semantic 'hub' and representations supported by occipital and middle/posterior temporal regions (Patterson et al., 2007; Lambon Ralph et al., 2017; Chen et al., 2017). In addition to demonstrations of altered ILF microstructure in semantic dementia (Agosta et al., 2010), recent studies report associations between ILF microstructure and individual differences in semantic learning (Ripollés et al., 2017) and memory (Hodgetts et al., 2017). Here, we found that participants with reduced diffusivity (i.e., lower MD values) in the ILF showed higher trait scores in both dimensions of EC. Critically, we found that the ILF supported both the general exploration of semantic information motivated by positive affect (EC as a feeling-of-interest) but also the search for specific information in order to close a knowledge gap (EC as an aversive feeling-of-deprivation) (Litman, 2005, 2008;

Loewenstein, 1994; Lauriola et al., 2015). One explanation for this may be that perhaps the

more that we learn, the more we are attuned to the gaps in our knowledge, and attending to

these gaps is tension-producing and enjoyable at the same time (Loewenstein, 1994). In

addition, the association between EC and ILF microstructure is in line with the literature on

the higher-order personality trait 'openness to experience', of which curiosity is one facet

(Woo et al., 2014). Privado et al. (2017) demonstrated that ILF microstructure was

associated with levels of trait 'openness'. Our findings extend this work by pinpointing that

the exploration and specific search for semantic information might be one critical factor that

carries the association between 'openness' and ILF microstructure.

Curiosity and Fornix

The hippocampus is a medial temporal lobe structure supporting the encoding and recall of

long-term memory (Burgess et al., 2002; Davachi, 2006; Eichenbaum et al., 2007; Murray et

al., 2018). Given that the hippocampus has been implicated in a number of processes critical

to curiosity, including exploration, reward seeking and novelty detection (O'Keefe and Nadel,

1978; Murray et al., 2017; Kumaran and Maguire, 2009; Voss et al 2017), we investigated

the relationship between curiosity and the fornix - the principal tract linking the hippocampus

with sites beyond the temporal lobe (Postans et al., 2014; Rudebeck et al., 2009). Regarding

the relationship between curiosity and fornix microstructure, we performed analyses

targeting the microstructure of the whole fornix, but also the anterior and posterior

hippocampal fornix segments that correspond to the functional subdivisions of the anterior

and posterior hippocampus, respectively (Christiansen et al., 2017; Saunders and Aggleton,

2007). Given current theoretical ideas, the anterior and posterior hippocampal fornix fibres

may reflect functional subdivisions of the anterior and posterior hippocampus reflecting gist-

based (schematic) and detailed (episodic) information, respectively (Robin and Moscovitch,

2017; Poppenk et al., 2013; Sheldon et al., 2019). Therefore, the present study investigated

whether the functional subdivisions of the fornix, connecting to the anterior and posterior

hippocampus, may potentially map onto diversive/interest and specific/deprivation curiosity,

respectively. Consistent with this hypothesis, we found that the posterior hippocampal fornix

(but not the anterior hippocampal fornix) showed a relationship with specific PC which is

described as the desire to reduce uncertainty by searching for a specific novel perceptual

information.

In contrast, we found that interest EC positively correlated with microstructure of the whole

fornix. Interest EC is described as the desire for diversive exploration and information

seeking which is accompanied by positive affect (Litman, 2008). Given these ideas on the

functional relevance of interest EC and theoretical ideas and evidence about anterior

hippocampal functions, we would have expected that interest EC would show a relationship

with anterior hippocampal fornix (i.e., which is strongly connected to anterior hippocampus).

The anterior hippocampus has, however, also been proposed to support gist-based and

schematic information (Poppenk et al., 2013; Robin and Moscovitch, 2017). Although

interest EC reflects the general, explorative search for semantic information, interest EC also

triggers search for detailed information rather than gist-based information. Therefore,

interest EC might also depend on more posterior hippocampal regions in which detailed

'cognitive maps' of the environment, as a means to obtain information, are formed and

transition in a graded fashion along the long axis to more gist-based global-contextual

'cognitive maps' (Aggleton, 2012; Graham et al., 2010; Murray et al., 2018; Poppenk et al.,

2013; Strange et al., 2014). In line with this idea of a hippocampal gradient, we found that

interest EC did not reach significance for correlations with the left anterior and the posterior

hippocampal fornix, but was found to significantly correlate with the whole fornix reflecting

the idea that perhaps interest EC depends on fornix fibres that stretch along the whole of the

long axis of the hippocampus aligned to EC's dependence on integrative conceptual and

detailed maps.

Our study involved questionnaires to tap into distinct curiosity traits. Other recent studies

that investigated individual differences in curiosity utilized measures of eye-movement as an

objective indicator of visual exploration. For instance, Baranes and colleagues (2015), found

curiosity-based enhancement of anticipatory gaze correlated with trait curiosity, and Risko et

al. (2012) used a scene-viewing task to demonstrate that participants' PC trait score

predicted the degree to which they explored the scenes. These studies using eye-

movements to investigate curiosity-based exploration and our present findings on fornix

microstructure highlight how individual differences in curiosity play a critical part in the

degree of exploration of one's environment.

Limitations and future directions

First, our correlation analyses cannot establish causality of one variable over the other.

Longitudinal studies would be needed to determine the causality of these relationships to

investigate whether trait curiosity shapes white matter connections, vice versa, or whether

both reinforce each other in a bidirectional manner. For instance, recent work on adaptive

myelination suggests that change in myelination through activity-dependent adaptation of an

initially hard-wired process is in response to experiences and contributes to learning

(Bechler et al., 2018). Second, interpreting the biological relevance of diffusion metrics from

white matter tracts, such as FA and MD, can be challenging. Whilst FA and MD are believed

to be inversely related where typically a high FA and low MD suggest greater white matter

microstructure (Vettel et al., 2017), we found that for the majority of microstructure-curiosity

correlations that only one of the two diffusion metrics significantly correlated with curiosity.

These dissociations could be due to the structure of the selected white matter tracts. For

example, evidence suggests that FA has been found to be less consistent along a given

tract (Yeatman et al., 2012) and other studies also found similar variability in FA and MD

measures of the fornix and ILF (e.g., Hodgetts et al., 2017). Inconsistences between FA and

MD measures could also be due to a number of biological properties such as axon diameter

and density, myelination and the arrangement of fibres in a given voxel (Beaulieu, 2002). For

instance, high FA has been found to reflect high myelin density and structured histological

orientation whereas high values of MD is more likely to reflect low myelin density and diffuse

histological orientation (Seehaus et al., 2015). This emphasizes the need to use more

sophisticated methods in future studies, such as Neurite Orientation Dispersion and Density

Imaging, a white matter index which is more informative about brain cellular microstructure

than FA and MD alone.

Conclusion

The present study found inter-individual variation in the microstructure of the fornix related to

interest EC and inter-individual variation in the microstructure of the ILF related to both

interest and deprivation EC. Furthermore, posterior hippocampal fornix microstructure was

associated with specific PC. In conclusion, our findings on the relationship between curiosity

traits and anatomical connections underlying well characterized brain networks provide a

foundation for future studies to examine the relationship between curiosity traits, curiosity

states and their neuroanatomical substrates. Our findings pave the way to further

understand inter-individual differences in curiosity and which aspects of curiosity benefit

language, memory and other cognitive processes cultivating a deeper knowledge and skill

set.

Author contributions

A.V., K.S.G., A.D.L. and M.J.G. contributed to the conception and design of the experiment. A.V. and A.C. contributed to data acquisition. All authors contributed to data analysis and interpretation. A.V. and M.J.G. drafted the manuscript and together with C.J.H., K.S.G. and A.D.L. revised the manuscript. A.D.L. and M.J.G. jointly supervised this work.

Funding

This work was supported by a departmental PhD studentship from the School of Psychology at Cardiff University to A.V., a PhD studentship from the Cardiff University Neuroscience and Mental Health Research Institute (NMHRI) to A.C., a Wellcome Strategic Award (104943/Z/14/Z) to C.J.H, A.C, K.G., A.D.L., and a COFUND fellowship funded by the Welsh Government and the European Commission and a Sir Henry Dale Fellowship (211201/Z/18/Z) funded by Wellcome and the Royal Society to M.J.G.

References

Aggleton JP, Wright NF, Rosene, DL, Saunders RC (2015) Complementary patterns of
direct amygdala and hippocampal projections to the macaque prefrontal cortex. Cerebral
Cortex 25(11):4351-4373.

Aggleton JP (2012) Multiple anatomical systems embedded within the primate medial
temporal lobe: implications for hippocampal function. Neuroscience & Biobehavioral
Reviews 36(7):1579-1596.

Agosta F, Henry RG, Migliaccio R, Neuhaus J, Miller BL, Dronkers NF, Brambata SM, Filippi
M, Ogar JM, Wilson SM, Gorno-Tempini ML (2010) Language networks in semantic
dementia. Brain 133(1):286-299.

Baranes A, Oudeyer PY, Gottlieb J (2015) Eye movements reveal epistemic curiosity in
human observers. Vision Research 117:81–90.

Beaulieu C (2002) The basis of anisotropic water diffusion in the nervous system - A
technical review. NMR in Biomedicine 15(7-8):435–455.

Bechler ME, Swire M, ffrench-Constant C (2018) Intrinsic and adaptive myelination—A
sequential mechanism for smart wiring in the brain. Developmental neurobiology 78(2):68-
79.

Berlyne DE (1954) A theory of human curiosity. British Journal of Psychology 3: 180-191.

Berlyne DE (1966) Curiosity and Exploration. Science 153(3731):25–33.

Burgess N, Maguire EA, O'Keefe J (2002) The human hippocampus and spatial and
episodic memory. Neuron 35(4):625-641.

Catani M, Thiebaut de Schotten M (2008) A diffusion tensor imaging tractography atlas for
virtual in vivo dissections. Cortex 44(8):1105–1132.

Chen Y, Chen K, Ding J, Zhang Y, Yang Q, Lv Y, Guo Q, Han Z (2017) Brain network for the
core deficits of semantic dementia: a neural network connectivity-behavior mapping
study. Frontiers in human neuroscience 11:267.

Christiansen K, Metzler-Baddeley C, Parker GD, Muhlert N, Jones DK, Aggleton JP, Vann

SD (2017) Topographic separation of fornical fibers associated with the anterior and

posterior hippocampus in the human brain: An MRI-diffusion study. Brain and Behavior

7(1):1-8.

Cohen MX, Schoene-Bake JC, Elger CE, Weber B (2009) Connectivity-based segregation of

the human striatum predicts personality characteristics. Nature Neuroscience 12(1):32-34.

Collins RP, Litman JA, Spielberger CD (2004) The measurement of perceptual curiosity.

Personality and Individual Differences 36(5):1127-1141.

Davachi L (2006) Item, context and relational episodic encoding in humans. Current opinion

in neurobiology 16(6):693-700.

Dell'Acqua F, Tournier JD (2018) Modelling white matter with spherical deconvolution: How

and why?. NMR in Biomedicine 32(4):e3945.

DeYoung CG (2014) Openness/Intellect: A dimension of personality reflecting cognitive

exploration. In: APA handbook of personality and social psychology: Personality processes

and individual differences (Cooper ML, Larsen RJ, eds), pp369-399. Washington, DC:

American Psychological Association.

Eichenbaum H, Yonelinas AP, Ranganath C (2007) The medial temporal lobe and

recognition memory. Annual Review of Neuroscience 30:123-152.

Galli G, Sirota M, Gruber MJ, Ivanof BE, Ganesh J, Materassi M, Thorpe A, Loaiza V,

Cappelletti M, Craik FIM (2018) Learning facts during aging: the benefits of

curiosity. Experimental aging research 44(4):311-328.

Gottlieb J, Oudeyer PY (2018) Towards a neuroscience of active sampling and curiosity.

Nature Reviews Neuroscience 19:758-770.

Graham KS, Barense MD, Lee AC (2010) Going beyond LTM in the MTL: a synthesis of

neuropsychological and neuroimaging findings on the role of the medial temporal lobe in

memory and perception. Neuropsychologia 48(4):831-853.

Gruber MJ, Gelman BD, Ranganath C (2014) States of Curiosity Modulate Hippocampus-

Dependent Learning via the Dopaminergic Circuit. Neuron 84(2):486-496.

Gruber MJ, Valji A, Ranganath C (2019) Curiosity and Learning: A Neuroscientific

Perspective. In: The Cambridge Handbook of Motivation and Learning. (Renninger KA, Hidi

SE, ed), pp 397-417. Cambridge University Press.

Herbet G, Zemmoura I, Duffau H (2018) Functional Anatomy of the Inferior Longitudinal

Fasciculus: From Historical Reports to Current Hypotheses. Frontiers in

neuroanatomy 12:77.

Hodgetts CJ, Postans M, Shine JP, Jones DK, Lawrence AD, Graham KS (2015)

Dissociable roles of the inferior longitudinal fasciculus and fornix in face and place

perception. Elife 4:e07902.

Hodgetts CJ, Postans M, Warne N, Varnava A, Lawrence AD, Graham KS (2017) Distinct

contributions of the fornix and inferior longitudinal fasciculus to episodic and semantic

autobiographical memory. Cortex 94:1-14.

Jones DK, Knösche TR, Turner R (2013) White matter integrity, fiber count, and other

fallacies: the do's and don'ts of diffusion MRI. Neuroimage 73:239-254.

Kang MJ, Hsu M, Krajbich IM, Loewenstein G, McClure SM, Wang JT, Camerer CF (2009)

The wick in the candle of leraning. Psychological Science 20(8):963-974.

Kumaran D, Maguire EA (2009) Novelty signals: a window into hippocampal information

processing. Trends in cognitive sciences 13(2):47-54.

Lambon Ralph M, Jefferies E, Patterson K, Rogers TT (2017) The neural and computational

bases of semantic cognition. *Nature Reviews Neuroscience 18*:42-55.

Lauriola M, Litman JA, Mussel P, De Santis R, Crowson HM, Hoffman RR (2015) Epistemic

curiosity and self-regulation. Personality and Individual Differences 83(1):202-207.

Leemans A, Jeurissen B, Sijbers J, Jones DK (2009) ExploreDTI: a graphical toolbox for

processing, analyzing, and visualizing diffusion MR data. Proceedings of the 17th Annual

Meeting of International Society for Magnetic Resonance in Medicine 17:3536.

Litman JA (2005) Curiosity and the pleasures of learning: Wanting and liking new

information. Cognition and Emotion 19:793-814

Litman JA (2008) Interest and deprivation factors of epistemic curiosity. Personality and

Individual Differences 44(7):1585-1595.

Litman JA, Jimerson TL (2004) The measurement of curiosity as a feeling of deprivation.

Journal of Personality Assessment 82(2):147-157.

Litman J, Spielberger C (2003) Measuring epistemic curiosity and its diversive and specific

components. Journal of Personality Assessment 80(1):75-86.

Loewenstein G (1994) The psychology of curiosity: A review and reinterpretation.

Psychological Bulletin 116(1):75-98.

Marvin CB, Shohamy D (2016) Curiosity and reward: Valence predicts choice and

information prediction errors enhance learning. Journal of Experimental Psychology: General

145(3):266-272.

McGillivray S, Murayama K, Castel AD (2015) Thirst for knowledge: The effects of curiosity

and interest on memory in younger and older adults. Psychology and Aging 30(4):835-841.

Murray EA, Wise SP, Graham KS (2017) The evolution of memory systems: ancestors,

anatomy, and adaptations. Oxford University Press.

Murray EA, Wise SP, Graham KS (2018) Representational specializations of the

hippocampus in phylogenetic perspective. Neuroscience letters 680:4-12.

O'Keefe J, Nadel L (1978) The hippocampus as a cognitive map. Oxford: Clarendon Press.

Parker GD, Marshall D, Rosin PL, Drage N, Richmond S, Jones DK (2013) Fast and fully

automated clustering of whole brain tractography results using shape-space analysis.

Proceedings of the International Society for Magnetic Resonance in Medicine. Salt Lake

City, USA, 778.

Pasternak O, Maier-Hein K, Baumgartner C, Shenton ME, Rathi Y, Westin CF (2014) The

estimation of free-water corrected diffusion tensors. In: Visualization and Processing of

Tensors and Higher Order Descriptors for Multi-Valued Data (Westin CF, Vilanova A,

Burgeth B, eds), pp. 249-270. Springer, Berlin, Heidelberg.

Pasternak O, Sochen N, Gur Y, Intrator N, Assaf Y (2009) Free water elimination and
mapping from diffusion MRI. Magnetic Resonance in Medicine 62(3):717-730.

Patterson K, Nestor PJ, Rogers TT (2007) Where do you know what you know? The
representation of semantic knowledge in the human brain. Nature Reviews
Neuroscience 8(12):976-987.

Poppenk J, Evensmoen HR, Moscovitch M, Nadel L (2013) Long-axis specialization of the
human hippocampus. Trends in cognitive sciences 17(5):230-240.

Postans M, Hodgetts CJ, Mundy ME, Jones DK, Lawrence AD, Graham KS (2014)
Interindividual variation in fornix microstructure and macrostructure is related to visual
discrimination accuracy for scenes but not faces. Journal of Neuroscience 34(36):12121-
12126.

Privado J, Román FJ, Saénz-Urturi C, Burgaleta M, Colom R (2017) Gray and white matter
correlates of the Big Five personality traits. Neuroscience, 349, 174-184.

Ranganath C, Ritchey M (2012) Two Cortical Systems for Memory-Guided Behaviour.
Nature Reviews. Neuroscience 13(10):713-26.

Rice GE, Lambon Ralph MA, Hoffman P (2015) The roles of left versus right anterior
temporal lobes in conceptual knowledge: an ALE meta-analysis of 97 functional
neuroimaging studies. Cerebral Cortex 25(11):4374-4391.

Ripollés P, Biel D, Peñaloza C, Kaufmann J, Marco-Pallarés J, Noesselt T, Rodríguez-
Fornells A (2017) Strength of Temporal White Matter Pathways Predicts Semantic Learning.
The Journal of Neuroscience 37(46):1720-17.

Risko EF, Anderson NC, Lanthier S, Kingstone A (2012) Curious eyes: Individual differences
in personality predict eye movement behavior in scene-viewing. Cognition 122(1):86-90.

Robin J, Moscovitch M (2017) Details, gist and schema: hippocampal–neocortical
interactions underlying recent and remote episodic and spatial memory. Current Opinion in
Behavioral Sciences 17:114-123.

Rudebeck SR, Scholz J, Millington R, Rohenkohl G, Johansen-Berg H, Lee AC (2009)

Fornix microstructure correlates with recollection but not familiarity memory. Journal of

Neuroscience 29(47):14987-14992.

Saunders RC, Aggleton JP (2007) Origin and topography of fibers contributing to the fornix

in macaque monkeys. Hippocampus 17(5):396-411.

Seehaus A, Roebroeck A, Bastiani M, Fonseca L, Bratzke H, Lori N, Vilanova A, Goebel R,

Galuske R (2015) Histological validation of high-resolution DTI in human post mortem

tissue. Frontiers in Neuroanatomy 9:98.

Sheldon S, Fenerci C, Gurguryan L (2019) A Neurocognitive Perspective on the Forms and

Functions of Autobiographical Memory Retrieval. Frontiers in Systems Neuroscience 13:4.

Silvia PJ, Kashdan TB (2009) Interesting things and curious people: Exploration and

engagement as transient states and enduring strengths. Social and Personality Psychology

Compass 3(5):785-797.

Smith SM (2002) Fast robust automated brain extraction. Human brain mapping 17(3):143-

155.

Stare CJ, Gruber MJ, Nadel L, Ranganath C, Gómez RL (2018) Curiosity-driven memory

enhancement persists over time but does not benefit from post-learning sleep. Cognitive

Neuroscience 9:100-115.

Strange BA, Witter MP, Lein ES, Moser EI (2014) Functional organization of the

hippocampal longitudinal axis. Nature Reviews Neuroscience 15(10):655.

Vettel JM, Cooper N, Garcia JO, Yeh F-C, Verstynen TD (2017) White Matter Tractography

and Diffusion-Weighted Imaging. eLS:1-9.

Voss JL, Bridge DJ, Cohen NJ, Walker JA (2017) A closer look at the hippocampus and

memory. Trends in cognitive sciences 21(8):577-588.

Wakana S, Caprihan A, Panzenboeck MM, Fallon JH, Perry M, Gollub RL, Hua K, Zhang J,

Jiang H, Dubey P, Blitz A, van Zijl P, Mori S (2007) Reproducibility of quantitative

tractography methods applied to cerebral white matter. Neuroimage 36(3):630-644.

Woo SE, Chernyshenko OS, Longley A, Zhang Z-X, Chiu C-Y, Stark SE (2014) Openness to

experience: Its lower level structure, measurement, and cross-cultural equivalence. Journal

of Personality Assessment 96(1):29-45.

Yeatman JD, Dougherty RF, Myall NJ, Wandell BA, Feldman HM (2012) Tract profiles of

white matter properties: automating fiber-tract quantification. PloS one 7(11):e49790.

    Figure Legends

Figure 1. Automated tractography reconstructions of the fornix, its anterior and
posterior hippocampal fornix fibres and the inferior longitudinal fasciculus (ILF). AND
(green), NOT (red), and SEED (blue) ROI gates for each of the tracts are displayed on the
sagittal midline plane. (**A**) Fornix tractography using AND and NOT gates. (**B**) Left ILF
tractography using SEED, AND and NOT gates. (**C**) Location of AND and NOT gates for
tractography of the anterior and posterior hippocampal fornix, respectively. (**D**) Posterior
hippocampal fornix tractography using one additional NOT gate placed between the head
and the body of the hippocampus to only include fornical fibres that connect with posterior
hippocampus (i.e., hippocampal body and tail). (**E**) Anterior hippocampal fornix tractography
using one additional AND gate placed between the head and body of the hippocampus (i.e.,
identical location as NOT gate in (D)) to include fibres that pass through this ROI gate to the
anterior hippocampus.

Figure 2.   Automated tractography reconstructions of anterior and posterior
hippocampal fornix fibres on coronal slices. Tractography of the fornix fibres projecting
to the posterior hippocampus (**A**). Tractography of fornix fibres projecting to the left anterior
hippocampus (**B**). Tractography of the fornix fibres projecting to the right anterior
hippocampus (**C**).

Figure 3. Inferior longitudinal fasciculus microstructure only shows relationship with
epistemic curiosity. These results were obtained from non-parametric permutation tests
that corrected for multiple comparisons across the two subscales within the Epistemic
Curiosity scale (EC). A significant relationship was found between MD of the inferior
longitudinal fasciculus (ILF) with deprivation- and interest EC (**A**, **B**, respectively). The line of
best fit is shown on each scatter plot with 50 data points.

Figure 4. Fornix microstructure shows relationship with interest epistemic curiosity. These results obtained from non-parametric permutation tests correcting for multiple comparisons across subscales within the Epistemic Curiosity scales (EC). A significant relationship was found between fractional anisotropy (FA) of the whole fornix and interest EC. The line of best fit is shown on the scatterplot with 51 data points.

Figure 5. Posterior hippocampal fornix microstructure shows relationship with aspects of perceptual curiosity. These results obtained from non-parametric permutation tests correcting for multiple comparisons across subscales within the Perceptual Curiosity scales (PC). A significant relationship was found between MD of the posterior hippocampal fornix (i.e., fornix fibres that project specifically into posterior hippocampus) and specific PC The line of best fit is shown on the scatterplot with 51 data points.