White matter pathways supporting individual differences in epistemic and perceptual curiosity

Ashvanti Valji[1], Alisa Priemysheva[1], Carl J. Hodgetts[1], Alison G. Costigan[1],

Greg D. Parker[2,1], Kim S. Graham[1], Andrew D. Lawrence[1], Matthias J. Gruber[1]

[1]Cardiff University Brain Research Imaging Centre (CUBRIC), School of Psychology, Cardiff University, Cardiff, United Kingdom, CF24 4HQ.

[2]Experimental MRI Centre (EMRIC), School of Bioscience, Cardiff University, Cardiff, United Kingdom, CF10 3AX.

Corresponding Authors:

Ashvanti Valji, ValjiA@cardiff.ac.uk;

Matthias J. Gruber, GruberM@cardiff.ac.uk;

Abstract

Across the lifespan, curiosity motivates us to learn, yet curiosity varies strikingly between

individuals. Such individual differences have been shown for two distinct dimensions of

curiosity: *epistemic curiosity* (EC), the desire to acquire conceptual knowledge, and

perceptual curiosity (PC), the desire for sensory information. It is not known, however,

whether both dimensions of curiosity depend on different brain networks and whether inter-

individual differences in curiosity depend on variation in anatomical connectivity within these

networks. Here, we investigated the neuroanatomical connections underpinning individual

variation in trait curiosity. Fifty-one female participants underwent a two-shell diffusion MRI

sequence and completed questionnaires measuring EC and PC. Using deterministic

spherical deconvolution tractography we extracted microstructural metrics (fractional

anisotropy (FA) and mean diffusivity (MD)) from two key white matter tracts: the fornix

(implicated in novelty processing, exploration, information seeking and episodic memory)

and the inferior longitudinal fasciculus (ILF) (implicated in semantic learning and memory). In

line with our predictions, we found that EC – but not PC – correlated with ILF microstructure.

Fornix microstructure, in contrast, correlated with both EC and PC, with posterior

hippocampal fornix fibres - associated with posterior hippocampal network connectivity -

linked to PC specifically. These findings suggest that differences in distinct dimensions of

curiosity map systematically onto specific white matter tracts underlying well characterized

brain networks. Furthermore, the results pave the way to study the anatomical substrates of

inter-individual differences in dimensions of trait curiosity that motivate the learning of

distinct forms of knowledge and skills.

Introduction

Curiosity is described as the desire for new information that motivates seeking out and

acquiring knowledge (Loewenstein, 1994; Litman, 2005). The momentary experience of

curiosity (state curiosity) can be seen as an emotional-motivational state that facilitates

exploration and knowledge acquisition (Silvia & Kashdan, 2009; Gottlieb & Oudeyer, 2018).

Consistent with this idea, studies have shown that states of high curiosity enhance long-term

memory (Kang et al., 2009; Gruber et al., 2014; McGillivray et al., 2015; Marvin & Shohamy,

2016; Stare et al., 2018; Galli et al., 2018). Furthermore, recent neuroimaging evidence

suggests that state curiosity enhances memory via increased activation in the mesolimbic

dopaminergic circuit including the hippocampus (Gruber et al., 2014; Kang et al., 2009).

Notably, the positive effects of state curiosity on memory have been found to greatly vary

between individuals in that individual variations observed in the midbrain and hippocampus

activity predict the magnitude of memory enhancements (Gruber et al., 2014).

Over the last decades, between-person differences in curiosity as a personality trait

(i.e. dispositional tendencies to experience and express curiosity) have been well

characterized. Based on Berlyne's (1954) suggestion that different types of curiosity are

aroused by opportunities for new knowledge or sensory stimulation, trait curiosity has been

split into two broad facets: curiosity as engagement with semantic knowledge - *epistemic*

curiosity (EC); or as engagement with sensory stimuli - *perceptual curiosity* (PC). Building on

Loewenstein's (1994) model of aversive curiosity, Litman and colleagues further proposed

that these two aspects of curiosity can be further separated into diversive/interest-based and

specific/deprivation-based curiosity. Diversive/interest curiosity is linked to positive affect

and is thought to energize and to direct exploration with the ultimate goal of stimulating one's

interest and reduce boredom. In contrast, specific/deprivation curiosity is accompanied by a

negative, frustrated feeling of information deprivation and uncertainty, associated with a

specific knowledge gap, that people are motivated to eliminate (Berlyne, 1966; Litman, 2005,

2008, Litman & Spielberger, 2003; Litman & Jimerson, 2004). Importantly, such inter-

individual differences in curiosity have been found to predict job performance and academic

achievement in the real world (Grossnickle, 2016; Kashdan & Yuen, 2007; Mussel, 2013).

The neuroanatomical substrates underpinning individual differences in trait curiosity

are unknown. Studies investigating higher-order personality traits subsuming curiosity,

however, provide a fruitful starting point to investigate the neuroanatomical connections

underlying trait curiosity (DeYoung, 2014; Woo et al., 2014). For example, Privado et al.

(2017) found an association between 'openness to experience' and microstructure of the

inferior longitudinal fasciculus (ILF), a ventral, temporo-occipital association tract implicated

in semantic learning and memory (Herbet et al., 2018; Hodgetts et al., 2015, 2017; Ripollés

et al., 2017). Additionally, Cohen et al. (2009) showed that individual differences in novelty

seeking were associated with microstructure of the fornix, a key pathway that connects the

hippocampus - involved in novelty detection, exploration, information seeking and episodic

memory (O'Keefe & Nadel, 1978; Kumaran & Maguire, 2009; Murray et al., 2017; Voss et

al., 2017) - to the thalamus, ventral striatum, amygdala and prefrontal cortex (Saunders &

Aggleton, 2007; Aggleton et al. 2015).

Here, we used multi-shell diffusion MRI and spherical deconvolution tractography to

investigate whether individual differences in ILF and fornix microstructural metrics (i.e.,

fractional anisotropy (FA) and mean diffusivity (MD)) would be associated with individual

differences in trait curiosity. Given the importance of ILF to semantic cognition (Jouen et al.,

2015; Chen et al., 2017; Hodgetts et al., 2017; Ripollés et al., 2017; Herbet et al., 2018), we

predicted an association between ILF microstructure and EC but not PC. In contrast, given

that hippocampal circuitry supports novelty detection, exploratory behaviour and information

seeking in many domains (O'Keefe & Nadel, 1978; Kumaran & Maguire, 2009; Otmakhova

et al., 2013; Murray et al., 2017; Voss et al., 2017) we predicted an association between

fornix microstructure and *both* EC and PC. Further, given evidence of a posterior (fine-

grained) to anterior (gist-based) gradient of representational specialization along the long-

axis of the hippocampus (Ranganath & Ritchey, 2012; Poppenk et al. 2013; Strange et al.,

2014; Murray et al., 2017), we predicted that microstructure of fornical fibres associated with

posterior and anterior hippocampus (Christiansen et al., 2017; Saunders & Aggleton, 2007)

would be more strongly associated with PC and EC, respectively.

Results

Epistemic curiosity – but not perceptual curiosity – correlates with ILF microstructure

ILF FA. We conducted a series of permutation tests that investigated the relationships

between trait curiosity scores and microstructure in *a-priori* selected anatomical tracts. For

each permutation test, we corrected for multiple comparisons for the two subscales

separately within the EC and PC scale. The first permutation test targeted ILF FA and EC.

We found that bilaterally averaged ILF FA did not significantly correlate with either subscale

of EC (deprivation EC, $r(50) = 0.143$, $p_{corr} = 0.243$, 95% CI [-0.105, 0.364]; interest EC, $r(50)$

$= 0.191$, $p_{corr} = 0.151$, 95% CI [-0.0734, 0.440]). A further permutation test was conducted on

bilaterally averaged ILF FA with the two subscales of PC, where again neither subscale

significantly correlated with bilateral ILF FA (specific PC, $r(50) = 0.109$, $p_{corr} = 0.329$, 95% CI

[-0.229, 0.427]; diverse PC, $r(50) = 0.207$; $p_{corr} = 0.122$, 95% CI [-0.109, 0.453]).

ILF MD. Targeting ILF MD, a permutation test revealed a significant negative correlation

between ILF MD and interest EC ($r(50) = -0.289$, $p_{corr} = 0.038$, 95% CI [-0.489, 0.007],

Figure 1A) and a significant negative correlation between ILF MD and deprivation EC ($r(50)$

$= -0.388$, $p_{corr} = 0.004$, 95% CI [-0.572, -0.124], **Figure 1B**). In contrast, bilateral ILF MD was

not significantly correlated with any subscale of PC (diverse PC, $r(50) = 0.020$, $p_{corr} =$

0.710, 95% CI [-0.260, 0.271], **Figure 1C**); specific PC, *r*(50) = -0.134, p_{corr} = 0.267, 95% CI

[-0.392, 0.157], **Figure 1D**).

Figure 1. Inferior longitudinal fasciculus microstructure only shows relationship with

epistemic curiosity. These results were obtained from non-parametric permutation tests

that corrected for multiple comparisons across the two subscales within the Epistemic

Curiosity scale (EC) and Perceptual Curiosity scale (PC). A significant negative correlation

was found between MD (mm^2 s^{-1}) of the inferior longitudinal fasciculus (ILF) with interest-

and deprivation EC (**A**, **B**, respectively). No significant correlations were found between ILF

MD (mm^2 s^{-1}) with diversive and specific PC (**C**, **D**, respectively). The line of best fit and 95%

confidence interval (CI) are shown on each scatter plot with 50 data points.

Neuropsychological and imaging evidence suggests that semantic knowledge is

represented bilaterally in the anterior temporal lobes (ATL) but may show subtle inter-

hemispheric (left > right) gradations for verbal stimuli (Rice et al., 2015; Hoffman & Lambon

Ralph, 2018). Therefore, we asked whether the significant correlation between bilateral ILF

MD and both EC subscales were driven specifically by the left as opposed to the right ILF.

Separate permutation tests were conducted for each subscale of EC with left ILF MD and

right ILF MD as the two variables of interest (i.e., correcting for multiple comparisons across

the two hemispheres). The first permutation test on deprivation EC found that both left and

right ILF MD significantly correlated with deprivation EC (left ILF: $r(50) = -0.341$, $p_{corr} =$

0.016, 95% CI [-0.566, -0.078]; right ILF: $r(50) = -0.358$, $p_{corr} = 0.012$, 95% CI [-0.564, -

0.106]). The second permutation test investigating whether interest EC correlates with left

and right ILF MD indicated a numerical negative relationship for both tracts but neither

reached significance with the adopted multiple comparisons correction (left ILF: $r(50) = -$

0.254, $p_{corr} = 0.066$, 95% CI [-0.491, 0.086]); right ILF: $r(50) = -0.267$, $p_{corr} = 0.051$, 95% CI [-

0.472, -0.056]).

In order to assess whether bilateral ILF MD correlations with subsets of EC were

significantly different from each other as well as subsets of PC, we conducted directional

Olkin's Z-tests (Cocor R package; Diedenhofen & Musch, 2015). For EC, we found that the

correlation between ILF MD and deprivation EC was not significantly different to the

correlation between ILF MD and interest EC ($z(50) = 0.849$, $p = 0.198$). Comparing EC and

PC subscales, however, we found that the correlation between ILF MD and deprivation EC

was significantly stronger than the correlation between ILF MD and specific PC ($z(50) =$

1.721, $p = 0.043$), and the correlation between ILF MD and diversive PC ($z(50) = 2.212$, $p =$

0.014). Furthermore, we found that the correlation between ILF MD and interest EC was

significantly stronger than the correlation between ILF MD and diversive PC ($z(50) = 2.407$,

p = 0.008), however, the correlation between ILF MD and interest EC was not significantly

stronger than the correlation between ILF MD and specific PC (z(50) = 1.172, p = 0.121).

Interest-based epistemic curiosity correlates with fornix microstructure

Fornix FA. Regarding fornix FA, permutation tests revealed a significant positive correlation

between interest EC and fornix FA (r(51) = 0.281, p_{corr} = 0.039, 95% CI [-0.008, 0.491],

Figure 2A). In contrast, deprivation EC showed no significant correlation with fornix FA

(r(51) = 0.155, p_{corr} = 0.214, 95% CI [-0.120, 0.422], **Figure 2B**). A second permutation test

was conducted on fornix FA with the two subscales of PC, diversive and specific, but neither

subscale significantly correlated with fornix FA (specific PC, r(51) = 0.111, p_{corr} = 0.328, 95%

CI [-0.266, 0.4252]; diversive PC, r(51) = 0.064, p_{corr} = 0.466, 95% CI [-0.204, 0.351]).

Fornix MD. Despite the earlier findings of a significant positive correlation between interest

EC and fornix FA, permutation tests revealed no significant negative correlation between

fornix MD and interest EC (r(51) = -0.110, p_{corr} = 0.332, 95% CI [-0.372, 0.171]) or

deprivation EC (r(51) = -0.029, p_{corr} = 0.574, 95% CI [-0.314, 0.296]). The second

permutation test, investigating the association between fornix MD and the two subscales of

PC, also showed that neither specific nor diversive PC significantly correlated with fornix MD

(specific PC, r(51) = -0.250, p_{corr} = 0.070, 95% CI [-0.499, 0.054]; diversive PC, (r(51) = -

0.159; p_{corr} = 0.214, 95% CI [-0.398, 0.113]).

Figure 2. Fornix microstructure shows relationship with aspects of epistemic curiosity. These results were obtained from non-parametric permutation tests correcting for multiple comparisons across subscales within the Epistemic Curiosity scales (EC). A significant positive correlation was found between fractional anisotropy (FA) of the whole fornix and interest EC **(A)** but not with deprivation EC **(B)**. The line of best fit and 95% confidence interval (CI) are shown on each scatter plot with 51 data points.

Specific perceptual curiosity shows an association with posterior hippocampal fornix microstructure

Recent accounts postulate a posterior-anterior gradient of representational granularity along the long axis of the hippocampus, linked to a gradient in anatomical connectivity (Aggleton, 2012; Strange et al., 2014), from 'fine' perceptual detail to 'course' or gist-like representations (Poppenk et al., 2013; Robin and Moscovitch, 2017; Sheldon et al., 2019). This account suggests that a stronger correlation might be evident between posterior hippocampal fornix and PC, and anterior hippocampal fornix and EC, respectively. To test this, we explored the relationship between *specific* PC (i.e., associated with detailed perceptual information seeking) and anterior/posterior hippocampal fornix MD. Conversely, to pinpoint how EC is associated with the anterior/posterior hippocampal fornix FA, we focussed our analyses on *interest* EC.

A first permutation test (corrected for multiple comparisons) targeted the three

individual fornix segmentations (i.e., left anterior, right anterior, bilateral posterior

hippocampal fornix). (Note that posterior hippocampal fornical fibres form the medial fornix

cannot easily be separated into separate hemispheres). We found that specific PC

significantly correlated with posterior hippocampal fornix MD (r(51) = -0.277, p_{corr} = 0.047,

95% CI [-0.528,0.056], **Figure 3B**), but it did not correlate significantly with left or right

anterior hippocampal fornix MD (left: (r(51 = -0.189, p_{corr} = 0.176, 95% CI [-0.451,0.062]),

Figure 3A; right: (r(51) = -0.028, p_{corr} = 0.610, 95% CI [-0.289,0.264]). This finding suggests

that specific PC might mainly be supported by fornical fibres that have connections to the

posterior hippocampus. Olkin's z-tests were employed to test whether the correlation

between specific PC and posterior hippocampal fornix MD was significantly different than the

correlation between specific PC and left/right anterior hippocampal fornix MD. The

correlation between posterior hippocampal fornix MD and specific PC was not significantly

different than the correlation between *left* anterior hippocampal fornix MD (z (51) = -0.934, p

= 0.175), however, it was significantly different than the correlation between *right* anterior

hippocampal fornix MD and specific PC (z (51) = -2.268, p = 0.012).

In contrast, although we found that interest EC significantly correlates with *whole*

fornix FA, the three distinct fornix segmentations did not reveal significant correlations with

interest EC after correcting for multiple comparisons (left anterior hippocampal fornix FA,

r(51) = 0.269, p_{corr} = 0.065, 95% CI [-0.029, 0.521], **Figure 3C**; right anterior hippocampal

fornix FA (r(51) = 0.080, p_{corr} = 0.479, 95% CI [-0.161, 0.307]; posterior hippocampal fornix

FA, r(51) = 0.272, p_{corr} = 0.062, 95% CI [-0.009, 0.479], **Figure 3D**). Olkin's z-test indicated

that the correlation between left anterior hippocampal fornix FA and interest EC was not

significantly different than the correlation between posterior hippocampal fornix FA and

interest EC (z (51) = 0.031, p = 0.488). In addition, Olkin's z-test indicated that the

correlation between right anterior hippocampal fornix FA and interest EC was not

significantly stronger than the correlation between posterior hippocampal fornix FA and

interest EC (*z* (51) = 1.443, *p* = 0.075).

Figure 3. Specific perceptual curiosity, but not interest epistemic curiosity, shows a

significant difference between correlations with anterior and posterior hippocampal

fornix microstructure. These results were obtained from non-parametric permutation tests

correcting for multiple comparisons across the three individual fornix segmentations. Specific

PC did not significantly correlate with MD (mm^2 s^{-1}) of the left anterior hippocampal fornix

(i.e., fornix fibres that project specifically into anterior hippocampus) **(A)** but was found to

negatively correlate with MD (mm^2 s^{-1}) of the posterior hippocampal fornix **(B)**. Interest EC

did not significantly correlate with FA of the left anterior hippocampal fornix (**C**), nor with FA

of the posterior hippocampal fornix **(D)**. The line of best fit and 95% confidence interval (CI)

are shown on each scatter plot with 51 data points.

In summary, we found that two individual subscales that tap into epistemic and

perceptual curiosity traits showed significant correlations with fornix microstructure. In

particular, we found that *whole* fornix FA significantly correlated with interest EC suggesting

that interest EC relates to microstructure in fornical fibres that connect with anterior and

posterior hippocampus. In contrast, specific PC specifically correlated with posterior

hippocampal fornix microstructure, which was significantly stronger compared to the

relationship with right anterior hippocampal fornix microstructure.

Discussion

Curiosity motivates us to seek out information and it facilitates knowledge acquisition

(Loewenstein, 1994; Litman, 2005; Silvia & Kashdan, 2009; Gottlieb & Oudeyer, 2018).

While a fledgling line of research has shown that curiosity states - the momentary

experience of curiosity - enhance hippocampus-dependent memory (for a review, see

Gruber et al., 2019), there is also a broad spectrum of variation in *stable* tendencies to

experience or express curiosity – trait curiosity. Importantly, trait curiosity has been shown to

predict real-world outcomes, such as academic achievement and job performance (Kashdan

& Yuen, 2007; Mussel, 2013). Here, we found that ILF microstructure correlated with both

interest and deprivation EC traits, but not with PC traits. Additionally, fornix microstructure

was associated with interest - but not deprivation - EC, and specific - but not diversive - PC.

In particular, while microstructure of the whole fornix correlated with interest EC, specific PC

correlated with posterior hippocampal fornix microstructure. These findings support the

notion that curiosity is a multifaceted motivational construct and that distinct aspects of

curiosity map onto specific white matter tracts underlying well-characterized brain networks

that support distinct representational systems (Murray et al., 2017).

Epistemic curiosity and ILF microstructure

The ILF, which connects ventral aspects of ATL, occipito-temporal, and occipital cortex

(Herbet et al., 2018; Panesar et al., 2018), appears critical for bidirectional interactions

between an ATL-based bilateral semantic 'hub' and representations supported by occipital

and middle/posterior temporal regions (Patterson et al., 2007; Lambon Ralph et al., 2017;

Chen et al., 2017). In addition to demonstrations of altered ILF microstructure in semantic

dementia (Agosta et al., 2010), recent studies report associations between bilateral ILF

microstructure and individual differences in semantic learning (Ripollés et al., 2017) and

memory (Horowitz-Kraus et al., 2014; Hodgetts et al., 2017). Here, we found that

participants with reduced diffusivity (i.e., lower MD values) in the ILF showed higher trait

scores in both dimensions of EC. Critically, we found that the ILF supported both the general

exploration of semantic information motivated by positive affect (EC as a feeling-of-interest)

but also the search for specific information in order to close a knowledge gap (EC as an

aversive feeling-of-deprivation) (Litman, 2005, 2008; Loewenstein, 1994; Lauriola et al.,

2015). One explanation for this may be that perhaps the more that we learn, the more we

are attuned to the gaps in our conceptual knowledge, and attending to these gaps is tension-

producing and enjoyable at the same time (Loewenstein, 1994). In addition, the association

between EC and ILF microstructure is in line with the literature on the higher-order

personality trait 'openness to experience', of which curiosity is one facet (Woo et al., 2014).

Privado et al. (2017) demonstrated that ILF microstructure was associated with levels of trait

'openness'. Our findings extend this work by pinpointing that the exploration and specific

search for semantic information might be one critical factor that carries the association
between 'openness' and ILF microstructure.

Curiosity and Fornix microstructure

The hippocampus is a medial temporal lobe structure supporting the encoding and recall of
long-term memory (Burgess et al., 2002; Davachi, 2006; Eichenbaum et al., 2007; Murray et
al., 2018). Given that the hippocampus has been implicated in a number of processes critical
to curiosity, including exploration, reward seeking and novelty detection (O'Keefe & Nadel,
1978; Otmakhova et al., 2013; Murray et al., 2017; Kumaran & Maguire, 2009; Voss et al
2017), we investigated the relationship between curiosity and microstructure of the fornix -
the principal tract linking the hippocampus with sites beyond the temporal lobe (Saunders &
Aggleton 2007; Aggleton et al., 2015). Regarding the relationship between curiosity and
fornix microstructure, we performed analyses targeting the microstructure of the whole
fornix, but also the anterior and posterior hippocampal fornix segments that correspond to
the functional subdivisions of the anterior and posterior hippocampus, respectively
(Christiansen et al., 2017; Saunders and Aggleton, 2007). Given current theoretical ideas,
the anterior and posterior hippocampal fornix fibres may reflect functional subdivisions of the
anterior and posterior hippocampus reflecting gist-based (schematic) and perceptually
detailed (episodic) information, respectively (Robin & Moscovitch, 2017; Poppenk et al.,
2013; Ranganath & Ritchey, 2012; Sheldon et al., 2019). Therefore, the present study
investigated whether the functional subdivisions of the fornix, connecting to the anterior and
posterior hippocampus, may potentially map onto diversive/interest and specific/deprivation
curiosity, respectively. Partially consistent with this hypothesis, we found that posterior
hippocampal fornix (but not the anterior hippocampal fornix) microstructure (FA) showed an
association with specific PC, which is described as the desire to reduce uncertainty by
searching for specific novel perceptual information. Of note, recent work has highlighted a
role for (posterior) HC circuitry in detailed visual exploration (Liu et al., 2017; Voss et al.,

2017) and Risko et al. (2012) used a scene-viewing task to demonstrate that participants'

PC trait score predicted the degree to which they explored visual scenes. These studies

using eye-movements to investigate hippocampal- and curiosity-based visual exploration

and our present findings on fornix microstructure highlight how individual differences in

curiosity may play a critical part in the degree of exploration of one's perceptual

environment, serving to accumulate information from the visual world, contributing to the

formation of detailed memory representations mediated by posterior hippocampal circuitry.

In contrast, we found that interest EC positively correlated with microstructure of the

whole fornix, rather than anterior hippocampal fornix specifically. Interest EC is described as

the desire for diversive exploration and information seeking, which is accompanied by

positive affect (Litman, 2008). Although interest EC reflects the reward-driven explorative

search for new knowledge, presumably involving interactions between anterior hippocampal

schematic or gist-based representations and reward/value representations mediated by

nucleus accumbens and ventromedial prefrontal cortex (Poppenk et al., 2013; Aggleton et

al., 2015), interest EC also triggers search for detailed information rather than gist-based

information, presumably involving more fine-grained posterior hippocampal representations.

Interest EC may therefore involve coordination along the entire hippocampal longitudinal

axis, in line with the graded and overlapping nature of long axis connectivity (Aggleton,

2012; Strange et al., 2014).

Limitations and future directions

First, our correlational analyses cannot establish causality in brain-behaviour relationships.

Longitudinal studies would be needed to determine whether trait curiosity shapes white

matter connections, vice versa, or whether both reinforce each other in a bidirectional

manner. For instance, recent work on adaptive myelination suggests that change in

myelination through activity-dependent adaptation of an initially hard-wired process occurs in

response to experiences and contributes to learning (Bechler et al., 2018). Second,

interpreting the biological relevance of tensor metrics from white matter tracts, such as FA

and MD, can be challenging. Whilst FA and MD are typically inversely related, where a high

FA and low MD suggest 'stronger' white matter connectivity (Vettel et al., 2017), we found

that for the majority of microstructure-curiosity correlations that only one of the two diffusion

metrics significantly correlated with curiosity, suggesting they are not redundant measures.

Dissociations between FA and MD measures could be due to a number of biological

properties such as axon diameter and density, myelination and the arrangement of fibres in

a given voxel (Beaulieu, 2002). For instance, high FA has been found to reflect high myelin

density and structured histological orientation whereas high values of MD are more likely to

reflect low myelin density and diffuse histological orientation (Seehaus et al., 2015). Future

work on the microstructural correlates of trait curiosity could apply advanced modelling

techniques, such as the "Neurite Orientation Dispersion and Density Imaging" model

(NODDI (Zhang et al., 2012)) for estimating biologically specific properties of the white

matter.

Our study involved self-report questionnaires to measure distinct curiosity traits.

While self-report questionnaires of personality have well known limitations (Vazire &

Carlson, 2010), such instruments, unlike task-based measures, are designed to maximise

consistent inter-individual differences, have high reliability and predict real world outcomes

(Grossnickle, 2016; Eisenberg et al., 2019; Enkavi et al., 2019). Nevertheless, future studies

should consider examining the link between task-evoked states of curiosity and trait

curiosity, and whether the same white matter tracts mediate state effects of curiosity in

different aspects of learning and memory.

Conclusion

The present study found inter-individual variation in the microstructure of the fornix related to

interest EC and inter-individual variation in the microstructure of the ILF related to both

interest and deprivation EC. Furthermore, posterior hippocampal fornix microstructure was

associated with specific PC. In conclusion, our findings on the relationship between curiosity

traits and anatomical connections underlying well characterized brain networks provide a

foundation for future studies to examine the relationship between curiosity traits, curiosity

states and their neuroanatomical substrates. Our findings pave the way to further

understand inter-individual differences in curiosity and which aspects of curiosity benefit

language, memory and other cognitive processes cultivating a deeper knowledge and skill

set.

Materials and Methods

Participants

Fifty-one healthy female adult undergraduate students, with a mean age of 20 years (SD ±

1, range = 19-24) participated. They provided written consent prior to participating in the

study, which was approved by the Cardiff University Research Ethics Committee, and

received a remuneration of approximately £25 for their participation.

Trait curiosity measures

Participants completed the *Epistemic Curiosity Scale (EC)* (Litman, 2008) and the

Perceptual Curiosity Scale (PC) (Collins et al., 2004), along with other measures not

relevant to the current study. The EC scale consists of five interest EC items and five

deprivation EC items with participants answering on a scale ranging from 1 (*almost never*) to

4 (*almost always*). The interest EC items are associated with behaviours that stimulate

positive affect, or involve learning something completely new (e.g. "I enjoy learning about

subjects that are unfamiliar to me"). In contrast, deprivation EC items describe behaviours

that reduce negative feelings of information deprivation and uncertainty (e.g. "I can spend

hours on a single problem because I just can't rest without knowing the answer"). The PC

scale (Collins et al., 2004) comprised of twelve items (6 diversive PC items and 6 specific

PC items) and again participants respond on a scale that ranged from 1 (*almost never*) to 4

(*almost always*). The diversive PC items describe exploratory behaviours in which one seeks

out new places and a broad range of sensory stimulation (e.g. "I like to discover new places

to go"), whereas specific PC describes exploration of novel, specific and sensorially

stimulating stimuli (e.g. "When I hear a strange sound, I usually try to find out what caused

it"). The Cronbach's alpha coefficients for the scales were all >= .70 suggesting good

internal consistency.

Imaging acquisition

Imaging data were obtained at CUBRIC, Cardiff University on a 3 Tesla MRI scanner

(Siemens Magnetom Prisma) with a 32-channel head coil. T1-weighted structural 3D images

were acquired using an MPRAGE sequence (orientation = sagittal; TR = 2250ms; TE =

3.06ms; TI = 900ms; flip angle = 9°; FOV = 256mm²; slice thickness = 1mm; voxel size =

1mm³; number of slices = 224; bandwidth = 230Hz/pixel; total acquisition time = 7 minutes

36 seconds).

Diffusion weighted images were acquired using a multi-shell sequence (orientation =

transversal/axial; TR = 9400ms; TE = 67.0ms; FOV = 256mm²; slice thickness = 2mm; voxel

size = 2mm³; number of slices = 80). Diffusion gradients were applied in (i) 30 isotropic

directions by using a diffusion-weighted factor b=1200sec/mm², (ii) in 60 isotropic directions

by using a diffusion-weighted factor b=2400sec/mm², and (iii) a volume without diffusion

gradients (b=0sec/mm²) (bandwidth = 1954Hz/pixel; total acquisition time = 15 minutes 51

seconds).

Diffusion MRI pre-processing

T1-weighted structural images were subjected to a 'brain-tissue only' mask using FSL's

Brain Extraction Tool (Smith, 2002). Using ExploreDTI (v4.8.3; Leemans et al., 2009) each

b-value image was then co-registered to the T1 structural image. Subsequently, all b-value

images were corrected for head motion and eddy currents within ExploreDTI. Tensor fitting

was conducted on the b-1200 data given the tensor model assumes hindered (Gaussian)

diffusion, and at lower b-values more of the signal is due to hindered rather than restricted

diffusion (Jones et al., 2013). To correct for voxel-wise partial volume artefacts arising from

free water contamination, the two-compartment 'Free Water Elimination' (FWE) procedure

was applied to the current b-1200 data – this improves reconstruction of white matter tracts

near the ventricles such as the fornix (Pasternak et al., 2009, 2014), yielding whole brain

voxel-wise free-water corrected FA and MD tissue maps.   Following FWE, corrected

diffusion tensor-derived structural metrics were computed. Fractional anisotropy (FA),

reflects the extent to which diffusion within biological tissue is anisotropic (constrained along

a single axis). MD (10^{-3} mm^2 s^{-1}) reflects overall degree of diffusivity (Vettel et al., 2017). The

resulting free water corrected FA and MD maps were inputs for the tractography analysis.

Tractography

As higher b-values allow for better fibre orientation estimations (Vettel et al., 2017), we

performed tractography on the b-2400 data using damped Richardson-Lucy spherical

deconvolution (dRL-SD). Spherical deconvolution provides a direct estimate of the

underlying distribution of fibre orientations in the brain and when applied to tractography

leads to accurate reconstructions of the major white matter pathway, and an improved ability

to describe complex white matter anatomy (Dell'Acqua & Tournier, 2018). The algorithm

extracted peaks in the fibre orientation density function (fODF) at the centre of each voxel,

where streamlines along the orientation of the fODF peaks were reconstructed using a step

size of 0.5mm. Streamline tracts were terminated if the direction of the pathway changed

through an angle greater than 45° or if the fODF threshold fell below 0.05.

In ExploreDTI, manual tractography was carried out using AND, NOT, and SEED

ROI gates on colour-coded FA maps to extract specific white matter tracts. AND gates

(**Figure 4** - green) were used to extract fibres that passed through the gate, NOT gates

(**Figure 4** - red) were used to exclude any fibres that passed through the gate, and finally

SEED gates (**Figure 4** - blue) were used as a starting point to extract fibres that passed

through this gate and then to include only those fibres that then passed through any added

AND gates. Manual tractography was carried out on a minimum of 15 subjects in order to

calculate a tract model to perform automated tractography on all 51 data sets (Explore DTI;

Parker et al., 2013). This procedure enables the construction of white matter tracts in space

in which streamlines belonging to particular anatomical features of interest consistently

project to distinct sub-regions, allowing the reconstruction of streamline data by observing

their projected positions (Parker et al., 2013). After running the automated tractography

software each tract was visually inspected, and any erroneous fibres were pruned using

additional NOT gates. These tract masks from the b=2400 data were then intersected with

the b=1200 free-water corrected FA and MD maps to derive free-water corrected tract-

specific measures of FA and MD values for statistical analysis.

Inferior Longitudinal Fasciculus tractography. The ILF (**Figure 4B**) was reconstructed using

a two-ROI approach in each hemisphere (Wakana et al., 2007). In the mid-saggital slice of

the brain, the coronal crosshair was placed posterior to the corpus callosum. In the coronal

plane a SEED gate was drawn around the entire cortex of interest. Next in the coronal view,

the last slice where the temporal lobe was separate from the frontal lobe was identified and

one AND gate was drawn around the temporal lobe. Any stray fibres not consistent with the

ILF pathway were removed with NOT gates. FA and MD of the right and left ILF were

summed and averaged to provide a bilateral measure for the main analyses.

Fornix tractography. The fornix (**Figure 4A**) was traced in line with the landmarks described

in Catani and Thiebaut de Schotten (2008). In the mid saggital slice of the brain, the coronal

crosshair was placed at the anterior commissure and moved approximately 6 voxels

posterior in the brain. In the coronal plane, one AND gate was drawn around the fornix

bundle where the anterior pillars enter the body of the fornix. Finally, NOT gates were drawn

around any protruding areas that were not part of the fornix.

Anterior and posterior hippocampal fornix tractography. In addition, we employed a method

adapted from prior work to reconstruct the anterior and posterior hippocampal fornix fibres

(Christiansen et al., 2017). Both anterior and posterior hippocampal fornix reconstructions

required the AND and NOT gates that were applied during whole fornix tractography. Some

NOT gates were augmented to enable better extraction of the anterior and posterior

hippocampal streamlines of the fornix. A standard landmark for the anterior-posterior

hippocampal boundary was proposed to be a small bundle of grey matter that outlines the

most anterior extent of the parahippocampal gyrus that is called uncal apex or uncus

(Poppenk et al., 2013). This landmark was identified for each hemisphere separately when

carrying out manual tractography of the anterior and posterior hippocampal fornix. In order to

perform this, the uncus was first localised at its anterior part and traced to its posterior

boundary. The first coronal slice in which the uncus was not visible anymore was used as

the landmark in order distinguish between fibres that project into anterior (head of the

hippocampus) and posterior hippocampus (body and tail of the hippocampus) (**Figure 4C**).

After the left and right hemispheric landmarks were identified, one NOT gate on each

hemisphere was drawn around the hippocampus to set boundaries for posterior

hippocampal fornix tracts, removing fibres that pass through these NOT gates (**Figure 4D**).

After the posterior hippocampal fornix was identified, the same coordinates of the anterior-

posterior hippocampal boundary landmark for each hemisphere were used to replace the

NOT gates with AND gates for the left and right anterior hippocampal fornix reconstruction

(**Figure 4E**). The posterior, left, and right anterior hippocampal fornix were saved as

separate tracts to aid subsequent automated tractography (**Figure 5**). Note that diffusion

tensor metrics of the whole fornix and those averaged across anterior and posterior

hippocampal fornix segments were highly correlated (FA, r(51) = 0.940, p < 0.001; MD, r(51)

= 0.942, p < 0.001) indicating that the anterior and posterior hippocampal fornix

reconstructions matched the whole fornix reconstructions.

Figure 4. Automated tractography reconstructions of the fornix, its anterior and

posterior hippocampal fornix fibres and the inferior longitudinal fasciculus (ILF). AND

(green), NOT (red), and SEED (blue) ROI gates for each of the tracts are displayed on the

sagittal midline plane. (**A**) Fornix tractography using AND and NOT gates. (**B**) Left ILF

tractography using SEED, AND and NOT gates. (**C**) Location of AND and NOT gates for

tractography of the anterior and posterior hippocampal fornix, respectively. (**D**) Posterior

hippocampal fornix tractography using one additional NOT gate placed between the head

and the body of the hippocampus to only include fornical fibres that connect with posterior

hippocampus (i.e., hippocampal body and tail). (**E**) Anterior hippocampal fornix tractography

using one additional AND gate placed between the head and body of the hippocampus (i.e.,

identical location as NOT gate in (D)) to include fibres that pass through this ROI gate to the

anterior hippocampus.

Figure 5. Automated tractography reconstructions of anterior and posterior

hippocampal fornix fibres on coronal slices. Tractography of the fornix fibres projecting

to the posterior hippocampus (**A**). Tractography of fornix fibres projecting to the left anterior

hippocampus (**B**). Tractography of the fornix fibres projecting to the right anterior

hippocampus (**C**).

For the questionnaire data, in the event of missing responses (2 participants failed to give a

response to one PC item), the mean value of the remaining items that were answered in the

full scale was calculated which then replaced the missing item score. For each curiosity

subscale (i.e., the two subscales of PC and EC), we calculated a total score for each

participant. Participants' data with diffusion tensor metrics +/– 3SD beyond the group mean

were considered as outliers and removed from respective analyses. This resulted in one

participant's data being removed from all analyses involving ILF MD and a different

participant's data being removed from analyses including bilaterally averaged ILF FA.

To test for associations between curiosity trait scores and microstructure of our

selected anatomical tracts, we conducted directional *Pearson's* correlations using MATLAB.

Since higher FA and lower MD is typically associated with 'stronger' white matter

connectivity (Vettel et al., 2017), we predicted a positive correlation between levels of

curiosity and FA and a negative correlation with MD.

To determine whether the *Pearson's* correlation coefficient *r* was statistically

significant, we performed non-parametric permutation tests that randomly permute the real

data between participants. Permutation tests were conducted separately for the two

microstructure metrics (i.e., FA and MD) and for the EC and PC scales. Importantly, we

corrected for multiple comparisons across the subscales within a curiosity scale (e.g.,

diversive- and specific PC). The steps were as follows: First, we performed *Pearson's*

correlations on the real data (i.e., correlations between the scores of the two curiosity

subscales and the microstructure measure (e.g., diversive PC with ILF MD and specific PC

with ILF MD)). Thereby, we obtained the empirical correlation coefficients reflecting the

relationship between the two curiosity subscales and a specific microstructure measure.

Second, within each curiosity subscale, we shuffled the curiosity scores across participants,

which resulted in pairs containing a curiosity score and a microstructure value that is

randomly assigned across participants. On these shuffled data, we then calculated surrogate

Pearson's coefficients for the two curiosity subscale scores and the microstructure metric,

and saved the maximum surrogate *Pearson's r* across the two correlations (i.e., subscale-

microstructure$_{max}$) (Groppe, Urbach & Kutas, 2011). Third, the second step was repeated

5000 times. Based on the 5000 permutations, we created a null distribution of all surrogate

subscale-microstructure$_{max}$ coefficient values and determined the alpha cut-off point

($p < 0.05$; one-sided; i.e., 4750th data point of the surrogate null distribution) in order to test

the statistical significance of the real *Pearson's* coefficients reflecting the relationship

between the two subscales and the microstructure measure. This approach allowed us to

correct for multiple comparisons across the two subscales within each curiosity scale. In

follow-up analyses for specific curiosity subscales (e.g., interest EC subscale), we also

performed follow-up permutation tests that corrected for multiple comparisons across both

hemispheres (e.g., left and right ILF MD). The 95% confidence intervals (CI) for each

correlation was derived using a bootstrapping method based on 1000 iterations. Olkin's z

test was used for the statistical comparison of dependent correlations, as implemented in the

r package 'cocor" (Diedenhofen & Musch, 2015).

Author contributions

A.V., K.S.G., A.D.L. and M.J.G. contributed to the conception and design of the experiment.

A.V. and A.C. contributed to data acquisition. All authors contributed to data analysis and

interpretation. A.V. and M.J.G. drafted the manuscript and together with C.J.H., K.S.G. and

A.D.L. revised the manuscript. A.D.L. and M.J.G. jointly supervised this work.

Conflict of interests: The authors declare no competing financial interests

Acknowledgements: We would like to thank the funders of this research as well as Ofer

Pasternak for providing the free-water correction pipeline, John Evans and Peter Hobden for

scanning support, Samuel Ridgeway and Bethany Coad for support in data collection, Sonya

Foley and Marie-Lucie Read for support in data pre-processing.

Funding

This work was supported by a departmental PhD studentship from the School of Psychology

at Cardiff University to A.V., a PhD studentship from the Cardiff University Neuroscience and

Mental Health Research Institute (NMHRI) to A.C., a Wellcome Strategic Award

(104943/Z/14/Z) to C.J.H, A.C, K.G., A.D.L., and a COFUND fellowship funded by the Welsh

Government and the European Commission and a Sir Henry Dale Fellowship

(211201/Z/18/Z) funded by Wellcome and the Royal Society to M.J.G.

References

Aggleton, J. P. (2012). "Multiple Anatomical Systems Embedded within the Primate Medial Temporal Lobe: Implications for Hippocampal Function." *Neuroscience and Biobehavioral Reviews 36 (7)*, 1579–96.

Aggleton, J.P., Wright, N.F., Rosene, D.L., & Saunders, R.C. (2015). Complementary patterns of direct amygdala and hippocampal projections to the macaque prefrontal cortex. *Cerebral Cortex 25* (11), 4351-4373.

Agosta, F., Henry, R. G., Migliaccio, R., Neuhaus, J., Miller, B. L., Dronkers, N. F., ... & Gorno-Tempini, M. L. (2010). Language networks in semantic dementia. *Brain*, *133*(1), 286-299.

Baranes, A., Oudeyer, P. Y., & Gottlieb, J. (2015). Eye movements reveal epistemic curiosity in human observers. *Vision Research, 117*, 81–90.

Beaulieu, C. (2002). The basis of anisotropic water diffusion in the nervous system - A technical review. *NMR in Biomedicine, 15(7–8)*, 435–455.

Bechler, M. E., Swire, M., & ffrench-Constant, C. (2018). Intrinsic and adaptive myelination—A sequential mechanism for smart wiring in the brain. *Developmental neurobiology*, *78*(2), 68-79.

Berlyne, D. E. (1954). A theory of human curiosity. *British Journal of Psychology, 3*, 180–191.

Berlyne, D. E. (1966). Curiosity and Exploration. *Science, 153(3731)*, 25–33.

Burgess, N., Maguire, E. A., & O'Keefe, J. (2002). The human hippocampus and spatial and episodic memory. *Neuron*, *35*(4), 625-641.

Catani, M., & Thiebaut de Schotten, M. (2008). A diffusion tensor imaging tractography atlas for virtual in vivo dissections. *Cortex, 44(8)*, 1105–1132.

Chen, L., Ralph, M. A. L., & Rogers, T. T. (2017). A unified model of human semantic knowledge and its disorders. *Nature human behaviour*, *1*(3), 0039.

Chen, Y., Chen, K., Ding, J., Zhang, Y., Yang, Q., Lv, Y., ... & Han, Z. (2017). Brain network

for the core deficits of semantic dementia: a neural network connectivity-behavior

mapping study. *Frontiers in human neuroscience*, *11*, 267.

Christiansen, K., Metzler-Baddeley, C., Parker, G. D., Muhlert, N., Jones, D. K., Aggleton, J.

P., & Vann, S. D. (2017). Topographic separation of fornical fibers associated with the

anterior and posterior hippocampus in the human brain: An MRI-diffusion study. *Brain*

and Behavior, 7(1), 1–8.

Cohen, M. X., Schoene-Bake, J. C., Elger, C. E., & Weber, B. (2009). Connectivity-based

segregation of the human striatum predicts personality characteristics. *Nature*

Neuroscience, 12(1), 32–34.

Collins, R. P., Litman, J. A., & Spielberger, C. D. (2004). The measurement of perceptual

curiosity. *Personality and Individual Differences, 36(5)*, 1127–1141.

Davachi, L. (2006). Item, context and relational episodic encoding in humans. *Current*

opinion in neurobiology, *16*(6), 693-700.

Dell'Acqua, F., & Tournier, J. D. (2018). Modelling white matter with spherical deconvolution:

How and why*?. NMR in Biomedicine*, e3945.

DeYoung C.G., (2014). Openness/Intellect: A dimension of personality reflecting cognitive

exploration. In: APA handbook of personality and social psychology: Personality

processes and individual differences (Cooper ML, Larsen RJ, eds), pp369-399.

Washington, DC: American Psychological Association.

Diedenhofen, B., & Musch, J. (2015). cocor: a comprehensive solution for the statistical

comparison of correlations. *PloS one*, *10*(3), e0121945.

doi:10.1371/journal.pone.0121945.

Eichenbaum, H., Yonelinas, A. P., & Ranganath, C. (2007). The medial temporal lobe and

recognition memory. *Annu. Rev. Neurosci.*, *30*, 123-152.

Eisenberg, I. W., Bissett, P. G., Enkavi, A. Z., Li, J., MacKinnon, D. P., Marsch, L. A., &

Poldrack, R. A. (2019). Uncovering the structure of self-regulation through data-driven

ontology discovery. *Nature communications*, *10*(1), 2319.

Enkavi, A. Z., Eisenberg, I. W., Bissett, P. G., Mazza, G. L., MacKinnon, D. P., Marsch, L.

A., & Poldrack, R. A. (2019). Large-scale analysis of test–retest reliabilities of self-

regulation measures. Proceedings of the National Academy of Sciences, 116(12), 5472-

5477.

Galli, G., Sirota, M., Gruber, M. J., Ivanof, B. E., Ganesh, J., Materassi, M., … Craik, F. I. M.

(2018). Learning facts during aging: the benefits of curiosity. *Experimental Aging*

Research, *44*, 311-328.

Gottlieb, J., & Oudeyer, P. Y. (2018). Towards a neuroscience of active sampling and

curiosity. *Nature Reviews Neuroscience*, *19*, 758-770.

Graham, K. S., Barense, M. D., & Lee, A. C. (2010). Going beyond LTM in the MTL: a

synthesis of neuropsychological and neuroimaging findings on the role of the medial

temporal lobe in memory and perception. *Neuropsychologia*, *48*(4), 831-853.

Grossnickle, E. M. (2016). Disentangling curiosity: Dimensionality, definitions, and

distinctions from interest in educational contexts. *Educational Psychology*

Review, *28*(1), 23-60.

Gruber, M. J., Gelman, B. D., & Ranganath, C. (2014). States of Curiosity Modulate

Hippocampus-Dependent Learning via the Dopaminergic Circuit. *Neuron, 84(2)*, 486–

496.

Gruber, M. J., Valji, A., & Ranganath, C. (2019). Curiosity and Learning: A Neuroscientific

Perspective. In: *The Cambridge Handbook of Motivation and Learning*. (Renninger KA &

Hidi SE, ed), pp 397-417. Cambridge University Press.

Herbet, G., Zemmoura, I., & Duffau, H. (2018). Functional anatomy of the inferior

longitudinal fasciculus: from historical reports to current hypotheses. *Frontiers in*

neuroanatomy, 12, 77.

Hodgetts, C. J., Postans, M., Shine, J. P., Jones, D. K., Lawrence, A. D., & Graham, K. S.

(2015). Dissociable roles of the inferior longitudinal fasciculus and fornix in face and

place perception. *Elife, 4*, e07902.

Hodgetts, C. J., Postans, M., Warne, N., Varnava, A., Lawrence, A. D., & Graham, K. S.

(2017). Distinct contributions of the fornix and inferior longitudinal fasciculus to episodic

and semantic autobiographical memory. *Cortex, 94*, 1–14.

Hoffman, P., & Lambon Ralph, M. A. (2018). From percept to concept in the ventral temporal

lobes: Graded hemispheric specialisation based on stimulus and task. *Cortex*, *101*, 107-

118.

Horowitz-Kraus, T., Wang, Y., Plante, E., & Holland, S. K. (2014). Involvement of the right

hemisphere in reading comprehension: a DTI study. *Brain research*, *1582*, 34-44.

Jones, D. K., Knösche, T. R., & Turner, R. (2013). White matter integrity, fiber count, and

other fallacies: the do's and don'ts of diffusion MRI. *Neuroimage, 73*, 239-254.

Jouen, A. L., Ellmore, T. M., Madden, C. J., Pallier, C., Dominey, P. F., & Ventre-Dominey,

J. (2015). Beyond the word and image: characteristics of a common meaning system for

language and vision revealed by functional and structural imaging. *NeuroImage*, *106*,

72-85.

Kang, M. J., Hsu, M., Krajbich, I. M., Loewenstein, G., McClure, S. M., Wang, J. T., &

Camerer, C. F. (2009). The wick in the candle of leraning. *Psychological Science, 20(8)*,

963–974.

Kashdan, T. B., & Yuen, M. (2007). Whether highly curious students thrive academically

depends on perceptions about the school learning environment: A study of Hong Kong

adolescents. *Motivation and Emotion*, *31*(4), 260–270.

Kumaran, D., & Maguire, E. A. (2009). Novelty signals: a window into hippocampal

information processing. *Trends in cognitive sciences, 13(2)*, 47-54.

Lambon Ralph, M., Jefferies, E., Patterson, K., & Rogers, T. T. (2017). The neural and

computational bases of semantic cognition. *Nature Reviews Neuroscience*, *18*(1), 42-55

Lauriola, M., Litman, J. A., Mussel, P., De Santis, R., Crowson, H. M., & Hoffman, R. R.

(2015). Epistemic curiosity and self-regulation. Personality and Individual Differences,

83(1), 202–207.

Leemans A, Jeurissen B, Sijbers J, Jones DK (2009). ExploreDTI: a graphical toolbox for

processing, analyzing, and visualizing diffusion MR data. Proceedings of the 17th

Annual Meeting of International Society for Magnetic Resonance in Medicine 17:3536.

Litman, J. A. (2005). Curiosity and the pleasures of learning: Wanting and liking new

information. *Cognition and Emotion, 19*, 793–814

Litman, J. A. (2008). Interest and deprivation factors of epistemic curiosity. *Personality and*

Individual Differences, 44(7), 1585–1595.

Litman, J. A., & Jimerson, T. L. (2004). The measurement of curiosity as a feeling of

deprivation. *Journal of Personality Assessment, 82(2)*, 147–157.

Litman, J., & Spielberger, C. (2003). Measuring epistemic curiosity and its diversive and

specific components. *Journal of Personality Assessment, 80(1)*, 75–86.

Liu, Z. X., Shen, K., Olsen, R. K., & Ryan, J. D. (2017). Visual sampling predicts

hippocampal activity. *Journal of Neuroscience*, *37*(3), 599-609.

Loewenstein, G. (1994). The psychology of curiosity: A review and reinterpretation.

Psychological Bulletin, 116(1), 75–98.

Marvin, C. B., & Shohamy, D. (2016). Curiosity and reward: Valence predicts choice and

information prediction errors enhance learning. *Journal of Experimental Psychology:*

General, 145(3), 266–272.

McGillivray, S., Murayama, K., & Castel, A. D. (2015). Thirst for knowledge: The effects of

curiosity and interest on memory in younger and older adults. *Psychology and Aging,*

30(4), 835–841.

Murray, E. A., Wise, S. P., & Graham, K. S. (2017). The evolution of memory systems:

ancestors, anatomy, and adaptations. Oxford University Press.

Murray, E. A., Wise, S. P., & Graham, K. S. (2018). Representational specializations of the
hippocampus in phylogenetic perspective. *Neuroscience letters*, *680*, 4-12.

Mussel, P. (2013). Introducing the construct curiosity for predicting job performance. *Journal*
of Organizational Behavior, *34*, 453–472.

O'Keefe, J. and Nadel, L. (1978) The hippocampus as a cognitive map, Oxford: Clarendon
Press.

Otmakhova, N., Duzel, E., Deutch, A. Y., & Lisman, J. (2013). The hippocampal-VTA loop:
the role of novelty and motivation in controlling the entry of information into long-term
memory. In *Intrinsically motivated learning in natural and artificial systems*(pp. 235-254).
Springer, Berlin, Heidelberg.

Panesar, S. S., Yeh, F. C., Jacquesson, T., Hula, W., & Fernandez-Miranda, J. C. (2018). A
quantitative tractography study into the connectivity, segmentation and laterality of the
human inferior longitudinal fasciculus. *Frontiers in neuroanatomy*, *12*:47

Parker, G. D., Marshall, D., Rosin, P. L., Drage, N., Richmond, S., & Jones, D. K. (2013).
Fast and fully automated clustering of whole brain tractography results using shape-
space analysis. *Proceedings of the International Society for Magnetic Resonance in*
Medicine. Salt Lake City, USA, 778.

Pasternak, O., Maier-Hein, K., Baumgartner, C., Shenton, M. E., Rathi, Y., & Westin, C. F.
(2014). The estimation of free-water corrected diffusion tensors. In: Visualization and
Processing of Tensors and Higher Order Descriptors for Multi-Valued Data (Westin CF.,
Vilanova A., Burgeth B., eds), pp. 249-270. Springer, Berlin, Heidelberg.

Pasternak, O., Sochen, N., Gur, Y., Intrator, N., & Assaf, Y. (2009). Free water elimination
and mapping from diffusion MRI. *Magnetic Resonance in Medicine 62(3)*, 717-730.

Patterson, K., Nestor, P. J., & Rogers, T. T. (2007). Where do you know what you know?
The representation of semantic knowledge in the human brain. *Nature Reviews*
Neuroscience, *8*(12), 976-987.

Poppenk, J., Evensmoen, H. R., Moscovitch, M., & Nadel, L. (2013). Long-axis
specialization of the human hippocampus. *Trends in cognitive sciences, 17(5)*, 230-240.

Privado, J., Román, F. J., Saénz-Urturi, C., Burgaleta, M., & Colom, R. (2017). Gray and
white matter correlates of the Big Five personality traits. *Neuroscience, 349*, 174–184.

Ranganath, C, & Ritchey, M. (2012). Two Cortical Systems for Memory-Guided Behaviour.
Nature Reviews. *Neuroscience 13 (10)*, 713–26.

Rice, G. E., Hoffman, P., Ralph, L., & Matthew, A. (2015). Graded specialization within and
between the anterior temporal lobes. *Annals of the New York Academy of*
Sciences, *1359*(1), 84-97.

Ripollés, P., Biel, D., Peñaloza, C., Kaufmann, J., Marco-Pallarés, J., Noesselt, T., &
Rodríguez-Fornells, A. (2017). Strength of Temporal White Matter Pathways Predicts
Semantic Learning. *The Journal of Neuroscience, 37(46)*, 1720–17.

Risko, E. F., Anderson, N. C., Lanthier, S., & Kingstone, A. (2012). Curious eyes: Individual
differences in personality predict eye movement behavior in scene-
viewing. *Cognition, 122(1)*, 86-90.

Robin, J., & Moscovitch, M. (2017). Details, gist and schema: hippocampal–neocortical
interactions underlying recent and remote episodic and spatial memory. *Current Opinion*
in Behavioral Sciences, 17, 114-123.

Saunders, R. C., & Aggleton, J. P. (2007). Origin and topography of fibers contributing to the
fornix in macaque monkeys. *Hippocampus, 17(5)*, 396-411.

Seehaus, A., Roebroeck, A., Bastiani, M., Fonseca, L., Bratzke, H., Lori, N., ... & Galuske,
R. (2015). Histological validation of high-resolution DTI in human post mortem
tissue. *Frontiers in Neuroanatomy, 9*, 98.

Sheldon, S., Fenerci, C., & Gurguryan, L. (2019). A Neurocognitive Perspective on the
Forms and Functions of Autobiographical Memory Retrieval. *Frontiers in Systems*
Neuroscience, 13.

Silvia, P. J., & Kashdan, T. B. (2009). Interesting things and curious people: Exploration and

engagement as transient states and enduring strengths. *Social and Personality*

Psychology Compass, 3(5), 785-797.

Smith, S. M. (2002). Fast robust automated brain extraction. *Human brain mapping, 17(3)*,

143-155.

Stare, C. J., Gruber, M. J., Nadel, L., Ranganath, C., & Gómez, R. L. (2018). Curiosity-

driven memory enhancement persists over time but does not benefit from post-learning

sleep. *Cognitive Neuroscience, 9*, 100-115.

Strange, B. A., Witter, M. P., Lein, E. S., & Moser, E. I. (2014). Functional organization of the

hippocampal longitudinal axis. *Nature Reviews Neuroscience, 15(10)*, 655.

Vazire, S., & Carlson, E. N. (2010). Self-knowledge of personality: Do people know

themselves?. *Social and personality psychology compass*, *4*(8), 605-620.

Vazire, S., & Carlson, E. N. (2010). Self-knowledge of personality: Do people know

themselves?. *Social and personality psychology compass*, *4*(8), 605-620.

Vettel, J. M., Cooper, N., Garcia, J. O., Yeh, F.-C., & Verstynen, T. D. (2017). White Matter

Tractography and Diffusion-Weighted Imaging. eLS, 1–9.

Voss, J. L., Bridge, D. J., Cohen, N. J., & Walker, J. A. (2017). A closer look at the

hippocampus and memory. *Trends in cognitive sciences, 21(8)*, 577-588.

Wakana, S., Caprihan, A., Panzenboeck, M. M., Fallon, J. H., Perry, M., Gollub, R. L., ... &

Blitz, A. (2007). Reproducibility of quantitative tractography methods applied to cerebral

white matter. *Neuroimage, 36(3)*, 630-644.

Woo, S. E., Chernyshenko, O. S., Longley, A., Zhang, Z.-X., Chiu, C.-Y., & Stark, S. E.

(2014). Openness to experience: Its lower level structure, measurement, and cross-

cultural equivalence. *Journal of Personality Assessment, 96(1)*, 29–45.

Yeatman, J. D., Dougherty, R. F., Myall, N. J., Wandell, B. A., & Feldman, H. M. (2012).

Tract profiles of white matter properties: automating fiber-tract quantification. *PloS one*,

7(11), e49790.

Zhang, H., Schneider, T., Wheeler-Kingshott, C. A., & Alexander, D. C. (2012). NODDI:

practical in vivo neurite orientation dispersion and density imaging of the human

brain. *Neuroimage*, *61*(4), 1000-1016.