Seven new Neocallimastigomycota genera from fecal samples of wild, zoo-housed, and domesticated herbivores: Description of *Ghazallomyces constrictus* gen. nov., sp. nov., *Aklioshbomyces papillarum* gen. nov., sp. nov., *Agriosomyces longus* gen. nov., sp. nov., *Capellomyces foraminis* gen. nov., sp. nov. and *Capellomyces elongatus* sp. nov., *Joblinomyces apicalis* gen. nov., sp. nov., *Khoyollomyces ramosus* gen. nov., sp. nov., and *Tahromyces munnarensis* gen. nov., sp. nov.

Radwa A. Hanafy[1], Vikram B. Lanjekar[2], Prashant K. Dhakephalkar [2], Tony M. Callaghan[3,] Sumit S. Dagar[2], Gareth W. Griffith[3], Mostafa S. Elshahed[1], and Noha H. Youssef[1*]

[1]Department of Microbiology and Molecular Genetics, Oklahoma State University, Stillwater, OK. [2]Bioenergy group, Agharkar Research Institute, Pune, India. [3]Institute of Biological, Environmental, and Rural Sciences (IBERS) Aberystwyth University, Aberystwyth, Wales, UK.

***Correspondence:** Noha H. Youssef: Noha@Okstate.edu.

ABSTRACT

We isolated and characterized sixty-five anaerobic gut fungi (AGF, Neocallimastigomycota)

strains from fecal samples of five wild (W), one zoo-housed (Z), and three domesticated (D)

herbivores in the US states of Texas (TX) and Oklahoma (OK), Wales (WA), and the Indian

states of Kerala (KE) and Haryana (HA). Phylogenetic assessment based on D1-D2 region of the

large rRNA subunit (LSU) identified seven distinct lineages, with strains recovered from Axis

Deer (W-TX) clustering within the *Orpinomyces-Neocallimastix-Pecoramyces-Feramyces* clade;

Boer Goat-domesticated Goat strains (W-TX, D-KE) clustering within the *Oontomyces-*

Anaeromyces-Liebetanzomyces clade; and domesticated Goat and Sheep strains (D-HA) as well

as Nilgiri Tahr strains (W-KE) forming two distinct clades associated with genus

Buwchfawromyces. The remaining three lineages, represented by strains recovered from

Mouflon-Boer Goat (W-TX), White Tailed Deer (W-OK), and Zebra-Horse (Z-OK, and D-WA),

displayed no specific suprageneric affiliation. All strains displayed monocentric thalli and

produced mono/uniflagellate zoospores with the exception of Axis Deer strains, which produced

polyflagellate zoospores. Isolates displayed multiple interesting microscopic features including

sporangia with tightly constricted necks and fine septa at the base (Axis Deer), papillated and

pseudo-intercalary sporangia (White-Tailed Deer), swollen sporangiophores and zoospores with

long flagella (Mouflon-Boer Goat), zoospore release through an apical pore followed by either

sporangial wall collapse (Axis Deer and Boer Goat-domesticated Goat) or sporangial wall

remaining intact after discharge (Zebra-Horse), multi-sporangiated thalli with branched

sporangiophores (Zebra-Horse), and short sporangiophores with subsporangial swellings (Nilgiri

Tahr). Internal transcribed spacer-1 region (ITS-1) sequence analysis indicated that Zebra-Horse

strains are representatives of the AL1 lineage, frequently encountered in culture-independent

surveys of the alimentary tract and fecal samples from hindgut fermenters. The other six

lineages, five of which were isolated from wild herbivores, have not been previously

encountered in such surveys. Our results significantly expand the genus level diversity within the

Neocallimastigomycota, and strongly suggest that wild herbivores represent a yet-untapped

reservoir of AGF diversity. We propose the creation of seven novel genera and eight novel

Neocallimastigomycota species to accommodate these strains, for which we propose the names

Agriosomyces longus (Mouflon and wild Boer Goat), *Aklioshbomyces papillarum* (White tailed

Deer), *Capellomyces foraminis* (wild Boar Goat) and *C. elongatus* (domesticated Goat),

Ghazallomyces constrictus (Axis Deer), *Joblinomyces apicalis* (domesticated Goat and Sheep),

Khoyollomyces ramosus (Zebra-Horse), and *Tahromyces munnarensis* (Nilgiri Tahr). The type

species are strains Axs-31, WT-2, MS-4, BGB-11, GFKJa1916, GFH683, ZS-33, and

TDFKJa193, respectively.

KEY WORDS: Anaerobic gut fungi, Neocallimastigomycota, Herbivores, 8 new taxa.

INTRODUCTION

Members of the anaerobic gut fungi (AGF, Phylum Neocallimastigomycota) colonize the

alimentary tract of mammalian and reptilian herbivores (Gruninger and others 2014; Ljungdahl

2008). Interest in the taxonomy, ecology, cell biology, and genomics of the

Neocallimastigomycota has been driven by their unique habitat, physiological preferences, and

evolutionary history, as well as by their possession of superior plant polymers degradation

capacities (Youssef and others 2013). Such characteristics render them a promising platform for

biofuel and biogas production from plant biomass (Ranganathan and others 2017; Young and

others 2018).

Currently, eleven different AGF genera have been described (Ariyawansa 2015; Barr and

others 1989 ; Breton and others 1990; Callaghan and others 2015; Dagar and others 2015; Gold

and others 1988; Hanafy and others 2017; Hanafy and others 2018; Heath and others 1983; Joshi

and others 2018; Li 2016; Ozkose and others 2001). However, it is reasonable to assume that

multiple novel, yet-uncultured AGF lineages remain to be isolated and characterized. The

inherent difficulty in isolating and maintaining these strictly anaerobic and senescence-prone

organisms severely hampers isolation and characterization efforts, and limits the number of

research groups dedicated to uncovering AGF diversity. Further, it is entirely plausible that

multiple AGF taxa are extremely fastidious, with complex nutritional requirements that are not

satisfied in current isolation protocols. Indeed, culture-independent diversity surveys utilizing the

internal transcribed spacer 1 (ITS-1) as a phylogenetic marker demonstrated that multiple novel

AGF lineages remain to be isolated and characterized (Kittelmann and others 2012;

Liggenstoffer and others 2010; Paul and others 2018; Mura and others 2019).

On the other hand, culturing efforts have occasionally recovered novel AGF strains that

bear no clear similarities to clades identified in culture-independent studies (Callaghan and

others 2015; Joshi and others 2018). This surprising observation could be attributed to

mismatches in the isolates' ITS-1 region to commonly utilized ITS-1 AGF primers (Callaghan

and others 2015), extremely narrow host range of some AGF taxa (Callaghan and others 2015),

or existence in extremely low relative abundance *in-situ*.

Moreover, it is important to note that while both culture-based and culture-independent

surveys have reported the presence of AGF communities in a relatively wide range of animal

hosts, such studies by no means represent an exhaustive catalogue of global AGF diversity in

nature. For example, due to logistical considerations, the great majority of studies have utilized

samples from domesticated herbivores, with efforts to isolate AGF strains from wild herbivores

being extremely rare (Nagpal and others 2011; Paul and others 2010; Tuckwell and others 2005;

Hanafy and others 2018).

In an effort to broaden the current Neocallimastigomycota global culture collection, we

conducted a multi-year isolation effort targeting novel AGF taxa in fecal samples from a wide

range of wild, zoo-housed, and domesticated herbivorous mammals. Here, we report on the

isolation and characterization of seven novel AGF genera, including the first cultured

representative of the hitherto uncultured AGF lineage AL1. The results expand the known AGF

genus-level diversity by >50% (from 11–18), and strongly suggest that wild undomesticated

herbivores represent a yet-untapped reservoir of novel AGF taxa.

MATERIALS AND METHODS

Samples— Fecal samples were obtained from Axis Deer (*Axis axis*), White Tailed Deer

(*Odocoileus virginianus*), Mouflon Sheep (*Ovis orientalis*), and Boer Goat (*Capra aegagrus*) in

two separate hunting expeditions in Sutton and Val Verde counties (TX), and Payne county (OK)

in October 2017 and April 2018 (TABLE 1). The hunting parties had all appropriate licenses,

and the animals were shot either on private land with the owner's consent or on public land

during the hunting season. Samples were also obtained from a Grevy's Zebra (*Equus grevyi*)

housed in the Oklahoma City Zoo in May 2018, with the sampling protocol approved by the

Oklahoma City Zoo and Botanical Garden's Scientific Review Committee. All fecal samples

were placed on ice on site, transferred to the laboratory within 24h of collection, where they were

immediately used as an inoculum for subsequent enrichment and isolation procedures.

In India, dried fecal samples were obtained from Nilgiri Tahr (*Nilgiritragus hylocrius*), and

domesticated but forest grazing Goat (*Capra aegagrus hircus*) in Munnar in the State of Kerala.

Fresh fecal samples were also obtained from domesticated Goats and Sheep (*Ovis aries*) in

Sonipat in the State of Haryana. Fresh fecal samples were transferred to the laboratory within

24h of collection, while dried fecal samples were transferred within 72 hours of collection. In

Wales, samples were obtained from two domesticated Horses in Llanbadarn Fawr, Ceredigion

County, and promptly transferred to the laboratory for processing.

Isolation procedures— In the USA, isolation of anaerobic fungal strains was conducted as

previously described (Hanafy and others 2017). Feces were suspended in rumen-fluid (RF)

media (Calkins and others 2016; Hanafy and others 2018) with either cellobiose, or 0.5%

cellobiose and switchgrass (0.5%) used as a substrate. Antibiotics (50 µg/mL kanamycin, 50

µg/mL penicillin, 20 µg/mL streptomycin, and 50 µg/mL chloramphenicol, respectively) were

added to inhibit growth of bacteria. Samples were serially diluted and incubated at 39 C for 24–

48 h. Dilutions showing visible signs of growth (clumping or floating plant materials and

production of gas bubbles) were then used for the preparation of roll tubes (Hungate 1969) on

RF-cellobiose agar media. Single colonies were picked into liquid RF-cellobiose media, and at

least three rounds of tube rolling and colony picking were conducted to ensure purity of the

obtained colonies. Strains were maintained by bi-weekly subculturing into RF-cellobiose media.

In India, the fecal samples were homogenised in anaerobic diluent (McSweeney and others 2005)

using BagMixer (Interscience, France). One ml of homogenate was inoculated into 9 ml fungal

culture medium (Joshi and others 2018) containing neutral detergent fibre (NDF) as the sole

carbon source, and serially diluted up to 10^{-3} dilution. The antibiotics benzylpenicillin and

streptomycin sulfate (final concentration 2 mg/ml) were used to inhibit the bacterial growth.

Following incubation at 39±1 C for 5–10 d, the tubes showing visible colonization of NDF were

used to isolate pure cultures of anaerobic fungi using serum roll bottle method as described

previously (Joshi and others 2018). The colonies differing in morphology were picked, grown in

liquid culture medium, and re-roll bottled until single culture was established.

Long-term storage was conducted by surface inoculation of RF-cellobiose agar media as

described previously (Calkins and others 2016), or by cryopreservation at -80 °C using 0.64 M

ethylene glycol as the cryoprotectant (Callaghan and others 2015). Cultures are available at

Oklahoma State University, Department of Microbiology and Molecular Genetics culture

collection, and at MACS Collection of Microorganisms (MCM), Agharkar Research Institute,

Pune, India. In Wales, isolation procedures were conducted as previously described in

(Callaghan and others 2015).

Morphological characterization— The colony morphology of 3d old cultures on roll bottles was

measured using a stereomicroscope (Leica M205 FA) equipped with a digital camera (Leica

DFC450 C), or directly from the roll tube. Samples for light and scanning electron microscopy

were obtained from liquid cultures at various stages of growth. Lactophenol cotton blue stain

was used for visualization of fungal structures using an Olympus BX51 microscope (Olympus,

Center Valley, Pennsylvania) equipped with a DP71 digital camera (Olympus), or a phase

contrast microscope equipped with a Canon DS126191 digital camera. For examination of nuclei

localization, samples were stained with 4′, 6 diamidino-2-phenylindole (DAPI, 10 μg/ml), as

previously described (Callaghan and others 2015; Hanafy and others 2018; Joshi and others

2018) and examined using a fluorescence Olympus BX51 microscope (Olympus, Center Valley,

Pennsylvania) equipped with a Brightline DAPI high-contrast filter set for DAPI fluorescence

and a DP71 digital camera (Olympus), or an Olympus BX53 differential interference contrast

(DIC) microscope equipped with a DP73 digital camera (Olympus). Scanning electron

microscopy was conducted with a FEI quanta scanning electron microscope (Hillsboro, Oregon,

USA), or a Carl Zeiss EVO MA15 (Hanafy and others 2017; Joshi and others 2018).

Phylogenetic analysis— Biomass was harvested and crushed in liquid N_2. DNA was extracted

from the ground fungal biomass using DNeasy PowerPlant Pro Kit (Qiagen Corp., Germantown,

MD, USA) according to the manufacturer's instructions, or using the CTAB DNA extraction

protocol (Joshi and others 2018). To assess phylogenetic relationships, the ITS-1 region, and the

D1/D2 region of the 28S rRNA (hereafter LSU) were amplified using the MN100 (5′-

TCCTACCCTTTGTGAATTTG-3′) / MNGM2 (5′-CTGCGTTCTTCATCGTTGCG-3′) pair,

and the NL1 (5′-GCATATCAATAAGCGGAGGAAAAG-3′) / NL4 (5′-

GGTCCGTGTTTCAAGACG G-3′) pair, respectively as previously described (Hanafy and

others 2017; Joshi and others 2018). The resulting PCR amplicon for the ITS-1 region was

cloned into a TOPO-TA cloning vector according to the manufacturer's instructions (Life

Technologies®, Carlsbad, CA) and several clones were Sanger sequenced, while the purified

LSU PCR amplicons were directly sequenced using the services of the Oklahoma State

University DNA core facility or a commercial provider (1st BASE, Singapore). The obtained

sequences were aligned to anaerobic fungal reference ITS-1 and LSU sequences downloaded

from GenBank using MAFFT aligner (Nakamura and others 2018) and the alignments were used

to construct maximum likelihood phylogenetic trees in MEGA7 (Kumar and others 2016), using

Chytriomyces sp. JEL176 as the outgroup. Bootstrap values were calculated on the basis of 100

replicates.

Ecological distribution— We queried GenBank and ITS-1 datasets (Kittelmann and others 2012;

Liggenstoffer and others 2010; Paul and others 2018) using reference ITS-1 sequences from

strains recovered in this study. The phylogenetic position of all closely related sequences (> 87%

sequence similarity) was evaluated by insertion into maximum likelihood trees. Taxonomy of

uncultured taxa followed the schemes outlined in prior publications (Kittelmann and others 2012;

Liggenstoffer and others 2010; Paul and others 2018)**.**

Accession numbers— Sequences generated in this study have been deposited in GenBank under

accession numbers MK881965-MK882046, MK775304, MK775310-MK775313, MK775315,

MK775321-MK775324, MK755326-MK755327, MK755330, MK755333. Alignments and

phylogenetic trees are available through TreeBase under study accession URL

http://purl.org/phylo/treebase/phylows/study/TB2:S24394

RESULTS

Isolation summary— Sixty-five different strains were obtained and characterized in this study

(TABLE 1). These isolates were obtained from fecal samples of five wild undomesticated

herbivores: Axis Deer (W-TX), White tailed Deer (W-OK), Mouflon (W-TX), Boer Goat (W-

TX), and Nilgiri Tahr (Munnar, Kerala, India), one zoo-housed Grevy's Zebra (Z-OK), two

domesticated Horses (D-WA), two domesticated Goats (D-HA and D-KE) and a domesticated

Sheep (D-HA) (TABLE 1). Morphological, microscopic and phylogenetic analysis described

below grouped these isolates into seven distinct clades (Labeled clades 1-7 in TABLE 1

according to alphabetical order of suggested genus names). Isolates representing three clades

were obtained from one host animal only: White Tailed Deer strains (Clade 2), Axis Deer strains

(Clade 4), and Nilgiri Tahr strains (clade 7); while isolates representing four clades were

identified in more than sample: Mouflon-Boer Goat strains (clade 1), Boer Goat-domesticated

Goat strains (clade 3), domesticated Sheep-Goat strains (clade 5), and Zebra-Horse strains (clade

6). No specific morphological or microscopic decipherable differences were identified between

different strains belonging to most of these clades, and one strain from each group was chosen

for detailed analysis (TABLE 1). The only two exceptions were: 1. Strains belonging to clade 3

(Boer Goat-domesticated Goat); where strains from wild Boer Goat (W-TX) displayed distinct

microscopic and phylogenetic differences from those obtained from the domesticated Goat (D-

KE) to warrant the detailed characterization and eventual description of two different strains

(TABLE 1 and detailed descriptions below), and 2. Strains belonging to clade 6 (Zebra-Horse)

where few microscopic, but negligible phylogenetic differences were observed between the 16

Zebra strains (Z-OK) and the 5 Horse strains (D-WA) identified. These differences are

highlighted below, but we do not believe they warrant the description of a new species, given the

negligible sequence divergences between these strains. Below, we provide a detailed

characterization of these seven novel groups.

Colony morphology and macroscopic growth characteristics—Clade 1 (Mouflon-Boer Goat)

strain MS-2 produced small brown circular colonies (0.2–1 mm diam.) on agar, and a thin

biofilm-like growth in liquid media (FIG. 1a). Clade 2 (White tailed Deer) strain WT-2 produced

beige circular colonies (0.5–2.5 mm diam.) with a brown central core of dense sporangial

structures and an outer ring of light gray hyphal growth. In liquid media, it produced heavy

growth of thick biofilms that firmly attached to the tube's glass surface (FIG. 1b). Clade 3 (Boar

Goat-domesticated Goat) strain BGB-11 produced small circular brown colonies (0.1–0.5 mm

diam.), with dark center of sporangia structures and a thin fungal biofilm in liquid media (FIG.

1c), while Boar Goat-domesticated Goat strain GFKJa1916 produced compact cottony off-white

colonies of 2–3 mm size, with a fluffy center of thick sporangial structures and surrounded by

radiating rhizoids (FIG. 1d). In liquid media, strain GFKJa1916 produced numerous fungal thalli

attached to the glass bottles on initial days of growth, which later developed into thin mat-like

structures. Clade 4 (Axis Deer) strain Axs-31 produced small circular white colonies (1–4 mm

diameter), with a brown central core of dense sporangial structures on agar and a thick fungal

biofilm-like growth in liquid media (FIG. 1e). Clade 5 (Domesticated Goat-Sheep) strain

GFH683 produced 1–2 mm sized colonies, having a dense dark central core of abundant

sporangial growth, surrounded by long and thin radiating rhizoids (FIG. 1g). In liquid media,

strain GFH683 produced numerous fungal thalli attached to the glass bottles on initial days of

growth, which later developed into thin mat-like structures. Clade 6 (Zebra-Horse) strains Zebra

strain ZS-33 produced small yellow to yellowish brown irregularly shaped colonies (FIG. 1f). In

liquid media, the fungal thalli were loose and exhibited a sand-like appearance resembling liquid

growth patterns generally observed with isolates belonging to the bulbous genera *Caecomyces*

and *Cyllamyces* (personal observation) (FIG. 1f). Finally, clade 7 (Nilgiri Tahr) strain

TDFKJa193 were smaller, approximately 1 mm in size, white in color with a compact and fluffy

center, surrounded by dotted circles of fungal thalli. In liquid media, the strain produced

numerous fungal thalli attached to the glass bottles on initial days of growth, which later

developed into thin mat-like structures (FIG. 1h).

Microscopic features—

Clade 1 (Mouflon-Boer Goat) strains— Strain MS2 produced small globose zoospores, with an

average diameter of 4 ± 1.1 μm (average ± standard deviation values from 29 zoospores, range

2.7–7.5 μm). Zoospores were mainly mono-flagellate with a flagellum length of 22 ± 3.8 μm

(average ± standard deviation values for 29 zoospores, range 16.6–30 μm), approximately 5–6

times longer than the zoospore body (FIG. 2a). Biflagellate zoospores (FIG. 2b) were rarely

encountered. Zoospores germinated into monocentric thalli with filamentous anuclear rhizoidal

systems (FIG. 2c-d). Both endogenous and exogenous sporangia were observed, which were

very homogenous and displayed no pleomorphism. Endogenous sporangia were globose, with a

diameter range of 15–65 μm (FIG. 2e-f). The rhizoid was swollen below the sporangial neck,

which was tightly constricted (FIG. 2e-f)**.** Exogenous sporangia were also consistently globose

and developed at the end of swollen sporangiophores (30–80 μm L X 5–10 μm W) (FIG. 2g-h).

The sporangial neck was constricted with a narrow neck port. Zoospores were released through

dissolution and rupturing of the sporangial wall (FIG. 2i).

Clade 2 (White-tailed Deer) strains— Strain WT-2 produced globose zoospores, with an average

diameter of 7.4 ± 2.4 μm (average ± standard deviation values for 35 zoospores, range 4.5–13

μm). Zoospores were mostly monoflagellate, with an average flagellum length of 22.8 ± 6.3 μm

(average ± standard deviation values for 35 zoospores, range 12–35 μm) (FIG. 3a). Zoospores

with two (FIG. 3b) to three (FIG. 3c) flagella were less frequently observed. Fungal thalli were

consistently monocentric with filamentous anucleate rhizoids (FIG. 3d). Germination of

zoospores produced two types of monocentric thalli, endogenous and exogenous. Endogenous

sporangia with single (FIG. 3e) and two adjacent rhizoidal systems (FIG. 3f) were observed.

Occasionally, pseudo- intercalary endogenous sporangia (sporangia present in the middle of two

main rhizoids) were encountered (FIG. 3g), Similar to what have previously been observed with

the genera *Oontomyces* (Dagar and other, 2015) and *Feramyces* (Hanafy and others, 2017).

Exogenous sporangia developed at the end of unbranched sporangiophores of varying length

from a few microns to 230 μm (FIG. 3h-j). No morphological differences were noticed between

endogenous and exogenous sporangia and their shapes ranged from ovoid (FIG. 3e-f), globose

(FIG. 3g-h), obpyriform (FIG. 3j-k and 3n-o), and ellipsoidal (FIG. 3i and 3l). Many, but not all,

sporangia were papillated with one (FIG. 3m-p) or two (FIG. 3q) papillae. These papillated

sporangia are similar to those previously observed in *Piromyces mae* (Li and others 1990). It is

believed that these papillae disintegrate to facilitate zoospore release. However, we were unable

to observe zoospore discharge through papillae in strain WT-2.

Clade 3 (Boer Goat-domesticated Goat) strains— Boer Goat Strain BGB-11 produced globose

zoospores, with an average diameter of 5.5 ± 0.97 μm (average ± standard deviation values for

40 zoospores, range 4–7 μm). The majority of zoospores were mono-flagellate with a flagellum

length of 19.6 ± 3.2 μm (average ± standard deviation values for 40 zoospores, range 15–25 μm),

(FIG. 4a). Occasionally, biflagellate zoospores were observed (FIG. 4b). Zoospores encystment

followed flagellar shedding (FIG. 4c). Zoospore cyst germinated producing germ tube (FIG. 4d)

that subsequently branched (FIG. 4e) into monocentric thalli with filamentous anucleate

rhizoidal systems (FIG. 4f-g).

The expansion of the zoospore cysts resulted in the formation of endogenous sporangia

that were ellipsoidal (FIG. 4h) and ovoid (FIG. 4i). In addition to endogenous sporangia,

exogenous sporangia were also observed at the end of unbranched sporangiophores ranging in

length between 20–150 μm (FIG. 4j-p). Some of the sporangiophores ended with sub-sporangial

swellings (FIG. 4l-m). Exogenous sporangia varied in shape from ovoid (FIG. 4k-l), ellipsoidal

with a single constriction (FIG. 4n), and globose (FIG. 4o-p). Zoospores were liberated through a

wide apical pore at the top of the sporangia followed by sporangial wall collapse (FIG. 4m, q-r).

Domesticated Goat strain GFKJa1916 on the other hand produced globose zoospores

(FIG. 5a), with an average diameter of 4–5 μm. The majority of zoospores were mono-flagellate

with a flagellum length of 15–20 μm. Bi- and tri-flagellate zoospores were also observed. Strain

GFKJa1916 zoospores germinated either endogenously or exogenously into a single monocentric

thallus, which was also confirmed by the presence of nuclei only in sporangia and their absence

in rhizoids (FIG. 5b-c). Endogenous sporangia varied in shape between cylindrical, elongate,

globose, sub-globose, ellipsoid & obovoid with sizes ranging between 8–60 μm wide & 10–140

μm long (FIG. 5d-g). Unlike Boer Goat Strain BGB-11, exogenous sporangia in the

domesticated Goat strain GFKJa1916 developed at the end of long thick sporangiophores (up to

300 μm in some cases) (FIG. 5h-l), and multisporangiate thalli were commonly observed with

two sporangia of either the same (FIG. 5j) or different (FIG. 5K-l) shape, similar to *Piromyces*

rhizinflatus (Ho and Barr 1995) and *Neocallimastix frontalis* (Barr and others 1995).

Clade 4 (Axis Deer) strains— Strain Axs-31 produced globose zoospores, with an average

diameter of 8.1 ± 1.3 μm (average ± standard deviation values for 35 zoospores, range 6–10.5

μm). All zoospores were polyflagellate, with (7–14) flagella and an average flagellum length of

23.5 ± 4.9 μm (average ± standard deviation values for 35 zoospores, range 16–31 μm) (FIG.

6a). Zoospores germinated into monocentric thalli with highly branched anucleate rhizoidal

systems (FIG. 6b-c).

Strain Axs-31 exhibited both endogenous and exogenous monocentric thallus

development. In endogenous thalli, zoospore cysts enlarged into new sporangia of different

shapes, including globose (FIG. 6d), tubular (FIG. 6e), clavate (FIG. 6f) and ellipsoidal (FIG.

6g). Endogenous sporangia displayed tightly constricted necks (point between sporangia and

rhizoids) with narrow ports (arrows in FIG. 6d-g).

During exogenous thallus development, zoospore cysts germinated from both ends.

Rhizoids developed on one side while sporangiophores developed on the opposite side. The

empty zoospore cyst remained as a persistent swollen structure at the base of sporangiophore

(FIG. 6h). Exogenous sporangia developed at the end of unbranched sporangiophores of varied

lengths. Short sporangiophores had an average length of 6–20 μm (FIG. 6h-i), while long

sporangiophores extended up to 200 μm (FIG. 6j). Some of the short sporangiophores had

eggcup-shaped appearance (FIG. 6k). Exogenous sporangia were ellipsoidal (FIG. 6j), ovoid

(FIG. 6k), globose (FIG. 6l), constricted ellipsoidal (FIG. 6m), pyriform (FIG. 6n), bowling pin-

shaped (FIG. 6o), and rhomboidal (FIG. 6p). Sporangial necks were constricted with narrow port

(FIG. 6m-p). At maturity, a fine septum developed at the base of the sporangium (FIG. 6n & p,

arrow). Zoospores were released through an apical pore followed by collapse of the sporangial

wall (FIG. 6q).

Clade 5 (Domesticated Goat and Sheep) strains— Strain GFH683 produced globose zoospores

(FIG. 8a-b), with an average diameter of 5–6 μm. The majority of zoospores were mono-

flagellate with 1, or 2 flagella (FIG. 7a-b). Flagellum length ranged between 20–22 μm.

Zoospores germinated to produce both endogenous and exogenous monocentric thalli (FIG. 7c-

f), as evident by presence of a single sporangium per thallus, nucleated sporangia but anucleate

rhizoids. Endogenous sporangia were globose, sub-globose, ovoid, and obovoid (FIG. 7g) with

sizes ranging between 8–40 μm wide & 10–40 μm long. Exogenous sporangia were terminal and

varied in shape between globose, ovoid, and obovoid with sporangiophores that varied in length

from 20–80 μm (FIG. 7h-i). Zoospores discharge occurred through gradual dissolution of a wide

apical portion of sporangial wall, resulting in formation of an empty cup-shaped sporangium

(FIG. 7j-l). Such zoospore liberation patterns, and empty cup shaped sporangia were earlier

documented for *Piromyces minutus* (Ho and Barr 1995).

Clade 6 (Zebra-Horse) strains— Strain ZS-33 produced spherical zoospores, with an average

diameter of 10.8 ± 3 μm (average ± standard deviation of 54 zoospores, range 6–17 μm). All

zoospores were uniflagellate, with an average flagellum length of 26 ± 6.5 μm (average ±

standard deviation of 54 zoospores, range 18–40 μm) (FIG. 8a). After shedding their flagella,

zoospores started to encyst (FIG. 8b) and germinate producing germ tube (FIG. 8c). Germ tube

branched and developed a highly branched anucleate rhizoidal system (FIG. 8d-e). Both narrow,

0.5–2.5 μm wide, and broad, 3–12.5 μm wide, hyphae were observed; intercalary swellings were

frequently encountered in the broad hyphae (arrow in FIG. 8f).

Both endogenous and exogenous sporangia were observed. Endogenous sporangia varied

in shape and size. Small endogenous sporangia were mainly subglobose (20–60 μm in diameter).

(FIG. 8g). Large endogenous (80–160 μm L X 35–65 μm W) sporangia were mainly ellipsoidal

(FIG. 8h). Exogenous sporangia size ranged between (80–270 µm L X 35–85 µm W) and

displayed a wider range of morphologies, e.g. heart-shaped (FIG. 8k), ovoid (FIG. 8l) and

pyriform (FIG. 8m). Sporangiophores ranged in length between 20–400 µm. Characteristically,

strain ZS-33 displayed a multisporangiate thallus: the majority of sporangiophores were

branched and bore two to four sporangia (FIG. 8i-j). Similar sporangial morphology has

previously been observed in members of the genus *Piromyces* (e.g. *P. rhizinflatus*), and

Caecomyces (e.g. *C. communis*) (Akin and others 1988; Akin and others 1989; Breton and others

1991). Unbranched sporangiophores with single sporangia were less frequently encountered

(approximately 30% of observed sporangiophores n=50, Fig. 8k-m).

Zoospores were liberated through a wide apical pore at the top of the sporangia. The

sporangial wall stayed intact after the discharge (FIG. 8n-p). Further, mature sporangia

frequently detached from hyphae or sporangiophores, probably serving as an additional mean of

fungal dispersal (FIG. 8q).

The type strain ZS-33 was obtained from Zebra fecal samples collected at the Oklahoma

City Zoo. No noticeable differences were observed between ZS-33, and all other strains (n=15)

obtained from Zebra fecal samples from the Oklahoma City Zoo. On the other hand, two distinct

microscopic differences were identified in strains from domesticated Horses in Llanbadarn Fawr,

Ceredigion County, Wales. First, multisporangiate thalli, copiously observed in ZS-33, were

extremely rare in Welch Horse strains, and second, Distinct-resting stages (FIC. 8r) were often

observed in Welch Horse strains, but never in Oklahoma City Zebra strains. Whether these

differences are distinct characteristics of each group of strains, or induced by variations in media

composition as well as growth and incubation procedures remain to be seen.

Clade 7 (Nilgiri Tahr) strains— Strain TDFKJa193 produced globose zoospores (FIG. 9a), with

an average diameter of 3–4 μm. The majority of zoospores were mono-flagellate with 1, 2, or 3

flagella (FIG. 9a). Flagellum length ranged between 12–15 μm. Strain TDFKJa193 exhibited

both endogenous and exogenous monocentric thallus development (FIG. 9b-e). Endogenous

sporangia were terminal, varied in shape between globose, ovoid & obovoid, and ranged in size

between 10–70 μm wide & 12–100 μm long (FIG. 9f-g). Some endogenous sporangia showed

sub-sporangial swellings (FIG. 9f). Endogenous sporangia with one or two main rhizoidal

systems (FIG. 9f) and with branched rhizoidal system (FIG. 9g) were also observed. Exogenous

sporangia, on the other hand, were globose, ovoid & obovoid and were observed at the end of

short sporangiophores (12–20 µm) (FIG. 9h-k). Some of the sporangiophores ended with sub-

sporangial swellings with (FIG 9 i-j) or without (FIG. 9h&k) constricted neck of 1–8 μm width

and 2–10 μm length. The presence of subsporangial swellings and short sporangiophores were

previously reported for *Piromyces mae* (Ho and Barr 1995) and *Buwchfawromyces eastonii*

(Callaghan and others 2015). Mature exogenous sporangia often showed the formation of a

septum at their base (FIG. 9k) similar to *Neocallimastix frontalis* (Ho and Barr 1995). Zoospores

liberation happened after irregular dissolution of the sporangial wall (FIG. 9l).

Phylogenetic analysis— Phylogenetic analysis using LSU (FIG. 10a) placed the isolated strains into

seven monophyletic and bootstrap-supported lineages that were distinct from all currently described

AGF genera. LSU sequence divergence estimates between various strains within a single clade

ranged between 0–1%. The closest cultured representative to each of the seven clades is shown in

TABLE 1. Within the LSU taxonomic framework, strains recovered from Axis Deer clustered within

the *Orpinomyces-Neocallimastix-Pecoramyces-Feramyces* suprageneric clade, while strains

recovered from Boer Goat and domesticated goat clustered within the *Oontomyces-Anaeromyces-*

Liebetanzomyces supragenus clade. On the other hand, strains recovered from Nilgiri Tahr, as well as

strains recovered from domesticated Goat and Sheep formed two distinct new clades associated with

the genus *Buwchfawromyces*. In contrast, strains recovered from Zebra-Horse, and Mouflon-Boer

Goat formed two distinct clades associated together but with no specific affiliation to any

suprageneric group. The remaining group represented by strains isolated from White Tailed Deer

displayed no specific affiliation to any currently characterized genera or supragenus groups within the

Neocallimastigomycota.

To investigate ITS-1 sequence variability often reported within a single AGF strain, the ITS-1

region was amplified, cloned, and sequenced from all type strains. ITS-1 sequence divergence

within type strains ranged between 0% for strain TDFKJa193 representative of the Nilgiri Tahr

strains clade, and 0–8.4% (average 3.4%) for strain MS-2 representative of the Mouflon-Boer

Goat strains clade. ITS-1-based analysis confirmed the monophyletic and distinct nature of all

seven lineages, but yielded a different topology (FIG. 10b), as consistently observed in prior

studies (Hanafy and others 2017; Wang and others 2017). Of special note, was the surprisingly

high ITS-1 sequence similarity of the Boer Goat-domesticated Goat clade represented by strains

BGB-11 and GFKJa1916 (FIG. 10B), to an *Anaeromyces* sp. isolate GA-04 (GenBank accession

number FJ912851.1, unpublished) and to *Anaeromyces robustus* (GenBank accession number

NR_148182.1 (Li and others 2016)). Average ITS-1 sequence divergence between various

clones of the Boer Goat-domesticated Goat clade and *Anaeromyces* sp. GA-04 was 1.2%, and

between various clones of the Boer Goat-domesticated Goat clade and *A. robustus* and was

4.2%. Unfortunately, the LSU sequence data from both *Anaeromyces* sp. GA-04 and *A. robustus*

are not available for further comparison. This may be attributed to the inaccurate identification

and reporting of these two cultures, an issue very well known in the anaerobic fungal taxonomy

using ITS-1. We noted that in the paper describing *Anaeromyces robustus*, the type strain seems

to deviate from the typical morphology of the genus *Anaeromyces*, e.g. multi-flagellate

zoospores, lack of sausage shaped hyphae, and whale-tail like sporangia (see Figure 148 in (Li

and others 2016)), casting doubts on the phylogenetic affiliation of *Anaeromyces robustus* to the

genus *Anaeromyces*. Regardless, that stark morphological differences exist between Boer Goat-

domesticated Goat strains and all other members of the genus *Anaeromyces*, e.g. monocentric

thalli as opposed to polycentric thalli, absence of hyphal constrictions as opposed to sausage-

shaped hyphae with multiple constrictions, and monoflagellate (2–4 flagella) versus uniflagellate

zoospores (TABLE 1, FIG. 4-5). Such differences, in addition to the high ITS-1 sequence

divergence values between the Boer Goat-domesticated Goat clade and other members of the

genus *Anaeromyces* (7.1–13.1% to *A. mucronatus* and 8.5–18.6% to *A. contortus*), strongly

support the distinction between these strains and the genus *Anaeromyces*.

Ecological distribution— We queried the GenBank nr database to determine whether

representatives of these seven novel clades were encountered in prior ITS-1-based culture-

independent AGF diversity surveys. Multiple sequences with high (95.3-100%) sequence

similarity to Zebra-Horse strain ZS-33 ITS-1 sequence were identified. These sequences were

recovered from fecal samples obtained from multiple animals housed in the Oklahoma City Zoo

(Liggenstoffer and others 2010), as well as in various locations (left and right dorsal colon,

caecum and right ventral colon) within the digestive tract of horses (Mura and others 2019). This

group has previously been assigned the alphanumeric designation (AL1) (Kittelmann and others

2012; Liggenstoffer and others 2010).

Surprisingly, ITS-1 sequences from the remaining six lineages (Axis Deer, White-Tailed

Deer, Mouflon-Boer Goat, Boer Goat-domesticated Goat, domesticated Goat-domesticated Sheep,

and Nilgiri Tahr) bore no close resemblance to all currently available ITS-1 sequence data (TABLE

1), with highest similarity being 83% in White-Tailed Deer to sequences from Bontebok, 84% in

Mouflon-Boer Goat to sequences from Bontebok, 88% in domesticated goat-domesticated sheep to

sequences from horse, 89% in Nilgiri Tahr strains to sequences from Okapi, 91% in Axis Deer to

sequences from Llama, and 91-92% in Boer Goat-domesticated Goat strains to sequences from cow.

As such, representatives of these novel lineages, the absolute majority of which (5/6) recovered from

fecal samples of wild non-domesticated herbivores, do not appear to correspond to any of the

alphanumerically designated uncultured groups previously identified in prior culture-independent

efforts.

TAXONOMY

Agriosomyces Hanafy, Vikram B. Lanjekar, Prashant K. Dhakephalkar, T.M. Callaghan, Dagar,

G.W. Griff, Elshahed, and N.H. Youssef, gen. nov.

MycoBank ID: MB830737

Typification. Agriosomyces longus. Hanafy, Vikram B. Lanjekar, Prashant K. Dhakephalkar,

T.M. Callaghan, Dagar, G.W. Griff, Elshahed, and N.H. Youssef,

Etymology: Agrioso= derived from the Greek word for wild; *myces* = the Greek name for fungus.

Obligate anaerobic fungus that produces small spherical monoflagellate zoospores with an

extremely long flagellum (22 ± 3.8 μm). Zoospores germinate into monocentric thalli with

filamentous anucleate rhizoidal systems. Both endogenous and exogenous globose sporangia are

observed, which are very homogenous and display no pleomorphism. Rhizoids are swollen

below the sporangial neck, which is tightly constricted. Zoospores are released through

dissolution and rupturing of the sporangial wall. The clade is defined by the sequences

MK882010-MK882013 (ITS-1) and MK881996 (D1-D2 28S rDNA).

Agriosomyces longus Hanafy, Vikram B. Lanjekar, Prashant K. Dhakephalkar, T.M. Callaghan,

Dagar, G.W. Griff, Elshahed, and N.H. Youssef, sp. nov.

MycoBank ID: MB830738.

Typification: The holotype is FIG. 2g in this manuscript, derived from the following: U.S.A.

TEXAS: Val verde county, 29.369′ N and 100.829′ W ~300 m ASL, 3d old culture of isolate

MS-2, originally isolated from freshly deposited feces of male Mouflon Sheep (*Ovis orientalis*),

April 2018, *Radwa Hanafy.* Ex-type strain: MS2, GenBank: MK881996 (D1-D2 28S rDNA).

Etymology: The species epithet "*longus*" refers to the extremely long flagellum observed in

zoospores of strain MS-2 (FIG. 2a).

Obligate anaerobic fungus that produces small globose monoflagellate zoospores with an

average diameter of 4 ± 1.1 μm. Zoospores are mainly mono-flagellate with a flagellum length

of 22 ± 3.8 μm, approximately 5-6 times longer than the zoospore body. Bi-flagellate zoospores

are rarely encountered. Zoospores germinate into monocentric thalli with filamentous anucleate

rhizoidal systems. Both endogenous and exogenous sporangia are observed, which display no

pleomorphism and both show globose morphology. In endogenous sporangia, the rhizoids are

swollen below the sporangial neck, which is tightly constricted. Exogenous sporangia develop at

the end of swollen sporangiophores, and the sporangial neck is constricted with a narrow neck

port. Zoospores are released through dissolution and rupturing of the sporangial wall. Produces

small brown spherical colonies on agar, and a thin biofilm-like growth in liquid media. The clade

is defined by the sequences MK882010-MK882013 (ITS-1) and MK881996 (D1-D2 28S

rDNA).

Additional specimens examined: Radwa Hanafy strain MS-4 (GenBank accession number of D1-

D2 28S rDNA amplicon MK881997) isolated from the same freshly deposited feces of male

Mouflon Sheep (*Ovis orientalis*) from which the type strain originated, April 2018, and strain

BGS-13 (GenBank accession number of D1-D2 28S rDNA amplicon MK881995) isolated from

freshly deposited feces of female wild Boer Goat *(Capra aegagrus),* April, 2018.

Aklioshbomyces Hanafy, Vikram B. Lanjekar, Prashant K. Dhakephalkar, T.M. Callaghan,

Dagar, G.W. Griff, Elshahed, and N.H. Youssef, gen. nov.

MycoBank ID: MB830735

Typification: Aklioshbomyces papillarum Hanafy, Vikram B. Lanjekar, Prashant K.

Dhakephalkar, T.M. Callaghan, Dagar, G.W. Griff, Elshahed, and N.H. Youssef.

Etymology: Aklioshb= derived from the Arabic word for grass-eaters (herbivores); *myces* = the

Greek name for fungus.

Obligate anaerobic fungus that produces globose monoflagellate zoospores. Zoospores germinate

into monocentric thalli with filamentous anucleate rhizoids. Exhibits both endogenous and

exogenous monocentric thallus development. Exogenous sporangia develop at the end of

unbranched sporangiophores of varying length. No morphological differences are observed

between endogenous and exogenous sporangia, with ovoid, globose, and obpyriform sporangial

shapes noted. The clade is defined by the sequences MK882038-MK882042 (ITS-1) and

MK882001 (D1-D2 28S rDNA).

Aklioshbomyces papillarum Hanafy, Vikram B. Lanjekar, Prashant K. Dhakephalkar, T.M.

Callaghan, Dagar, G.W. Griff, Elshahed, and N.H. Youssef, sp. nov.

MycoBank ID: MB830736

Typification: The holotype is FIG. 3m in this manuscript, derived from the following: U.S.A.

OKLAHOMA: Payne county, 36.145′ N and 97.007′ W ~300 m ASL, 3d old culture of isolate

WT-2, originally isolated from freshly deposited feces of female White-Tailed Deer (*Odocoileus*

virginianus)*,* October 2017, *Radwa Hanafy.* Ex-type strain: WT-2, GenBank: MK882001 (D1-

D2 28S rDNA).

Etymology: The species epithet *"papillarum"* refers to the papillae observed on the majority of

strain WT-2 sporangia (FIG. 3m-q).

Obligate anaerobic fungus that produces globose monoflagellate zoospores with an average

diameter of 7.4 ± 2.4 μm. The majority of zoospores are monoflagellate, with zoospores with two

to three flagella less frequently observed. Fungal thalli are consistently monocentric with

filamentous anucleate rhizoids. Germination of zoospores produces two types of monocentric

thalli, endogenous and exogenous. Endogenous sporangia with single and two adjacent rhizoidal

systems are observed. Pseudo-intercalary endogenous sporangia are occasionally observed.

Sporangiophores carrying exogenous sporangia exhibit varying length from a few microns to 230

μm. Endogenous and exogenous sporangia are ovoid, globose, obpyriform, and ellipsoidal.

Sporangia are mostly papillated with one or two papillae. Produces beige circular colonies with a

brown central core of dense sporangial structures and an outer ring of light gray hyphal growth

on agar, and heavy growth of thick biofilms that firmly attached to the tube's glass surface in

liquid media. The clade is defined by the sequences MK882038-MK882042 (ITS-1) and

MK882001 (D1-D2 28S rDNA).

Additional specimens examined: Radwa Hanafy strains WT-1 (MK882000), WT-3 (MK881998),

WT-4 (MK881999), WT-41(MK882002), WTS-51 (MK882006), WTS-52 (MK882003), WTS-

53 (MK882004), and WTS-54 (MK882005), (GenBank accession number of D1-D2 28S rDNA

amplicon in parenthesis) isolated from the same freshly deposited feces of female White-Tailed

Deer from which the type strain originated, October 2017.

Capellomyces Hanafy, Vikram B. Lanjekar, Prashant K. Dhakephalkar, T.M. Callaghan, Dagar,

G.W. Griff, Elshahed, and N.H. Youssef, gen. nov.

MycoBank ID: MB830739.

Typification. Capellomyces foraminis. Hanafy, Vikram B. Lanjekar, Prashant K. Dhakephalkar,

T.M. Callaghan, Dagar, G.W. Griff, Elshahed, and N.H. Youssef.

Etymology: Capella= derived from the Latin word for Goat; *myces* = the Greek name for fungus.

Obligate anaerobic fungus that produces monoflagellate (1–3 flagella) zoospores. Zoospores

germinate into monocentric thalli with filamentous anucleate rhizoidal systems. Both

endogenous and exogenous sporangia are observed, with varying shapes and sizes. The clade is

defined by the sequences MK882007-MK882009 (ITS-1), and MK881975 (D1-D2 28S rDNA).

Capellomyces foraminis Hanafy, Vikram B. Lanjekar, Prashant K. Dhakephalkar, T.M.

Callaghan, Dagar, G.W. Griff, Elshahed, and N.H. Youssef, sp. nov.

MycoBank ID: MB830740

Typification: The holotype is FIG. 4n in this manuscript, derived from the following: U.S.A.

U.S.A. TEXAS: Val verde county, 29.369′ N and 100.829′ W ~300 m ASL, 3d old culture of

isolate BGB-11, originally isolated from freshly deposited feces of a female Boer Goat (*Capra*

aegagrus) April 2018, *Radwa Hanafy.* Ex-type strain: BGB-11, GenBank: MK881975 (D1-D2

28S rDNA).

Etymology: The species epithet "*foraminis*" refers to the wide apical pore at the top of the

sporangia through which zoospores are discharged.

Obligate anaerobic fungus that produces spherical monoflagellate zoospores. Zoospores start to

encyst after shedding their flagella. Zoospore cyst germinates, producing germ tube that

subsequently branches into monocentric thalli with filamentous anucleate rhizoidal systems.

Endogenous and exogenous sporangia are produced. Endogenous sporangia are ellipsoidal or

ovoid. Exogenous sporangia are formed at the end of un-branched sporangiophores (20–150

μm). Some of the sporangiophores exhibit sub-sporangial swellings. Exogenous sporangia are

ovoid, ellipsoidal with a single constriction, and globose. Zoospores are liberated through a wide

apical pore at the top of the sporangia followed by sporangial wall collapse. Colonies are small

(0.1–0.5 mm diameter) circular and brown, with dark center of sporangia structures on agar.

Produces thin fungal biofilm in liquid media. The clade is defined by the sequences MK882007-

MK882009 (ITS-1), and MK881975 (D1-D2 28S rDNA).

Additional species examined. Radwa Hanafy, strains BGB-2 (MK881974), BGC-12

(MK881976), BGS-11 (MK881977), and BGS-12 (MK881978) isolated from the same freshly

deposited feces of a female Boer Goat (*Capra aegagrus*), April 2018, from which the type strain

originated, April 2018 (D1-D2 28S rDNA amplicon GenBank accession number in parenthesis).

Capellomyces elongatus. Hanafy, Vikram B. Lanjekar, Prashant K. Dhakephalkar, T.M.

Callaghan, Dagar, G.W. Griff, Elshahed, and N.H. Youssef.

MycoBank: MB830869.

Typification: The holotype is FIG. 5k in this manuscript, derived from the following: India,

KERALA, town of Munnar, 10.219′ N and 77.106′ E ~2100 m ASL, 3d old culture of isolate

GFKJa1916, originally isolated from freshly deposited feces of a domesticated but forest grazing

Goat (*Capra aegagrus*), *Sumit S. Daggar.* Ex-type strain: GFKJa1916, GenBank: ITS-1

(MK775315), D1-D2 28S rDNA (MK775304).

Etymology: The species epithet "*elongatus*" refers to the characteristic long sporangiophore of

exogenous sporangia.

Obligate anaerobic fungus that produces globose monoflagellate (1, 2, or 3 flagella) zoospores.

Zoospore cyst germinate both endogenously and exogenously to produce monocentric thalli with

filamentous anucleate rhizoidal systems. Endogenous sporangia are cylindrical, elongate,

globose, sub-globose, ellipsoid & obovoid with sizes ranging between 8–60 μm wide & 10–140

μm long. Exogenous sporangia are formed at the end of developed at the end of long thick

sporangiophores (up to 300 µm). Multisporangiate thalli are commonly observed with two

sporangia of either the same or different shapes. Colonies are compact of 2–3 mm size, cottony

and off-white in color with a compact and fluffy center made up of thick sporangia type

structures, and surrounded by radiating rhizoids. Produces numerous fungal thalli that attach to

the glass bottles on initial days of growth, which later develop into thin mat-like structures in

liquid media. The clade is defined by the sequence MK775304 (D1-D2 28S rDNA).

Additional species examined. None

Ghazallomyces Hanafy, Vikram B. Lanjekar, Prashant K. Dhakephalkar, T.M. Callaghan, Dagar,

G.W. Griff, Elshahed, and N.H. Youssef, gen. nov.

MycoBank ID: MB830733

Typification: Ghazallomyces constrictus Hanafy, Vikram B. Lanjekar, Prashant K.

Dhakephalkar, T.M. Callaghan, Dagar, G.W. Griff, Elshahed, and N.H. Youssef.

Etymology: Ghazallo = derived from the Arabic word for Deer (Ghazalla); *myces* = the Greek

name for fungus.

Obligate anaerobic fungus that produces polyflagellate zoospores. Zoospores germinate into

monocentric thalli with highly branched anucleate rhizoidal systems. Exhibits both endogenous

and exogenous monocentric thallus development. Sporangia produced from endogenous and

exogenous thalli development are pleomorphic, exhibiting a wide range of sporangial shapes.

During exogenous thallus development, zoospore cysts germinate from both ends, with rhizoids

developing from one side and sporangiophore developing from the opposite side. The empty

zoospore cyst remains as a persistent swollen structure at the base of unbranched sporangiophore

that exhibit wide variations in lengths. Zoospores are released through an apical pore followed

by collapse of the sporangial wall. The clade is defined by the sequences MK882043 (ITS-1) and

MK881971 (D1-D2 28S rDNA).

Ghazallomyces constrictus Hanafy, Vikram B. Lanjekar, Prashant K. Dhakephalkar, T.M.

Callaghan, Dagar, G.W. Griff, Elshahed, and N.H. Youssef, sp. nov.

MycoBank ID: MB830734

Typification: The holotype is FIG. 6h in this manuscript, derived from the following: U.S.A.

TEXAS: Sutton county, 30.591′ N and 100.138′ W ~300 m ASL, 3d old culture of isolate Axs-

31, originally isolated from freshly deposited feces content of female Axis Deer (*Axis axis),* Apr.

2018, *Radwa Hanafy.* Ex-type strain: Axs-31, GenBank: MK881971 (D1-D2 28S rDNA).

Etymology: The species epithet *"constrictus"* refers to the observed constricted necks (point

between sporangia and rhizoids) in the species endogenous sporangia (FIG. 6d-g).

Obligate anaerobic fungus that produces globose polyflagellate zoospores with 7–14 flagella.

Zoospores germinate into monocentric thalli with highly branched anucleate rhizoidal systems.

Exhibits both endogenous and exogenous monocentric thallus development. Endogenous

sporangia produced from zoospore cyst enlargement develop into different shapes including

globose, tubular, clavate, and ellipsoidal. Endogenous sporangia display tightly constricted necks

(point between sporangia and rhizoids) with narrow ports. Exogenous sporangia develop at the

end of unbranched sporangiophores of varied lengths. Both short (6–20μm) and long (up to

200μm) sporangiophores are observed. The exogenous sporangia display ellipsoidal, ovoid,

globose, constricted ellipsoidal, pyriform, bowling pin-like, and rhomboidal shapes. Sporangial

necks are constricted with narrow port. A fine septum develops at the base of the sporangium at

maturity. Zoospores are released through an apical pore followed by collapse of the sporangial

wall. Produces small circular white colonies (1–4 mm diameter) with a brown central core of

dense sporangial structures on agar, and a thick fungal biofilm growth in liquid media. The clade

is defined by the sequences MK882043 (ITS-1) and MK881971 (D1-D2 28S rDNA).

Additional specimens examined: Radwa Hanafy strains ADC-2 (MK881965), ADS-14

(MK881966), AXS-33 (MK881967), AXS-34 (MK1881968), ADS-12 (MK881969), AXS-32

(MK881970), ADS-11 (MK881972), and ADS-21 (MK881973) (GenBank accession number of

D1-D2 28S rDNA amplicon in parenthesis) isolated from the same freshly deposited feces of

female Axis Deer (*Axis axis)* from which the type strain originated, April 2018.

Joblinomyces Hanafy, Vikram B. Lanjekar, Prashant K. Dhakephalkar, T.M. Callaghan, Dagar,

G.W. Griff, Elshahed, and N.H. Youssef, gen. nov.

MycoBank ID: MB830867

Typification. Joblinomyces apicalis. Hanafy, Vikram B. Lanjekar, Prashant K. Dhakephalkar,

T.M. Callaghan, Dagar, G.W. Griff, Elshahed, and N.H. Youssef.

Etymology: Joblino= honoring Keith N. Joblin for his contributions to the field of anaerobic

fungi; *myces* = the Greek name for fungus.

Obligate anaerobic fungus that produces globose monoflagellate zoospores. Both endogenous

and exogenous sporangia are observed with varying shapes and sizes. Sporangiophores of

exogenous sporangia vary in length. Exogenous sporangia have short and frequently swollen

sporangiophores. Zoospores discharge occurs through gradual dissolution of a wide apical

portion of sporangial wall, resulting in formation of an empty cup-shaped sporangium. The clade

is defined by the sequences MK910278 (ITS-1) and MK910268 (D1-D2 28S rDNA).

***Joblinomyces apicalis*.** Hanafy, Vikram B. Lanjekar, Prashant K. Dhakephalkar, T.M.

Callaghan, Dagar, G.W. Griff, Elshahed, and N.H. Youssef.

MycoBank ID: MB830868

Typification: The holotype is FIG. 7c in this manuscript, derived from the following: India,

HARYANA, city of Sonipat, 28.988′ N and 76.941′ E ~220 m ASL, 3d old culture of isolate

GFH683, originally isolated from freshly deposited feces of a domesticated goat (*Capra*

aegagrus hircus), *Sumit Dagar.* Ex-type strain: GFH683, GenBank: MK910268 (D1-D2 28S

rRNA).

Etymology: The species epithet *"apicalis"* refers to the zoospore discharge through the

dissolution of a wide apical portion of the sporangial wall.

Obligate anaerobic fungus that produces globose monoflagellate zoospores with 1, or 2 flagella.

Zoospores germinate to produce both endogenous and exogenous monocentric thalli.

Endogenous sporangia vary in shape between globose, sub-globose, ovoid, and obovoid with

sizes ranging between 8–40 μm wide & 10–40 μm long. Exogenous sporangia are terminal and

vary in shape between globose, ovoid, obovoid. Sporangiophores vary in length from 20–80 µm.

Zoospores discharge occur through gradual dissolution of a wide apical portion of sporangial

wall, resulting in formation of an empty cup-shaped sporangium. Produces 1–2 mm sized

colonies with a dense dark central core of abundant sporangial growth, surrounded by long and

thin radiating rhizoids. In liquid media, it produces numerous fungal thalli that attach to the glass

bottles on initial days of growth, and later develop into thin mat-like structures. The clade is

defined by the sequences MK910278-MK910282 (ITS-1) and MK910268-MK910272 (D1-D2

28S rDNA).

Additional specimens examined: Sumit Dagar strains GFH681 (MK910263-MK910267) and GFH682

(MK775330) (GenBank accession number of D1-D2 28S rDNA amplicon in parenthesis) isolated

from the same freshly deposited domesticated goat feces from which the type strain originated, and

SFH683 (MK775333) isolated from freshly deposited domesticated sheep feces in the city of Sonipat,

Haryana, India.

Khoyollomyces Hanafy, Vikram B. Lanjekar, Prashant K. Dhakephalkar, T.M. Callaghan, Dagar,

G.W. Griff, Elshahed, and N.H. Youssef, gen. nov.

MycoBank ID: MB830741

Typification. Khoyollomyces ramosus. Hanafy, Vikram B. Lanjekar, Prashant K. Dhakephalkar,

T.M. Callaghan, Dagar, G.W. Griff, Elshahed, and N.H. Youssef.

Etymology: Khyollo= derived from the Arabic word for horses; *myces* = the Greek name for

fungus.

Obligate anaerobic fungus that produces spherical uniflagellate zoospores. Zoospores encyst and

develop a highly branched anucleate rhizoidal system. Both endogenous and exogenous

sporangia are observed. Small endogenous sporangia are subglobose and large endogenous

sporangia are ellipsoidal. Exogenous sporangia displayed a wider range of shapes. The majority

of sporangiophores are branched and bear two to four sporangia. Unbranched sporangiophores

bearing a single sporangium are less frequently encountered. Zoospores are liberated through a

wide apical pore at the top of the sporangia. Mainly found in the digestive tracts of equids. The

clade is defined by the sequences MK882019 (ITS-1), and MK881981 (D1-D2 28S rDNA).

Khoyollomyces ramosus. Hanafy, Vikram B. Lanjekar, Prashant K. Dhakephalkar, T.M.

Callaghan, Dagar, G.W. Griff, Elshahed, and N.H. Youssef, sp. nov.

MycoBank ID: MB830742.

Typification: The holotype is FIG. 8j in this manuscript, derived from the following: U.S.A.

OKLAHOMA: Oklahoma City, 35.524′ N and 97.472′ W ~300 m ASL, 3d old culture of isolate

ZS-33, originally isolated from freshly deposited feces of a Grevy's Zebra *(Equus grevyi),* May,

2018, *Radwa Hanafy.* Ex-type strain: ZS-33, GenBank: MK881981 (D1-D2 28S rDNA).

Etymology: The species epithet *"ramosus"* (Latin for branched) refers to the observed branched

sporangiophores bearing two to four sporangia in *K. ramosus* type strain ZS-33 (FIG. 8i-k).

Obligate anaerobic fungus that produces spherical uniflagellate zoospores. Zoospores encyst and

germinate producing germ tube that develops into a highly branched anucleate rhizoidal system.

Both narrow, 0.5–2.5µm wide, and broad hyphae, 3–12.5µm wide, are produced; intercalary

swellings are frequently encountered in the broad hyphae. Both endogenous and exogenous

sporangia were observed. Endogenous sporangia vary in shape and size, with small endogenous

sporangia mainly subglobose (20–60 µm in diameter) while large endogenous (80–160µm L X

35–65µm W) sporangia mainly ellipsoidal. Exogenous sporangia ranged in size between (80–

270µm L X 35–85µm W)) and display a wide range of morphologies, e.g. heart-shaped, ovoid,

and pyriform. Displays a multisporangiate thallus, with the majority of sporangiophores being

branched and bearing two to four sporangia. Unbranched sporangiophores with single sporangia

are less frequently encountered (approximately 30% of observed sporangiophores). Zoospores

are liberated through a wide apical pore at the top of the sporangia. The sporangia stay intact

after the discharge. Mature sporangia frequently detach from hyphae or sporangiophores.

Produces small yellow to yellowish brown irregularly shaped colonies on agar. In liquid media,

the fungal growth is loose and exhibited a sand-like appearance. The clade is defined by the

sequences MK882019 (ITS-1), and MK881981 (D1-D2 28S rDNA).

Additional specimens examined: Radwa Hanafy strains ZC-31 (MK881979), ZC-32

(MK881980), ZC-33 (MK881981), ZC-41 (MK881982), ZC-42 (MK881983), ZC-43

(MK881984), ZC-51 (MK881985), ZC-53 (MK881986), ZS-21 (MK881987), ZS-22

(MK881988), ZS-31 (MK881989), ZS-32 (MK881990), ZS-41 (MK881992), ZS-42

(MK881993), and ZS-43 (MK881994) (GenBank accession number of D1-D2 28S rDNA

amplicon in parenthesis) isolated from the same freshly deposited Zebra feces from which the

type strain originated, May 2018. Tony Callaghan strains: HoCal4.A2, HoCal4.A2.2, HoCal4.A4

isolated from fresh horse feces (Llanbadarn, nr. Aberystwyth; 52.4156,-3.8878), August 2013.

Two further cultures (Tmc003.6a, TMC3.6b) were isolated from a different horse at the same

site, November 2013.

Tahromyces Hanafy, Vikram B. Lanjekar, Prashant K. Dhakephalkar, T.M. Callaghan, Dagar,

G.W. Griff, Elshahed, and N.H. Youssef, gen. nov.

MycoBank ID: MB830865

Typification. Tahromyces munnarensis. Hanafy, Vikram B. Lanjekar, Prashant K.

Dhakephalkar, T.M. Callaghan, Dagar, G.W. Griff, Elshahed, and N.H. Youssef.

Etymology: Tahro= referring to the Nilgiri Tahr from which the species was isolated; *myces =*

the Greek name for fungus.

Obligate anaerobic fungus that produces globose monoflagellate zoospores. Both endogenous

and exogenous sporangia are observed with varying shapes and sizes. Endogenous sporangia

with one or two main rhizoidal systems and with branched rhizoidal system are frequently

observed. Exogenous sporangia have short and frequently swollen sporangiophores. Sporangial

necks are frequently constricted. Septa often form at the base of mature exogenous sporangia.

Zoospores liberation happens after irregular dissolution of the sporangial wall. The clade is

defined by the sequences MK775321 (ITS-1), and MK775310 (D1-D2 28S rDNA).

Tahromyces munnarensis Hanafy, Vikram B. Lanjekar, Prashant K. Dhakephalkar, T.M.

Callaghan, Dagar, G.W. Griff, Elshahed, and N.H. Youssef.

MycoBank ID: MB830866

Typification: The holotype is FIG. 9h in this manuscript, derived from the following: India,

KERALA, town of Munnar, 10.219′ N and 77.106′ E ~2100 m ASL, 3d old culture of isolate

TDFKJa193, originally isolated from freshly deposited feces of a Nilgiri Tahr *(Nilgiritragus*

hylocrius), Sumit Dagar. Ex-type strain: TDFKJa193, GenBank: ITS-1 accession number

(MK775321), D1-D2 28S rDNA (MK775310).

Etymology: The species epithet *"munnarensis"* refers to the town that the type species was

isolated from.

Obligate anaerobic fungus that produces globose monoflagellate zoospores with 1, 2, or 3

flagella. Both endogenous and exogenous sporangia were observed. The sporangia vary in size

between 12–100 μm in length and 10–70 μm in width, and display a wide range of morphologies

like globose, ovoid, and obovoid. Sporangiophores are short (12–20 μm) with frequent

subsporangial swellings. Sporangial necks (1–8 μm width and 2–10 μm length) are frequently

constricted. Septa often form at the base of mature exogenous sporangia. Zoospores liberation

happens after irregular dissolution of the sporangial wall. Produces colonies that are small (1

mm), white in color with a compact and fluffy center, surrounded by dotted circles of fungal

thalli. In liquid media, it produces numerous fungal thalli attaching to the glass bottles on initial

days of growth, and later developing into thin mat-like structures. The clade is defined by the

sequence MK775310 (D1-D2 28S rDNA).

Additional specimens examined: Sumit Dagar strains TDFKJa1924 (MK775323), TDFKJa1926

(MK775322), and TDFKJa1927 (MK775324) (GenBank accession number of D1-D2 28S rDNA

amplicon in parenthesis) isolated from the same Nilgirir Tahr feces from which the type strain

originated.

DISCUSSION

Here, we report on the isolation and characterization of multiple novel AGF strains from a

concerted sampling effort of domesticated, zoo-housed, and wild animals from North America,

Europe, and Asia. We propose seven new AGF genera to accommodate these novel strains,

hence expanding the AGF genus-level diversity by more than 50% (From 11 to 18). All newly

described taxa produced filamentous, monocentric thalli, similar to seven of the eleven currently

described genera. Six of the seven novel genera described here produce mono/uniflagellate

zoospores, similar to eight of the eleven currently described taxa. As such, filamentous taxa with

moncentric thalli and monflagellate zoospore appear to be the most common thallus morphology

and zoospore flagellation patterns in the Neocallimastigomycota predominant within currently

described AGF genera. It is interesting to note that for decades, microscopic-based identification

of AGF strains typically assigned isolates with such morphology to the genus *Piromyces* (Ho and

others 1993). We note broad similarities between the microscopic features of *Aklioshbomyces*

papillarum and P. mae (papillated sporangia), and *Joblinomyces* and *P. Minutus* (zoospores

release through a wide apical portion of sporangial wall, resulting in formation of an empty cup-

shaped sporangium). Unfortunately, the absnece of sequence data and extant cultures of these

previously described "*Piromyces*" taxa prevents futher investigation into this issue (Ho and

others 1993).

The current isolates were obtained in a multi-year effort to describe novel AGF strains

from a wide range of animal hosts in the United States, India, Wales. The majority of novel taxa

described here (5/7 genera, 6/8 species) originated from wild undomesticated animals (Axis

Deer, White Tailed Deer, Mouflon, Boer Goat, and Nilgiri Tahr), underscoring their potential as

novel, yet-untapped reservoir of AGF diversity. Such novelty, which has recently been

postulated (Hanafy and others 2018) could be attributed to higher variability in the quality and

quantity of ingested plant material, and the significant daily and seasonal fluctuations in feeding

frequencies.

Culture-independent surveys utilizing ITS-1 as a phylogenetic marker (Kittelmann and others

2012; Liggenstoffer and others 2010), and subsequent meta-analysis (Kittelmann and others 2012;

Kittelmann and others 2013) have identified multiple novel yet-uncultured AGF genus-level lineages.

This study has been successful in isolating the first representatives of novel group AL1 from Zebra

and Horse fecal samples. In a prior survey of AGF in zoo-housed animals (Liggenstoffer and others

2010), members of this lineage were encountered in approximately half of the animal hosts examined

(18/35). AL1-affiliated sequences were more predominant in hindgut fermenters (7/9 hosts), and

comprised a relatively high proportion of the AGF community in multiple hosts, e.g. 99.9% in three

different Zebra individuals, 56.7 and 68.3% in two horses, and 29.6% in a Grant's Gazelle). By

comparison, they were only encountered in 11/26 foregut fermenters, where they constituted 0.01%

to 38% of the AGF community in these animals. Further a recent seminal spatial analysis that

analyzed AGF community in samples directly obtained from various locations along horses digestive

tracts (Mura and others 2019) identified AL1 group as a prominent component of the AGF

community in the right ventral (88%), and left dorsal (98%) colons in horses. As such, this novel

genus appears to exhibit a preference for hindgut fermenters of the family Equidae. The reason for

such preferences, and the general preference of some fungal taxa to specific hosts remains unclear

(Callaghan and others 2015; Dagar and others 2015).

Surprisingly, comparative analysis of ITS-1 sequences indicated that the Axis Deer, White

tailed Deer, Mouflon-Boer Goat, Boer Goat-domesticated Goat, Nilgiri Tahr, and domesticated Goat-

sheep groups appear to be completely novel, and previously unencountered in prior culture-based or

culture-independent studies. ITS-1 sequences from these four isolates did not display mismatches to

common ITS-1 primers, did not have an atypical length that could hinder its amplification or

detection via PCR, and were readily amplified from pure-cultures' genomic DNA. As such, we posit

that the lack of prior observation of these taxa is biologically relevant, and is indicative of their

relatively specific host preference and/or predominance in wild, rather than domesticated herbivores.

Indeed, although 30 animals were screened in the current study, three of these seven novel genera

were isolated only from a single host (White tailed Deer, Axis Deer, and Nilgiri Tahr), while the

other four were isolated from only two hosts (Mouflon and Boer Goat, Boer Goat and domesticated

Goat, domesticated Goat and Sheep, and Zebra-domesticated Horse) (TABLE 1).

Collectively, the steady identification of novel taxa in culture-based and culture-independent

surveys, as well as the sparse overlap between these studies strongly suggests that the scope of AGF

diversity in nature is much broader than currently estimated (Kittelmann and others 2012; Paul and

others 2018). Compared to the prokaryotic component of the rumen and herbivorous gut, the diversity

of the rumen mycobiome remains woefully understudied. To provide a more thorough understanding

of the AGF diversity in nature, concerted efforts that systematically assess the AGF diversity and

community structure in various spatial (e.g. across various compartments of the herbivorous gut),

temporal (e.g. across the lifespan of an animal), and geographic dimensions in a wide range of

domesticated and wild herbivores is needed. Much remains to be understood regarding the diversity

and community structure of AGF within various locations of the gastrointestinal tract of an animal

host, interspecies stochastic differences between AGF communities in animal subjects, temporal age-

related progression of AGF in animal hosts, and the response of the AGF community to various

factors e.g. feeding patterns, antibiotic administration, animal disease, and co-housing arrangements

and combinations thereof.

ACKNOWLEDGMENTS We thank Jim and Tammy Austin for providing fecal samples,

Britny Johnson for technical assistance, Dr. Karthick Balasubramanian for his valuable help, Drs.

Rebecca Snyder and Jennifer D'Agostino for providing fecal samples from the Oklahoma City

Zoo, and to the Kerala Forest Department, Munnar for sampling permission (WL10-

51713/2017). This work has been funded by the NSF-DEB Grant number 1557102 to NHY and

MSE, Department of Biotechnology (DBT) project no. BT/PR15694/PBD/26/506/2015 to SSD.

LITERATURE CITED

Akin D, Borneman W, Windham W. 1988. Rumen fungi: morphological types from Georgia

cattle and the attack on forage cell walls. Biosystems 21 385–391.

Akin D, Lyon C, Windham W, Rigsby L. 1989. Physical degradation of lignified stem tissues by

ruminal fungi. Applied and Environmental Microbiology 55:611–616.

Ariyawansa HA, Hyde KD, Jayasiri SC, Buyck B, Thilini Chethana KW, Dai DQ, Dai YC,

Daranaama DA, Jayawardena RS, Lücking R, Ghobad-Nejhad M, Niskanen T, Thambugala KM,

Voigt K, Zhao RL, Li G-J, Doilom M, Boonmee S, Yang, ZL, Cai Q, Cui Y-Y, Bahkali AH,

Chen J, Cui BK, Chen JJ, Dayarathne MC, Dissanayake AJ, Ekanayaka AH, Hashimoto A,

Hongsanan S, Jones EBG, Larsson E, Li WJ, Li Q-R, Liu JK, Luo ZL, Maharachchikumbura

SSN, Mapook A, McKenzie EHC, Norphanphoun C, Konta S, Pang KL, Perera RH,

Phookamsak R, Phukhamsakda C, Pinruan U, Randrianjohany E, Singtripop C, Tanaka K, Tian

CM, Tibpromma S, Abdel-Wahab MA, Wanasinghe DN, Wijayawardene NN, Zhang JF, Zhang

H, Abdel-Aziz FA, Wedin M, Westberg M, Ammirati JF, Bulgakov T, Lima DX, Callaghan TM,

Callac P, Chang CH, Coca LF, Dal-Forno M, Dollhofer V, Fliegerová K, Greiner K, Griffith

GW, Ho H-M, Hofstetter V, Jeewon R, Kang JC, Kirk PM, Kytövuori I, Lawrey JD, Xing J, Li

H, Liu ZY, Liu XZ, Liimatainen K, Lumbsch HT, Matsumura M, Moncada B, Nuankaew S,

Parnmen S, de Azevedo Santiago ALCM, Sommai S, de Souza CAF, de Souza-Motta CM, Su

HY, Suetrong S, Wang Y, Wei S-F, Wen T-C, Yuan HS, Zhou LW, Réblová M, Fournier J,

Camporesi E, Luangsa-ard JJ, Tasanathai K, Khonsanit A, Thanakitpipattana D, Somrithipol S,

Diederich P, Millanes AM, Common RS, Stadler M, Yan JY, Li X, Lee HW, Nguyen TTT, Lee

HB, Battistin B, Marsico O, Vizzini A, Vila J, Ercole E, Eberhardt U, Simonini G, Wen H-A,

Chen X-H, Miettinen O, Spirin V, Hernawati. 2015. Fungal diversity notes 111–252 - taxonomic

and phylogenetic contributions to fungal taxa. Fungal Diversity 75:27– 274.

Barr D, Yanke L, Bae H, McAllister T, Cheng K. 1995. Contributions on the morphology and

taxonomy of some rumen fungi from Canada. Mycotaxon 54:203–214.

Barr DJ, Kudo H, Jakober KD, Cheng KJ. 1989. Morphology and development of rumen fungi:

Neocallimastix sp., *Piromyces communis*, and *Orpinomyces bovis* gen.nov., sp.nov. Canadian

Journal of Botany 67 2815–2824.

Breton A, Bernalier A, Dusser M, Fonty G, Gaillard-Martinie B, Guitlot J. 1990. *Anaeromyces*

mucronatus nov. gen., nov. sp. A new strictly anaerobic rumen fungus with polycentric thallus.

FEMS Microbiology Letters 70:177–182.

Breton A, Dusser M, Gaillard-Martine B, Guillot J, Millet L, Prensier G. 1991. *Piromyces*

rhizinflata nov. sp., a strictly anaerobic fungus from faeces of the Saharian ass: a morphological,

metabolic and ultrastructural study. FEMS Microbiology Letters 82:1–8.

860
Calkins S, Elledge NC, Hanafy RA, Elshahed MS, Youssef NH. 2016. A fast and reliable

procedure for spore collection from anaerobic fungi: Application for RNA uptake and long-term

storage of isolates. Journal of Microbiological Methods 127:206–213.

Callaghan TM, Podmirseg SM, Hohlweck D, Edwards JE, Puniya AK, Dagar SS, Griffith GW.

2015. *Buwchfawromyces eastonii* gen. nov., sp. nov.: a new anaerobic fungus

(Neocallimastigomycota) isolated from buffalo faeces. Mycokeys 9:11–28.

Dagar SS, Kumar S, Griffith GW, Edwards JE, Callaghan TM, Singh R, Nagpal AK, Puniya AK.

2015. A new anaerobic fungus (*Oontomyces anksri* gen. nov., sp. nov.) from the digestive tract

of the Indian camel (*Camelus dromedarius*). Fungal Biology 19:731–737.

Gold JJ, Heath IB, Bauchop T. 1988. Ultrastructural description of a new chytrid genus of

caecum anaerobe, *Caecomyces equi* gen. nov., sp. nov., assigned to the Neocallimasticaceae

BioSystems 21:403–415.

Gruninger RJ, Puniyab AK, Callaghanc TM, Edwardsc JE, Youssef N, Dagare SS, Fliegerova K,

Griffith GW, Forster R, Tsang A et al. 2014. Anaerobic Fungi (Phylum Neocallimastigomycota):

Advances in understanding of their taxonomy, life cycle, ecology, role, and biotechnological

potential. FEMS Microbiology Ecology. 90:1–17.

881
Hanafy RA, Elshahed MS, Liggenstoffer AS, Griffith GW, Youssef NH. 2017. *Pecoramyces*

ruminantium, gen. nov., sp. nov., an anaerobic gut fungus from the feces of cattle and sheep.

Mycologia 109:231– 243.

Hanafy RA, Elshahed MS, Youssef NH. 2018. *Feramyces austinii*, gen. nov., sp. nov., an

anaerobic gut fungus from rumen and fecal samples of wild Barbary sheep and fallow deer.

Mycologia 110:513–525.

Heath BI, Bauchop T, Skipp RA. 1983. Assignment of the rumen anaerobe *Neocallimastix*

frontalis to the Spizellomycetales (Chytridiomycetes) on the basis of its polyflagellate zoospore

ultrastructure. Canadian Journal of Botany 61:295–307.

Ho YW, Bar DJS, Abdullah N, Jalaludin S, Kudo H. 1993a. A new species of *Piromyces* from

the rumen of deer in Malaysia. Mycotaxon 47:285–293.

Ho YW, Barr DJS. 1995. Classification of anaerobic gut fungi from herbivores with emphasis on

rumen fungi from malaysia. Mycologia 87:655–677.

Hungate RE. 1969. A roll tube method for cultivation of strict anaerobes. Methods in

Microbiology 3:117–132.

Joshi A, Lanjekar VB, Dhakephalkar PK, Callaghan TM, Griffith GW, Dagar SS. 2018.

Liebetanzomyces polymorphus gen. et sp. nov., a new anaerobic fungus (Neocallimastigomycota)

isolated from the rumen of a goat. Mycokeys 40:89–110.

Kittelmann S, G. E. Naylor, J. P. Koolaard, Janssen PH. 2012. A Proposed Taxonomy of

Anaerobic Fungi (Class Neocallimastigomycetes) Suitable for Large-Scale Sequence-Based

Community Structure Analysis. PLoS ONE 7:e36866.

Kittelmann S, H. Seedorf, W. A. Walters, J. C. Clemente, R. Knight, J. I. Gordon, Janssen PH.

2013. Simultaneous Amplicon Sequencing to Explore Co-Occurrence Patterns of Bacterial,

Archaeal and Eukaryotic Microorganisms in Rumen Microbial Communities. PLoS ONE

8:e47879.

Kumar S, Stecher G, Tamura K. 2016. MEGA7: Molecular evolutionary genetics analysis

version 7.0 for bigger datasets. Molecular Biology and Evolution 33:1870–1874.

Li GJ, Hyde KD, Zhao RL, Hongsanan S, Abdel-Aziz FA, Abdel-Wahab MA, Alvarado P,

Alves-Silva G, Ammirati JF, Ariyawansa HA, Baghela A, Bahkali AH, Beug M, Bhat DJ,

Bojantchev D, Boonpratuang T, Bulgakov TS, Camporesi E, Boro MC, Ceska O, Chakraborty D,

Chen JJ, Chethana KWT, Chomnunti P, Consiglio G, Cui BK, Dai DQ, Dai YC, Daranagama

DA, Das K, Dayarathne MC, De Crop E, De Oliveira RJV, de Souza CAF, de Souza JI,

Dentinger BTM, Dissanayake AJ, Doilom M, Drechsler-Santos ER, Ghobad-Nejhad M, Gilmore

SP, Góes-Neto A, Gorczak M, Haitjema CH, Hapuarachchi KK, Hashimoto A, He MQ, Henske

JK, Hirayama K, Iribarren MJ, Jayasiri SC, Jayawardena RS, Jeon SJ, Jerônimo GH, Jesus AL,

Jones EBG, Kang JC, Karunarathna SC, Kirk PM, Konta S, Kuhnert E, Langer E, Lee HS, Lee

HB, Li WJ, Li XH, Liimatainen K, Lima FX, Lin CG, Liu JK, Liu XZ, Liu ZY, Luangsa-ard JJ,

Lücking R, Lumbsch HT, Lumyong S, Leaño EM, Marano AV, Matsumura M, McKenzie EHC,

Mongkolsamrit S, Mortimer PE, Nguyen TTT, Niskanen T, Norphanphoun C, O'Malley MA,

Parnmen S, Pawłowska J, Perera RH, Phookamsak R, Phukhamsakda C, Pires-Zottarelli CLA,

Raspé O, Reck MA, Rocha SCO, de Santiago ALCMA, Senanayake IC, Setti L, Shang QJ,

Singh SK, Sir EB, Solomon KV, Song J, Srikitikulchai P, Stadler M, Suetrong S, Takahashi H,

Takahashi T, Tanaka K, Tang LP, Thambugala KM, Thanakitpipattana D, Theodorou MK,

Thongbai B, Thummarukcharoen T, Tian Q, Tibpromma S, Verbeken A, Vizzini A, Vlasák J,

Voigt K, Wanasinghe DN, Wang Y, Weerakoon G, Wen HA, Wen TC, Wijayawardene NN,

Wongkanoun S, Wrzosek M, Xiao YP, Xu JC, Yan JY, Yang J, Da Yang S, Hu Y, Zhang JF,

Zhao J, Zhou LW, Peršoh D, Phillips AJL, Maharachchikumbura SSN. 2016. Fungal diversity

notes 253–366: taxonomic and phylogenetic contributions to fungal taxa. Fungal Diversity 78:1–

237.

Li J, Health B, Bauchop T. 1990. *Piromyces mae* and *Piromyces dumbonica*, two new species of

uniflagellate anaerobic chytridiomycete fungi from the hindgut of the horse and elephant.

Canadian Journal of Botany 68:1021–1033.

Liggenstoffer AS, Youssef NH, Couger MB, Elshahed MS. 2010 Phylogenetic diversity and

community structure of anaerobic fungi (Phylum Neocallimastigomycota) in ruminant and non-

ruminant herbivores. The ISME Journal 4:1225–1235.

Ljungdahl LG. 2008 The cellulase/hemicellulase system of the anaerobic fungus *Orpinomyces*

PC-2 and aspects of its use. Annals of the New York Academy of Sciences 1125 308–321.

McSweeney CS, Denman SE, Mackie RI. 2005. Rumen bacteria. In: Makkar HPS, McSweeney

CS, eds. Methods in Gut Microbial Ecology for Ruminants. Dordrecht, Netherlands: Springer. p.

23–37.

Mura E, Edwards J, Kittelmann S, Kaerger K, Voigt K, Mrázek J, Moniello G, Fliegerova K.

2019. Anaerobic fungal communities differ along the horse digestive tract. Fungal Biology

123:240–246.

Nagpal R, Puniya AK, Sehgal JP, Singh K. 2011. In vitro fibrolytic potential of anaerobic rumen

fungi from ruminants and non-ruminant herbivores. Mycoscience 52:31–38.

Nakamura T, Yamada KD, Tomii K, Katoh K. 2018. Parallelization of MAFFT for large-scale

multiple sequence alignments. Bioinformatics 34:2490–2492.

Ozkose E, Thomas BJ, Davies DR, Griffith GW, Theodorou MK. 2001. *Cyllamyces aberensis*

gen. nov. sp.nov., a new anaerobic gut fungus with branched sporangiophores isolated from

cattle. Canadian Journal of Botany 79:666–673.

Paul SS, Bu D, Xu J, Hyde KD, Yu Z. 2018. A phylogenetic census of global diversity of gut

anaerobic fungi and a new taxonomic framework. Fungal Diversity 89:253–266.

Paul SS, Deb SM, Punia BS, Singh D, Kumar R. 2010. Fibrolytic potential of anaerobic fungi

(*Piromyces* sp.) isolated from wild cattle and blue bulls in pure culture and effect of their

addition on in vitro fermentation of wheat straw and methane emission by rumen fluid of

buffaloes. Journal of the Science of Food and Agriculture 90:1218–1226.

Ranganathan A, Smith OP, Youssef NH, Struchtemeyer CG, Atiyeh HK, Elshahed MS. 2017.

Utilizing anaerobic fungi for two-stage sugar extraction and biofuel production from

lignocellulosic biomass. Frontiers in Microbiology 8:635.

Tuckwell DS, Nicholson MJ, McSweeney CS, Theodorou MK, Brookman JL. 2005. The rapid

assignment of ruminal fungi to presumptive genera using ITS-1 and ITS2 RNA secondary

structures to produce group-specific fingerprints. Microbiology 151:1557–1567.

986
Wang X, Liu X, Groenewald JZ. 2017. Phylogeny of anaerobic fungi (phylum

Neocallimastigomycota), with contributions from yak in China. Antonie Van Leeuwenhoek

110:87–103.

Young D, Dollhofer V, Callaghan TM, Reitberger S, Lebuhn M, Benz JP. 2018. Isolation,

identification and characterization of lignocellulolytic aerobic and anaerobic fungi in one- and

two-phase biogas plants. Bioresoure Technology 268:470–479.

Youssef NH, Couger MB, Struchtemeyer CG, Liggenstoffer AS, Prade RA, Najar FZ, Atiyeh

HK, Wilkins MR, Elshahed MS. 2013. Genome of the anaerobic fungus *Orpinomyces* sp. C1A

reveals the unique evolutionary history of a remarkable plant biomass degrader. Applied and

Environmental Microbiology 79:4620–4634.

1000

1001

1002

1003

LEGENDS AND FOOTNOTES.

FIG. 1. Macroscopic features of colony morphology on agar media and fungal biomass in liquid

media for (a) *Agriosomyces longus* strain MS2, brown circular colonies and thin fungal biofilm

in liquid medium. (b) *Aklioshbomyces papillarum* strain WT-2, beige circular colonies with a

brown central core of sporangia and thick fungal biofilm in liquid medium that firmly attaches to

the tube's glass surface, note the inverted tube shows how the fungus thalli attached to the tube.

(c) *Capellomyces foraminis* strain BGB-11, small brown circular colonies with dark center and

thin fungal biofilm in liquid medium. (d) *Capellomyces elongatus* strain GFKJa1916, compact

cottony off-white circular colonies with a fluffy central core of sporangia surrounded by

radiating sporangia and thin fungal biofilm in liquid medium that attach to the tube's glass. (e)

Ghazallomyces constrictus strain Axs-31, white circular colonies on roll tubes and thick fungal

biofilm in liquid medium. (f) *Joblinomyces apicalis* strain GFH683, beige circular colonies with

a brown central core of sporangia and thin fungal biofilm in liquid medium. (g) *Khyollomyces*

ramosus strain ZS-33, yellowish brown colonies of irregular shape on agar medium and loose

fungal thalli with sand-like appearance. (h) *Tahromyces munnarensis* strain TDFKJa193 white

circular colonies with fluffy center, surrounded by dotted circles of fungal thalli and thick fungal

biofilm in liquid medium that attach to the tube's glass.

FIG. 2. Microscopic features of *Agriosomyces longus* (Clade 1, Mouflon-Boer Goat) strain

MS2. Light (e-h) and scanning electron (a and i) micrographs are shown. DAPI staining for

nuclei visualizing using a fluorescence microscope equipped with a Brightline DAPI high-

contrast filter set (c). Overlay image is shown in (d). (a) A monoflagellate zoospore. (b) A

biflagellate zoospore. (c-d) Monocentric thalli, with nuclei occurring in sporangia, not in rhizoids

or sporangiophores. (e-f) Endogenous globose sporangia with tightly constricted necks and

subsporangial swellings (arrows). (g-h) Exogenous globose sporangia, note the swollen

sporangiophores. (i) An empty sporangium after zoospore release and rupturing of the sporangial

wall. Bar: a=5 μm, b=20 μm c-i=50 μm.

FIG. 3. Microscopic features of *Aklioshbomyces papillarum* (Clade 2, White-Tailed Deer) strain

WT-2. Light (a-e, and g-q) and scanning electron (f) micrographs are shown. Light microscopy

pictures were examined after staining with lactophenol cotton blue (a-c, e, and g-q), as well as

following nuclei staining with DAPI (d). (a) A monoflagellate zoospore. (b) A biflagellate

zoospore. (c) A triflagellate zoospore. (d) Monocentric thalli, with nuclei occurring in sporangia,

not in rhizoids or sporangiophores. (e-g) Endogenous sporangial development: (e) Ovoid

sporangium with single rhizoidal system, (f) Ovoid sporangium with two main rhizoidal systems,

(g) Globose pseudo-intercalary sporangium, between two main rhizoidal systems. (h-k)

Exogenous sporangial development: (h) Globose sporangium on a very short sporangiophore, (i)

Ellipsoidal sporangium, (j) Obpyriform sporangium on a long sporangiophore, (k) Obpyriform

sporangium. (l) Ellipsoidal sporangium. (m) Sporangia with lateral single papilla, (n-p)

Sporangia with terminal single papilla. (q) Sporangium with two papillae. Bar: a-c, e, f, n & o

=20 μm, g-k, p & q =50 μm, d, l & m =100 μm.

FIG. 4. Microscopic features of *Capellomyces foraminis* (Clade 3, Boer Goat) strain BGB-11.

Light (a-i, l, o & p) and scanning electron (j, k, m, n, q &r) micrographs are shown. Light

microscopy pictures were examined after staining with lactophenol cotton blue (a-e, h, i, & o), as

well as following nuclei staining with DAPI (f). Overlay image is shown in (g). (a) A

monoflagellate zoospore. (b) A biflagellate zoospore. (c) Zoospore cyst, arrow points to the shed

flagellum. (d) Germinating zoospore cyst producing a germ tube (arrow). (e) Rhizoidal system

development. (f-g) Monocentric thalli with nuclei occurring in sporangia, not in rhizoids or

sporangiophores. (h-i) Endogenous sporangial development: (h) Ellipsoidal sporangium with two

main rhizoidal systems (arrows), (i) Ovoid sporangium with single rhizoidal system. (j-q)

Exogenous sporangial development: (j) Exogeous sporangium with a short sporangiophore (Sp),

note the empty zoospore cyst (Zc). (k) Ovoid sporangium with a long sporangiophore, (l) Ovoid

sporangium on a long sporangiophore ends with sub-sporangial swelling (arrow), (m) Collapsed

empty sporangium on a long sporangiophore ends with sub-sporangial swelling (arrow), (n)

Constricted ellipsoidal sporangium, (o-p) Globose Sporangia. (q) Zoospores ae released through

apical pore. (r) An empty sporangium following zoospores release. Abbreviations: (Sp),

sporangiophore; (Zc), zoospore cyst. Bar: a-j, l, o & p =20 μm, k, m, n, q & r =50 μm.

Fig 5. Microscopic features of *Capellomyces elongatus* (Clade 3, domesticated Goat) strain

GFKJa1916**.** Differential interference contrast (a, d-f and i-l), scanning (g-h), phase contrast (b)

and fluorescence (c) micrographs. (a) A monoflagellate zoospores. (b-c) Monocentric thalli;

nuclei were observed in sporangia, not in rhizoids or sporangiophores. (d-g) Endogenous

sporangia. (d) Globose endogenous sporangium with one main rhizoidal systems. (e-f)

Endogenous sporangia with multiple rhizoidal systems. (g) Endogenous sporangium on wheat

straw fibers. (h-l) Exogenous sporangia: (h) ovoid-shaped sporangium with long sporangiophore.

(i) Multiple ovoid, and globose exogenous sporangia with long sporangiophores. (j)

Multisporangiate thallus with two sporangia (same shape). (k-l) Multisporangiate thallus with

two sporangia (different shape).

Scale bar = 20 μM.

FIG. 6. Microscopic features of *Ghazallomyces constrictus* (Clade 4, Axis Deer) strain Axs-31.

Light (a-h, f, k and m-p) and scanning electron (i, j, l and q) micrographs are shown. Light

microscopy pictures were examined after staining with lactophenol cotton blue (a, d–h, k, and m-

p), as well as following nuclei staining with DAPI (b). Overlay image is shown in (c). (a) A

polyflagellate zoospore. (b-c) Monocentric thalli, with nuclei occurring in sporangia, not in

rhizoids or sporangiophores. (d-g) Endogenous sporangia with tightly constricted necks (arrows):

(d) Young globose sporangium, (e) Young tubular sporangium, (f) Mature clavate sporangium,

(g) and Mature ellipsoidal sporangium. (h-p) Exogenous sporangia: (h) Young sporangium on a

short flattened sporangiophore (Sp), note the persistent empty zoospore cyst (Zc) and the

rhizoidal system (R), (i) Ovoid sporangium on short sporangiophore, (j) Ellipsoidal sporangium

on long sporangiophore, (k) ovoid sporangium on an eggcup-shaped sporangiophore (arrow), (l)

Globose sporangium, (m) Constricted ellipsoidal sporangium with tightly constricted neck

(arrow) on long sporangiophore, (n) Pyriform sporangium, note the fine septum at the base of

sporangium (arrow), (o) Bowling pin-shaped sporangium, (p) Rhomboidal sporangium with

constricted neck and fine septum (arrows), note the persistent empty zoospore cyst (Zc). (q)

Zoospores are released through apical pore followed by collapse of the sporangial wall.

Abbreviations: (Sp), sporangiophore; (Zc), zoospore cyst; (R), rhizoid. Bar: a=20μm, b-q=50

μm.

FIG. 7. Microscopic features of *Joblinomyces apicalis* (Clade 5, domesticated Goat and Sheep)

strain GFH683. Phase contrast (b, c and e), fluorescence (d, f), scanning electron (g-h, and m)

differential interference (k-l) micrographs. (a) A monoflagellate zoospore. (b) A biflagellate

spherical zoospore. (c-f) Monocentric thalli; nuclei were observed in sporangia, not in rhizoids or

sporangiophores, note the empty cup-shaped sporangium after zoospore release (arrow). (g)

Ovoid endogenous sporangium. (h-i) Exogenous sporangia: (h) Ovoid sporangium with short

sporangiophore. (i) Subglobose sporangium with long sporangiophore. (j-l) Zoospore release: (j)

Dissolution of the apical portion of sporangial wall (arrow). (k-l) Cup-shaped sporangia with

wide apical pores and intact sporangial walls (arrow). (g) Colonization of rice straw fibers by

fungal rhizoids and emanating.

Scale bar = 20 µM.

FIG. 8. Microscopic features of *Khyollomyces ramosus* (Clade 6, Zebra-Horse) strain ZS-33 (a-

q) and distinct resting stage structure from strain HoCal4.A2.2 (r). Light (a-e, g, i-j, l & q) and

scanning electron (f, h, k, m &n-p) micrographs are shown. DAPI staining for nuclei visualizing

using a fluorescence microscope equipped with a Brightline DAPI high-contrast filter set (d).

Overlay image is shown in (e). (a) A uniflagellate zoospore. (b) Zoospore cyst after shedding of

the flagellum. (c) Germinating zoospore cyst producing a germ tube (arrow). (d-e) Monocentric

thalli, with nuclei occurring in sporangia, not in rhizoids or sporangiophores. (f) hyphal

structures with intercalary swellings in wide hyphae (arrows). (g-h) Endogenous sporangial

development: (g) Young subglobose sporangium with single rhizoidal system, (i) Mature

ellipsoidal sporangium with two main rhizoidal systems. (i-m) Exogenous sporangial

development: (i) Multisporangiate thallus with two sporangia, (j) Multisporangiate thallus with

four sporangia, (k) Heart-shaped sporangium. (l) Ovoid sporangium (labeled S) on a wide

flattened sporangiophore (labeled Sp), (m) Pyriform sporangium. (n) Zoospores are released

through apical pore. (o-p) Empty sporangia with intact sporangial walls after zoospores

discharge. (q) Mature sporangia detached from hyphae or sporangiophores. (r) Resting stages

from strain HoCal4.A2.2. Bar: a-c & f-g =20 µm, d-e, k &m-q =50 µm and h-j &l =100 µm.

FIG. 9. Microscopic features of *Tahromyces munnarensis* (Clade 7, Nilgiri Tahr) strain

TDFKJa193. Differential interference contrast (a, f and i-l), phase contrast (b, d, g and h),

fluorescence (c and e) micrographs. (a) A monoflagellate and triflagellate zoospores. (b-e)

Monocentric thalli; nuclei were observed in sporangia, not in rhizoids or sporangiophores. (f-g)

Endogenous sporangia. (f) Ovoid endogenous sporangium with two rhizoidal systems, note the

sub-sporangial swelling (arrow). (g) Endogenous sporangium with branched rhizoids. (h-l)

Exogenous sporangia: (h) Globose sporangium with short swollen sporangiophore (arrow). (i-j)

Exogenous sporangia with sub-sporangial swellings and constricted necks (arrow). (k) Ovoid

sporangium with septum at the sporangial base (arrow). (l) Zoospore release through dissolution

of a wide apical pore. Scale bar = 20 µM.

FIG. 10. Phylogenetic affiliation of the 7 newly described genera to other AGF genera based on

the sequences of the D1–D2 domains of nuc 28S rDNA gene (a), and partial ITS-1 sequences

(b). Sequences were aligned in MAFFT (Nakamura and others 2018) and the alignment was used

to construct phylogenetic trees in MEGA7 (Kumar and others 2016) using a maximum likelihood

approach. Bootstrap values from 100 replicates are shown for nodes with more than 70%

bootstrap support.

lone e .1)	ITS1: MK775315	colonies (2-3 mm) with a compact and fluffy center surrounded by radiating rhizoids; in broth, numerous fungal thalli attach to the glass bottles initially, and later developed into thin mat-like structures (Fig. 1e)	flagellate, globose; size, 4–5 μm diameter, flagellum length, 15-20 μm (Fig 6a)	filamentous (Fig. 6b-c). Sporangia: endogenous (Fig. 6d-g) & exogenous (Fig. 6h-l)	sporangiophore in case of exogenous development; multisporangiate thallus with two sporangia of same or different shapes
o lone na 3.1)	LSU: MK881971, ITS1: MK882043-MK882046	On agar: White circular colonies with a brown central sporangial core (1-4 mm diam.); In liquid: Thick fungal biofilm growth (Fig. 1a)	Polyflagellated, globose, 6-10.5μm in dimeter, 7-14 flagella, flagellum length:16-31μm (Fig. 2a)	Monocentric/ filamentous (Fig. 2b-c). Sporangia: endogenous (Fig. 2d-g) & exogenous (Fig. 2h-p)	Persistent zoospore cyst, Sporangia with tightly constricted necks, narrow ports and fine septum at the base
lone e 4.1)		On agar: small (1-2 mm) colonies with a dense dark central core of abundant sporangial growth, surrounded by long and thin radiating rhizoids; in broth, numerous fungal thalli attach to the glass bottles initially, an later develop into thin mat-like structures (Fig. 1g)	Mono- or bi-flagellate; globose; 5–6 μm diameter; flagellum length, 20-22 μm (Fig. 8a-b)	Monocentric/ filamentous (Fig. 8c-f). Sporangia: endogenous (Fig. 8g) & exogenous (Fig. 8h-i)	Zoospores are discharged after dissolution of a wide apical portion of sporangial wall, resulting in formation of empty cup-shaped sporangia
lone elle .1)/ lade	LSU: MK881981, ITS1: MK882019	On agar:Yellow to yellowish-brown of irregular shapes; In liquid: Loose fungal thalli with sand-like appearance (Fig. 1f).	Uniflagellated, spherical, 6-17 μm in diameter, flagellum length:18-40μm (Fig.7a)	Monocentric/ filamentous (Fig. 7d-e). Sporangia: endogenous (Fig. 7g-h) & exogenous (Fig. 7i-m)	Multisporangiated thalli & branched sporangiophores
one)	LSU: MK775310, ITS1: MK775321	On agar: small (1 mm) white colonies, with a compact and fluffy center, surrounded by dotted circles of fungal thalli; in broth, numerous fungal thalli attach to the glass bottles initially, and later develop	Mono-, bi-or tri-flagellate; globose; 3–4 μm diameter; flagellum length, 12-15 μm (Fig 9a)	Monocentric/ filamentous (Fig. 9b-e). Sporangia: endogenous (Fig.9f-g) & exogenous (Fig. 9h-l)	Distinct subsporangial swelling with or without constricted neck on a short sporangiophore

(n)

(o)

(p)

(q)

(r)

A

- 83
- 93
- Orpinomyces
- 97 — *Ghazallomyces* (Clade 4, Axis Deer)
- 100 — Pecoramyces
- 98 — Neocallimastix
- 99
- 96 — Feramyces
- 85
- 100 — Caecomyces
- 88
- Cyllamyces
- 99
- Piromyces
- 91
- 86 — *Joblinomyces* (Clade 5, domesticated Goat-Sheep)
- 80 — Buchfawromyces
- 89 — *Tahromyces* (Clade 7, Tahr)
- 100
- 96 — *Khyollomyces* (Clade 6, Zebra-Horse)
- 92
- 100 — *Agriosomyces* (Clade 1, Mouflon-Boer Goat)
- 100 — Liebetanzomyces
- 89
- 75 — *Capellomyces foraminis*
- 90
- *Capellomyces elongatus*   *Capellomyces* (Clade 3, Boer Goat-domesticated Goat)
- 100 — Oontomyces
- Anaeromyces
- 93
- *Aklioshbomyces* (Clade 2, White-Tail Deer)
- 89 — DQ536493.1 *Chytriomyces* sp. WB235A isolate AFTOL-ID 1536

B

- 74 — *Aklioshbomyces* (Clade 2)
- 80
- 100 — Orpinomyces
- 83 — Pecoramyces
- 95 — *Ghazallomyces* (Clade 4)
- 99 — *Agriosomyces* (Clade 1)
- 100 — Feramyces
- 99 — Neocallimastix
- Neocallimastix
- 73
- 94 — *Joblinomyces* (Clade 5)
- 95
- 100 — Cyllamyces
- 77
- 82 — Caecomyces
- 81 — Piromyces
- 98 — Liebetanzomyces
- 79
- 91 — NR 148182.1 *Anaeromyces robustus* JMRC SF:12178
- KU057354.1 *Anaeromyces robustus*
- 97 — *Capellamyces elongatus* isolate GFKJa1916 (MK775315)
- *Capellamyces foraminis* isolate BGB11- clone C3 (MK882007)
- *Capellamyces foraminis* isolate BGB11- clone C1 (MK882009)   *Capellomyces* (Clade 3)
- 84 — *Capellamyces foraminis* isolate BGB11- clone C4 (MK882008)
- FJ912851.1 *Anaeromyces* sp. GA 04
- 95 — Anaeromyces
- 100 — Oontomyces
- 74
- 100 — *Tahromyces* (Clade 7)
- 99 — Buchfawromyces
- 98 — *Khyollomyces* (Clade 6)
- AY349118.1 *Chytriomyces* sp. JEL176