Signatures of selection in a recent invasion reveals adaptive divergence in a highly vagile

invasive species

Adam P. A. Cardilini[1†], Katarina C. Stuart[2†], Phillip Cassey[3], Mark F. Richardson[4,5], William

Sherwin[2], Lee A. Rollins[2,4*], Craig D.H. Sherman[4*]

[1]Deakin University, School of Life and Environmental Sciences, Waurn Ponds, VIC, 3216,

Australia

[2]Evolution & Ecology Research Centre, UNSW Sydney, Sydney NSW 2051 AUSTRALIA

[3]Centre for Applied Conservation Science, and School of Biological Sciences, University of

Adelaide, SA 5005, Australia

[4]Deakin University, Centre for Integrative Ecology, School of Life and Environmental

Sciences, Waurn Ponds, VIC, 3216, Australia

[5]Deakin University, Genomics Centre, School of Life and Environmental Sciences, Waurn

Ponds, VIC, 3216, Australia

[†]Joint first Authors

*Joint last authors

Corresponding Authors: Craig Sherman (craig.sherman@deakin.edu.au); Lee Ann Rollins

(l.rollins@unsw.edu.au)

Abstract

A detailed understanding of population genetics in invasive populations helps us to identify drivers of successful introductions. Here, we investigate putative signals of selection in Australian populations of invasive common starlings, *Sturnus vulgaris*, and seek to understand how these have been influenced by introduction history. We use reduced representation sequencing to determine population structure, and identity Single Nucleotide Polymorphisms (SNPs) that are putatively under selection. We found that since their introduction into Australia, starling populations have become genetically differentiated despite the potential for high levels of dispersal, and that selection has facilitated their adaptation to the wide range of environmental conditions across their geographic range. Isolation by distance appears to have played a strong role in determining genetic substructure across the starling's Australian range. Analyses of candidate SNPs that are putatively under selection indicate that aridity, precipitation, and temperature may be important factors driving adaptive variation across the starling's invasive range in Australia. However, we also note that the historic introduction regime may leave footprints on sites flagged as being under adaptive selection, and encourage critical interpretation of selection analyses.

Keywords: *Sturnus vulgaris*, common starling, invasive species, local adaptation, population genetics

Introduction

Understanding the proximate molecular mechanisms of natural selection is a central goal in evolutionary biology. Invasive species have long presented an opportunity to elucidate evolutionary mechanisms underlying selection (Baker and Stebbins 1965). Rapid adaptation is commonplace in invasive populations (Lee 2002; Rollins, et al. 2013) and is proposed to be important to their long-term success (Dlugosch and Parker 2008). The short evolutionary timescale over which this adaptation must take place provides opportunity to characterise the ecological and evolutionary processes that contribute to successful establishment and spread (Lee 2002). Understanding these proximate mechanisms enables us to predict how evolutionary adaptation will influence range distribution, or how populations (both native and invasive) may respond to environmental change (Bay, et al. 2018).

Investigating the genetics underlying evolutionary change in invasive populations gives us the opportunity to understand how a range of demographic and adaptive processes shape populations (e.g. bottlenecks, population expansions and adaptation) (reviewed in Sherman, et al. 2016). This information helps us understand invasion dynamics (Tepolt, et al. 2009) and may illuminate the adaptive response of species to a novel or changing environment (Riquet, et al. 2013). Using these data, we can identify signatures of molecular evolution in alien species (Ashburner, et al. 2000; Franks and Munshi-South 2014) by identifying loci putatively under selection when considering population genetic statistics such as F_{ST}, or loci that are significantly associated with environmental variables (Foll and Gaggiotti 2008; Gunther and Coop 2013; Loh, et al. 2015).

Human introduced populations often have discrete (but potentially numerous and repetitious) geographic introductory sites, which will have variable numbers of founding individuals from potentially different source populations (Dlugosch and Parker 2008). What

may then result is an amalgamation of subpopulations that, depending on the age of the

populations and degree of connectivity, have the potential to be genetically differentiated

geographically due to legacy genotypes (genotypes of founding individuals). Further,

bottlenecks encourage stochastic evolution through processes such as genetic drift. During

rapid range expansion, allelic surfing can create signatures at neutral loci that resemble

selection, particularly at range edges (Klopfstein, et al. 2006; Lotterhos and Whitlock 2015;

Hoban, et al. 2016; reviewed in Sherman et al. 2016). Until these discrete introductions make

contact, it is possible that these processes are occurring concurrently and independently at

different introduction sites and may leave genetic footprints. It is important to understand to

what extent historic invasion regime influences patterns of adaptive genetic variation in

invasive populations, particularly those with multiple and spatially separate introductions.

The common starling (*Sturnus vulgaris*) invasion in Australia represents an excellent

system in which to assess the interaction between adaptive selection and historic

introduction regime in shaping population structure. Starlings are a highly successful invasive

species that has established populations around the world (Higgins, et al. 2006). Starlings

were introduced into Australia from England during the 1850's onwards at several locations,

including Melbourne (Victoria, VIC), Brisbane (Queensland, QLD), Adelaide (South Australia,

SA) and Hobart (Tasmania, TAS) (Fig. 1a), with the total individuals introduced numbering in

the hundreds (Jenkins 1977; Higgins, et al. 2006). They quickly spread from their introduction

sites on the south-eastern coastline to occupy a large range in south-eastern Australia (Long

1981, Rollins, et al. 2009; Rollins, et al. 2011), across habitats with significant environmental

gradients (Woolnough, et al. 2005). They have now persisted in Australia for approximately

60 generations since their introduction (Higgins, et al. 2006; Rollins, et al. 2016).

Starlings are predominantly found in agricultural landscapes relying on open fields to

forage (Feare 1985; Whitehead, et al. 1995). The starling native range covers a wide range

of environments spanning the Palearctic, including temperate forests and grasslands, as well

as both hotter and colder climates at their native range edges (Higgins, et al. 2006). Starlings

are a highly dispersive species, able to disperse upwards of 1000km from their natal site

(Waterman, et al. 2008). Nevertheless, across their Australian range, starlings have two

possible major barriers to movement. The hot, arid areas of Australia present a challenge to

the species; central Australia is unlikely to be colonised by starlings and the large desert

expanse of the Nullarbor Plain (between SA and WA) is known to restrict the species'

movement (Woolnough, et al. 2005). Second, large contiguous areas of forest and elevated

mountain ranges along the Eastern coast of the continent are likely to act as a barrier to

starling movement, restricting gene flow (Fig. 1b). High elevation is suspected to play an

important role in shaping starling demography within starling invasive populations (Ross

1983), because the species has an upper elevation limit of 2000m (Higgins, et al. 2006).

These biogeographical barriers may therefore play a major role in determining patterns of

gene flow across Australia.

Interestingly, while starlings in the native Palearctic and invasive North American

ranges display migratory behaviour, this has been lost in Australian starlings (Cabe 1999,

Rollins, *et al.* 2009). While starlings may move in excess of 100's of kms from their natal site

(Feare 1984), a lack of largescale Australia-wide interbreeding is evident in their underlying

genetic population structure. Previous genetic surveys based on microsatellite markers

showed that starlings in Australia form four distinct genetic clusters (Rollins, et al. 2009). The

genetic cluster representing the range-edge at the western most extent of the starlings' range

in Western Australia (WA) contained novel genetic variants relative to range-core and native

individuals, and evidence of genetic drift and selection was identified in this cluster (Rollins,

et al. 2016). This stands in contrast to starling populations of North America, also introduced

in the late 19[th] Century, for which high dispersal and continental wide interbreeding appears

to have led to genetic homogenisation, minimal population substructure, and lower F_{ST}

values between sites, when compared to the Australian starlings (Cabe 1998, 1999). The

Australian invasion is characterised by lower genetic diversity than that of the native range

(Rollins, et al. 2011). Nevertheless, there is clear evidence of morphological variation in

response to temperature and rainfall gradients across the invasive range in Australia

(Cardilini, et al. 2016).

Given the wide range of environments that Australian starlings inhabit and the

approximately 60 generations since introduction, the invasive starling provides an excellent

model for investigating the population dynamics and adaptive variation of a highly dispersive

invasive species. Further, the multiple Australian introductions enables us to examine

whether introduction regime may leave genetic footprints in modern day populations. Here,

we use a reduced representation sequencing approach to generate a SNP data set to

assess the roles the Australian landscape and starlings' invasion history have played in

shaping the genetic structure and patterns of adaptive genetic variation found across the

Australian range. We aim to test two non-mutually exclusive hypothesis: that genetic

differentiation will be related to geographic distance, and that genetic differentiation will be

related to environment gradients. To do this, we compare population structure to underlying

environmental differences and geographic distances separating sites. Finally, we then use

three different analyses to test for adaptive genetic variation across the starlings geographic

range in Australia.

Methods

Sample collection and sexing

We collected a total of 568 starling samples from 24 localities across the invasive range in Australia (Fig. 1b, Table 1). Sample collection occurred in two separate phases. First, we used a subset of pre-extracted DNA samples (n = 150) collected by Rollins *et al.* (2009), from localities across South Australia and Western Australia from 2003 – 2007. Second, we collected 418 samples between May 2011 and October 2012 from 18 locations across the south-eastern portion of the Australian range (Fig. 1b). These birds were collected in three ways: *i)* by trapping, *ii)* as carcasses collected from 3[rd] party hunters or landowners that humanely killed birds on their property, or *iii)* chicks collected directly from nests. In the latter case, only one individual per family (adult or chick) was included in the dataset to avoid the inclusion of closely related individuals. We recorded Global Position System (GPS) coordinates upon collection. When 3[rd] party hunters did not record GPS coordinates, the closest corresponding address was recorded and GPS coordinates were identified using Google maps. We took a muscle sample from the thigh or breast and stored it in 70% ethanol prior to DNA extraction. We extracted DNA with a Gentra PureGene Tissue Kit, following the manufacturer's protocol. All samples were sexed using molecular methods (Griffiths, et al. 1998).

Fig. 1: A) Australian starling distribution map according to eBird data (accessed Nov 2018),

with introduction sites marked and first major introduction date listed adjacently, and current

known range edge marked in red. B) Map of southeastern Australia. Points denote starling

collection localities. Point shape indicates the environmental cluster that the point belonged

to, as identified by a PCA analysis of environmental variables; circle = arid, diamond = semi-

arid, square = non-arid. Point colour indicates the genetic cluster that the population

belonged to as identified by population Bayesian fastSTRUCTURE analysis; pink =

Munglinup cluster, yellow = southern Australia cluster, orange = southwest New South Wales

cluster, and blue = northern Australia cluster. The blue line represents The Great Dividing

Range (continental divide). Sample site number key corresponds to that listed in Table 1.

Environmental data collection

For each collection locality, we extracted climatic variables from Bioclim data sets using the

RASTER package in R (Hijmans 2016). Climatic variables included: mean diurnal range

(Bio02), isothermality (seasonal temperature variation) (Bio03), temperature seasonality

(Bio04), maximum temperature of the warmest month (Bio05), minimum temperature of the

coldest month (Bio06), temperature annual range (Bio07), precipitation of the wettest month

(Bio13), precipitation of the driest month (Bio14) and precipitation seasonality (Bio15). We

extracted the aridity index (AI) data from the CGIAR-CSI Global-Aridity and Global-PET

Geospatial Database (Zorner, et al. 2008). We downloaded monthly average normalised

difference vegetation index (NDVI) values from the Australian Bureau of Meteorology for the

period 1997-2011. We averaged monthly datasets across years (e.g. mean NDVI January

1997-2011) and then across all months to create a mean NDVI raster file from which we

extracted mean NDVI (mnNDVI) for each sample. We calculated variability in day length

(varDL) per sample using the 'daylength()' function in the R package GEOSPHERE (Hijmans,

et al. 2015). We defined the position of each collection locality as the central longitude and

latitude of all samples collected from the same collection locality. Using an R function that we

wrote, we extracted elevation data (Elev) for each sample collection locality from Google

maps (see supplementary file for function). We calculated the average of each environmental

variable at each collection locality and used these for further analysis. Using the R package

SP (Bivand, et al. 2013), we calculated geographic distance variables as Euclidean distance

(km) of each locality to the nearest coastline and Euclidean distance of each locality to

nearest known site of introduction. To identify a reduced set of variables that explained

environmental variation across the introduced range, we conducted a principal components

analysis (PCA) including all environmental variables and run in R using the PRCOMP function

(RCoreTeam 2015). We calculated the pairwise distance between points on a plot of PC1

and PC2 to calculate pairwise environmental distances between collection localities, which

were subsequently used in the analysis of isolation by environment (IBE matrix) (Wang 2013;

Wang, et al. 2013; Leydet, et al. 2018). Finally, using the R package SP (Bivand, et al. 2013)

we calculated pairwise distances as the greater circle distances between collection localities

to test for isolation by distance (IBD matrix).

These environmental data were used to examine IBE, IBD, and the relationship

between geographic and environmental distances across sampling sites (see "Genetic,

environmental and geographic relationships"), and in the identification of candidate loci under

selection (see "Detecting putative loci under selection").

208 Library construction, sequencing and SNP calling

The Cornell University Institute of Biotechnology Genomics Facility conducted library

construction and Genotyping-by-Sequencing (GBS) (Elshire, et al. 2011), using the

restriction enzyme *PstI*. Each individual received a unique barcode before multiplexing 96

individuals per lane (Illumina HiSeq 2000; 6 lanes, 100 bp single-end reads).

We processed raw sequence data using TASSEL 3.0 and called SNPs for all

individuals using the UNEAK pipeline (Bradbury, et al. 2007; Lu, et al. 2013; Glaubitz, et al.

2014). We used default values for parameters and sequence read (tags) that were detected

at least five times across all samples were retained. We compared tags (aligned 64 bp

reads) using network filtering (identifies tag pairs by filtering out repeats, sequencing errors,

and paralogs) and considered those with a 1 bp mismatch to be candidate SNPs, using an

error tolerance rate of 0.03 (Lu, et al. 2013). After these initial filtering steps, 6,102,279 reads

remained and we combined and converted into Tags on Physical Map format (creating a

pseudo chromosome), which acted as a reference. SNPs were called for each individual and

we identified a total of 291,189 SNP loci. When calling SNPs against the reference pseudo-

chromosome, we only considered tags recorded in at least three individuals. We used a

maximum genotype depth of 60 (~10X the average genotype depth of the dataset) for all

data sets to filter out genotypes that were possibly overrepresented (due to being in

repetitive genomic regions) (Almeida, et al. 2014). We removed any locus that was missing

in more than 90% of samples and, subsequently, removed all samples that had less than

10% of the total number of loci. We tested our data for conformity to Hardy Weinberg

Equilibrium (HWE) (VCFTOOLS, v0.1.12b) (Danecek, et al. 2011) and, after correcting *p*-

values for multiple comparisons using FDR 0.05 (Benjamini and Hochberg 1995), only loci

identified out of HWE in five or more collection localities were to be removed; no loci were

found to be out of HWE in five or more collection localities.

We examined a range of filtering metrics (detailed methods in Supplementary

Material, Fig. S1) and chose a final dataset with a minor allele frequency (MAF) of 0.05, a

minimum genotype depth of three and maximum level of missingness of 50%, resulting in the

inclusion of 16,177 SNP loci across 499 individuals (data accessible at

https://github.com/katarinastuart/Sv1_StarlingGBS/tree/master/Data). We chose this dataset

because including tags with large proportions of missing data can obscure signal, while

stringent missingness filtration by definition reduces diversity, leading to an underestimation

of population structure (Huang and Knowles 2016). Regarding MAF, higher thresholds are

proven to inhibit population substructure detection, a problem that also persists when

singletons are retained (i.e. low MAF threshold) (Linck and Battey 2019). Hereafter this SNP

dataset will be called the genome-wide dataset.

Population structure analysis

Population structure based on genome-wide SNP loci

Global F_{ST} and pairwise F_{ST} values were estimated in the R package HIERFSTAT (Goudet

2005), as were observed heterozygosity [H_O] and within population gene diversity [H_S].

Upper and lower confidence intervals (95%) were obtained for F_{ST} from 10,000 bootstraps, *p*-

values were corrected for multiple comparisons using FDR 0.05 (Benjamini and Hochberg

1995). Collection locality F_{ST} (F_{ST} of one population against all other populations) and

Tajima's D (Tajima 1989) values within all collection localities were calculated in the R

package POPGENOME (Pfeifer, et al. 2014).

We determined the number of distinct genetic groups in Australia using a Bayesian

assignment test in the program FASTSTRUCTURE (v1.0) (Raj, et al. 2014). Additionally, we

analysed population structure of a neutral data set, to ensure it was reasonable to use the

genome-wide dataset without removing SNPs contributing to non-neutral variation (detailed

methods in Supplementary Material).

Population structure based on outlier SNP loci

We identified F_{ST} outlier loci in BAYESCAN (v2.1) (Fischer, et al. 2011), which uses a Bayesian

approach to compare the F_{ST} values of a locus across subpopulations (Foll and Gaggiotti

2008). BAYESCAN was run for two different groupings of the samples. On the first run, we

allocated individuals to one of 24 subpopulations that related to their collection localities.

BAYESCAN is prone to high levels of false positives; however, Lotterhos and Whitlock (2014)

showed that this effect can be reduced by increasing the prior odds value. BAYESCAN was

run with default parameters except that the prior odds value was set to 100. Only loci with a

Bayes factor greater than three were kept as outliers because values above this indicate

substantial evidence of selection (Jeffreys 1961). We determined the number of distinct

genetic groups determined by the outlier loci dataset identified with BayeScan using

FASTSTRUCTURE (detailed methods in Supplementary Material). Additionally, to validate the

outlier loci data set, we generated a random subset of an equivalent number of SNPs from

the genome-wide dataset and analysed these as above (detailed methods in Supplementary

Material).

Genetic, environmental and geographic relationships

We used Multiple Matrix Regression with Randomization analysis (MMRR) (Dryad

doi:10.5061/dryad.kt71r) to determine if geographic distance between localities predicts

environmental distance. Then, we used MMRR analysis to determine if geographic distance

between localities (IBD matrix) predicts genetic distance (isolation by distance, IBD), or if

environmental distance (IBE matrix) predicts genetic distance (isolation by environment;

IBE). MMRR was implemented by comparing a pairwise genetic distance matrix with IBD and

IBE distance matrices following Wang (2013). Briefly, random permutations are conducted

between the rows and columns of the dependant (genetic distance) matrix to produce the

resulting matrix and its coefficients (while other matrices remain fixed) - a null distribution is

formed from this permutation, and observed values may be compared to it to determine if

there is any relation between matrix distances (Legendre, et al. 1994).

Detecting putative loci under selection

We investigated adaptive variation across the starling's range using candidate loci identified

by BAYSCAN (F_{ST} outlier loci) as outlined above, and through two different methods of

environmental association analyses (BAYENV2 and RDA). Bayescan was run with as above.

While measures of population differentiation such as F_{ST} allow for SNP's putatively under

selection to be identified, measured differentiation may simply be a result of neutral variation

(demographic effects, drift etc.), and such an approach does not directly allow possible

selective drivers to be identified (e.g. environment) (Feng, et al. 2015). Hence it is common

for scans for selection to also include environmental association analysis (Pluess, et al.

2016), such as logistic regression. The nature of adaptive drivers though means a diverse

array of factors may be subtly influencing genomic sites, so many regressions may need to

be performed if there are a number of environmental drivers of interest. To address this,

constrained ordination methods are fast becoming a popular method when conducting

genotype-environment associations with multivariate and multilocus datasets, boasting

detection power coupled with low false-positive and high true-positive rates (Forester, et al.

2018), and therefore are invaluable for investigating adaptation (Capblancq, et al. 2018).

Under the conditions of genotype-environment associations, constrained ordination

techniques such as RDA represent parsimonious models that attempt to capture total

population variation specifically explained by constraining predictor (e.g. environmental)

variables. Thus, RDA environmental association analysis was conducted alongside the

traditional allele frequency association tests. However, RDA cannot be conducted on

matrices with missing data, so imputation on missing values is necessary – a particular

problem for dataset like GBS that are characterised by high levels of missingness. It is

possible that imputation might lead to type I and type II errors, hence we used this method

alongside the two more traditional analyses (BAYSCAN and BAYENV2) to test for signals of

selection.

We used BAYENV2 (v2.0; Gunther and Coop 2013) to test for association of minor

allele frequencies with environmental variables. Bayenv2 calculates the correlation between

allele frequencies and environmental variables and was used to tests for associations

between all SNPs and three environmental variables (Coop, et al. 2010; Gunther and Coop

2013). The three environmental variables that were tested included Aridity, Bio05 (maximum

temperature of the warmest month), and Bio 15 (precipitation seasonality). These variables

were chosen because temperature and precipitation have previously been shown to

influence starling phenotype (Cardilini, et al. 2016). We estimated the covariate matrix using

a random subset of 2,000 loci. Runs were set at 100,000 iterations and we kept any locus

with a Bayes factor greater than three as being potentially under selection (Gunther and

Coop 2018). From the candidate SNPs identified by BAYENV2, we extracted the SNP that

most highly correlated with each of Aridity, Bio05, and Bio 15, and plotted allele frequency

against the environmental variables score.

We used the R package VEGAN (v3) (Oksanen, et al. 2018) to conduct redundancy

analysis (RDA) to determine if any particular SNP locus is heavily loaded onto predictor

axes, from which we may infer the occurrence of selection. The genome-wide SNP dataset

was loaded against the climatic variables associated with our sampling sites. The SNP

genotype dataset was processed into 012 format (count reflects the number of the minor

allele) using VCFtools, and missing SNP genotypes (average missingness across all sites:

44.3%) were imputed based on probabilities for each given genotype at that genome locality

using a custom script (detailed methods in Supplementary Material). To reduce

multicollinearity, from the complete list of environmental variables, we retained seven

predictors with relatively low variance inflation factors (range: 1.9—6.3) to reduce

multicollinearity: (Elev, Bio03 (isothermality); Bio06 (minimum temperature of the coldest

month); Bio13 (precipitation of the wettest month); Bio15 (precipitation seasonality); mnNDVI;

and varDL. We used the function ANOVA.CCA in VEGAN to confirm the model's significance.

Once the SNP data were loaded against the RDA axes, candidates for selection were

determined to be those that lay more than three standard deviations away from the mean

(Forester, et al. 2018).

We extracted the associated 64 bp tags for loci identified by at least one of the

methods of BAYESCAN, BAYENV2, and RDA. These tags were searched against the annotated

starling genome available on NCBI (GCF_001447265.1) and functional descriptions

extracted, following the protocol described in Richardson and Sherman (Richardson and

Sherman 2015). Briefly, 64 bp tags of outlier loci were searched against the starling

predicted RNAs using Blastn with a e-value cutoff of 1.0×10^{-3} (Camacho, et al. 2009). Gene

names were extracted from the blast output for the best hit matches per outlier loci (smallest

e-value, highest % identity and length match) and these gene names were matched to

Uniprot IDs and associated functional descriptions extracted (Bairoch, et al. 2008).

Results

Population environmental distribution

A PCA analysis of environmental parameters grouped collection localities into three distinct

clusters (Fig. 1b; Supplementary Material, Fig. S2, Table S1). These clusters defined three

distinct environmental regions: (i) Western Australian and South Australian collection

localities (arid, blue); (ii) eastern Australia collection localities inland of The Great Dividing

Range (semi-arid, red); and (iii) collection localities on the coastal side of The Great Dividing

Range (non-arid, green) (Supplementary Material, Table. S1).

Population structure analysis

Population structure based on genome-wide SNP loci

Analysis of the patterns of genetic diversity revealed that populations at the expansion front

in Western Australia (Munglinup and Condingup) displayed relatively lower levels of genetic

diversity compared to populations in the east of the introduced range (Table 1). Further,

Munglinup was the only population to have a positive Tajima's D (low frequency of rare

alleles). The global level of genetic differentiation across collection localities was low, but

significant with an average F_{ST} of 0.027 (95% CI 0.025 – 0.030) (Supplementary Material,

Table S2). Pairwise comparisons of F_{ST} between collection locations revealed significant

genetic structuring between 263 out of 276 pairwise comparisons (Supplementary Material,

Table S2). Collection locality (location vs all others) pairwise F_{ST} values revealed that those

populations with the lowest levels of genetic diversity also showed the greatest level of

genetic differentiation to all other populations (Table 1).

The Bayesian structure analysis based on the full dataset of 16,177 SNPs showed

that K=2, and K=4 had similar levels of support (Supplementary Material, Fig. S3a), while

K=4 had the most support when considering only neutral genome-wide loci (Supplementary

Material, Fig. S3a). For the former, the four genetic clusters were Munglinup, southern

Australia, southwest New South Wales, and northern Australia (Fig. 1b, Fig. 2). Within the

FASTSTRUCTURE plot, the Munglinup and southwest New South Wales clusters were

relatively distinct though admixture is still present, while the two larger genetic clusters were

less well defined suggesting more admixture (Fig 3). Subsequent FASTSTRUCTURE analysis

found no substructure within any of the four genetic clusters, and analysis on the neutral

dataset produced a very similar likelihood plot (Supplementary Material, Fig. S3b).

Population structure based on outlier SNP loci

Out of the 16,177 SNPs, BAYESCAN analyses identified 89 outlier loci (Table 2) as candidates

for selection (log10(BF) > 3), while all other loci were considered neutral. These loci made up

the outlier data set. As expected, pairwise F_{ST} values calculated for outlier loci were

considerably higher than those calculated for the genome-wide dataset, with an average F_{ST}

of 0.122 (95% CI 0.101 – 0.130, non-overlapping with the CI for genome-wide SNP loci

calculated F_{ST}) and most locality F_{ST} comparisons for the outlier dataset were significant

(256/276; Supplementary Material, Table S2). FASTSTRUCTURE analyses of the outlier

dataset identified two distinct genetic clusters which covered the larger clusters from the

genome wide SNP dataset, with the range edge cluster integrated into the eastern cluster,

and the inner NSW cluster split amongst the two new large outlier clusters (Fig. 2;

Supplementary Material, Fig. S3c), whereas we found no genetic structure in the random

subset of loci (Supplementary Material, Fig. S3d). Subsequent FASTSTRUCTURE analysis

found no substructure in the two main genetic clusters (data not presented).

Fig. 2. fastSTRUCTURE plots indicating the genetic structure of Australian starling populations. Each column represents an individual sample, with the colours indicating the estimated likelihood that the individual belongs to a particular genetic group. Collection localities are arranged from west to northeast following geographic distribution. The vertical black lines demarcate state political boundaries indicated in the abbreviations above each plot, and the vertical grey lines delimit collection localities. The black bars at the top of the graph and their corresponding names identify genetic clusters. Each panel is derived from a different dataset with a) indicating genetic structure of the genome-wide dataset when K = 4 and b) indicating the genetic structure of the outlier dataset when K = 2. Labels indicate the distinct genetic clusters reported by the fastSTRUCTURE analysis for each panel; panel a) Munglinup (A), southern Australia (B), southwest New South Wales (C), northern Australia (D); panel b) southern Australia (A), northern Australia (B). Symbols and colour used at the bottom of the figure are based on Fig. 1b.

Table 1. Summary statistics for each starling collection locality. N = number of samples collected from each location, GC = the genetic cluster from whole-genome dataset (1 = Munglinup cluster, 2 = southern Australia cluster, 3 = southwest New South Wales cluster, 4 = northern Australia cluster), OGC = the genetic cluster from outlier dataset (1 = Western and Southern Australia, 2 = Eastern and Northern Australia), H_O = Observed Heterozygosity, H_S = the within population gene diversity, Locality F_{ST} = fixation index for each population against all other individuals, % Poly SNPs = the percentage of polymorphic SNPs. We do not report a Tajima's D value for Streaky Bay due to low sample size.

Collection Locality	Sample site number	State	N	GC	OGC	Env. Cluster	H_O	H_S	Locality F_{ST}	% Poly. SNPs	Tajima's D
Munglinup	1	WA	18	1	1	1	0.092	0.225	0.094	55.2	0.137
Condingup	2	WA	16	2	1	1	0.092	0.257	0.051	66.1	-0.129
Coorabie	3	SA	19	2	1	1	0.137	0.265	0.028	80.9	-0.395
Streaky Bay	4	SA	9	2	1	1	0.069	0.267	0.137	35.9	NA
Tumby Bay	5	SA	22	2	1	1	0.114	0.269	0.030	83.3	-0.134
McLarenvale	6	SA	20	2	1	1	0.141	0.266	0.032	86.0	-0.423
Warrnambool	7	VIC	24	2	1	3	0.117	0.271	0.026	88.6	-0.286
Bendigo	8	VIC	25	2	1	2	0.159	0.273	0.027	93.4	-0.449
Wonthaggi	9	VIC	22	2	1	3	0.119	0.272	0.030	81.6	-0.292
Orbost	10	VIC	22	2	1	3	0.148	0.270	0.028	89.0	-0.468
King Island	11	TAS	14	2	1	3	0.135	0.272	0.027	80.9	-0.395
Sheffield	12	TAS	21	2	1	3	0.098	0.265	0.037	79.2	-0.106
Cygnet	13	TAS	17	2	1	3	0.095	0.262	0.037	72.5	-0.190
Hay	14	NSW	26	3	1	2	0.158	0.259	0.025	86.2	-0.581
Albury	15	NSW	24	3	1	2	0.200	0.259	0.038	89.5	-0.899
Nowra	16	NSW	25	4	2	3	0.133	0.268	0.030	89.2	-0.278
Maitland	17	NSW	25	4	2	3	0.149	0.266	0.034	91.7	-0.391
Dubbo	18	NSW	20	3	2	2	0.165	0.257	0.030	82.9	-0.671
Nyngan	19	NSW	25	4	2	2	0.126	0.273	0.023	84.7	-0.391
Austral Eden	20	NSW	21	4	2	3	0.142	0.273	0.024	87.0	-0.349
Tamworth	21	NSW	11	4	2	2	0.129	0.271	0.029	64.6	-1.409
Moree	22	NSW	24	4	2	2	0.166	0.271	0.028	91.6	-0.546
Lismore	23	NSW	22	4	2	3	0.123	0.270	0.024	86.9	-0.393
Lemon Tree	24	QLD	24	4	2	2	0.141	0.272	0.024	89.0	-0.418

424 **Table 2**. Summary of candidate SNPs under putative selection, as detected by Fst outlier (Bayescan) and environmental association
425 (Bayenv2 and RDA) approaches. Bayenv2 analysis is broken down into the three tested environmental variables Aridity, Bio05 (maximum
426 temperature of the warmest month), and Bio 15 (precipitation seasonality). RDA is broken down into three key RDA axes. Displayed are
427 number of unique SNPs identified for each method (diagonal), total number of SNPs identified for each method (bottom line) and SNPs
428 common to each pairwise comparison of approaches (bottom matrix).

Approach	Bayescan	Aridity	Bio05	Bio15	Axis 1	Axis 2	Axis 3
Bayescan	69						
Bayenv2 (Aridity)	2	23					
Bayenv2 (Bio05)	8	0	156				
Bayenv2 (Bio15)	1	0	0	40			
RDA (Axis 1)	8	0	1	1	23		
RDA (Axis 2)	1	0	0	2	0	24	
RDA (Axis 3)	0	3	9	0	0	0	40
Total: 445 (375 unique)	**89**	**28**	**174**	**43**	**32**	**27**	**52**

Genetic, environmental and geographic relationships

Geographic distance did significantly correlate with environmental distance across the

starling's Australian range (Fig. 3), so IBE and IBD were calculated in the same MMRR

model to account for confounding measures of geographic distance and environmental

distance. When genetic distance was tested against geographic distance and environmental

distance in an MMRR, only IBD was found to explain any variation in genetic differentiation

observed between collection localities (Fig. 4a). IBE was not found to be a significant

predictor of genetic distance (Fig. 3). When the genetic distance of the outlier dataset was

tested against geographic distance and environmental distance in an MMRR, only IBD was

found to explain any variation between collection localities (Fig. 4b). IBE was not found to be

a significant predictor of genetic distance (Fig. 3).

Fig. 3. Results of Multiple Matrix Regression with Randomization analysis (MMRR) capturing genetic, environmental and geographic relationships in Australian starlings. Analysis results on the outside of the triangle were performed on the genome-wide dataset ($r^2 = 0.229$), while analysis results listed inside the triangle and with an asterix (*) indicates analysis that were performed on outlier loci data set ($r^2 = 0.396$). Analysis results of geographic distance and environmental distance is not specific to a genetic data set ($r^2 = 0.117$). Results coloured red indicate significant results.

Detecting putative loci under selection

Using three different methods to identify SNPs putatively under selection, we found a total of 375 different SNPs. BAYESCAN analyses of F_{ST} identified 89 outlier loci (Table 2). BAYENV2 identified positive associations between 245 SNPs and three environmental variables including Aridity (N = 28 SNPs), Bio05 (temperature) (N = 174 SNPs), and Bio15 (precipitation) (N = 43 SNPs) (Table 2). The minor allele frequency of the locus most strongly associated with Aridity (ID = 68063014) showed a negative relationship (C = -0.272, SE = 0.070, t = -3.883, P < 0.001, r^2 = 0.380; Fig. 5a). The minor allele frequencies of the SNP (ID = 233190026) most strongly associated with Bio05 showed a strong positive relationship (C = 0.018, SE = 0.004, t = 4.116, P < 0.001, r^2 = 0.409; Fig. 5b). The relationship between Bio05 and the minor allele frequency for this SNP is closely aligned with geography, with the Munglinup cluster and almost all collection localities from the southern Australia cluster being fixed at this locus. Loci associated with Bio15 (ID = 196163030) showed a positive relationship (C = 0.008, SE = 0.002, t = 3.607, P = 0.002, r^2 = 0.343; Fig. 5c), with most individuals from the northern Australia cluster being fixed at this locus.

Fig. 4. Plots of isolation by distance (IBD) in Australian starlings with Mantel correlation and p-value indicated. Each point represents a pairwise relationship between collection localities and the line represents the linear relationship between genetic distance and geographic distance. Plot a) represents IBD of the full dataset, while plot b) represents IBD of the outlier dataset. Both genetic distance and geographic distance were $\log_{10}$ transformed.

Fig. 5. The relationship between starling collection locality allele frequency and a locus strongly associated with a) Aridity (SNP ID 68063014), b) Bio05 (SNP ID 233190026), and c) Bio15 (SNP ID 196163030). The black line in each panel shows the linear relationship between the environmental variable and allele frequency, with the grey ribbon indicating the 95% confidence interval. Each point represents a collection locality with its shape indicating the genetic cluster that it belonged to; circle = Munglinup cluster, triangle = southern Australia cluster, square = southwest NSW, and cross = northern Australia cluster.

The RDA model was significant ($F_{7,491}$ = 1.2228, P < 0.001; Fig. 6), and reported

population clustering largely concurring with the results of the PCA environmental

parameters (Supplementary Material, Fig. S2). RDA identified 111 candidate SNPs that

showed strong association with the local environment (first 3 axes reported significance and

were used for SNP identification; $F_{1,491}$ = 2.0516, P < 0.001; $F_{1,491}$ = 1.2855, P < 0.001; $F_{1,491}$

= 1.1708, P < 0.008). The genetic variation between the westernmost starlings (WA and SA)

and NSW starlings correlated with a number of variables: Bio03 (Isothermality); Bio06 (Min

Temperature of Coldest Month); Bio15 (Precipitation Seasonality) ; Elev (Elevation); mnNDVI

(mean vegetation cover) (Fig. 6). Meanwhile, genotype differences between southern

starlings (VIC and TAS), and northern starlings (NSW and QLD) correlated with varDL

(Variability in Day Length) and Bio13 (Precipitation of Wettest Month) (Fig. 6).

From the 375 unique SNPs identified across all three methods (Table 2), a total of 25

proteins were identified when mapped to the starling reference genome (Supplementary

Material, Table S3). Of these 25 mapped proteins, 7 loci were associated with BAYESCAN F_{ST}

outliers, 11 were associated with BAYENV2 environmental variables, and 8 were associated

with candidate loci identified through the RDA environmental association (Supplementary

Material, Table S3). Among the annotated SNPs, we see a range of biological functions

traditionally associated with adaptation in invasive species, including processes such as

immune system responses (*C1QC* and *STAB1*) (Colautti, et al. 2004), temperature tolerance

(*HSPA9*), stress (*TRAF1*) (Zerebecki and Sorte 2011), and dietary alterations (*CTRL*) (Spit,

et al. 2014).

Fig. 6: The ordination plot of redundancy analysis (RDA) of 16177 SNP data for 499
starlings, collected from 24 sample sites across 5 Australian states. Individuals are coloured
by location, black arrows represent selected environmental variables. The relative placement
of environmental predictor arrows to individual and location sample clustering represents the
correlation between that environmental variable and the genetic differences seen in that
cluster.

Discussion

This study found that since their introduction into Australia, starling populations have become

genetically differentiated despite the potential for high levels of dispersal, and that selection

has facilitated their adaptation to the wide range of environmental conditions across their

geographic range. Isolation by distance appears to have played a strong role in determining

genetic substructure across the starling's Australian range, but neutral population structure

does not seem to be affected by environmental factors (IBE). Analyses of candidate SNPs

that are putatively under selection indicate that aridity, precipitation, and temperature may be

important factors driving selective change across the starling's invasive range in Australia,

indicating genetic differentiation in response to environment may be occurring at a limited

number of loci. We discuss some of our findings alongside a similar study on the invasive

North American starlings and explore population similarities and differences across these

replicated invasions. We do note that some loci that are candidates for selection suggest that

the faint genetic footprint left by the historic regime may hide, or be mistaken as, putative

adaptation.

Population structure

Population structure of Australian starlings

Our genome-wide SNP data set identified between two and four distinct genetic clusters.

When the genome wide data set was analysed at K=4, two of the genetic clusters covered

large geographic regions; the first encompassing southern Australia (WA to TAS), and the

other encompassing northern Australia (NSW and southern QLD). Munglinup and arid inland

NSW represented two small distinct clusters, indicating higher population isolation in these

areas. The population genetic structure of Australian starlings identified with the genome-

wide dataset of SNPs was similar to that identified in previous work, but contained some key

differences (Rollins, et al. 2009; Rollins, et al. 2011). Microsatellite data support the presence

of four populations: two small, distinct incursions into WA and two large populations, one in

SA, and another including VIC, TAS, and NSW (Rollins et al. 2009). It is likely that

microsatellite variation is less sensitive given the reduced power of the data set to

characterise differences, when compared to the SNP data set.

Interestingly, Munglinup represented the only population that reported a positive

Tajima's D (Table 1), which suggests demographic (e.g. a sudden population contraction) or

selection effects. Munglinup represents the western-most range edge, and it was likely

established from a small founding number of individuals with limited on-going gene flow and

potential selection, thereby providing conditions for more rapid divergence. Despite the fact

that Munglinup is only approximately 150km away from the next sampling site (Condingup),

we see distinct genetic differences at the range edge. Similar results were found in previous

studies (Rollins et al. 2011), which support the idea that the Munglinup subpopulation is a

result of a later, separate incursion into WA, and not a result of dispersion of Condingup

individuals, demonstrating that expanding ranges edges may be the site of complex

demographic processes.

Population structure analysis using the outlier loci dataset revealed only two

population clusters. A restricted outlier dataset (though prone to false positives) may allow for

the detection of adaptive variation , and has been used in many prior studies to further

resolve population structure than just using neutral loci (Hess, et al. 2013; Milano, et al.

2014; Benestan, et al. 2015). In this study, the genome-wide dataset identifying four genetic

clusters while the outlier loci identified two genetic clusters (Fig. 2). What the results may

indicate is two main genetic clusters may be differentiated by adaptive differences, whereas

demographic effects may be responsible for the structure seen using neutral loci.

The two large genetic clusters identified in the genome-wide and outlier loci dataset

largely overlap, and may be explained by two competing phenomena: 1) a combination of

geographic isolation and/or historic genetic diversity from different source populations, or 2)

adaptive selection driving genetic differentiation. Strong IBD, and the fact that the higher

subpopulation structure resolution of the genome-wide dataset determined that

subpopulations were not centred around unique introduction sites, points to the latter

explanation. Further, previous studies have not reported any IBD within regions centred

around introduction sites within the range (Rollins et al. 2009), indicating significant

exchange between localities and suggesting that the historic introduction regime is an

unlikely factor driving primary population substructure.

Comparison of Population structure of Australian and North American starlings

Replicated starling introductions in Australia and North America allow us to explore how

invasive populations may develop diverse genetic characteristics in response to different

environments. Although analyses indicate clearly defined environmental regions across the

starling's range in Australia, environmental parameters examined in this study explained little

of the genome-wide genetic structure observed in Australian starling populations

(Supplementary Material, Fig. S3). A non-significant environmental association using the

genome-wide SNP dataset may indicate that high levels of vagility are obscuring signals of

IBE. It is also possible that previously observed morphological association with

environmental gradients is due to plasticity or is driven by genetic change at only a small

number of loci. This later explanation is more likely, as geographic distance showed a strong

relationship with, and explained a large amount of the variation in, genetic difference

between collection localities, and also correlated with environmental distance. These findings

contrast with similar analysis conducted on the North American starlings, which found

geographic distance was a poor predictor of genetic differentiation across the species range,

and that environment may be playing a greater role in population similarity and clustering

(Hofmeister, et al. 2019). Some but not all starling populations migrate across North America,

and this variation in migratory strategy may result in a stronger signature of IBE.

Collection localities within regions of Australia show low levels of genetic

differentiation, nevertheless F_{ST} values are higher than those values reported in the larger

(square km coverage) North American starling distribution (Hofmeister, et al. 2019). This

further supports the notion that there are greater constraints to gene flow across the

Australian continent than in North America. We note that the genetic division between VIC

and NSW reported by both genome-wide and outlier loci SNP data sets straddles two key

environmental barriers; the Great Dividing Range (an area of high elevation), and inland

NSW (extreme aridity). Despite the fact that this is a highly dispersive species, it is possible

that these environmental features restrict gene flow. While further work is required to

examine the underlying causes for starling demographic and invasion history differences, this

system demonstrates both the malleability of invasion regimes and the importance of

studying replicated global invasions.

Putatively adaptive variation

There is an expectation that local adaptation in highly mobile species (such as starlings) will

be rare (Kvistad, et al. 2015). High levels of gene flow are often seen in highly mobile

species, which acts to homogenise genetic differences between populations, while stabilising

selection results in the loss of rare variants (North, et al. 2011). However, locally adapted

variants have been identified in species that are highly mobile (Nielsen, et al. 2009; Limborg,

et al. 2012; Bourret, et al. 2013; Hess, et al. 2013; Matala, et al. 2014; Milano, et al. 2014;

Benestan, et al. 2015), and in introduced species (Rohfritsch, et al. 2013; Chown, et al.

2015), which indicates that local adaptation can occur despite high levels of gene flow.

Over all analyses, several hundred unique candidate loci (375) for local adaptation

were identified, and from this, a total of 25 proteins were identified (Supplementary Material,

Table S3). Support for the occurrence of adaptive selection across the starling's range

comes from the functional nature of the polymorphisms detected (Supplementary Material,

Table S2) (Dudaniec, et al. 2018; Leydet, et al. 2018). Many of the biological functions are

shared with the functions of putative loci under selection identified from the 15,038 SNPs

found in in the North American starling study (Hofmeister, *et al.* 2019), and two genes were

reported as candidates for selection in both studies (*COL18A*, tissue morphogenesis; *DEF6*,

cell morphology regulation; Supplementary Material, Table S3). This overlap adds strength to

the notion that both invasive populations are undergoing concurrent rapid adaptation.

Interestingly, we identify many more SNPs that are associated with temperature than

precipitation or aridity. In contrast, Hofmeister *et al.* (2019) found a more even distribution

across a similarly diverse array of environmental predictors in environmental association

tests. While an excess of temperature associated SNPs may be (in part) a result of linkage,

one plausible biological explanation may be that annual temperatures of the starling's native

range are more similar to that of North America than Australia. Further, as invasive starling

populations are strongly associated with suburban and rural areas (Zufiaurre, et al. 2016), it

is possible that temperature may exert a selective pressure that is less affected by human-

mediated environmental disturbances than other selective forces (e.g. aridity). Further

analysis on the role anthropogenic land alteration has on starling spread and evolution is an

important future direction for this research.

Among the other proteins associated with Bio05 are a range of biological functions

plausibly linked to temperature, for instance a calcium channel protein (CACNA1C;

Supplementary Material, Table S2) that plays an important role in synaptic plasticity and

hence behaviour, with modification of this gene being associated with changed learning

strategies in mice (Koppe, et al. 2017). The importance of behavioural flexibility is a well-

established in invasion biology (Wright, et al. 2010), and starlings are known for high relative

brain sizes, a measure that has been linked to greater invasion success (Sol and Lefebvre

2000).

There was a particularly strong association between temperature and the candidate

SNP associated with the genomic region that codes for Tcdd-Inducible Poly(Adp-Ribose)

Polymerase (TIPARP). This protein is involved in pathways controlling androgen and

estrogen metabolic process (and hence sexual characteristics), as well as the development

of blood, skeleton and various organ structures (Supplementary Material, Table 3) (Ame, et

al. 2004). Given the large number of functions associated with TIPARP, there could be

multiple reasons it is associated with temperature, including variation in exposure to

xenobiotic chemicals across a temperature gradient, or because temperature variation is

highly dependent on latitude, and correlated with other seasonal fluctuations, both of which

cue sexual maturation in starlings (Dawson 2005).

However, closer examination of the plots of allelic frequency across environmental

variables reveals that environmental associations may be confounded by local signatures of

founding populations. Sampling sites that are fixed for a single allele may not be randomly

distributed, but rather may appear exclusively (or predominantly) within a single genetic

group. For example, the locus strongly associated with Bio15 (Fig. 5c) is fixed in eight

localities, six of which are in the northern Australia genetic cluster. Similar patterns can be

seen in Fig. 5b, while Fig. 5a represents a site where fixation does not occur within the same

subpopulation. It is possible that the alternative allele is not present in those areas due to

founding effects at introduction, and therefore there may be other loci linked to environmental

variables whose relationship is obscured by the legacy of limited diversity in some founding

populations. While genomic approaches provide unprecedented sensitivity and power for

testing for signatures of selection, caution should be applied when interpreting results for

invasive species, or populations that have recently undergone rapid expansion, as neutral

loci may then mimic signature of selection (Lotterhos and Whitlock 2014) (Riquet, et al.

2013). Characterising the native range genomic landscape for an invasive species would

provide invaluable information to help discern the nature of polymorphisms within their

invasive range, allowing us to identify if they are existing rare polymorphisms, or novel in the

invasive range. Lastly, our study demonstrates the utility of using multiple approaches when

searching for loci putatively under selection. We report some overlap between candidate loci

identified through the three methods we used. However, many of the loci were reported for

only one of the exploratory methods, and all of the annotated proteins identified were unique

to their identification method.

In Australian starling populations, we showed that geographic distance plays a strong

role in the genetic structure of populations, and that environmental barriers are likely

reinforcing the observed patterns of genetic structure. Our genetic evidence also suggests

that the contemporary Australian population of common starlings is most likely made up of

two large and spatially distinct populations, plus one smaller southwestern NSW group, and

one smaller genetic group existing at the Western-most range edge. These patterns are very

different from those observed in a similar study on North American starlings, a demonstration

that this species has the capacity for adaptive invasion strategies. Strong IBD and

environmental associations for individual loci are highly suggestive of adaptive genetic

variation, and doubtless these selective pressures will continue to drive population

differentiation.

Acknowledgements

PC was supported by an ARC Future Fellowship (FT0991420). LR was supported by an ARC Discovery Early Career Research Award (DE150101393). Thank you to Katherine L. Buchanan and Katie Hyma for their assistance with this project and manuscript. Thank you to Natalie R. Hofmeister for the useful and insightful discussions regarding these data. Thank you to the four anonymous reviewers for the constructive comments they provided, helping us to improve our manuscript. AC would like to acknowledge the lives of the 499 starlings who were killed and used for this research, and acknowledge an additional 236 starlings who were killed in conjunction with this project.

Data Accessibility

Scripts and data are archived on GitHub, and will be made public upon publication:

https://github.com/katarinastuart/Sv1_StarlingGBS

References

Almeida P, Goncalves C, Teixeira S, Libkind D, Bontrager M, Masneuf-Pomarede I, Albertin W, Durrens P, Sherman DJ, Marullo P, et al. (2014). A Gondwanan imprint on global diversity and domestication of wine and cider yeast Saccharomyces uvarum. Nature Communications, 5.

Ame JC, Spenlehauer C, de Murcia G. (2004). The PARP superfamily. Bioessays, 26, 882-893.

Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K, Dwight SS, Eppig JT, et al. (2000). Gene Ontology: tool for the unification of biology. Nature Genetics, 25, 25-29.

Bairoch A, Bougueleret L, Altairac S, Amendolia V, Auchincloss A, Puy GA, Axelsen K, Baratin D, Blatter MC, Boeckmann B, et al. (2008). The Universal Protein Resource (UniProt). Nucleic Acids Research, 36, D190-D195.

Baker HG, Stebbins GL. (1965). The Genetics of Colonizing Species. New York.: Academic Press.

Bay RA, Harrigan RJ, Le Underwood V, Gibbs HL, Smith TB, Ruegg K. (2018). Genomic signals of selection predict climate-driven population declines in a migratory bird. Science, 359, 83-86.

Benestan L, Gosselin T, Perrier C, Sainte-Marie B, Rochette R, Bernatchez L. (2015). RAD
genotyping reveals fine-scale genetic structuring and provides powerful population
assignment in a widely distributed marine species, the American lobster (Homarus
americanus). Molecular Ecology, 24, 3299-3315.

Benjamini Y, Hochberg Y. (1995). Controlling the false discovery rate - a practical and
powerful approach to multiple testing. Journal of the Royal Statistical Society Series B-
Statistical Methodology, 57, 289-300.

Bivand RS, Edzer p, Virgilio GR. (2013). Applied Spatial Data Analysis with R. New York:
Springer

Bourret V, Kent MP, Primmer CR, Vasemagi A, Karlsson S, Hindar K, McGinnity P, Verspoor
E, Bernatchez L, Lien S. (2013). SNP-array reveals genome-wide patterns of geographical
and potential adaptive divergence across the natural range of Atlantic salmon (Salmo salar).
Molecular Ecology, 22, 532-551.

Bradbury PJ, Zhang Z, Kroon DE, Casstevens TM, Ramdoss Y, Buckler ES. (2007).
TASSEL: software for association mapping of complex traits in diverse samples.
Bioinformatics, 23, 2633-2635.

Cabe PR. (1999). Dispersal and population structure in the European Starling. Condor, 101,
451-454.

Cabe PR. (1998). The effects of founding bottlenecks on genetic variation in the European
starling (Sturnus vulgaris) in North America. Heredity, 80, 519-525.

Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. (2009).
BLAST plus : architecture and applications. Bmc Bioinformatics, 10.

Capblancq T, Luu K, Blum MGB, Bazin E. (2018). Evaluation of redundancy analysis to
identify signatures of local adaptation. Molecular Ecology Resources, 18, 1223-1233.

Cardilini APA, Buchanan KL, Sherman CDH, Cassey P, Symonds MRE. (2016). Tests of
ecogeographical relationships in a non-native species: what rules avian morphology?
Oecologia, 181, 783-793.

Chown SL, Hodgins KA, Griffin PC, Oakeshott JG, Byrne M, Hoffmann AA. (2015).
Biological invasions, climate change and genomics. Evolutionary Applications, 8, 23-46.

Colautti RI, Ricciardi A, Grigorovich IA, MacIsaac HJ. (2004). Is invasion success explained
by the enemy release hypothesis? Ecology Letters, 7, 721-733.

Coop G, Witonsky D, Di Rienzo A, Pritchard JK. (2010). Using Environmental Correlations to
Identify Loci Underlying Local Adaptation. Genetics, 185, 1411-1423.

Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE, Lunter
G, Marth GT, Sherry ST, et al. (2011). The variant call format and VCFtools. Bioinformatics,
27, 2156-2158.

Dawson A. (2005). The effect of temperature on photoperiodically regulated gonadal
maturation, regression and moult in starlings - potential consequences of climate change.
Functional Ecology, 19, 995-1000.

Dlugosch KM, Parker IM. (2008). Founding events in species invasions: genetic variation,
adaptive evolution, and the role of multiple introductions. Molecular Ecology, 17, 431-449.

Dudaniec RY, Yong CJ, Lancaster LT, Svensson EI, Hansson B. (2018). Signatures of local
adaptation along environmental gradients in a range-expanding damselfly (Ischnura
elegans). Molecular Ecology, 27, 2576-2593.

Elshire RJ, Glaubitz JC, Sun Q, Poland JA, Kawamoto K, Buckler ES, Mitchell SE. (2011). A
Robust, Simple Genotyping-by-Sequencing (GBS) Approach for High Diversity Species. Plos
One, 6, e19379.

Feare CJ. (1985). The Starling: Issue 7 of Shire natural history.

Feng XJ, Jiang GF, Fan Z. (2015). Identification of outliers in a genomic scan for selection
along environmental gradients in the bamboo locust, Ceracris kiangsu. Scientific Reports, 5.

Fischer MC, Foll M, Excoffier L, Heckel G. (2011). Enhanced AFLP genome scans detect
local adaptation in high-altitude populations of a small rodent (Microtus arvalis). Molecular
Ecology, 20, 1450-1462.

Foll M, Gaggiotti O. (2008). A Genome-Scan Method to Identify Selected Loci Appropriate
for Both Dominant and Codominant Markers: A Bayesian Perspective. Genetics, 180, 977-
993.

Forester BR, Lasky JR, Wagner HH, Urban DL. (2018). Comparing methods for detecting
multilocus adaptation with multivariate genotype-environment associations. Molecular
Ecology, 27, 2215-2233.

Franks SJ, Munshi-South J. (2014). Go forth, evolve and prosper: the genetic basis of
adaptive evolution in an invasive species. Molecular Ecology, 23, 2137-2140.

Glaubitz JC, Casstevens TM, Lu F, Harriman J, Elshire RJ, Sun Q, Buckler ES. (2014).
TASSEL-GBS: A High Capacity Genotyping by Sequencing Analysis Pipeline. Plos One, 9,
9034.

Goudet J. (2005). HIERFSTAT, a package for R to compute and test hierarchical F-
statistics. Molecular Ecology Notes, 5, 184-186.

Griffiths R, Double MC, Orr K, Dawson RJG. (1998). A DNA test to sex most birds.
Molecular Ecology, 7, 1071-1075.

Gunther T, Coop G. (2013). Robust Identification of Local Adaptation from Allele
Frequencies. Genetics, 195, 205-220.

Gunther T, Coop G. 2018. A short manual for BayEnv2.
https://bitbucket.org/tguenther/bayenv2_public/src/default/bayenv2_manual.pdf.

Hess JE, Campbell NR, Close DA, Docker MF, Narum SR. (2013). Population genomics of
Pacific lamprey: adaptive variation in a highly dispersive species. Molecular Ecology, 22,
2898-2916.

Higgins PJ, Peter JM, Cowling SJ. (2006). Handbook of Australian, New Zealand & Antarctic
Birds. Volume 7 Boatbill to Starlings. Melbourne: Oxford University Press.

Hijmans RJ. 2016. Raster Geographic Data Analysis and Modeling. . https://cran.r-
project.org/web/packages/raster/index.html.

Hijmans RJ, Williams E, Vennes C. 2015. Geosphere: spherical trigonometry. https://CRAN.
R-project. org/package= geosphere.

Hoban S, Kelley JL, Lotterhos KE, Antolin MF, Bradburd G, Lowry DB, Poss ML, Reed LK,
Storfer A, Whitlock MC. (2016). Finding the Genomic Basis of Local Adaptation: Pitfalls,
Practical Solutions, and Future Directions. American Naturalist, 188, 379-397.

Hofmesiter NR, Werner SJ, Lovette IJ (2019) Environment but not geography explains
genetic variation in the invasive and largely panmictic European starling in North America. In.
Biorxiv.

Huang HT, Knowles LL. (2016). Unforeseen Consequences of Excluding Missing Data from
Next-Generation Sequences: Simulation Study of RAD Sequences. Systematic Biology, 65,
357-365.

Jeffreys H. (1961). Theory of probability. Oxford: Oxford University Press.

Jenkins CFH. (1977). The Noah's ark syndrome. Western Australia: The Zoological Gardens
Board.

Klopfstein S, Currat M, Excoffier L. (2006). The fate of mutations surfing on the wave of a
range expansion. Molecular Biology and Evolution, 23, 482-490.

Koppe G, Mallien AS, Berger S, Bartsch D, Gass P, Vollmayr B, Durstewitz D. (2017).
CACNA1C gene regulates behavioral strategies in operant rule learning. Plos Biology, 15.

Kvistad L, Ingwersen D, Pavlova A, Bull JK, Sunnucks P. (2015). Very Low Population
Structure in a Highly Mobile and Wide-Ranging Endangered Bird Species. Plos One, 10,
e0143746.

Lee CE. (2002). Evolutionary genetics of invasive species. Trends in Ecology & Evolution,
17, 386-391.

Legendre P, Lapointe FJ, Casgrain P. (1994). Modeling brain evolution from behaviour - a
permutational regression approach. Evolution, 48, 1487-1499.

Leydet KP, Grupstra CGB, Coma R, Ribes M, Hellberg ME. (2018). Host-targeted RAD-Seq
reveals genetic changes in the coral Oculina patagonica associated with range expansion
along the Spanish Mediterranean coast. Molecular Ecology, 27, 2529-2543.

Limborg MT, Helyar SJ, de Bruyn M, Taylor MI, Nielsen EE, Ogden R, Carvalho GR,
Bekkevold D, Consortium FPT. (2012). Environmental selection on transcriptome-derived
SNPs in a high gene flow marine fish, the Atlantic herring (Clupea harengus). Molecular
Ecology, 21, 3686-3703.

Linck E, Battey CJ. (2019). Minor allele frequency thresholds strongly affect population
structure inference with genomic data sets. Molecular Ecology Resources, 19, 639-647.

Loh PR, Tucker G, Bulik-Sullivan BK, Vilhjalmsson BJ, Finucane HK, Salem RM, Chasman
DI, Ridker PM, Neale BM, Berger B, et al. (2015). Efficient Bayesian mixed-model analysis
increases association power in large cohorts. Nature Genetics, 47, 284-290.

Long JL. (1981). Introduced Birds of the World. New York: Universe Books.

Lotterhos KE, Whitlock MC. (2014). Evaluation of demographic history and neutral
parameterization on the performance of F-ST outlier tests. Molecular Ecology, 23, 2178-
2192.

Lotterhos KE, Whitlock MC. (2015). The relative power of genome scans to detect local
adaptation depends on sampling design and statistical method. Molecular Ecology, 24, 1031-
1046.

Lu F, Lipka AE, Glaubitz J, Elshire R, Cherney JH, Casler MD, Buckler ES, Costich DE.
(2013). Switchgrass Genomic Diversity, Ploidy, and Evolution: Novel Insights from a
Network-Based SNP Discovery Protocol. Plos Genetics, 9, e1003215.

Matala AP, Ackerman MW, Campbell MR, Narum SR. (2014). Relative contributions of
neutral and non-neutral genetic differentiation to inform conservation of steelhead trout
across highly variable landscapes. Evolutionary Applications, 7, 682-701.

Milano I, Babbucci M, Cariani A, Atanassova M, Bekkevold D, Carvalho GR, Espineira M,
Fiorentino F, Garofalo G, Geffen AJ, et al. (2014). Outlier SNP markers reveal fine-scale
genetic structuring across European hake populations (Merluccius merluccius). Molecular
Ecology, 23, 118-135.

Nielsen EE, Hemmer-Hansen J, Poulsen NA, Loeschcke V, Moen T, Johansen T,
Mittelholzer C, Taranger GL, Ogden R, Carvalho GR. (2009). Genomic signatures of local
directional selection in a high gene flow marine organism; the Atlantic cod (Gadus morhua).
Bmc Evolutionary Biology, 9, 3128–3150.

North A, Pennanen J, Ovaskainen O, Laine AL. (2011). Local adaption in a changing world:
The roles of gene-flow, mutation, and sexual reproduction. Evolution, 65, 79-89.

Oksanen J, Guillaume BF, Friendly M, Kindt R, Legendre P, McGlinn D, Minchin PR, O'Hara
RB, Simpson GL, Solymos P, et al. 2018. vegan: Community Ecology Package. .
https://CRAN.R-project.org/package=vegan.

Pfeifer B, Wittelsburger U, Ramos-Onsins SE, Lercher MJ. (2014). PopGenome: An Efficient
Swiss Army Knife for Population Genomic Analyses in R. Molecular Biology and Evolution,
31, 1929-1936.

Pluess AR, Frank A, Heiri C, Lalaguee H, Vendramin GG, Oddou-Muratorio S. (2016).
Genome-environment association study suggests local adaptation to climate at the regional
scale in Fagus sylvatica. New Phytologist, 210, 589-601.

Raj A, Stephens M, Pritchard JK. (2014). fastSTRUCTURE: Variational Inference of
Population Structure in Large SNP Data Sets. Genetics, 197, 573-U207.

RCoreTeam. 2015. A language and environment for statistical computing. R Foundation for
Statistical Computing. Vienna, Austria.

Richardson MF, Sherman CDH. (2015). De Novo Assembly and Characterization of the
Invasive Northern Pacific Seastar Transcriptome. Plos One, 10, e0142003.

Riquet F, Daguin-Thiebaut C, Ballenghien M, Bierne N, Viard F. (2013). Contrasting patterns
of genome-wide polymorphism in the native and invasive range of the marine mollusc
Crepidula fornicata. Molecular Ecology, 22, 1003-1018.

Rohfritsch A, Bierne N, Boudry P, Heurtebise S, Cornette F, Lapegue S. (2013). Population
genomics shed light on the demographic and adaptive histories of European invasion in the
Pacific oyster, Crassostrea gigas. Evolutionary Applications, 6, 1064-1078.

Rollins LA, Moles AT, Lam S, Buitenwerf R, Buswell JM, Brandenburger CR, Flores-Moreno
H, Nielsen KB, Couchman E, Brown GS, et al. (2013). High genetic diversity is not essential
for successful introduction. Ecology and Evolution, 3, 4501-4517.

Rollins LA, Woolnough AP, Sinclair R, Mooney NJ, Sherwin WB. (2011). Mitochondrial DNA
offers unique insights into invasion history of the common starling. Molecular Ecology, 20,
2307-2317.

Rollins LA, Woolnough AP, Wilton AN, Sinclair R, Sherwin WB. (2009). Invasive species
can't cover their tracks: using microsatellites to assist management of starling (Sturnus
vulgaris) populations in Western Australia. Molecular Ecology, 18, 1560-1573.

Ross HA. (1983). Genetic differentiation of starling (Sturnus-vulgaris-aves) populations in
New Zealand and Great Britain Journal of Zoology, 201, 351-362.

Sherman CDH, Lotterhos KE, Richardson MF, Tepolt CK, Rollins LA, Palumbi SR, Miller AD.
(2016). What are we missing about marine invasions? Filling in the gaps with evolutionary
genomics. Marine Biology, 163, 198.

Sol D, Lefebvre L. (2000). Behavioural flexibility predicts invasion success in birds
introduced to New Zealand. Oikos, 90, 599-605.

Spit J, Zels S, Dillen S, Holtof M, Wynant N, Broeck JV. (2014). Effects of different dietary
conditions on the expression of trypsin- and chymotrypsin-like protease genes in the
digestive system of the migratory locust, Locusta migratoria. Insect Biochemistry and
Molecular Biology, 48, 100-109.

Tajima F. (1989). Statistical-method for testing the neutral mutation hypothesis by DNA
polymorphism. Genetics, 123, 585-595.

Tepolt CK, Darling JA, Bagley MJ, Geller JB, Blum MJ, Grosholz ED. (2009). European
green crabs (Carcinus maenas) in the northeastern Pacific: genetic evidence for high
population connectivity and current-mediated expansion from a single introduced source
population. Diversity and Distributions, 15, 997-1009.

Wang IJ. (2013). Examining the full effects of landscape heterogeneity on spatial genetic
variation: A multiple matrix regression approach for quantifying geographic and ecological
isolation. Evolution, 67, 3403-3411.

Wang IJ, Glor RE, Losos JB. (2013). Quantifying the roles of ecology and geography in
spatial genetic divergence. Ecology Letters, 16, 175-182.

Waterman M, Fuller CO, Murray MD. (2008). Studies of roosting common starlings Sturnus
vulgaris in South Australia. Corella, 32, 25–29.

Whitehead SC, Wright J, Cotton PA. (1995). Winter field use by the European starling
Sturnus vulgaris - habitat preferences and the availability of prey Journal of Avian Biology,
26, 193-202.

Woolnough AP, Gray GS, Lowe TJ, Kirkpatrick WE, Rose K, Martin GR. (2005). Distribution
and abundance of pest animals in Western Australia: A survey of institutional knowledge.
Western Australia: Vertebrate Pest Research Section, Department of Agriculture.

Wright TF, Eberhard JR, Hobson EA, Avery ML, Russello MA. (2010). Behavioral flexibility
and species invasions: the adaptive flexibility hypothesis. Ethology Ecology & Evolution, 22,
393-404.

Zerebecki RA, Sorte CJB. (2011). Temperature Tolerance and Stress Proteins as
Mechanisms of Invasive Species Success. Plos One, 6, e14806.

Zorner RJ, Trabucco A, Bossio DA, Verchot LV. (2008). Climate change mitigation: A spatial
analysis of global land suitability for clean development mechanism afforestation and
reforestation. Agriculture Ecosystems & Environment, 126, 67-80.

Zufiaurre E, Abba A, Bilenca D, Codesido M. (2016). Role of landscape elements on recent
distributional expansion of European starlings (Sturnus vulgaris) in agroecosystems of the
Pampas, Argentina. Wilson Journal of Ornithology, 128, 306-313.