Delta phase resets mediate non-rhythmic temporal prediction

Jonathan Daume[a]*, Peng Wang[a], Alexander Maye[a], Dan Zhang[b] and Andreas K. Engel[a]

[a] Department of Neurophysiology and Pathophysiology, University Medical Center Hamburg-Eppendorf, Hamburg, 20246, Germany

[b] Department of Psychology, School of Social Sciences, Tsinghua University, Beijing, 100084, China

---

* Correspondence: j.daume@uke.de; twitter: @jonathan_daume

1 Abstract

The phase of neural oscillatory activity aligns to the predicted onset of upcoming
stimulation. Whether such phase alignments represent phase resets of underlying neural
oscillations or just rhythmically evoked activity, and whether they can be observed in a
rhythm-free visual context, however, remains unclear. Here, we recorded the
magnetoencephalogram while participants were engaged in a temporal prediction task judging
the visual or tactile reappearance of a uniformly moving stimulus. The prediction conditions
were contrasted with a control condition to dissociate phase adjustments of neural oscillations
from stimulus-driven activity. We observed stronger delta band inter-trial phase consistency
(ITPC) in a network of sensory, parietal and frontal brain areas, but no power increase
reflecting stimulus-driven or prediction-related processes. Delta ITPC further correlated with
prediction performance in the cerebellum and visual cortex. Our results provide evidence that
phase alignments of low-frequency neural oscillations underlie temporal predictions in a non-
rhythmic visual and crossmodal context.

15 Keywords

16   Temporal prediction; crossmodal prediction; neural oscillations; delta band; beta band;
17 inter-trial phase coherence; spectral power; phase reset; cerebellum; magnetoencephalography

Introduction

Neural oscillations reflect alternating states of higher or lower neural excitability,
modulating the efficiency by which coupled neurons engage in mutual interactions (Buzsáki,
2006). As a result, neural communication and information processing has been shown to
occur in a phase-dependent manner (Engel et al., 2001; Fries, 2005) reflected, for example, by
fluctuations in perception thresholds correlating with the phase of ongoing oscillations
(VanRullen, 2016). Based on these assumptions, oscillations were also linked to temporal
predictions of upcoming relevant information (Arnal and Giraud, 2012; Engel et al., 2001;
Rimmele et al., 2018). Studies have shown that animals can utilize predictive aspects of
environmental stimuli in a way that reaction times are reduced (Gould et al., 2011; Lakatos et
al., 2008; Rohenkohl and Nobre, 2011; Stefanics et al., 2010) or stimulus processing is
enhanced (Cravo et al., 2013; Wilsch et al., 2015). By means of top-down induced phase
resets of neural oscillations, phases of high excitability might be adjusted towards the
expected onset of relevant upcoming stimulation in order to optimize relevant behavior
(Schroeder and Lakatos, 2009).

Due to the rhythmic and therefore temporally highly predictable nature of many auditory
stimuli such as speech or music, particularly in the auditory domain, many studies gathered
evidence that oscillations reset and thereby adjust their phase towards rhythmic stimuli of
various frequencies (Doelling and Poeppel, 2015; Giraud and Poeppel, 2012). Also in the
visual domain, studies showed that neural oscillations align to temporal structure rhythmic
visual input (Breska and Deouell, 2017b; Cravo et al., 2013; Gomez-Ramirez et al., 2011;
Lakatos et al., 2008; Saleh et al., 2010). Other studies, however, reported a specific effect for
visual temporal predictions only in the alpha band (8 – 12 Hz), although sensory input was
provided at lower frequencies (Rohenkohl and Nobre, 2011; Samaha et al., 2015).

Moreover, whether temporal predictions indeed involve phase resets of endogenous
neural oscillations remains a matter of debate (Breska and Deouell, 2017a; Doelling et al.,
2019; Novembre and Iannetti, 2018). Despite their ecological relevance, using rhythms for the
investigation of an involvement of oscillations in temporal predictions entails methodological
and conceptual challenges. Rhythmic input leads to a continuous stream of regularly bottom-
up evoked potentials, which are – at least – difficult to distinguish from top-down phase
adjusted neural oscillations within the same frequency. Rather than by phase resets of
endogenous neural oscillations, phase alignments across trials could therefore also be caused
by stimulus-evoked potentials that just appear to be rhythmic during rhythmic stimulation
(Doelling et al., 2019; Novembre and Iannetti, 2018; Zoefel et al., 2018).

Temporal prediction processes have further been shown to be reflected by slow buildups
of neural activity, which ramps up until the predicted time point is reached; also called
contingent negative variation (CNV; Breska and Deouell, 2017a; Macar et al., 1999). In a
rhythmic temporal prediction context, such slowly ramping activity between subsequent
stimulus pairs would also lead to significant phase-locking of low-frequency activity across
trials, which again would be very difficult to be distinguished from phase-locking of phase-
aligned endogenous neural oscillations. Conclusive evidence that temporal predictions
involve phase resets of endogenous oscillations rather than stimulus-driven or prediction-
evoked potentials is still lacking.

In addition, using only auditory rhythmic stimulation precludes the opportunity to link
phase adjustments to a more general neural mechanism that predicts the temporal structure of
any external input. If phase adjustments form the basis of tracking the temporal regularities of
any relevant information, neural oscillations should align also to predictable temporal
regularities that are inferred from input that does not itself comprise auditory rhythmic or
discrete components, such as, for instance, uniform visual motion.

For this reason, we set out to investigate whether phase adjustments of neural activity can
be observed for predictable visual motion stimuli. We measured magnetoencephalography
(MEG) while healthy participants watched a visual stimulus continuously moving across the
screen until it disappeared behind an occluder (Fig. 1A). We manipulated the time for the
stimulus to reappear on the other side of the occluder. The task was to judge whether the
stimulus reappeared too early or too late based on the speed of the stimulus earlier to
disappearance. Hence, participants were required to temporally predict the correct time point
of reappearance to be able to accomplish the task. Participants further performed a control
task, in which the task was to judge the luminance of the reappearing stimulus instead of its
timing. Importantly, physical appearance of both conditions was exactly the same in all
aspects of the stimulation. Any purely stimulus-driven, bottom-up activity should therefore
level out between the two conditions.

Moreover, since it has been shown that sensory stimulation can lead to crossmodal phase
adjustments also in relevant but unstimulated other modalities (Lakatos et al., 2007; Mercier
et al., 2013), we further introduced a third condition in which a tactile instead of a visual
stimulus was presented at reappearance. By contrasting it to the luminance matching control
condition, we sought to determine whether phase alignments can be observed in regions
associated with tactile stimulus processing, when sensory information was in fact only
provided to the visual system.

We hypothesized that in the two temporal prediction tasks, as compared to the luminance
matching control task, we would observe stronger inter-trial phase consistency (ITPC) within
time windows between disappearance and expected reappearance. These phase alignments
should particularly be observed at low frequencies, e.g., in the delta band, matching the
temporal scale of the disappearance window (on average 1.5 s). Importantly, if such enhanced
ITPC reflected phase resets of ongoing neural oscillations, we should not observe any
changes in delta power during temporal predictions, as the amplitude of phase-resetting
endogenous oscillations should not be altered. On the other hand, when stimulus-driven or
prediction-evoked neural activity lead to the observed phase alignments, observed ITPC
differences should be accompanied with differences in total delta power during temporal
predictions, representing the evoked neural activity in each trial. Further, if the phase of
neural oscillations indeed codes for the time point of the expected reappearance in each
participant, participants showing a more consistent judgment of reappearance timing – as
represented by a steep slope of the psychometric function – should have stronger ITPC during
temporal predictions than participants who performed less accurately. If evoked neural
activity underlies temporal predictions, these correlations should as well be accompanied by
correlations between delta power and the steepness of the psychometric curve within the same
brain region.

Results

Behavioral results

Participants did not receive feedback about the correctness of their response. This ensured that participants relied on their individual and subjective "right on time" (ROT) impression in the temporal prediction conditions and "point of subjective equivalence" (PSE) in the luminance matching condition. Across participants, there was no statistically significant bias towards "too early/darker" or "too late/brighter" responses in the visual temporal prediction (Δt (ROT_V) = 13.15 ± 155.20 ms; $t(22)$ = .41; p = .69; Cohen's d = .09) or in the luminance matching task (ΔRGB (PSE) = -1.29 ± 4.54 RGB; $t(22)$ = -1.36; p = .19; Cohen's d = -.28), respectively (Fig. 1B). In the tactile temporal prediction task, participants showed a significant bias towards "too early" responses (Δt (ROT_T) = 99.80 ± 150.00 ms; $t(22)$ = 3.19; p = .004; Cohen's d = .67).

To assess whether reaction times were dependent on the timing of the reappearing stimulus (Fig. 1C), we fitted a mixed-effect model to reaction times from all trials using the categorial variable *condition* (with the luminance matching task as reference level) and *timing difference* as well as their interaction as predictors. Since in the temporal prediction conditions we expected reaction times to be slowest for timing differences around zero and faster for high timing differences, we used a second-order polynomial term for *timing differences* (see Methods). Across all timing differences, reaction times were significantly faster in the tactile temporal prediction task as compared the luminance matching task (β = -0.26; t = 14.21; p < 0.001), but not significantly different between the visual temporal prediction and the luminance matching task (β = -0.03; t = -1.34; p = 0.18). Across all conditions, reaction times linearly decreased with increase timing difference (β = -0.04; t = -2.48; p = 0.02) as well as showed a quadratic relationship with timing difference (β = 0.02; t = 2.42; p = 0.02). Importantly, as indicated by the interaction results, timing difference had a stronger negative linear (β = -0.13; t = -10.53; p < 0.001) and stronger negative quadratic influence on reaction times from the visual (β = -0.11; t = -8.18; p < 0.001) as well as a stronger negative quadratic influence on reactions times from the tactile temporal prediction task (β = -0.10; t = -7.52; p < 0.001) as compared to those from the luminance matching task (see Figure 1C and supplementary table S1 for the complete model output).

Figure 1. Experimental design and behavioral results. (**A**) A stimulus moved towards the center of the screen until it disappeared behind an occluder. The task was to judge whether the stimulus reappeared *too early* or *too late*. In the luminance matching condition, task was to judge whether the luminance became *brighter* or *darker*. Importantly, physical stimulation was exactly the same as in the visual prediction task. In the tactile temporal prediction task, at reappearance a tactile stimulus was presented contralateral to the disappearance of the visual stimulus. (**B**) Psychometric functions and individual ROT/PSE estimates. A timing difference of 0 refers to the objectively correct reappearance of the stimulus after 1,500 ms. Analogously, a luminance difference of 0 refers to equal luminance after reappearance provided in RGB values (see Methods). Colored areas depict standard errors of the mean (SEM). (**C**) Log-transformed and standardized reaction times for all timing differences (mean ± SEM). P = proportion; LM = luminance matching; t = time; l = luminance; RGB = red-green-blue.

Temporal prediction was associated with reduced beta power in sensory regions

Analyzing the neural data, we were first interested in investigating which frequency bands showed modulated spectral power during windows of temporal predictions in order to narrow down frequency bands of interest for further analyses. For that, we tested an average of spectral power across all sensors and conditions against a pre-stimulus baseline window. As a first step, we obtained a general overview of power modulations at each event in the experimental paradigm. Due to the jittered stimulation built into the design (see Materials and Methods), we computed cluster-based permutations statistics in three separate time windows (Fig. 2A) centered on: (a) the onset of the moving stimulus ("Movement"), (b) disappearance of the stimulus behind the occluder ("Disappearance"), and (c) reappearance of the stimulus ("Reappearance").

In time bins around movement onset as well as reappearance of the stimulus, but not around disappearance, clusters of frequencies in the delta and theta range showed a statistically significant increase of spectral power as compared to the baseline window. All time windows further depicted a significant decrease of spectral power in frequencies within the beta and gamma range (all cluster p-values < .008). Importantly, even with using a liberal cluster alpha level of .05 (one-sided), we did not find a statistically significant modulation of delta power during the disappearance window. This was also not the case when reducing the test to sensors from occipital regions only (see Fig. S1).

Since we were most interested in examining modulations associated with temporal predictions, i.e., during the disappearance window, we further compared spectral power estimates between the temporal prediction tasks and the luminance matching task in all sensors within the disappearance window while ignoring the other windows. We restricted our analysis to the classical beta band ranging from 13 to 30 Hz, showing the strongest modulation as compared to baseline during the disappearance window. Cluster-based permutation statistics revealed reduced beta power during visual temporal prediction in occipital sensors during all time-bins of the disappearance window (cluster-p = .01). Source level statistics revealed a statistically significant decrease of beta power in a cluster of bilateral occipital voxels (cluster-p = .01). Beta power was further reduced during tactile prediction in a cluster of occipital as well as left lateralized frontocentral sensors (cluster-p = .002). At source level, a significant power reduction in the beta band was most strongly apparent in parts of bilateral visual as well as left-lateralized somatosensory cortex (cluster-p = .01).

Figure 2. Power modulations during temporal prediction. (**A**) Spectral power averaged across sensors, conditions, and participants. Each window was centered on the different events within the paradigm and normalized with pre-stimulus baseline. Time 0 refers to the onset of each event. Cluster-based permutation statistics revealed significant power modulations as compared to baseline (unmasked colors). See also Fig. S1. (**B,C**) Difference between the two temporal prediction and the luminance matching task, respectively, within the beta band (13 – 30 Hz) in time bins around stimulus disappearance. Black dots indicate sensors of the clusters showing significant differences between the conditions. At source level, cluster-based permutation statistics revealed cluster of voxels showing significant differences between the conditions (colored voxels). LM = luminance matching.

Delta inter-trial phase consistency was enhanced during temporal prediction

For the analysis of ITPC, we followed a similar approach. First, we tested ITPC
differences to baseline in the three time windows for an average across all sensors and
conditions using cluster-based permutation statistics. ITPC was significantly increased across
a range of different frequencies in time bins around movement onset, disappearance and
reappearance of the stimulus (all cluster-$p < .001$; Fig. 3A). For time windows centered on
movement onset as well as reappearance significant ITPC increases were strongest in the
delta to alpha range. At disappearance of the stimulus, significant ITPC increases were
observed up to the low beta range with strongest increases in the delta band.

Hence, the delta band showed no increase in power but the strongest increase in ITPC as
compared to baseline during the disappearance window for an average across all conditions
(see Fig. 2A, 3A, and S1). For further statistical comparisons between conditions, we
therefore restricted our analyses to an average of frequencies between 0.5 to 3 Hz (for
condition-specific delta band ITPC differences to baseline during disappearance, see Fig. S2).
For a better estimation of when differences in ITPC between the conditions became apparent,
we enlarged the analysis of ITPC to time bins ranging from -1,900 ms to 1,900 ms centered
on the disappearance of the stimulus. Note that in this enlarged analysis window the timing of
the movement onset as well as the reappearance of the stimulus strongly jittered across trials.
The effect of these events on ITPC estimates were thus strongly reduced (see Fig. S3).

We found two clusters that showed significantly stronger ITPC during visual temporal
predictions as compared to luminance matching (Fig. 3B). One cluster included sensors from
right temporal, frontal and occipital regions in time bins from -400 to 1,900 ms (cluster $p <$
.001). The second cluster included left frontotemporal sensors in time bins ranging from 0 to
1,900 ms (cluster $p = .01$) Source level analysis revealed that for an average of the time
window from -400 to 1,900 ms ITPC differences between the two conditions were strongest
in right-lateralized central and inferior frontal voxels (cluster $p < .001$).

ITPC was also significantly enhanced in bilateral temporal sensors during tactile
temporal predictions, evolving around -400 ms in right temporal sensors and shifting towards
left hemisphere with ongoing disappearance time (cluster $p < .001$; Fig. 3C). In this contrast,
however, differences in ITPC were more strongly apparent also in frontal and central sensors.
Besides strongest differences in ITPC again in right superior parietal and inferior frontal
voxels, source level analysis also revealed strong differences in bilateral somatosensory
voxels for the contrast of tactile prediction to luminance matching (cluster $p < .001$).

Figure 3D depicts absolute delta ITPC estimates for all three conditions in the enlarged
disappearance time window. Values were averaged across participants and all the sensors that
exhibited the top 20% of t values in the ITPC contrast between visual temporal prediction and
luminance matching between 0 and 1,500 ms (see Fig. 3B; similar results were obtained for
sensors showing the top 10% or 5% of t values, see Fig. S3D). ITPC initially increased for all
three conditions, but dropped down to stimulus movement level shortly afterwards in the
luminance matching condition. ITPC in the visual as well as tactile temporal prediction tasks
stayed elevated throughout the entire disappearance window.

Figure 3. ITPC during temporal prediction as compared to luminance matching. (**A**) ITPC estimates averaged across sensors, conditions, and participants. Masked colors refer to non-significant ITPC modulations as compared to baseline (cluster-based permutation statistics). (**B,C**) Difference in ITPC between the visual or tactile prediction and the luminance matching task, respectively, within the delta band (0.5 – 3 Hz). For clarity, only every second time bin was plotted. Black dots indicate sensors of the clusters showing significant differences. On source level, clusters of voxels showing significant differences between the conditions are colored. See also Fig. S2 and S3 (**D**) Time course of absolute delta ITPC estimates within each condition for time bins centered around disappearance of the stimulus (time 0; mean ± SEM). ITPC estimates were averaged across channels that showed the top 20% of t-values for the comparison of the visual prediction with the luminance matching task (see topography). LM = luminance matching.

Control analyses on delta power differences between conditions

In contrast to ITPC, delta power did not significantly increase with disappearance of the stimulus in an average across conditions and channels as compared to baseline (see Fig. 2A and S1). Nevertheless, to examine whether channels showing the strongest differences in ITPC between conditions also show differences in delta power, we averaged delta power within the channels showing the strongest ITPC differences (same as in Fig. 3D) and

compared power values from each of the two temporal prediction conditions with the
luminance matching condition, respectively, within the same enlarged window of -1.900 to
1.900 ms around stimulus disappearance (Fig. 4).

Figure 4A shows the time courses of total delta power in each condition. The overall
pattern of the delta power time courses was largely different to the pattern of the ITPC time
course in each of the conditions (compare to Fig 3D). In the visual temporal condition task as
well as in the luminance matching task, delta power did not increase around disappearance of
the stimulus and did not differ in any of the time bins between the two conditions during the
disappearance window (even for uncorrected t-tests). Also in the tactile condition, delta power
did not increase at around disappearance of the stimulus as observed for ITPC in this
condition. However, it strongly increased in late time windows, showing significant
differences in the tactile conditions as compared to the luminance matching task in time bins
between 700 and 1900 ms after disappearance (cluster $p < .001$, dashed black line in Fig. 4A).

The combined pattern of an early delta ITPC increase and a late delta power increase in
this condition could speak in favor of a CNV underlying the processes of temporal prediction.
A CNV describes activity that is building up until the expected time point of an upcoming
event is reached. After this time point, the build-up process sharply terminates (see, e.g.
Breska and Deouell, 2017a; Macar et al., 1999). In such a scenario, ITPC would be increased
as soon as the slow build-up process initiates (here at disappearance), but power increases
might become observable only later in the prediction process. A phase-reset of ongoing
oscillations, on the other hand, should not lead to an increase in delta power during the
disappearance window.

To further investigate whether a CNV could explain the observed pattern of ITPC and
power time courses, we computed additional control analyses. If a CNV would explain the
increase in total power in the tactile condition, it should be locked to the disappearance of the
stimulus and be present in each temporal prediction trial. Consequently, it should also be
removed when subtracting the ERF from each trial in the time domain, before computing
delta power in each single trial (i.e., when computing induced power). However, as the upper
panel in Figure 4B shows, even after removing the ERF from each trial, delta power in the
tactile condition was still strongly increased as compared to the luminance matching task in
late disappearance time windows (cluster p < .001). Delta ITPC, on the other hand, was
completely removed after subtracting the ERF (Fig. 4B lower panel).

Figure 4C depicts the delta power time course of the ERF itself in each condition. Delta
power of the ERF increased for both, the tactile as well as the visual temporal prediction task,
as compared to the control task. Similar to the ITPC effect, this increase already started in
time bins shortly before disappearance (both cluster $p < .001$). Moreover, the strength of the
increase in power in the visual temporal prediction task resembles the increase in the tactile
task, and was not much stronger in the tactile task as observed for total power (Fig. 4A).

As a next step, we computed two mixed-effect regression models to examine the effect of
delta power on ITCP. In one model, we used the variables *condition* and *time* as well as their
interaction as predictors for ITPC only (Fig. 4D upper panel). In the other, we also added
delta power as a covariate to the model in order to adjust for the variance explained by delta
power (lower panel). After adding delta power as covariate, predicted ITPC values were
reduced in the tactile prediction condition during late time windows of disappearance.
However, they were still significantly different between both the visual and the tactile
temporal prediction as compared to the luminance matching task, respectively, in all time bins
during disappearance (see supplementary Table S2 for a complete model output).

Figure 4. Delta power control analyses. (**A**) Time course of total delta power (0.5 – 3 Hz) in all conditions for the channels showing the strongest delta ITPC effect (same as in Fig. 3D). Time point 0 again refers to the complete disappearance of the stimulus. Colored lines depict uncorrected p-values below 0.05 from comparisons of the respective temporal prediction condition with the luminance matching task in each time bin. Dashed black lines depict p-values that survived the cluster-based permutation test. (**B**) The upper panel depicts the time course of induced delta power in each condition after a condition-wise subtraction of the ERF from each trial in the time domain. The lower panel depicts ITPC in each condition after ERF subtraction. (**C**) Delta power time course of the ERF, i.e., after averaging all trials in each condition in the time domain first. (**D**) Predicted delta ITPC values from mixed-effects regression models with an interaction term of *condition* and *time* as predictors for ITPC. Upper panel: without adjusting for delta power; lower panel: with adjusting for delta power by adding power as a covariate to the model. For a better comparability, standardized ITPC values were back-transformed to the original scale prior to plotting.

Delta ITPC, but not delta power, correlated to behavioral performance

We further hypothesized that if phase alignments of neural oscillations were indeed associated with temporal predictions, a participant who judged the reappearance of the stimulus within her individual subjectively correct ROT framework in a consistent manner should also exhibit stronger ITPC during temporal predictions, as a consistent timing judgement across trials should involve a similar phase across trials. The consistency of judgements can be inferred from the steepness of the psychometric function – the steeper the psychometric function, the more consistent the answers of the participant. We computed Pearson correlations of source level delta ITPC with the steepness of the psychometric function across participants and found statistically significant positive correlations in the visual (cluster *p* = .003) as well as in the tactile temporal prediction task (cluster *p* = .002; Fig. 5). Strongest correlations were found in the cerebellum and right lateralized early visual areas

in both tasks. No clusters showing significant positive or negative correlations were observed
in the luminance matching task (all cluster *p* > .1). If such correlations between phase
alignments and behavior are related to evoked neural activity during temporal predictions,
however, we should also observe similar correlation also between delta power and behavior.
Hence, we averaged delta power within the voxels that showed the correlations between ITPC
and behavior and computed Pearson correlations between this average and the steepness of
the slope in each condition. We found no significant correlation between delta power and
behavior in the visual (r = 0.31, *p* = 0.15) nor in the tactile temporal prediction task (r = 0.16,
p = 0.47).

Figure 5. Correlation of ITPC to behavior. (A,B) Correlation of individual ITPC estimates with the individual
steepness of the psychometric function within all voxels, shown in (A) for the visual prediction, and in (B) for the
tactile prediction condition. ITPC estimates were averaged within the delta band and time windows of 0 to 1,000
ms centered on the disappearance of the stimulus. Only the clusters of voxels showing significant correlations are
colored. In the scatter plots, each dot represents one participant and ITPC estimates were averaged across all
voxels within the clusters of significant correlations. There was no significant correlation observed for the
luminance matching condition or between delta power and behavior.

ITPC did not correlate with eye movements

A potential confound for the observed effects in ITPC could be that participants tracked
the moving stimulus with their eyes to be able to judge the correct time point of reappearance.
Thus, consistent horizontal eye movements with the speed of the stimulus might lead to
enhanced ITPC in the delta band. To make sure that differences in eye movements do not
explain the observed differences in ITPC between the conditions, we analyzed horizontal eye
movements recorded by an eye tracker (ET) during the MEG measurement. Figure 6A depicts
condition-wise horizontal eye positions averaged across all participants and centered on the
disappearance of the stimulus, showing no systematic differences between the conditions.

Moreover, if horizontal eye movements would explain the effects in ITPC, we should observe
the same effects between the conditions when we compute ITPC for the ET data. Differences
in ITPC between the two temporal prediction conditions and the luminance matching
condition are depicted Figure 6B and C. Using cluster-based permutation statistics, we did not
observe any time-frequency cluster that revealed significant differences between the
conditions (all cluster $p > .1$).

Further, we tested whether there are any significant correlations between individual ITPC
values obtained from the MEG data and from the ET data. We averaged ITPC values from a
time window of 0 to 1.000 ms and again used the top 20% of channels showing the strongest
effect for ITPC for the MEG data (for channels see Fig. 3D), and did not observe significant
correlations between the ITPC values obtained from MEG and ET data in the temporal
prediction tasks (Fig. 6D). The strongest correlation was found in the luminance matching
condition, which suggests that the ITPC differences found in the MEG data cannot be
explained by horizontal eye movements during temporal predictions.

Figure 6. Analysis of horizontal eye movements. (**A**) Condition-wise eye positions centered on stimulus
disappearance (time 0 s) and averaged across all participants. A visual angle of 0° refers to the fixation dot (1°
visual angle roughly corresponds to 1 cm on the screen). Colored areas depict SEM. (**B,C**) Differences in ITPC
between (**B**) the visual prediction and (**C**) the tactile prediction condition to the luminance matching condition in
low frequencies and time bins around disappearance of the stimulus (time 0 s). Utilizing cluster-permutation
statistics, no clusters of significant differences were observed between the conditions. (**D**) Condition-wise
correlations between ITPC estimates obtained from the eye tracker data and the MEG sensors across all
participants. ET = eye tracker.

Discussion

Our task design enabled us to disentangle phase resets of ongoing neural oscillations from evoked event-related potentials. We found that phase alignments, but not power, were stronger in the context of temporal predictions than in a task where temporal structure was less relevant. This supports the hypothesis that phase adjustments of ongoing neural oscillations, and not stimulus-driven or prediction-evoked activity, form the neuronal basis of temporal prediction processes and suggest that this framework can be extended to predictions that have to be inferred from stimulation that does not itself comprise rhythmic and discrete components. The strength of the observed phase adjustments further correlated with the ability to consistently judge the temporal reappearance of the stimulus across participants, suggesting also a functional relevance of the observed phase adjustments for temporal predictions.

Cross-modal temporal predictions are reflected by a beta power reduction in both sensory systems

It has been suggested that temporal predictions of upcoming events might be mediated by neuronal oscillations in the delta and beta frequency range (Arnal and Giraud, 2012). The enhanced phase consistency of delta oscillations as well as the power modulations in the beta band observed in the current study are in line with this hypothesis. However, earlier reports on beta power modulations during temporal predictions are inconsistent. On the one hand, studies found that beta power was even increased shortly before the onset of the expected stimulus in auditory (Arnal et al., 2015) and visual rhythmic stimulation (Saleh et al., 2010). On the other hand, van Ede et al. (van Ede et al., 2011) found that predicting the onset of a tactile stimulus was specifically associated with a reduction of beta power in contralateral tactile areas and accompanied by faster reaction times. The authors suggest that a reduction in beta power might signal preparatory processes in the sensory system that expects the upcoming event.

The observed decrease in beta power in task-relevant sensory regions in the current study largely match the results reported by van Ede et al. (van Ede et al., 2011). During visual temporal predictions, beta band power was reduced in visual sensory regions as compared to the visual control condition during the entire disappearance time. During crossmodal predictions, in which temporal information was provided to the visual system, but reappearance was expected in the tactile domain, beta band power was decreased in both, visual as well as tactile regions.

Since also in the luminance matching condition participants expected to perceive a visual stimulus, preparatory processes alone cannot explain this reduction in beta power. This is especially the case in the crossmodal condition, in which no visual stimulus was expected, but stronger decreases in beta were also observed in visual areas. Moreover, since we observed beta decreases also in tactile regions at the time of visual stimulus disappearance, the decrease could not solely be an effect of external stimulation.

Beta decreases observed during temporal predictions might therefore relate to more than only preparatory processes to an upcoming stimulus. Cross-modal decreases in beta band activity in both the temporal information providing visual as well as the stimulation expecting tactile system might reflect that both sensory modalities are continuously involved in temporal prediction processes, not only in processes preparing for the upcoming stimulation. We found no significant increases in beta power during temporal predictions. Whether decreases in beta power are associated with non-rhythmic temporal predictions while

407   increases might reflect temporal predictions during rhythmic stimulation, remains subject to
408   future research.

Enhanced ITPC cannot be explained by event-related increases in neural activity

In earlier investigations of phase adjustments to external predictive stimulation,
participants were mostly presented with streams of auditory rhythmic input. Rhythmic and
discrete input, however, also causes strongly evoked brain activity within the same frequency
range, which makes it difficult to disentangle streams of evoked activity from entrained
endogenous neural oscillations (Novembre and Iannetti, 2018; Zoefel et al., 2018). Our results
provide evidence that phase alignments of low-frequency fluctuations observed during
temporal predictions cannot solely be explained by stimulus-driven, bottom-up evoked brain
activity (see also, Doelling et al., 2019; Kösem et al., 2018). In the current study, we aimed at
reducing stimulus-evoked brain responses to a minimum by presenting participants with a
continuously moving stimulus instead of several discrete stimuli. We were particularly
interested in the time point at which the stimulus transiently disappeared behind an occluder
(as opposed to sharp onsets and offsets in discrete rhythmic stimulation). At disappearance,
we did not observe an increase in low-frequency power as compared to pre-stimulus baseline
in any of the conditions, which could explain an increase of phase alignments after
disappearance of the stimulus. Moreover, by using an experimental design in which physical
stimulation at disappearance was exactly the same during temporal predictions as well as the
control condition, we controlled for brain responses that could have been driven by bottom-
up, stimulus-processing activity and would therefore not be specific to temporal predictions.
Importantly, delta ITPC, but not power, was stronger during temporal predictions at and after
disappearance of the stimulus, suggesting that delta phase alignments during temporal
predictions cannot be solely related to brain responses evoked by the offset of the visual
movement.

It has been further suggested that a CNV, i.e., activity that is ramping up until the
expected time point is reached, might underlie enhanced phase alignments during temporal
predictions (Breska and Deouell, 2017a). CNVs have often been observed in timing tasks
(e.g., Macar et al., 1999; Pfeuty et al., 2003; Praamstra et al., 2006), and such ramping
activity initialized by temporal predictions would, besides an increase in power, also lead to
increased phase alignments as reflected by enhanced ITPC during temporal predictions. These
increases in activity are therefore not caused by the physical stimulation itself but specifically
related to temporal predictions. As described above, the observed pattern of an early delta
ITPC increase and a late delta power increase in our tactile prediction condition could speak
in favor of a CNV underlying the processes of temporal prediction (see Fig. 3D and 4A).
However, there are several aspects that argue against an involvement of a CNV in our data.

First of all, if in our data a CNV underlay temporal predictions, we should have observed
a late power increase also in the visual temporal prediction task, in which participants also
focused temporal predictions but saw the exact same physical stimulation as in the control
task. Even using uncorrected t-tests, however, we did not observe total delta power
differences between the two conditions in any of the time bins after stimulus disappearance.
Since we see strong ITPC increases in both temporal prediction condition, but a delta power
increase only in the tactile condition, it is unlikely that CNV-like activity would explain the
phase alignments observed in *both* temporal predictions tasks.

Moreover, by subtracting the ERF from each single trial, all activity that is phase-locked
to the disappearance of the stimulus should be removed from the data, that is, all activity
related to a phase-reset of oscillations *as well as* all activity reflecting event-related potentials.
However, also after subtracting the ERF, the strong delta increase in the tactile condition was

still observable. This suggests that the increase in power in the tactile condition was not
associated to the temporal prediction processes locked to the disappearance of the stimulus.
Since the reappearance of the stimulus strongly jittered in relation to the time point of
disappearance and the tactile condition was the only condition in which a sharp-onsetting
tactile stimulus was presented, it is likely that this delta power increase in late windows was
caused by the presentation of the tactile stimulus. In contrast to power, however, delta ITPC
was completely removed after subtracting the ERF, which confirms that subtracting the ERF
reliably removed all disappearance-locked activity.

Further, as stated above, averaging across all trials, i.e., computing the ERF, would
capture all activity from each trial which is locked to disappearance of the stimulus, i.e.,
phase-reset oscillatory activity and/or event-related potentials. In contrast, unlocked activity
should be removed by the averaging. If a CNV caused the late power increase in the tactile
condition, this pattern of a late increase in power should also be observable for the power time
course of the ERF. A phase-reset of oscillatory activity, on the other hand, would rather cause
an ERF power time course that shows differences already at early time windows of the
disappearance, as is the case in our data. The time course of delta power of the ERFs,
therefore, speak against a CNV representing the power increase but, rather, for oscillatory
activity that resets its phase at disappearance.

Therefore, instead of a CNV causing phase alignments of slow fluctuations across trials
(as described above) the opposite might hold, i.e., phase resets of oscillatory activity might
actually, after averaging, lead to results erroneously suggesting a CNV. If so, studies that
observed a CNV after averaging, could have in fact also extracted all oscillatory activity that
has reset its phase after the onset of a temporal cue. Only if a CNV was present in single trial
data, and not only after averaging, such event-related slow fluctuations would indeed relate to
single trial temporal predictions. In our data, however, we did not observe a temporal
prediction related increase in *total* delta power, which is computed on single trial time courses
before averaging. An increase in power was only visible *after* averaging all trials in the time
domain first. Thus, the lack of a power increase in total power together with a CNV-like
power increase after averaging across trials suggests that neural oscillations reset their phase
according to the temporal structure of the stimulation, but did not alter in amplitude on a
single trials basis.

Taken together, we observed strong ITPC differences between the conditions but no
(total) power differences that could be explained by event-related potentials such as a CNV.
Instead of evoked or CNV-like activity, our results therefore suggest that the phase
alignments observed during temporal predictions are associated to neural oscillations that
adjusted their phase to the temporal structure of the stimulation in order to predict the
reappearance of the upcoming stimulation.

Neural oscillations at low frequencies adapt to the temporal structure of non-rhythmic
visual motion stimulation

Earlier studies have observed that neural oscillations entrain towards rhythmic sensory
input to track the low-frequency temporal regularities of the stimulation, especially in the
auditory domain (Giraud and Poeppel, 2012). Such phase entrainment does not only occur in
the delta band but can flexibly adapt to the frequency of the external input also at higher
frequencies such as the theta or the alpha band during auditory stimulation (Doelling and
Poeppel, 2015). However, in the visual system, evidence for the tracking of temporally
predictive input by neural oscillations is not as clear. On the one hand, studies showed that the
phase of neural oscillations is involved in temporal predictions of low-frequency visual input
(Breska and Deouell, 2017a; Cravo et al., 2013; Saleh et al., 2010; Wilsch et al., 2015). On

the other hand, studies suggested that temporal predictions in the visual system were specific
to the alpha band, although sensory input was provided at lower frequencies (Rohenkohl and
Nobre, 2011; Samaha et al., 2015). Rohenkohl and Nobre (Rohenkohl and Nobre, 2011), for
instance, used rhythmically presented visual stimuli at 2.5 and 1.25 Hz moving across the
screen until it disappeared behind an occluder. Nevertheless, neural oscillations exclusively
from the alpha band showed modulated activity associated with temporal predictions during
the disappearance time. They found no phase locking of oscillations in lower frequencies.

In the current study, we provide further evidence that neural oscillations from the delta
band show enhanced phase alignment during visual temporal predictions across trials. In
order to adapt to the temporal regularity of the presented visual stimulus, delta frequencies in
a wide network of parietal and frontal brain areas exerted more consistent phase resets at
around the time point of disappearance of a monotonically moving stimulus as compared to a
luminance matching control condition. The strength of this phase adjustment in each
participant correlated with the consistency in judging a reappearance of the visual stimulus as
too early or too late. This was the case only in the temporal prediction tasks, which underlines
the behavioral relevance of the observed phase adjustments for temporal predictions.

Importantly, our study suggests that the mechanism of phase adjustments for temporal
predictions can be extended to external stimulation that does not as such involve rhythmic or
discontinuous stimulation. We found that low-frequency oscillations can adjust their phase
also to the temporal structure of external stimulation that had to be inferred from uniform
visual motion. This is also in line with recent studies reporting enhanced performance as well
as an involvement of delta phase for non-rhythmic, yet predictable stimulation in the auditory
(Herbst and Obleser, 2019) as well as the visual domain (Breska and Deouell, 2017a; but see
Obleser et al., 2017 and Breska and Deouell, 2017b for a discussion about the rhytmicity of
their non-rhythmic visual stimulation). While both studies involve onsets of discrete stimuli,
they show that delta phase was involved in temporal prediction processes during stimulation
that was not itself purely rhythmic. By showing that the phase of neural oscillations also align
to a rhythm-free, non-discrete, unimodal visual as well as crossmodal visuotactile stimulation,
our results further indicate that the framework of phase adjustments during temporal
predictions might be generalized also to other, if not all, forms of temporally predictive
external stimulation.

Phase resets occurred in a network of frontoparietal and sensory brain areas

We observed enhanced ITPC values in a network of mostly frontal and parietal brain
areas during visual as well as crossmodal temporal predictions. Similarly, Besle et al. (2011)
observed significant phase entrainment to audiovisual stimulation in a wide network of
distributed areas including parietal and inferior frontal areas. These observations support the
notion that brain areas involved in temporal predictions may constitute a frontoparietal timing
network (Coull and Nobre, 2008; Rimmele et al., 2018).

Further, we found enhanced ITPC values also in early somatosensory areas contralateral
to the disappearance of the purely visual stimulus during crossmodal temporal predictions,
despite the fact that prediction-relevant information was provided only by a moving visual
stimulus. This supports evidence reported earlier showing that stimulation within one
modality can crossmodally reset the phase of ongoing low-frequency in other modalities,
which might be an important mechanism for multisensory integration processes (Lakatos et
al., 2007; Mercier et al., 2013).

Similarly, we expected to find enhanced ITPC during temporal predictions in early visual
areas. In fact, increased delta ITPC as compared to baseline were also observed in occipital

sensors (see Fig. S2), but they were not significantly different between the conditions.
However, we found that voxels in early visual areas showed strong correlations between
individual ITPC estimates and the steepness of the psychometric function in both temporal
prediction tasks, but not in the luminance matching task. This suggests that consistent phase
resets of delta oscillations within visual areas might have supported consistent timing
judgments with the participants' subjective timing frameworks. This indicates an involvement
also of the visual system in processes related to temporal prediction.

Moreover, strong correlations between ITPC and behavior were also observed in the
cerebellum, supporting earlier reports on a involvement of the cerebellum in temporal
prediction processes (Breska and Ivry, 2016). Roth and coworkers (Roth et al., 2013), for
instance, found that cerebellar patients were significantly impaired in recalibrating sensory
temporal predictions of a reappearing visual stimulus. This finding is of particular interest as
we adapted the authors' experimental paradigm for the use in the current study. Theirs and
our results therefore indicate that the cerebellum might be crucially involved in accurate and
consistent judgments of temporal regularities deployed in perceiving object motion.

Conclusions

We provide evidence that the phase of neural oscillations can adjust to the temporal
regularities of external stimulation and do not arise as a byproduct of stimulus-driven or
prediction-related evoked potentials. Such phase alignments could provide a key mechanism
that predicts the onset of upcoming events in order to optimize processing of relevant
information and thereby adapt behavior. We show that temporal information provided to one
modality leads to phase adjustments in another modality when crossmodal temporal
predictions are necessary, providing further evidence that such crossmodal phase resets could
be the neuronal basis of multisensory integration processes. Moreover, phase alignments were
observed for unimodal visual as well as crossmodal visuotactile non-rhythmic and non-
discrete stimulation, suggesting a generalizability of phase resets as a mechanism for temporal
predictions to all forms of external stimulation. Taken together, our results provide important
further insights into the neural mechanisms that might be utilized by the brain to predict the
temporal onsets of upcoming events.

Materials and Methods

Participants

Twenty-three healthy volunteers (mean age ± standard deviation (SD): 27.13 ± 4.30 years; 20 females; all right-handed) took part in the study. They gave informed written consent and were monetarily compensated with 13 €/hour for participation. All volunteers had normal or corrected-to-normal vision, normal touch, as well as no background of psychiatric or neurological disorder. The ethics committee of the Medical Association Hamburg approved the study protocol (PV5073), and the experiment was carried out in accordance with the approved guidelines and regulations.

Experimental procedure

The experimental paradigm used in the current study was adopted from an earlier report investigating visual temporal predictions in cerebellar patients (Roth et al., 2013). Our experiment consisted of three conditions: a *visual* temporal prediction task, a crossmodal (*tactile*) temporal prediction task, and a *luminance matching* (control) task. The trials of all conditions started with the presentation of a randomly generated, white noise occluder (size: 7.5° x 11.3° (h x w)) that was smoothed with a Gaussian filter (imgaussfilt.m in MATLAB) and presented in the middle of the screen against a grey background screen (luminance: 44 cd/m^2; corresponds to 115 red-green-blue (RGB) values in our setting; see Figure 1A). At the center of the occluder, a red fixation dot was presented. We instructed participants to fixate this dot throughout the entire trial. After 1500 ms, an oval stimulus (size: 3.5° x 1.0°) set on in the periphery of the screen, moving towards the occluder with a speed of 6.9 °/s. The luminance of the stimulus differed in all trials between 120 to 161 cd/m^2 (6 steps, counterbalanced, corresponds to 170 to 220 RGB). For half of the participants, the stimulus started on the left side of the occluder and moved from left side towards the right side. For the other half, the stimulus started on the right side and moved from right to left. The direction of movement was kept constant for each participant throughout the entire experiment. In each trial, the starting point of the stimulus differed such that the stimulus took 1,000 to 1,500 ms to disappear completely behind the occluder from starting point, randomly jittered with 100 ms (counterbalanced). The size of the occluder and the speed of the stimulus were chosen so that the stimulus would need exactly 1,500 ms to reappear on the other side of the occluder. However, we manipulated the timing and the luminance of the reappearing stimulus. In each trial, the reappearance of the stimulus differed between ±17 to ±467 ms (randomly jittered, but counterbalanced in steps of 50 ms; corresponds to ±1 to ±28 frames with a jitter of 3 frames at 60 Hz) from the correct reappearance time of 1,500 ms. Hence, the stimulus was covered by the occluder for 1,033 to 1,967 ms and was reappearing at 20 different time points. In the visual prediction task as well as in the luminance matching task, we also manipulated the luminance of the reappearing stimulus relative the luminance the stimulus had before disappearance in each trial (jittered, but counterbalanced between ±1 to ±40 cd/m^2, also using 20 different values; corresponds to ±1 to ± 28 RGB in steps of 3 RGB to make it similar to the timing manipulation). After reappearance, the stimulus moved to the other side of the screen for 500 ms with the same speed until it set off the screen. The occluder was presented throughout the entire trial.

By manipulating the timing as well as the luminance in both conditions, we made sure that both, the visual temporal prediction as well as the luminance matching task had the exact equal physical appearance throughout all trials. They only differed in their cognitive set. In

the visual temporal prediction task, we asked participants to judge whether the stimulus was reappearing *too early* or *too late* based on the speed the stimulus had earlier to the occluder (which was kept constant throughout the entire experiment). In the luminance matching task, participants were asked to judge whether the luminance of the reappearing visual stimulus became *brighter* or *darker* as compared to the stimulus earlier to disappearance. Participants answered by pressing one of two buttons with their index or middle finger of the hand contralateral to the reappearing stimulus.

The tactile temporal prediction task was equal to the visual temporal prediction task, with the only difference that a tactile stimulus instead of a visual was presented at the time of reappearance to the right or left index finger (depending on which side the stimulus was expected to reappear behind the occluder). The tactile stimulus was presented by means of a Braille piezostimulator (QuaeroSys, Stuttgart, Germany; 2 x 4 pins, each 1 mm in diameter with a spacing of 2.5 mm), pushing up all eight pins for 200 ms. At that time, nothing happened on the screen. Participants gave their answer with the same hand as in the other two conditions (i.e., with the hand that was not stimulated by the Braille stimulator). Response mapping of the two buttons was counterbalanced across all participants. As soon as participants gave their answer, the fixation dot turned dark grey for 100 ms to indicate that the response was registered. However, participants did not receive trial-wise feedback about the correctness of their response. After a short delay of 200 ms, the white-noise occluder was randomly re-shuffled to signal the start of a new trial.

All three conditions were presented block-wise. At the beginning of each block, participants were informed about the current task. The order of presentation of the conditions was kept constant for each participant, but was randomized across participants (counterbalanced). At the end of each block, they were informed about the overall accuracy of their answers within the last block and were allowed to rest as long as they wanted. Each participant performed two sessions at two different recording days. The experimental procedure was kept constant across both sessions, i.e., movement direction, response mapping, as well as condition order did not change in the second session for individual participants. Each session comprised twelve blocks, i.e., four blocks per condition. Each block consisted of 60 trials, resulting in a total number of 480 trials per condition or 1,440 trials in total. Due to technical difficulties, for one participant we only acquired data from one session with a total number of 720 trials.

At the beginning of each recording day, participants performed a short training of all conditions to get familiar with the overall experimental procedure and the stimulus material. This training took part in the same environment as the subsequent recording session. At the end of the second recording day, participants filled a questionnaire asking for any specific strategy they might have used for the temporal prediction task.

We used MATLAB R2014b (MathWorks, Natick, USA; RRID: SCR_001622) and Psychtoolbox (Brainard, 1997) (RRID: SCR_002881) on a Dell Precision T5500 with Ubuntu 64-bit operating system (Version: 16.04.5 LTS) for stimulus presentation. The visual stimuli were projected onto a matte backprojection screen at 60 Hz with a resolution of $1,920 \times 1,080$ pixels positioned 65 cm in front of participants. To mask the sound of the Braille stimulator during tactile stimulation, we presented participants with auditory pink noise at sampling rate of 48 kHz and volume of 85 dB using MEG-compatible in-ear headphones (SRM-252S, STAX Limited, Fujimi, Japan) during all experimental blocks.

Data acquisition and pre-processing

MEG was recorded at a sampling rate of 1,200 Hz using a 275-channel whole-head system (CTF MEG International Services LP, Coquitlam, Canada) situated in a dimly lit, sound attenuated and magnetically shielded chamber. We additionally recorded electrical eye, muscle and cardiac activity with Ag/AgCl-electrodes in order to have a better estimate for endogenous artefacts. Online head localizations (Stolk et al., 2013) were used to navigate participants back to their original head position prior to the onset of a new experimental block if their movements exceeded five mm from their initial position. The initial head position from the first recording day was saved so that participants could be navigated back to their initial head position also during the second recording day. This assured comparable head positions of each participant across sessions. Five malfunctioning channels were either not recorded or excluded from further analysis for all participants. To further control for eye movement artifacts, eye movements were tracked with an MEG-compatible EyeLink 1000 Long Range Mount system (SR Research, Osgoode, Canada).

We analyzed reaction time data using R (R Core Team, 2014) (RRID: SCR_001905) and RStudio (RStudio Inc., Boston, USA; RRID: SCR_000432). Trials with reaction times longer than three standard deviations were excluded from analysis. Due to the right-skewed nature of reaction times, reaction time data were first log-transformed and then standardized across all trials.

All other data were analyzed using MATLAB R2016b with FieldTrip (Oostenveld et al., 2011) (RRID: SCR_004849), the MEG and EEG Toolbox Hamburg (METH, Guido Nolte; RRID: SCR_016104), or custom made scripts. The physiological continuous recording of each session was first cut into epochs of variable length. Each trial was cut 1,250 ms earlier to stimulus movement onset and 1,250 ms after offset of the reappeared stimulus. Trial length therefore varied between 4,717 and 6,183 ms. To prevent that the timing in a given trial was not exactly as intended, e.g., by short movement interruptions of the stimulus, we removed trials which contained MEG marker timings that differed from the intended timing of the moving stimulus in the trial by at least one frame (17 ms). Thus, we excluded on average 1.2 trials in each participant and each session (range: 0 – 24 trials).

Moreover, trials containing strong muscle artifacts or jumps were detected by semi-automatic procedures implemented in FieldTrip and excluded from analysis. The remaining trials were filtered with a high-pass filter at 0.5 Hz, a low-pass filter at 170 Hz, and three band-stop filters at 49.5–50.5 Hz, 99.5–100.5 Hz and 149.5–150.5 Hz and subsequently down-sampled to 400 Hz.

We performed an independent component analysis (infomax algorithm) to remove components containing eye-movements, muscle, and cardiac artefacts. Components were identified by visual inspection of their time course, variance across samples, power spectrum, and topography. On average, 25.7 ± 8.6 components were rejected in each participant and each session. All trials were again visually inspected and trials containing artefacts that were not detected by the previous steps were removed.

As a final step, using procedures described by Stolk *et al.* (Stolk et al., 2013) and online (http://www.fieldtriptoolbox.org/example/how_to_incorporate_head_movements_in_MEG_a nalysis/) we identified trials in which the head position of the participant differed by 5 mm from the mean circumcenter of the head position from the whole session (on average: 2.6 trials per participant and session, range: 0 – 86 trials) and excluded them from further analysis. 670.2 ± 26.7 trials of the total of 720 trials remained from pre-processing on average per participant in each session.

Quantification and statistical analysis

In the current experiment, we introduced a control condition that was physically identical to our temporal prediction tasks (until reappearance in the tactile condition) in order to account for processes that are not directly related temporal predictions. Hence, for most of our statistical analyses, we were interested in comparing the two temporal prediction tasks with the luminance matching control task, respectively, and not in comparing the two temporal prediction tasks with each other. Therefore, instead of computing an analysis of variance across all three conditions, we directly computed two separate t-tests for the comparison of the visual or the tactile temporal prediction with the luminance matching task, respectively, and accounted for multiple comparisons by adjusting the alpha level.

Psychometric curve

We did not provide participants with feedback about the correctness of their response. Hence, participants responded within their individual framework of a "subjectively correct" reappearance timing or a "subjectively equal" luminance of the stimulus, respectively. To obtain these subjective points of "right-on-time" (ROT) in the temporal prediction tasks or the "points of subjective equality" (PSE) in the luminance matching task, we fitted a psychometric curve to the behavioral data of each participant from all trials in each condition. First, for each timing difference or luminance difference, respectively, we computed the proportion of "too late" or "brighter" answers for each participant. Then, we fitted a binomial logistic regression (psychometric curve) using the glmfit.m and gmlval.m functions provided in MATLAB. The fitted timing or luminance difference value at 50% proportion "too late" or "brighter" answers was determined as ROT or PSE for each participant, respectively. To test for a significant bias towards one of the answers, we tested the ROT or PSE from all participants against zero using one-sample t-tests ($\alpha = .05 / 3 = .017$). The steepness of the psychometric function was computed as the reciprocal of the difference between fitted timing or luminance difference values at 75% and 25% proportion "too late" or "brighter" answers, respectively.

Mixed regression model for reaction times

To test whether reaction times were dependent on the timing difference of the reappearing stimulus, we fitted a random intercept and slope mixed model to reaction times from all trials using the categorial variable *condition* (with the luminance matching task as reference level) and *timing difference* as well as their interaction as fixed effects. Since in the temporal prediction conditions we expected reaction times to be slowest for timing differences around zero and faster for high timing differences, we used a second-order polynomial term for *timing differences*. Subject ID was used as grouping variable to model an individual intercept for each participant, and *timing difference* was modeled with random slope. We used R including the *lme4* package for computing the mixed-effect model, and the package *parameters* to compute p-values using the "Kenward" option, which estimates p-values for fixed effects using the Kenward-Roger approach (Kenward and Roger, 1997).

Spectral power

We decomposed the MEG recordings into time-frequency representations by convolving the data with complex Morlet's wavelets (Cohen, 2014). The recording of each trial and channel was convolved with 40 complex wavelets, logarithmically spaced between 0.5 to 100 Hz. With increasing frequency, the number of cycles for each wavelet logarithmically increased from two to ten cycles. For all analyses of the MEG data, we considered

subjectively correct trials only, i.e., trials in which participants answered correctly based on
their individual ROT. To correct for trial count differences between the tasks, we stratified the
number of trials for each participant for the three different conditions by randomly selecting
as many trials for each condition as the number available from the condition with lowest trial
count.

Since the temporal dependencies between the movement onset, disappearance behind the
occluder and reappearance of the stimulus varied strongly between trials, averaging across
trials would heavily smear the power estimates of the different stages within each trial. To
obtain an estimate of spectral power modulations related to the different events in our
experimental paradigm, we cut each trial further into four separate, partly overlapping
windows (see Figure 2A): a "Baseline" window from -550 to -50 ms earlier to movement
onset; a "Movement" window from -50 to 950 ms relative to the movement onset; a
"Disappearance" window from -350 to 950 ms relative to complete disappearance of the
stimulus behind the occluder; and a "Reappearance" window from -350 to 450 ms relative to
the (first frame) reappearance of the stimulus. Spectral power estimates were then averaged
across all trials belonging to the same condition in each window and binned into time
windows of 100 ms (centered on each full deci-second). All power estimates were normalized
using the pre-stimulus baseline window from -500 to -200 ms earlier to movement onset.

For all statistical analyses on sensor level, we first flipped all sensors of participants, who
saw the stimulus moving from right to left, at the sagittal midline, i.e., the anterior-posterior
axis. This made sure that lateralized activity due to the lateralized stimulation was comparable
across groups. From this on, we considered all participants as if for everyone the stimulus was
moving from the left to the right side. Channels that did not have a counterpart on the
opposite site were excluded from further analyses. In order to obtain an overview of the
spectral power modulations related to the different events within the trials, we then averaged
the power estimates across all channels and conditions (grand average) and tested each time-
frequency pair of the Movement, Disappearance and Reappearance windows against the pre-
stimulus baseline using paired-sample t-tests. We controlled for multiple comparisons by
employing cluster-based permutation statistics as implemented in FieldTrip(Maris and
Oostenveld, 2007). In this procedure, neighboring time-frequency bins with an uncorrected p-
value below 0.05 are combined into clusters, for which the sum of t-values is computed. A
null-distribution is created through permutations of data across participants ($n = 1,000$
permutations), which defines the maximum cluster-level test statistics and corrected p-values
for each cluster. For each window, a separate cluster-permutation test was performed ($\alpha = .05$;
liberally chosen to observe all ongoing power modulations; see Results section).

Since we were most interested in differences between the conditions during the
disappearance time, we subsequently compared the spectral power estimates averaged within
the beta range (13–30 Hz; see Results section) at each time point within the disappearance
window and all channels from the visual or tactile temporal prediction task with the
luminance matching task. We again employed cluster-permutation statistics, this time by
clustering neighboring channels and time points. We used a one-sided $\alpha = .025 / 2 = .0125$,
since negative and positive clusters were tested separately, and to adjust for the two separate
comparisons between the conditions (used throughout the study unless stated differently).

To estimate spectral power in source space, we computed separate leadfields for each
recording session and participant based on each participant's mean head position in each
session and individual magnetic resonance images. We used the single-shell volume
conductor model (Nolte, 2003) with a 5,003 voxel grid that was aligned to the MNI152
template brain (Montreal Neurological Institute, MNI; http://www.mni.mcgill.ca) as
implemented in the METH toolbox. Cross-spectral density (CSD) matrices were computed
from the complex wavelet convolved data in steps of 100 ms in the same time windows as

outlined above. To avoid biases in source projection, common adaptive linear spatial filters
(DICS beamformer (Gross et al., 2001)) pointing into the direction of maximal variance were
computed from CSD matrices averaged across all time bins and conditions for each
frequency.

All time-frequency resolved CSD matrices were then multiplied with the spatial filters to
estimate spectral power in each of the 5,003 voxels and normalized with the pre-stimulus
baseline window. Analogous to sensor space, we first flipped all voxels at the y-axis
(anterior-posterior axis) for the half of participants that saw the stimulus moving from right to
left earlier to further statistical analysis. We then averaged across all time bins within the
disappearance window and utilized cluster-based permutation statistics to identify clusters of
voxels that show statistical difference in beta power between each of the temporal prediction
tasks and the luminance matching task.

Inter-trial phase consistency

We computed ITPC estimates from the complex time-frequency representations obtained
from the wavelet convolution as described in the *Spectral power* section above. In each time
sample and trial, the phase of the complex data was extracted (using the function angle.m in
MATLAB). ITPC was then computed across all subjectively correct and stratified trials
within each of the four time windows in all frequencies as

831
$$ITPC_{tf} = \left| n^{-1} \sum_{r=1}^{n} e^{ik_{tfr}} \right|$$

where n is the number of trials and k the phase angle in trial r at time-frequency point tf
(Cohen, 2014). In other words, ITPC is the length of the mean vector from all phase vectors
with length 1 across all trials at a given time-frequency point. Values for ITPC can vary
between 0 and 1, where 0 means that at a given time-frequency point there is no phase
consistency across trials at all and 1 means all trials show the exact same phase. Similar to
spectral power, we averaged ITPC estimates again in bins of 100 ms and plotted all time
windows averaged across all channels and conditions to obtain a general overview of ITPC
estimates at all events during the trial.

Since we were most interested in ITPC related to stimulus disappearance behind the
occluder, we subsequently computed ITPC in a longer time window from -1,900 ms to 1,900
ms centered around time of complete stimulus disappearance behind the occluder. Thus, we
took advantage of the fact that the onset of other events within each trial, such as the
movement onset and the reappearance of the stimulus, strongly jittered across all trials and
strong contributions of these events to ITPC could thereby be reduced (see Fig. S3). For
statistical analysis, we first averaged ITPC estimates within a frequency band of 0.5 to 3 Hz
(see Results) and then computed cluster-based permutation statistics across all 100 ms time
bins within the 3,800 ms long window and all sensors between each of the temporal
prediction tasks and the luminance matching task.

ITPC on source level was computed using the same leadfields and common beamformer
filters as for spectral power (see above). The complex time-frequency representations
obtained from the wavelet convolution within the 3,800 ms long window on sensor level were
multiplied with the filters to obtain the time-frequency representations in each of the 5,003
voxels. ITPC was computed for each time sample and frequency and then averaged within the
time window showing statistically significant difference between the temporal prediction
tasks and the luminance matching task on sensor level and within the pre-defined frequency
band of 0.5 to 3 Hz. Cluster-based permutation statistics were employed to identify clusters of

voxels showing statistically significant differences in ITPC between the conditions on source
level.

Correlations between condition-wise source level ITPC estimates and the steepness of
each individual's psychometric function were computed using Pearson correlations in each of
the 5,003 voxels within the grid. For this analysis, we averaged ITPC estimates from time
bins of 0 to 1,000 ms with respect to the disappearance of the stimulus within the pre-defined
delta band of 0.5 to 3 Hz. Multiple comparisons were accounted for by using cluster-based
permutation statistics as implemented in FieldTrip ($\alpha = .025 / 3 = .008$)

Delta power control analyses and mixed models

For control analyses of delta power differences between the conditions, we computed
delta power using the same wavelet convolution approach as described for ITPC for the
enlarged time windows between -1,900 ms to 1,900 locked to stimulus disappearance. To
obtain total delta power, we computed power in each single trial first and then averaged
power within the delta band ($0.5 - 3$ Hz) and the respective channels showing the strongest
ITPC effect (see Fig. 3D) for each time bin and condition. Induced power was obtained by
first averaging all trials in each condition and channel in the time domain, i.e., by computing
an ERF in each channel and condition, and then subtracting this average from all single trials
in each channel and condition separately. After subtracting the ERF, power was estimated as
described for total power above. Delta power of the ERF itself was estimated by applying a
wavelet convolution to the ERF, i.e., the average across trials, in each condition and channel
and subsequently averaging power estimates within the delta band and the respective
channels. All time courses were baseline corrected with a pre-disappearance window of -
1,500 to -500 ms relative to disappearance in each condition.

To further examine the effect of delta power on ITPC, we computed random intercept
and random slope mixed-effects models using *condition* and *time* as well as their interaction
as fixed effects for predicting ITPC. One model also included delta power as an additional
covariate, the other one did not. We first averaged delta ITPC as well as delta power ($0.5 - 3$
Hz) from each condition and each time bin (-1.900 ms $- 1.900$ ms) within the channels
showing the strongest effect for ITPC (see Fig. 3D). ITPC as well as the baseline-corrected
power values were standardized across all data for an easier interpretation of the model
estimates. *Subject ID* was used as grouping variable to model an individual intercept for each
participant, and *time* was modeled as random slope. To ensure a flexible relationship between
time and ITPC, we modeled *time* using natural cubic splines with 10 degrees of freedom. For
plotting, we computed ITPC values as predicted by the interaction between *condition* and
time and back-transformed the values to the original scale for an easier evaluation. As for the
reaction time model, we used R including the *lme4* package for computing the mixed-effect
model, the package *parameters* to compute p-values using the "Kenward" option, as well as
the package *splines* for generating the natural cubic splines.

Acknowledgements

We thank Florian Göschl, Tessa Rusch, Marina Fiene and Guido Nolte for valuable
discussions. This work was funded by grants from the DFG (SFB TRR 169/B1 and SFB
936/A3 to A.K.E.).

Author contributions

Conceptualization, J.D., A.K.E, P.W., A.M.; Methodology, J.D., A.K.E.; Software, J.D.;
Formal Analysis, J.D.; Investigation, J.D.; Writing – Original Draft, J.D.; Writing – Review
& Editing, A.K.E., P.W., A.M., D.Z.; Visualization, J.D.; Funding Acquisition, A.K.E.;
Supervision, A.K.E.; Project Administration, A.K.E., D.Z.; Resources, A.K.E.

Competing interests

The authors declare no competing interests.

Data availability

Custom scripts and data will be made available upon full submission of the manuscript.
909

References

Arnal, L.H., and Giraud, A.L. (2012). Cortical oscillations and sensory predictions. Trends Cogn. Sci. *16*, 390–398. doi: 10.1016/j.tics.2012.05.003.

Arnal, L.H., Doelling, K.B., and Poeppel, D. (2015). Delta-beta coupled oscillations underlie temporal prediction accuracy. Cereb. Cortex *25*, 3077–3085. doi: 10.1093/cercor/bhu103.

Besle, J., Schevon, C.A., Mehta, A.D., Lakatos, P., Goodman, R.R., McKhann, G.M., Emerson, R.G., and Schroeder, C.E. (2011). Tuning of the human neocortex to the temporal dynamics of attended events. J. Neurosci. *31*, 3176–3185. doi: 10.1523/JNEUROSCI.4518-10.2011.

Brainard, D.H. (1997). The psychophysics toolbox. Spat. Vis. *10*, 433–436.

Breska, A., and Deouell, L.Y. (2017a). Neural mechanisms of rhythm-based temporal prediction: delta phase-locking reflects temporal predictability but not rhythmic entrainment. PLOS Biol. *15*, e2001665. doi: 10.1371/journal.pbio.2001665.

Breska, A., and Deouell, L.Y. (2017b). Dance to the rhythm, cautiously: isolating unique indicators of oscillatory entrainment. PLOS Biol. *15*, e2003534. doi: 10.1371/journal.pbio.2003534.

Breska, A., and Ivry, R.B. (2016). Taxonomies of timing: where does the cerebellum fit in? Curr. Opin. Behav. Sci. *8*, 282–288. doi: 10.1016/j.cobeha.2016.02.034.

Buzsáki, G. (2006). Rhythms of the brain (New York: Oxford University Press.).

Cohen, M. (2014). Analyzing neural time series data: theory and practice (Cambridge: MIT Press).

Coull, J.T., and Nobre, A.C. (2008). Dissociating explicit timing from temporal expectation with fmri. Curr. Opin. Neurobiol. *18*, 137–144. doi: 10.1016/j.conb.2008.07.011.

Cravo, A.M., Rohenkohl, G., Wyart, V., and Nobre, A.C. (2013). Temporal expectation enhances contrast sensitivity by phase entrainment of low-frequency oscillations in visual cortex. J. Neurosci. *33*, 4002–4010. doi: 10.1523/JNEUROSCI.4675-12.2013.

Doelling, K.B., and Poeppel, D. (2015). Cortical entrainment to music and its modulation by expertise. Proc. Natl. Acad. Sci. *112*, E6233–E6242. doi: 10.1073/pnas.1508431112.

Doelling, K.B., Assaneo, M.F., Bevilacqua, D., Pesaran, B., and Poeppel, D. (2019). An oscillator model better predicts cortical entrainment to music. Proc. Natl. Acad. Sci. 201816414. doi: 10.1073/pnas.1816414116.

van Ede, F., de Lange, F., Jensen, O., and Maris, E. (2011). Orienting attention to an upcoming tactile event involves a spatially and temporally specific modulation of sensorimotor alpha- and beta-band oscillations. J. Neurosci. *31*, 2016–2024. doi: 10.1523/JNEUROSCI.5630-10.2011.

Engel, A.K., Fries, P., and Singer, W. (2001). Dynamic predictions: oscillations and synchrony in top-down processing. Nat. Rev. Neurosci. *2*, 704–716. doi: 10.1038/35094565.

Fries, P. (2005). A mechanism for cognitive dynamics: neuronal communication through neuronal coherence. Trends Cogn. Sci. *9*, 474–480. doi: 10.1016/j.tics.2005.08.011.

Giraud, A.-L., and Poeppel, D. (2012). Cortical oscillations and speech processing: emerging computational principles and operations. Nat. Neurosci. *15*, 511–517. doi: 10.1038/nn.3063.

Gomez-Ramirez, M., Kelly, S.P., Molholm, S., Sehatpour, P., Schwartz, T.H., and Foxe, J.J.
(2011). Oscillatory sensory selection mechanisms during intersensory attention to
rhythmic auditory and visual inputs: a human electrocorticographic investigation. J.
Neurosci. *31*, 18556–18567. doi: 10.1523/JNEUROSCI.2164-11.2011.

Gould, I.C., Rushworth, M.F., and Nobre, A.C. (2011). Indexing the graded allocation of
visuospatial attention using anticipatory alpha oscillations. J. Neurophysiol. *105*, 1318–
1326. doi: 10.1152/jn.00653.2010.

Gross, J., Kujala, J., Hamalainen, M., Timmermann, L., Schnitzler, A., and Salmelin, R.
(2001). Dynamic imaging of coherent sources: studying neural interactions in the human
brain. Proc. Natl. Acad. Sci. *98*, 694–699. doi: 10.1073/pnas.98.2.694.

Herbst, S.K., and Obleser, J. (2019). Implicit temporal predictability enhances pitch
discrimination sensitivity and biases the phase of delta oscillations in auditory cortex.
Neuroimage *203*, 116198. doi: 10.1016/j.neuroimage.2019.116198.

Kenward, M.G., and Roger, J.H. (1997). Small sample inference for fixed effects from
restricted maximum likelihood. Biometrics doi: 10.2307/2533558.

Kösem, A., Bosker, H.R., Takashima, A., Meyer, A., Jensen, O., and Hagoort, P. (2018).
Neural entrainment determines the words we hear. Curr. Biol. *28*, 2867-2875.e3. doi:
10.1016/j.cub.2018.07.023.

Lakatos, P., Chen, C.-M., O'Connell, M.N., Mills, A., and Schroeder, C.E. (2007). Neuronal
oscillations and multisensory interaction in primary auditory cortex. Neuron *53*, 279–292.
doi: 10.1016/j.neuron.2006.12.011.

Lakatos, P., Karmos, G., Mehta, A.D., Ulbert, I., and Schroeder, C.E. (2008). Entrainment of
neuronal oscillations as a mechanism of attentional selection. Science *320*, 110–113. doi:
10.1126/science.1154735.

Macar, F., Vidal, F., and Casini, L. (1999). The supplementary motor area in motor and
sensory timing: evidence from slow brain potential changes. Exp. Brain Res. *125*, 271–
280. doi: 10.1007/s002210050683.

Maris, E., and Oostenveld, R. (2007). Nonparametric statistical testing of EEG- and MEG-
data. J. Neurosci. Methods *164*, 177–190. doi: 10.1016/j.jneumeth.2007.03.024.

Mercier, M.R., Foxe, J.J., Fiebelkorn, I.C., Butler, J.S., Schwartz, T.H., and Molholm, S.
(2013). Auditory-driven phase reset in visual cortex: human electrocorticography reveals
mechanisms of early multisensory integration. Neuroimage *79*, 19–29. doi:
10.1016/j.neuroimage.2013.04.060.

Nolte, G. (2003). The magnetic lead field theorem in the quasi-static approximation and its
use for magnetoencephalography forward calculation in realistic volume conductors.
Phys. Med. Biol. *48*, 3637–3652. doi: 10.1088/0031-9155/48/22/002.

Novembre, G., and Iannetti, G.D. (2018). Tagging the musical beat: neural entrainment or
event-related potentials? Proc. Natl. Acad. Sci. *115*, E11002–E11003. doi:
10.1073/pnas.1815311115.

Obleser, J., Henry, M.J., and Lakatos, P. (2017). What do we talk about when we talk about
rhythm? PLOS Biol. *15*, e2002794. doi: 10.1371/journal.pbio.2002794.

Oostenveld, R., Fries, P., Maris, E., and Schoffelen, J.-M. (2011). FieldTrip: Open source
software for advanced analysis of MEG, EEG, and invasive electrophysiological data.
Comput. Intell. Neurosci. *2011*, 1–9. doi: 10.1155/2011/156869.

Pfeuty, M., Ragot, R., and Pouthas, V. (2003). When time is up: cnv time course

differentiates the roles of the hemispheres in the discrimination of short tone durations.
Exp. Brain Res. *151*, 372–379. doi: 10.1007/s00221-003-1505-6.

Praamstra, P., Kourtis, D., Kwok, H.F., and Oostenveld, R. (2006). Neurophysiology of
implicit timing in serial choice reaction-time performance. J. Neurosci. *26*, 5448–5455.
doi: 10.1523/JNEUROSCI.0440-06.2006.

R Core Team (2014). R: a language and environment for statistical computing. R Found. Stat.
Comput. Vienna Austria.

Rimmele, J.M., Morillon, B., Poeppel, D., and Arnal, L.H. (2018). Proactive sensing of
periodic and aperiodic auditory patterns. Trends Cogn. Sci. *22*, 870–882. doi:
10.1016/j.tics.2018.08.003.

Rohenkohl, G., and Nobre, A.C. (2011). Alpha oscillations related to anticipatory attention
follow temporal expectations. J. Neurosci. *31*, 14076–14084. doi:
10.1523/JNEUROSCI.3387-11.2011.

Roth, M.J., Synofzik, M., and Lindner, A. (2013). The cerebellum optimizes perceptual
predictions about external sensory events. Curr. Biol. *23*, 930–935. doi:
10.1016/j.cub.2013.04.027.

Saleh, M., Reimer, J., Penn, R., Ojakangas, C.L., and Hatsopoulos, N.G. (2010). Fast and
slow oscillations in human primary motor cortex predict oncoming behaviorally relevant
cues. Neuron *65*, 461–471. doi: 10.1016/j.neuron.2010.02.001.

Samaha, J., Bauer, P., Cimaroli, S., and Postle, B.R. (2015). Top-down control of the phase of
alpha-band oscillations as a mechanism for temporal prediction. Proc. Natl. Acad. Sci.
112, 8439–8444. doi: 10.1073/pnas.1503686112.

Schroeder, C.E., and Lakatos, P. (2009). Low-frequency neuronal oscillations as instruments
of sensory selection. Trends Neurosci. *32*, 9–18. doi: 10.1016/j.tins.2008.09.012.

Stefanics, G., Hangya, B., Hernádi, I., Winkler, I., Lakatos, P., Ulbert, I., Hernadi, I.,
Winkler, I., Lakatos, P., and Ulbert, I. (2010). Phase entrainment of human delta
oscillations can mediate the effects of expectation on reaction speed. J. Neurosci. *30*,
13578–13585. doi: 10.1523/JNEUROSCI.0703-10.2010.

Stolk, A., Todorovic, A., Schoffelen, J.-M., and Oostenveld, R. (2013). Online and offline
tools for head movement compensation in MEG. Neuroimage *68*, 39–48. doi:
10.1016/j.neuroimage.2012.11.047.

VanRullen, R. (2016). Perceptual cycles. Trends Cogn. Sci. *20*, 723–735. doi:
10.1016/j.tics.2016.07.006.

Wilsch, A., Henry, M.J., Herrmann, B., Maess, B., and Obleser, J. (2015). Slow-delta phase
concentration marks improved temporal expectations based on the passage of time.
Psychophysiology *52*, 910–918. doi: 10.1111/psyp.12413.

Zoefel, B., ten Oever, S., and Sack, A.T. (2018). The involvement of endogenous neural
oscillations in the processing of rhythmic input: more than a regular repetition of evoked
neural responses. Front. Neurosci. *12*, 1–13. doi: 10.3389/fnins.2018.00095.

1037