Allelic polymorphism shapes community function in evolving *Pseudomonas*

***aeruginosa* populations**

Sheyda Azimi[a,b], Aled E. L. Roberts[c], Shengyun Peng[b], Joshua S. Weitz[a,b,c], Alan McNally[e],

Samuel P. Brown[a,b] and Stephen P. Diggle[a,b*]

[a] Center for Microbial Dynamics & Infection, Georgia Institute of Technology, Atlanta,

U.S.A.; [b]School of Biological Sciences, Georgia Institute of Technology, Atlanta, U.S.A.; [c]

Microbiology & Infectious Diseases Group, Institute of Life Science, Swansea University

Medical School, Swansea, U.K.; [d] School of Physics, Georgia Institute of Technology,

Atlanta, U.S.A; [e] Institute of Microbiology and Infection, College of Medical and Dental

Sciences, University of Birmingham, Birmingham, U.K.

*Correspondence: stephen.diggle@biosci.gatech.edu

Keywords: diversity, quorum sensing, *Pseudomonas aeruginosa*, community function

Abstract

Pseudomonas aeruginosa is an opportunistic pathogen that chronically infects the lungs of

individuals with cystic fibrosis (CF) by forming antibiotic resistant biofilms. Emergence of

phenotypically diverse isolates within CF *P. aeruginosa* populations has previously been

reported, however, the impact of heterogeneity on social behaviors and community function

is poorly understood. Here we describe how this heterogeneity impacts on behavioral traits

by evolving the strain PAO1 in biofilms grown in a synthetic sputum medium for 50 days.

We measured social trait production and antibiotic tolerance and used a metagenomic

approach to analyze and assess genomic changes over the duration of the evolution

experiment. We found that (i) evolutionary trajectories were reproducible in independently

evolving populations; (ii) over 60% of genomic diversity occurred within the first 10 days of

selection. We then focused on quorum sensing (QS), a well-studied *P. aeruginosa* trait that is

commonly mutated in strains isolated from CF lungs. We found that at the population level

(i) evolution in sputum medium selected for decreased production of QS and QS-dependent

traits; (ii) there was a significant correlation between *lasR* mutant frequency, the loss of

protease and the 3O-C12-HSL signal, and an increase in resistance to clinically relevant β-

lactam antibiotics, despite no previous antibiotic exposure. Overall, our findings provide

insights into the effect of allelic polymorphism on community functions in diverse *P.*

aeruginosa populations. Further, we demonstrate that *P. aeruginosa* population and

evolutionary dynamics can impact on traits important for virulence and can lead to increased

tolerance to β-lactam antibiotics.

Introduction

The cystic fibrosis (CF) lung is a spatially complex and inflamed environment that provides

beneficial growth conditions for a number of bacterial species, including the opportunistic

pathogen *Pseudomonas aeruginosa* [1-5]. Chronic infection of CF lungs with highly adapted

and antibiotic resistant biofilms of *P. aeruginosa* is a major cause of lung function decline,

which results in a concomitant increase in morbidity and mortality in individuals with CF [3,

4, 6-11]. Longitudinal studies of chronic CF infections with *P. aeruginosa* have revealed that

patients become infected at a young age with an environmental or transmissible isolate that

evolves and adapts over time to the lung environment [12-18]. Studies on *P. aeruginosa*

populations isolated from individual lungs, have demonstrated divergent evolution, resulting

in heterogeneous populations of *P. aeruginosa* within patients [16, 17, 19]. This genetic

adaptation and diversification is likely to impact on levels of pathogenicity and the efficacy of

antibiotic treatment [16, 20], and could potentially impact how other species of microbe

colonize the CF lung [4, 21-27].

Studies on explanted CF lungs have shown that the spatial structure found within lungs, and

physical separation of infecting isolates, plays a role in generating the vast phenotypic and

genotypic heterogeneity seen within *P. aeruginosa* populations in individual patients [16, 17,

28, 29]. Major adaptations of *P. aeruginosa* to the CF lung include alginate production, loss of

quorum sensing (QS), hypermutability and increased resistance to antimicrobials [7, 8, 14, 30,

31]. Heterogeneity in *P. aeruginosa* populations has also been explained by early divergent

evolution and adaptation to differential ecological niches [32] and recombination between

isolates residing in the airways [16, 33], however, the exact mechanisms leading to

heterogeneity have yet to be fully elucidated. Further, whilst it is accepted that genomic

heterogeneity arises in CF chronic lung infections, it remains unknown how genotypic changes

shape community functions within whole *P. aeruginosa* populations. Understanding what

drives community structure and function remains a key goal in microbial ecology, because

overall community function is determined by all the individuals in the population.

In this study, we hypothesized that the combination of wild-type and mutated alleles in

populations shapes community functions, which could result in clinically-relevant outcomes

such as increased or decreased antibiotic tolerance and traits important for virulence. To test

this, we first evolved *P. aeruginosa* PAO1 in biofilms on plastic beads [34] for up to 50 days.

This bead biofilm system has previously been successfully used to study genetic adaptation

and phenotypic diversity of *Burkholderia cenocepacia* and *P. aeruginosa* to different

environmental conditions [35]. Our study differs from this previous work, in that our major

focus was on community function within populations rather than on specific isolates and

defining ecological niches. We grew our bead associated biofilms in a synthetic CF sputum

media (SCFM), which recapitulates the chemical environment found in CF sputum [36-38].

Our study therefore generated phenotypically and genotypically heterogeneous populations of

P. aeruginosa in a spatially structured environment chemically relevant to CF sputum. We

utilized these heterogenous evolved populations to study the changes in the functional

community phenotypes instead of the traditional approach of working with single evolved

isolates. We used a metagenomic approach to assess genetic alterations within evolving

populations, and we monitored fluctuations in allele frequency during the selection process. To

determine the impact of genomic heterogeneity within populations on various phenotypes, we

assessed collective phenotypic traits (community function) of the evolved populations.

One of the most commonly described adaptations of *P. aeruginosa* to the CF lung is the loss

of the *las* QS system, predominantly through point mutations, frameshifts and deletions in the

lasR gene [17, 39-41]. We used our evolved populations to specifically focus on the impact of

lasR mutation frequency on QS phenotypes. The *lasR* gene encodes the LasR transcriptional

regulator, which binds the QS signal *N*-(3-oxo-dodecanoyl)-L-homoserine lactone (3O-C12-

HSL) [42-45]. LasR-bound 3O-C12-HSL controls the transcription of approximately 10 % of

the *P. aeruginosa* genome, including a number of genes involved in social behaviors,

pathogenesis, antibiotic resistance and biofilm formation [46-49]. Despite a number of

previous studies which describe the changes and adaptation of various lineages of *P.*

aeruginosa in CF lungs [14, 50-53], it remains unclear how polymorphisms in the *lasR* gene

impacts on community function within evolving heterogenous *P. aeruginosa* populations. This

is because most previous studies that focused on within host adaptation of *P. aeruginosa* used

single colonies isolated from temporal CF sputum samples and not whole populations [12, 14,

15, 19, 54].

Overall we found that (i) evolutionary trajectories were reproducible between independently

evolving populations and that over 60% of genomic changes in populations occurred within

the first 10 days of selection; (ii) after 30 days of evolution in SCFM, the evolved communities

displayed an increase in *lasR* mutant frequency and a decrease in QS-dependent traits; (iii)

there was a significant correlation between *lasR* mutant frequency, the loss of social traits and

an increase in tolerance to β-lactam antibiotics. Our findings provide insights into how allelic

polymorphism and population heterogeneity in general, can impact on phenotypes and

community functions within evolving *P. aeruginosa* populations. Further, we demonstrate that

changes in *P. aeruginosa* population dynamics can alter factors associated with virulence and

provide explanations for increased antibiotic tolerance, even in situations when antibiotics have

not been used.

Results

Genomic variation in evolving biofilm populations over 50 days selection in SCFM. We

evolved the *P. aeruginosa* strain PAO1 for 50 days ($\approx$ 800 generations) in biofilms using a

previously described biofilm bead method [34], and a growth medium that chemically mimics

CF sputum (SCFM), and where the physiology of *P. aeruginosa* is similar to when grown in

human sputum [36-38]. Our experimental evolution approach contained four-independent

replicate lines (Fig. S1). We collected and stored biofilm evolved populations after 10, 20, 30,

40 and 50 days of evolution (Rounds 1 to 5: R1 – R5). We used the Illumina MiSeq platform

to deep sequence evolved populations in order to determine genomic changes through time.

We also sequenced our laboratory PAO1 ancestral strain, and after *de novo* genome assembly

of this strain, we mapped the sequence reads of the evolved populations to the ancestor in order

to detect SNPs [55]. Our SNP calling analysis, combined with an analysis of allele frequency,

revealed that in all four independent replicate lines, an average of 282 ± 13 SNPs occurred in

the populations after 10 days of selection (Fig. 1A). We found that around 60 % of these SNPs,

were present through all other rounds of selection (Fig. 1B; Fig. S2). This suggests that the

evolutionary trajectories of biofilm growth in SCFM are similar in independently evolving

populations, and that the major genetic heterogeneity in evolving populations occurs during

the early phases of selection.

SNP frequency in genes involved in social traits fluctuates over time. In our evolution

experiment, we found emergence of polymorphisms in 45 genes involved in various

physiological functions (Fig. S3; Table S1). We found that between 10-25% of SNPs were

fixed (frequency of 1) in the populations over 50 days of selection across all four replicate

evolution lines (Fig. S4). When we focused on the allele frequency and not the number of

positions altered in each coding region, we found that the frequency of SNPs in genes involved

in different traits changed during the course of the experiment (Fig. 2; Fig. S3; Fig. S4 and Fig.

S5).   The genes highlighted in Fig. 2 are genes that have previously been shown to be

commonly mutated in *P. aeruginosa* CF isolates [9, 15, 16, 29, 30, 56].

The allele frequencies of nonsynonymous SNPs in *ccoN2* (PA1557), and synonymous SNPs

in *pvdJ* (PA2400) and *tufA* (PA4265) became fixed in the population at a frequency of 1, while

the frequency of nonsynonymous SNPs in *phzC2* (PA1901) and *pvdD* (PA2399) fluctuated

between 0.4-0.5 in different rounds of selection. We detected a number of SNPs occurring in

mutS (PA3620), *pilQ* (PA5040), *pilN* (PA5043) at 10, 20 and 30 days of selection in all four

independent evolved lines. We observed an increase in *lasR* (PA1430) mutant allele frequency

between 30-40 days of selection (Fig. 2; Fig. S3 and Fig. S5).

Accumulation of SNPs shapes community functions in evolved *P. aeruginosa* populations.

We next examined the production of phenotypic social traits in evolved populations in order to determine changes in community function of the genetically heterogenous evolving populations. We measured levels of biofilm formation, QS signals, total protease and the siderophores pyoverdine and pyochelin. We observed a small but significant increase in biofilm formation by evolved populations when compared to the PAO1 ancestor (Fig. 3A). After 30 days of selection, production of total protease (Fig. 3B) and the 3O-C12-HSL QS signal (Fig. 3C) decreased in evolved populations, however, the levels of C4-HSL signal (Fig. 3C), and pyochelin and pyoverdine (Fig. 3D) did not follow this trend, and any changes were generally not significantly different from the values of the ancestor strain. We also observed an increase in colony morphology types (morphotypes) in the evolved populations starting after 20 days of selection (Fig. S6).

To determine whether the emergence and increase in frequency of SNPs in the populations impact upon community function, we used a linear regression model. We focused on changes in the allele frequency detected in the QS regulator LasR. We chose *lasR* because the genes and phenotypes it regulates in *P. aeruginosa* are well understood [30, 40, 44, 47, 57] and also, *lasR* mutants are regularly isolated from CF sputum [13, 50, 58]. We detected a non-synonymous SNP (V208G) in the DNA binding domain of LasR between 30-40 days of selection (Fig. S5). Interestingly, SNPs in the same position (V208) have been identified in *P. aeruginosa* isolates collected from CF sputum samples [40]. V208 is adjacent to the D209 residue in the LasR DNA-binding domain [59, 60], suggesting an impact on the structure of LasR and its DNA binding affinity. We assessed the impact of *lasR* V208G SNP frequency on changes on two QS-dependent phenotypic traits; production of 3O-C12-HSL signal and total protease. We found a significant negative correlation between the frequency of the V208G *lasR*

SNP in the whole-evolved populations and the total protease activity of evolved populations.

We found that 87% (R^2=0.8704, F=20.15; p=0.0206) of the decreased protease activity was

correlated with the accumulation of the *lasR* mutation in the populations (Fig. 4A). We used

the same analysis to determine whether *lasR* mutant accumulation impacts on 3O-C12-HSL

production within populations. We found that only 53% of the decreased 3O-C12-HSL levels

(R^2=0.5363, F=3.469; p=0.1594) can be explained by the accumulation of *lasR* mutation in the

populations (Fig. 4B). However, when we only included the last 30 days of selection in our

analysis, the decrease in 3O-C12-HSL levels could be fully (100%) explained by the

accumulation of *lasR* mutants (R^2=1.0, p=0.0034) (Fig. 4B). Our analysis did not show any

correlation between the frequency of the *lasR* SNP and changes in the production of C4-HSL

(Fig. 4C) or biofilm formation (Fig. 4D).

Accumulation of *lasR* mutants in evolved populations leads to increased tolerance to β-

lactam antibiotics. Previously it has been shown that *lasR* mutants display increased β-

lactamase activity, and therefore increased resistance to β-lactam antibiotics (such as

ceftazidime) [61] which are routinely used in CF clinics. To determine possible links between

the loss of *lasR* function and changes in tolerance to routinely used antibiotics, we first assessed

the antimicrobial susceptibility levels of the evolved populations. When we tested levels of

antimicrobial susceptibility to six routinely used antibiotics for chronic CF lung infection, after

30 days of selection the evolved populations showed an increased tolerance (indicated by a

decrease in the zone of inhibition) to three antibiotics; ceftazidime, piperacillin/tazobactam and

meropenem which are all β-lactam class antibiotics (Fig. 5; Fig. S7). We then tested for

correlations between frequencies of the *lasR* V208G SNP and resistance to ceftazidime and

piperacillin/tazobactam. We observed a positive and significant correlation between

accumulation of *lasR* mutants and the increased resistance to both ceftazidime and

piperacillin/tazobactam (Fig. 5A and B). We also tested whether the increase in tolerance could

be explained by an increase in biofilm formation. We found there was no significant

correlation, suggesting that increases in biofilm formation do not necessarily translate to an

increase in drug tolerance (Fig. S7C).

To confirm that our evolved populations contained *lasR* mutants with increased β-lactam

tolerance and that *lasR* frequency affects community function, we first selected and sequenced

an evolved isolate (BB8) from the final round of selection (50 days) with no protease activity

and which demonstrated increased antibiotic tolerance compared to the ancestor PAO1 (Fig.

5C). The sequencing of isolate BB8 revealed the same SNP in the *lasR* gene that we detected

in the evolved populations (V208G). We complemented BB8 with an intact copy of the PAO1

lasR gene *in trans* on a plasmid and then assessed β-lactam tolerance levels. We observed a

decrease in tolerance to ceftazidime and piperacillin/tazobactam in the *lasR* complemented

BB8 isolate (Fig. 5C). Finally we tested whether changes in the starting frequency of *lasR*

mutants in mixed populations impacted on community functions. We found that when BB8

was mixed with wild type cells, this resulted in an increased tolerance to ceftazidime (Fig. 5C)

and a reduction in 3O-C12-HSL but not C4-HSL in mixed populations (Fig. S8).

Discussion

Despite a number of recent studies focused on adaptive changes in evolved populations of *P.*

aeruginosa in environments designed to mimic CF sputum [35, 62-66], there remain significant

gaps in knowledge about how genomic and phenotypic diversity impacts upon on interactions

within populations and on community-level phenotypes. In our current study, we focused on

understanding how the evolution and coexistence of multiple lineages of *P. aeruginosa* shape

community functions. To generate diversity, we performed a 50-day selection experiment,

evolving PAO1 as biofilms in an artificial sputum medium (SCFM) [36-38]. We observed (i)

a rise in genetic diversity after 10 days of selection in biofilms grown in SCFM; (ii) up to 25%

of SNPs became fixed in the population and carried in evolved populations through 50 days of

selection; (iii) emergence and accumulation of SNPs in genes involved in motility, respiration,

DNA mismatch repair, transcription and QS emerged between 10 and 40 days of selection and

(iv) accumulation of *lasR* SNPs correlated with decreased protease activity, 3O-C12-HSL

production and increased tolerance to β-lactam antibiotics despite no prior treatment with any

antibiotic.

The divergent evolution of *P. aeruginosa* during chronic infection of CF lungs has been the

focus of numerous studies on longitudinal collections of sputum samples. Analysis of single *P.*

aeruginosa isolates from these collections has identified genomic signatures for adaptation to

the CF lung environment [9, 14, 29, 50, 53, 67, 68]. Similar studies on collections of *P.*

aeruginosa isolates sourced from single sputum samples, have shown considerable phenotypic

and genetic diversification of *P. aeruginosa* strains which group into clades [16, 17], and which

may colonize different ecological niches in CF lungs [28, 69]. All of these studies focused on

the diversification, adaptation and heterogeneity of *P. aeruginosa* during chronic CF lung

infection. However, the impact that this heterogeneity may have on collective phenotypes has

been largely over-looked. We hypothesized that the combination of wild-type and mutated

alleles in populations shapes collective phenotypes. We observed that the accumulation and

frequency of various SNPs in heterogenous populations of *P. aeruginosa,* significantly impacts

the collective phenotypes of evolved populations (Fig. 2; Fig. 3). Although we observed that

the majority of genomic changes occurred during the first 10 days of selection, the functional

outcome was only impactful after 20-30 days, where we observed an increase in biofilm

formation and colony morphotypes and a significant decrease in total protease activity and

production of 3O-C12-HSL by evolved populations.

The biofilm bead system has previously been used to examine the evolution of *P. aeruginosa*

in biofilms [35], although there are significant differences between this and our study. First,

the environmental conditions were different, with the previous study growing in a minimal

media (M63) whilst we used SCFM. Secondly, we evolved PAO1 whereas the previous study

evolved PA14 [35]. Whilst both are popular model strains for *P. aeruginosa* study, they differ

in a number of factors including virulence [70]. Third, we focused primarily on the emergent

properties of the whole community rather than individual isolates, which we show has

important implications for signaling and antimicrobial tolerance. Finally, the previous study

studied the emergence of *mutL* and *mutS* genes which may have contributed to the emergence

of different individual morphotypes [35]. We observed low frequency mutations in the *mutS*

gene after 30 days (Fig. S3). As these were synonymous mutations, they did not appear to

significantly increase mutation rate after 30 days of selection.

Our study also shows similarities with another recent study. Here the authors evolved PA14 in

SCFM supplemented with mucin to create a structured environment [62]. The main findings

of this study was that mucin promoted diversification of *P. aeruginosa* between and not within

populations. The authors found increases in tolerance to antibiotics in evolved populations and

changes in growth, motility, pyocyanin production and biofilm formation. They did not present

genomic evidence that underpinned these changes [62].

Our study identified genomic changes often observed in *P. aeruginosa* isolates taken from CF

lungs including changes in secretion genes (*hcpA, hcpB, vgrG2a, vgrG4b*), motility genes (*pilQ*

and *pilN*), iron aquisition genes (*pvdD, pvdJ*) and biofilm formation (*cdrA*). We also observed

a consistent change in all lines and rounds of selection in anaerobic respiratory pathways (Fig.

2; Fig. S3). *ccoN2* (PA1557) encodes the cytochrome C oxidase subunit (ccb3 type). Increased

expression of *ccoN2* in small colony variant has been previously noted during microaerophilic

conditions [71, 72].

One of best-known signatures of *P. aeruginosa* adaptation to the CF lung environment, is

mutation in the QS regulator LasR. Several studies on *P. aeruginosa* strains isolated from CF

sputum samples, found significant genomic changes in *lasR,* including truncation, deletion,

frame shifts and SNPs, with many resulting in a loss of function [30, 40, 41, 50, 73]. Although

there can be a high percentage of *lasR*-deficient isolates collected from patients, studies have

also shown differential frequencies of functional and intact *lasR* alleles within patients [13,

74]. Considering the importance of *lasR*-dependent social interactions for the fitness of *P.*

aeruginosa, mutation-frequency of *lasR* within populations of *P. aeruginosa* could

significantly impact on QS-dependent phenotypes and fitness at the population level [75-77].

Previously we have demonstrated that an increase in the frequency of QS social cheats (*lasR*

mutants) in defined populations of wild-type and *lasR* mutants, leads to a reduction in

cooperation and virulence in mouse models of infection [75]. These studies showed that even

simple mixed genotype populations of *P. aeruginosa* can have a significant impact on

community function, virulence and the outcome of infection.

In our current study, we observed a negative correlation between an increase in the frequency

of the *lasR* V208G SNP in evolved populations and levels of protease activity and 3O-C12-

HSL production (Fig. 4). Mutation of *lasR* in *P. aeruginosa* isolates from CF lungs has also

previously been shown to be important for increased tolerance to β-lactam antibiotics such as

ceftazidime [19, 56, 61]. Here we examined whether the increase of *lasR* mutants in our

evolved populations had an impact on levels of antimicrobial tolerance. Despite no prior

treatments with antibiotics, we observed an increased tolerance to three antibiotics from the β-

lactam family (Fig. 5). In contrast, we found no correlation between increased biofilm

production by evolved populations and antibiotic resistance (Fig. S7). Resistance of *P.*

aeruginosa to antibiotics in chronic infections such as CF or wounds, is often thought to be

due to specific mechanisms such as efflux pumps or via production of excess polysaccharides

such as alginate. Our findings suggest that the accumulation and frequency of genetic variants

that might not traditionally be associated with resistance to drugs (e.g. QS mutants) within a

heterogenous population, can alter phenotypes within populations that can result in important

clinical repercussions.

Our work suggests that in the future, we should consider metagenomic and metaphenotypic

assessments of *P. aeruginosa* populations collected from CF patients, rather than focusing on

single colonies. This is because the phenotype of populations is dictated by the frequencies of

various alleles in the populations. Focusing on just single isolates sourced from infections or

long-term evolution experiments, results in particular strains being characterized with certain

phenotypes, which misrepresents what is found in the population as a whole. It becomes

particularly problematic in studies focusing on single colonies from longitudinal samples, and

when genomic sequencing predicts how a strain genetically evolves over time during an

infection. Our findings may also be particularly relevant when considering whether a *P.*

aeruginosa infection is resistant or sensitive to antibiotic treatments. Our findings may extend

to other infections caused by *P. aeruginosa* such as non-healing chronic wounds and they may

also be relevant to other species of bacteria.

Materials and methods

Bacterial strains and growth conditions. For our experimental evolution, we used the PAO1

(University of Nottingham) strain of *P. aeruginosa.* For SCFM we followed the protocol

provided in [36, 38]. Briefly for the buffer base we prepared NaH_2PO_4 (1.3 mM), Na_2HPO4

(1.2 mM), KNO_3 (0.348 mM), K_2SO_4 (0.271 mM), NH_4Cl (2.28 mM), KCl (14.9 mM), NaCl

(51.8 mM) was prepared in 10mM of MOPS at pH=6.8, then the amino acids were added[36,

38] . The Dextrose (3mM), L-lactic acid (9.3 mM), $CaCl_2*2 H_2O$ (1.75 mM), $MgCl_2* 6H_2O$

(0.606 mM) and $Fe.SO_4*7H_2O$ (0.0036 mM) was added fresh every time the media was

prepared.

Long term experimental evolution. To assess how genomic diversity impacts on *P.*

aeruginosa populations, we generated a diverse population using a long-term evolution

experimental approach. We evolved the *P. aeruginosa* strain PAO1 in biofilms using plastic

beads [34, 35, 78] (9x6 mm width) suspended in SCFM, in order to mimic a biofilm life cycle

and a chemical environment similar to that found in CF lung sputum. To start the experimental

evolution process, we first grew PAO1 on an LB agar plate. Then we inoculated a single colony

of PAO1 into 3 ml of fresh SCFM [38], and incubated for up to 6 h to grow up to mid-log

phase. We stored this mid-log phase cells as the ancestral PAO1 and compared all further

phenotypic and genomic properties of the evolved populations to this. We inoculated mid-log

phase cells to $OD_{600} \approx 0.05$ into 4 tubes (in order to evolve separate independent lines named

A-D) containing 3 ml of SCFM and a plastic bead. We incubated cultures for 24 h at 37 °C/200

rpm. After 24 h of incubation, we transferred bacterial covered beads into fresh tubes

containing 3 ml of SCFM and a new bead. After each round the biofilms that formed on beads

were composed of approximately 10^8 cells. We then incubated again for 24 h at 37 °C/200 rpm.

We continued the bead transfers for 50 days and stored biofilm portion of populations (attached

to plastic beads) every 10 days (Rounds 1 to 5: R1-R5) (Fig. S1). We did not transfer or store

the planktonic fractions at any point during the selection experiment.

Deep sequencing of evolved populations. We extracted Genomic DNA from evolved

populations after 18 h growth in SCFM; using DNeasy® Blood & Tissue Kit (QIAGEN) by

following the manufacturer's instructions. We prepared sequencing libraries using the

NexteraXT protocol (Illumina), and sequenced in 24-plex on the Illumina MiSeq platform to

obtain an approximate calculated level of coverage of 220-600× for each evolved population.

A *de novo* assembly of the ancestral strain genome (*P. aeruginosa* PAO1 ancestor) was

obtained using Spades with the –careful flag, and annotated using Prokka. We mapped reads

of the evolved populations against the ancestral PAO1 genome using BWA[79]. For SNP

calling, the sequences were summarized using a MATLAB script for base quality, genomic

position and mapping quality [55] script using bowtie2 [80], SAMtools [81] and BAMtools.

To determine the allele frequency, we applied the breseq consensus model [82] to each of the

samples collected. Later, for the variant sites, we used the composition in each nucleotide as

the surrogacy for the allele frequency, and we excluded any allele frequency below 10% in

each evolved population. Our deep sequence analysis was not designed to detect any insertions

or deletions (INDELS), only SNPs. It is likely that INDELs were present in genes in our

evolved populations.

Measurement of biofilms formed on beads. To determine the levels of biofilm formation by

each evolved population, we grew biofilms on plastic beads as previously described [34, 78].

For each set of biofilm assays, we directly inoculated a 10 µl loop of frozen evolved

populations (approximately 10^6 cells) into 3 ml of SCFM and incubated at 37°C/200 rpm for

16 h. Then we measured the OD_{600}, and diluted in 3ml SCFM to $OD_{600} \approx 0.05$. Then 3 plastic

beads were added to each tube. After 24 h growth at 37°C/200 rpm and biofilm formation. The

beads were then washed 3× with 10 ml of PBS to remove any residual planktonic cells not

bound to the plastic beads. Then we transferred each plastic bead into 1ml of PBS and sonicated

the beads for 10 mins, using a bath sonicator to detach biofilm forming cells from the beads.

We then serially diluted the cells and plated out onto LB agar plates for colony forming unit

(CFU) calculations. To assess other phenotypic traits, a cell free supernatant of biofilm forming

cells was prepared from liquid part of the cultures and corrected values for $OD_{600} \approx 1$.

Preparation of cell free supernatants. To assess the levels of protease, QS signals and siderophore production during biofilm formation, we collected 3 ml of the SCFM used in the biofilm assays. We use the media surrounding the bead biofilms; and measured the OD_{600} and adjusted it to 1 for all the cultures with SCFM. We then filtered the supernatants using 0.22 μm filters and used these cell free supernatants to measure phenotypic traits.

Total protease activity. To assess the total protease activity of evolved populations, we used skimmed milk agar plates. We inoculated 10 μl of cell free supernatant from each evolved population onto skimmed milk agar plates (1.2% Bacto Agar, 0.015% of skimmed milk) alongside 10μl of 10μg/ml of proteinase K and supernatant of PAO1 as controls. The zone of clearance was scored based on appearance and measured with a ruler (in mm). We then imaged each plate using an Epson scanner at 800dpi. We then compared it to the zone of clearance produced by the PAO1 ancestor [83].

Siderophore production. To measure the levels of the two main siderophores produced by *P. aeruginosa*, we used the cell free supernatants. 100 μl aliquots of cell free supernatant from evolved populations and the PAO1 ancestor was transferred into a black clear bottom 96 well plate (Corning). We measured the emission as Relative Fluorescent Units (RFU) using a Multi microplate reader (Tecan Infinite® M200 Pro). We measured the wavelengths at excitation of 400 nm/emission460 nm for Pyoverdine and 350/430 nm for Pyochelin [30, 84, 85]. We corrected the values for pyoverdine and pyochelin to the absorption at (OD_{600}) of the original cultures.

Measurement of C4-HSL and 3O-C12-HSL produced by evolved populations. The cell free supernatants were used to determine the concentration of QS signals. We used two *E. coli* bioreporter strains to measure production of the two main signal molecules by the evolved population. The *E. coli* reporters pSB536 and pSB1142 were used to detect C4-HSL and 3O-

C12-HSL respectively [86]. We calculated signal levels based on standard curves fitted to the

concentrations of synthetic 3O-C12-HSL and C4-HSL standards (Sigma) [87, 88].

Mixed isolate experiments. For mixed constructed populations, we grew PAO1, the BB8

evolved isolate and BB8 complemented with *lasR* (BB8:*lasR*) and empty vector (BB8:vector)

in SCFM to $OD_{600} \approx 0.5$. We then mixed PAO1 and BB8 at two different starting frequencies

(50:50 and 30:70) in 3 ml of SCFM. The cultures were then incubated for 16 h at 37°C/200

rpm. We measured and adjusted the OD_{600} of each culture to $OD_{600} \approx 1$ and prepared a cell

free supernatant for QS signal activity measurements.

Antibiotic susceptibility assay. To determine the antibiotic susceptibility in evolved

populations, we followed the British Society for Antimicrobial Chemotherapy (BSAC)

guidelines (Version.14, 05-01-2015) using Isosensitest agar plates (Oxoid). We tested the

susceptibility of evolved populations, PAO1 ancestral strain and the NCTC (10662) PAO1

strain to Gentamicin (10 μg), Meropenem (10 μg), Ciprofloxacin (1 μg), Ceftazidime (30 μg),

Piperacillin/Tazobactam (85 μg) and Amikacin (30 μg) (Oxoid). The zone of inhibition and

clearance in this method was compared to the available zone of inhibition breakpoints for

susceptibility (mm) for each tested antibiotic based on BSAC guidelines.

Determining colony morphology diversity in evolved populations. To determine the

diversity in colony morphology in the biofilm evolved population, we used a Congo Red based

agar media (1% agar, 1xM63 salts (3g monobasic KHPO4, 7g K2PO4, 2g NH4.2SO4, pH

adjusted to 7.4), 2.5mM magnesium chloride, 0.4 mM calcium chloride, 0.1% casamino acids,

0.1% yeast extracts, 40 mg/L Congo red solution, 100 μM ferrous ammonium sulphate and

0.4% glycerol) [83]. We recovered the evolved populations from beads and serially diluted the

populations and then inoculated onto CRA plates alongside the PAO1 ancestor. We incubated

the plates overnight at 37°C, and for a further 4 days at 22°C. The colonies were imaged using

an Epson scanner at 800dpi.

Complementation of *lasR*. To complement an evolved isolate (BB8) containing a *lasR*

mutation with a functional *lasR* allele, we amplified a 920 bp product comprising *lasR* and 200

bp upstream of the *lasR* start codon that includes its native promoter, from genomic DNA

isolated from wildtype ancestral PAO1. We cloned this PCR product into the shuttle vector

pME6032, which replicates in both *E. coli* and *P.aeruginosa*, [89] by Gibson assembly [90]

using the commercially available NEBuilder HiFi DNA Assembly Cloning Kit (New England

Biolabs, Ipswich, MA). We introduced the *lasR* complementation construct into BB8 by

electroporation [91] and plating on selective media containing 300 ug/mL tetracycline. We

then tested susceptibility of the *lasR* complemented strains to ceftazidime and

piperacillin/tazobactam using the BSAC method. For mixed constructed populations, we grew

PAO1, the BB8 evolved isolate and BB8 complemented with *lasR* (BB8:*lasR*) and empty

vector (BB8:vector) in SCFM to $OD_{600} \approx 0.5$. We then mixed PAO1 and BB8 at two different

starting frequencies (50:50 and 30:70) in 1 ml of PBS and tested for antibiotic tolerance by

following the British Society for Antimicrobial Chemotherapy (BSAC) guidelines (Version.14,

05-01-2015) using Isosensitest agar plates (Oxoid).

Statistical analysis. For statistical analysis of the phenotypic assays, we used GraphPad Prism

8.0. For analysis of SNP frequency, we used R package 3.6. We used the Interactive Venn [92]

to analyze shared SNPs within and between evolved populations.

Acknowledgements. For funding we thank the Human Frontier Science Program

(RGY0081/2012) and Georgia Institute of Technology; The Cystic Fibrosis Foundation

(DIGGLE18I0) to SPD; Cystic Fibrosis Foundation for a Fellowship to SA (AZIMI18F0);

CF@latna for a Fellowship to SA (3206AXB). The National Heart Lung Blood Institute

(R56HL142857) and The Simons Foundation (396001) to SPB. We acknowledge Jacob

Thomas for help with *lasR* complementation and Freya Harrison and James Gurney for helpful

comments on the work.

Figure legends

Figure 1. The evolutionary trajectory of *P. aeruginosa* during biofilm growth in SCFM.

The evolution trajectories were similar in four independent replicate lines, and the major

genetic heterogeneity in the evolved populations occurred during the first 10 days of selection.

(A) SNP calling analysis showed that the four independently evolved populations had more

than 80% similarity (267 shared SNPs) in genomic changes in the populations over the 50 days

of selection. (B) The Venn diagram shows the shared SNPs between all 4 replicate lines, over

50 days of selection. Numbers in brackets, (i.e 1354 for 10 days), represent the combined

number of SNPs in each of the four evolved lines. The majority of genetic variation within the

populations occurred within the first 10 days of selection. Each circle represents the total

number SNPs for each independently evolved line.

Figure 2. Allele frequency of SNPs changes over the course of selection. Emergence of

nonsynonymous SNPs in genes involved in social traits (QS: *lasR*), oxidative respiration and

DNA mismatch repair, motility and iron chelation occurred after 20 and 30 days of selection;

and fixed SNPs in genes such as elongation factor *tufA,* Cytochrome c oxidase subunit (*ccoN2*)

and pyoverdine biosynthesis protein *pvdJ* emerged after 10 days of selection and were fixed in

the populations over 50 days (A-D represents replicate evolving lines).

Figure 3. Phenotypes of evolved populations of PAO1. (A) Increase in biofilm formation by

evolved populations of PAO1 (Kurskal-Wallis, Dunn's multiple comparison to PAO1, $p < 0.05$,

n=5. Errors bars indicate the standard error of the mean, SEM); (B) The total protease activity

was reduced at 30 days of selection; (C) Production of 3O-C12-HSL significantly decreased at

30 days of selection (Kruskal-Wallis, uncorrected Dunn's test multiple comparison to PAO1,

$p < 0.05$, n=5. Errors bars indicate the SEM) and there were no significant changes in the levels

of C4-HSL; (D) There were no significant changes in siderophore production in evolved

populations compared to the PAO1 ancestor (Kruskal-Wallis, uncorrected Dunn's test multiple

comparison to PAO1, $p > 0.05$, n=5). Each dot represents the mean value of one independent

experiment for each evolved population at each round of selection. All experiments were

performed in 3 technical replicates.

Figure 4. Loss of protease activity and a decrease in QS signal production in evolved

populations can be explained by an accumulation of *lasR* SNPs. A linear regression model

showed that the loss of total protease activity and the decrease in 3O-C12-HSL levels can be

explained by the accumulation of *lasR* SNPs. (A) There is a significant negative correlation

between *lasR* SNP frequency and total protease activity of the evolved populations ($R^2 =$

0.7298, $p = 0.04$; n=8); (B) There is no significant correlation between 3O-C12-HSL levels

and accumulation of the *lasR* SNP in the evolved populations ($R^2 = 0.5363$, $p = 0.1594$; n=8).

However, if only changes in 3O-C12-HSL levels after 30 days of selection is considered, there

is a strong correlation between emergence and accumulation of *lasR* SNP in evolved

populations (small inset: $R^2 = 1.0$; $p = 0.0034$; n=8); (C) There are no significant correlations

between accumulation of *lasR* SNPs in the evolved populations and levels of C4-HSL ($R^2 =$

0.1187, $p = 0.5702$; n=8); (D) There is no significant correlation between *lasR* frequency and

biofilm formation ($R^2 = 0.0004$, $p = 0.9062$; n=8).

Figure 5. Increased resistance to ceftazidime and piperacillin/tazobactam in evolved

populations is correlated with *lasR* SNP frequency. There is an increased resistance,

indicated by a decrease in the zone of inhibition (ZOI) to ceftazidime (A) and

piperacillin/tazobactam (B) in evolved populations after 30 days of selection (R3). The red

lines represent the ZOI breakpoint (mm) for each antibiotic (BSAC guidelines). Below the lines

indicates resistance. There is a positive correlation between increased resistance and the

frequency of the *lasR* SNP in the evolved populations (n=8, Pearson correlation. Each dot

represents the mean value for the zone of inhibition at each round of selection in all 8

independent experiments. (C) Complementation of a 50 day evolved isolate (BB8) containing

a SNP in *lasR* with an intact *lasR in trans,* restores sensitivity to ceftazidime and

piperacillin/tazobactam. Differential frequencies of B8 mixed with ancestral PAO1

significantly increased tolerance to ceftazidime (n=5, Two way ANOVA, multiple comparison

to PAO1, Fisher's LSD test, p <0.05. Errors bars indicate SEM).

Supplementary figure legends

Figure S1. Long term evolution of PAO1 in biofilms and SCFM. PAO1 was grown in SCFM

and inoculated into four independent replicate lines in the presence of one plastic bead (6×7

mm) to enable the bacteria to form biofilms on the surface. Every 24 hours the beads were

transferred to a new tube with fresh SCFM media and a new bead. After 10 days, the bacterial

populations were sonicated off the beads and stored as round 1 (R1). To start the next 10 days

of selection, a 10 µl loop of bacteria (approximately 10^6 cells) was used. This whole process

was repeated 5 times to give 50 days (rounds 1 to 5) of selection.

Figure S2. Number of SNPs shared in all evolved lines during 50 days of evolution in

SCFM. We tested the number and percentages of the SNPS that occurred and are shared

between various evolution lines at different rounds of selections. There are 207 SNPs that are

shared between all 4 lines of selection (A-D) at day 10, 205 on day 20, 206 on day 30, 172 on

day 40 and 169 on day 50 (upper panels). We observed that the majority of SNPs occurred

during the first 10 days of selection and are shared and carried over to subsequent rounds of

selection in all evolution lines.

Figure S3. Number of SNPs in genes over 50 days of selection in SCFM. The emergence

of polymorphisms in 45 genes involved in various physiological functions (Evolved line A

shown as a representative for all evolved lines). The shaded boxes represent the abundance of

SNPs in each gene.

Figure S4. Abundance of differential allele frequencies in each evolved population over

five rounds of selection. Different SNPS emerged after 10 days of selection at varying

frequencies within the populations. Each round of selection contained varying frequencies of

SNPs. Over 10-25% of SNPs were fixed in populations after 10 days of selection, across all

four independently evolving populations (A-D).

Figure S5. Number of SNPs in various genes after 50 days of selection. We found high

levels of similarity in number and frequency of SNPs in all independent evolved lines after 50

days of selection. Each data point is representative of SNPs in each gene, at various frequencies

in all four evolved lines.

Figure S6. Increase in diversity of colony morphologies in evolving populations over time.

Diversity of colony morphology increased over time in evolutionary lines. We used Congo Red

Agar plates to highlight differences in morphologies between haplotypes.

Figure S7. Increased resistance to β-lactam antibiotics in evolved populations. (A) There

is a decrease in the ZOI to all β-lactam antibiotics after 30 days of selection in evolved

populations. (B) There is an increase in resistance to meropenem after 30 days of selection in

the evolved population, which significantly correlates to accumulation of the *lasR* SNP in the

populations. However, the increased resistance does not pass the threshold to be considered

clinically resistant to meropenem. (C) There is no correlation between the levels of biofilm

formation or antibiotic resistance at the population level.

Figure S8. Increase in *lasR* mutant frequency changes the functional phenotype of

synthetic mixed populations. Varying frequencies of BB8 isolate (V208G *lasR* allele) with

the PAO1 ancestor, significantly alters the levels of 3O-C12-HSL and has no impact C4-HSL

levels (Multiple comparison, One Way ANOVA, n=5).

Table S1. SNP positions and altered allele frequencies in four evolved lines over 5 rounds of

selection.

Table S2. SNPs in the 50 day evolved isolate BB8.

References

1. Adler, F.R. and T.G. Liou, *The Dynamics of Disease Progression in Cystic Fibrosis.* PLoS One, 2016. **11**(6): p. e0156752.
2. Cantin, A.M., et al., *Inflammation in cystic fibrosis lung disease: Pathogenesis and therapy.* J Cyst Fibros, 2015. **14**(4): p. 419-30.
3. Elborn, J.S., *Cystic fibrosis.* Lancet, 2016. **388**(10059): p. 2519-2531.
4. Jorth, P., et al., *Direct Lung Sampling Indicates That Established Pathogens Dominate Early Infections in Children with Cystic Fibrosis.* Cell Rep, 2019. **27**(4): p. 1190-1204 e3.
5. Diggle, S.P. and M. Whiteley, *Microbe Profile: Pseudomonas aeruginosa: opportunistic pathogen and lab rat.* Microbiology, 2019.
6. Frederiksen, B., C. Koch, and N. Hoiby, *Antibiotic treatment of initial colonization with Pseudomonas aeruginosa postpones chronic infection and prevents deterioration of pulmonary function in cystic fibrosis.* Pediatr Pulmonol, 1997. **23**(5): p. 330-5.
7. Burns, J.L., et al., *Longitudinal assessment of Pseudomonas aeruginosa in young children with cystic fibrosis.* J Infect Dis, 2001. **183**(3): p. 444-52.
8. Bjarnsholt, T., et al., *Pseudomonas aeruginosa biofilms in the respiratory tract of cystic fibrosis patients.* Pediatr Pulmonol, 2009. **44**(6): p. 547-58.
9. Hogardt, M. and J. Heesemann, *Adaptation of Pseudomonas aeruginosa during persistence in the cystic fibrosis lung.* Int J Med Microbiol, 2010. **300**(8): p. 557-62.
10. Doring, G., I.G. Parameswaran, and T.F. Murphy, *Differential adaptation of microbial pathogens to airways of patients with cystic fibrosis and chronic obstructive pulmonary disease.* FEMS Microbiol Rev, 2011. **35**(1): p. 124-46.
11. Filkins, L.M. and G.A. O'Toole, *Cystic Fibrosis Lung Infections: Polymicrobial, Complex, and Hard to Treat.* PLoS Pathog, 2015. **11**(12): p. e1005258.
12. Jelsbak, L., et al., *Molecular epidemiology and dynamics of Pseudomonas aeruginosa populations in lungs of cystic fibrosis patients.* Infect Immun, 2007. **75**(5): p. 2214-24.
13. Wilder, C.N., G. Allada, and M. Schuster, *Instantaneous within-patient diversity of Pseudomonas aeruginosa quorum-sensing populations from cystic fibrosis lung infections.* Infect Immun, 2009. **77**(12): p. 5631-9.
14. Folkesson, A., et al., *Adaptation of Pseudomonas aeruginosa to the cystic fibrosis airway: an evolutionary perspective.* Nat Rev Microbiol, 2012. **10**(12): p. 841-51.
15. Marvig, R.L., et al., *Convergent evolution and adaptation of Pseudomonas aeruginosa within patients with cystic fibrosis.* Nat Genet, 2015. **47**(1): p. 57-64.
16. Darch, S.E., et al., *Recombination is a key driver of genomic and phenotypic diversity in a Pseudomonas aeruginosa population during cystic fibrosis infection.* Sci Rep, 2015. **5**: p. 7649.
17. Williams, D., et al., *Divergent, coexisting Pseudomonas aeruginosa lineages in chronic cystic fibrosis lung infections.* Am J Respir Crit Care Med, 2015. **191**(7): p. 775-85.
18. Wee, B.A., et al., *Whole genome sequencing reveals the emergence of a Pseudomonas aeruginosa shared strain sub-lineage among patients treated within a single cystic fibrosis centre.* BMC Genomics, 2018. **19**(1): p. 644.
19. Smith, E.E., et al., *Genetic adaptation by Pseudomonas aeruginosa to the airways of cystic fibrosis patients.* Proc Natl Acad Sci U S A, 2006. **103**(22): p. 8487-92.

20. Khaledi, A., et al., *Transcriptome Profiling of Antimicrobial Resistance in*
Pseudomonas aeruginosa. Antimicrob Agents Chemother, 2016. **60**(8): p. 4722-33.
21. Baldan, R., et al., *Adaptation of Pseudomonas aeruginosa in Cystic Fibrosis airways*
influences virulence of Staphylococcus aureus in vitro and murine models of co-
infection. PLoS One, 2014. **9**(3): p. e89614.
22. O'Toole, G.A., *Cystic Fibrosis Airway Microbiome: Overturning the Old, Opening*
the Way for the New. J Bacteriol, 2018. **200**(4).
23. Ahlgren, H.G., et al., *Clinical outcomes associated with Staphylococcus aureus and*
Pseudomonas aeruginosa airway infections in adult cystic fibrosis patients. BMC
Pulm Med, 2015. **15**: p. 67.
24. Limoli, D.H., et al., *Pseudomonas aeruginosa Alginate Overproduction Promotes*
Coexistence with Staphylococcus aureus in a Model of Cystic Fibrosis Respiratory
Infection. MBio, 2017. **8**(2).
25. Radlinski, L., et al., *Pseudomonas aeruginosa exoproducts determine antibiotic*
efficacy against Staphylococcus aureus. PLoS Biol, 2017. **15**(11): p. e2003981.
26. O'Brien, S. and J.L. Fothergill, *The role of multispecies social interactions in shaping*
Pseudomonas aeruginosa pathogenicity in the cystic fibrosis lung. FEMS Microbiol
Lett, 2017. **364**(15).
27. Limoli, D.H. and L.R. Hoffman, *Help, hinder, hide and harm: what can we learn*
from the interactions between Pseudomonas aeruginosa and Staphylococcus aureus
during respiratory infections? Thorax, 2019.
28. Jorth, P., et al., *Regional Isolation Drives Bacterial Diversification within Cystic*
Fibrosis Lungs. Cell Host Microbe, 2015. **18**(3): p. 307-19.
29. Marvig, R.L., et al., *Within-host microevolution of Pseudomonas aeruginosa in*
Italian cystic fibrosis patients. BMC Microbiol, 2015. **15**: p. 218.
30. Jiricny, N., et al., *Loss of social behaviours in populations of Pseudomonas*
aeruginosa infecting lungs of patients with cystic fibrosis. PLoS One, 2014. **9**(1): p.
e83124.
31. Bartell, J.A., et al., *Evolutionary highways to persistent bacterial infection.* Nat
Commun, 2019. **10**(1): p. 629.
32. Markussen, T., et al., *Environmental heterogeneity drives within-host diversification*
and evolution of Pseudomonas aeruginosa. MBio, 2014. **5**(5): p. e01592-14.
33. Didelot, X., et al., *Within-host evolution of bacterial pathogens.* Nat Rev Microbiol,
2016. **14**(3): p. 150-62.
34. Poltak, S.R. and V.S. Cooper, *Ecological succession in long-term experimentally*
evolved biofilms produces synergistic communities. ISME J, 2011. **5**(3): p. 369-78.
35. Flynn, K.M., et al., *Evolution of Ecological Diversity in Biofilms of Pseudomonas*
aeruginosa by Altered Cyclic Diguanylate Signaling. J Bacteriol, 2016. **198**(19): p.
2608-18.
36. Palmer, K.L., et al., *Cystic fibrosis sputum supports growth and cues key aspects of*
Pseudomonas aeruginosa physiology. J Bacteriol, 2005. **187**(15): p. 5267-77.
37. Turner, K.H., et al., *Essential genome of Pseudomonas aeruginosa in cystic fibrosis*
sputum. Proc Natl Acad Sci U S A, 2015. **112**(13): p. 4110-5.
38. Palmer, K.L., L.M. Aye, and M. Whiteley, *Nutritional cues control Pseudomonas*
aeruginosa multicellular behavior in cystic fibrosis sputum. J Bacteriol, 2007.
189(22): p. 8079-87.
39. Varga, J.J., et al., *Genotypic and phenotypic analyses of a Pseudomonas aeruginosa*
chronic bronchiectasis isolate reveal differences from cystic fibrosis and laboratory
strains. BMC Genomics, 2015. **16**: p. 883.

40. Feltner, J.B., et al., *LasR Variant Cystic Fibrosis Isolates Reveal an Adaptable*
Quorum-Sensing Hierarchy in Pseudomonas aeruginosa. MBio, 2016. **7**(5).
41. Kostylev, M., et al., *Evolution of the Pseudomonas aeruginosa quorum-sensing*
hierarchy. Proc Natl Acad Sci U S A, 2019. **116**(14): p. 7027-7032.
42. Latifi, A., et al., *A hierarchical quorum-sensing cascade in Pseudomonas aeruginosa*
links the transcriptional activators LasR and RhIR (VsmR) to expression of the
stationary-phase sigma factor RpoS. Mol Microbiol, 1996. **21**(6): p. 1137-46.
43. Gambello, M.J. and B.H. Iglewski, *Cloning and characterization of the Pseudomonas*
aeruginosa lasR gene, a transcriptional activator of elastase expression. J Bacteriol,
1991. **173**(9): p. 3000-9.
44. Pearson, J.P., E.C. Pesci, and B.H. Iglewski, *Roles of Pseudomonas aeruginosa las*
and rhl quorum-sensing systems in control of elastase and rhamnolipid biosynthesis
genes. Journal of Bacteriology, 1997. **179**(18): p. 5756-5767.
45. Gilbert, K.B., et al., *Global position analysis of the Pseudomonas aeruginosa*
quorum-sensing transcription factor LasR. Mol Microbiol, 2009. **73**(6): p. 1072-85.
46. Diggle, S.P., et al., *Evolutionary theory of bacterial quorum sensing: when is a signal*
not a signal? Philos Trans R Soc Lond B Biol Sci, 2007. **362**(1483): p. 1241-9.
47. Bjarnsholt, T., et al., *Quorum sensing and virulence of Pseudomonas aeruginosa*
during lung infection of cystic fibrosis patients. PLoS One, 2010. **5**(4): p. e10115.
48. Bartell, J.A., et al., *Reconstruction of the metabolic network of Pseudomonas*
aeruginosa to interrogate virulence factor synthesis. Nat Commun, 2017. **8**: p. 14631.
49. Whiteley, M., S.P. Diggle, and E.P. Greenberg, *Progress in and promise of bacterial*
quorum sensing research. Nature, 2017. **551**(7680): p. 313-320.
50. Ciofu, O., et al., *Genetic adaptation of Pseudomonas aeruginosa during chronic lung*
infection of patients with cystic fibrosis: strong and weak mutators with
heterogeneous genetic backgrounds emerge in mucA and/or lasR mutants.
Microbiology, 2010. **156**(Pt 4): p. 1108-19.
51. Diaz Caballero, J., et al., *Selective Sweeps and Parallel Pathoadaptation Drive*
Pseudomonas aeruginosa Evolution in the Cystic Fibrosis Lung. MBio, 2015. **6**(5): p.
e00981-15.
52. Feliziani, S., et al., *Mucoidy, Quorum Sensing, Mismatch Repair and Antibiotic*
Resistance in Pseudomonas aeruginosa from Cystic Fibrosis Chronic Airways
Infections. Plos One, 2010. **5**(9).
53. Nguyen, D. and P.K. Singh, *Evolving stealth: Genetic adaptation of Pseudomonas*
aeruginosa during cystic fibrosis infections. Proceedings of the National Academy of
Sciences of the United States of America, 2006. **103**(22): p. 8305-8306.
54. Marvig, R.L., et al., *Genome analysis of a transmissible lineage of pseudomonas*
aeruginosa reveals pathoadaptive mutations and distinct evolutionary paths of
hypermutators. PLoS Genet, 2013. **9**(9): p. e1003741.
55. Lieberman, T.D., et al., *Genetic variation of a bacterial pathogen within individuals*
with cystic fibrosis provides a record of selective pressures. Nat Genet, 2014. **46**(1):
p. 82-7.
56. LaFayette, S.L., et al., *Cystic fibrosis-adapted Pseudomonas aeruginosa quorum*
sensing lasR mutants cause hyperinflammatory responses. Sci Adv, 2015. **1**(6).
57. Rumbaugh, K.P., J.A. Griswold, and A.N. Hamood, *Contribution of the regulatory*
gene lasR to the pathogenesis of Pseudomonas aeruginosa infection of burned mice. J
Burn Care Rehabil, 1999. **20**(1 Pt 1): p. 42-9.
58. Storey, D.G., et al., *Pseudomonas aeruginosa lasR transcription correlates with the*
transcription of lasA, lasB, and toxA in chronic lung infections associated with cystic
fibrosis. Infection and Immunity, 1998. **66**(6): p. 2521-2528.

59.   Chowdhury, N. and A. Bagchi, *Molecular insight into the activity of LasR protein*
from Pseudomonas aeruginosa in the regulation of virulence gene expression by this
organism. Gene, 2016. **580**(1): p. 80-87.
60.   Kafle, P., et al., *Molecular Insights into the Impact of Oxidative Stress on the*
Quorum-Sensing Regulator Protein LasR. Journal of Biological Chemistry, 2016.
291(22): p. 11776-11786.
61.   D'Argenio, D.A., et al., *Growth phenotypes of Pseudomonas aeruginosa lasR mutants*
adapted to the airways of cystic fibrosis patients. Mol Microbiol, 2007. **64**(2): p. 512-
33.
62.   Schick, A. and R. Kassen, *Rapid diversification of Pseudomonas aeruginosa in cystic*
fibrosis lung-like conditions. Proc Natl Acad Sci U S A, 2018. **115**(42): p. 10714-
10719.
63.   Fung, C., et al., *Gene expression of Pseudomonas aeruginosa in a mucin-containing*
synthetic growth medium mimicking cystic fibrosis lung sputum. J Med Microbiol,
2010. **59**(Pt 9): p. 1089-100.
64.   Behrends, V., et al., *Direct assessment of metabolite utilization by Pseudomonas*
aeruginosa during growth on artificial sputum medium. Appl Environ Microbiol,
2013. **79**(7): p. 2467-70.
65.   Melnyk, A.H., et al., *Evolution of Cost-Free Resistance under Fluctuating Drug*
Selection in Pseudomonas aeruginosa. mSphere, 2017. **2**(4).
66.   Wong, A., N. Rodrigue, and R. Kassen, *Genomics of adaptation during experimental*
evolution of the opportunistic pathogen Pseudomonas aeruginosa. PLoS Genet, 2012.
8(9): p. e1002928.
67.   Bielecki, P., et al., *Ex vivo transcriptional profiling reveals a common set of genes*
important for the adaptation of Pseudomonas aeruginosa to chronically infected host
sites. Environ Microbiol, 2013. **15**(2): p. 570-87.
68.   Ciofu, O., et al., *Occurrence of hypermutable Pseudomonas aeruginosa in cystic*
fibrosis patients is associated with the oxidative stress caused by chronic lung
inflammation. Antimicrob Agents Chemother, 2005. **49**(6): p. 2276-82.
69.   Manfredi, P. and U. Jenal, *Bacteria in the CF Lung: Isolation Drives Diversity.* Cell
Host Microbe, 2015. **18**(3): p. 268-9.
70.   Diggle, S.P. and M. Whiteley, *Microbe Profile: Pseudomonas aeruginosa:*
opportunistic pathogen and lab rat. Microbiology, 2020. **166**(1): p. 30-33.
71.   Kirisits, M.J., et al., *Characterization of colony morphology variants isolated from*
Pseudomonas aeruginosa biofilms. Appl Environ Microbiol, 2005. **71**(8): p. 4809-21.
72.   Arai, H., *Regulation and Function of Versatile Aerobic and Anaerobic Respiratory*
Metabolism in Pseudomonas aeruginosa. Front Microbiol, 2011. **2**: p. 103.
73.   Wang, C., et al., *Transcription-associated mutation of lasR in Pseudomonas*
aeruginosa. DNA Repair, 2016. **46**: p. 9-19.
74.   Tai, A.S., et al., *Genotypic Diversity within a Single Pseudomonas aeruginosa Strain*
Commonly Shared by Australian Patients with Cystic Fibrosis. Plos One, 2015.
10(12).
75.   Rumbaugh, K.P., et al., *Kin selection, quorum sensing and virulence in pathogenic*
bacteria. Proc Biol Sci, 2012. **279**(1742): p. 3584-8.
76.   Kohler, T., A. Buckling, and C. van Delden, *Cooperation and virulence of clinical*
Pseudomonas aeruginosa populations. Proc Natl Acad Sci U S A, 2009. **106**(15): p.
6339-44.
77.   Rumbaugh, K.P., et al., *Quorum sensing and the social evolution of bacterial*
virulence. Curr Biol, 2009. **19**(4): p. 341-5.

78.   Traverse, C.C., et al., *Tangled bank of experimentally evolved Burkholderia biofilms*
765   *reflects selection during chronic infections.* Proc Natl Acad Sci U S A, 2013. **110**(3):
766   p. E250-9.
767   79.   Li, H. and R. Durbin, *Fast and accurate short read alignment with Burrows-Wheeler*
768   *transform.* Bioinformatics, 2009. **25**(14): p. 1754-60.
769   80.   Langmead, B. and S.L. Salzberg, *Fast gapped-read alignment with Bowtie 2.* Nat
770   Methods, 2012. **9**(4): p. 357-9.
771   81.   Li, H., et al., *The Sequence Alignment/Map format and SAMtools.* Bioinformatics,
772   2009. **25**(16): p. 2078-9.
773   82.   Barrick, J.E., et al., *Identifying structural variation in haploid microbial genomes*
774   *from short-read resequencing data using breseq.* BMC Genomics, 2014. **15**: p. 1039.
775   83.   Mayer-Hamblett, N., et al., *Pseudomonas aeruginosa in vitro phenotypes distinguish*
776   *cystic fibrosis infection stages and outcomes.* Am J Respir Crit Care Med, 2014.
777   **190**(3): p. 289-97.
778   84.   Kummerli, R., et al., *Phenotypic plasticity of a cooperative behaviour in bacteria.* J
779   Evol Biol, 2009. **22**(3): p. 589-98.
780   85.   Ankenbauer, R., S. Sriyosachati, and C.D. Cox, *Effects of siderophores on the growth*
781   *of Pseudomonas aeruginosa in human serum and transferrin.* Infect Immun, 1985.
782   **49**(1): p. 132-40.
783   86.   Winson, M.K., et al., *Multiple N-Acyl-L-Homoserine Lactone Signal Molecules*
784   *Regulate Production of Virulence Determinants and Secondary Metabolites in*
785   *Pseudomonas-Aeruginosa.* Proceedings of the National Academy of Sciences of the
786   United States of America, 1995. **92**(20): p. 9427-9431.
787   87.   Cornforth, D.M., et al., *Combinatorial quorum sensing allows bacteria to resolve*
788   *their social and physical environment.* Proc Natl Acad Sci U S A, 2014. **111**(11): p.
789   4280-4.
790   88.   Popat, R., et al., *Conflict of interest and signal interference lead to the breakdown of*
791   *honest signaling.* Evolution, 2015. **69**(9): p. 2371-83.
792   89.   Heeb, S., et al., *Small, stable shuttle vectors based on the minimal pVS1 replicon for*
793   *use in gram-negative, plant-associated bacteria.* Molecular Plant-Microbe
794   Interactions, 2000. **13**(2): p. 232-237.
795   90.   Gibson, D.G., et al., *Enzymatic assembly of DNA molecules up to several hundred*
796   *kilobases.* Nature Methods, 2009. **6**(5): p. 343-U41.
797   91.   Choi, K.H., A. Kumar, and H.P. Schweizer, *A 10-min method for preparation of*
798   *highly electrocompetent Pseudomonas aeruginosa cells: application for DNA*
799   *fragment transfer between chromosomes and plasmid transformation.* J Microbiol
800   Methods, 2006. **64**(3): p. 391-7.
801   92.   Heberle, H., et al., *InteractiVenn: a web-based tool for the analysis of sets through*
802   *Venn diagrams.* Bmc Bioinformatics, 2015. **16**.
803

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Step 1

PAO1 Ancestral strain

Single colony

3ml of SCFM

6 h at 37°C/200 rpm

Step 2

Dilute mid-log phase cells into 4 lines

Plastic bead added

Step 3

3ml of SCFM

Bead transfer
Passage #1

24 h at 37°C/200 rpm

3ml of SCFM

24 h at 37°C/200 rpm

Bead transfer
Passage #2

3ml of SCFM

8 more passages

Bead transfer
Up to Passage #10
Round 1 (R1: 10 days)
Sonicated the cells
off the beads and
stored in -80 °C

A loop of the frozen stock from each line (A-D) was plated out on LB agar plates; all colonies were harvested and these colony homogenates were inoculated to 3ml of SCFM to start the next round of selection up to 5 rounds (50 days).

Figure S1

Figure S2

Figure S3

Figure S4

Figure S5

Figure S6

Figure S7

Figure S8

Figure 5