Yap suppresses T cell function and infiltration in the tumor microenvironment

Eleni Stampouloglou[1], Anthony Federico [2,3], Emily Slaby [4], Stefano Monti [2,3], Gregory L. Szeto[4,5*] and Xaralabos Varelas[1*]

[1] Department of Biochemistry, Boston University School of Medicine, Boston, MA 02118, USA

[2] Department of Medicine, Division of Computational Biology, Boston University School of Medicine, Boston, MA 02118, USA

[3] Bioinformatics Program, Boston University, Boston, MA 02115, USA.

[4] Department of Chemical, Biochemical, and Environmental Engineering, University of Maryland Baltimore County, Baltimore, MD 21250, USA.

[5] Marlene and Stewart Greenebaum Comprehensive Cancer Center, University of Maryland, Baltimore, MD 21201, USA.

* Correspondence:

Xaralabos Varelas, Department of Biochemistry, Boston University School of Medicine, Boston, Massachusetts 02118, USA

Email: xvarelas@bu.edu

Phone: 617-358-4575

&

Gregory L. Szeto, Department of Chemical, Biochemical, and Environmental Engineering, University of Maryland Baltimore County, Baltimore, MD 21250, USA

Email: gszeto@umbc.edu

Phone: 410-455-8157

ABSTRACT

A major challenge for cancer immunotherapy is sustaining T cell activation and recruitment in immunosuppressive solid tumors. Here we report that Yap levels are sharply induced upon CD4$^+$ and CD8$^+$ T cell activation and that Yap functions as an immunosuppressive factor and inhibitor of effector differentiation. Loss of Yap results in enhanced T cell activation, differentiation and function, which translates *in vivo* to an improved ability for T cells to infiltrate and repress tumors. Gene expression analyses of tumor-infiltrating T cells following Yap deletion implicates Yap as a mediator of global T cell responses in the tumor microenvironment and as a key negative regulator of T cell tumor infiltration and patient survival in diverse human cancers. Collectively, our results indicate that Yap plays critical roles in T cell biology, and suggest that inhibiting Yap activity improves T cell responses in cancer.

INTRODUCTION

CD8$^+$ and CD4$^+$ T cells are central players in the adaptive immune system. T cells elicit targeted, antigen-specific responses for direct killing of an infected or transformed cell, shaping and regulating the immune response in host defense (Kumar et al., 2018). Most mature T cells circulate in a resting, naïve state, and upon cognate antigen recognition, T cells become activated, proliferate clonally, and differentiate into effector T cells. Naïve CD8$^+$ T cells differentiate into cytotoxic T cells, while CD4$^+$ T cells differentiate into an array of different types of helper T cells (i.e., Th1, Th2, Th17 or Treg) depending on microenvironmental cues (Williams and Bevan, 2007). Each phenotype is defined by expression of signature transcription factors and effector cytokines leading to distinct functions (Luckheeram et al., 2012).

T cell activation also upregulates negative feedback mechanisms, such as inhibitory receptors, which restrain their action to minimize pathogenic inflammation and autoimmunity (Kumar et al., 2018; Murakami and Riella, 2014). This network of immunosuppressive factors is frequently co-opted in chronic infections and cancer, leading to terminally differentiated and exhausted T cells that lose effector function and ability to infiltrate disease sites (Wherry and Kurachi, 2015). The finding that revitalization of exhausted, dysfunctional T cells can restore the immune response has revolutionized cancer therapy with the use of checkpoint inhibition (Pauken and Wherry, 2015). Chimeric antigen receptor T cells (CAR-T), engineered for enhanced antigen recognition and co-stimulation also demonstrate promising clinical efficacy (Brentjens et al., 2013; Lee et al., 2015; Maude et al., 2014). However, both immunotherapies are effective for only a fraction of patients (Feng et al., 2013; Herbst et al., 2014; Rizvi et al., 2015; Topalian et al., 2012; Zhou et al., 2018). Major challenges to extending the efficacy of immunotherapy to more cancer patients include sustaining T cell activation and achieving T cell infiltration in the immunosuppressive microenvironment of solid tumors (Balatoni et al., 2018; Brown and Mackall, 2019; Cogdill et al., 2017; Darvin et al., 2018; Gajewski et al., 2013; Galon et al., 2006; Galon et al., 2012; Hamid et al., 2011; Mlecnik et al., 2011; Pages et al., 2005). Improved understanding of mechanisms controlling T cell differentiation and function is critical to overcoming these barriers.

Yap is a key effector of the Hippo signaling pathway, directing transcriptional programs that control stem cell biology by integrating microenvironmental and cell intrinsic cues (Varelas, 2014). Yap regulation leads to control of cell death and survival, proliferation and cell fate determination, and dysregulated Yap activity contributes to disease, most notably cancer (Piccolo et al., 2014). While the dynamics of Yap regulation coupled with differentiation are well characterized in stem cells and tissue specific progenitor cells, less is known about the role of Yap in T cells. The Hippo pathway has been implicated in coupling CD8[+] T cell clonal expansion to terminal differentiation through upregulation of kinase LATS1 and Yap degradation (Thaventhiran et al., 2012). Yap has also been shown to be important for Treg function, as have the Hippo pathway kinases MST1 and MST2 (Du et al., 2014; Li et al., 2015; Ni et al., 2018; Shi et al., 2018). The MST1/2 kinases have also been implicated in thymocyte egress and antigen recognition, lymphocyte polarization, adhesion and trafficking, survival, differentiation and proliferation (Dong et al., 2009; Katagiri et al., 2006; Katagiri et al., 2009; Mou et al., 2012; Nehme et al., 2012; Nishikimi et al., 2014; Tang et al., 2015; Ueda et al., 2012; Zhou et al., 2008). While these reports provide evidence for Hippo-Yap signaling as a mediator of T cell responses, the role of Yap in regulating CD4[+] and CD8[+] T cell differentiation and function remains understudied.

Here we report that Yap is a critical regulator of T cell fate and function, and that loss of Yap enhances T cell responses in cancer. We show that Yap is induced upon T cell activation and that T cell-specific deletion of *YAP* enhanced T cell activation and function. Yap deletion in T cells also inhibited growth of B16F10 tumors in mice and increased CD4[+] and CD8[+] T cell infiltration in the tumor. RNA-sequencing analysis of Yap-deleted tumor infiltrated lymphocytes (TILs) showed upregulation of key signaling events in T cell activation, differentiation and function. Yap-regulated gene expression signatures correlate with T cell infiltration and patient survival across multiple human cancers in The Cancer Genome Atlas (TCGA). Our results implicate Yap as a major inhibitor of T cell activation and function, and suggest that Yap inhibition can increase tumor infiltration and survival in cancer immunotherapy patients.

RESULTS

Yap inhibits CD4$^+$ and CD8$^+$ T cell activation

To study the role of Yap in T cells, we started by analyzing Yap levels in isolated primary CD4$^+$ and CD8$^+$ T cells from mice that were either unstimulated or activated with anti-CD3/CD28 coated beads for 24 hours. We observed that *Yap* RNA levels were dramatically increased upon stimulation of CD4$^+$ and CD8$^+$ T cells **(Figure 1A-B)**, suggesting a role for Yap in T cell activation. To gain insight into Yap function in T cells, we generated a mouse model whereby Yap deletion and EYFP expression are induced under the control of the CD4 promoter (Yap-loxP/loxP: LSL-EYFP; CD4-Cre, herein referred to as Yap-cKO). No systemic defects or gross phenotypes were observed in Yap-cKO mice housed in a barrier facility. Deletion of the *Yap* gene was achieved in both CD4$^+$ and CD8$^+$ T cells at the CD4$^+$CD8$^+$ double-positive stage of T cell development, as confirmed by EYFP expression **(Supplementary Figure 1A-B)**.

We next measured the surface levels of activation markers (CD69 and CD25) in Yap-cKO T cells. CD4$^+$ and CD8$^+$ T cells were isolated from wild type (WT) and Yap-cKO mouse spleens, activated, and levels of CD69 and CD25 were measured 24 hours later. We found increased levels of CD25 in Yap-cKO CD4$^+$ T cells (**Figure 1C-D**) and CD69 in Yap-cKO CD8$^+$ T cells (**Figure 1E-F**) compared to WT cells. CD4$^+$ and CD8$^+$ T cell proliferation was also measured, but no significant differences were observed between WT and Yap-cKO cells 72 hours post-stimulation (**Supplementary Figure 1E-F**). A major mechanism by which Yap functions is via association with the TEAD family of transcription factors in the nucleus, leading to recruitment to DNA that enables transcriptional regulation (Zhao et al., 2008). Verteporfin (also known as Visudine) is a small molecule that has been shown to repress the binding of Yap to TEADs, blocking nuclear Yap activity (Liu-Chittenden et al., 2012). We tested whether verteporfin treatment of WT CD4$^+$ and CD8$^+$ T cells affected the levels of the early activation markers CD71 and CD69 following activation. We observed that verteporfin treatment of CD4$^+$ and CD8$^+$ T cells increased CD71 (**Figure 1E-F**) and CD69 (**Figure 1G-H**) levels in a concentration-dependent manner, and that these increases were sustained up to 3 days post-activation. Consistent with the phenotype observed with Yap-cKO cells, verteporfin treatment did not significantly affect proliferation of

WT T cells 3 days post stimulation (**Supplementary Figure 1G**). Together, these data indicate that Yap is induced upon T cell activation and functions as a suppressor of T cell activation.

Yap inhibits CD4$^+$ T cell differentiation

Upon encountering their cognate antigen and receiving appropriate co-stimulation, naïve CD4$^+$ T cells become activated and differentiate into several subsets, including Th1, Th2 and Th17, to contribute to protective immunity or immunopathology depending on microenvironmental signals (Luckheeram et al., 2012). Signature transcription factors and effector cytokines define each subset: Th1 is defined by expression of T-BET and IFNγ, Th2 by GATA3 and IL-4, and TH17 by RORγt and IL-17. Given the wealth of evidence for Yap in stem cell regulation and the coordination of Yap inactivation with cell differentiation, we hypothesized that deletion of Yap alters the ability of CD4$^+$ T cells to differentiate. To test this hypothesis, we isolated naïve CD4$^+$ T cells from WT and Yap-cKO mice and cultured them under Th1, Th17 and Th2 polarizing conditions (Geng et al., 2017; Krebs et al., 2016). We found that Yap deleted CD4$^+$ T cells showed significantly enhanced Th1, Th17, and Th2 differentiation compared to WT cells, demonstrated by increased intracellular IFNγ, IL-17, and GATA3 expression, respectively (**Figure 2A-C**). Therefore, Yap acts as an inhibitor of CD4$^+$ T cell differentiation, and loss of Yap increases T cell responsiveness to polarizing cytokine and activation signals to drive differentiation.

Deletion of Yap in T cells promotes T cell infiltration into developing tumors and blocks tumor growth

Since we observed that Yap inhibits T cell activation and differentiation in vitro, we aimed to study the effect of Yap deletion in vivo. For this purpose, we chose to utilize the subcutaneous B16F10 mouse tumor model, due to its poorly immunogenic phenotype and highly immunosuppressive microenvironment that exhibits low T cell infiltration (Celik et al., 1983; Geiger et al., 1993; Turk et al., 2004; Wang et al., 1998). Since Yap deletion leads to enhanced activation and differentiation, we expected the possibility of increased immune infiltration and potentially tumor growth inhibition with B16F10 challenge in the Yap-cKO mice. We subcutaneously injected WT and Yap-cKO mice with B16F10 cells and monitored tumor growth. Yap deletion in T cells resulted in superior anti-tumor immunity, as evidenced by the significant delay in tumor growth in Yap-cKO compared to WT mice (**Figure 3A-B**). Given the strong correlation between CD8$^+$ T

cell tumor infiltration and patient survival and response to immunotherapy (Gajewski, 2007; Galluzzi et al., 2018; Galon et al., 2013; Galon et al., 2006; Galon et al., 2012; Hamid et al., 2011; Mlecnik et al., 2016; Mlecnik et al., 2011; Pages et al., 2005; Pages et al., 2018), we next investigated the extent of T cell infiltration in B16F10 tumors developed in WT versus Yap-cKO mice. Immunofluorescence microscopy analysis revealed that tumors from Yap-cKO mice were significantly more infiltrated with CD8$^+$ T cells at both the tumor center and tumor edge compared to WT mice (**Figure 3C**). Flow cytometry revealed more CD4$^+$ and CD8$^+$ T cells infiltrating Yap-cKO mouse tumors compared to WT ones (**Figure 3D-F**). Yap deficient T cells therefore have the capacity to limit tumor growth, which our observations suggest results from a combination of increased potential for activation, differentiation, effector function and tumor infiltration.

To directly compare the intrinsic tumor-infiltrating potential of WT vs Yap-cKO T cells in vivo, we isolated equal numbers of tdTomato$^+$ CD8$^+$ T cells from WT mice and EYFP$^+$ CD8$^+$ T cells from Yap-cKO mice and mixed these cells at a 1 to 1 ratio. Cell mixtures were intravenously injected into WT mice the same day as subcutaneous injection of B16F10 cells. Absolute numbers of tdTomato$^+$ (WT) and EYFP$^+$ (Yap-cKO) CD8$^+$ T cells were then measured in tumors after 15 days. Yap-cKO CD8$^+$ T cells were significantly infiltrated in tumors **(Figure 3G),** with nearly 30% of all CD8$^+$ tumor-infiltrating T cells being EYFP$^+$ Yap-cKO T cells compared to almost no tdTomato$^+$ WT T cells being observed in tumors **(Figure 3H)**.

Yap regulates global T cell responses in the local tumor microenvironment

We next aimed to define the Yap-regulated signaling networks in T cells that impacted tumor growth and T cell infiltration. CD4$^+$ and CD8$^+$ tumor infiltrating lymphocytes (TILs) were isolated from B16 tumors grown in WT or Yap-cKO mice, as well as CD4$^+$ and CD8$^+$ T cells from corresponding tumor draining lymph nodes (TDLNs), and analyzed by RNA-Seq **(Supplementary Tables 1-3)**. A large number of genes were differentially expressed in CD4$^+$ and CD8$^+$ TILs isolated from Yap-cKO mice compared to WT mice **(Figure 4A-B and Supplementary Figure 2A-B)**. Notably, T cells in the TDLNs showed markedly fewer gene expression changes with Yap deletion than those observed with TILs (**Figure 4C-D**). These data are consistent with Yap function being coordinated with T cell activation, and suggest that the

enhanced anti-tumor responses observed in Yap-cKO T cells are mediated by changes in cellular responses to local tumor microenvironment signals.

Hyper-enrichment analyses (Federico and Monti, 2019) of the gene expression changes identified in TILs from Yap-cKO mice showed an induction of genes related to T cell activation, differentiation, survival and migration (**Figure 4E-F**) (**Supplementary Table 3**). Yap-cKO TILs show enrichment of genes associated with T cell receptor signaling, genes that encode major co-stimulatory molecules, and genes downstream of TCR signaling **(Supplementary Figure 3A-B)**, suggesting enhanced responsiveness to TCR signals. Cytokine and cytokine receptor signaling gene sets are also enriched in Yap-cKO TILs, suggesting that cytokine production as well as responsiveness to cytokine receptor engagement are enhanced. Further, chemokine and chemokine receptor signaling gene sets are enriched in Yap-cKO TILs **(Supplementary Figure 3C-F)**, which likely contributes to their improved tumor infiltrating capacities. Subset defining transcription factors and cytokines associated with each of the major CD4$^+$ T cell phenotypes were all upregulated in Yap KO CD4$^+$ TILs **(Supplementary Figure 3G-H)**, suggesting that Yap deletion leads to enhanced naive CD4$^+$ T cell differentiation. Using unique upregulated genes after differentiation to each of the major T helper subsets (Stubbington et al., 2015), we show that Yap-cKO CD4$^+$ TILs are more skewed towards a Th2 and Treg phenotype compared to WT CD4$^+$ TILs **(Supplementary Figure 4A-D)**. As expected, gene set enrichment profiles identified in Yap-cKO TILs also included "Yap1 and TAZ stimulated gene expression" and "signaling by Hippo", which were repressed in both CD4$^+$ and CD8$^+$ TILs (**Figure 4G-H**).

Comparison of gene expression changes identified in CD4$^+$ and CD8$^+$ Yap-cKO TILs with clinical data from TCGA (Weinstein et al., 2013) showed a significant correlation with gene expression associated with T cell infiltration across a variety of human cancers (**Figure 4I**). Genes altered in Yap-cKO TILs were also significantly associated with patient survival across several cancers (**Figure 4J-L**), with increased Yap activity in TILs correlating with poor prognosis, suggesting that Yap function in T cells contributes to aggressive human cancer development and cancer immunosuppression.

DISCUSSION

We present the first comprehensive study on Yap function in both CD4[+] and CD8[+] T cells and report that Yap functions as a novel immunosuppressive factor. We found that disrupting Yap activity leads to enhanced T cell activation, augmented potential for differentiation, and increased tumor infiltration. Two major signals are necessary for T cell activation: signaling from engagement of the T cell receptor (TCR) with its cognate antigen:major histocompatibility (MHC) complex, and a second signal from the co-stimulatory receptor CD28 binding to ligands CD80 and CD86 (Bretscher, 1999; Smith-Garvin et al., 2009). Through recruitment of key effector kinases, phosphatases, and scaffold proteins, transcriptional programs are induced leading to production of cytokines and cytokine receptors, which importantly include IL-2 and CD25, the α chain of the heterotrimeric high-affinity IL-2 receptor. IL-2 binding to the IL-2 receptor, together with TCR signaling and co-stimulation elicit transcriptional changes resulting in proliferation and differentiation (Curtsinger et al., 1999). RNA-Sequencing analysis of WT vs Yap-cKO TILs revealed that CD3, CD28, CD80 and CD86 receptor molecules, kinases, phosphatases, and scaffold proteins are all upregulated with Yap deletion, offering a mechanism for enhanced activation of CD4[+] and CD8[+] T cells.

Yap has both TEAD-dependent and TEAD-independent mechanisms of action. We demonstrated that enhanced CD4[+] and CD8[+] T cell activation could be observed *in vitro* using either Yap-cKO cells or treatment with verteporfin, a reported inhibitor of Yap-TEAD activity, during activation. The dose-dependent effect of verteporfin suggests that Yap suppresses T cells at least partially through TEAD-dependent mechanisms. Interestingly, the effect of Yap deletion or TEAD-dependent inhibition by verteporfin induces *in vitro* activation of T cells, but not proliferation. This is consistent with verteporfin-induced increases in early activation markers. Previous studies have demonstrated proliferation has a sharp, switch-like threshold for T cell receptor signal to elicit proliferation (Au-Yeung et al., 2014), and our experimental conditions provide optimal signaling above this threshold. Expression levels of activation markers have been directly linked to TCR signal strength, such as CD69 levels being directly regulated by affinity and dose of TCR ligand (Allison et al., 2016) and CD71 levels being directly linked to mTOR activation level

(Zheng et al., 2007). Our observations suggest that the effects of Yap may link TCR signal strength to negative feedback via a TEAD-dependent mechanism.

The rapid induction of Yap post-activation suggests that there are no Yap-specific inducers in the tumor, but rather that the signals driving Yap expression are shared canonical signals of T cell activation. Therefore, Yap is likely part of a normal negative feedback mechanism for T cell activation similar to inhibitory checkpoint molecules, and inhibition of Yap must be timed before or during T cell activation. Our observations interestingly also indicate that Yap plays a prominent role only following activating signals from the microenvironment. This is highlighted by the large number of genes impacted in Yap-deleted CD4$^+$ and CD8$^+$ T cells isolated from TILs as compared to TDLNs, and is consistent with our data showing Yap levels increase after T cell activation. These data suggest that therapeutic inhibition of Yap in T cells may have fewer side effects compared to strategies such as checkpoint blockade, since the regulatory activity of Yap is synchronized with T cell activation and appears specific to sites of active T cell priming. Yap connects a variety of extracellular stimuli into intracellular cues that informs the cell of its own structural features (actin cytoskeleton, polarity, cell shape) as well as its location and surroundings (mechanical, adhesion, ECM), instructing cellular survival, proliferation, differentiation and fate. Therefore, a better understanding of how Yap is regulated by these signals in T cells may reveal new insights into immunomodulatory mechanisms.

Regulation of T cell activation by the microenvironment through polarizing cytokines allows for diverse, context-specific differentiation and functional diversity in CD4$^+$ T cells (Luckheeram et al., 2012). We observe that Yap-cKO CD4$^+$ T cells have increased capacity to differentiate towards Th1, Th2, and Th17 phenotypes. These observations are important as they indicate Yap does not preferentially bias T cell differentiation, but instead permits responses to local microenvironmental cues. This strongly suggests that Yap inhibition may better enable responses to current immunotherapy strategies. Clues into Yap-regulated events are embedded in our RNA-Sequencing of Yap-cKO TILs, which show broad changes in T cell responses. Genes regulated by Yap include several key effectors of T cell activation, including those controlling JAK-STAT signaling that modulate the polarization of naïve CD4s into T helper subsets (Seif et al., 2017). Notably, Yap-regulated genes are also enriched for those regulated by the TGFβ and Wnt pathways, which have

important pleotropic roles in T cell biology (Batlle and Massague, 2019; Chae and Bothwell, 2018). Given the known convergence of Yap with these immunomodulatory pathways (Varelas and Wrana, 2012) it is likely Yap directs their transcriptional targets and signal strength.

An important finding of our studies is that Yap-cKO mice show delayed tumor growth in conjunction with increased infiltration and expression of effector molecules. The well-characterized B16F10 melanoma model generates tumors that are characterized as immune deserts (Celik et al., 1983; Chen and Mellman, 2017; Geiger et al., 1993; Turk et al., 2004; Wang et al., 1998). In immune deserts, T cells are completely excluded from the tumor microenvironment due to suppressed tumor immunogenicity and insufficient T cell priming, co-stimulation, and activation (Gajewski et al., 2013). Patients with immune desert tumors show reduced survival, and are not responsive to immune checkpoint inhibitor (ICI) strategies. The ability for Yap-cKO CD4$^+$ and CD8$^+$ T cells to become activated and infiltrate tumors that normally inhibit T cell infiltration is therefore very notable and important, as these phenotypes are key for combating tumor growth in human cancer patients (Gajewski, 2015). Adoptive transfer of Yap-cKO CD8$^+$ T cells showed that CD8$^+$ T cells have intrinsically enhanced tumor infiltration capacity compared to WT host CD8$^+$ T cells. These data suggest that one of the early events in Yap-mediated tumor immunosuppression may be exclusion of CD8$^+$ T cells from the tumor. In mice with Yap-cKO T cells, tumor cell killing by early infiltration of CD8$^+$ T cells may elicit 1) tumor antigen release for enhanced intratumoral T cell priming and 2) increased intratumoral activation of CD8$^+$ T cells, causing chemokine secretion and recruitment of CD4$^+$ T cells and other immune cells. The significant correlation of CD8$^+$ and CD4$^+$ Yap-cKO gene signatures with tumor T cell infiltration in TCGA data suggest that Yap represses CD8$^+$ and CD4$^+$ T cell migration and tumor infiltration in diverse human cancers, and highlights Yap in T cells as a target for broad cancer therapy.

Collectively, our data show that Yap is an important immunosuppressor and suggest that inhibition of Yap activity in T cells could have important clinical implications in T cell therapies against cancer and other diseases. Insights from our study therefore motivate future investigation of YAP inhibition to enhance anti-tumor responses and the broader role of Yap in T cell regulation in other contexts.

MATERIALS & METHODS

Mouse Strains and Genotyping

Yap-loxP/loxP mice provided by Dr. Jeff Wrana and previously described (Reginensi et al., 2013) were backcrossed to the C57BL/6 background for 10 generations, and bred with the Tg(Cd4-cre)1Cwi (Jax: 022071) (Lee et al., 2001) and LSL-EYFP (Jax: 006148) (Srinivas et al., 2001) lines to derive Yap-loxP/loxP; LSL-EYFP; CD4-Cre mice. For adoptive cell transfers, CD4-cre mice were crossed with LSL-tdTomato mice (Jax: 007914) (Madisen et al., 2010). All experiments were performed with 6-10 weeks old mice. Animal protocols and study designs were approved by Boston University School of Medicine and UMBC. Mice were maintained in pathogen-free facilities at BUMC and UMBC and were PCR genotyped using published protocols (Lee et al., 2001; Madisen et al., 2010; Reginensi et al., 2013; Srinivas et al., 2001).

Cell culture and mouse tumor challenges

B16F10 cells were cultured in DMEM supplemented with glucose, L-glutamine, sodium pyruvate, 10% FBS, penicillin and streptomycin. Cells were split once they reached 70% confluency and were not used for mouse challenge past a fifth passage. T cells were cultured in RPMI 1640 supplemented with 10% FBS, 1 mM sodium pyruvate, 50 µM β-ME, penicillin, streptomycin, 2 mM L-glutamine, 100 mM non-essential amino acids, 5 mM HEPES free acid and β-mercaptoethanol. For tumor inoculations, 5×10^4 B16F10 cells were injected subcutaneously on the right flank of each mouse on day 0. Tumor volume was estimated using the formula $(L \times W^2)/2$. Survival endpoint was reached once the tumors measured 500mm^3, around day 15. Mice were euthanized by isoflurane inhalation and subsequent cervical dislocation and tumors were harvested for further prospecting.

T cell isolation

Spleens from WT or YAP-cKO mice were pushed through a 70µm mesh (Falcon) using an insulin syringe plunger and washed with PBS. Cells were treated with ACK red blood cell lysis buffer and splenocyte single cell suspensions were prepared for magnetic separation or stained for sorting by flow cytometry. CD4$^+$ or CD8$^+$ T cell enrichment was performed using magnetic beads (Miltenyi

Biotec or STEMCELL Technologies). Naïve CD4$^+$ T cells were isolated using a naive CD4$^+$ T cell isolation kit (Miltenyi Biotec).

Adoptive cell transfers

For the adoptive cell transfers, EYFP$^+$ Yap-cKO CD8$^+$ T cells were mixed 1:1 with dtTomato$^+$ WT CD8$^+$ T cells and injected intravenously into 8-week-old WT C57BL/6 mice (Taconic) on day 0. Mice also received a subcutaneous injection of 5x10^4 B16F10 cells on the same day. On day 15 of tumor growth, tumors were harvested and stained with dead cell dye, CD45, CD3, CD4 and CD8 antibodies for flow cytometric analysis.

Tumor digestion

B16 tumors from WT and YAP-cKO mice were dissected, mechanically disrupted and digested in serum free media containing 2mg/ml collagenase type I (Worthington) and DNase I (Sigma) for 30 minutes at 37°C in a rotator. Tumor digests were then passed through a 70μm mesh (Falcon) using an insulin syringe plunger and washed with PBS. Cells were treated with ACK red blood cell lysis buffer (Gibco) and tumor single cell suspensions were prepared for staining and analysis by flow cytometry. For determining absolute numbers of tumor infiltrating T cells by flow cytometry, AccuCount fluorescent particles (Spherotech) were added to the tumor digests.

Flow cytometry

Isolated splenocytes or tumor digests were washed with PBS and stained with the LIVE/DEAD fixable near-IR dead cell stain kit (Invitrogen). Cells were then washed with stain buffer (BD), resuspended in stain buffer containing Fc block (BD) and incubated for 5 minutes at 4°C. Surface antibodies were added in predetermined concentrations and cells were incubated for 30 minutes at 4°C or 15 minutes at room temperature, before being washed with BD stain buffer and resuspended in PBS for flow cytometric analysis. For intracellular cytokine staining, cells were fixed and permeabilized using the BD Cytofix/Cytoperm Fixation/Permeabilization kit (BD) after dead cell dye and surface staining. For transcription factor staining, the eBioscience Foxp3/ Transcription factor staining buffer set was used (eBioscience). Flow cytometry analyses were performed on BD LSRII at Boston University School of Medicine Flow Cytometry Facility or at the University of

Maryland School of Medicine Center for Innovative Biomedical Resources, Flow Cytometry Shared Service and analyzed by FlowJo (TreeStar).

T cell activation and proliferation assays

CD4$^+$ or CD8$^+$ T cells were isolated using magnetic beads as described above. Cells were stained using the CellTrace Violet or CFSE Cell Proliferation Kit (Life Technologies). Briefly, purified cells were washed with PBS and incubated with CellTrace dye for 20 minutes at 37°C protected from light. After 20 minutes, complete RMPI medium was added to the cell suspension and the cells were incubated 5 minutes further before being washed and resuspended in complete RPMI medium. Cells were cultured in 96 well plates at 1x10^5 cells per well and were stimulated using αCD3/CD28 dynabeads (Gibco) at a 1:1 ratio with T cells. On days 1 and 3, cells were stained with dead cell dye as well as lineage and activation markers CD3, CD4, CD8, CD69, CD71, and CD25 and proliferation was measured at the same time.

CD4$^+$ T cell in vitro differentiation

For CD4$^+$ T cell in vitro differentiations into Th1, Th2 and Th17 subsets, naïve CD4$^+$ T cells were enriched using magnetic beads (Miltenyi Biotec). Purified cells were plated at 1x10^5 cells per well on a 96 well plate coated with 10μg/ml αCD3 (Biolegend), and cultured with 2μg/ml soluble αCD28 (Biolegend). The following conditions were specific to each differentiation regime: Th1, 10ng/ml IL-12 and 10μg/ml αIL-4; Th2, 50ng/ml IL-4, 10μg/ml αIFNγ and 10μg/ml αIL-12; Th17, 50ng/ml IL-6, 20ng/ml IL-1β, 5ng/ml IL-23, 1ng/ml TGFβ, 12μg/ml αIFNγ and 10μg/ml αIL-4. Cells were cultured for 5 days, before being stimulated with 50ng/ml PMA (Sigma, P1585) and 1μg/ml ionomycin (Sigma, I0634) for 6 hours at 37 °C in the presence of Golgistop (monensin, BD) or Golgiplug (brefeldin, BD) added after the first 30 minutes of stimulation. Cells were stained with dead cell dye, surface markers (CD3, CD4, CD8, CD25) and intracellular cytokines (IFNγ, IL-17) or transcription factors (GATA3) as described above.

Immunofluorescence microscopy

Harvested B16 tumors were fixed overnight in PLP fixative, followed by incubation in 15% and 30% sucrose. Tumors were embedded in OCT and frozen. Cryosections were cut at 5μm thickness

and stored at -20°C. On the day of staining, slides were air dried for 1 hour and fixed in acetone for 10 minutes. Slides were washed with PBS and blocked with PBS containing 10% donkey serum, 0.05% sodium azide, 0.5% triton X-100 and 0.2% BSA for 1 hour at room temperature. Slides were stained with rat anti-mouse CD8 (clone CT-CD8a, Fisher) for 2 hours at room temperature, and were subsequently washed with PBS containing 1% Tween. Secondary antibody was diluted in blocking buffer and applied for 1 hour at room temperature (donkey anti-rat alexa 647, Jackson Immuno Research Labs). Slides were washed and mounted in ProLong antifade reagent with DAPI (Life Technologies). Images were acquired using an AxioObserver D1 equipped with a X-Cite 120LED System.

Quantitative real-time PCR (qPCR)

RNA was extracted using Rneasy Mini Kit (Qiagen), and 1μg was used to generate cDNA using an iScript cDNA Synthesis Kit (Biorad). Taqman primers for mouse GAPDH and YAP (Life Tech) were mixed with cDNA and Taqman Universal Master Mix II (Life Tech) and C_T values were normalized to unstimulated controls.

Sample preparation for RNA-Seq

B16 tumors from WT and YAP-cKO mice were digested as described above. Tumors were subsequently stained with dead cell dye and antibodies against CD45, CD3, CD4, and CD8, and $CD4^+$ and $CD8^+$ TILs were sorted from each tumor using a BDFACSARIA instrument. $CD4^+$ and $CD8^+$ T cells were sorted from WT and YAP-cKO tumor draining lymph nodes, as well. Cells were sorted into TRIzol LS reagent (Invitrogen) and RNA was isolated using a miRNeasy micro kit (Qiagen).

Transcriptomic analyses and gene expression signature extraction

RNA quality was evaluated using Agilent Bioanalyzer 2100 Eukaryote Total RNA Pico chips. RNA-Seq libraries were prepared using the SMART-Seq v4 Ultra Low Input RNA Kit (Takara, 634889) from total RNA, following the manufacturer's protocol. Libraries were then sequenced on a HiSeq 4000 using 75-bp paired end reads to an average depth of $22,445,650 \pm 240,398$ reads (SEM). Transcript abundance estimates were quantified using Salmon to mouse reference transcriptome from assembly GRCm38 (mm10), aggregated to gene level for UCSC-annotated

genes using tximport, and DESeq2 was used to calculate normalized counts and differential expression (Love et al., 2014; Patro et al., 2017). CD4 and CD8 up/down gene signatures were generated through differential expression analysis via DESeq2. Differentially expressed genes (DEG) between CD4 vs. WT and CD8 vs. WT were defined as $\log2(FC) > 1$ (up) or $\log2(FC) < -1$ (down) and FDR < 0.05. The top 25 differentially expressed genes for CD4$^+$ and CD8$^+$ T cells were visualized with a heatmap combined with a barplot annotation. The heatmap cells represent the log normalized expression values for each sample and include the top 25 upregulated and top 25 downregulated genes. Each row is accompanied by a bar representing the log-fold change in gene expression (KO/WT). Plots were generated using the *ComplexHeatmap* software package available in R.

Analysis of gene expression signatures in TCGA datasets

The activation of CD4/8 up/down signatures was calculated with Gene Set Variation Analysis (GSVA) (Hanzelmann et al., 2013) in primary tumor samples across multiple TCGA RNA-Seq datasets. Signature activation was summarized by the sum of activation of the upregulated signature and inactivation of the downregulated signature. For example, $CD4_{sig}$ activity = $GSVA(CD4_{up\ sig}) - GSVA(CD4_{down\ sig})$. Heatmaps for all TCGA analyses were generated with the *pheatmap* software package available in R. TCGA data includes count matrices generated with STAR2/HTSeq downloaded from Genomic Data Commons (GDC) Data Portal. Each count matrix was normalized by relative log expression (RLE) with DESeq2. This analysis was constrained to samples where survival information was available and immune infiltration could be estimated with Tumor Immune Estimation Resource (TIMER) (Li et al., 2016). The average T Cell infiltration per sample was estimated with TIMER in each TCGA dataset, represented as a single heatmap cell. This average was measured separately for CD4 T Cell and CD8 T Cell infiltration. Additionally, these values were summed to observe their additive effect.

For correlation of signatures with T cell infiltration, CD4/8 signature activation was correlated with the sum of CD4$^+$ T cell and CD8$^+$ T cell infiltration in each TCGA dataset estimated by TIMER (Li et al., 2016). The heatmap is colored by the correlation coefficient and the text of each cell is the adjusted p-value of the correlation. For survival analysis, the CD4/8 signature activation was used to stratify patients across TCGA datasets into high or low activated groups separated by

the mean. For each data set, Kaplan-Meier survival plots were generated and the results were summarized with a heatmap. The color of each cell corresponds to the following categories. Dark blue in the graphs signifies the average survival probability for patients with low activity of the signature is higher while red signifies the average survival probability for patients with high activity of the signature is higher. Light blue and light orange signify the same groups with higher survival probability respectively, however the difference is not significant. The text of each cell is the p-value for the survival estimation. The distribution of p-values arising from the multiple survival analyses for each signature across TCGA datasets was compared to a uniform distribution using a Kolmogorov-Smirnov test.

ACKNOWLEDGEMENTS

We thank Dr. Jeffrey Wrana for sharing the Yap-loxP mice. We acknowledge support from the Boston University Flow Cytometry core and the University of Maryland School of Medicine Center for Innovative Biomedical Resources, Flow Cytometry Shared Service and the Institute for Genome Sciences for RNA sequencing. X.V. was funded by a grant from the NIH National Heart Lung and Blood Institute (R01HL124392). G.L.S. is funded by in part by the UMGCC P30 grant under award number P30 CA134274 from the National Cancer Institute, NIH. EMS was supported in part by the Nathan Schnaper Program (NIH R25CA186872). We disclose that ES, GLS, and XV are inventors on a pending patent application related to the targeting Yap as an immunomodulatory therapy.

FIGURE LEGENDS

Figure 1. Yap RNA expression is induced upon T cell activation resulting in suppression of T cell activation. WT and YAP-cKO CD4$^+$ and CD8$^+$ T cells were isolated from mouse spleens and stimulated with αCD3 and αCD28 coated magnetic beads to test expression of YAP mRNA and activation markers. Activation marker expression was also tested for WT CD4$^+$ and CD8$^+$ T cells treated with increasing concentrations of Verteporfin under IL-2, αCD3 and αCD28 stimulation. Statistical differences were determined by using a Student's t-test (a-d) or one-way repeated-measures ANOVA with post-hoc test for linear trend with increasing verteporfin (e-h).

a. YAP mRNA expression by CD4$^+$ T cells isolated from WT or YAP-cKO mice 24 hours post CD3/CD28 stimulation (n=3/group).

b. YAP mRNA expression by CD8$^+$ T cells isolated from WT or YAP-cKO mice 24 hours post CD3/CD28 stimulation (n=3/group).

c. CD25 expression on WT and YAP-cKO CD4$^+$ T cells 24 hours post CD3/CD28 stimulation (n=3/group).

d. CD69 expression on WT and YAP-cKO CD8$^+$ T cells 24 hours post CD3/CD28 stimulation (n=3/group).

e. CD71 expression on WT CD4$^+$ T cells three days post IL-2 and CD3/CD28 stimulation and increasing concentration of verteporfin (n=4/group).

f. CD71 expression on WT CD8$^+$ T cells three days post IL-2 and CD3/CD28 stimulation and increasing concentration of verteporfin (n=4/group).

g. CD69 expression on WT CD4$^+$ T cells three days post IL-2 and CD3/CD28 stimulation and increasing concentration of verteporfin (n=4/group).

h. CD69 expression on WT CD8$^+$ T cells three days post IL-2 and CD3/CD28 stimulation and increasing concentration of verteporfin (n=4/group).

Figure 2. Deletion of YAP in CD4$^+$ T cells results in increased IFNγ, IL-17 and GATA3 expression under Th1, Th17 and Th2 polarizing conditions, respectively. Naïve CD4$^+$ T cells from WT and YAP-cKO mice were isolated using magnetic beads. WT and YAP KO T cells were cultured under Th1, Th17 or Th2 skewing conditions for 5 days in the presence of CD3 and CD28 antibodies. On Day5, IFNγ, IL-17 and GATA3 expression was measured using flow cytometry. Statistical differences were determined by using a Student's t-test.

a. IFNγ and IL-17 expression in WT and YAP-cKO CD4$^+$ T cells under Th1 polarizing conditions (n=3/group).

b. IFNγ and IL-17 expression in WT and YAP-cKO CD4$^+$ T cells under Th17 polarizing conditions (n=3/group).

c. IFNγ and GATA3 expression in WT and YAP-cKO CD4$^+$ T cells under Th2 polarizing conditions (n=3/group).

Figure 3. T cell specific deletion of YAP results in reduced B16 tumor growth and enhanced T cell tumor infiltration. Mice were challenged with 5x10^4 B16F10 cells injected subcutaneously on the right flank. Some mice received adoptive cell transfer of WT and YAP-cKO CD8$^+$ T cells. Tumor growth was monitored over the course of 15 days, until the maximum size of the tumors reached 500mm^3. Tumors were harvested for immunofluorescence or flow cytometric analysis. Statistical differences were determined by using a Student's t-test.

a. B16 tumor growth curve of WT and YAP-cKO mice (n=9/group).

b. Tumor weight on Day 15 from B16 tumors derived from WT and YAP-cKO mice (n=9/group).

c. CD8$^+$ T cell immunofluorescence on day 15 of tumor growth.

d. Absolute numbers of CD3$^+$ TILs from WT and YAP-cKO B16 tumors. Tumors were harvested on Day 15, stained with antibodies against CD45, CD3, CD4 and CD8 and analyzed using flow cytometry (n=5/WT mice, n=4/ YAP-cKO mice).

e. Absolute numbers of CD4$^+$ TILs from WT and YAP-cKO B16 tumors, prepared as in Fig4d (n=5/WT mice, n=4/ YAP-cKO mice).

f. Absolute numbers of CD8$^+$ TILs from WT and YAP-cKO B16 tumors, prepared as in Fig4d (n=5/WT mice, n=4/ YAP-cKO mice).

g. Absolute numbers of dTom$^+$ and EYFP$^+$ WT vs YAP-cKO CD8$^+$ T cells in C57BL/6 tumors. WT dTom$^+$ CD8$^+$ T cells were mixed 1:1 with YAP-cKO EYFP$^+$ CD8$^+$ T cells prior to being injected into WT C57BL/6 mice. Subsequently, mice were injected subcutaneously with B16F10 melanoma cells and absolute number of infiltrating T cells was determined on D15 by flow cytometry (n=5/group).

h. Percentage of dTom$^+$ and EYFP$^+$ CD8$^+$ T cells out of total tumor infiltrating CD8$^+$ T cells (n=5/group).

Figure 4. RNA-Seq analysis of WT and CD4Cre YAP KO CD4$^+$ and CD8$^+$ B16 TILs uncovers YAP as a negative regulator of T cell activation, differentiation, cytokine signaling and migration.

a. WT vs YAP-cKO CD4$^+$ TIL DEG heatmap.

b. WT vs YAP-cKO CD8$^+$ TIL DEG heatmap.

c. WT vs YAP-cKO CD4$^+$ TDLN DEG heatmap.

d. WT vs YAP-cKO CD8$^+$ TDLN DEG heatmap.

e. Hyper-enrichment analysis shows enrichment of induced gene sets observed in CD4$^+$ YAP-cKO TILs.

f. Hyper-enrichment analysis shows enrichment of induced gene sets observed in CD8$^+$ YAP-cKO TILs.

g. Hyper-enrichment analysis shows enrichment of repressed gene sets observed in CD4$^+$ YAP-cKO TILs.

h. Hyper-enrichment analysis shows enrichment of repressed gene sets observed in CD8$^+$ YAP-cKO TILs.

i. Gene expression changes identified in YAP-cKO CD4$^+$ and CD8$^+$ TILs correlate with genes reflecting tumor infiltration across many cancers in TCGA data. The heatmap is colored by the coefficient and the text of each cell represents the adjusted p-value of the correlation.

j. Gene expression changes identified in YAP-cKO CD4$^+$ and CD8$^+$ T cells correlate with patient survival data available in TCGA across several cancers. Blue: the average survival probability for patients with low activity of the signature is higher. Red: the average survival probability for patients with high activity of the signature is higher. Yellow: neither group has a higher probability of survival. The text of each cell is the p-value for the survival estimation. The distribution of p-values arising from the multiple survival analyses for each signature across TCGA datasets was compared to a uniform distribution using a Kolmogorov-Smirnov test.

k. Kaplan-Meier Survival analysis showing the average survival probability of patients with lung adenocarcinoma that show low versus high YAP activity derived from the YAP-cKO CD4$^+$ gene expression signature.

l. Kaplan-Meier Survival analysis showing the average survival probability of patients with lung adenocarcinoma that show low versus high YAP activity derived from the YAP-cKO CD8$^+$ gene expression signature.

Supplementary Figure 1. Yap deletion or inhibition does not affect T cell proliferation. CD4$^+$ and CD8$^+$ T cells were isolated from WT or YAP-cKO mice and screened of EYFP expression as well as activation marker expression and proliferation after αCD3 and αCD28 stimulation. Proliferation was also tested for WT CD4$^+$ and CD8$^+$ T cells treated with increasing concentrations of Verteporfin under IL-2, αCD3 and αCD28 stimulation. Statistical differences were determined by using a Student's t-test, with ns indicating not-significant.

a. EYFP expression by flow cytometry on CD4$^+$ cells isolated from WT or YAP-cKO mouse spleens.
b. EYFP expression by flow cytometry on CD8$^+$ cells isolated from WT or YAP-cKO mouse spleens.
c. CD69 expression on WT and YAP-cKO CD4$^+$ T cells 24 hours post CD3/CD28 stimulation (n=3/group).
d. CD25 expression on WT and YAP-cKO CD8$^+$ T cells 24 hours post CD3/CD28 stimulation (n=3/group).
e. WT vs YAP-cKO CD4$^+$ T cell proliferation (n=3/group).
f. WT vs YAP-cKO CD8$^+$ T cell proliferation (n=3/group).
g. Proliferation of DMSO vs verteporfin treated WT CD4$^+$ and CD8$^+$ T cells (representative of 4 independent experiments).

Supplementary Figure 2. Top 25 upregulated and downregulated genes with YAP deletion in CD4$^+$ and CD8$^+$ TILs.
a. WT vs YAP-cKO CD4$^+$ TILs top and bottom DEG heatmap.
b. WT vs YAP-cKO CD8$^+$ TILs top and bottom DEG heatmap.

Supplementary Figure 3. Expression of genes related to T cell activation and co-stimulation, chemokines and chemokine receptors, T helper subset defining cytokines and transcription factors are upregulated in YAP-cKO CD4$^+$ and CD8$^+$TILs.
a. Log$_{10}$(normalized RNA-Seq counts +1) of T cell activation related genes in WT versus YAP-cKO CD8$^+$ TILs.

b. Log$_{10}$(normalized RNA-Seq counts +1) of T cell activation related genes in WT versus YAP-cKO CD4$^+$ TILs.

c. Log$_{10}$(normalized RNA-Seq counts +1) of chemokine genes in WT versus YAP-cKO CD8$^+$ TILs.

d. Log$_{10}$(normalized RNA-Seq counts +1) of chemokine receptor genes in WT versus YAP-cKO CD8$^+$ TILs.

e. Log$_{10}$(normalized RNA-Seq counts +1) of chemokine genes in WT versus YAP-cKO CD4$^+$ TILs.

f. Log$_{10}$(normalized RNA-Seq counts +1) of chemokine receptor genes in WT versus YAP-cKO CD4$^+$ TILs.

g. Log$_{10}$(normalized RNA-Seq counts +1) of T helper subset defining cytokines in WT versus YAP-cKO CD4$^+$ TILs.

h. Log$_{10}$(normalized RNA-Seq counts +1) of T helper subset defining transcription factors in WT versus YAP-cKO CD4$^+$ TILs.

Significant differences were determined by a Student's *t* test; *, $p<0.05$; **, $p<0.01$; ***, $p<0.001$.

Supplementary Figure 4. YAP-cKO vs WT CD4$^+$ TILs are skewed towards TH2 and Treg signatures compared to WT.

a. Heatmap of statistically significant differentially expressed TH1 genes in WT vs YAP-cKO CD4$^+$ TILs.

b. Heatmap of statistically significant differentially expressed TH2 genes in WT vs YAP-cKO CD4$^+$ TILs.

c. Heatmap of statistically significant differentially expressed TH17 genes in WT vs YAP-cKO CD4$^+$ TILs.

d. Heatmap of statistically significant differentially expressed Treg genes in WT vs YAP-cKO CD4$^+$ TILs.

REFERENCES

Allison, K.A., Sajti, E., Collier, J.G., Gosselin, D., Troutman, T.D., Stone, E.L., Hedrick, S.M., and Glass, C.K. (2016). Affinity and dose of TCR engagement yield proportional enhancer and gene activity in CD4+ T cells. Elife *5*.

Au-Yeung, B.B., Zikherman, J., Mueller, J.L., Ashouri, J.F., Matloubian, M., Cheng, D.A., Chen, Y., Shokat, K.M., and Weiss, A. (2014). A sharp T-cell antigen receptor signaling threshold for T-cell proliferation. Proc Natl Acad Sci U S A *111*, E3679-3688.

Balatoni, T., Mohos, A., Papp, E., Sebestyen, T., Liszkay, G., Olah, J., Varga, A., Lengyel, Z., Emri, G., Gaudi, I.*, et al.* (2018). Tumor-infiltrating immune cells as potential biomarkers predicting response to treatment and survival in patients with metastatic melanoma receiving ipilimumab therapy. Cancer immunology, immunotherapy : CII *67*, 141-151.

Batlle, E., and Massague, J. (2019). Transforming Growth Factor-beta Signaling in Immunity and Cancer. Immunity *50*, 924-940.

Brentjens, R.J., Davila, M.L., Riviere, I., Park, J., Wang, X., Cowell, L.G., Bartido, S., Stefanski, J., Taylor, C., Olszewska, M.*, et al.* (2013). CD19-targeted T cells rapidly induce molecular remissions in adults with chemotherapy-refractory acute lymphoblastic leukemia. Science translational medicine *5*, 177ra138.

Bretscher, P.A. (1999). A two-step, two-signal model for the primary activation of precursor helper T cells. Proceedings of the National Academy of Sciences of the United States of America *96*, 185-190.

Brown, C.E., and Mackall, C.L. (2019). CAR T cell therapy: inroads to response and resistance. Nature reviews Immunology *19*, 73-74.

Celik, C., Lewis, D.A., and Goldrosen, M.H. (1983). Demonstration of immunogenicity with the poorly immunogenic B16 melanoma. Cancer research *43*, 3507-3510.

Chae, W.J., and Bothwell, A.L.M. (2018). Canonical and Non-Canonical Wnt Signaling in Immune Cells. Trends Immunol *39*, 830-847.

Chen, D.S., and Mellman, I. (2017). Elements of cancer immunity and the cancer-immune set point. Nature *541*, 321-330.

Cogdill, A.P., Andrews, M.C., and Wargo, J.A. (2017). Hallmarks of response to immune checkpoint blockade. British journal of cancer *117*, 1-7.

Curtsinger, J.M., Schmidt, C.S., Mondino, A., Lins, D.C., Kedl, R.M., Jenkins, M.K., and Mescher, M.F. (1999). Inflammatory cytokines provide a third signal for activation of naive CD4+ and CD8+ T cells. Journal of immunology (Baltimore, Md : 1950) *162*, 3256-3262.

Darvin, P., Toor, S.M., Sasidharan Nair, V., and Elkord, E. (2018). Immune checkpoint inhibitors: recent progress and potential biomarkers. Experimental & molecular medicine *50*, 165.

Dong, Y., Du, X., Ye, J., Han, M., Xu, T., Zhuang, Y., and Tao, W. (2009). A cell-intrinsic role for Mst1 in regulating thymocyte egress. Journal of immunology (Baltimore, Md : 1950) *183*, 3865-3872.

Du, X., Shi, H., Li, J., Dong, Y., Liang, J., Ye, J., Kong, S., Zhang, S., Zhong, T., Yuan, Z.*, et al.* (2014). Mst1/Mst2 regulate development and function of regulatory T cells through modulation of Foxo1/Foxo3 stability in autoimmune disease. Journal of immunology (Baltimore, Md : 1950) *192*, 1525-1535.

Federico, A., and Monti, S. (2019). hypeR: Geneset enrichment analysis based on hyper-geometric test. Bioconductor
.

Feng, Y., Roy, A., Masson, E., Chen, T.T., Humphrey, R., and Weber, J.S. (2013). Exposure-response relationships of the efficacy and safety of ipilimumab in patients with advanced melanoma. Clinical cancer research : an official journal of the American Association for Cancer Research *19*, 3977-3986.

Gajewski, T.F. (2007). Failure at the effector phase: immune barriers at the level of the melanoma tumor microenvironment. Clinical cancer research : an official journal of the American Association for Cancer Research *13*, 5256-5261.

Gajewski, T.F. (2015). The Next Hurdle in Cancer Immunotherapy: Overcoming the Non-T-Cell-Inflamed Tumor Microenvironment. Seminars in oncology *42*, 663-671.

Gajewski, T.F., Woo, S.R., Zha, Y., Spaapen, R., Zheng, Y., Corrales, L., and Spranger, S. (2013). Cancer immunotherapy strategies based on overcoming barriers within the tumor microenvironment. Current opinion in immunology *25*, 268-276.

Galluzzi, L., Chan, T.A., Kroemer, G., Wolchok, J.D., and Lopez-Soto, A. (2018). The hallmarks of successful anticancer immunotherapy. Science translational medicine *10*.

Galon, J., Angell, H.K., Bedognetti, D., and Marincola, F.M. (2013). The continuum of cancer immunosurveillance: prognostic, predictive, and mechanistic signatures. Immunity *39*, 11-26.

Galon, J., Costes, A., Sanchez-Cabo, F., Kirilovsky, A., Mlecnik, B., Lagorce-Pages, C., Tosolini, M., Camus, M., Berger, A., Wind, P.*, et al.* (2006). Type, density, and location of immune cells within human colorectal tumors predict clinical outcome. Science (New York, NY) *313*, 1960-1964.

Galon, J., Pages, F., Marincola, F.M., Angell, H.K., Thurin, M., Lugli, A., Zlobec, I., Berger, A., Bifulco, C., Botti, G.*, et al.* (2012). Cancer classification using the Immunoscore: a worldwide task force. Journal of translational medicine *10*, 205.

Geiger, J.D., Wagner, P.D., Cameron, M.J., Shu, S., and Chang, A.E. (1993). Generation of T-cells reactive to the poorly immunogenic B16-BL6 melanoma with efficacy in the treatment of spontaneous metastases. Journal of immunotherapy with emphasis on tumor immunology : official journal of the Society for Biological Therapy *13*, 153-165.

Geng, J., Yu, S., Zhao, H., Sun, X., Li, X., Wang, P., Xiong, X., Hong, L., Xie, C., Gao, J.*, et al.* (2017). The transcriptional coactivator TAZ regulates reciprocal differentiation of TH17 cells and Treg cells. Nature immunology *18*, 800-812.

Hamid, O., Schmidt, H., Nissan, A., Ridolfi, L., Aamdal, S., Hansson, J., Guida, M., Hyams, D.M., Gomez, H., Bastholt, L.*, et al.* (2011). A prospective phase II trial exploring the association between tumor microenvironment biomarkers and clinical activity of ipilimumab in advanced melanoma. Journal of translational medicine *9*, 204.

Hanzelmann, S., Castelo, R., and Guinney, J. (2013). GSVA: gene set variation analysis for microarray and RNA-seq data. BMC Bioinformatics *14*, 7.

Herbst, R.S., Soria, J.C., Kowanetz, M., Fine, G.D., Hamid, O., Gordon, M.S., Sosman, J.A., McDermott, D.F., Powderly, J.D., Gettinger, S.N.*, et al.* (2014). Predictive correlates of response to the anti-PD-L1 antibody MPDL3280A in cancer patients. Nature *515*, 563-567.

Katagiri, K., Imamura, M., and Kinashi, T. (2006). Spatiotemporal regulation of the kinase Mst1 by binding protein RAPL is critical for lymphocyte polarity and adhesion. Nature immunology *7*, 919-928.

Katagiri, K., Katakai, T., Ebisuno, Y., Ueda, Y., Okada, T., and Kinashi, T. (2009). Mst1 controls lymphocyte trafficking and interstitial motility within lymph nodes. The EMBO journal *28*, 1319-1331.

Krebs, C.F., Turner, J.E., Paust, H.J., Kapffer, S., Koyro, T., Krohn, S., Ufer, F., Friese, M.A., Flavell, R.A., Stockinger, B.*, et al.* (2016). Plasticity of Th17 Cells in Autoimmune Kidney Diseases. Journal of immunology (Baltimore, Md : 1950) *197*, 449-457.

Kumar, B.V., Connors, T.J., and Farber, D.L. (2018). Human T Cell Development, Localization, and Function throughout Life. Immunity *48*, 202-213.

Lee, D.W., Kochenderfer, J.N., Stetler-Stevenson, M., Cui, Y.K., Delbrook, C., Feldman, S.A., Fry, T.J., Orentas, R., Sabatino, M., Shah, N.N.*, et al.* (2015). T cells expressing CD19 chimeric antigen receptors for acute lymphoblastic leukaemia in children and young adults: a phase 1 dose-escalation trial. Lancet (London, England) *385*, 517-528.

Lee, P.P., Fitzpatrick, D.R., Beard, C., Jessup, H.K., Lehar, S., Makar, K.W., Perez-Melgosa, M., Sweetser, M.T., Schlissel, M.S., Nguyen, S.*, et al.* (2001). A critical role for Dnmt1 and DNA methylation in T cell development, function, and survival. Immunity *15*, 763-774.

Li, B., Severson, E., Pignon, J.C., Zhao, H., Li, T., Novak, J., Jiang, P., Shen, H., Aster, J.C., Rodig, S.*, et al.* (2016). Comprehensive analyses of tumor immunity: implications for cancer immunotherapy. Genome Biol *17*, 174.

Li, J., Du, X., Shi, H., Deng, K., Chi, H., and Tao, W. (2015). Mammalian Sterile 20-like Kinase 1 (Mst1) Enhances the Stability of Forkhead Box P3 (Foxp3) and the Function of Regulatory T Cells by Modulating Foxp3 Acetylation. The Journal of biological chemistry *290*, 30762-30770.

Liu-Chittenden, Y., Huang, B., Shim, J.S., Chen, Q., Lee, S.J., Anders, R.A., Liu, J.O., and Pan, D. (2012). Genetic and pharmacological disruption of the TEAD-YAP complex suppresses the oncogenic activity of YAP. Genes & development *26*, 1300-1305.

Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol *15*, 550.

Luckheeram, R.V., Zhou, R., Verma, A.D., and Xia, B. (2012). CD4(+)T cells: differentiation and functions. Clinical & developmental immunology *2012*, 925135.

Madisen, L., Zwingman, T.A., Sunkin, S.M., Oh, S.W., Zariwala, H.A., Gu, H., Ng, L.L., Palmiter, R.D., Hawrylycz, M.J., Jones, A.R.*, et al.* (2010). A robust and high-throughput Cre reporting and characterization system for the whole mouse brain. Nat Neurosci *13*, 133-140.

Maude, S.L., Frey, N., Shaw, P.A., Aplenc, R., Barrett, D.M., Bunin, N.J., Chew, A., Gonzalez, V.E., Zheng, Z., Lacey, S.F.*, et al.* (2014). Chimeric antigen receptor T cells for sustained remissions in leukemia. The New England journal of medicine *371*, 1507-1517.

Mlecnik, B., Bindea, G., Angell, H.K., Maby, P., Angelova, M., Tougeron, D., Church, S.E., Lafontaine, L., Fischer, M., Fredriksen, T.*, et al.* (2016). Integrative Analyses of Colorectal Cancer Show Immunoscore Is a Stronger Predictor of Patient Survival Than Microsatellite Instability. Immunity *44*, 698-711.

Mlecnik, B., Tosolini, M., Kirilovsky, A., Berger, A., Bindea, G., Meatchi, T., Bruneval, P., Trajanoski, Z., Fridman, W.H., Pages, F.*, et al.* (2011). Histopathologic-based prognostic factors of colorectal cancers are associated with the state of the local immune reaction. Journal of clinical oncology : official journal of the American Society of Clinical Oncology *29*, 610-618.

Mou, F., Praskova, M., Xia, F., Van Buren, D., Hock, H., Avruch, J., and Zhou, D. (2012). The Mst1 and Mst2 kinases control activation of rho family GTPases and thymic egress of mature thymocytes. The Journal of experimental medicine *209*, 741-759.

Murakami, N., and Riella, L.V. (2014). Co-inhibitory pathways and their importance in immune regulation. Transplantation *98*, 3-14.

Nehme, N.T., Schmid, J.P., Debeurme, F., Andre-Schmutz, I., Lim, A., Nitschke, P., Rieux-Laucat, F., Lutz, P., Picard, C., Mahlaoui, N.*, et al.* (2012). MST1 mutations in autosomal recessive

primary immunodeficiency characterized by defective naive T-cell survival. Blood *119*, 3458-3468.

Ni, X., Tao, J., Barbi, J., Chen, Q., Park, B.V., Li, Z., Zhang, N., Lebid, A., Ramaswamy, A., Wei, P.*, et al.* (2018). YAP Is Essential for Treg-Mediated Suppression of Antitumor Immunity. Cancer discovery *8*, 1026-1043.

Nishikimi, A., Ishihara, S., Ozawa, M., Etoh, K., Fukuda, M., Kinashi, T., and Katagiri, K. (2014). Rab13 acts downstream of the kinase Mst1 to deliver the integrin LFA-1 to the cell surface for lymphocyte trafficking. Science signaling *7*, ra72.

Pages, F., Berger, A., Camus, M., Sanchez-Cabo, F., Costes, A., Molidor, R., Mlecnik, B., Kirilovsky, A., Nilsson, M., Damotte, D.*, et al.* (2005). Effector memory T cells, early metastasis, and survival in colorectal cancer. The New England journal of medicine *353*, 2654-2666.

Pages, F., Mlecnik, B., Marliot, F., Bindea, G., Ou, F.S., Bifulco, C., Lugli, A., Zlobec, I., Rau, T.T., Berger, M.D.*, et al.* (2018). International validation of the consensus Immunoscore for the classification of colon cancer: a prognostic and accuracy study. Lancet (London, England) *391*, 2128-2139.

Patro, R., Duggal, G., Love, M.I., Irizarry, R.A., and Kingsford, C. (2017). Salmon provides fast and bias-aware quantification of transcript expression. Nature methods *14*, 417-419.

Pauken, K.E., and Wherry, E.J. (2015). Overcoming T cell exhaustion in infection and cancer. Trends Immunol *36*, 265-276.

Piccolo, S., Dupont, S., and Cordenonsi, M. (2014). The biology of YAP/TAZ: hippo signaling and beyond. Physiological reviews *94*, 1287-1312.

Reginensi, A., Scott, R.P., Gregorieff, A., Bagherie-Lachidan, M., Chung, C., Lim, D.S., Pawson, T., Wrana, J., and McNeill, H. (2013). Yap- and Cdc42-dependent nephrogenesis and morphogenesis during mouse kidney development. PLoS Genet *9*, e1003380.

Rizvi, N.A., Hellmann, M.D., Snyder, A., Kvistborg, P., Makarov, V., Havel, J.J., Lee, W., Yuan, J., Wong, P., Ho, T.S.*, et al.* (2015). Cancer immunology. Mutational landscape determines sensitivity to PD-1 blockade in non-small cell lung cancer. Science (New York, NY) *348*, 124-128.

Seif, F., Khoshmirsafa, M., Aazami, H., Mohsenzadegan, M., Sedighi, G., and Bahar, M. (2017). The role of JAK-STAT signaling pathway and its regulators in the fate of T helper cells. Cell communication and signaling : CCS *15*, 23.

Shi, H., Liu, C., Tan, H., Li, Y., Nguyen, T.M., Dhungana, Y., Guy, C., Vogel, P., Neale, G., Rankin, S.*, et al.* (2018). Hippo Kinases Mst1 and Mst2 Sense and Amplify IL-2R-STAT5 Signaling in Regulatory T Cells to Establish Stable Regulatory Activity. Immunity *49*, 899-914 e896.

Smith-Garvin, J.E., Koretzky, G.A., and Jordan, M.S. (2009). T cell activation. Annual review of immunology *27*, 591-619.

Srinivas, S., Watanabe, T., Lin, C.S., William, C.M., Tanabe, Y., Jessell, T.M., and Costantini, F. (2001). Cre reporter strains produced by targeted insertion of EYFP and ECFP into the ROSA26 locus. BMC Dev Biol *1*, 4.

Stubbington, M.J., Mahata, B., Svensson, V., Deonarine, A., Nissen, J.K., Betz, A.G., and Teichmann, S.A. (2015). An atlas of mouse CD4(+) T cell transcriptomes. Biology direct *10*, 14.

Tang, F., Gill, J., Ficht, X., Barthlott, T., Cornils, H., Schmitz-Rohmer, D., Hynx, D., Zhou, D., Zhang, L., Xue, G.*, et al.* (2015). The kinases NDR1/2 act downstream of the Hippo homolog MST1 to mediate both egress of thymocytes from the thymus and lymphocyte motility. Science signaling *8*, ra100.

Thaventhiran, J.E., Hoffmann, A., Magiera, L., de la Roche, M., Lingel, H., Brunner-Weinzierl, M., and Fearon, D.T. (2012). Activation of the Hippo pathway by CTLA-4 regulates the expression of Blimp-1 in the CD8+ T cell. Proceedings of the National Academy of Sciences of the United States of America *109*, E2223-2229.

Topalian, S.L., Hodi, F.S., Brahmer, J.R., Gettinger, S.N., Smith, D.C., McDermott, D.F., Powderly, J.D., Carvajal, R.D., Sosman, J.A., Atkins, M.B.*, et al.* (2012). Safety, activity, and immune correlates of anti-PD-1 antibody in cancer. The New England journal of medicine *366*, 2443-2454.

Turk, M.J., Guevara-Patino, J.A., Rizzuto, G.A., Engelhorn, M.E., Sakaguchi, S., and Houghton, A.N. (2004). Concomitant tumor immunity to a poorly immunogenic melanoma is prevented by regulatory T cells. The Journal of experimental medicine *200*, 771-782.

Ueda, Y., Katagiri, K., Tomiyama, T., Yasuda, K., Habiro, K., Katakai, T., Ikehara, S., Matsumoto, M., and Kinashi, T. (2012). Mst1 regulates integrin-dependent thymocyte trafficking and antigen recognition in the thymus. Nature communications *3*, 1098.

Varelas, X. (2014). The Hippo pathway effectors TAZ and YAP in development, homeostasis and disease. Development *141*, 1614-1626.

Varelas, X., and Wrana, J.L. (2012). Coordinating developmental signaling: novel roles for the Hippo pathway. Trends Cell Biol *22*, 88-96.

Wang, J., Saffold, S., Cao, X., Krauss, J., and Chen, W. (1998). Eliciting T cell immunity against poorly immunogenic tumors by immunization with dendritic cell-tumor fusion vaccines. Journal of immunology (Baltimore, Md : 1950) *161*, 5516-5524.

Weinstein, J.N., Collisson, E.A., Mills, G.B., Shaw, K.R., Ozenberger, B.A., Ellrott, K., Shmulevich, I., Sander, C., and Stuart, J.M. (2013). The Cancer Genome Atlas Pan-Cancer analysis project. Nature genetics *45*, 1113-1120.

Wherry, E.J., and Kurachi, M. (2015). Molecular and cellular insights into T cell exhaustion. Nat Rev Immunol *15*, 486-499.

Williams, M.A., and Bevan, M.J. (2007). Effector and memory CTL differentiation. Annual review of immunology *25*, 171-192.

Zhao, B., Ye, X., Yu, J., Li, L., Li, W., Li, S., Yu, J., Lin, J.D., Wang, C.Y., Chinnaiyan, A.M.*, et al.* (2008). TEAD mediates YAP-dependent gene induction and growth control. Genes & development *22*, 1962-1971.

Zheng, Y., Collins, S.L., Lutz, M.A., Allen, A.N., Kole, T.P., Zarek, P.E., and Powell, J.D. (2007). A role for mammalian target of rapamycin in regulating T cell activation versus anergy. J Immunol *178*, 2163-2170.

Zhou, D., Medoff, B.D., Chen, L., Li, L., Zhang, X.F., Praskova, M., Liu, M., Landry, A., Blumberg, R.S., Boussiotis, V.A.*, et al.* (2008). The Nore1B/Mst1 complex restrains antigen receptor-induced proliferation of naive T cells. Proceedings of the National Academy of Sciences of the United States of America *105*, 20321-20326.

Zhou, H., Luo, Y., Zhu, S., Wang, X., Zhao, Y., Ou, X., Zhang, T., and Ma, X. (2018). The efficacy and safety of anti-CD19/CD20 chimeric antigen receptor- T cells immunotherapy in relapsed or refractory B-cell malignancies:a meta-analysis. BMC cancer *18*, 929.

Figure 1

Figure 2

Figure 3

Figure 4

Supplementary Figure 1

Supplementary Figure 2

Supplementary Figure 3

Supplementary Figure 4