Abundance and diversity of resistomes differ between healthy human oral cavities and gut

Victoria R. Carr[1], Elizabeth Witherden[1], Sunjae Lee[1], Saeed Shoaie[1,2], Peter Mullany[3], Gordon B.

Proctor[1], David Gomez-Cabrero[1,4,5], David L. Moyes[1]

[1] Centre for Host-Microbiome Interactions, Faculty of Dentistry, Oral and Craniofacial Sciences,

King's College London, UK

[2] Science for Life Laboratory, KTH—Royal Institute of Technology, SE-171 21 Stockholm, Sweden

[3] Microbial Diseases, Eastman Dental Institute, University College London, UK

[4] Translational Bioinformatics Unit, NavarraBiomed, Departamento de Salud-Universidad Pública

de Navarra, Pamplona, 31008, Navarra, Spain

[5] Unit for Computational Medicine, Karolinska Institutet (L8:05). Karolinska University

Hospital, SE-171 76 Stockholm, Sweden.

Correspondence and requests for materials: victoria.carr@kcl.ac.uk, david.moyes@kcl.ac.uk

Keywords

resistome; antimicrobial resistance; microbiome; metagenomics; oral; gut

Abstract

The global threat of the "antimicrobial resistance apocalypse" that has arisen in recent years has

driven the use of high-throughput sequencing techniques to monitor the profile of resistance genes,

known as the "resistome", in microbial populations. The human oral cavity contains a poorly

explored reservoir of these genes, and little is known about their abundance and diversity, or how

their profile compares with antimicrobial resistance genes in the gut. Here we analyse the resistome

profiles of 788 oral cavities worldwide and compare these profiles with paired stool samples from

shotgun metagenomic data. We find country and body site-specific differences in the prevalence of

antimicrobial resistance genes, classes and mechanisms in oral and stool samples. However, the

abundance of these antimicrobial resistance classes do not correlate with antibiotic prescription

rates. A greater similarity was found in interpersonal resistomes between the same body sites than

intrapersonal resistomes across different body sites. Between individuals, the oral cavity contains

the highest and lowest abundances of specific antimicrobial resistance genes, but a lower diversity

of resistance genes compared to the gut, which is likely influenced by differences in microbial

composition and exposure to antimicrobial selection pressures. Co-occurrence analysis shows

contrasting ARG-species associations between saliva and stool samples. This is the first study to

date that characterises the oral cavity resistome worldwide, identifying its distinctive signatures

compared to the gut.Maintenance and persistence of antimicrobial resistance is likely to vary across

different body sites. Thus, we highlight the importance of characterising the resistome across body

sites to uncover the antimicrobial resistance potential in the human body.

Introduction

In recent years, antimicrobial resistance (AMR) has been highlighted as one of the biggest threats to

global health, food production and economic development[1]. Given this rapidly developing global

crisis, it is imperative that the current gaps in our understanding of the distribution, spread and

associations of all AMR factors are filled. AMR is most often conferred through the expression of

antimicrobial resistance genes (ARGs) that reduce a microbe's susceptibility to the effects of an

antimicrobial compound. As such, monitoring the abundance and diversity of these ARG profiles, or

the "resistome", has huge potential to increase our understanding of the spread and persistence of

AMR within a population. High-throughput next-generation sequencing technologies are beginning

to be used as tools for screening ARGs for potential surveillance of antimicrobial resistance

worldwide. Shotgun metagenomic data mapped against dedicated ARG reference databases are

providing a wealth of insight into the resistomes of human[2–7] and animal guts[8,9] as well as the wider

environment[10–13]. However, no large studies have, to date, attempted to characterise the resistome

profiles of the human oral cavity. Commensal microbes from the oral cavity harbouring ARGs have

considerable potential to lead to antimicrobial resistant infections at other body sites. For example,

ß-lactam, clindamycin, and erythromycin resistant strains of oral streptococci have caused

infections at distal body sites such as infective endocarditis[14].

Metagenomic studies of the oral cavity indicate that this site potentially contains a diverse range of

ARGs, including those encoding resistance to tetracycline, amoxycillin and gentamicin in saliva and

plaque samples[15,16]. Thus, oral ARGs appear to be natural features of the human oral cavity. The

presence of an oral resistome containing aminoglycoside, ß-lactam, macrolide, phenicol and

tetracycline ARGs in isolated Amerindian communities and ancient humans, indicates that the

presence of these genes is not dependent on antibiotic exposure and is an inherent feature of the oral

microbiome[17,18].

The oral microbial community faces unique ecological pressures, such as mechanical force,

nutritional availability, pH levels, oxidative stress and redox potential. Despite these continually

changing conditions, these communities have been shown to be relatively stable even after short-

term antibiotic exposure. Horizontal gene transfer (HGT) has been documented as an important

mechanism for the transfer and acquisition of ARGs within and between oral bacterial species[19,20].

The erythromycin resistance *mefA* and *mefE* genes have been found on the *MEGA* mobile genetic

element associated with Tn*916*-like conjugative transposons (also called integrative conjugative

elements ICE), and this has been implicated in conjugative transfer between viridans group

streptococci (VGS) and other streptococci[21]. Thus, the oral microbiome contains a long-standing

and mobile population of ARGs and is a significant reservoir for ARGs to be transferred to

pathogenic microbes.

Here, we derive and compare the oral and the gut resistomes from 788 and 386 shotgun

metagenomes, respectively, from healthy individuals from China[22], Fiji[23], the Philippines[24], Western

Europe[25–27] and the US[28]. We found country-specific differences in the proportion of saliva, dental

plaque and stool samples containing ARGs, ARG classes and mechanisms. We made up to 415

comparisons of oral resistomes with paired gut resistomes derived from stool shotgun metagenomes

from the same individuals, showing the oral resistome contains the highest and lowest abundances

of ARGs, but a lower diversity of ARGs than the gut resistome. Overall, these results demonstrate

the requirement for wider AMR surveillance studies at different body sites, including the oral

cavity, to understand the composition of the resistome across different human microbial habitats.

86   **Results**

87 **Country and body site-specific differences in resistomes**

88
To establish the incidence of ARGs in oral as well as stool metagenomes collected from various

regions, metagenomes were mapped and quantified against the Comprehensive Antibiotic

Resistance Database (CARD)[29]. Saliva samples were only available from China, Fiji, the

Philippines and Western Europe. To account for the differences in read depths across different

datasets, the samples were subsampled to the same number of reads across cohorts for absolute

ARG incidence measures. The percentages of saliva samples that contain at least one ARG for each

class and mechanism from these cohorts were evaluated. To account for varying read depth across

samples, the samples were subsampled to the same number of sequences. Saliva samples from

China, Fiji, the Philippines and Western Europe contain 20, 14, 23 and 17 ARG classes, respectively

(Fig. 1a). Further ARG classes are found in Philippines saliva samples, but most of this variability

originates from one individual: a farmer from Zambal who has carbapenem, fosfomycin, rifamycin,

and triclosan ARG classes[24]. All or almost all saliva samples from every cohort contain cephamycin,

fluoroquinolone, lincosamide, macrolide, streptogramin and tetracycline ARGs, and a high

percentage (above 50 %) of saliva samples from all cohorts contain pleuromutilin ARGs. Unlike

most cohorts, all saliva samples from China contain aminoglycoside ARGs represented by one

ARG, *APH(3')-Ia,* and also a high proportion of these samples contain glycylcycline represented by

one ARG, *tet(A)* (Supplementary Fig. 1a). The peptide ARG class is only found in saliva from

Chinese and Philippines individuals. Mechanisms of antimicrobial resistance including antibiotic

efflux, inactivation, target alteration and target protection are present in all saliva samples across all

cohorts (Fig. 1b), whilst the antibiotic target replacement mechanism is found in China, Philippines

and Western Europe but not in Fiji. The mechanism of reduced permeability to antibiotic is only

found in saliva from the same farmer in Zambal.

Dental plaque metagenomic data were only available from China and the US. The percentages of

the China and US plaque samples containing at least one ARG class and mechanism were

compared, and found to consist of 16 and 18 ARG classes, respectively (Fig. 1c). A greater

percentage of Chinese compared to US plaque samples contain pleuromutilin and

sulfonamide/sulfone ARGs. Similarly to saliva, all or almost all plaque samples from both cohorts

contain lincosamide, macrolide, streptogramin and tetracycline ARGs, with a high percentage

(above 50 %) of these containing cephamycin, fluoroquinolone, glycylcycline and pleuromutilin

ARGs. Notably, fluoroquinolone and tetracycline ARG classes in dental plaque are comprised of

fewer ARGs compared to saliva samples (Supplementary Fig. 1b). Antibiotic efflux, inactivation,

target alteration, target protection and antibiotic target replacement mechanisms are present in all

samples across both cohorts (Fig. 1d).

Stool samples were available from all locations, apart from the Philippines. Stool samples from

China, Fiji, the US and Western Europe were found to contain 31, 30, 17 and 30 ARG classes,

respectively (Fig. 1e). All or almost all stool samples contain cephalosporin, cephamycin,

5   5

diaminopyrimidine, lincosamide, macrolide, streptogramin and tetracycline ARGs, although most of

these ARG classes are found in lower percentages of Fiji stool samples. Compared to oral samples,

stool samples contain a lower proportion of the fluoroquinolone ARG class but higher proportions

of cephalosporin and diaminopyrimidine ARG classes exclusively consisting of *CblA-1* and *dfrF*

ARGs, respectively (Supplementary Fig. 1c). In addition to the US stool samples containing the

fewest number of ARG classes, they also contain a low proportion (less than 50 %) of

fluoroquinolones, penams, penems, peptide and phenicols ARG classes compared to China, Fiji and

Western Europe. Antibiotic efflux, inactivation, target alteration, target protection and antibiotic

target replacement mechanisms are present in all samples across all cohorts, but the reduced

permeability to antibiotic mechanism is not found in the US (Fig. 1f).

To determine whether there are differences in ARG composition between oral and gut samples as

well as between countries, the ARG incidence profiles, i.e. resistotypes, for every sample were

summarised using Principal Coordinates Analysis. Resistotypes were extracted using hierarchical

clustering and silhouette analysis[30]. Four resistotypes in total were identified (Fig. 2a, b). Oral

samples are mainly found in two major resistotypes, R2 and R4. R2 is comprised of mainly buccal

mucosa and saliva, and R4 contains mainly dental plaque and dorsum of tongue. All stool samples

are found in only two major resistotypes, R1 and R3. R1 consists of mostly stool and saliva from

Fiji, whereas R3 contains mainly stool from China, Fiji, US and Western Europe.

To evaluate whether the resistome is related to antibiotic prescription rates, the abundance of ARGs

for every ARG class was compared to the "Defined Daily Doses Per 1000 individuals" of

equivalent drug classes for each region. Prescription data were derived from ResistanceMap

(https://resistancemap.cddep.org/). This comparison indicates that overall ARG class abundance

does not follow a significant linear relationship with antibiotic prescriptions for any country and

body site (Fig. 2c, Supplementary Table 1).

The availability of longitudinal oral and stool samples from US individuals who had not taken

antimicrobial agents over two years afforded us the ability to investigate the stability of resistomes

without antibiotics. Hierarchical clustering reveals that the same individuals and body sites cluster

together, verifying that resistomes at all sites remain stable over a prolonged period with no

antimicrobial selection pressure (Supplementary Fig. 2).

Comparisons between oral and gut resistomes reveal differences in ARG composition,

abundance and diversity

To further investigate the differences between oral and gut resistome profiles, the abundance and

diversity between oral and gut samples from China, Fiji, US and Western Europe were evaluated

and compared. The total abundance, measured as the total Reads Per Kilobase Million (RPKM), of

all ARGs in gut samples is lower than in oral (buccal mucosa, dental plaque, dorsum of tongue and

saliva) samples across all pairwise comparisons in all cohorts (Fig. 3a). The overall abundance is

similar between paired US oral sites, with buccal mucosa and dental plaque having a slightly higher

abundance than the tongue dorsum samples (Supplementary Fig. 3). Oral samples contain a higher

relative abundance of ARGs coding for fluoroquinolone efflux pumps,

lincosamide/streptogramin/pleuromutilin resistance, macrolide efflux pumps and

macrolide/lincosamide resistance than stool samples (Fig. 3b, Supplementary Fig. 4). These classes

are mostly dominated by one of two ARGs across all cohorts, such as *patB* (coding for part of the

PatA-PatB efflux pump) in the fluoroquinolone efflux pump class (Supplementary Fig. 5a, b, c, d).

Stool contains a higher proportion of tetracyclines and "Other" ARGs across all cohorts (Fig. 3b,

Supplementary Fig. 4). These "Other" ARGs are mostly found in aminocoumarins,

aminoglycosides, cephalosporins, diaminopyrimidines, penams, penems and peptides across all

cohorts (Supplementary 5a, b, c, d).

The abundance of individual ARGs were compared between sample types using differential analysis

with DESeq2[31]. A meta-analysis strategy was implemented to combine results from all regions.

Stool samples are enriched with more ARGs compared to oral samples, but oral samples have

enriched ARGs of highest and lowest abundances compared to stool samples across all regions (Fig.

3c, Supplementary Fig. 6). *macB, OXA-85* and *tetA(60)* ARGs in dental plaque have the highest log

fold changes (20.8, 21.0 and 21.1 respectively), whilst *patB* and *RlmA(II)* in plaque, and *RlmA(II)* in

saliva have the lowest log fold changes (2.9, 2.6 and 2.3 respectively) compared to stool samples

(Fig. 3c). Highest log fold changes are seen in *cmlA6, Lactobacillus reuteri cat-TC, macB, TEM-1*

and *tet32* from saliva (20.6, 20.5,19.8, 20.5 and 20.5 respectively), and *cmlA6, macB, OXA-85,*

pmrA, tetA(60) and *tet(G)* from dental plaque (20.8, 20.6, 20.7, 20.7, 20.8 and 20.8 respectively)

compared to stool samples in China that are not enriched across all cohorts (Supplementary Fig. 7).

As well as differences between oral and gut, differentially abundant ARGs were found between

different sites in the oral cavity (Supplementary Fig. 6). For example, between the US dorsum of

tongue and plaque samples, and between the US dorsum of tongue and buccal mucosa, all ARGs are

enriched in the dorsum of the tongue. Similarly, between dental plaque and buccal mucosa, all

ARGs are enriched in dental plaque. Most of these ARGs confer resistance to cephamycin,

fluoroquinolone, MLS (macrolide/lincosamide/streptogramin) and tetracycline antibiotics. From

China, there are more significantly abundant ARGs in saliva than plaque with resistance to

aminoglycoside, cephalosporin, fluoroquinolone (*pmrA* and *patB*), lincosamides, macrolides (*macB*

and *mefA* ARGs), MLS (in particular to the *Erm 235* ribosomal RNA methyltransferase family) and

tetracycline antibiotics. Overall, stool samples are enriched with more alternative ARGs and ARG classes compared to oral samples, but with the highest and lowest enrichments of individual ARGs originating from oral samples.

To investigate ARG diversity further, the ARG richness was evaluated between pairwise comparisons of sample types for each cohort with ARG richness defined as the number of unique ARGs per sample. Although there are no significant difference between saliva and stool samples from Fiji and Western Europe, both Chinese and US samples have significant differences in ARG richness. China and US stool samples have a significantly higher ARG richness than paired China plaque and saliva, and paired US plaque and buccal mucosa (Mann-Whitney, paired, two-sided t-test, p-value < 0.05) (Fig. 4). In contrast, the US dorsum of tongue contains a significantly higher ARG richness than US stool. Between oral sites, Chinese saliva has a greater ARG richness than paired dental plaque (Mann-Whitney, paired, two-sided t-test, p-value < 0.05). In addition, US dorsum of tongue has a higher ARG richness than both plaque and buccal mucosa, whilst plaque has a greater ARG richness than buccal mucosa (Mann-Whitney, paired, two-sided t-test, p-value < 0.05). It is important to note that while ARG richness only measures the gene incidence regardless of expression, multiple ARGs have the potential to be involved in the expression of a single efflux pump complex, meaning ARG richness may overestimate this potential expression. Therefore, to determine the impact of this overestimation, the analysis was repeated to exclude ARGs that regulate or are part of an efflux pump complex. The differences in ARG richness have the same significance across all paired samples and countries (Supplementary Fig. 8).

Oral and gut ARG profiles associate with species

Spearman's correlation analysis between ARG and species abundances were conducted to predict the origin of ARGs. Only significant correlations are found in saliva and stool from China, and saliva from the Philippines. The *CfxA* beta-lactamase family, *RlmA(II), tetQ, tetA(46), pgpB, patB*

and *pmrA* ARGs are all strongly correlated with specific species found in both countries (Fig. 5a, b).

The strongest co-occurrence can be found in saliva samples from China between *APH(3')-Ia* and a

Komagataella pastoris strain, *Lactococcus lactis*, *Enterococcus faecalis* and *Leuconostoc*

mesenteroides, whilst *pgpB* correlates with *Porphyromonas gingivalis*. The highly abundant ARG

RlmA(II) in Chinese saliva is associated with *Gemella haemolysans*, *Veillonella parvula* and

Streptococcus mitis/oralis/pneumoniae. In contrast to saliva, there are fewer species associated with

a greater number of ARGs in stool samples from China. *E. coli* in Chinese stool samples is co-

associated with many ARGs that encode multidrug efflux pumps and ARGs from *E. coli including*

ampC beta-lactamase, *acrA* and *mdfA* (Supplementary Fig. 9). Thus, this analysis has the potential

to be a predictive tool of ARG origin in metagenomes. However, it can only be applied where an

ARG or taxon is found in a high proportion (in this case, at least half) of the samples to ensure

Spearman's correlation does not falsely rank many zero-values.

240   **Discussion**

Antimicrobial resistance is one of the most serious health problems of recent times. The advent of

high-throughput sequencing technologies has enabled us to analyse resistomes throughout a

microbiome. In this study, we provide key insights into the resistomes of different intraoral sites

from healthy individuals across diverse geographical locations and compare their composition to

paired gut resistomes. At a population level, there are both country and body site-specific

differences in the prevalence of ARGs, ARG classes and resistance mechanisms. . It is possible that

differences in extraction protocols and batch effects may have a greater bias towards some ARGs

over others. Therefore, we do not make direct statistical comparisons between cohorts. For China,

the Philippines, US and Western Europe, the abundance of ARG classes does not correlate with

antibiotic prescription rates. A possible reason is for this is the prescription data does not include

over-the-counter antimicrobial use, which is especially prevalent in China and the Philippines, and

thus may underestimate antimicrobial use[32,33]. In addition, antibiotics are widely used in husbandry

and the fishing industry with poorly understood impacts on AMR incidence and dynamics in

humans[34–36]. Prescription levels for a particular antibiotic are unlikely to be of significant value in

the surveillance of AMR in the regional community. Instead, determining the population resistome

is significantly more informative[37].

The abundance and diversity of ARGs at different body sites is also of interest. Although, there are

significantly more distinct ARGs in stool compared to oral samples, those ARGs present at the

highest relative abundances exist in oral samples. There are several potential reasons why this may

be. It is notable that many sites in the oral cavity (e.g. plaque and tongue dorsum) host highly

complex and robust microbial biofilm structures. It has been posited that the compact structure of

microbes within oral biofilms is a conducive environment for the aquistion of ARGs and their HGT

within biofilms[38]. Likewise, the generally protective nature of biofilms against antimicrobial drugs

may favour ARG acquisition. It is notable that the dorsum of tongue contains a higher diversity of

these genes than other oral sites. This may be explained by the unique papillary structure on the

dorsum of the tongue which acts as complex microbiological niche favouring the deposition of oral

debris and microbes[39], thus giving rise to a richer microbial community with potentially greater

numbers of transient microbes. Another reason could be the difference in species resident in the gut

compared to the oral cavity. *E. coli* strains in Chinese stool samples are predicted to contain a

variety of ARGs, especially of the multidrug class, whereas species found in Chinese saliva were

estimated to contain fewer ARGs. Due to stringent constraints of the correlation analysis, however,

it was not possible to predict the origins for ARGs for all cohorts.

Antibiotic use leading to acquisition of ARGs is another potential factor. Pharmacokinetics of orally

administered antibiotics suggests that the oral cavity and oesophagus would be only briefly exposed

to the antibiotic during swallowing, whilst the gut is exposed over a more prolonged period.As

antibiotics transit through the intestinal tract, they are gradually absorbed via the intestinal

epithelium into the bloodstream. Therefore, microbes in the gut exposed to antibiotics for a longer

period of time due to their increased bioavailability will receive higher antibiotic selection pressures

than those in the oral cavity[40]. The incidences where the oral cavity is likely to acquire ARGs from

selective pressures of local antibiotics are from topical antibiotics for periodontal infections or

orally-administered antibiotics being absorbed into the bloodstream.

The differences in ARG profiles across body sites has significant implications for the

characterisation and interpretation of resistome studies. Previous shotgun metagenomic studies have

focused almost exclusively on the resistome from the human gut[2–4,6]. While the gut may be a diverse

reservoir of ARGs, whether these genes are particularly prevalent or have the potential for

expression sufficient to drive resistant infections at other body sites is not clear[41]. It is therefore

imperative that to test potential applications of non-culture based metagenomic AMR surveillance,

we need to characterise the resistome at different body sites with different pharmacokinetic

exposures to antimicrobials. This information can then be integrated with culture-based

susceptibility tests, culturomics[42] and functional metagenomic screens[43] to determine the expression

potential of these ARGs. In doing so, we will obtain a more complete picture of the state of AMR

within a population.

Methods

Metagenomic sequence data

12   12

A total of 1174 publicly available metagenomic samples covering the US, China, Fiji, the

Philippines and Western Europe (France and Germany), all sequenced using Illumina HiSeq 2000,

were analysed. Longitudinal US samples were excluded from the majority of the study after the first

time point to ensure each sample was independent, unless specified otherwise. All metagenomes

passed over half the quality control metrics in FastQC

(https://www.bioinformatics.babraham.ac.uk/projects/fastqc/) with these pass rates calculated in

MultiQC[44]. These samples include 1) longitudinal data across two years with various timepoints

from the Human Microbiome Project 1 (referred to as US)[28] containing buccal mucosa (n = 164: 33

with one, 34 with two, 19 with three and 1 with six timepoints); dorsum of tongue (n = 188: 22 with

one, 43 with two, 24 with three and 2 with four timepoints); dental plaque (n = 191: 24 with one, 41

with two, 21 with three, 1 with four and 3 with six timepoints); stool (n = 156: 13 with one, 33 with

two, 21 with three, 2 with four and 1 with six timepoints), 2) healthy control samples from a

Chinese rheumatoid arthritis study[22] containing dental plaque (n = 32); saliva (n = 33); stool (n =

72), 3) saliva (n = 136) and stool (n = 137) samples from Fiji[23], 4) saliva samples (n = 23) from

healthy hunter-gatherers and traditional farmers from the Philippines[24], and 5) saliva (n = 21) and

stool (n = 21) samples from Western Europe (5 saliva and 5 stool samples from Germany[25,27], and 16

saliva and 16 stool samples from France[26,27]).

Raw paired-end metagenomic reads from Chinese and Philippines samples were downloaded from

the EBI (https://www.ebi.ac.uk/metagenomics/). Paired-end metagenomic samples from US were

downloaded from https://portal.hmpdacc.org/. Raw paired-end metagenomic reads from Fiji

(project accession PRJNA217052), France and Germany (project accession PRJEB28422) were

downloaded from the NCBI. All US, China, Fiji and Philippines samples, and stool samples from

France and Germany, were collected and sequenced as described in the following cited studies[22–

[26,28]. Saliva samples from France and Germany were collected and sequenced as described in the

following cited study[27]. Metadata for the samples can be found in Supplementary Table 2.

Processing metagenomic data

The raw reads for all samples were trimmed using AlienTrimmer 0.4.0[45] with parameters *-k 10 -l 45*

-m 5 -p 40 -q 20 and Illumina contaminant oligonucleotides

(https://gitlab.pasteur.fr/aghozlan/shaman_bioblend/blob/18a17dbb44cece4a8320cce8184adb99665

83aaa/alienTrimmerPF8contaminants.fasta). Human contaminant sequences were removed from all

samples by discarding reads that mapped against a human reference genome (downloaded from

Human Genome Resources at NCBI on 27th February 2017) using Bowtie2 2.2.3[46] with parameters

-q -N 1 -k 1 --fr --end-to-end --phred33 --very-sensitive --no-discordant. The quality of the raw

reads and the filtered reads of each sample was evaluated using the Fastqc 0.11.3

(https://github.com/s-andrews/FastQC).

Extracting ARGs

All processed metagenomes were mapped against *nucleotide_fasta_protein_homolog_model* from

the antimicrobial resistance database CARD 3.0.0[29] using KMA 1.2.6. Hits were extracted where

the template coverage was greater than 90 %. The metagenomes were mapped against these hits

using Bowtie2 2.2.5 with parameter *–very-sensitive-local*. Mapped reads were filtered from

unmapped reads, sorted and indexed using Samtools 1.9[47]. Statistics for the number of reads

mapped for each ARG were extracted using Bedtools 2.28.0[48].

Abundance of ARGs

The reads per kilobase of read per million (RPKM) was calculated for every sample as the number

of reads divided by the total number of library reads per million, then divided by the gene length in

kilobases. The relative abundance of ARGs for each country and sample type was calculated by

dividing the RPKM by the sum of RPKM for each country and sample type. The relative abundance

of ARGs for each sample and sample type was calculated by dividing the RPKM by the sum of the

RPKM for each sample. Differential abundance of ARGs between paired sample types from each

country were calculated using the DESeq2 1.20.0 package[31] as recommended by Jonsson et al.[49].

ARGs that were significantly differentially abundant (adjusted p-value < 0.05) across study cohorts

for paired sample comparisons were extracted using a meta-analysis random effects model with the

metafor 2.1-0 package[50].

Comparison of ARG class abundance and antibiotic prescription rate

The mean RPKM and standard error for every ARG class was compared against the antibiotic

prescription rate measured as the Defined Daily Dose (DDD) Per 1000 individuals in 2015 from

China, the Philippines, Western Europe (France and Germany) and the US. This data was derived

from ResistanceMap (https://resistancemap.cddep.org/) accessed on 19[th] February 2019. No

antibiotic use data was available for Fiji. Linear regression was conducted on the log transformed

mean RPKM versus the DDD Per 1000 for all ARG classes.

Percentage of samples containing ARGs, ARG classes and ARG mechanisms

To show whether the percentages of samples containing an ARG class were consistent across the

same number of reads, metagenomes were first subsampled using seqtk 1.2

(https://github.com/lh3/seqtk) with parameter seed *-s100.* 6.9 million reads were subsampled from

18 saliva samples with the lowest number of reads greater than 6.9 million reads, from each cohort:

China, Fiji, the Philippines and Western Europe. 18 million reads were subsampled from 18 dental

plaque with the lowest number of reads greater than 18 million reads from both China and US

cohorts. 16.9 million reads were subsampled from 18 stool samples with the lowest number of reads

greater than 16.9 million reads from China, Fiji, the US and Western Europe cohorts. These were

mapped to CARD 3.0.0 as described in *Extracting ARGs*. R 3.5.1 was used for all downstream

analysis. Each ARG was annotated with "Drug Class" and "Resistance Mechanism" using CARD

3.0.0 metadata. Percentages of samples containing an ARG, ARG class and mechanism were

calculated from these samples. 95 % confidence intervals (CIs) were evaluated from percentages

extracted from bootstrapping samples 100 times for each cohort and sample type.

Principal Coordinates Analysis

807 metagenomic files that contained at least 1 millions reads and were not longitudinal US

samples were first subsampled to 1 million reads using seqtk 1.2 (https://github.com/lh3/seqtk) with

parameter seed *-s100*. These were mapped to CARD 3.0.0 as described in *Extracting ARGs*. 17

subsampled samples did not contain ARGs leaving the number of samples to 790. Principal

Coordinates Analysis was applied to the binary distance between ARG presence or absence profiles

for each sample (excluding longitudinal US samples) using the vegan 2.5-2 package (https://cran.r-

project.org/web/packages/vegan/index.html). Resistotypes were extracted using hierarchical

clustering of the Euclidean distance between principal coordinates with eigenvalues above zero.

Silhouette analysis was used to determine the optimum number of resistotypes using the cluster

2.0.7.1 package (https://cran.r-project.org/web/packages/cluster/index.html). The number of

resistotypes is defined by the number of clusters with the largest silhouette width.

ARG diversity

To ensure the ARG richness could be compared statistically across different sample types from the

same individuals[51], the metagenomes (excluding longitudinal US samples) were subsampled using

seqtk 1.2 with seed *-s100* . Paired samples from the same individuals in each of the following

groups containing two sample types were subsampled to a number rounded down by two significant

figures from the lowest number of reads in the group.

China dental plaque vs. saliva: 3.5 million reads were sampled from China dental plaque (n = 31)

and paired saliva (n = 31) samples.*China stool vs. saliva:* 3.5 million reads were sampled from

China stool (n = 31) and paired saliva (n = 31) samples.*China stool vs. dental plaque:* 14 million

reads were sampled from China stool (n = 30) and paired dental plaque (n = 30) samples.*US buccal*

mucosa vs. dental plaque: 1 million reads were sampled from US buccal mucosa (n = 78) and

paired dental plaque (n = 78) samples.*US buccal mucosa vs. dorsum of tongue:* 1 million reads were

sampled from US buccal mucosa (n = 86) and paired dorsum of tongue (n = 86) samples.*US buccal*

mucosa vs. stool: 1 million reads were sampled from and US buccal mucosa (n = 64) and paired

stool (n = 64) samples. *US dental plaque vs. dorsum of tongue:* 4.2 million reads were sampled

from US dental plaque (n = 89) and paired dorsum of tongue (n = 89) samples.*US dental plaque vs.*

stool: 4.2 million reads were sampled from US dental plaque (n = 68) and paired stool (n = 68)

samples.*US dorsum of tongue vs. stool:* 14 million reads were sampled from US dorsum of tongue

(n = 67) and paired stool (n = 67) samples. *Fiji saliva vs. stool:* 1.2 million reads were sampled

from Fiji saliva (n = 132) and paired stool (n = 132) samples. Fiji samples containing fewer than 1.2

million reads were excluded from the analysis.*Western Europe saliva vs. stool:* 3.1 million reads

were sampled from Western Europe saliva (n = 21: 5 from Germany and 16 from France) and paired

stool (n = 20: 5 from Germany and 16 from France) samples. These were mapped to the CARD

database as described in Methods *Extracting ARGs*.

Once the metagenomes were subsampled, ARGs extracted and filtering by coverage, the ARG

diversity per sample was measured as the ARG richness, recommended previously by Bengtsson-

Palme et al.[60]. For every sample, the ARG richness was calculated as the number of unique ARGs.

To account for multiple ARGs coding for an efflux pump complex, the ARG richness was

calculated excluding ARGs that regulate or are part of an efflux pump complex. The ARG richness

between samples in each group was tested for statistical significance with a Mann-Whitney, paired,

two-sided t-test.

Correlation analysis

MetaPhlAn2 2.6.0[53] was used to extract taxonomic composition from all samples. Spearman's

correlation was applied to relative abundances of reads mapped to ARG and MetaPhlAn2 species

profiles for paired samples. ARGs and species that were not found in more than half of samples for

each country were removed, to alleviate the bias from potential joint ranking of zero values by

Spearman's rank. The rho and p-values were calculated using the *stats* package in R and the p-

values were adjusted with Benjamini-Hochberg where FDR < 5 %. Correlations were found from

China saliva and paired stool samples, and Philippines saliva samples. No significant correlations

could be found from Fiji, Western Europe or US samples.

Data Availability

ARG data, figures, tables and script to run analysis are available at https://github.com/blue-

moon22/resistomeData

High-resolution figures are available at https://figshare.com/s/5d020d6f52a7685f341f

Code Availability

R package for resistome analysis is available at https://github.com/blue-moon22/resistomeAnalysis

449   **References**

1. *Antimicrobial resistance: global report on surveillance*. (World Health Organization, 2014).

2. Rose, G. *et al.* Antibiotic resistance potential of the healthy preterm infant gut microbiome. *PeerJ* **5**, e2928 (2017).

3. Feng, J. *et al.* Antibiotic Resistome in a large-scale healthy human gut microbiota deciphered by metagenomic and network analyses. *Environmental Microbiology* n/a-n/a doi:10.1111/1462-2920.14009

4. Forslund, K. *et al.* Country-specific antibiotic use practices impact the human gut resistome. *Genome Res.* **23**, 1163–1169 (2013).

5. Rahman, S. F., Olm, M. R., Morowitz, M. J. & Banfield, J. F. Machine Learning Leveraging Genomes from Metagenomes Identifies Influential Antibiotic Resistance Genes in the Infant Gut Microbiome. *mSystems* **3**, e00123-17 (2018).

6. Hu, Y. *et al.* Metagenome-wide analysis of antibiotic resistance genes in a large cohort of human gut microbiota. *Nature Communications* **4**, 2151 (2013).

7. Pärnänen, K. *et al.* Maternal gut and breast milk microbiota affect infant gut antibiotic resistome and mobile genetic elements. *Nat Commun* **9**, (2018).

8. Thomas, M. *et al.* Metagenomic characterization of the effect of feed additives on the gut microbiome and antibiotic resistome of feedlot cattle. *Scientific Reports* **7**, 12257 (2017).

9. Munk, P. *et al.* Abundance and diversity of the faecal resistome in slaughter pigs and broilers in nine European countries. *Nature Microbiology* **3**, 898 (2018).

10. Pehrsson, E. C., Forsberg, K. J., Gibson, M. K., Ahmadi, S. & Dantas, G. Novel resistance functions uncovered using functional metagenomic investigations of resistance reservoirs. *Front. Microbiol.* **4**, (2013).

11. Ma, L. *et al.* Catalogue of antibiotic resistome and host-tracking in drinking water deciphered by a large scale survey. *Microbiome* **5**, 154 (2017).

12. Noyes, N. R. *et al.* Characterization of the resistome in manure, soil and wastewater from dairy and beef production systems. *Scientific Reports* **6**, 24645 (2016).

13. Bengtsson-Palme, J., Boulund, F., Fick, J., Kristiansson, E. & Larsson, D. G. J. Shotgun metagenomics reveals a wide array of antibiotic resistance genes and mobile elements in a polluted lake in India. *Front. Microbiol.* **5**, (2014).

14. Knoll, B., Tleyjeh, I. M., Steckelberg, J. M., Wilson, W. R. & Baddour, L. M. Infective Endocarditis Due to Penicillin-Resistant Viridans Group Streptococci. *Clin Infect Dis* **44**, 1585–1592 (2007).

15. Diaz-Torres, M. L. *et al.* Determining the antibiotic resistance potential of the indigenous oral microbiota of humans using a metagenomic approach. *FEMS Microbiol Lett* **258**, 257–262 (2006).

16. Seville, L. A. *et al.* Distribution of tetracycline and erythromycin resistance genes among human oral and fecal metagenomic DNA. *Microb. Drug Resist.* **15**, 159–166 (2009).

17. Clemente, J. C. *et al.* The microbiome of uncontacted Amerindians. *Science Advances* **1**, e1500183 (2015).

18. Warinner, C. *et al.* Pathogens and host immunity in the ancient human oral cavity. *Nature Genetics* **46**, 336–344 (2014).

19. Roberts, A. P. & Kreth, J. The impact of horizontal gene transfer on the adaptive ability of the human oral microbiome. *Front Cell Infect Microbiol* **4**, (2014).

20. Witherden, E. A., Bajanca-Lavado, M. P., Tristram, S. G. & Nunes, A. Role of inter-species recombination of the ftsI gene in the dissemination of altered penicillin-binding-protein-3-mediated resistance in Haemophilus influenzae and Haemophilus haemolyticus. *J. Antimicrob. Chemother.* **69**, 1501–1509 (2014).

21. Chaffanel, F., Charron-Bourgoin, F., Libante, V., Leblond-Bourget, N. & Payot, S. Resistance Genes and Genetic Elements Associated with Antibiotic Resistance in Clinical and Commensal Isolates of Streptococcus salivarius. *Appl. Environ. Microbiol.* **81**, 4155–4163 (2015).

22. Zhang, X. *et al.* The oral and gut microbiomes are perturbed in rheumatoid arthritis and partly normalized after treatment. *Nature Medicine* **21**, 895–905 (2015).

23. Brito, I. L. *et al.* Mobile genes in the human microbiome are structured from global to individual scales. *Nature* **535**, 435–439 (2016).

24. Lassalle, F. *et al.* Oral microbiomes from hunter-gatherers and traditional farmers reveal shifts in commensal balance and pathogen load linked to diet. *Mol. Ecol.* **27**, 182–195 (2018).

25. Voigt, A. Y. *et al.* Temporal and technical variability of human gut metagenomes. *Genome Biology* **16**, 73 (2015).

26. Zeller, G. *et al.* Potential of fecal microbiota for early-stage detection of colorectal cancer. *Molecular Systems Biology* **10**, 766 (2014).

27. Schmidt, T. S. B. *et al.* Extensive transmission of microbes along the gastrointestinal tract. *eLife* **8**, e42693 (2019).

28. Human Microbiome Project Consortium. A framework for human microbiome research. *Nature* **486**, 215–221 (2012).

29. McArthur, A. G. *et al.* The comprehensive antibiotic resistance database. *Antimicrob. Agents Chemother.* **57**, 3348–3357 (2013).

30. Rousseeuw, P. J. Silhouettes: A graphical aid to the interpretation and validation of cluster analysis. *Journal of Computational and Applied Mathematics* **20**, 53–65 (1987).

31. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biology* **15**, 550 (2014).

32. Chang, J. *et al.* Sale of antibiotics without a prescription at community pharmacies in urban China: a multicentre cross-sectional survey. *J. Antimicrob. Chemother.* **72**, 1235–1242 (2017).

33. Auta, A. *et al.* Global access to antibiotics without prescription in community pharmacies: A systematic review and meta-analysis. *Journal of Infection* (2018). doi:10.1016/j.jinf.2018.07.001

34. Liu, X., Steele, J. C. & Meng, X.-Z. Usage, residue, and human health risk of antibiotics in Chinese aquaculture: A review. *Environ. Pollut.* **223**, 161–169 (2017).

35. Cheng, G. *et al.* Antibiotic alternatives: the substitution of antibiotics in animal husbandry? *Front. Microbiol.* **5**, (2014).

36. Stanton, T. B. A call for antibiotic alternatives research. *Trends in Microbiology* **21**, 111–113 (2013).

37. Hendriksen, R. S. *et al.* Global monitoring of antimicrobial resistance based on metagenomics analyses of urban sewage. *Nature Communications* **10**, 1124 (2019).

38. Roberts, A. P. & Mullany, P. Oral biofilms: a reservoir of transferable, bacterial, antimicrobial resistance. *Expert Review of Anti-infective Therapy* **8**, 1441–1450 (2010).

39. Roldán, S., Herrera, D. & Sanz, M. Biofilms and the tongue: therapeutical approaches for the control of halitosis. *Clin Oral Invest* **7**, 189–197 (2003).

40. Palleja, A. *et al.* Recovery of gut microbiota of healthy adults following antibiotic exposure. *Nature Microbiology* **3**, 1255 (2018).

41. Piddock, L. J. V. Assess drug-resistance phenotypes, not just genotypes. *Nature Microbiology* **1**, 16120 (2016).

42. Greub, G. Culturomics: a new approach to study the human microbiome. *Clinical Microbiology and Infection* **18**, 1157–1159 (2012).

43. Mullany, P. Functional metagenomics for the investigation of antibiotic resistance. *Virulence* **5**, 443–447 (2014).

44. Ewels, P., Magnusson, M., Lundin, S. & Käller, M. MultiQC: summarize analysis results for multiple tools and samples in a single report. *Bioinformatics* **32**, 3047–3048 (2016).

45. Criscuolo, A. & Brisse, S. AlienTrimmer: A tool to quickly and accurately trim off multiple short contaminant sequences from high-throughput sequencing reads. *Genomics* **102**, 500–506 (2013).

46. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat. Methods* **9**, 357–359 (2012).

47. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–2079 (2009).

48. Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics* **26**, 841–842 (2010).

49. Jonsson, V., Österlund, T., Nerman, O. & Kristiansson, E. Statistical evaluation of methods for identification of differentially abundant genes in comparative metagenomics. *BMC Genomics* **17**, 78 (2016).

50. Viechtbauer, W. Conducting Meta-Analyses in R with the metafor Package. *Journal of Statistical Software* **36**, 1–48 (2010).

51. Bengtsson-Palme, J., Larsson, D. G. J. & Kristiansson, E. Using metagenomics to investigate human and environmental resistomes. *J Antimicrob Chemother* **72**, 2690–2703 (2017).

52. Bengtsson-Palme, J. The diversity of uncharacterized antibiotic resistance genes can be predicted from known gene variants—but not always. *Microbiome* **6**, 125 (2018).

53. Truong, D. T. *et al.* MetaPhlAn2 for enhanced metagenomic taxonomic profiling. *Nat Meth* **12**, 902–903 (2015).

Acknowledgements

The research was supported by the Centre for Host-Microbiome Interactions, King's College London, the Biotechnology and Biological Sciences Research Council (BBSRC) (Grant No. BB/M009513/1) awarded to D.L.M., and Engineering and Physical Sciences Research Council (EPSRC), BBSRC and UK Research and Innovation (UKRI) (Project No. EP/S001301/1) awarded to S.S.. The authors would like to thank Professor William Wade for constructive criticism of the manuscript. S.S. acknowledges Uppmax for providing assistance for computational infrastructure.

Author contributions

V.C. and D.L.M. conceived the presented idea. V.C. and D.G-C. conducted the bioinformatics and data analysis. V.C. and D.L.M. wrote the manuscript with support from D.G-C., E.W., G.B.P., P.M., S.L. and S.S.. D.L.M. and D.G-C. oversaw the project.

Figures

Fig. 1 Percentage of saliva samples from individuals that contain **a** an ARG class and **b** an ARG mechanism, from China (n = 18), Fiji (n = 18), the Philippines (n = 18) and Western Europe (n = 18). Percentage of dental plaque samples from individuals that contain **c** an ARG class and **d** an ARG mechanism, from China (n = 18) and the US (n = 18). Percentage of stool samples from individuals that contain, **e** an ARG class and **f** an ARG mechanism, from China (n = 18), Fiji (n = 18), the US (n = 18) and Western Europe (n = 18). Error bars are 95 % confidence intervals (CIs) that were evaluated from percentages extracted from bootstrapping samples 100 times.

Fig. 2 a Principal Coordinates Analysis of the incidence (presence/absence) of ARGs for all samples where each sample is represented by a point. Samples are labelled as Resistotype clusters, evaluated from hierarchical clustering of binary distance between ARG incidence profiles. Number of clusters was selected with the highest silhouette width using silhouette analysis. US buccal mucosa (n = 86), China dental plaque (n = 29), US dental plaque (n = 80), US dorsum of tongue (n = 91), China saliva (n = 33), Fiji saliva (n = 129), Philippines saliva (n = 22), Western Europe saliva (n = 21), China stool (n = 72), Fiji stool (n = 136), US stool (n = 70), and Western Europe stool (n = 21). **b** Percentage of Resistotypes that contain samples from a body site and geographical location. **c** Mean and standard error (error bars) of reads per kilobase of read per million (RPKM) of ARGs for each ARG class against the Defined Daily Doses Per 1000 individuals in 2015 from China, the Philippines, Western Europe (France and Germany) and the US. (Fiji antibiotic use data unavailable.) Mean RPKM calculated from China dental plaque (n = 32), China saliva (n = 33), China stool (n = 72), Philippines saliva (n = 23), US buccal mucosa (n = 87), US dental plaque (n = 90), US dorsum of tongue (n = 91), US stool (n = 70), Western Europe saliva (n = 21) and Western Europe stool (n = 21).

Fig. 3 Resistome abundance for China, Fiji, Western Europe and US. **a** Absolute abundance in log10 of reads per kilobase of read per million (RPKM) of ARGs for paired samples between China stool (n = 30) and dental plaque (n = 30), China stool (n = 31) and saliva (n = 31), Fiji saliva (n = 132) and stool (n = 132), US stool (n = 68) and dental plaque (n = 68), US stool (n = 69) and dorsum of tongue (n = 69), US stool (n = 64) and buccal mucosa (n = 64), and Western Europe saliva (n = 21) and stool (n = 21).. **b** Relative abundance of reads labelled by the top ten most abundant ARG classes across all geographical locations or "Other" classes for each sample type from China saliva (n = 33), dental plaque (n = 32) and stool (n = 72), Fiji saliva (n = 136) and stool (n = 137), Western Europe saliva (n = 21) and stool (n = 21), and US buccal mucosa (n = 87), dorsum of tongue (n = 91), dental plaque (n = 90) and stool (n = 70). **c** Estimated average log2 fold change of ARGs between paired dental plaque and stool, and saliva and stool samples using random effects meta-analysis across study cohorts (p-value < 0.05). Error bars are 95 % confident intervals from meta-analysis. ARGs selected for meta-analysis where adjusted p-value < 0.05 from differential abundance analysis of US and China dental plaque and stool, and China, Fiji and Western Europe saliva and stool samples.

Fig. 4 ARG richness for paired samples between China dental plaque (n = 31) and saliva (n = 31), China stool (n = 30)
and dental plaque (n = 30), China stool (n = 31) and saliva (n = 31), Fiji saliva (n = 132) and stool (n = 132), US buccal
mucosa (n = 78) and dental plaque (n = 78), US buccal mucosa (n = 86) and dorsum of tongue (n = 86), US dental
plaque (n = 89) and dorsum of tongue (n = 89), US buccal mucosa (n = 64) and stool (n = 64), US dental plaque (n =
68) and stool (n = 68), US dorsum of tongue (n = 67) and stool (n = 67), and Western Europe saliva (n = 21) and stool
(n = 21) with Mann-Whitney, paired, two-sided t-test (p-value < 0.05 as *, < 0.01 as **, < 0.005 as ***)
508
509

510 **Fig. 5** Spearman's correlation of ARG and species abundance from saliva samples represented as heatmaps of *rho*
511 between 0 and 1, only where adjusted p-value < 0.05. Each heatmap represents **a** China (n = 31) and **b** Philippines
512 saliva (n = 23). Rows and columns are clustered by hierarchical clustering of Euclidean distance. Columns are coloured
513 by phylum. P-values are adjusted by Benjamini-Hochberg multiple test correction.