Title:

Combining statistical and mechanistic models to unravel the drivers of mortality within a rear-edge beech population

Authors:

Cathleen Petit-Cailleux[1], Hendrik Davi[1], François Lefevre[1], Joseph Garrigue[2], Jean-André Magdalou[2], Christophe Hurson[2,3], Elodie Magnanou[2,4], and Sylvie Oddou-Muratorio[1].

Adresses

[1]INRA, UR 629 Ecologie des Forêts Méditerranéennes, URFM, Avignon, France

[2]Réserve Naturelle Nationale de la Forêt de la Massane, France

[3]Fédération des Réserves Naturelles Catalanes, Prades, France

[4]Sorbonne Université, CNRS, Biologie Intégrative des Organismes Marins, BIOM, F-66650 Banyuls-sur-Mer, France

*Corresponding author:

Sylvie Oddou-Muratorio

INRA, Unité Ecologie des Forêts Méditerranéennes

Domaine Saint Paul, Site Agroparc

84914 Avignon Cedex 9, France

Fax: +33 (0)4 32 72 29 02

sylvie.muratorio@inra.fr

Abstract

As several studies report an increasing decline of forests, a major issue in ecology is to better understand and predict tree mortality. The interactions between the different factors and physiological processes driving tree mortality, as well as the individual between-trees variability of mortality risk remain to be characterised.

This study is based on an exceptional yearly individual monitoring of 4327 European beeches (*Fagus sylvatica*) since 2002 in a rear-edge population within a natural reserve. We combined two types of approaches. Statistical models were used to quantify the effects of climate, competition, tree size and health on mortality. Carbon reserves, hydraulic conductance and late frosts were simulated using a process-based model to disentangle the mechanisms driving temporal and inter-individual variations in mortality.

The mortality rate at population level was associated to drought indices in statistical models, and driven by a combination of conductance loss, carbon reserve depletion and late frost damages in the process-based simulations. In statistical models, the individual probability of mortality decreased with mean growth, and increased with crown defoliation, budburst earliness, fungi presence and competition. Interaction effects between tree size and defoliation were significant, the probability of mortality being higher for a small-defoliated tree than for a tall one. Finally, the process-based model predicted a higher conductance loss and a higher frequency of late frosts for earlier trees together with a higher level of carbon reserve, while the ability to defoliate crown was found to limit the impact of hydraulic stress and allow carbon reserve accumulation.

We discuss the convergences and divergences obtained between statistical and process-based approaches and we highlight the importance of combining them to disentangle the processes underlying mortality, and to account for individual variability in vulnerability.

Keywords: *Fagus sylvatica*, logistic regression, process-based model, CASTANEA, Longitudinal analysis, defoliation.

Introduction

Global change has been repeatedly reported to generate forest decline and tree mortality, both in terms of background, non-catastrophic mortality(Van Mantgem et al. 2009, Lorenz and Becher 2012) and of massive, catastrophic mortality due to extreme, pulse events (Allen et al. 2010; Lorenz and Becher 2012; Mueller et al. 2005). To predict how such new regime of tree mortality will impact forest structure, composition and ecosystem services (Anderegg et al. 2015; Choat et al. 2018), we need to better decipher the respective roles of the various drivers and mechanisms underlying tree mortality.

Studying mortality poses several well-recognized challenges. It is triggered by several factors and involves several underlying and interacting physiological processes. The factors triggering mortality include extreme, pulse climatic events (i.e. repeated drought, storm, flood, heavy snow, late frost, wildfire) or sudden changes in biotic interactions (i.e. emerging pests, invasive species), but also long-term climatic or biotic perturbations (i.e. recurrent water deficits, changes in competition at the community level) (Maraun et al. 2003; McDowell et al. 2011). Moreover, these factors can have interactive effects: for instance, drought may increase tree vulnerability to pests (Durand-Gillmann et al. 2014; Anderegg et al. 2015) or stand vulnerability to fire (Brando et al. 2014). Finally, a single factor triggering mortality may involve several underlying physiological processes, with several thresholds leading to mortality and potentially feedbacks between them (McDowell et al. 2011). This is exemplified by drought, which is usually considered to trigger mortality through the combination of hydraulic failure and carbon starvation (Adams et al. 2017; Anderegg et al. 2012; McDowell et al. 2011). Hydraulic failure is the loss of conductance resulting from major xylem embolism, i.e. the formation of water vapour or air bubbles within the conductive vessels due to high water tension from soil to canopy (Tyree and Sperry 1989). To avoid hydraulic failure, trees can close their stomata before reaching the vessel species-specific embolism point (Cowan and Farquhar 1977). However, stomata closure mechanically reduces photosynthetic activity, forcing the tree to ensure basal metabolism using carbon reserves, which can eventually deplete them particularly during long drought, leading to mortality through carbon starvation. Many experimental studies on drought suggest that hydraulic failure is the most frequent cause of tree mortality, and at least, often the initial step towards cascading processes leading to mortality (Choat et al. 2018). Nevertheless, the relationship between carbon and water fluxes and the role of biotic factors during and after drought are far from being resolved (Adams et al. 2017; Feng et al. 2018; McDowell et al. 2011; Meir, Mencuccini, and Dewar 2015; O'Brien et al. 2014).

Another challenge when studying mortality is that the physiological processes governing tree vulnerability may vary in space and time. For instance, vulnerability may vary among individual trees within a population according to (i) the local spatial heterogeneity of resources availability,

especially soil water (Nourtier et al. 2014); (ii) the heterogeneity in individual life history, and in particular the legacies of past stresses on tree morphology and anatomy (Vanoni et al. 2016); (iii) the inter-individual variation of physiological response to stresses, which depends on ontogenic, plastic, and genetic effects controlling the expression of traits (Anderegg 2015; Vitasse et al. 2009). Vulnerability may also vary through time during the year for a given individual/population, not only because of temporal climatic variation but also through individual variation in phenological processes. This is well illustrated by the risk of late frost damages, which tightly depend on the coincidence between temporal patterns of budburst phenology, and the climatic sequence of low temperatures. Although relatively large safety margins were found regarding the risk of late frost damages during budburst across many European temperate tree and shrub species (Bigler and Bugmann 2018), these safety margins may reduce with climate change, due to the advance in the budburst date of trees (Augspurger 2009). When young leaves have been damaged, some species can reflush, i.e. produce another cohort of leaves (reflush, Augspurger 2009; Menzel, Helm, and Zang 2015), but the time required to reflush leads to a shorter growing season (Lenz et al. 2013), and eventually to mortality if trees do not have enough reserves to reflush.

Available approaches to investigate the multiple drivers and processes underlying tree mortality can be classified into two broad categories: statistical, phenomenological approaches vs process-based, mechanistic approaches. Statistical approaches use forest inventory data to test how endogenous factors (e.g. related to tree/stand size and growth rate) and exogenous factors (the biotic and abiotic environment, including management) affect stand- or individual- mortality rates. In the last decades, many statistical studies compared mortality rates among species or populations over large climatic gradients and demonstrated the overall positive effect of drought severity on mortality, although usually explaining only a limited proportion of the variance in mortality rates (Allen et al. 2010; Greenwood et al. 2017). Moreover, mortality rates were predicted with a higher accuracy when individual covariates for tree growth, size and/or competition were included in the statistical models, highlighting the importance of inter-individual variability in the threshold to mortality (Hülsmann, Bugmann, and Brang 2017; Monserud 1976). Recent statistical studies attempted to include functional traits involved in the response to stress as additional covariates to improve the accuracy of mortality prediction. For instance, Carnicer et al. (2011) showed that defoliation trends are consistent with mortality trends in southern European forests. Benito Garzón et al. (2018) found that mortality increased in populations with negative hydraulic safety margins for 15 species out of the 25 studied. Overall, the main advantage of statistical approaches is their ability to account for a potentially high number of factors and processes triggering mortality, and for individual variability in the threshold to mortality. However, one should keep in mind that statistical models hardly handle co-linearity and non-linearity of effects and non-randomization inherent to

natural population designs. In addition, the accuracy of statistical predictions dramatically decreases

outside the studied area (Hülsmann , Bugmann, and Brang 2017).

On the other hand, biophysical and ecophysiological process-based models initially developed

to simulate carbon and water fluxes in forest ecosystems can also be used to investigate the

environmental drivers and physiological processes triggering tree mortality. For example, using the

model CASTANEA, Davi & Cailleret (2017) showed that mortality of *Abies alba* in southern

France resulted from the combination of drought-related carbon depletion and pest attacks. Using

six different models, Mcdowell et al. (2013) found that mortality depended more on the duration of

hydraulic stress rather than on a specific physiological threshold. A main advantage of process-

based models is their ability to predict mortality outside the study area (i.e. under new combinations

of forcing variables in a changing environment). Their main limitation is their large number of

parameters, which makes it difficult to use them in populations where the model has not been

precisely calibrated and validated. Moreover, biophysical and ecophysiological process-based

models generally do not well take into account individual effects at tree level. Hence, statistical and

process-based approaches appear as complementary, and many authors call for studies comparing or

combining them (Hawkes 2000; O'Brien et al. 2017; Seidl et al. 2011).

An iconic tree of pan-European forests, the European beech (*Fagus sylvatica* L Crantz)

combines a widespread distribution and a high-predicted sensitivity to climate change (Cheaib et al.

2012; Kramer et al. 2010). In particular, bioclimatic niche models predict a future reduction of this

species at the rear-edge of its distribution over the next few decades, in response to reduced

precipitations (Cheaib et al. 2012; Kramer et al. 2010). This prediction is consistent with this

species' well-known sensitivity to summer droughts. For instance, more frequent extreme drought

events have been associated in beech to decreased growth (Dittmar, Zech, and Elling 2003; Jump,

Hunt, and Penuelas 2006; Knutzen et al. 2017), altered physiological performances (Bréda et al.

2006) and increased defoliation (Penuelas and Boada 2003). However, the low mortality rate

observed so far in beech has led some authors to propose that this species presents a higher heat

stress tolerance and metabolic plasticity as compared to other tree species (García-Plazaola et al.

2008). This apparent paradox between a low mortality and a high sensitivity to climate makes of

beech an interesting model species to study mortality.

In this study, we used a combination of statistical regression models, and the process-based

model CASTANEA to investigate patterns of mortality within a population located at the warm and

dry ecological margin of European beech (Supplementary Fig. S1). Growth, decline, mortality and

budburst characteristics have been monitored for 14 years in a set of 4327 adult trees. First, we

tested the impact of climatic variables on population-level mortality rate with a beta-regression

model, simultaneously using CASTANEA to investigate which physiological mechanisms drove

mortality at population scale. Then, we used logistic regression to characterise the respective effects of exogenous (competition) and endogenous factors (size, growth, crown defoliation, fungi presence) on the individual probability of mortality, simultaneously using CASTANEA to simulate trees with different endogenous characteristics and to investigate differences of physiological responses between individuals.

Materials and Methods

Study site

La Massane (42° 28' 41" N, 3° 1' 26" E) is a forest of 336 ha located in the French eastern Pyrenees; its elevation ranges from 600 m to 1127 m above sea level. Located at the South of beech distribution area, the forest is at the junction of Mediterranean and mountainous climate with mean annual rainfall of 1260 mm (ranging from 440 to 2000 mm) and mean annual temperature of 11°C (with temperature from -10°C to 35°C) (Supplementary Fig. S2). No logging operation was allowed since 1886 and the forest was classified as a reserve in 1974. European beech is the dominant tree in the canopy representing about 68% of basal area of the forest. Beech is in mixture with downy oak (*Quercus pubescens* Wild), maples (*Acer opalus Mill., Acer campestris L., Acer monspessulanum L.*), and holly (*Ilex aquifolium L.)*. A 10-ha fenced plot was remoted from cow grazing since 1956. All trees from this protected plot were geo-referenced and individually monitored since 2002 (Supplementary Fig. S3).

We estimated the soil water content (SWC) through soil texture, soil depth and percentage of coarse elements measured in two soil pits. Secondly, we estimated the mean Leaf Area Index (LAI) by using hemispherical photographs (Canon 5D with Sigma 8mm EXDG fisheye). We computed the LAI and clumping index following the methodology described in (Davi et al. 2009).

Monitoring of tree characteristics

This study is based on the monitoring of 4327 beech trees in the protected plot from 2002 to 2016. Beech sometimes produces stump shoots resulting in multiple stems at a single position (coppice). Here, every stem of all coppices was individually monitored and called "tree" in the following.

Tree mortality was recorded twice a year, in spring and autumn, from 2003 to 2016. We considered that a tree died at year n-1 when its buds failed to burst at spring of year n, and that a tree died at year n when budburst occurred at spring but no leaves were remaining at autumn of year n, and no budburst occurred at year n+1. All the 4327 trees were alive at year 2003 (Supplementary Fig. S4). We computed the annual mortality rate (τ) for each year *n* as:

$$\tau_n = \frac{N_{dead,n}}{N_{alive,n-1}} \text{ (eq. 1),}$$

where $N_{dead,n}$ (respectively $N_{alive,n}$) is the number of dead (respectively alive) trees at year n.

Several variables where measured or computed at tree level (Table 1). Diameter at breast

height (i.e. 1.30 m above ground) was measured in 2002 and 2012 (respectively DBH and

DBH2012). Here we focus on the impact of climate change on mature trees, hence only trees with

DBH greater than 10 cm in 2002 were retained for the analysis. Individual growth was measured by

the relative increment in basal area (MBAI) between 2002 and 2012, estimated as: $MBAI_i=$

$(\pi(DBH2012_i - DBH_i)/4)/N_{yearsAlive, i}$, where $N_{yearsAlive, i}$ is the number of year where individual i was

observed being alive. For a subset of 1199 trees, height in 2002 was estimated.

A strong individual variation in budburst phenology is documented in La Massane (Gaussen

1958; Perci du Sert 1982), with ~10% beeches initiating budburst about 2 weeks before all the

others. Here, budburst was thus treated as a binary qualitative variable, and our sample included 237

such "early" beeches.

The presence of major defoliated branches and leaves was recorded each year between 2003

and 2016 (except 2010) as a qualitative measure (DEF=1 for presence; DEF=0 for absence). These

annual measures were cumulated and weighted over the observed period as:

$$DEFw = \frac{\sum_{i=1}^{Nalive}(DEF_i)}{N_{alive}} \text{ (eq 2),}$$

where Nalive is the number of years where a tree was observed as alive (year 2010 was not taken

into account). DEFw is an integrative, ratio ordered variable combining the recurrence of

defoliation, and the ability to recover from defoliation.

The presence of fructification of the saproxylic fungus *Oudemansiella mucida* was recorded

as a qualitative measure (Fungi = 1 for presence; Fungi = 0 for absence). Because once observed,

the fructification persists all the subsequent years, we analysed it as a binary variable.

Competition around each focal beech stem was estimated by four variables. Firstly, we used

the number of stems in the coppice as an indicator of within-coppice competition (Nstem).

Secondly, we used the competition index introduced by Martin and Ek (1984), which accounts

simultaneously for the diameter (DBH) and the distance (d_{ij}) of each competitor *j* to the competed

individual *i*:

$$\text{Compet}_{i,dmax} = \frac{1}{DBH_i}\sum_{j=1}^{n} DBH_j \exp\left[\frac{-16d_{ij}}{DBH_i + DBH_j}\right] \text{ (eq 3),}$$

where, *n* is the total number of competitors in a given radius dmax (in m) around each focal

individual i. Only trees j with $DBH_j > DBH_i$ are considered as competitors. This index was

previously shown to describe more accurately the competition than indices relying on diameter only

(Stadt et al. 2007). We computed three competition indices: the intra-specific competition index

(Compet$_{intra}$) which only accounts for the competition of beech stems not belonging to the coppice

of the focal tree; the intra-specific competition index (Compet$_{intra+}$) which accounts for all beech

stems belonging or not to the coppice of the focal tree; and the total competition index (Compet$_{tot}$),

which accounts for all stems and species. We considered that stems located less than 3 m away from

the focal stem belonged to the same coppice. All the indices were computed at all distances from

1m (or 3m for Compet$_{intra+}$) to 50 m from the target tree, with 1m step. We retained d$_{max}$= 15m,

because all indices reached a plateau after this value, suggesting that in a radius greater than 15m,

the increasing number of competitors is compensated by the distance.

Table 1: Quantitative (A) and qualitative (B) variables monitored at individual level. All the
variables were measured in 4327 trees, except for H_2002 (1199 trees).
230

Code	Quantitative variable	mean	min	max	unit
DBH	Diameter at breast height measured in 2002	21.65	10	116	cm
DBH_2012	Diameter at breast height measured in 2012	22.84	10	116	cm
MBAI	Mean basal area increment between 2002 and 2012.	0.13	0	2.8	cm-2.year^{-1}
H_2002	Height measured in 2002	8.8	2	26	m
DEFw	Defoliation mark cumulated and weighted by the number of living year over the period	0.14	0	1	
Nstem	Number of stems observed in the coppice	1.54	1	11	
Compet intra	Intra-specific competition index	2.74	0	11.44	
Compet intra+	Intra-specific competition index accounting also for competition within the coppice	2.35	0	10.052	
Compet$_{tot}$	Total competition index, where we accounted for all stems from every speciesB	4.63	0.12	19.98	

Code	Qualitative variable	Level	Number of trees	
Fungi	Presence (1) or absence (0) of the saproxylic fungus	1	414	
		0	3913	
Budburst	Beeches initiating budburst about 2 weeks before all the others (1) or not (0)	1	237	
		0	4090	

Climate data

Local climate was daily monitored on site since respectively 1976 for temperature and 1960

for precipitation and mean relative humidity. In order to obtain a complete climatic series, we used

the quantile mapping and anomaly method in the R package "meteoland" (De Caceres et al. 2018),

considering the 8-km-resolution-SAFRAN reanalysis (Vidal et al. 2010) as reference. From the

corrected climate series, 14 bioclimatic variables were computed using the "dismo" R package

(Hijmans et al. 2017) (Online Appendix 1). We also derived several drought-related variables from

the standardised precipitation-evapotranspiration index (SPEI) using the R package "SPEI"

(Beguería and Vicente-Serrano 2017). We modified the calibration of evapotranspiration (ETP),

adapted to crop in SPEI package, and used the ETP estimation provided by CASTANEA. The SPEI

variables were the maximum, minimum, mean and the cumulated SPEI from June to August,

computed over one, three and twelve months. The number of late frost days (i.e. the number of days

with negative temperature occurring after beech budburst) and a late frost index (the cumulative

amount of negative temperatures occurring after beech budburst) were also computed. In total, 28

climatic variables were considered (Online Appendix 1).

Statistical model of annual mortality rate at population level

We used beta-regression models to investigate the effect of climatic variables on the annual

mortality rate at population-level. Beta-regression models predict a response variable varying

between [0,1], and account as for features like heteroscedasticity or asymmetry, which are

commonly obtained in time-series of annual mortality rates. To face the problem of explaining 14

observations with 28 climatic variables highly correlated among each other's, we first grouped them

by category (precipitation, temperature, frost, SPEI) and selected between one and four variable per

category (Online Appendix 1). We ended-up with the following complete model for mortality at

year n (with n varying from 2004 to 2016):

$$\tau_n = MeanTColdestQ_n + MeanTWarmestQ_n + PDriestM_n + PWettestM_n + NLF_n +$$

$$SPEI12\_mean_n + SPEI1\_min_n + SPEI3\_JJA_n \text{ (model I)},$$

where MeanTDriestQ is the mean temperature of the driest quarter; PDriestM and PColdestQ are

the mean precipitation of the driest and coldest quarter respectively; NLF is the number of late frost

days; $SPEI12\_mean_n$ and $SPEI1\_min_n$, $SPEI\_max_n$ are the mean and minimum SPEI values

computed over respectively 12 and 1 month-references; and SPEI3_JJA is is the mean SPEI

computed over June, July and August (Online Appendix 1).

Then, we used a stepwise, backwards procedure to select the best model based on Akaike's

Information Criterion (AIC, Akaike 1987), and to test for interactions between factors (Online

Appendix 1). Beta-regression was fitted with the R package betareg (Cribari-Neto and Zeileis

2010). We investigated model validity by checking the leverage points (i.e. points having a greater

weight than expected randomly) with the Cooks distance (Cook distance < 0.5 indicate not

leverage). We tested the goodness-of-fit with the Brier test score (Brier 1950). To explore the

sensitivity and specificity of the model, we use also the receiver operating characteristic (ROC)

curve.

Statistical model of mortality at individual level

We used logistic regression models to investigate how tree characteristics affect the individual

probability of mortality ($P_{mortality}$). This approach is appropriate for a binary response variable, and a

mixture of qualitative and quantitative explanatory variables not necessarily normally distributed

(Hosmer and Lemeshow 2000). We considered the following complete logistic regression models

for $P_{mortality}$:

$P_{mortality}$ = [DEFcum + Fungi + Budburst+ MBAI + (Nstem **OR** $Compet_{intra}$ **OR** $Compet_{intra+}$

OR $Compet_{inter}$)] ×(DBH+DBH²) (model II)

where defoliation (DEFcum), growth (MBAI), size (DBH) and competition (Nstem and the Compet

indices) factors were quantitative variables, while fungus presence (Fungi) and budburst phenology

were qualitative variables. We included both a linear and quadratic effect of DBH by specifying this

factor as a polynomial of degree 2. Interaction effects of each factors with this polynomial were

included.

To select the best competition-related variables, we first ran model II with each competition

term successively (Online Appendix 3). Then, we used a stepwise procedure to select the most

parsimonious model based on AIC. When two models had similar AIC (delta < 2) (Arnold 2010),

the one with less variable (most parsimonious) was selected.

Collinearity resulting from correlations between predictor variables is expected to affect

statistical significance of correlated variables by increasing type II errors (Schielzeth 2010). To

evaluate this risk, we first checked for correlation among factors included in model II (Fig. S5). We

also computed the generalized variation inflation factor (GVIF) with the R package "car" (Fox and

Weisberg 2011). A threshold of $GVIF^{1/2dof}$ < 2 is commonly accepted to considers that variables are

not too much correlated and do not make the model unstable. Note that here, we aim at identifying

the climatic factors and the tree characteristics involved in mortality and not to maximize the

prediction of mortality.

As classically done in logistic regression, results were expressed in terms of odd ratios, also

called relative risk. The odd ratio, is a statistical measure, expressing the degree of dependency

between variables (eq: 4 a,b). For instance, the odds-ratio for mortality as a function of Budburst

are:

$$Oddsratio_{earlysvsnormal} = \frac{Odds_{early}}{Odds_{normal}} \qquad (eq\ 4,a),$$

$$With\ Odds_{early} = \frac{P_{mortality}(early)}{1-P_{mortality}(early)}\ (eq\ 4,b),$$

We computed odd ratios with "questionr" the R package (Barnier, Briatte, and Larmarange 2018).

The interactions were visualized with the package "jtools" (Long 2018).

Simulations with CASTANEA process-based model

306 **CASTANEA model:** CASTANEA is a process-based model used to simulate carbon and water

307 fluxes in forest ecosystems with no spatial-explicit representation of trees (Dufrêne et al. 2005). A

tree is abstracted as six functional elements: leaves, branches, stem, coarse, fine roots and reserve (corresponding to non-structural carbohydrates). The canopy is divided into five layers of leaves. Photosynthesis is half-hourly estimated for each canopy layer using the Farquhar et al. (1980) model analytically coupled to the stomatal conductance model proposed by Ballet al. (1987). Maintenance respiration is estimated as proportional to the nitrogen content of the considered organs (Ryan 1991). Growth respiration is estimated from growth increment combined with a construction cost specific to the type of tissue (De Vries, Brunsting, and Van Laar 1974). Transpiration is also hourly calculated using the Monteith (1965) equations. The dynamics of soil water content (SWC; in mm) is estimated daily using a three-layers bucket model. Soil drought drives stomata closure via a linear decrease in the slope of the Ball et al. (1987) relationship, when relative SWC is under 40% of field capacity (Granier, Biron, and Lemoine 2000; Sala and Tenhunen 1996). In the carbon allocation sub-model (Davi et al., 2009; Davi & Cailleret 2017), the allocation coefficients between compartments (fine roots, coarse roots, wood, leaf and reserves) are estimated daily depending on the sink force and the phenology constraints. More details are provided in Dufrêne et al. (2005). CASTANEA model was originally developed and validated at stand-scale for beech (Davi et al. 2005).

In this study, we used the CASTANEA version described in Davi and Cailleret (2017) with two major modifications. First, for budburst, we used the one-phase UniForc model, which described the cumulative effect of forcing temperatures on bud development during the ecodormancy phase (Chuine, Cour, and Rousseau 1999; Gauzere et al. 2017). We computed the number of late frost days (NLF) as the sum of late frost days experimented after budburst initiates.

Second, from the calculation of daily midday water potential and beech vulnerability curve to embolism (eq5), we estimated the Percentage of Loss of Conductance (PLC) (eq:6) and simulated the defoliation by a loss of LAI. The leaf water potential Ψ_{leaf} was computed as:

$$\Psi_{leaf}(t+1) = \Psi_{soil}(t+1) - \frac{TR}{3600} \times R_{SoilToLeaves} + \frac{\Psi_{leaf}(t)}{\Psi_{soil}(t+1) + TR \times R_{SoilToLeaves}}$$
$$\times e^{\frac{deltaT}{R_{SoilToLeaves} \times Cap_{SoilToLeaves}}} \ (eq: 5),$$

where soil water potential (Ψ_{soil} MPa) was calculated from daily soil water content (Campbell 1974); leaf water potential was estimated hourly (deltaT=3600s) from simulated sapflow (TR in mmol.m^{-2}.leaf^{-1}). We simulated TR following the soil-to-leaves hydraulic pathway model used in (Loustau et al. 1990). We used one resistance (RsoilToleaves in MPa.m^{-2}.s^{-1}.Kg^{-1}) and one capacitance (CapSoilToleaves in kg.m^{-2}.Mpa^{-1}) along the pathway. The Campbell (1974) resistance ($R_{soilToleaves}$) was assessed using midday and predawn water potentials found in the literature.

PLC computation was adapted from Pammenter and Willigen (1998) formula:

$$PLC = \frac{1}{1+e^{slope(\Psi_{minSoil}-\Psi_{P50})}} \text{ (eq:6),}$$

with $\Psi_{min\ soil}$ (MPa) the simulated midday water potential in the soil and Ψ_{50} (MPa) the species-specific potential below which 50% of the vessels are embolized, slope is a constant fixed to 50. We have added an option in the model that allows to simulate branch mortality. We simulate branch mortality and defoliation induced by this mortality from the emboli level of xylem vessels, assuming that branch mortality is proportional to PLC. The LAI is thus reduced during the season by this process. To qualify the effect of this process, we compared simulations with branch mortality modelling and simulations without.

We used PLC, NLF and carbon reserves as proxies of tree responses to drought and late frost stresses. In addition to investigate the interaction between these stresses without a priori, we computed a compound vulnerability index (CVI) combining PLC, NLF and carbon reserve with an identical weight:

$$CVI_n = \left(\frac{PLC_n}{max(PLC_{period})} + \frac{NLF_n}{max(NLF_{period})}\right) - \frac{Reserve_n}{max(Reserve_{period})} \quad \text{(eq:7),}$$

with n, the focal year. The composite index could vary between two and minus one.

Simulation design: First, we simulated a population of 100 trees representing the variability in individual characteristics observed in La Massane (height-diameter allometry, DBH, leaf area index and early/late budburst). The values of these individual characteristics were randomly drawn in a Gaussian law with mean and standard deviation measured in the whole population (Online appendix 3). We also simulated a range of environmental conditions representing the variability: in SWC and tree density randomly drawn in uniform law (Online appendix 3).

Second, we simulated 16 individuals corresponding to a complete cross design with four size categories (5, 15, 30, 40 cm in DBH), two budburst type (about 10 % early and about 90% 2 weeks later), and two defoliation level (Defoliated or Non-defoliated).

Results

Statistical model at population level

The total mortality rate reached 23% of the monitored trees between 2004 and 2016. The highest annual mortality rates were observed in 2006 (3.3%) and 2010 (2.9%); the lowest in 2008 (0.7%) and 2009 (0.9%).

The best beta-regression model for annual population-level mortality rate explained 74% of its variation among years (Table 2). The best model had both a high validity and goodness-of-fit (online appendix A1). It included spring and summer SPEI (SPEI_JJA), the precipitation of the

driest month (PdriestM) and the interaction between them (Table 1). SPEI_JJA and PdriestM decreased the relative mortality rate by 1.15 and 3.03 times (p-value < 0.01), respectively, while their interaction increased the relative mortality rate by 5.37 times (p-value < 0.01). The effect of PdriestM corresponded to an effect of summer droughts for the first part of the studied period (2004-2010) and to an effect of winter drought in the most recent years (2009, 2011-2016) (online appendix A1). The three highest observed annual mortality rates (>2%) coincide with the driest month of the year being in summer.

Table 2: Parameter estimates and related statistics for the best model I (beta regression at population level). β is the maximum likelihood estimate, with its estimated error (SE), z-value, and associated p-value (Pr(>|z|). OR is the odds ratio. This table shows only the factors with significant effects (see Appendix A1 for complete results).

Variables	β	SE	z value	Pr(>|z|)	OR
SPEI3_JJA	-0.15	0.054	-2.71	8.10e-03	0.87
PDriestM	-1.10	0.28	-3.89	1.10e-04	0.33
SPEI3_JJA : PdriestM	1.69	0.31	5.37	1.15e-07	5.37

Process based model at population scale

CASTANEA simulated in 2006 an important increase in PLC, with a population mean of 31% and a range from 14% to 64% among individuals (Fig. 1a). Four other years had a high mean PLC (>10%): 2009, 2010, 2012 and 2015, with respective PLC of 18%, 18%, 14% and 14%.

Simulated carbon reserves decreased in years 2006 (51 gC.m²), 2010 (186 gC.m²) and 2015 (175 gC.m²). After a peak in 2011, with 354 gC.m², carbon reserves continuously decreased until 2015 (Fig. 1b).

Finally, simulated trees suffered seven years of late frost during the period, with a peak in 2005 where individuals suffered from 4 to 8 days of late frost depending on individual phenology. Two other years of high late frost occurred in 2010 and 2012, with respectively from 5 to 9 days and from 5 to 7 days of late frost (Fig. 1c).

None of these indicators was significantly correlated to annual variation of ring width or mortality rate. But, the composite vulnerability index was significantly correlated to the annual mortality rate observed between 2004 and 2016 (r= 0.58, p-value 0.04) (Fig. 1d).

399 **Figure 1: Proxies of physiological stresses simulated by CASTANEA at population-level from**
400 **2004 to 2016: (a) percentage of loss of conductance (PLC); (b) biomass of reserve; (c) number**
401 **of late frost day;. (d) Composite vulnerability index integrating a+b+c.** The black line is the
402 mean of simulation, and the grey envelope represent the variation from the 1st to the 3rd quartile.
403 The yellow line is the mortality rate observed in La Massane.
404

Statistical model of mortality at individual level

All the variables listed in model II had a significant main effect on the probability of mortality (Table 3) and were retained in the best model. This model explained 49 % of the observed mortality (i.e. dead individuals truly observed dead) and had both a high validity and goodness-of-fit (Online appendix 3 for details). Among the competition indices, N_{stem} was selected as it led to the fit with the lower AIC (Online Appendix 3). The relative probability of mortality increased by 1000, 1.13, 2.25 and 1.75 times, respectively for trees with a high defoliation, a higher density of coppice stems, an earlier budburst and the presence of fungi (Table 3). By contrast, trees with high mean growth had 1.06 times lower probability of mortality. The polynomial of degree 2 correspond to a U-shape and traduced a higher relative probability of mortality for both the smaller and the bigger trees (Table 3, Online appendix 3).

Table 3: Parameter estimates and related statistics for the model II (logistic regression at individual level). β is the maximum likelihood estimate, with its estimated error (SE), z-value, and associated p-value (Pr(>|z|)). OR is the odds ratio.

Variables	β	SE	z value	Pr(>|z|)	OR
DEFw	6.91	0.26	26.29	< 2.2 e-16	1.0 e03
DBH	-11.59	5.15	-2.25	2.42 e-02	9.22 e-06
DBH²	22.25	4.95	4.50	6.86 e-06	4.60 e09
MBAI	-5.91e-02	1.06 e-02	-5.55	2.80 e-08	0.94
Budburst (Early)	0.81	0.17	4.69	2.69 e-06	2.25
Nstem	0.12	3.60 e-02	3.42	6.24 e-04	1.13
Fungi (Presence)	0.56	0.14	3.94	8.10 e-05	1.75
DEFw:poly(DBH)1	-55.17	14.81	-3.73	1.95 e-04	1.10 e-24
DEFw:poly(DBH)2	32.09	16.81	1.91	5.63 e-02	8.64 e13
poly(DBH)1:MBAI	0.37	0.45	0.83	0.41	1.45
poly(DBH)2:MBAI	-0.83	0.44	-1.89	5.90 e-02	0.43

Interactions effects on mortality between size and defoliation were significant (Fig. 2a). At equal level of defoliation, the probability of mortality increased significantly more for smaller trees (Fig. 2a, Table 3). At equal mean growth, the relative probability of mortality for smaller trees was lower than for bigger trees. This interaction is only true for lower mean growth (MBAI <25 cm².y^{-1}) (Fig. 2b). Only the strength of the effects changed for alternative models considering the stem not belonging to a coppice and the model with DBH as qualitative variable (Online appendix 3).

Figure 2: Interaction effects in model II (a) between dimeter (DBH) and weighted defoliation (DEFw). (b) between DBH and the mean growth in basal area (MBAI). Regression lines are plotted for three values of DBH, corresponding to +/- 1 standard deviation (10.7cm) from the mean (22 cm). Confidence interval at 95% are shown around each regression line.

Process based model at individual scale

Simulations for the different individuals showed that tree vulnerability was driven by the intensity of climatic stress and by tree individual characteristics. The process-based model allowed to identify the factors that increased individual tree vulnerability during a drought year (2006) and a frost year (2010) and those that enhance their efficiency during a good year (2008). Drought-induced embolism was higher for early budburst trees and for bigger trees. The ability to defoliate

was limiting the risk of cavitation (Fig. 3a), but The ability to defoliate increased the risk of carbon

starvation (Fig. 3b). The drought induced risk of carbon starvation (i.e. low reserve) was also higher

for bigger trees and early trees. The resilience measured by carbon reserve in 2008 is higher for

smaller, early, and non-defoliated trees (Fig. 3b). The positive effect of earlier budburst remains

high even a year of frost in 2010 (Fig. 3b).

Figure 3: Physiological proxies of vulnerability simulated for eight trees differing in size,
defoliation and budburst phenology. We focus on three key years: 2006 (severe drought); 2008
(no stress); 2010 (late frosts). Colours indicate the initial DBH at the beginning of the simulation:
5cm (light blue) versus 40 cm (dark green). Triangle (respectively round) symbols indicate
individuals with early (respectively "normal") budburst. Empty (respectively full) symbols indicate
individuals able (respectively not able) to defoliate.

Discussion

A relatively high population-level rate of mortality in response to drought and frost

The annual mortality rates observed in this study ranged between 0.7 and 3.3%. The statistical

model explained 74% of its variation among years, which is higher than usually observed (30-40%)

(Greenwood et al. 2017). Their best abiotic predictors were the mid-term drought from June to

August (SPEI_JJA) and the amount of precipitations in the driest month (PdriestMonth).

Meanwhile, simulations suggested that a combination of late frost sensitivity, loss of hydraulic

conductance and reduction of carbon reserve drove population-level mortality rate.

The annual mean rate of mortality estimated here (2%) is in the upper range of the (few)

mortality estimates available for beech. Hülsmann et al. (2016) reported annual mean rates of

mortality of 1.4%, 0.7% and 1.5% in unmanaged forests of Switzerland, Germany and Ukraine,

with a maximum of mortality rate of 2.2%. Archambeau et al. (2019) estimated even lower mortality rates (mean annual value=0.0038%, range = 0.00374% to 0.00382%) from European forest inventory data (including managed and unmanaged forests). Overall, these mortality rates are low as compared to other tree species; for instance, according to the French national forest inventory, the average mortality is 0.1% for Beech against 0.3% in average and 0.4% for spruce or 0.2% for silver fir (IFN 2016). The relatively high value observed here may result from the absence of management, combined with the population location at the dry, warm species margin (Fig. S1), where most population extinctions are expected in Europe (Thuiller et al. 2005).

Moreover, our results suggest that the high observed mortality is primarily driven by summer droughts, including both the pulse effects of severe drought (PdriestMonth) but also long-term effects of repeated droughts (through SPEI_JJA). The major role of SPEI on beech mortality was also found of Archambeau et al. (2019), while Hülsmann et al. (2016) found a major role of competition (except for stroms in Ukraine). A long-term effect of drought was also found by Davi and Cailleret (2017) in *Abies alba Mill.*, or by Carnicer et al., (2011) in 12 European tree species, but our results additionally suggest positive interaction of long-term and pulse effects of drought on increasing risk of mortality.

Another major result from this study is that besides drought, other sources of stresses such as late frosts cannot be neglected to predict patterns of mortality in beech. The benefit of integrating drought- and frost-related stresses was particularly clear when we combined the number of late frost days, the percentage of loss of conductance and the level of carbon reserves into a composite vulnerability index. Indeed, this composite vulnerability index yielded a good correlation between predicted and observed mortality at population level. In particular, simulations showed that in 2010 (a year without drought), the high-observed mortality rate coincides with an extreme late frost event. This is consistent with the emerging consensus that mortality at dry, warm margins is not due either to carbon starvation or to hydraulic failure, but rather results from a balance between stresses vulnerability and carbon assimilation (e.g. McDowell et al. 2011; Sevanto et al. 2014). These results are also consistent with study of Vanoni et al (2016), which showed that both drought and frost could contribute to beech mortality.

In future developments, the composite vulnerability index could be refined by weighting its different components. In addition, in this study, we could not account for the temporal dynamics of mortality, such as the existence of positive or negative post-effects across years. But whenever possible(for instance in dendrochronological studies), such effects could be included, especially since there may be a lag between the weakening of a tree and its final death as already shown for beech tree in Vanoni (2016) and silver fir in Davi & Cailleret (2017).

Inter-individual variation in the vulnerability to drought and frost

The main strength of this study is the large number of trees individually monitored, which provided us with unusual large sample size to test for the effects of factors modulating individual vulnerability to mortality in beech. Firstly, we confirmed that mean growth has a positive effect on survival, as previously shown (Cailleret and Davi 2011; Gao et al. 2018). Moreover, interaction effects between growth and size were detected, such that for weak mean growth, bigger trees were less vulnerable to mortality than smaller trees. The probability of mortality also decreased faster with increasing mean growth for small than big trees. The variation of the relationship between mean growth and mortality according to tree size was already reported in beech seedlings (Collet and Le Moguedec 2007) and other species (Kneeshaw et al. 2006; Lines, Coomes, and Purves 2010).

Secondly, we found that defoliation increased the risk of mortality. This result was expected from previous studies (Dobbertin and Brang 2001, Carnicer et al. 2011), although the consequences of defoliation are still debated for beech. Senf et al. (2018) showed that defoliation was associated to tree decline, while Bauch et al., (1996) and Pretzsch (1996) found that the growth of highly defoliated beech trees did not decrease and could even increase in some cases. Such a positive effect of defoliation is not trivial, as drought-driven defoliation, occurring typically during summer due to vessels embolism in some branches, is expected to mechanically decrease carbon reserves during the current year. Our simulations comparing trees able or not to defoliate shed light on the multiple effects of defoliation, by showing that defoliation indeed decreased carbon reserves in good years but could also limit the embolism and the decrease of reserves during very dry years. We also observed significant interaction between defoliation and trees size on defoliation, suggesting that small trees were more vulnerable to defoliation than big trees. However, we cannot rule out this this effect is due in part to the qualitative method used to survey defoliation, which does not take into account the percentage of the crown loss. Hence, defoliation may be biased with respect to size, such that small defoliated trees will on average have a higher proportion of canopy loss, and therefore be more impacted than big tree.

Thirdly, both statistical and process-based approaches found that early individuals were more vulnerable. By contrast, (Robson et al. 2013) report no evidence of higher vulnerability of early trees, but a higher growth of early individuals, as also found in our simulation where early trees accumulate more reserves during good years. This discrepancy may be due to the fact that our studied population is at the rear-edge of beech distribution, where earliest budburst dates are observed due to higher temperature and may expose trees to a higher risk of late frost. Moreover, the presence of very early trees in the studied population makes it somehow peculiar, although

similar cases were observed (Gaussen 1958; Perci du Sert 1982). The higher vulnerability of early trees could also result from a faster depletion of their carbon reserve (Fig. S6), probably because they develop faster their canopy, which also mechanically increase their water needs, and the risk of embolism. Hence, the relationships between phenology and mortality deserve further investigation, especially since the spatio-temporal variation of budburst patterns under climate change may combine into complex spatio-temporal patterns of stresses (Vanoni et al. 2016).

Regarding the effect of size, the results differed between the statistical approach, where large trees died less than small ones, and the simulations, which predicted a greater vulnerability of larger trees. There may be several explanations to this discrepancy. The first reason is that CASTANEA model simulates an average individual, without effect of competition for light and water, and may thus underestimate the background mortality due to competition (which is higher, although weak, in small trees). In addition, CASTANEA also does not account for individual dominance status, which can affect the current carbon balance of a tree and hence the capacity to mitigate the stress. Indeed, in the observed populations, big trees are more likely to be dominant, with a better access to light resources promoting carbon accumulation, as compared to small trees, which are more likely to be suppressed. A second reason is that tree size may covary in environment quality in the studied population, such that big trees have more chance to occur on better soils, while environment do not covary with tree size in the simulations. Therefore, the size effect observed through the statistical approach may reveal a confounding effect of spatial soil heterogeneity, not taken into account in the process-based model. A measurement of water availability at individual tree level would be necessary but was out of the scope of this study.

Combining statistical and process-based approaches to disentangle the drivers of tree vulnerability

This study highlighted the interest of combining statistical and process-based approaches of mortality. These two approaches illustrate the classical compromise between fine understanding of physiological mechanisms driving mortality with complex and expensive process-based models, versus high precision in local mortality predictions with statistical models requiring less data but having also a weaker ability of generalization. Most often, studies adopt either of the two approaches, and generally statistical approaches prevails (Hülsmann et al. 2016; Seidl et al. 2011). However, the two approaches are tightly complementary, and combining them is a powerful approach to decipher the respective roles of the drivers and mechanisms underlying tree mortality and understand their variability among individuals or years (Hawkes 2000; O'Brien et al. 2017; Seidl et al. 2011).

In this study, the statistical model at individual scale, which aggregates the defoliation effect over the whole period including dry and good years, showed a predominant negative effect of defoliation, while the process-based model showed that during a severe drought, defoliation somehow limits the risk of hydraulic failure. These results overall suggest that the studied period may include so many recurrent dry years that the benefit of defoliation becomes negligible. Another contradiction was the higher vulnerability of early individuals to late frost demonstrated by the process-based model, while no effect of late frost was detected using the statistical approach at population level. However, the low proportion of early individuals in the studied population (~10 %) can explain this apparent contradiction. The major benefits of our approach combining different approaches (statistical, process-based) at different scales (population, individual) is that it allows to disaggregate ecological patterns observed at an upper scale (population, multi-year period), and to get back to patterns observed at a lower scale where processes operate (individual, year). This ability to aggregate/disaggregate patterns is acknowledged as a powerful approach to understand apparent contradictions between patterns observed at different scales (Clark et al. 2011) .

There are however some limitations to the approaches that were used here. First, none of them could fully account for the non-independence of climatic effects on mortality between years. Indeed, the effect of a climatic event at year n may depends on the effects of previous years, as already observed for beech, where successive drought lead to a growth decline (Jump et al. 2006; Knutzen et al. 2017; Vanoni et al. 2016) or even a modification in sap flow (Hesse et al. 2019). Moreover, the processes driving mortality may change through time as the most sensitive individuals are progressively eliminated, and/or the surviving trees become less and less sensitive (i.e. acclimation Niinemets 2010). Besides accounting for these limitations, another extension of the present study would be to combine statistical and process-based approaches at a larger spatial scale. This would allow investigating whether the respective drought and late frost sensitivity differ among the rear, core and leading edge of species distribution, as suggested by Cavin and Jump (2017).

Author Contributions

JAM, JG, CH and EM provided the data. CP performed the wood core analyses. CP, FL and SOM designed and ran the statistical models. CP and HD ran the process-based model. CP drafted the manuscript, and all the authors contributed to improve it.

Acknowledgments

We are grateful to Maxime Cailleret, Bruno Fady, and Nicolas Martin for discussions and comments on a previous version of this manuscript. We thank Nicolas Mariotte INRA URFM

618 Avignon for wood core sampling, and Frédéric Guibal for their analyses. The study was partly

619 funded by the EU ERA-NET BiodivERsA projects TIPTREE (BiodivERsA2-2012-15), the ANR

620 project MeCC (ANR-13-ADAP-0006). +GENTREE

References

Adams, Henry D. et al. 2017. "A Multi-Species Synthesis of Physiological Mechanisms in Drought-Induced
Tree Mortality." *Nature Ecology and Evolution* 1(9):1285–91.

Akaike, Hirotugu. 1987. "Factor Analysis and AIC." Pp. 371–86 in. Springer, New York, NY.

Allen, Craig D. et al. 2010. "A Global Overview of Drought and Heat-Induced Tree Mortality Reveals
Emerging Climate Change Risks for Forests." *Forest Ecology and Management* 259(4):660–84.

Anderegg, W. R. L. et al. 2012. "From the Cover: The Roles of Hydraulic and Carbon Stress in a Widespread
Climate-Induced Forest Die-Off." *Proceedings of the National Academy of Sciences* 109(1):233–37.

Anderegg, William R. L. 2015. "Spatial and Temporal Variation in Plant Hydraulic Traits and Their
Relevance for Climate Change Impacts on Vegetation." *New Phytologist* 205(3):1008–14.

Anderegg, William R. L. et al. 2015. "Tree Mortality from Drought, Insects, and Their Interactions in a
Changing Climate." *New Phytologist* 208(3):674–83.

Archambeau, Juliette et al. 2019. "Similar Patterns of Background Mortality across Europe Are Mostly
Driven by Drought in European Beech and a Combination of Drought and Competition in Scots Pine."
Under Review (265171).

Arnold, Todd W. 2010. "Uninformative Parameters and Model Selection Using Akaike's Information
Criterion." *Journal of Wildlife Management* 74(6):1175–78.

Augspurger, Carol K. 2009. "Spring 2007 Warmth and Frost: Phenology, Damage and Refoliation in a
Temperate Deciduous Forest." *Functional Ecology* 23(6):1031–39.

Ball, J. Timothy, Ian E. Woodrow, and Joseph A. Berry. 1987. "A Model Predicting Stomatal Conductance
and Its Contribution to the Control of Photosynthesis under Different Environmental Conditions." Pp.
221–24 in *Progress in photosynthesis research*. Springer.

Barnier, Julien, François Briatte, and Joseph Larmarange. 2018. "Questionr: Functions to Make Surveys
Processing Easier."

Bauch, Josef. 1986. "Characteristics and Response of Wood in Declining Trees from Forests Affected by
Pollution." *IAWA Journal* 7(4):269–76.

Beguería, Santiago and Sergio M. Vicente-Serrano. 2017. "SPEI: Calculation of the Standardised
Precipitation-Evapotranspiration Index."

Benito Garzón, Marta et al. 2018. "The Legacy of Water Deficit on Populations Having Experienced
Negative Hydraulic Safety Margin." *Global Ecology and Biogeography* 27(3):346–56.

Bigler, Christof and Harald Bugmann. 2018. "Climate-Induced Shifts in Leaf Unfolding and Frost Risk of
European Trees and Shrubs." *Scientific Reports* 8(1):1–10.

Brando, Paulo Monteiro et al. 2014. "Abrupt Increases in Amazonian Tree Mortality Due to Drought-Fire
Interactions." *Proceedings of the National Academy of Sciences of the United States of America*
111(17):6347–52.

Bréda, Nathalie, Roland Huc, André Granier, and Erwin Dreyer. 2006. "Temperate Forest Trees and Stands
under Severe Drought▯: A Review of Ecophysiological Responses , Adaptation Processes and Long-
Term Consequences." *Annals of Forest Science* 63(6):625–44.

Brier, Glenn W. 1950. "Verification of Forecasts Expressed in Terms of Probability." *Monthey Weather
Review* 78(1):1–3.

De Caceres, Miquel, Nicolas Martin-StPaul, Marco Turco, Antoine Cabon, and Victor Granda. 2018.
"Estimating Daily Meteorological Data and Downscaling Climate Models over Landscapes."
Environmental Modelling and Software 186–96.

Cailleret, Maxime et al. 2017. "A Synthesis of Radial Growth Patterns Preceding Tree Mortality." *Global
Change Biology* 23(4):1675–90.

Cailleret, Maxime and Hendrik Davi. 2011. "Effects of Climate on Diameter Growth of Co-Occurring Fagus
Sylvatica and Abies Alba along an Altitudinal Gradient." *Trees - Structure and Function* 25(2):265–76.

Campbell, Gaylon S. 1974. "A Simple Method for Determining Unsaturated Conductivity from Moisture
Retention Data." *Soil Science* 117(6):311–14.

Carnicer, Jofre et al. 2011. "Widespread Crown Condition Decline, Food Web Disruption, and Amplified
Tree Mortality with Increased Climate Change-Type Drought." *Proceedings of the National Academy
of Sciences* 108(4):1474–78.

Cavin, Liam and Alistair S. Jump. 2017. "Highest Drought Sensitivity and Lowest Resistance to Growth
Suppression Are Found in the Range Core of the Tree Fagus Sylvatica L. Not the Equatorial Range
Edge." *Global Change Biology* 23(1):362–79.

Cheaib, Alissar et al. 2012. "Climate Change Impacts on Tree Ranges: Model Intercomparison Facilitates
Understanding and Quantification of Uncertainty." *Ecology Letters* 15(6):533–44.

Choat, Brendan et al. 2018. "Triggers of Tree Mortality under Drought." *Nature* 558(7711):531–39.

Chuine, Isabelle, P. Cour, and D. D. Rousseau. 1999. "Selecting Models to Predict the Timing of Flowering
of Temperate Trees: Implications for Tree Phenology Modelling." *Plant, Cell and Environment*
22(1):1–13.

Clark, James S. et al. 2011. "Individual-Scale Variation, Species-Scale Differences: Inference Needed to
Understand Diversity." *Ecology Letters* 14(12):1273–87.

Collet, Catherine and Gilles Le Moguedec. 2007. "Individual Seedling Mortality as a Function of Size,
Growth and Competition in Naturally Regenerated Beech Seedlings." *Forestry* 80(4):359–70.

Cowan, I. R. and G. D. Farquhar. 1977. "Stomatal Function in Relation to Leaf Metabolism and
Environment." *Symposia of the Society for Experimental Biology* 31:471–505.

Cribari-Neto, Francisco and Achim Zeileis. 2010. "Beta Regression in R." *Journal of Statistical Software*
34(2).

Davi, H. et al. 2005. "Modelling Carbon and Water Cycles in a Beech Forest. Part II.: Validation of the Main
Processes from Organ to Stand Scale." *Ecological Modelling* 185(2–4):387–405.

Davi, H., C. Barbaroux, C. Francois, and E. Dufrêne. 2009. "The Fundamental Role of Reserves and
Hydraulic Constraints in Predicting LAI and Carbon Allocation in Forests." *Agricultural and Forest
Meteorology* 149(2):349–61.

Davi, Hendrik and Maxime Cailleret. 2017. "Assessing Drought-Driven Mortality Trees with Physiological
Process-Based Models." *Agricultural and Forest Meteorology* 232:279–90.

Dittmar, Christoph, Wolfgang Zech, and Wolfram Elling. 2003. "Growth Variations of Common Beech
(Fagus Sylvatica L.) under Different Climatic and Environmental Conditions in Europe—a
Dendroecological Study." *Forest Ecology and Management* 173(1–3):63–78.

Dobbertin, Matthias and Peter Brang. 2001. "Crown Defoliation Improves Tree Mortality Models." *Forest
Ecology and Management* 141(3):271–84.

Dufrêne et al. 2005. "Modelling Carbon and Water Cycles in a Beech Forest Part I□: Model Description and
Uncertainty Analysis on Modelled NEE." *Ecological Modelling* 185:407–36.

Durand-Gillmann, Marion, Maxime Cailleret, Thomas Boivin, Louis Michel Nageleisen, and Hendrik Davi.
2014. "Individual Vulnerability Factors of Silver Fir (Abies Alba Mill.) to Parasitism by Two
Contrasting Biotic Agents: Mistletoe (Viscum Album L. Ssp. Abietis) and Bark Beetles (Coleoptera:
Curculionidae: Scolytinae) during a Decline Process." *Annals of Forest Science* 71(6):659–73.

Farquhar, G. D., S. von Caemmerer, and J. A. Berry. 1980. "A Biochemical Model of Photosynthetic CO_2
Assimilation in Leaves of C3 Species." *Planta* 149(1):78–90.

Feng, Xue et al. 2018. "The Ecohydrological Context of Drought and Classification of Plant Responses."
Ecology Letters, November 1, 1723–36.

Fox, John and Sanford Weisberg. 2011. *An {R} Companion to Applied Regression*. Second. Thousand Oaks
{CA}: Sage.

Gao, Shan et al. 2018. "Dynamic Responses of Tree-Ring Growth to Multiple Dimensions of Drought."
Global Change Biology 24(11):5380–90.

García-Plazaola, José Ignacio, Raquel Esteban, Koldobika Hormaetxe, Beatriz Fernández-Marín, and José
María Becerril. 2008. "Photoprotective Responses of Mediterranean and Atlantic Trees to the Extreme
Heat-Wave of Summer 2003 in Southwestern Europe." *Trees* 22(3):385–92.

Gaussen, H. 1958. "Le Hêtre Aux Pyrénées Espagnoles." Pp. 185–91 in *Actas del tercer congreso
internacional de estudios pirenaicos, Gerona*.

Gauzere, J. et al. 2017. "Integrating Interactive Effects of Chilling and Photoperiod in Phenological Process-
Based Models. A Case Study with Two European Tree Species: Fagus Sylvatica and Quercus Petraea."
Agricultural and Forest Meteorology 244–245:9–20.

Gillner, Sten, Nadja Rüger, Andreas Roloff, and Uta Berger. 2013. "Low Relative Growth Rates Predict
Future Mortality of Common Beech (Fagus Sylvatica L.)." *Forest Ecology and Management* 302:372–
78.

Granier, A., P. Biron, and D. Lemoine. 2000. "Water Balance, Transpiration and Canopy Conductance in Two
Beech Stands." *Agricultural and Forest Meteorology* 100(4):291–308.

Greenwood, Sarah et al. 2017. "Tree Mortality across Biomes Is Promoted by Drought Intensity, Lower
Wood Density and Higher Specific Leaf Area" edited by J. Chave. *ECOLOGY LETTERS* 20(4):539–53.

Hawkes, Corinna. 2000. "Woody Plant Mortality Algorithms: Description, Problems and Progress."
Ecological Modelling 126(2–3):225–48.

Hesse, Benjamin D., Michael Goisser, Henrik Hartmann, and Thorsten E. E. Grams. 2019. "Repeated
Summer Drought Delays Sugar Export from the Leaf and Impairs Phloem Transport in Mature Beech"
edited by M. Dannoura. *Tree Physiology* 39(2):192–200.

Hijmans, Robert J., Steven Phillips, John Leathwick, Jane Elith, and Maintainer Robert J. Hijmans. 2017.
"Package 'Dismo.'" *Circles* 9(1):1–68.

Hosmer, David W. and Stanley Lemeshow. 2000. *Applied Logistic Regression*.

Hülsmann, Lisa et al. 2016. "Does One Model Fit All? Patterns of Beech Mortality in Natural Forests of
Three European Regions." *Ecological Applications* 26(8):2463–77.

Hülsmann, Lisa, Harald Bugmann, and Peter Brang. 2017. "How to Predict Tree Death from Inventory Data
– Lessons from a Systematic Assessment of European Tree Mortality Models - SUPP." *Canadian*
Journal of Forest Research (April):cjfr-2016-0224.

Hülsmann, Lisa, Harald Bugmann, Maxime Cailleret, and Peter Brang. 2018. "How to Kill a Tree: Empirical
Mortality Models for 18 Species and Their Performance in a Dynamic Forest Model." *Ecological*
Applications 28(2):522–40.

IGN. 2016. *La Mortalité*. France. https://inventaire-forestier.**ign**.fr/IMG/pdf/2018_**mortalite**.pdf

Jump, Alistair S., Jenny M. Hunt, and Josep Penᵤuelas. 2006. "Rapid Climate Change-Related Growth
Decline at the Southern Range Edge of Fagus Sylvatica." *Global Change Biology* 12(11):2163–74.

Kneeshaw, Daniel D., Richard K. Kobe, K. David Coates, and Christian Messier. 2006. "Sapling Size
Influences Shade Tolerance Ranking among Southern Boreal Tree Species." *Journal of Ecology*
94(2):471–80.

Knutzen, Florian, Choimaa Dulamsuren, Ina Christin Meier, and Christoph Leuschner. 2017. "Recent
Climate Warming-Related Growth Decline Impairs European Beech in the Center of Its Distribution
Range." *Ecosystems* 20(8):1494–1511.

Kramer, Koen et al. 2010. "Modelling Exploration of the Future of European Beech (Fagus Sylvatica L.)
under Climate Change-Range, Abundance, Genetic Diversity and Adaptive Response." *Forest Ecology*
and Management 259(11):2213–22.

Lebourgeois, F., N. Bréda, E. Ulrich, and A. Granier. 2005. "Climate-Tree-Growth Relationships of
European Beech (Fagus Sylvatica L.) in the French Permanent Plot Network (RENECOFOR)." *Trees*
19:385–401.

Lenz, Armando, Günter Hoch, Yann Vitasse, and Christian Körner. 2013. "European Deciduous Trees
Exhibit Similar Safety Margins against Damage by Spring Freeze Events along Elevational Gradients."
New Phytologist 200(4):1166–75.

Lines, Emily R., David A. Coomes, and Drew W. Purves. 2010. "Influences of Forest Structure, Climate and
Species Composition on Tree Mortality across the Eastern US" edited by A. Hector. *PLoS ONE*
5(10):e13212.

Long, Jacob A. 2018. "Jtools: Analysis and Presentation of Social Scientific Data."

Lorenz, Martin and Georg Becher. 2012. *Forest Condition in Europe*.

Loustau, D., A. Granier, F. El Hadj Moussa, M. Sartore, and M. Guedon. 1990. "Evolution Saisonnière Du
Flux de Sève Dans Un Peuplement de Pins Maritimes." *Annales Des Sciences Forestières* 47(6):599–
618.

Van Mantgem, Phillip J. et al. 2009. "Widespread Increase of Tree Mortality Rates in the Western United
States." *Science* 323(5913):521–24.

Maraun, Mark, Jörg-Alfred Salamon, Katja Schneider, Matthias Schaefer, and Stefan Scheu. 2003. "Oribatid
Mite and Collembolan Diversity, Density and Community Structure in a Moder Beech Forest (Fagus
Sylvatica): Effects of Mechanical Perturbations." *Soil Biology and Biochemistry* 35(10):1387–94.

Martin, George L. and Alan R. Ek. 1984. "A Comparison of Competition Measures and Growth Models for
Predicting Plantation Red Pine Diameter and Height Growth." *Forest Science* 30(3):731–43.

Mcdowell, Nate G. et al. 2013. "Evaluating Theories of Drought-Induced Vegetation Mortality Using a
Multimodel-Experiment Framework." *New Phytologist* 200(2):304–21.

McDowell, Nate G. et al. 2011. "The Interdependence of Mechanisms Underlying Climate-Driven
Vegetation Mortality." *Trends in Ecology & Evolution* 26(10):523–32.

Meir, Patrick, Maurizio Mencuccini, and Roderick C. Dewar. 2015. "Drought-Related Tree Mortality:
Addressing the Gaps in Understanding and Prediction." *New Phytologist* 207(1):1443–47.

Menzel, Annette, Raimund Helm, and Christian Zang. 2015. "Patterns of Late Spring Frost Leaf Damage and
Recovery in a European Beech (Fagus Sylvatica L.) Stand in South-Eastern Germany Based on
Repeated Digital Photographs." *Frontiers in Plant Science* 6:110.

Monserud, Robert A. 1976. "Simulation of Forest Tree Mortality." *Forest Science* 22(4):438–44.

Monteith, J. L. 1965. "Evaporation and Environment. The State and Movement of Water in Living
Organisms. Symposium of the Society of Experimental Biology, Vol. 19 (Pp. 205-234)."

Mueller, Rebecca C. et al. 2005. "Differential Tree Mortality in Response to Severe Drought: Evidence for
Long-Term Vegetation Shifts." *Journal of Ecology* 93(6):1085–93.

Niinemets, Ülo. 2010. "Responses of Forest Trees to Single and Multiple Environmental Stresses from
Seedlings to Mature Plants: Past Stress History, Stress Interactions, Tolerance and Acclimation." *Forest
Ecology and Management* 260(10):1623–39.

Nourtier, Marie et al. 2014. "Transpiration of Silver Fir (Abies Alba Mill.) during and after Drought in
Relation to Soil Properties in a Mediterranean Mountain Area." *Annals of Forest Science* 71(6):683–95.

O'Brien, Michael J. et al. 2017. "A Synthesis of Tree Functional Traits Related to Drought-Induced Mortality
in Forests across Climatic Zones" edited by J. Firn. *Journal of Applied Ecology* 54(6):1669–86.

O'Brien, Michael J., Sebastian Leuzinger, Christopher D. Philipson, John Tay, and Andy Hector. 2014.
"Drought Survival of Tropical Tree Seedlings Enhanced by Non-Structural Carbohydrate Levels."
Nature Climate Change 4(8):710–14.

Pammenter NW and Vander Willigen C. 1998. "A Mathematical and Statistical Analysis of the Curves
Illustrating Vulnerability of Xylem to Cavitation." *Tree Physiology* 18(Equation 1):589–593.

Penuelas, Josep and Martí Boada. 2003. "A Global Change-Induced Biome Shift in the Montseny Mountains
(NE Spain)." *Global Change Biology* 9(2):131–40.

Perci du Sert, Th. 1982. *Relations Entre La Phenologie et La Morphologie Du Hêtre Dans Le Massif Des
Albères*.

Pretzsch, Hans. 1996. "Growth Trends of Forests in Southern Germany." Pp. 107–31 in *Growth Trends in
European Forests*. Berlin, Heidelberg: Springer Berlin Heidelberg.

Robson, T. Matthew, Erwin Rasztovits, Pedro J. Aphalo, Ricardo Alia, and Ismael Aranda. 2013. "Flushing
Phenology and Fitness of European Beech (Fagus Sylvatica L.) Provenances from a Trial in La Rioja,
Spain, Segregate According to Their Climate of Origin." *Agricultural and Forest Meteorology* 180:76–
85.

Ryan, Michael G. 1991. "Effects of Climate Change on Plant Respiration." *Ecological Applications*
1(2):157–67.

Sala, A. and J. D. Tenhunen. 1996. "Simulations of Canopy Net Photosynthesis and Transpiration in Quercus
Ilex L. under the Influence of Seasonal Drought." *Agricultural and Forest Meteorology* 78(3–4):203–
22.

Schielzeth, Holger. 2010. "Simple Means to Improve the Interpretability of Regression Coefficients."
Methods in Ecology and Evolution 1(2):103–13.

Seidl, Rupert et al. 2011. "Modelling Natural Disturbances in Forest Ecosystems: A Review." *Ecological*
Modelling 222(4):903–24.

Senf, Cornelius et al. 2018. "Canopy Mortality Has Doubled in Europe's Temperate Forests over the Last
Three Decades." *Nature Communications* 9(1):4978.

Sevanto, Sanna, Nate G. Mcdowell, L. Turin Dickman, Robert Pangle, and William T. Pockman. 2014. "How
Do Trees Die? A Test of the Hydraulic Failure and Carbon Starvation Hypotheses." *Plant, Cell and*
Environment 37(1):153–61.

Stadt, Kenneth J. et al. 2007. "Evaluation of Competition and Light Estimation Indices for Predicting
Diameter Growth in Mature Boreal Mixed Forests." *Annals of Forest Science* 64(64):477–90.

Thuiller, Wilfried, Sandra Lavorel, M. B. Araujo, Martin T. Sykes, and I. Colin Prentice. 2005. "Climate
Change Threats to Plant Diversity in Europe." *Proceedings of the National Academy of Sciences*
102(23):8245–50.

Tyree, M. T. and J. S. Sperry. 1989. "Vulnerability of Xylem to Cavitation and Embolism." *Annual Review of*
Plant Physiology and Plant Molecular Biology 40(1):19–36.

Vanoni, Marco, Harald Bugmann, Magdalena Nötzli, and Christof Bigler. 2016. "Drought and Frost
Contribute to Abrupt Growth Decreases before Tree Mortality in Nine Temperate Tree Species." *Forest*
Ecology and Management Journal 382:51–63.

Vidal, Jean-Philippe, Eric Martin, Laurent Franchistéguy, Martine Baillon, and Jean-Michel Soubeyroux.
2010. "A 50-Year High-Resolution Atmospheric Reanalysis over France with the Safran System."
International Journal of Climatology 30(11):1627–44.

Vitasse, Yann et al. 2009. "Leaf Phenology Sensitivity to Temperature in European Trees: Do within-Species
Populations Exhibit Similar Responses?" *Agricultural and Forest Meteorology* 149(5):735–44.

De Vries, F. W. T. Pennin., A. H. M. Brunsting, and H. H. Van Laar. 1974. "Products, Requirements and
Efficiency of Biosynthesis a Quantitative Approach." *Journal of Theoretical Biology* 45(2):339–77.

848