Combining statistical and mechanistic models to identify the drivers of mortality within a rear-edge beech population

Authors:

Cathleen Petit-Cailleux[1], Hendrik Davi[1], François Lefèvre[1], Joseph Garrigue[2], Jean-André Magdalou[2], Christophe Hurson[2,3], Elodie Magnanou[2,4], and Sylvie Oddou-Muratorio[1].

Adresses

[1]INRAE, URFM, Avignon, France

[2]Réserve Naturelle Nationale de la Forêt de la Massane, France

[3]Fédération des Réserves Naturelles Catalanes, Prades, France

[4]Sorbonne Université, CNRS, Biologie Intégrative des Organismes Marins, BIOM, F-66650 Banyuls-sur-Mer, France

Keywords: *Fagus sylvatica* L., logistic regression, process-based model, dynamic vegetation model, CASTANEA, longitudinal analysis, defoliation.

Abstract

Since several studies have been reporting an increase in the decline of forests, a major issue in ecology is to better understand and predict tree mortality. The interactions between the different factors and the physiological processes giving rise tree mortality, as well as the individual variability in mortality risk, still need to be better assessed.

This study investigates mortality in a rear-edge population of European beech (*Fagus sylvatica* L.) using a combination of statistical and process-based modelling approaches. Based on a survey of 4323 adult beeches since 2002 within a natural reserve, we first used statistical models to quantify the effects of competition, tree growth, size, defoliation and fungi presence on mortality. Secondly, we used an ecophysiological process-based model (PBM) to separate out the different mechanisms giving rise to temporal and inter-individual variations in mortality by simulating depletion of carbon stocks, loss of hydraulic conductance and damage due to late frosts in response to climate.

The combination of all these simulated processes was associated with the temporal variations in the population mortality rate. The individual probability of mortality decreased with increasing mean growth, and increased with increasing crown defoliation, earliness of budburst, fungi presence and increasing competition, in the statistical model. Moreover, the interaction between tree size and defoliation was significant, indicating a stronger increase in mortality associated to defoliation in smaller than larger trees. Finally, the PBM predicted a higher conductance loss together with a higher level of carbon reserves for trees with earlier budburst, while the ability to defoliate the crown was found to limit the impact of hydraulic stress at the expense of the accumulation of carbon reserves.

We discuss the convergences and divergences obtained between statistical and process-based approaches and we highlight the importance of combining them to characterize the different processes underlying mortality, and the factors modulating individual vulnerability to mortality.

Introduction

Global changes have been repeatedly reported to be the cause of forest decline and tree mortality, both in terms of background, non-catastrophic mortality (Van Mantgem et al. 2009, Lorenz and Becher 2012) and of massive, catastrophic mortality due to extreme, 'pulse' events (Allen et al. 2010; Lorenz and Becher 2012; Mueller et al. 2005). To predict how such a new regime of trees mortality will impact upon forest structure, composition and ecosystem services (Anderegg et al. 2015a; Choat et al. 2018), we need to better understand the respective roles of the various drivers and mechanisms underlying tree mortality.

Studying mortality poses several challenges, in particular because it is triggered by several factors and involves several interacting physiological processes. The factors triggering mortality include extreme, pulse climatic events (i.e. repeated drought, storms, floods, heavy snow, late frosts, wildfires) or sudden changes in biotic interactions (i.e. emerging pests, invasive species), but also long-term climatic or biotic perturbations (i.e. recurrent water deficits, changes in competition at the community level) (Maraun et al. 2003; McDowell et al. 2011). Moreover, these factors can have interactive effects. For instance, drought may increase trees' vulnerability to pests (Durand-Gillmann et al. 2014; Anderegg et al. 2015b) or predispose them to wildfires (Brando et al. 2014). Finally, a single factor triggering mortality may involve several underlying physiological processes, with several thresholds leading to mortality and potential feedback between them (McDowell et al. 2011). This is exemplified by drought, which is usually considered to trigger mortality through the combination of hydraulic failure and carbon starvation (Adams et al. 2017; Anderegg et al. 2012; McDowell et al. 2011).

Another challenge when studying mortality is that the physiological processes governing tree vulnerability may vary in space and time. For instance, vulnerability may vary among individual trees within a population according to (i) the spatial heterogeneity in available resources, especially soil water (Nourtier et al. 2014); (ii) the heterogeneity in an individual tree's life history, and in particular the effects of past stresses on tree morphology and anatomy (Vanoni et al. 2016); (iii) the inter-individual variation of physiological responses to stresses, which depends on ontogenic, plastic, and genetic effects controlling the expression of traits (Anderegg 2015a; Vitasse et al. 2009). Vulnerability may also vary through time for a given individual/population, not only because of temporal climatic variation but also through individual variations in phenological processes. This is well illustrated by the risk of late frost damage, which is closely related to the coincidence between

temporal patterns of budburst phenology, and the climatic sequence of low temperatures. Although relatively large safety margins were found regarding the risk of late frost damage during budburst across many European temperate tree (Bigler and Bugmann 2018), these safety margins may reduce with climate change, due to earlier budburst (Augspurger 2009). When young leaves have been damaged, some species can reflush, i.e. produce another cohort of leaves (Augspurger 2009; Menzel, Helm, and Zang 2015), but the time required to reflush may reduce the length of the growing season (Lenz et al. 2013), and may lead to mortality if trees do not have enough reserves to do this.

Available approaches to investigate the multiple drivers and processes underlying tree mortality can be classified into two broad categories: statistical, phenomenological approaches versus process-based, mechanistic approaches. Statistical approaches use forest inventory data to test how tree characteristics (e.g. related to tree size and growth rate, biotic and abiotic environment, including management) affect population- or individual-level mortality. By comparing species or populations over areas with large climatic variations, such studies have demonstrated the overall effect of drought severity on mortality, although usually explaining only a limited proportion of the variance observed in mortality (Allen et al. 2010; Greenwood et al. 2017). Moreover, probabilities of mortality have been predicted with a higher accuracy when individual covariates for tree growth, size and/or competition were included in the statistical models, highlighting the importance of inter-individual variability in the threshold for mortality (Hülsmann, Bugmann, and Brang 2017; Monserud 1976). Recent statistical studies have attempted to include functional traits involved in the response to stress as additional covariates to improve the accuracy of mortality prediction, such as defoliation (Carnicer et al. 2011) or hydraulic safety margins (Benito-Garzón et al. 2018). Overall, the main advantage of statistical approaches is their ability to account for a potentially high number of factors and processes triggering mortality and for individual variability in the threshold for mortality. However, these statistical models barely deal with the usually low temporal resolution of mortality data, missing information on the cause of tree death, and non-randomization inherent in natural population designs. In addition, the accuracy of statistical predictions can decrease outside the studied area (Hülsmann, Bugmann, and Brang 2017).

On the other hand, biophysical and ecophysiological process-based models (PBMs), initially developed to simulate carbon and water fluxes in forest ecosystems, are also useful to investigate the environmental drivers and physiological processes triggering tree mortality. For example, using

the PBM CASTANEA, Davi & Cailleret (2017) showed that mortality of silver fir in southern France

resulted from the combination of drought-related carbon depletion and pest attacks. Using six

different PBMs, Mc Dowell et al. (2013) found that mortality depended more on the duration of

hydraulic stress than on a specific physiological threshold. A main advantage of PBMs is their ability

to understand how physiological processes drive mortality and to predict mortality under new

combinations of forcing variables in a changing environment. However, they need a large number

of parameters to be calibrated. Most often, calibration is made using the average parameter value

known at species level, and therefore does not account for possible inter-individual variability of

ecophysiological processes, and for its effect on response trajectory (Berzaghi et al. 2019).

Moreover, biophysical and ecophysiological PBMs generally do not take into account individual tree

characteristics (i.e. related to ontogenic, plastic and/or genetic variation). Hence, statistical and

process-based approaches appear as complementary, and many authors have called for studies

comparing or combining them (Hawkes 2000; O'Brien et al. 2017; Seidl et al. 2011).

The European beech (*Fagus sylvatica* L.) combines a widespread distribution (from northern

Spain to southern Sweden and from England to Greece) and an expected high sensitivity to climate

change (Cheaib et al. 2012; Kramer et al. 2010). Bioclimatic niche models predict a future reduction

of this species at the rear edge of its range over the next few decades (Cheaib et al. 2012; Kramer

et al. 2010). Its growth is highly sensitive to droughts (Dittmar, Zech, and Elling 2003; Jump, Hunt,

and Penuelas 2006; Knutzen et al. 2017), which increase defoliation (Penuelas and Boada 2003).

However, the low mortality rate observed so far in beech has led some authors to propose that this

species presents a higher heat stress tolerance and metabolic plasticity when compared to other

tree species (García-Plazaola et al. 2008). This apparent paradox between a low mortality and a high

sensitivity to climate makes beech an interesting model species to study.

In this study, we used a combination of statistical regression model and the PBM CASTANEA

(Figure 1), to investigate the drivers of mortality within a population located at the warm and dry

ecological margin for European beech (42° 28' 41" N, 3° 1' 26" E; Supplementary Figure S1).

Mortality, decline (crown defoliation and fungi presence), size, growth, competition and budburst

were characterised in a set of 4323 adult trees over a 15 year-period from 2002 to 2016. CASTANEA

was used to simulate the number of late frost days, the percentage loss of conductance (PLC) and

the biomass of carbon reserves in response to stress. Specifically, we addressed the following

questions: (1) How do climatic factors and physiological processes drive temporal variation in the

mortality rate? (2) How do factors varying at tree-level modulate the individual tree's probability of

mortality? (3) How do physiological mechanisms modulate the vulnerability of individuals?

**Figure 1: Combining process-based and statistical models to study variables and processes
involved in tree mortality**. The square boxes indicate the measured factors and response variables
considered in statistical models. Boxes with rounded corners indicate stress-related output variables
simulated with the PBM CASTANEA. The blue box on the left delineates the input variables of
CASTANEA. At the top, grey arrows indicate the relationships considered to link stress-related
output variables simulated by CASTANEA with observed mortality rate at population level. At the
bottom, the black arrows indicate the relationships considered in the statistical model for the
probability of mortality at individual level (solid lines: expected positive effect; dashed lines:
expected negative effect; non-linear effects were expected for size). Moreover, the effects of size,
early budburst and defoliation on the individual probability of mortality were also investigated using
the PBM. PLC – percentage loss of conductance.

Materials and Methods

Study site

La Massane is a forest of 336 ha located in the French eastern Pyrenees ranging from 600 to 1127 m.a.s.l. Located in the south of the beech range, the forest is at the junction of Mediterranean and mountainous climates with a mean annual rainfall of 1260 mm (ranging from 440 to 2000 mm) and mean annual temperature of 11°C (with daily temperature ranging from -10°C to 35°C) (Supplementary Figure S2). No logging operations have been allowed since 1886 and the forest was classified as a reserve in 1974. European beech is the dominant tree in the canopy representing about 68% of basal area of the forest. Beech is in mixture with downy oak (*Quercus pubescens* Willd), maple (*Acer opalus* Mill*., Acer campestris* L., *Acer monspessulanum* L.), and holly (*Ilex aquifolium* L.). A 10 ha fenced plot has excluded grazing from cows since 1956. All trees from this protected plot have been geo-referenced and individually monitored since 2002 (Supplementary Figure S3).

We estimated the soil water capacity (SWCa) through soil texture, soil depth and percentage of coarse elements measured in two soil pits. Secondly, we estimated the mean Leaf Area Index (LAI) by using hemispherical photographs (Canon 5D with Sigma 8mm EXDG fisheye). We computed the LAI and clumping index following the methodology described by Davi et al. (2009). SWCa and LAI were measured at population level.

Individual tree measurements

This study is based on the characterisation of twelve variables in 4323 beech trees in the protected plot over the period from 2002 to 2016 (Table 1). Note that beech sometimes produces stump shoots resulting in multiple stems from a single position (coppice), but that here, every stem of all the coppices was individually monitored and subsequently referred to "tree".

Table 1: Quantitative (a) and categorical variables (b) measured at individual level. All the variables were measured in 4323 trees, except for H_{2002} (1199 trees). The "Cat" column indicates the category (i.e. size, growth, competition, decline, phenology). The "YMeas." column indicates the year of measurement; note that all the variables were measured only once, so when two dates are given they indicate the period over which the variable is computed.

(a)

Code	Variable	Cat	YMeas.	mean	min	max	unit
DBH_{2002}	Diameter at breast height measured in 2002	Size	2002	21.7	10.0	116.0	cm
DBH_{2012}	Diameter at breast height measured in 2012	Size	2012	22.8	10.0	116.0	cm
MBAI	Mean basal area increment between 2002 and 2012.	Growth	2002 -2012	4.7	0.0	95.0	cm². year⁻¹
H_{2002}	Height measured in 2002	Size	2002	8.8	2.0	26.0	m
DEFw	Cumulated and weighted defoliation score	Decline	2003-2016	0.1	0.0	1.0	-
Nstem	Number of stems observed in the coppice	Compet	2002	1.5	1.0	11.0	-
$Compet_{intra}$	Intra-specific competition index	Compet	2002	2.7	0.0	11.4	-
$Compet_{intra+}$	Intra-specific competition index accounting for within-coppice competition	Compet	2002	1.0	0.0	12.7	-
$Compet_{tot}$	Total competition index, accounting for within-coppice competition	Compet	2002	4.6	0.1	20.0	-

(b)

Code	Variable	Cat	YMeas.	Level	Number of trees
Fungi	Presence (1) or absence (0) of the saproxylic fungus	Decline	2003-2016	1: 0:	414 3913
Budburst	Early (1) or late (0) budburst	Phenology	2002	1: 0:	237 4090

Tree mortality was recorded every year from 2003 to 2016, based on two observations (in autumn, based on defoliation and in spring, based on budburst). A tree was considered to have died at year n when (1) budburst occurred in the spring of year n but (2) no leaves remained in the autumn of year n, and (3) no budburst occurred in year n+1. All the 4323 trees were alive in year 2003 (Supplementary Figure S4). We computed the annual mortality rate (τ_n) for each year n as:

$$\tau_n = \frac{N_{dead,n}}{N_{alive,n-1}} \text{ (Equation 1)},$$

where $N_{dead,n}$ (respectively $N_{alive,n}$) is the number of dead (respectively alive) trees in year n.

Diameter at breast height (DBH) was measured 1.30 m above ground level in 2002 and 2012. As we focused on the drivers of mature tree mortality, only trees with DBH_{2002} greater than 10 cm

were retained for analysis. Individual growth was measured by the mean increment in basal area (MBAI) between 2002 and 2012, estimated as:

$$MBAI = (\pi(DBH_{2012} - DBH_{2002})^2/4)/NyearsAlive_i \qquad \text{(Equation 2)},$$

where $NyearsAlive_i$ is the number of years where individual i was observed being alive. Height in 2002 was estimated for a subset of 1199 trees.

A bimodal pattern in budburst phenology had been previously reported in La Massane (Gaussen 1958; Perci du Sert 1982). Some trees were observed to systematically initiate budburst about two weeks before all the others. Here, the monitoring allowed budburst phenology to be surveyed as a binary categorical variable, distinguishing trees with early budburst from the others.

The presence of defoliated major branches was recorded each year between 2003 and 2016 (except 2010) as a categorical measure (DEF = 1 for presence; DEF = 0 for absence). These annual measures were cumulated and weighted over the observation period for each individual in the following quantitative variable:

$$DEFw_i = \frac{\sum_{j=1}^{NyearsAlive_i} DEF_j}{NyearsAlive_i} \qquad \text{(Equation 3)},$$

Year 2010 was not included in $NyearsAlive_i$. DEFw integrates (without disentangling) the recurrence of defoliation and the ability to recover from defoliation. The presence of fructification of the saproxylic fungus *Oudemansiella mucida* (Schrad.) was recorded as a categorical measure (Fungi = 1 for presence; Fungi = 0 for absence). Given that once observed, the fructification persists throughout the subsequent years, we analysed it as a binary variable.

Competition around each focal beech stem was estimated by the number of stems in the coppice (*Nstem*) as an indicator of within-coppice competition. We also computed competition indices accounting simultaneously for the diameter (DBH$_{2002}$) and the distance (d_{ij}) of each competitor *j* to the competed individual *i*, following Martin and Ek (1984):

$$Compet_{i,dmax} = \frac{1}{DBH_{2002i}} \sum_{j=1}^{N_{compet}} DBH_{2002j} \exp\left[\frac{-16d_{ij}}{DBH_{2002i}+DBH_{2002j}}\right] \qquad \text{(Equation 4)},$$

where N_{compet} is the total number of competitors in a given radius dmax (in m) around each focal individual *i*. Only trees with DBH$_{2002}$> DBH$_{2002\ i}$ are considered as competitors. Such indices were shown to describe more accurately the competition than indices relying on diameter only (Stadt et al. 2007). We computed this competition index in three ways. The intra-specific competition index Compet$_{intra}$ only accounts for the competition of beech stems not belonging to the coppice of the focal tree. The intra-specific competition index Compet$_{intra+}$ accounts for all beech stems belonging,

or not, to the coppice of the focal tree. The total competition index $Compet_{tot}$ accounts for all stems

and species. We considered that stems located less than 3 m away from the focal stem belonged to

the same coppice. The three indices were first computed at all distances from 1 m (or 3 m for

$Compet_{intra}$) to 50 m from the target tree, with 1 m steps. We retained d_{max} = 15 m in subsequent

analyses, because all indices plateaued after this threshold value, suggesting that in a radius greater

than 15 m, the increasing number of competitors is compensated for by distance.

Climate data

Local climate has been daily monitored *in situ* since 1976 and 1960 for temperature and

precipitation/mean relative humidity, respectively. In order to obtain a complete climatic series

(from 1959 to 2016), we used the quantile mapping and anomaly method in the R package

"meteoland" (De Caceres et al. 2018), considering the 8-km-resolution-SAFRAN reanalysis (Vidal et

al. 2010) as reference.

From the corrected climate series, we derived the daily climatic input variables for CASTANEA,

which are the minimum, mean and maximum temperatures (in °C), the precipitation (mm), the wind

speed (m.s^{-1}), the mean relative humidity (%) and the global radiation (MJ.m^{-2}).

Simulations with CASTANEA

Model overview: CASTANEA is a PBM initially developed to simulate carbon and water fluxes in

forest ecosystems with no spatial-explicit representation of trees (Dufrêne et al. 2005). A tree is

abstracted as six functional elements: leaves, branches, stem, coarse roots, fine roots and reserves

(corresponding to non-structural carbohydrates). The canopy is divided into five layers of leaves.

Photosynthesis is half-hourly calculated for each canopy layer using the model of Farquhar et al.

(1980), analytically coupled to the stomatal conductance model proposed by Ball et al. (1987).

Maintenance respiration is calculated as proportional to the nitrogen content of the considered

organs (Ryan 1991). Growth respiration is calculated from growth increment combined with a

construction cost specific to the type of tissue (De Vries, Brunsting, and Van Laar 1974).

Transpiration is hourly calculated using the Monteith (1965) equations. The dynamics of soil water

content (SWCo; in mm) is calculated daily using a three-layer bucket model. Soil drought drives

stomata closure via a linear decrease in the slope of the Ball et al. (1987) relationship, when relative

SWCo is under 40% of field capacity (Granier, Biron, and Lemoine 2000; Sala and Tenhunen 1996).

In the carbon allocation sub-model (Davi et al., 2009; Davi & Cailleret 2017), the allocation

coefficients between compartments (fine roots, coarse roots, wood, leaf and reserves) are

calculated daily depending on the sink force and the phenological constraints. CASTANEA model was
originally developed and validated at stand-scale for beech (Davi et al. 2005).

Focal processes and output variables: In this study, we focussed on three response variables
simulated by CASTANEA: (1) the biomass of reserves (BoR) as an indicator of vulnerability to carbon
starvation; (2) the percentage loss of conductance (PLC) as an indicator of vulnerability to hydraulic
failure; and (3) the number of late frost days (NLF) as an indicator of vulnerability to frost damage.
Note that we did not simulate mortality with CASTANEA because the thresholds in PLC, NLF and BoR
triggering mortality are unknown. These variables were simulated using the CASTANEA version
described in Davi and Cailleret (2017) with two major modifications. First, for budburst, we used the
one-phase UniForc model, which describes the cumulative effect of forcing temperatures on bud
development during the ecodormancy phase (Chuine, Cour, and Rousseau 1999; Gauzere et al.
2017). We simulated damage due to late frosts (see details in Appendix 1) and considered that trees
were able to reflush after late frosts. We calculated NLF as the sum of late frost days experienced
after budburst initiation.

Second, we implemented a new option in CASTANEA to compute PLC following the formula of
Pammenter and Willigen (1998):

$$\text{PLC} = \frac{1}{1+e^{\text{slope}(\Psi_{\text{leaf}}-\Psi_{P50})}} \qquad \text{(Equation 5)},$$

291 with Ψ_{leaf} (MPa) the simulated midday leaf water potential, Ψ_{50} (MPa) the species-specific
292 potential below which 50% of the vessels are embolized, and *slope* a constant fixed to 50.

293 The leaf water potential Ψ_{leaf} was calculated as:

$$\Psi_{\text{leaf}}(t+1) = \Psi_{\text{soil}}(t+1) - \frac{\text{TR}}{3600} \times \text{R}_{\text{SoilToLeaves}} + \frac{\Psi_{\text{leaf}}(t)}{\Psi_{\text{soil}}(t+1) + \text{TR} \times \text{R}_{\text{SoilToLeaves}}}$$

$$\times\, e^{\frac{\text{deltaT}}{\text{R}_{\text{SoilToLeaves}} \times \text{Cap}_{\text{SoilToLeaves}}}}$$

(Equation 6),

where the soil water potential (Ψ_{soil} MPa) was calculated from daily SWCo (Campbell 1974). Ψ_{leaf}
was calculated hourly (deltaT = 3600s) based on the sap flow (TR in mmol.m^{-2}.leaf^{-1}) simulated
following the soil-to-leaves hydraulic pathway model of Loustau et al. (1990). We used a single
resistance ($R_{\text{SoilToleaves}}$ in MPa.m^2.s^1.kg^{-1}, following Campbell 1974) and a single capacitance
($Cap_{\text{SoilToleaves}}$ in kg.m^{-2}.MPa^{-1}) along the pathway. $R_{\text{SoilToleaves}}$ was assessed using midday and predawn
water potentials found in the literature.

We added a binary option in CASTANEA to simulate branch mortality and defoliation as a
function of PLC. If the PLC at year *n* was >0, the LAI at year *n* was reduced by the PLC value for trees
able to defoliate (option "Defoli-able"). Otherwise, PLC has no consequences on LAI.

Simulation design: The aim of the first simulations was to investigate whether response variables
simulated by CASTANEA correlated with patterns of observed mortality at population scale. We
simulated a population of 100 trees representing the variability in individual characteristics
observed in La Massane in terms of height-diameter allometry, DBH, leaf area index and budburst
phenology (Appendix 1). We also simulated a range of environmental conditions representing the
observed variability in SWCa and tree density. We also used this first simulation to validate
CASTANEA based on the correlation between simulated and observed ring width (Appendix 1).
For this first simulation, the values of focal output variables (PLC, NLF and BorR) were averaged
across the 100 trees. We also computed a cumulated vulnerability index (CVI) for each year *n*
combining the simulated PLC, NLF and BoR as follows:

317
$$\mathrm{CVI_n} = \left(\frac{\mathrm{PLC_n}}{\max(\mathrm{PLC})} + \frac{\mathrm{NLF_n}}{\max(\mathrm{NLF})} \right) - \frac{\mathrm{BoR_n}}{\max(\mathrm{BoR})} \quad \text{(Equation 7)},$$

Note that each term is weighted by its maximal value across all years, so that the contribution of
the three drivers to vulnerability is balanced. The possible range of CVI is [-1; 2].

The second simulation aimed at investigating the differences in physiological responses
between individuals with different characteristics. We simulated eight individuals corresponding to
a complete cross design with two size categories (5 and 40 cm in DBH), two budburst types (early
and normal), and two defoliation levels (option "Defoli-able" activated or not).

Statistical models of mortality to explore individual drivers of mortality
We used logistic regression models to investigate how tree characteristics affect the individual
probability of mortality ($P_{mortality}$). This approach is appropriate for a binary response variable and a
mixture of categorical and quantitative explanatory variables, which are not necessarily normally
distributed (Hosmer and Lemeshow 2000). We considered the four following complete logistic
regression models:

$P_{mortality}$ = [DEFw + Fungi + Budburst+ MBAI + (N_{stem} OR $Compet_{intra}$ OR $Compet_{intra+}$ OR
$Compet_{tot}$)] ×($DBH_{2002} + DBH_{2002}^2$) (Equation 8)

where the predictors defoliation (DEFw), growth (MBAI), size (DBH$_{2002}$) and competition (Nstem or

the Compet indices) were quantitative variables, and the presence of fungi (Fungi) and budburst

phenology (Budburst) were categorical variables. We included both a linear and quadratic effect of

DBH$_{2002}$ by specifying this effect as a polynomial of second degree. Interaction effects of the previous

predictors with this polynomial were included.

All variables were scaled before fitting the model. To select the best competition-related

variables, we first fitted the model described by equation 8 with each competition term successively

(Appendix 3). Then, we used the R package "MuMin" to compare and select the most parsimonious

models based on AIC. When two models had similar AIC (delta < 2) (Arnold 2010), the one with

fewer variables (most parsimonious) was selected. With our model we seek to understand factors

related to mortality and not to prediction, so we did not select a variable as recommended by Heinze

et al. (2018) and Lederer et al. (2019). Model validity was checked based on the leverage points (i.e.

points having a greater weight than expected by chance) with the Cook's distance (Cook distance <

0.5 indicate no leverage). We evaluated the goodness-of-fit with the Brier test score (Brier 1950).

We evaluated the sensitivity and specificity of the model using the receiver operating characteristic

(ROC) curve.

Collinearity resulting from correlations between predictor variables is expected to affect the

statistical significance of correlated variables by increasing type II errors (Schielzeth 2010). To

evaluate this risk, we first checked for correlation among predictors included in equation 9 (Figure

S5). We also computed the variation inflation factor (VIF) with the R package "car". A threshold of

GVIF$^{1/2df}$ < 2 is commonly accepted to show that variables are not excessively correlated and do not

make the model unstable.

We expressed the results in terms of odds ratios, also called relative risk, indicating the degree

of dependency between variables. For instance, the odds ratio for mortality as a function of

budburst characteristics (early vs normal) is:

$$\text{OddsRatio}_{\text{Early.vs.Normal}} = \frac{\text{Odds}_{\text{Early}}}{\text{Odds}_{\text{Normal}}} \qquad \text{(Equation 9)},$$

$$\text{With Odds}_{\text{Early}} = \frac{P_{\text{mortality}}(\text{Early})}{1 - P_{\text{mortality}}(\text{Early})} \quad \text{and Odds}_{\text{Normal}} = \frac{P_{\text{mortality}}(\text{Normal})}{1 - P_{\text{mortality}}(\text{Normall})}.$$

We computed odds ratios with "questionr" the R package (Barnier, Briatte, and Larmarange 2018).

The interactions were visualized with the package "jtools" (Long 2018).

Results

Temporal variations in population mortality rate in relation to physiological vulnerability simulated with CASTANEA.

We found a significant positive correlation between observed and simulated ring widths (p-value << 0.01). Although CASTANEA tended to overestimate growth at the beginning of the simulated period, and simulated a decreasing trend in growth over time that was not visible in the observed data. This is likely to be due to a bad estimation of population density prior to the monitoring period (see details in Appendix 1).

The total mortality rate between 2004 and 2016 was 23% (Figure 2; Table S1). After 2004 (2.6%), two peaks of high mortality were observed, in 2006-2007 (3.3% in 2006) and in 2010 (2.9%). The lowest mortality rate was observed in 2008 (0.8%).

CASTANEA simulated inter-annual variations in the percent loss of conductance (PLC): the mean PLC value varied among years, from 1% in 2004 and 2005 to 31% in 2006 (Figure 2a). The mean simulated biomass of carbon reserves (BoR) varied among years, from 51 $gC.m^{-2}$ in 2006 to 354 $gC.m^{-2}$ in 2011. Finally, the number of late frost days (NLF) varied among years, from 0.2 in 2016 to 7.56 days in 2010 (Figure 2c). The variation in the cumulative vulnerability index (CVI) integrated these different responses (Figure 2d), showed a peak in 2006 (drought), in 2010 (late frost) and in 2012 (combination of frost and drought).

None of the response variables simulated by CASTANEA (NLF, PLC, BoR) was significantly correlated to annual variation in mortality rate. The only significant correlation was observed between CVI and the annual mortality rate (r = 0.58, p-value = 0.04). Hence, inter-annual variations in CVI were a good predictor of the mortality rate, except in year 2007. Besides the stress-related variables simulated with CASTANEA, we also investigated the effects of climatic variables on inter-annual variations in mortality rates using a beta-regression model (Appendix 2). We considered drought indices computed from climatic series. This approach confirmed the effect of drought on mortality.

389 **Figure 2: Stress-related output variable simulated with CASTANEA from 2004 to 2016: (a)**
390 **percentage loss of conductance (PLC); (b) biomass of reserves in gC.kg^{-1} (BoR); (c) number of late**
391 **frost days (NLF);. (d) Cumulated vulnerability index (CVI) integrating a,b and c.** The black line is the
392 mean of simulation, and the grey area represents the inter-individual variation from the 1st to the
393 3rd quartile. The yellow line is the mortality rate observed in La Massane.
394

Inter-individual variation in the probability of mortality

All the variables listed in equation 8 were retained in the best logistic regression model and had a significant main effect on the probability of mortality (Table 2). This model explained 49% of the observed mortality and had both a good validity and goodness-of-fit (Appendix 3). However, the GVIF$^{1/2df}$ score for DBH is superior to 2 meaning that the collinearity with other variables is high, which is certainly due to the high number of interaction tested and not significant. Defoliation had the strongest linear effect on mortality: the relative probability of mortality increased by 1020 times for a one-unit increase in DEFw. Then, the relative probability of mortality was 2.28 higher for trees with earlier budburst as compared to others, and 1.88 higher for trees bearing fungi fructifications as compared to others. Among the competition-related variables, N_{stem} was selected as it was associated with the lowest AIC. The relative probability of mortality increased with increasing N_{stem}, and decreased with increasing MBAI. Regarding the effect of tree size, the polynomial of degree 2 corresponded to a U-shape and traduced a higher relative probability of mortality for both the smaller and the larger trees (In addition, this calibration is based on the mean of the individuals' 2, Appendix 3).

Table 2: Effects of tree characteristics on the individual tree's probability of mortality. Variables are defined in Table 1. Effects were estimated with a logistic regression model (equation 8). β is the maximum likelihood estimate, with its estimated error (SE), z-value, and associated p-value. OR is the odds ratio.

Variables	β	SE	z value	p-value	OR
DEFw	6.93	0.26	26.30	<0.0001	$1.02 \ 10^3$
Fungi	0.63	0.16	3.96	<0.0001	1.88
Budburst	0.82	0.17	4.71	<0.0001	2.26
MBAI	-0.45	0.08	-5.58	<0.0001	0.64
Nstem	0.13	0.04	3.44	<0.0001	1.14
DBH_{2002}	-9.19	8.16	-1.13	0.26	$1.02 \ 10^{-4}$
DBH_{2002}^2	21.32	9.32	2.29	0.02	$1.81 \ 10^9$
$DEFw:DBH_{2002}$	-49.71	15.51	-3.20	$5.91 \ 10^{-4}$	$2.57 \ 10^{-22}$
$DEFw:DBH_{2002}^2$	32.97	17.08	1.93	0.05	$2.08 \ 10^{14}$
$Fungi:DBH_{2002}$	-11.33	8.69	-1.30	0.19	$1.21 \ 10^{-5}$
$Fungi:DBH_{2002}^2$	-0.44	8.51	-0.05	0.96	0.65
$Budburst:DBH_{2002}$	0.27	10.41	0.03	0.98	1.31
$Budburst:DBH_{2002}^2$	-3.11	12.19	-0.26	0.80	$4.46 \ 10^{-2}$
$MBAI:DBH_{2002}$	2.89	3.37	0.86	0.39	18.10
$MBAI:DBH_{2002}^2$	-6.18	3.34	-1.85	0.06	$2.07 \ 10^{-3}$
$Nstem:DBH_{2002}$	1.32	4.12	0.32	0.75	3.76
$Nstem:DBH_{2002}^2$	0.00	4.88	0.00	1.00	1.00

Interaction effects between size and defoliation on mortality were significant: the relative
probability of mortality increased more rapidly with DEFw for small rather than larger trees, and at
an equal level of defoliation, the probability of mortality was always higher for smaller trees (Figure
3a, Table 2). Interaction effects between size and growth on mortality were also significant: the
decrease in the relative probability of mortality with increasing mean growth was evident mostly
for small trees (Figure 3b). These results were robust for the choice of the competition variable
(N_{stem} versus competition indices), for the choice of the size variable (height instead of diameter)
and for the consideration of size (DBH_{2002}) as a quantitative versus categorical variable (Appendix
3). Finally, we obtained convergent results with an alternative approach (survival analysis) which
account simultaneously for both levels of variability (individual and temporal) in our data set
(Appendix 4).

Individual variation in vulnerability simulated with CASTANEA
Simulations with CASTANEA showed that inter-individual differences in tree size, phenology,
and defoliation, together with the intensity of climatic stress, affected the physiological responses
to stress. The magnitude of the individual effects on tree vulnerability differed during a drought year
(2006), a frost year (2010) and a good year (2008, 2014 or 2016; Figure 4). The loss of conductance
was higher for trees with early budburst and for larger trees, but this effect was only evident in
drought years (Figure 4a). Moreover, during drought, the ability to defoliate decreased the risk of
cavitation (Figure 4a) but increased the risk of carbon starvation (Figure 4b). By contrast, phenology
only poorly affected the biomass of reserve (BoR): even during a frost year, trees with earlier
budburst did not reduce their BoR, due to their ability to reflush (Figure 4b). BoR was always lower
for large tree, even without stress. This was expected, because there is no explicit competition for
light in CASTANEA. Hence large trees and small trees have a relatively similar photosynthesis when
it is scaled by soil surface (large trees photosynthesise slightly more because they have a stronger
LAI). Large trees, on the other hand, have a larger living biomass and thus a higher level of
respiration, which leads to lower reserves.

Figure 3: Interaction effects in the logistic regression model at individual-level (a) between diameter (DBH_{2002}) and weighted defoliation (DEFw). (b) between DBH_{2002} and the mean growth in basal area (MBAI). Regression lines are plotted for three values of DBH_{2002}, corresponding to ± 1 standard deviation (10.7 cm) from the mean (22 cm). Confidence intervals at 95% are shown around each regression line.

Figure 4: Physiological proxies of vulnerability simulated for eight trees differing in size, defoliation and budburst phenology. We focus on three key years**:** 2006 (drought); 2008 (no stress); 2010 (late frosts). Colours indicate the DBH at the beginning of the simulation: 5 cm (light blue) versus 40 cm (dark green). Triangles (respectively round) indicate individuals with early (respectively "normal") budburst. Empty (respectively full) indicate individuals able (respectively not able) to defoliate.

470 **Table 3: Summary of the main effects of the studied variables on mortality.**
471 **NLF: number of late frost days; PLC= percentage of loss conductance; BoR: Biomass of reserve**
472

Temporal variations in the population mortality rate.		
Simulated variables	**Related climate stress**	**Process-based model**
NLF	late frost	Not directly correlated with the observed mortality rate
PLC	drought	Not directly correlated with the observed mortality rate
BoR	late frost and drought	Not directly correlated with the observed mortality rate
CVI	PLC + NLF - BoR	Positively correlated with the observed mortality rate
Inter-individual variations in tree's probability of mortality		
Variables	**Statistical model**	**Process-based model**
Crown defoliation	Associated to a strong increase in mortality, especially for small trees	Associated to a lower BoR but also to a lower PLC
Size (DBH)	The smallest and the largest trees had a higher mortality	Large trees had always a lower BoR, and a higher PLC in drought year
Growth (MBAI)	Fast-growing trees had a lower mortality (evident only for small trees)	Not tested
Budburst phenology	Tree with earlier budburst had a higher mortality	Tree with earlier budburst had a lower PLC in drought year
Competition (Nstem)	Increasing competition was associated to a higher mortality	Not tested
Presence of fungi	Tree with fungi had a higher mortality	Not tested

473
474

475 *Discussion*

476 By combining statistical and process-based models (Figure 1, Table 3), this study shed new

477 light on the inter-annual and inter-individual variability of mortality in a drought- and frost-prone

478 beech population. We showed that mortality in this marginal population is triggered by a

479 combination of climatic factors, and that the vulnerability to drought and frost is modulated by

480 several individual characteristics (defoliation, vegetative phenology, growth, size, competitors).

481 **The rate of mortality increased in response to drought and late frosts**

482 The annual mortality rates observed in this study ranged between 0.7 and 3.3% (mean value

483 = 2%). This is at the upper range of the few mortality estimates available for beech. Hülsmann et al.

484 (2016) reported annual mean rates of mortality of 1.4%, 0.7% and 1.5% in unmanaged forests of

485 Switzerland, Germany and Ukraine, with a maximum mortality rate of 2.2%. Archambeau et al.

(2019) estimated even lower mortality rates (mean annual value = 3.8 10^{-3}%, range = 3.7 10^{-3}% to
3.8 10^{-3}%) from European forest inventory data (including managed and unmanaged forests).
Overall, these mortality rates are low when compared to other tree species; for instance, according
to the French national forest inventory, the average mortality is 0.1% for beech against 0.3% on
average for other species and 0.4% for spruce or 0.2% for silver fir (IFN 2016). The relatively high
value observed here may result from the absence of management, combined with the population
location being at the dry, warm margin of species distribution (Figure S1), where most population
extinctions are expected in Europe (Thuiller et al. 2005). However, we cannot rule out that the size
threshold in inventories, which differ between studies, also affects these different mortality
estimates (e.g., a higher frequency of smaller trees increases the mortality rate).

We showed that inter-annual variations in the observed mortality rate at population-level
were significantly associated with variations in the cumulative vulnerability index (CVI) integrating
the number of late frost days (NLF), the percentage loss of conductance (PLC) and the biomass of
carbon reserves (BoR) simulated by CASTANEA. This association was not found when the three
response variables simulated by CASTANEA were considered separately, highlighting that patterns
of mortality in beech are driven by a combination of drought and late-frost stresses. In particular,
simulations showed that in 2010 (a year without drought), the high mortality rate coincided with an
extreme late frost event. This is consistent with the study of Vanoni et al (2016), which showed that
both drought and frost could contribute to beech mortality. Our results also support the emerging
consensus that mortality at dry, warm margins is not due either to carbon starvation or hydraulic
failure, but is rather the result of a balance of all these responses (e.g. McDowell et al. 2011; Sevanto
et al. 2014).

In future developments, the CVI could be refined in several ways. Its different components
could be weighted based on ecophysiological knowledge. The CVI could also benefit from taking into
account the temporal dynamics of mortality, such as the existence of positive or negative post-
effects across years. The low number of observations in this study compelled us to ignore these
lagged effects, which probably explains why the CVI failed to predict the high mortality observed in
2007. Indeed, the high mortality in 2007 is probably due to the lagged effect of the 2006 drought.
Such lags between the weakening of a tree and its final death were shown for beech in Vanoni
(2016) and silver fir in Davi & Cailleret (2017).

The vulnerability to drought and frost varied among individuals

516
The large number of trees individually monitored in this study provided us with an
exceptionally large sample size to test for the effects of factors modulating individual vulnerability
to climatic hazards (drought and late frost) and to biotic pressures (competition and the presence
of a fungus). Firstly, we found that a higher mean growth was associated with a lower probability of
mortality, as previously demonstrated (Cailleret and Davi 2011; Gao et al. 2018). This decrease in
mortality with increasing mean growth was evident mostly for small trees as already reported in
beech seedlings (Collet and Le Moguedec 2007) and other species (Kneeshaw et al. 2006; Lines,
Coomes, and Purves 2010), but not in adult beech trees to our knowledge.

Secondly, we found that increased defoliation was associated with increasing mortality. This
result was expected from previous studies (Dobbertin and Brang 2001, Carnicer et al. 2011),
although the consequences of defoliation are still being debated for beech. Senf et al. (2018)
showed that defoliation was associated with tree decline, while Bauch et al., (1996) and Pretzsch
(1996) found that the growth of highly defoliated beech trees did not decrease and could even
increase in some cases. Our simulations comparing trees able, or not able, to defoliate, shed light
on the multiple effects of defoliation on mortality. These simulations showed that defoliation indeed
decreased carbon reserves in good years but could also limit the loss of hydraulic conductance
during dry years. We also observed a significant interaction between defoliation and tree size on
mortality, showing that small trees were more vulnerable to mortality in response to defoliation
than large trees. However, we cannot rule out that this effect is due in part to the categorical
method used to survey defoliation, which does not take into account the percentage of crown loss.
Hence, defoliation may be biased with respect to size, such that small and defoliated trees will on
average have a higher proportion of canopy loss, and therefore be more impacted than large and
defoliated trees.

Thirdly, both statistical and process-based approaches found that trees with early budburst
were more prone to die. By contrast, Robson et al. (2013) showed that trees with early budburst
were not more vulnerable to mortality, but rather grew better, consistent with our simulations
where trees with early budburst accumulate more reserves during good years. This discrepancy may
be due to the location of our studied population at the rear-edge of beech distribution, where earlier
budburst dates are observed due to higher temperature and may expose trees to a higher risk of
late frost. It may be that the presence of trees with very early budburst in the studied population
makes it somewhat unusual, although similar cases have been observed elsewhere (Gaussen 1958;

Perci du Sert 1982). In CASTANEA simulations, the higher vulnerability of early trees resulted rather
from a higher risk of hydraulic failure than from a higher impact of late frosts. This is because trees
with early budburst have a longer vegetation season and they develop their canopies faster, which
also increases their water needs due to the increase of transpiration. Altogether, the relationships
between phenology and mortality deserve further investigation, especially since the spatio-
temporal variation of budburst patterns under climate change may produce complex spatio-
temporal patterns of stresses (Vanoni et al. 2016).

Regarding the effect of size, the results differed between the statistical approach, where large
trees died less than small ones, and the simulations, which predicted a greater vulnerability to
drought of large trees. There may be several explanations for this discrepancy. The first reason is
that CASTANEA simulates an average tree without explicit competition for light and water; hence
not accounting for the higher observed background mortality in small trees as compared to large
ones (Figure S7). In addition, CASTANEA also does not account for individual dominance status,
which can affect the current carbon balance of a tree and hence its capacity to mitigate stress. In
the studied population, large trees are more likely to be dominant, with better access to light
resources promoting carbon accumulation, as compared to small trees, which are more likely to be
suppressed. Another reason is that tree size may vary with environmental factors in the studied
population, such that large trees have a tendency to occur on better soils. Therefore, the size effect
observed through the statistical approach may reveal the confounding effect of spatial soil
heterogeneity, not taken into account in the PBM. A measurement of water availability at individual
tree level would be necessary to address this issue but was out of the scope of this study.

Combining statistical and process-based approaches to identify the causes of tree
vulnerability
These two approaches illustrate the classical compromise between a fine understanding of
physiological mechanisms driving mortality, with complex and expensive PBMs, versus high
precision in local mortality predictions, with statistical models requiring less data, but having a
weaker ability to generalize proximal causes. Most often, studies adopt either of the two
approaches, and generally statistical approaches prevail (Hülsmann et al. 2016; Seidl et al. 2011).
However, the two approaches are highly complementary, and combining them allows the
deciphering of the respective roles of the drivers and mechanisms underlying tree mortality and
understanding their variability among individuals or years (Hawkes 2000; O'Brien et al. 2017; Seidl

et al. 2011). The two approaches can be compared as this study does at the individual level, or they

can be combined as when we analysed the correlation between the observed mortality rate and

simulated stress response variables. An upper level of integration would be inverse modelling,

where observed mortality rates could be used to infer the physiological thresholds (e.g. in BoR, PLC

and NLF) likely to trigger mortality (Davi & Cailleret 2017; Cailleret et al., 2020).

This study also illustrated a classical difficulty in combining statistical and process-based

approaches, related to the difference between observed variables and PBM parameters. For

instance, the comparison of defoliated and non-defoliated trees does not have exactly the same

meaning when using CASTANEA and the statistical approach. In CASTANEA, we compared trees, able

versus unable to defoliate, while these average trees shared on average the same edaphic

conditions. In the statistical approach, we compared trees with different levels of defoliation, but

which also probably did not share the same edaphic and biotic conditions. Defoliation was thus also

likely to be an indicator of the fertility of the environment, such that on shallow soils, defoliation

was stronger and the probability of mortality increased. Hence, the correlation does not necessarily

involve a causal relationship between defoliation and mortality.

The major benefits of our approach combining different approaches (statistical, process-

based) at different scales (population, individual) is that it should ultimately allow to disentangle

ecological patterns observed at an upper scale (population, multi-year period), and get back to

patterns observed at a lower scale where processes operate (individual, year). This ability to

aggregate/disaggregate patterns is acknowledged as a powerful approach to understand apparent

contradictions between patterns observed at different scales (Clark et al. 2011). There are however

some limitations to the approaches we used here. First, none of them could fully account for the

non-independence of climatic effects on mortality between years. Indeed, the effect of climatic

variables at a given year may depend on other variables expressed in previous years. This was

observed in beech, where several drought years finally led to a growth decline (Jump et al. 2006;

Knutzen et al. 2017; Vanoni et al. 2016) or a modification in sap flow (Hesse et al. 2019). Moreover,

the processes driving mortality may change through time as the most sensitive individuals are

progressively eliminated, and/or the surviving trees become less and less sensitive (i.e. acclimation

Niinemets 2010). Finally, the statistical model at the individual level could not fully make use of the

repeated measurements of mortality over the years, partly because other individual variables were

measured only once over the study period (except defoliation). Survival analyses could

unfortunately not fully address this limitation (Appendix 4), and the development of a finely tuned
Bayesian approach was out of the scope of this study. Besides methodological improvements,
another extension to the present study would be to combine statistical and process-based
approaches at a larger spatial scale, among populations across climatic gradients. This would allow
the investigation of whether the respective drought and late frost sensitivity differ between the
rear, core and leading edge of species distribution, as suggested by Cavin and Jump (2017).

Data accessibility
617 The data set analysed in this preprint is available online under the zenodo repository
618 (https://doi.org/10.5281/zenodo.3519315). Raw data can be obtained from JG, JAM and CH.

Supplementary material
620 The process-based model CASTANEA is an open-source software available on capsis website:
621 http://capsis.cirad.fr/
622 Supplementary materials (Figures and Tables) for this preprint are available on bioRxiv (XXX).

Author Contributions
624 JAM, JG, CH and EM measured and mapped all the trees. CPC performed the wood core analyses.
625 CPC, FL and SOM designed and ran the statistical models. CPC and HD ran the PBM. CPC drafted the
626 manuscript, and all authors contributed to its improvement.

Acknowledgments
We are grateful to M. Cailleret, B. Fady, and N. Martin Saint Paul for discussions and comments on
a previous version of this manuscript. We thank E.Walker and F.Bonneu for statistical discussions
and advices, N. Mariotte for wood core sampling, and F. Guibal for their analyses. SOM and HD were
funded by the EU ERA-NET BiodivERsA projects TIPTREE (BiodivERsA2-2012-15) and the ANR project
MeCC (ANR-13-ADAP-0006). CP received funding from the European Union's Horizon 2020 research
and innovation programme under grant agreement No. 676876 (GenTree).

Conflict of interest disclosure
635 The authors of this preprint declare that they have no financial conflict of interest with the content
636 of this article. SOM is one of the PCIEcology recommenders.

References
638 Adams, Henry D. et al. 2017. "A Multi-Species Synthesis of Physiological Mechanisms in Drought-
639   Induced Tree Mortality." Nature Ecology and Evolution 1(9):1285–91.

Akaike, Hirotugu. 1987. "Factor Analysis and AIC." Pp. 371–86 in. Springer, New York, NY.

Allen, Craig D. et al. 2010. "A Global Overview of Drought and Heat-Induced Tree Mortality Reveals Emerging Climate Change Risks for Forests." Forest Ecology and Management 259(4):660–84.

Anderegg, W. R. L. et al. 2012. "From the Cover: The Roles of Hydraulic and Carbon Stress in a Widespread Climate-Induced Forest Die-Off." Proceedings of the National Academy of Sciences 109(1):233–37.

Anderegg, William R. L. 2015a. "Spatial and Temporal Variation in Plant Hydraulic Traits and Their Relevance for Climate Change Impacts on Vegetation." New Phytologist 205(3):1008–14.

Anderegg, William R. L. et al. 2015b. "Tree Mortality from Drought, Insects, and Their Interactions in a Changing Climate." New Phytologist 208(3):674–83.

Archambeau, Juliette et al. 2019. "Similar Patterns of Background Mortality across Europe Are Mostly Driven by Drought in European Beech and a Combination of Drought and Competition in Scots Pine." Agricultural and Forest Meteorology, 2020, vol. 280, p. 107772.

Arnold, Todd W. 2010. "Uninformative Parameters and Model Selection Using Akaike's Information Criterion." Journal of Wildlife Management 74(6):1175–78.

Augspurger, Carol K. 2009. "Spring 2007 Warmth and Frost: Phenology, Damage and Refoliation in a Temperate Deciduous Forest." Functional Ecology 23(6):1031–39.

Ball, J. Timothy, Ian E. Woodrow, and Joseph A. Berry. 1987. "A Model Predicting Stomatal Conductance and Its Contribution to the Control of Photosynthesis under Different Environmental Conditions." Pp. 221–24 in Progress in photosynthesis research. Springer.

Barnier, Julien, François Briatte, and Joseph Larmarange. 2018. "Questionr: Functions to Make Surveys Processing Easier."

Bauch, Josef. 1986. "Characteristics and Response of Wood in Declining Trees from Forests Affected by Pollution." IAWA Journal 7(4):269–76.

Beguería, Santiago and Sergio M. Vicente-Serrano. 2017. "SPEI: Calculation of the Standardised Precipitation-Evapotranspiration Index."

Benito Garzón, Marta et al. 2018. "The Legacy of Water Deficit on Populations Having Experienced Negative Hydraulic Safety Margin." Global Ecology and Biogeography 27(3):346–56.

Berzaghi, F., Wright, I. J., Kramer, K., Oddou-Muratorio, S., Bohn, F. J., Reyer, C. P., ... & Hartig, F. 2020. Towards a new generation of trait-flexible vegetation models. *Trends in Ecology & Evolution*, *35*(3), 191-205.

Bigler, Christof and Harald Bugmann. 2018. "Climate-Induced Shifts in Leaf Unfolding and Frost Risk of European Trees and Shrubs." Scientific Reports 8(1):1–10.

Brando, Paulo Monteiro et al. 2014. "Abrupt Increases in Amazonian Tree Mortality Due to Drought-Fire Interactions." Proceedings of the National Academy of Sciences of the United States of America 111(17):6347–52.

Bréda, Nathalie, Roland Huc, André Granier, and Erwin Dreyer. 2006. "Temperate Forest Trees and Stands under Severe Drought : A Review of Ecophysiological Responses , Adaptation Processes and Long-Term Consequences." Annals of Forest Science 63(6):625–44.

Brier, Glenn W. 1950. "Verification of Forecasts Expressed in Terms of Probability." Monthey Weather Review 78(1):1–3.

De Caceres, Miquel, Nicolas Martin-StPaul, Marco Turco, Antoine Cabon, and Victor Granda. 2018. "Estimating Daily Meteorological Data and Downscaling Climate Models over Landscapes." Environmental Modelling and Software 186–96.

Cailleret, Maxime et al. 2017. "A Synthesis of Radial Growth Patterns Preceding Tree Mortality." Global Change Biology 23(4):1675–90.

Cailleret, Maxime and Hendrik Davi. 2011. "Effects of Climate on Diameter Growth of Co-Occurring Fagus Sylvatica and Abies Alba along an Altitudinal Gradient." Trees 25(2):265–76.

Cailleret, M., Bircher, N., Hartig, F., Hülsmann, L., & Bugmann, H. (2020). Bayesian calibration of a growth dependent tree mortality model to simulate the dynamics of European temperate - forests. Ecological Applications, 30(1), e02021

Campbell, Gaylon S. 1974. "A Simple Method for Determining Unsaturated Conductivity from Moisture Retention Data." Soil Science 117(6):311–14.

Carnicer, Jofre et al. 2011. "Widespread Crown Condition Decline, Food Web Disruption, and Amplified Tree Mortality with Increased Climate Change-Type Drought." Proceedings of the National Academy of Sciences 108(4):1474–78.

Cavin, Liam and Alistair S. Jump. 2017. "Highest Drought Sensitivity and Lowest Resistance to Growth Suppression Are Found in the Range Core of the Tree Fagus Sylvatica L. Not the Equatorial Range Edge." Global Change Biology 23(1):362–79.

Cheaib, Alissar et al. 2012. "Climate Change Impacts on Tree Ranges: Model Intercomparison Facilitates Understanding and Quantification of Uncertainty." Ecology Letters 15(6):533–44.

Choat, Brendan et al. 2018. "Triggers of Tree Mortality under Drought." Nature 558(7711):531–39.

Chuine, Isabelle, P. Cour, and D. D. Rousseau. 1999. "Selecting Models to Predict the Timing of Flowering of Temperate Trees: Implications for Tree Phenology Modelling." Plant, Cell and Environment 22(1):1–13.

Clark, James S. et al. 2011. "Individual-Scale Variation, Species-Scale Differences: Inference Needed to Understand Diversity." Ecology Letters 14(12):1273–87.

Collet, Catherine and Gilles Le Moguedec. 2007. "Individual Seedling Mortality as a Function of Size, Growth and Competition in Naturally Regenerated Beech Seedlings." Forestry 80(4):359–70.

Cowan, I. R. and G. D. Farquhar. 1977. "Stomatal Function in Relation to Leaf Metabolism and Environment." Symposia of the Society for Experimental Biology 31:471–505.

Cribari-Neto, Francisco and Achim Zeileis. 2010. "Beta Regression in R." Journal of Statistical Software 34(2).

Davi, H. et al. 2005. "Modelling Carbon and Water Cycles in a Beech Forest. Part II.: Validation of the Main Processes from Organ to Stand Scale." Ecological Modelling 185(2–4):387–405.

Davi, H., C. Barbaroux, C. Francois, and E. Dufrêne. 2009. "The Fundamental Role of Reserves and Hydraulic Constraints in Predicting LAI and Carbon Allocation in Forests." Agricultural and Forest Meteorology 149(2):349–61.

Davi, Hendrik and Maxime Cailleret. 2017. "Assessing Drought-Driven Mortality Trees with Physiological Process-Based Models." Agricultural and Forest Meteorology 232:279–90.

Dittmar, Christoph, Wolfgang Zech, and Wolfram Elling. 2003. "Growth Variations of Common Beech (Fagus Sylvatica L.) under Different Climatic and Environmental Conditions in Europe—a Dendroecological Study." Forest Ecology and Management 173(1–3):63–78.

Dobbertin, Matthias and Peter Brang. 2001. "Crown Defoliation Improves Tree Mortality Models." Forest Ecology and Management 141(3):271–84.

Dufrêne et al. 2005. "Modelling Carbon and Water Cycles in a Beech Forest Part I : Model Description and Uncertainty Analysis on Modelled NEE." Ecological Modelling 185:407–36.

Durand-Gillmann, Marion, Maxime Cailleret, Thomas Boivin, Louis Michel Nageleisen, and Hendrik Davi. 2014. "Individual Vulnerability Factors of Silver Fir (Abies Alba Mill.) to Parasitism by Two Contrasting Biotic Agents: Mistletoe (Viscum Album L. Ssp. Abietis) and Bark Beetles (Coleoptera: Curculionidae: Scolytinae) during a Decline Process." Annals of Forest Science 71(6):659–73.

Farquhar, G. D., S. von Caemmerer, and J. A. Berry. 1980. "A Biochemical Model of Photosynthetic CO_2 Assimilation in Leaves of C3 Species." Planta 149(1):78–90.

Feng, Xue et al. 2018. "The Ecohydrological Context of Drought and Classification of Plant Responses." Ecology Letters, November 1, 1723–36.

Gao, Shan et al. 2018. "Dynamic Responses of Tree-Ring Growth to Multiple Dimensions of Drought." Global Change Biology 24(11):5380–90.

García-Plazaola, José Ignacio, Raquel Esteban, Koldobika Hormaetxe, Beatriz Fernández-Marín, and José María Becerril. 2008. "Photoprotective Responses of Mediterranean and Atlantic Trees to the Extreme Heat-Wave of Summer 2003 in Southwestern Europe." Trees 22(3):385–92.

Gaussen, H. 1958. "Le Hêtre Aux Pyrénées Espagnoles." Pp. 185–91 in Actas del tercer congreso internacional de estudios pirenaicos, Gerona.

Gauzere, J. et al. 2017. "Integrating Interactive Effects of Chilling and Photoperiod in Phenological Process-Based Models. A Case Study with Two European Tree Species: Fagus Sylvatica and Quercus Petraea." Agricultural and Forest Meteorology 244–245:9–20.

Gillner, Sten, Nadja Rüger, Andreas Roloff, and Uta Berger. 2013. "Low Relative Growth Rates Predict Future Mortality of Common Beech (Fagus Sylvatica L.)." Forest Ecology and Management 302:372–78.

Granier, A., P. Biron, and D. Lemoine. 2000. "Water Balance, Transpiration and Canopy Conductance in Two Beech Stands." Agricultural and Forest Meteorology 100(4):291–308.

Greenwood, Sarah et al. 2017. "Tree Mortality across Biomes Is Promoted by Drought Intensity, Lower Wood Density and Higher Specific Leaf Area" edited by J. Chave. ECOLOGY LETTERS 20(4):539–53.

Heinze, Georg. Christine Wallisch et al., Variable selection – A review and recommendations for the practicing statistician. Biometrical Journal 60 (2018), S. 431-449

Hawkes, Corinna. 2000. "Woody Plant Mortality Algorithms: Description, Problems and Progress." Ecological Modelling 126(2–3):225–48.

Hesse, Benjamin D., Michael Goisser, Henrik Hartmann, and Thorsten E. E. Grams. 2019. "Repeated Summer Drought Delays Sugar Export from the Leaf and Impairs Phloem Transport in Mature Beech" edited by M. Dannoura. Tree Physiology 39(2):192–200.

Hijmans, Robert J., Steven Phillips, John Leathwick, Jane Elith, and Maintainer Robert J. Hijmans. 2017. "Package 'Dismo.'" Circles 9(1):1–68.

Hosmer, David W. and Stanley Lemeshow. 2000. Applied Logistic Regression.

Hülsmann, Lisa et al. 2016. "Does One Model Fit All? Patterns of Beech Mortality in Natural Forests of Three European Regions." Ecological Applications 26(8):2463–77.

Hülsmann, Lisa, Harald Bugmann, and Peter Brang. 2017. "How to Predict Tree Death from Inventory Data – Lessons from a Systematic Assessment of European Tree Mortality Models - SUPP." Canadian Journal of Forest Research (April):cjfr-2016-0224.

Hülsmann, Lisa, Harald Bugmann, Maxime Cailleret, and Peter Brang. 2018. "How to Kill a Tree: Empirical Mortality Models for 18 Species and Their Performance in a Dynamic Forest Model." Ecological Applications 28(2):522–40.

IGN. 2016. La Mortalité. France. https://inventaire-forestier.ign.fr/IMG/pdf/2018_mortalite.pdf

Jump, Alistair S., Jenny M. Hunt, and Josep Peñuelas. 2006. "Rapid Climate Change-Related Growth Decline at the Southern Range Edge of Fagus Sylvatica." Global Change Biology 12(11):2163–74.

Kneeshaw, Daniel D., Richard K. Kobe, K. David Coates, and Christian Messier. 2006. "Sapling Size Influences Shade Tolerance Ranking among Southern Boreal Tree Species." Journal of Ecology 94(2):471–80.

Knutzen, Florian, Choimaa Dulamsuren, Ina Christin Meier, and Christoph Leuschner. 2017.
"Recent Climate Warming-Related Growth Decline Impairs European Beech in the Center of
Its Distribution Range." Ecosystems 20(8):1494–1511.
Kramer, Koen et al. 2010. "Modelling Exploration of the Future of European Beech (Fagus Sylvatica
L.) under Climate Change-Range, Abundance, Genetic Diversity and Adaptive Response."
Forest Ecology and Management 259(11):2213–22.
Lebourgeois, F., N. Bréda, E. Ulrich, and A. Granier. 2005. "Climate-Tree-Growth Relationships of
European Beech (Fagus Sylvatica L.) in the French Permanent Plot Network (RENECOFOR)."
Trees 19:385–401.
Lederer, D. J., S. C. Bell et al., Control of Confounding and Reporting of Results in Causal Inference
Studies. Guidance for Authors from Editors of Respiratory, Sleep, and Critical Care Journals.
Ann Am Thorac Soc 16 (2019), S. 22-28.
Lenz, Armando, Günter Hoch, Yann Vitasse, and Christian Körner. 2013. "European Deciduous
Trees Exhibit Similar Safety Margins against Damage by Spring Freeze Events along
Elevational Gradients." New Phytologist 200(4):1166–75.
Lines, Emily R., David A. Coomes, and Drew W. Purves. 2010. "Influences of Forest Structure,
Climate and Species Composition on Tree Mortality across the Eastern US" edited by A.
Hector. PLoS ONE 5(10):e13212.
Long, Jacob A. 2018. "Jtools: Analysis and Presentation of Social Scientific Data."
Lorenz, Martin and Georg Becher. 2012. Forest Condition in Europe.
Loustau, D., A. Granier, F. El Hadj Moussa, M. Sartore, and M. Guedon. 1990. "Evolution
Saisonnière Du Flux de Sève Dans Un Peuplement de Pins Maritimes." Annales Des Sciences
Forestières 47(6):599–618.
Van Mantgem, Phillip J. et al. 2009. "Widespread Increase of Tree Mortality Rates in the Western
United States." Science 323(5913):521–24.
Maraun, Mark, Jörg-Alfred Salamon, Katja Schneider, Matthias Schaefer, and Stefan Scheu. 2003.
"Oribatid Mite and Collembolan Diversity, Density and Community Structure in a Moder
Beech Forest (Fagus Sylvatica): Effects of Mechanical Perturbations." Soil Biology and
Biochemistry 35(10):1387–94.
Martin, George L. and Alan R. Ek. 1984. "A Comparison of Competition Measures and Growth
Models for Predicting Plantation Red Pine Diameter and Height Growth." Forest Science
30(3):731–43.
McDowell, Nate G. et al. 2013. "Evaluating Theories of Drought-Induced Vegetation Mortality
Using a Multimodel-Experiment Framework." New Phytologist 200(2):304–21.
McDowell, Nate G. et al. 2011. "The Interdependence of Mechanisms Underlying Climate-Driven
Vegetation Mortality." Trends in Ecology & Evolution 26(10):523–32.
Meir, Patrick, Maurizio Mencuccini, and Roderick C. Dewar. 2015. "Drought-Related Tree
Mortality: Addressing the Gaps in Understanding and Prediction." New Phytologist
207(1):1443–47.
Menzel, Annette, Raimund Helm, and Christian Zang. 2015. "Patterns of Late Spring Frost Leaf
Damage and Recovery in a European Beech (Fagus Sylvatica L.) Stand in South-Eastern
Germany Based on Repeated Digital Photographs." Frontiers in Plant Science 6:110.
Monserud, Robert A. 1976. "Simulation of Forest Tree Mortality." Forest Science 22(4):438–44.
Monteith, J. L. 1965. "Evaporation and Environment. The State and Movement of Water in Living
Organisms. Symposium of the Society of Experimental Biology, Vol. 19 (Pp. 205-234)."
Mueller, Rebecca C. et al. 2005. "Differential Tree Mortality in Response to Severe Drought:
Evidence for Long-Term Vegetation Shifts." Journal of Ecology 93(6):1085–93.

Niinemets, Ülo. 2010. "Responses of Forest Trees to Single and Multiple Environmental Stresses
from Seedlings to Mature Plants: Past Stress History, Stress Interactions, Tolerance and
Acclimation." Forest Ecology and Management 260(10):1623–39.
Nourtier, Marie et al. 2014. "Transpiration of Silver Fir (Abies Alba Mill.) during and after Drought
in Relation to Soil Properties in a Mediterranean Mountain Area." Annals of Forest Science
71(6):683–95.
O'Brien, Michael J. et al. 2017. "A Synthesis of Tree Functional Traits Related to Drought-Induced
Mortality in Forests across Climatic Zones. Journal of Applied Ecology 54(6):1669–86.
O'Brien, Michael J., Sebastian Leuzinger, Christopher D. Philipson, John Tay, and Andy Hector.
2014. "Drought Survival of Tropical Tree Seedlings Enhanced by Non-Structural Carbohydrate
Levels." Nature Climate Change 4(8):710–14.
Pammenter NW and Vander Willigen C. 1998. "A Mathematical and Statistical Analysis of the
Curves Illustrating Vulnerability of Xylem to Cavitation." Tree Physiology 18(Equation 1):589–
593.
Penuelas, Josep and Martí Boada. 2003. "A Global Change-Induced Biome Shift in the Montseny
Mountains (NE Spain)." Global Change Biology 9(2):131–40.
Perci du Sert, Th. 1982. Relations Entre La Phenologie et La Morphologie Du Hêtre Dans Le Massif
Des Albères.
Pretzsch, Hans. 1996. "Growth Trends of Forests in Southern Germany." Pp. 107–31 in Growth
Trends in European Forests. Berlin, Heidelberg: Springer Berlin Heidelberg.
Robson, T. Matthew, Erwin Rasztovits, Pedro J. Aphalo, Ricardo Alia, and Ismael Aranda. 2013.
"Flushing Phenology and Fitness of European Beech (Fagus Sylvatica L.) Provenances from a
Trial in La Rioja, Spain, Segregate According to Their Climate of Origin." Agricultural and
Forest Meteorology 180:76–85.
Ryan, Michael G. 1991. "Effects of Climate Change on Plant Respiration." Ecological Applications
1(2):157–67.
Sala, A. and J. D. Tenhunen. 1996. "Simulations of Canopy Net Photosynthesis and Transpiration in
Quercus Ilex L. under the Influence of Seasonal Drought." Agricultural and Forest
Meteorology 78(3–4):203–22.
Schielzeth, Holger. 2010. "Simple Means to Improve the Interpretability of Regression
Coefficients." Methods in Ecology and Evolution 1(2):103–13.
Seidl, Rupert et al. 2011. "Modelling Natural Disturbances in Forest Ecosystems: A Review."
Ecological Modelling 222(4):903–24.
Senf, Cornelius et al. 2018. "Canopy Mortality Has Doubled in Europe's Temperate Forests over
the Last Three Decades." Nature Communications 9(1):4978.
Sevanto, Sanna, Nate G. Mcdowell, L. Turin Dickman, Robert Pangle, and William T. Pockman.
2014. "How Do Trees Die? A Test of the Hydraulic Failure and Carbon Starvation
Hypotheses." Plant, Cell and Environment 37(1):153–61.
Stadt, Kenneth J. et al. 2007. "Evaluation of Competition and Light Estimation Indices for
Predicting Diameter Growth in Mature Boreal Mixed Forests." Annals of Forest Science
64(64):477–90.
Thuiller, Wilfried, Sandra Lavorel, M. B. Araujo, Martin T. Sykes, and I. Colin Prentice. 2005.
"Climate Change Threats to Plant Diversity in Europe." Proceedings of the National Academy
of Sciences 102(23):8245–50.
Tyree, M. T. and J. S. Sperry. 1989. "Vulnerability of Xylem to Cavitation and Embolism." Annual
Review of Plant Physiology and Plant Molecular Biology 40(1):19–36.

Vanoni, Marco, Harald Bugmann, Magdalena Nötzli, and Christof Bigler. 2016. "Drought and Frost
Contribute to Abrupt Growth Decreases before Tree Mortality in Nine Temperate Tree
Species." Forest Ecology and Management Journal 382:51–63.
Vidal, Jean-Philippe, Eric Martin, Laurent Franchistéguy, Martine Baillon, and Jean-Michel
Soubeyroux. 2010. "A 50-Year High-Resolution Atmospheric Reanalysis over France with the
Safran System." International Journal of Climatology 30(11):1627–44.
Vitasse, Yann et al. 2009. "Leaf Phenology Sensitivity to Temperature in European Trees: Do
within-Species Populations Exhibit Similar Responses?" Agricultural and Forest Meteorology
149(5):735–44.
De Vries, F. W. T. Pennin., A. H. M. Brunsting, and H. H. Van Laar. 1974. "Products, Requirements
and Efficiency of Biosynthesis a Quantitative Approach." Journal of Theoretical Biology
45(2):339–77.
886

Appendices

Four supplementary appendices are available on bioRxiv (BIORXIV/2019/645747):

Appendix 1: CASTANEA model, calibration and simulation design

Appendix 2: Beta-regression model for the temporal variations in the rate of mortality at population
level

Appendix 3: Logistic regression models for the probability of mortality at tree-level

Appendix 4: Survival analysis the probability of mortality at tree- and year-levels