Scalable and cost-effective ribonuclease-based rRNA depletion for bacterial transcriptomics

Yiming Huang[1,2†]**, Ravi U Sheth**[1,2†]**, Andrew Kaufman**[1]**, Harris H Wang**[1,3,*]

[1] Department of Systems Biology, Columbia University, New York, NY, USA.
[2] Integrated Program in Cellular, Molecular, and Biomedical Studies, Columbia University, New York, NY, USA.
[3] Department of Pathology and Cell Biology, Columbia University, New York, NY, USA
[†] These authors contributed equally to this work.
* Correspondence to: hw2429@columbia.edu

Abstract

Bacterial RNA sequencing (RNA-seq) is a powerful approach for quantitatively delineating the global transcriptional profiles of microbes in order to gain deeper understanding of their physiology and function. Cost-effective bacterial RNA-seq requires efficient physical removal of ribosomal RNA (rRNA), which otherwise dominates transcriptomic reads. However, current methods to effectively deplete rRNA of diverse non-model bacterial species are lacking. Here, we describe a new probe and ribonuclease based strategy for bacterial rRNA removal. We implemented the method using either chemically synthesized oligonucleotides or amplicon-based single-stranded DNA probes and validated the technique on three novel gut microbiota isolates from three distinct phyla. We further showed that different probe sets can be used on closely related species. We provide a detailed methods protocol and an online tool to generate custom probe libraries. This approach lays the groundwork for large-scale and cost-effective bacterial transcriptomics studies.

Keywords: RNA-seq, transcriptomics, rRNA depletion, microbiome

Background

Bacterial RNA-seq provides global and in-depth transcriptional profiling of a given species or community that can yield quantitative and mechanistic insights into microbial function and ecology, and holds exciting potential for routine use in microbiology studies and large-scale screening applications[1-7]. In bacteria, the 16S and 23S ribosomal RNA (rRNA) are the most abundant RNA molecules in the cell and typically account for more than 90% of the total RNA[8]. These rRNA therefore need to be experimentally removed from total RNA during library preparation steps to ensure cost-effective bacterial RNA-seq with sufficient coverage of coding sequences of the transcriptome. Mammalian RNA-seq has a similar challenge of rRNA removal, but solutions are more tractable since 1) most mammalian mRNAs are polyadenylated and can thus be easily selected or captured[9, 10], or 2) the same species (e.g. human, with the same rRNA sequences) are generally profiled such that sequence-specific depletion methods can be streamlined[11-14]. In contrast, bacterial mRNAs do not have a poly(A) sequence, which makes enrichment strategies infeasible, and 16S and 23S rRNA sequences are diverse between species, which makes universal depletion approaches much more challenging. A number of commercially available rRNA depletion methods exist[8, 15-18] (Supplementary Table 1), but they typically rely on targeting short conserved rRNA regions or specific rRNA modifications such that only full length rRNA can be removed. Recent methods for RNA barcoding have enabled dramatically reduced cost in RNA-seq by pooling multiple samples together at the total RNA stage and performing downstream library preparation reactions in a pooled format[19]. However, since these protocols generate fragmented RNA (including rRNA), typical rRNA removal kits cannot be used. While sequence-independent rRNA depletion methods such as those that rely on DSN nuclease have been described[20], they suffer from low efficiency and sequence-specific or GC biases.

The Ribo-Zero kit (Illumina, Inc.) is a popular commercial rRNA depletion kit that is compatible with fragmented RNA and has been used effectively on some (e.g. [19]) but not all bacteria. Unfortunately, this kit was recently discontinued by the manufacturer and no commercial replacement is available. Motivated by the lack of a reliable alternative, we set out to develop a new bacterial rRNA depletion method with the following criteria: 1) high efficiency 16S/23S rRNA depletion, 2) scalable to diverse bacterial species, 3) compatibility with fragmented or barcoded RNA samples, and 4) low cost per reaction. Correspondingly, we developed a ribonuclease (RNase) H based method for bacterial rRNA depletion described here. We show that our approach efficiently depletes bacterial rRNA with minimal off-target effects to the transcriptome and demonstrate the use of the method on bacterial species from three different phyla. Moreover, we show that species-specific depletion probes can be generated by chemical synthesis or from PCR amplicons (with different cost considerations) and that probe-sets can be utilized on species with similar 16S/23S sequences.

Results

A RNase H based rRNA depletion strategy

RNase H based methods are widely used for mammalian rRNA depletion[11, 13]. We thus aimed to implement and optimize a similar strategy for bacterial rRNA depletion. Given the diversity of bacterial 16S and 23S sequences, our method requires a user-customizable approach to design new probe sequences. Probes for a species of interest can be generated via two strategies (**Fig. 1a**). First, in a chemical synthesis approach, probes can be designed from 16S and 23S sequences by splitting sequences into ~50nt single-stranded (ss) DNA oligonucleotides that are synthesized de novo. Second, in a PCR amplicon-based approach, universal primers (the forward primer is 5'-phosphorylated) can be used to amplify 16S and 23S amplicons from genomic DNA and the resulting product can be converted to ssDNA by lambda exonuclease, which

80 selectively digests 5'-phosphorylated strands of dsDNA(21). The generated ssDNA probes (oligo
81 or amplicon) are subsequently hybridized to total RNA. The ssDNA:ssRNA duplexes are
82 selectively degraded with RNase H, and the remaining ssDNA probes in the reaction are then
83 degraded by DNase I, thus finally yielding enriched mRNAs (**Fig. 1b**).
84

85
86
Fig. 1: Workflow for bacterial RNase H based rRNA depletion. a) Probes used for depletion can be
either designed and chemically synthesized from known rRNA sequences (oligo-based) or generated by
PCR from genomic DNA with 5'-phosphorylated forward primers and subsequent lambda exonuclease
digestion (amplicon-based). b) Probes are then hybridized to total RNA and the rRNA bound by the ssDNA
probes is degraded by RNase H. Finally, all remaining probes are degraded by DNase I, resulting in
enriched mRNAs.
93

Optimizing RNase H reaction parameters

94
We first optimized the conditions for probe hybridization and RNase H degradation of
bacterial rRNA. To test reaction conditions, we took total RNA isolated from a novel human gut
microbial isolate, *Bacteroides dorei*, and split the same RNA across multiple conditions. 16S and
23S sequences were determined from a draft genome of the isolate and a probe library composed
of 87 distinct ~50nt oligonucleotides that tiled the 16S and 23S sequences was designed and
chemically synthesized (Methods, Supplementary Table 2). Total RNA from this strain was
subjected to fragmentation and adapter ligation (with sample barcoding) as per the standard
RNAtag-seq protocol (Methods)(19). We replaced the typical Ribo-Zero rRNA depletion step with
our RNase H depletion reaction, and the resulting RNA after depletion was subjected to reverse
transcription, second adapter ligation, PCR amplification, and Illumina sequencing (to a low
coverage for initial optimization, Methods). The resulting sequencing reads were analyzed using
a standard RNA-seq analysis pipeline and the proportion of reads originating from rRNA of total
mapped reads was determined (Methods).
We tested two different RNase H enzymes, a traditional RNase H (New England Biolabs)
and a thermostable Hybridase™ RNase H (Lucigen), with their corresponding buffer and
temperature conditions at a fixed probe-to-RNA of 1:1 based on previous mammalian RNase H
depletion protocols(13) (Methods). In addition, we tested various reaction times (Fig. 2a). For both
RNase H enzymes, the proportion of rRNA reads decreased after depletion, with longer
incubation times yielding improved depletion efficiency for Hybridase RNase H but not NEB
RNase H. To understand the contribution of probe-to-RNA ratio to overall depletion efficiency, we
fixed the reaction time to 30 minutes and tested different probe-to-RNA ratios (Fig. 2b). For both

RNase H enzymes, increasing probe-to-RNA ratios improved depletion efficiency, yielding low
levels of reads mapped to rRNA (NEB 5:1 probe: 12.90%, Hybridase 5:1 probe: 6.82%); overall
the Hybridase RNase H enzyme outperformed the NEB RNase H enzyme at all ratios. Based on
these results, we adopted the following depletion enzyme and conditions for subsequent studies:
Hybridase RNase H with its corresponding reaction buffers, a probe-to-RNA ratio of 5:1, and a
reaction duration of 30 minutes. While even higher probe-to-RNA ratios or longer reaction times
led to better performance, we chose to use the minimal conditions need to yield satisfactory
depletion in order to minimize potential off-target degradation.
124

125
126
127 ***Fig. 2**: **Optimization of RNase H reaction conditions**. **a)** Proportion of rRNA-aligning reads of total*
128 *mapped reads (% rRNA reads), for un-depleted and RNase H depleted samples for two RNase H enzymes*
129 *and various reaction times with probe-to-RNA fixed to 1:1, or **b)** various probe-to-RNA ratios with reaction*
130 *time fixed to 30 minutes. The same RNA sample isolated from Bacteroides dorei was split for rRNA*
131 *depletion across different reaction conditions.*
132

Validating RNase H-based rRNA depletion

Having determined an optimal set of reaction conditions, we next set out to validate the
rRNA depletion performance of the method across diverse bacterial species and assess any
potential technical artifacts the method may introduce. We obtained a new RNA sample for the *B.*
dorei strain, and also obtained RNA from two additional distinct novel gut microbiota isolates for
testing, *Collinsella aerofaciens* and *Dorea longicatena.* Collectively these three isolates represent
three distinct phyla (Bacteroidetes, Actinobacteria, Firmicutes, respectively). We performed
library preparation and rRNA depletion as before, utilizing the optimal reaction conditions as well
as a chemically synthesized oligonucleotide probe library designed for each strain based on draft
genomes from Illumina assemblies (Supplementary Table 2). The resulting RNA libraries as well
as the same RNA with no rRNA depletion were sequenced at a high coverage (Method). In all
three strains, our method significantly depleted rRNA efficiently bringing 94.0±2.0% of reads
mapping to rRNA down to 26.5±6.2% of reads mapping to rRNA, which represents a 13.0±4.1
fold enrichment of non-rRNA reads across the three strains (Fig. 3a, Supplementary Figure 1).
We also compared the performance of our method to the Ribo-Zero bacterial kit, which we applied
to the same RNA samples. In 2 of 3 cases, our technique yielded higher performance than Ribo-
Zero (10.0, 17.6 and 11.5-fold enrichment of non-rRNA reads by our RNase H method versus 3.6,
9.6 and 15.3-fold enrichment by Ribo-Zero, respectively). The overall efficiency was uniform
across the three strains, highlighting the consistent performance of strain-specific probe sets. In
addition, for all non-rRNA reads, depleted samples showed consistent overall mappability
(79.2±12.5% for depleted samples, 75.8±8.1% for un-depleted samples, Supplementary Figure
2) and a higher proportion of reads mapped to CDSs (93.6±1.4% for depleted samples, 83.8±2.9%
for un-depleted samples, Supplementary Figure 2), indicating our method minimally affects overall

library quality. Together these results demonstrate that the RNase H based rRNA depletion
method has favorable performance across three diverse strains.
Next, to determine if rRNA depletion may have altered the transcriptome due to off-target
depletion, we compared the depleted expression values to un-depleted expression values for all
annotated CDSs. Importantly, since both samples were generated from the same RNA sample,
any difference reflects bias due to the depletion method (in addition to technical bias associated
with sample preparation). We mapped at least 600K reads to annotated CDSs for each sample,
to ensure high coverage for expression quantification (Supplementary Table 3). Overall, the
depleted samples showed high consistency with un-depleted samples in terms of transcriptome
correlation (R^2 = 0.948, 0.973, 0.944, respectively, Fig. 3b) and expression distribution (quantile-
quantile plot, points lie on a straight line if there is no global shift in expression profile) (Fig.
3c).Together, these results validate that the RNase H based rRNA depletion method minimally
perturbs the underlying transcriptome.
169

170
171
Fig. 3: *Application of RNase H based rRNA depletion with oligos to three diverse gut microbiota*
species. *a) Proportion of rRNA-aligning reads of total mapped reads (% rRNA reads), for un-depleted,*
Ribo-Zero depleted, and RNase H depleted samples under optimized reaction condition across three
microbiota species from different phyla. The same RNA samples used for Ribo-Zero depletion were pooled
*together to yield the RNA used for RNase H reaction. **b, c)** Consistency in transcriptome between rRNA*
*depleted samples and un-depleted samples in terms of expression correlation **(b)** and expression*
*distribution **(c)**. TPM indicates transcripts per million for each CDS. **c)** Points lie on a straight line in Quantile-*
Quantile (Q-Q) plots if there is no global shift in the distribution of expression profile between depleted
samples and un-depleted samples.
181
Amplicon-based probe generation and validation

The chemically synthesized oligo probe strategy provides efficient rRNA depletion but
requires synthesis of a specific probe library. While the cost of the probe library can be amortized
over many reactions and the total cost is minimal compared to purchasing a commercial kit
(Supplementary Table 1), for small scale studies it may be desirable to generate probes on a
smaller scale for a lower overall cost. We therefore set out to validate a PCR amplicon and ssDNA
generation strategy to generate probes from a microbial DNA template derived from universal
primers. In theory, this strategy could also allow for the generation of complex probe sets from
templates of multiple different species. Using universal primers where the forward primer is 5'-
phosphorylated (Supplementary Table 4), we amplified the 16S and 23S regions from *B. dorei*
and subjected the amplicons to lambda exonuclease digestion, which selectively digests the 5'-
phosphorylated strands(21) to yield complementary ssDNA probes (amplicon) (Methods). We
then repeated the testing of rRNA depletion efficiency on the same *B. dorei* RNA as in Fig. 3,
utilizing the same reaction conditions and a 5:1 or a 10:1 amplicon probe-to-RNA ratio (Fig. 4a).
The 5:1 ratio amplicon probes showed similar depletion efficiency (16.1% of reads mapping to
rRNA) as the oligo probes, while the 10:1 ratio showed increased depletion efficiency (4.0%
mapping to rRNA), which demonstrate that amplicon-based probes can also effeciently deplete
rRNA. As before, we compared the expression values of depleted samples with the amplicon
probes to expression values from the un-depleted samples generated from the same RNA (Fig.
4b-c). The amplicon-based depleted samples showed high consistency with un-depleted samples
(R^2 = 0.936). Together, these results demonstrate that in addition to chemically synthesized
oligonucleotides, ssDNA probes generated from PCR amplicons can yield efficient rRNA
depletion with minimal perturbation of the transcriptome.
205

208   ***Fig. 4****: **RNase H based rRNA depletion with amplicons**. a) Proportion of rRNA-aligning reads of total*
209   *mapped reads (% rRNA reads), for un-depleted, Ribo-Zero depleted, and RNase H depleted samples with*
210   *chemically synthesized oligonucleotides (oligo) or amplicon-based ssDNA (amplicon) probes. **b, c)** Scatter*
211   *plot **(b)** and Quantile-Quantile (Q-Q) plot **(c)** for depleted sample using amplicon probes with 5:1 probe-to-*
212   *RNA ratio and un-depleted sample.*
213

Applicability of probes to closely related species

The cost of generating new probe libraries for novel species may impose practical
experimental constraints for studies that involve transcriptomic profiling of many different
microbial strains. We hypothesized that probe libraries developed for a specific strain may be
applied also to closely related species with similar 16S and 23S sequences. A single probe library

that can be scalably re-applied to many potential strains could thus significantly reduce the overall
cost. We therefore tested the *B. dorei* oligo probe pool on two closely related *Bacteroides* species,
Bacteroides uniformis ATCC 8492 and *Bacteroides vulgatus* ATCC 8482 with the same library
preparation and optimal reaction conditions as before. In addition, we tested the same probe
library on distantly related species, the *C. aerofaciens* and *D. longicatena* strains tested before.
The *B. dorei* probe library was able to efficiently deplete *B. uniformis* and *B. vulgatus* rRNA (Fig.
5a) with minimal perturbation to their transcriptomes (R^2 = 0.984, 0.983, respectively,
Supplementary Figure 3) as before. As expected, the probes did not deplete rRNA from the
distantly related organisms (*C. aerofaciens* and *D. longicatena*).
To understand the correlation between probe-to-target sequence similarity and rRNA
depletion efficiency, we quantified the percent of mismatches in target rRNA sequences
compared to *B. dorei* rRNA sequences. Species with less than 10% mismatches overall could be
efficiently depleted (Fig. 5b). We performed a similar analysis for each individual oligo probe, and
the correlation of probe mismatches to depletion efficiency of the cognate target sequence
suggests that probes with less than 25% mismatches can be used to efficiently deplete local
regions of rRNA (Supplementary Figure 4). Taken together, these results demonstrate scalability
of our designed probe libraries for reuse in closely related species.
236

237
238
Fig. 5*: **Oligo probes can be applied to closely related species**. **a)** Proportion of rRNA-aligning reads of*
total mapped reads (% rRNA reads), for un-depleted and RNase H depleted samples using a oligo probe
library designed for B. dorei across closely related species (B. uniformis and B. vulgatus) and distantly
related species (C. aerofaciens and D. longicatena). ***b)*** *Scatter plot between probe-to-target sequence*
similarity and fold enrichment of non-rRNA reads for five species. Probe-to-target sequence similarity was
calculated as the average of percentages of base with mismatches in 16S and 23S rRNA alignments.
245
246 **Discussion**
247
To facilitate use of this rRNA depletion technique by the research community, we have
included a detailed step-by-step protocol for the method (Supplementary Methods). In addition,
to streamline design of oligonucleotide pools, we have created a simple tool to generate probe
libraries from target 16S and 23S sequences (https://github.com/hym0405/RNaseH_depletion).
The tool can also perform simple calculations of probe identity to various different 16S and 23S
sequences to evaluate the ability of pools to be applied to different sequences.
We developed two separate strategies for generating probe libraries that perform with
similar efficiency. While chemically synthesizing probes requires less hand-on time, it constitutes
a high up-front cost (~$450/probe set) but generally yields enough material for a large number of

reactions (>12,000 reactions). Alternatively, in-house amplicon-based ssDNA probes do not require a significant up-front cost, but do require more time (~8 hour protocol end-to-end) to generate and quality control. Based purely on cost (Supplementary Figure 5), synthesized probes become cost-effective after ~90 reactions compared to amplicon probes. Importantly however, we note that the cost per reaction in both cases ($4.98 for amplicon probes, ~$0.04 for oligo probes and $12.90 for other reagents per depletion reaction, Supplementary Table 5) is significantly lower than commercially available kits (e.g. ~$80 per reaction for the discontinued Ribo-Zero rRNA Removal Kits, Supplementary Table 1).

Beyond the application to single-species bacterial rRNA depletion shown here, the method may have utility for depletion of unwanted sequences for various transcriptomics applications. For example, our approach could be applied to non-model organisms for rRNA depletion. In addition, extension of this strategy to more complex probe pools could enable high efficiency rRNA depletion for metatranscriptomics experiments. Chemically synthesized oligo pools could be designed for specific environments based on the expected genera present, or custom probe pools could be generated on a per-sample basis by pooled amplification of 16S and 23S rRNA from a complex microbial community.

Finally, we note that for specific applications, further optimization to reaction times, probe-to-RNA ratios, and RNase H reaction temperature could even further improve efficiency. We chose to use conservative reaction conditions that still achieved significant depletion to avoid any potential off-target effects. However, future work could further validate and improve upon different reaction conditions in the context of specific samples to yield accurate transcriptome data.

Conclusion

We have developed a new RNase H based ribosomal RNA depletion method for bacterial transcriptomics that is compatible with barcoded or fragmented RNA libraries. We validated that the method has high efficiency for rRNA depletion (~13-fold enrichment of non-rRNA reads) and minimal off-target perturbation to the transcriptome on three diverse bacterial species. We showed that the method is compatible with chemically synthesized oligo probes or amplicon-based ssDNA probes generated from genomic DNA. Finally, we demonstrated that probe pools can be successfully applied to closely related species. This technique is expected to have broad utility in routine bacterial transcriptomics experiments and emerging large-scale bacterial RNA-seq studies.

Methods

Bacterial strains and culture condition. *Bacteroides dorei*, *Collinsella aerofaciens* and *Dorea longicatena* are novel isolates derived from a fecal sample obtained from a healthy human adult subject, obtained during the course of unrelated experiments. This work was approved and conducted under Columbia University Medical Center Institutional Review Board protocol AAAR0753, and written informed consent was obtained from the subject. *Bacteroides uniformis* strain ATCC 8492 and *Bacteroides vulgatus* strain ATCC 8482 were obtained from American Type Culture Collection, Manassas, VA, USA. All bacterial strains were grown in Gifu Anaerobic Medium Broth, Modified [HyServe 05433] under anaerobic conditions (5% H_2, 10% CO_2, 85% N_2) in a Coy Labaratory Products anaerobic chamber.

Bacterial RNA and DNA extraction. Bacterial RNA was obtained utilizing the RNAsnap method(22). Briefly, 1mL of saturated culture was centrifuged and the cell pellet was stored at -80°C. The pellet was then suspended in 500µL RNAsnap mix (95% formamide, 18mM EDTA, 0.025% SDS and 1% B-mercaptoethanol), lysed using ~200µL of 0.1mm Zirconia Silica beads [Biospec 11079101Z] using a BioSpec mini bead beater [Biospec 1001] for 2 minutes in a deep

well 96-well plate [Axygen P-DW-20-C]. The plate was centrifuged and the clarified lysate was
cleaned up using a Zymo ZR-96 RNA Clean & Concentrator kit [Zymo R1080] per manufacturer's
instructions. Bacterial DNA was obtained using our standard automated DNA extraction pipeline
based on a scaled down version of the Qiagen MagAttract PowerMicrobiome DNA/RNA Kit
[Qiagen 27500-4-EP], detailed fully in (23).

Genome sequencing and assembly of novel isolates. To obtain draft genome references for
novel isolates, paired-end libraries were constructed using a low-volume Nextera sequencing
protocol(24) and sequenced on Illumina Nextseq 500/550 platform to obtain about 6 million
paired-end reads with a read length of 2x75bp for each novel isolate. The reads were processed
using Cutadapt v2.1(25) with following parameters "-a file:[Nextera_adapter.fa] -A
file:[Nextera_adapter.fa] --minimum-length 25:25 -u 15 -u -5 -U 15 -U -5 -q 15 --max-n 0 --pair-
filter=any" to remove first 15 bases, last 5 bases and adapters. Nextera adapter sequences used
for trimming are provided in the Supplementary Table 6. To further improve the performance of
genome assembling, PacBio long-read sequencing was performed for each isolate by SNPsaurus.
PacBio long reads and remaining Illumina reads were assembled using Unicycler(26) in hybrid
mode with default parameters to generate draft assembly for each isolate. The draft assemblies
were further assessed using CheckM(27) to ensure their high quality (completeness >99% and
contamination <1%).

Draft genome assembly annotation. Draft genomes for novel isolates (*Bacteroides dorei*,
Collinsella aerofaciens and *Dorea longicatena)* were *de novo* assembled as described above,
and genome assemblies for *Bacteroides uniformis* strain ATCC 8492 and *Bacteroides vulgatus*
strain ATCC 8482 were downloaded from NCBI (RefSeq assembly accession:
GCF_000154205.1 and GCF_000012825.1). Genome annotation was performed using Prokka
v1.13.3(28) with following parameters "--rnammer --rfam" for all 5 bacterial strains, and sequences
of predicted 16S and 23S rRNAs were extracted from genome using in-house script for
downstream rRNA reads alignment (only one 16S and one 23S rRNA were used for strains with
multiple rRNA annotations since their sequences are almost identical).

RNA-seq library preparation and sequencing. The RNA-seq library was constructed following
standard RNAtag-seq protocol with minor modifications(19). Briefly, 4µg of total RNA was
fragmented using 2X FastAP buffer [ThermoFisher EF0651], depleted of DNA and
dephosphorylated using TURBO™ DNase [ThermoFisher AM2239] and FastAP [ThermoFisher
EF0651], and then fragmented RNA after 2X SeraPure SPRI beads cleanup(29, 30) was ligated
to barcoded first adapter [NEB M0437M]. Ligation product was further subjected to size selection
and purification using Zymo RNA Clean and Concentrator-5 [Zymo R1015]. Barcoded RNA
samples were previously subjected to rRNA depletion using Illumina Ribo-Zero rRNA Removal
Kit (Bacterial) per the manufacturer's instructions, and we replaced the typical Ribo-Zero rRNA
depletion step with our RNase H depletion reaction. The same RNA samples were pooled
together to yield the RNA used for the RNase H reaction. Barcoded RNA was firstly quantified
using Qubit RNA HS Assay Kit [ThermoFisher Q32855] per manufacturer's instructions, and
100ng of barcoded RNA was subjected to RNase H based rRNA depletion for condition
optimization experiment while 500ng of barcoded RNA was subjected to RNase H based rRNA
depletion to validate depletion performance across diverse bacterial species and with different
types of probes. To generate library with no rRNA depletion as control, 16ng of barcoded RNA
(for optimization experiment) and 50ng RNA of barcoded RNA (for validation experiment) was
directly diluted into 14µL and subjected to downstream library preparation. rRNA depleted RNA
was subjected to reverse transcription [ThermoFisher 18090010], degradation of the RNA [NEB
M0297S] and ligation of second adapter [NEB M0437M]. Ligation product was further amplified
with primers that contain Illumina P5 and P7 adapters and sample indexes, and the PCR reactions
were stopped during exponential amplification (typically ~15 cycles). The PCR products were

subjected to gel electrophoresis on E-Gel™ EX Agarose Gels, 2% [ThermoFisher G402002] and expected DNA bands (around 300bp-600bp) were excised from gel and extracted by Wizard™ SV Gel and PCR Cleanup System [Promega A9282] following the manufacturer's instructions to remove PCR primers and adapter dimers from sequencing library. Resulting libraries were sequenced to a low coverage in optimization experiment and a high coverage in validation experiment (Supplementary Table 7). Sequences of all adapters and primers used in library preparation are provided in the Supplementary Table 8; we note that in our implementation of RNAtag-seq we have swapped the orientation of R1 and R2 primers such that RNA barcodes are sequenced in read 2 and not read 1.

For the condition optimization experiment (100ng of RNA as input for RNase H reaction) except the un-depleted control, libraries were sequenced on Illumina MiSeq platform (reagent kits: v2 50-cycles, single-end mode) at 12 pM loading concentration with 10% PhiX spike-in (PhiX Control V3 [Illumina FC-110-3001]) following the manufacturer's instructions.

For the depletion performance validation experiment (500ng of RNA as input for RNase H reaction) and the un-depleted control in optimization experiment, libraries were sequenced on Illumina NextSeq 500/550 platform (reagent kits: High Output v2 75-cycles, single-end mode) at 1.8 pM loading concentration with 2.5% PhiX spike-in (PhiX Control V3 [Illumina FC-110-3001]) following the manufacturer's instructions.

ssDNA probe generation. Probes for a species of interest can be generated via two strategies: chemically synthesized oligonucleotide (oligo) or amplicon-based ssDNA (amplicon).

For oligo probes, we developed a tool to automatically design oligo probe libraries for given 16S and 23S sequences: ribosomal RNA sequence was spilt into multiple ~50nt DNA oligos covering the entire length of its reverse complement. The oligo probe libraries for 5 strains were chemically synthesized from Integrated DNA Technologies Inc. in plate format with normal desalting purification, and equimolarly pooled together to generate oligo probe mixes used in this study.

For amplicon probes, universal primers in which the forward primer is 5'-phosphorylated (8F/1541R for 16S amplicon and 10F/2756R for 23S amplicon, Supplementary Table 4) were used to amplify 16S and 23S amplicons from bacterial genomic DNA, and the resulting PCR product was subjected to gel electrophoresis (E-Gel™ EX Agarose Gels, 2% [ThermoFisher G402002]), and gel extraction (Wizard™ SV Gel and PCR Cleanup System [Promega A9282]) following the manufacturer's instructions, then 16S and 23S amplicons were amplified again using the same primers with gel extract as template. Amplicon products were purified using 1X SeraPure SPRI beads(29, 30) and then subjected to a lambda exonuclease [ThermoFisher EN0562] digestion, which selectively digests the 5'-phosphorylated strands resulting in amplicon ssDNA probes complementary to ribosomal RNAs. Finally, 16S amplicon probe and 23S amplicon probe were mixed in equimolar amounts to generate amplicon probe mixes used in this study. A detailed step-by-step protocol for the method is included in Supplementary Methods.

RNase H reaction optimization. To optimize RNase H reaction conditions, we added oligo probe mix to 100ng of fragmented RNA after first adapter ligation with various probe-to-RNA ratios (1:1, 5:1 or 10:1), the mixture was incubated in hybridization buffer (200mM NaCl, 100mM Tris-HCl pH7.5) in a final volume of 5µL at 95°C for 2 minutes, the temperature was slowly ramped down (-0.1°C/sec) to 45°C (for Hybridase RNase H) or 22°C (for NEB RNase H), and the mixture was incubated at 45°C (for Hybridase RNase H) or 22°C (for NEB RNase H) for additional 5 minutes. After probe hybridization, we added 5µL preheated Hybridase RNase H reaction mix (10U Hybridase Thermostable RNase H [Lucigen H39500], 0.5µmol Tris-HCl pH 7.5, 1µmol NaCl and 0.2µmol MgCl$_2$) or 5µL NEB RNase H reaction mix (10U NEB RNase H [NEB M0297S] and 1uL 10X NEB RNase H reaction buffer) to the RNA-probe mixture and incubated the mixture at 45°C

(for Hybridase RNase H) or 37°C (for NEB RNase H) for various depletion times (10 minutes, 30
minutes and 60 minutes). After RNase H digestion, 15µL DNase I reaction mix (4U TURBO
DNase [ThermoFisher AM2239] and 2.5µL 10X TURBO DNase Buffer) was added, and the
mixture was incubated at 37°C for 30 minutes to degrade ssDNA probes. RNA after depletion
was purified with 2X SeraPure SPRI beads(29, 30) and eluted into 14µL for downstream library
construction.

RNase H based ribosomal RNA depletion. Based on the results of optimization experiments,
we selected reaction conditions of Hybridase RNase H enzyme, 5:1 probe-to-RNA ratio and 30
minutes reaction duration. To validate the performance of this method across diverse bacterial
species and with different types of probes, we used 500ng of RNA as input and scaled the
hybridization reaction to 15µL. We added 2500ng ssDNA probe mix (oligo or amplicon) to 500ng
of fragmented RNA after first adapter ligation, the mixture was incubated in hybridization buffer
(200mM NaCl, 100mM Tris-HCl pH7.5) in a final volume of 15µL at 95°C for 2 minutes, the
temperature was slowly ramped down (-0.1°C/sec) to 45°C, and the mixture was incubated at
45°C for additional 5 minutes. After probe hybridization, we added 5µL preheated Hybridase
RNase H reaction mix that contains 15U Hybridase Thermostable RNase H [Lucigen H39500],
0.5µmol Tris-HCl pH 7.5, 1µmol NaCl and 0.4µmol MgCl2 to the RNA-probe mixture and
incubated the mixture at 45°C for 30 minutes. After RNase H digestion, 30µL DNase I reaction
mix (6U TURBO DNase [ThermoFisher AM2239] and 5µL 10X TURBO DNase Buffer) was added,
and the mixture was incubated at 37°C for 30 minutes to degrade ssDNA probes. RNA after
depletion was purified with 2X SeraPure SPRI beads(29, 30) and eluted into 14µL for downstream
library construction. A detailed step-by-step protocol for the method is included in Supplementary
Methods.

RNA-seq data analysis. Raw sequencing reads were first processed using Cutadapt v2.1(25)
with following parameters "-a file:[RNAtagSeq_adapter.fa] -u 5 --minimum-length 20 --max-n 0 -q
20" to remove first 5 bases and adapters before alignment. RNAtag-seq adapter sequences used
for trimming are provided in the Supplementary Table 6. To remove reads originated from PhiX
spike-in, we aligned remaining reads to the Coliphage phi-X174 genome (downloaded from NCBI,
sequence accession: NC_001422.1) using Bowtie2 v2.3.4 in single-end and "--very-sensitive"
mode, and only unmapped reads were used for downstream alignment. We then aligned
remaining reads to 16S and 23S rRNAs of corresponding strains using Bowtie2 v2.3.4(31) in
single-end mode to estimate the proportion of reads originating from rRNA. The reads not mapped
to rRNA were also outputted by "--un" option in Bowtie2 and further aligned to corresponding
genome reference using Bowtie v1.2.2(32) in single-end mode with following parameters "-a -m
1 --best --strata". All alignments were subsequently converted to BAM format, sorted and indexed
using SAMtools v1.9(33) for further analysis.

To evaluate the performance of rRNA depletion, following read-level metrics were calculated
(reads with multiple alignments suppressed in BAM file were included in the following calculations):
proportion of reads originating from rRNA, i.e. rRNA reads (%), was defined as: number of reads
mapped to rRNA / number of total mapped reads; and fold enrichment of non-rRNA reads was
defined as: fold change of (1 - proportion of reads originating from rRNA).

To evaluate the transcriptomic consistency between depleted samples and un-depleted samples,
the number of reads uniquely mapped to each CDS (reads with multiple alignments were
excluded) was calculated using featureCounts v1.6.2(34) without restraint on strandness (-s 0),
we then quantified the expression level of each CDS by transcripts per million (TPM) using in-
house script. We used $\log_{10}(TPM + 0.1)$ as input to calculate the Pearson correlation, Spearman
correlation and R-squared for gene-level correlation measurement, and visualized the quantiles

of TPM distributions in quantile-quantile (Q-Q) plots to evaluate whether rRNA depletion could
cause global shift in transcriptome (CDSs with TPM > 2000 were removed in Q-Q plot).

Probe-to-target sequence similarity analysis. To test the scalability of probes to other species,
we quantified probe-to-target sequence similarity by the number of mismatches in their sequence
alignment. Firstly, 16S and 23S rRNA sequences of *B. dorei* and target species were aligned
respectively using MUSCLE v3.8.31(35) with default setting, then we counted the number of
mismatches (including gaps) in their alignment, and the percent of mismatches in rRNAs was
calculated as: number of mismatches in rRNA alignment / total length of rRNA alignment
(including gaps). In addition, for each individual probe in *B. dorei* oligo library, we counted the
number of mismatches in the ~50nt tile of rRNA alignment that this probe covers (including gaps),
and percent of mismatches in each probe was calculated as: the number of mismatches in the
tile (including gaps) / total length of the tile (including gaps). In addition, depletion efficiency, i.e.
rRNA fold change, of that local rRNA sequence was calculated as the average fold change of
reads counts (normalized to non-rRNA mapped reads) for each base in the tile.

Code availability. Code utilized in this study can be accessed at
https://github.com/hym0405/RNaseH_depletion

467
468 <u>**Declarations**</u>
469
470 **Data availability**
471
472 Raw sequencing data is available at NCBI SRA under PRJNA542677.
473
474 **Competing interests**
475
476 H.H.W. , Y.H. and R.U.S. are inventors on a provisional patent application filed by the Trustees
477 of Columbia University in the City of New York regarding this work.
478
479 **Funding**
480
481 H.H.W. acknowledges funding from the NIH (1R01AI132403, 1R01DK118044), Burroughs
482 Welcome Fund (PATH 1016691), Bill & Melinda Gates Foundation (INV-000609), and the
483 Schaefer Research Scholars Program for this work. R.U.S. is supported by a Fannie and John
484 Hertz Foundation Fellowship and a NSF Graduate Research Fellowship (DGE-1644869).
485
486 **Authors' contributions**
487
488 Y.H. and R.U.S. designed the method, performed experiments and analyzed all data. A.K.
489 assisted with experiments. H.H.W. supervised the project. All authors wrote and approved the
490 manuscript.
491
492 **Acknowledgements**
493
494 We thank Nathan Johns and Peter Sims for helpful discussions.
495
496 <u>**References**</u>
497

1.   Hibberd MC, Wu M, Rodionov DA, Li X, Cheng J, Griffin NW, et al. The effects of
micronutrient deficiencies on bacterial species from the human gut microbiota. Sci Transl
Med. 2017;9(390).

2.   Dey N, Wagner VE, Blanton LV, Cheng J, Fontana L, Haque R, et al. Regulators of gut
motility revealed by a gnotobiotic model of diet-microbiome interactions related to travel.
Cell. 2015;163(1):95-107.

3.   Charbonneau MR, O'Donnell D, Blanton LV, Totten SM, Davis JC, Barratt MJ, et al.
Sialylated Milk Oligosaccharides Promote Microbiota-Dependent Growth in Models of
Infant Undernutrition. Cell. 2016;164(5):859-71.

4.   Lawley B, Sims IM, Tannock GW. Whole-transcriptome shotgun sequencing (RNA-seq)
screen reveals upregulation of cellobiose and motility operons of Lactobacillus ruminis
L5 during growth on tetrasaccharides derived from barley beta-glucan. Appl Environ
Microbiol. 2013;79(18):5661-9.

5.   Creecy JP, Conway T. Quantitative bacterial transcriptomics with RNA-seq. Curr Opin
Microbiol. 2015;23:133-40.

6.   Bush EC, Ray F, Alvarez MJ, Realubit R, Li H, Karan C, et al. PLATE-Seq for genome-
wide regulatory network analysis of high-throughput screens. Nat Commun.
2017;8(1):105.

7.   Garrido D, Ruiz-Moyano S, Lemay DG, Sela DA, German JB, Mills DA. Comparative
transcriptomics reveals key differences in the response to milk oligosaccharides of infant
gut-associated bifidobacteria. Sci Rep. 2015;5:13517.

8.   Giannoukos G, Ciulla DM, Huang K, Haas BJ, Izard J, Levin JZ, et al. Efficient and
robust RNA-seq process for cultured bacteria and complex community transcriptomes.
Genome Biol. 2012;13(3):R23.

9.   Mortazavi A, Williams BA, McCue K, Schaeffer L, Wold B. Mapping and quantifying
mammalian transcriptomes by RNA-Seq. Nat Methods. 2008;5(7):621-8.

10.   Marioni JC, Mason CE, Mane SM, Stephens M, Gilad Y. RNA-seq: an assessment of
technical reproducibility and comparison with gene expression arrays. Genome Res.
2008;18(9):1509-17.

11.   Morlan JD, Qu K, Sinicropi DV. Selective depletion of rRNA enables whole transcriptome
profiling of archival fixed tissue. PLoS One. 2012;7(8):e42882.

12.   Herbert ZT, Kershner JP, Butty VL, Thimmapuram J, Choudhari S, Alekseyev YO, et al.
Cross-site comparison of ribosomal depletion kits for Illumina RNAseq library
construction. BMC Genomics. 2018;19(1):199.

13.   Adiconis X, Borges-Rivera D, Satija R, DeLuca DS, Busby MA, Berlin AM, et al.
Comparative analysis of RNA sequencing methods for degraded or low-input samples.
Nat Methods. 2013;10(7):623-9.

14.   Sooknanan R, Pease J, Doyle K. Novel methods for rRNA removal and directional,
ligation-free RNA-seq library preparation. Nat Methods. 2010;7(10):I-Ii.

15.   He SM, Wurtzel O, Singh K, Froula JL, Yilmaz S, Tringe SG, et al. Validation of two
ribosomal RNA removal methods for microbial metatranscriptomics. Nat Methods.
2010;7(10):807-U58.

16.   Petrova OE, Garcia-Alcalde F, Zampaloni C, Sauer K. Comparative evaluation of rRNA
depletion procedures for the improved analysis of bacterial biofilm and mixed pathogen
culture transcriptomes. Sci Rep. 2017;7:41114.

17.   He S. Ribosomal RNA Removal Methods for Microbial Transcriptomics. Metagenomics
for Microbiology: Elsevier; 2015. p. 39-53.

18.   Dunman PM, Murphy E, Haney S, Palacios D, Tucker-Kellogg G, Wu S, et al.
Transcription profiling-based identification of Staphylococcus aureus genes regulated by
the agr and/or sarA loci. J Bacteriol. 2001;183(24):7341-53.

19.   Shishkin AA, Giannoukos G, Kucukural A, Ciulla D, Busby M, Surka C, et al.
Simultaneous generation of many RNA-seq libraries in a single reaction. Nat Methods.
2015;12(4):323-U60.
20.   Yi H, Cho YJ, Won S, Lee JE, Jin Yu H, Kim S, et al. Duplex-specific nuclease efficiently
removes rRNA for prokaryotic RNA-seq. Nucleic Acids Res. 2011;39(20):e140.
21.   Higuchi RG, Ochman H. Production of single-stranded DNA templates by exonuclease
digestion following the polymerase chain reaction. Nucleic Acids Res. 1989;17(14):5865.
22.   Stead MB, Agrawal A, Bowden KE, Nasir R, Mohanty BK, Meagher RB, et al. RNAsnap:
a rapid, quantitative and inexpensive, method for isolating total RNA from bacteria.
Nucleic Acids Res. 2012;40(20):e156.
23.   Ji BWS, R. U.; Dixit, P. D.; Wang, H. H.; Vitkup, D. . Quantifying spatiotemporal
dynamics and noise in absolute microbiota abundances using replicate sampling.
bioRxiv. 2019;310649.
24.   Baym M, Kryazhimskiy S, Lieberman TD, Chung H, Desai MM, Kishony R. Inexpensive
multiplexed library preparation for megabase-sized genomes. PLoS One.
2015;10(5):e0128036.
25.   Martin M. Cutadapt removes adapter sequences from high-throughput sequencing
reads. EMBnet journal. 2011;17(1):10-2.
26.   Wick RR, Judd LM, Gorrie CL, Holt KE. Unicycler: Resolving bacterial genome
assemblies from short and long sequencing reads. PLoS Comput Biol.
2017;13(6):e1005595.
27.   Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. CheckM: assessing
the quality of microbial genomes recovered from isolates, single cells, and
metagenomes. Genome Res. 2015;25(7):1043-55.
28.   Seemann T. Prokka: rapid prokaryotic genome annotation. Bioinformatics.
2014;30(14):2068-9.
29.   Rohland N, Reich D. Cost-effective, high-throughput DNA sequencing libraries for
multiplexed target capture. Genome Res. 2012;22(5):939-46.
30.   Faircloth BG, T. Homemade AMPure XP beads Ethan-omics: Ethan-omics; 2011 [
31.   Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods.
2012;9(4):357-9.
32.   Langmead B, Trapnell C, Pop M, Salzberg SL. Ultrafast and memory-efficient alignment
of short DNA sequences to the human genome. Genome Biol. 2009;10(3):R25.
33.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence
Alignment/Map format and SAMtools. Bioinformatics. 2009;25(16):2078-9.
34.   Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. Bioinformatics. 2014;30(7):923-30.
35.   Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high
throughput. Nucleic Acids Res. 2004;32(5):1792-7.
587