# NEUROGENETIC BASIS OF PERSONALITY

Sofie L. Valk[1,2*]; Felix Hoffstaedter[1,2]; Julia Camilleri[1,2]; Peter Kochunov[3]; B.T. Thomas Yeo[4,5,6,7]; Simon B. Eickhoff[1,2]

[1] *Institute of Neuroscience and Medicine (INM-7: Brain and Behaviour), Research Centre Jülich, 52425 Jülich, Germany*
[2] *Institute of Systems Neuroscience, Heinrich Heine University Düsseldorf, 40225 Düsseldorf, Germany*
[3] *Maryland Psychiatric Research Center, University of Maryland School of Medicine, Baltimore, Maryland, USA.*
[4] *Department of Electrical and Computer Engineering, Clinical Imaging Research Centre, Institute for Health and Memory Networks Program, National University of Singapore, Singapore, Singapore.*
[5] *Athinoula A. Martinos Center for Biomedical Imaging, Massachusetts General Hospital, Charlestown, MA, USA.*
[6] *Centre for Cognitive Neuroscience, Duke-NUS Medical School, Singapore, Singapore.*
[7] *NUS Graduate School for Integrative Sciences and Engineering, National University of Singapore, Singapore, Singapore.*

*Corresponding author:
S.L.Valk: svalk@fz-juelich.de

49 **Summary**
50

Personality traits are influenced by genetic and environmental factors and are related to

mental health. Previous neuroimaging studies have reported associations between personality

and brain structure, while at the same time, studies on imaging genetics have repeatedly

demonstrated heritability of brain structure. However, to date, it is unknown whether the

association between personality and brain macrostructure can be explained by shared genetic

factors. Here we report a large-scale twin study (Human Connectome Project), finding genetic

correlation between personality traits and brain structure. Agreeableness was genetically

correlated with the posterior-mid temporal, midcingulate, and dorsolateral frontal cortices,

while extraversion showed a genetic relation to temporoparietal junction. Quantitative

functional decoding indicated regions associated with personality are implicated in (socio-

)cognitive and language processing. Last, assessment of genetic correlation of personality and

adaptive behavior suggests overlapping neurogenetic substrates of personality and adaptive

function and problems. Together, these observations provide first evidence of a direct genetic

link between complex behavioral traits such as personality, local brain morphometry, and

function of the respective brain regions.

67
68
69

Personality traits quantify individual differences in goals, cognition, and emotion, that lead to

behavior[1]. It is a broad construct conceptualized in specific traits to quantify variability in

human behaviors. The most prominent factor approach of personality is the Five-factor

personality structure (NEO- Five-Factor-Inventory – NEO-FFI)[2,3], operationalizing

personality into dimensions or traits of agreeableness (friendly/compassionate vs.

challenging/detached), conscientiousness (efficient/organized vs. easy-going/careless),

extraversion (outgoing/ energetic vs. solitary/reserved), neuroticism (sensitive/nervous vs.

secure/confident), and openness (inventive/curious vs. consistent/cautious). These traits have

been linked to the quality of social relationships[4], job performance[5], and risk for mental

disorders[6,7]. For example, neuroticism has been related to a higher risk for psychiatric

disorders such as depression, stress and anxiety[8-10], whereas ADHD has been related to

extraversion[11]. At the same time, Big Five traits are five broad factors, combining more

specific aspects of personality[12]. Personality facets, which decompose personality traits to

more detailed descriptions of individual-difference[12], are used to further qualify inter-

individual difference in personality. For example, extraversion entails both social as well as

non-social components and can be subdivided in facets of sociability, positive affect, and

activity, whereas openness relates to intellect, but also unconventionality[12].

Personality traits as well as facets are highly heritable with about 50% of the variance

attributable to additive genetic factors[13-17] [18]. In the last decade genome-wide association

studies (GWAS) have also discovered several genetic variants associated with these

traits[11,19,20]. Recent work has also identified genetic correlations between personality and

psychiatric disorders. For example, extraversion was related to attention-deficit/hyperactivity

disorder, neuroticism to internalizing behavior, and openness to schizophrenia[11]. In addition

to developments in behavioral genetics, the relationship between personality and biology has

also been investigated using magnetic resonance imaging (MRI)[21-24]. While most studies

found some phenotypic relationship between brain structure and variations in personality

traits, the findings are inconsistent and often conflicting[21,23,24]. Possible causes for this

variability could be small sample sizes, variability in gender effects, as well as

methodological variations. Yet, despite often-conflicting reports, neurobehavioral correlates

of personality are often reported in multi-modal regions of the frontal, cingulate, and

temporo-parietal cortices implicated in higher cognitive processing. These regions have been

found to greatly expand over the course of primate evolution, with complex connectivity

patterns supporting cognitive flexibility[25]. Importantly, quantitative indices of brain structure

are also heritable[26-30]. This raises the question whether shared variations in personality traits

and brain structure are genetically driven, and if so, if the neurogenetic markers of personality

traits are located within regions associated with higher-order processing.

Here, we investigate the genetic correlation of personality traits and brain morphology

to probe the genetic basis of their relationship. We seek to answer the following three

questions: (i) Can we replicate previous observations of heritability of personality traits as

well as brain structure? (ii) Is the relationship between personality traits and brain structure

driven by genetic processes? (iii) Is there a genetic link between personality traits and

adaptive behavior at both the behavioral and brain level?

We assessed (co-)heritability of NEO-FFI personality traits and brain structure in a

large, well-characterized sample of young participants drawn from the Human Connectome

Project. This sample included monozygotic and dizygotic twins, siblings, and unrelated

individuals aged 22-37 which enabled the analysis of shared genetic variance between

personality and brain structure. We captured variations in brain morphometry using an atlas-

based approach[31] leading to a more homogeneous and biological meaningful representation

than common summary measures based on macro-anatomy[32]. Analysis of heritability and co-

heritability was performed using maximum likelihood variance-decomposition methods using

Sequential Oligogenic Linkage Analysis Routines (www.solar-eclipse-genetics.org; Solar

Eclipse 8.4.0.) Heritability ($h^2$) is the total additive genetic variance and genetic ($\rho_g$)

correlations were estimated using bivariate polygenic analyses. To qualify the function of the

brain regions observed genetically related to personality in terms of associated tasks and

behavior, we performed quantitative function decoding based on the BrainMap database[33].

**Results**
127
128    *Heritability and genetic correlation of NEO-FFI personality score (Figure 1)*
129
First, we assessed heritability and genetic correlations of personality traits using maximum

likelihood variance-decomposition methods with Solar[34]. We observed all NEO-FFI scores to

be heritable: agreeableness (h2=0.31, p<0.001), conscientiousness (h2=0.43, p<0.001),

extraversion (h2=0.45, p<0.001), neuroticism (h2=0.35, p<0.001), and openness (h2=0.57,

p<0.001). Openness and conscientiousness ($\rho_g$= -0.27, p<0.005) had significant negative

genetic correlations. Extraversion showed positive genetic correlation with agreeableness ($\rho_g$

= 0.40, p<0.003. Personality facets, constructed based on previous literature [12], where all

heritable and co-heritable with the NEO-FFI traits they were theorized to be subcomponents

of (see further; Supplementary Figure 1).

139
140    **FIGURE 1. INTER-INDIVIDUAL DIFFERENCES IN PERSONALITY ARE UNDER GENETIC
141    CONTROL.** A). Heritability of personality scores; B). Phenotypic and genetic correlation of
142    personality traits. Only findings that survive Bonferroni correction are reported.
143

*Genetic correlation of personality traits and brain structure (Figure 2)*

As all personality traits were significantly heritable in our current sample, we next confirmed

cortical thickness to be heritable in our parcel-based approach (Supplementary Figure 2). This

enabled us to perform genetic correlation analyses between personality traits and cortical

structure. Agreeableness showed genetic correlation (FDR p<0.05) with cortical thickness in

left posterior superior/mid temporal sulcus (rho=0.03; $\rho_g$ = 0.60, p<0.0005; $\rho_e$ = -0.15,

p<0.01), right superior frontal cortex extending to mid cingulate (rho=-0.04; $\rho_g$ = -0.55,

p<0.0004; $\rho_e$ =0.18, p<0.005), and right dorsolateral frontal cortex (rho=-0.02; $\rho_g$ = -0.51,

p<0.0006; $\rho_e$ =0.21, p<0.0004). In a post-hoc analysis, we verified our observations at the

level of agreeable facets, to investigate which personality facets genetic correlations at the

level of personality traits were driven by specific personality facets (Supplementary Figure 3).

This follow-up analysis revealed that the right superior frontal cortex related both to pro-

sociality and non-antagonistic traits, whereas left posterior superior/mid temporal sulcus and

right dorsolateral frontal cortex showed genetic correlation with non-antagonistic traits only.

Second, we observed a genetic correlation (FDR p<0.01) between extraversion and left TPJ

thickness (rho=0.1, p<0.001; $\rho_g$ = 0.49 p<0.00005; $\rho_e$ = -0.11, p<0.05). Follow up analysis of

the relationship between facets of extraversion and left TPJ thickness also showed significant

genetic correlation with 'sociability' facet of extraversion (see supplementary Figure 3).

Observations of genetic correlations remained after controlling for general cognitive ability

(Supplementary figure 4). We did not observe significant genetic correlations between

cortical thickness and Conscientiousness, Neuroticism and Openness. Last, we performed a

post-hoc analysis on the clusters that showed a significant phenotypic correlation between

brain structure and personality traits to assess whether these phenotypical correlations were

driven by genetic correlations. For agreeableness, we observed significant genetic correlation

between left superior dorsolateral frontal cortex ($\rho_g$ = -0.35 p<0.0038; $\rho_e$ = 0.01, p=ns). Other

clusters of agreeableness, as well as neuroticism and openness only showed genetic

correlations at trend levels (see further Supplementary Figure 5).

172
**FIGURE 2. SHARED GENETIC BASIS FOR PERSONALITY TRAITS AND BRAIN STRUCTURE.** A-E).
Upper cortical surface display: phenotypic correlation as reference for genetic correlation
analysis. Lower cortical surface display: Genetic correlation of personality trait and cortical
thickness.

FDR-corrected findings have a black outline, uncorrected findings (p<0.005) are shown semi-
transparent. For our both phenotypic and genetic correlation analysis between NEO-FFI traits
and brain structure, we performed whole brain analysis.
180

*Functional decoding of neurogenetic correlates of personality traits (Figure 3)*

To enable inference about processes that are sustained by the regions showing significant

genetic correlation with personality traits, we performed quantitative functional decoding

using the BrainMap database[35]. Here we observed TPJ, implicated in extraversion, to be

activated above chance by (socio-) cognitive tasks and hence should be implicated in

cognitive and social cognitive processing (p<0.05). Next to socio-cognitive processing, the

TPJ, was implicated in cognition related to reasoning and language. Of the parcels related to

agreeableness, the left posterior superior-mid temporal sulcus was related to various forms of

language processing, such as phonology, syntax, speech, and semantics (p<0.05), whereas the

left superior frontal / mid cingulate parcel was related to action imagination and execution, as

well as cognition and perception, and the dorsolateral prefrontal cortex was found implicated

in cognitive tasks involving memory and reasoning.

193
194  **FIGURE 3. NEUROGENETIC CORRELATES OF PERSONALITY TRAITS FUNCTIONALLY RELATE**
195  **TO (SOCIO-) COGNITIVE AND LANGUAGE PROCESSING.** Quantitative functional decomposition
196  of neurogenetic markers of personality based on the FDR-corrected clusters of phenotypical
197  relationships of Figure 2.
198

*Neurogenetic overlap between personality traits and adaptive behavior (Figure 4)*

Last, based on previous behavioral reports of genetic correlation between personality and

mental health, we sought to explore whether we could observe a neurogenetic overlap

between personality traits and mental health. As a proxy of mental health, we used self-report

data on adaptive function and problems, scored according to DSM-oriented scales. First, we

performed phenotypical and genetic correlation analyses between personality traits and sub-

clinical DSM-oriented markers of adaptive functions and problems at the behavioral level.

Following we explored whether regions that showed a genetic correlation with personality

traits also related to adaptive function and problems.

In our behavioral analysis, we observed considerable genetic correlation between personality

traits and markers of adaptive behavior. Agreeableness related negatively to anti-social

problems. Conscientiousness showed negative genetic correlation with AD/H, depressive, and

inattention problems, whereas extraversion related negatively to avoidant behavior.

Neuroticism showed a positive genetic correlation to a broad range of adaptive functions

including AD/H problems, anxiety, avoidant, depressive, hyperactive, and inattention

problems (see Further Supplementary table 1). Follow up analysis of personality facets in

relation to adaptive function and problems followed these patterns (Supplementary Figure 6).

Genetic correlations between adaptive behavior and regions associated with personality traits

followed behavioral patterns, with TPJ showing genetic correlation with AD/H (rho=0.04; $\rho_g$

= 0.23, p<0.05; $\rho_e$ = -0.06, p<n.s.) as well as avoidant problems (rho=-0.08; $\rho_g$ = -0.23,

p<0.05; $\rho_e$ = 0.04, p<n.s.) and right dorsolateral frontal cortex relating to hyperactivity

problems (rho=0.02; $\rho_g$ = 0.25, p<0.05; $\rho_e$ =-0.09, p<n.s.).

224
**FIGURE 4. NEO-FFI PERSONALITY TRAITS AND ADAPTIVE BEHAVIOR SHOW GENETIC**
**CORRELATION IN BEHAVIORAL AND BRAIN.** A). Phenotypic and genetic correlation between
NEO-FFI personality traits and ASR, DSM oriented scale[36], only correlations that survive
Bonferroni corrections are displayed; B). Post-hoc relation of adaptive behavior to brain
regions significantly associated with personality traits, * indicates a phenotypic/genetic
correlation below 0.05.
231

233
234
235

**Discussion**

Personality is an individuals' characteristic set of cognitive, emotional and behavioral patterns. Previous work has shown that personality is heritable and correlates with variation in brain structure [11,17,21-24]. Here, we examined the neurogenetic basis of personality and brain structure using large-scale structural imaging and pedigree structure of the HCP. Our analysis showed that brain-behavior relationships in personality are in part driven by shared genetic effects. We observed genetic correlations of agreeableness with cortical thickness in posterior superior/mid temporal sulcus, superior frontal/mid cingulate, and dorsolateral frontal cortex and of extraversion with cortical thickness in the TPJ. Structural regions that had a genetic link with personality traits were observed implicated in (socio-) cognitive, emotion, and language processing as revealed by quantitative functional decoding using the BrainMap database. Together our results provide evidence that the correlation between brain structure and personality traits is, in part, driven by shared genetic processes anchored in regions involved in (socio-)cognitive and language processes. Our observations suggest a direct genetic link between variations in local brain morphometry, function of the respective brain regions, and complex behavioral traits such as personality.

In the current study focused on the relationship between individual differences in personality traits and local anatomy, using compressed surface-based MRI data based on the parcellation scheme of Schaefer et al. Using compressed features of structural MRI has been suggested to both to improve signal-to-noise of brain measures, assuming the cortex can be subdivided in cortical areas that bridge different features such as connectivity and structure (cf. Eickhoff et al.[32] and Genon et al.[33]), as well as to optimize analysis scalability in our genetic analyses. The Schaefer parcellation is derived using functional MRI data from ~1500 subjects, by integrating local approaches that detect abrupt transitions in functional connectivity patterns and global approaches that cluster similar functional connectivity patterns[31]. Previous

research found that the Schaefer parcellation showed convergence with partitions based on

structure alone[37]. A combination of within-area micro circuitry, proxied by brain

morphometry, and between-area connectivity enables each area to perform a unique set of

computations[38]. Therefore, a parcellation approach that considers both local and global

connectivity benefits structural image analysis, as it reduces signal-to-noise both within

individual as across individuals and make control for multiple comparisons more

straightforward[33]. Based on the findings in our study we suggest our approach might be a

fruitful first exploratory step to further investigate the shared genetic basis of brain and

behavior, and locate neural mechanisms of interest. Future studies can then verify these

results by exploring more specific genetic mechanisms, as well as neuroanatomical features.

For example, there is converging evidence that inter-individual differences in topological

integrative features of multi-modal areas are more important for higher-order cognitive

functions, such as intelligence, than local features only[39].

The present results indicate a genetic correlation between complex behavioral traits and brain

structure. The strength of genetic correlations is strongly dependent on the heritability of each

of the correlated markers. In our sample, between 30% and 60% (on average 42%) of

variance in personality traits was explained by additive genetic factors, with similar

heritability estimates for meta-traits and facets. This is in line with previous studies using twin

and family samples[15] as well as genome-wide approaches[11]. A recent meta-analysis[17]

confirmed that an average 40% of the variance in personality traits is of genetic origin. Also,

conform previous studies[28,40-43], we observe high ($h2>0.4$) heritability of cortical thickness,

with highest values in primary sensory areas. The heritability patterns highlight high

heritability of cortical thickness in unimodal cortices, whereas variance in association cortices

is on average less influenced by genetic factors [28,30,40-42,44,45].

Genetic correlation analyses between brain morphometry and personality traits revealed a

genetic link of agreeableness and extraversion to brain structure. Here, we observed a genetic

correlation between agreeableness and cortical thickness in posterior superior-mid temporal

sulcus, superior frontal/mid cingulate, and dorsolateral frontal cortex. Agreeableness is the

ability to trust others, be cooperative, and is linked to prosocial behavior[46,47] as well as to non-

antagonistic traits.   Posterior superior-mid temporal areas, including Wernicke's area, are

consistently implicated in language processing[48], as also shown by our functional decoding

analysis, and are part of the default mode network[31,49]. Superior frontal/mid cingulate areas,

are associated by functional decoding with imagination and execution of action[50], whereas

dorsolateral region is associated by functional decoding with reasoning and memory[51]. Both

cingulate as well as dorsolateral regions are involved in higher-order reasoning, meta-

cognition, and cognitive control[52-54].   Indeed, the genetic relationship between agreeableness

and regions involved in higher-order reasoning is in line with theoretical accounts that the

ability to reflect on ones actions is essential for being able to be cooperative and prosocial[52].

At the same time, we observed a moderate relationship of dorsolateral regions with

hyperactivity behaviors. Previous meta-analyses in patient populations have linked

abnormalities in dorsolateral prefrontal functioning to Ad/H[55]. Overall, neurogenetic overlap

between behavioral functions such as trust and cooperation, as well as problems of

hyperactive behavior, and brain systems involved in communication and reasoning highlight

functional specificity of genetic control over brain and behavior.

In case of extraversion we observed similar functional specificity. Extraversion was

genetically associated with thickness of the TPJ. Regions in the temporo-parietal junction

have repeatedly been associated with both social cognition and language[56-59]. Indeed, this was

confirmed by our functional decoding analysis, which showed that these areas were

predominantly related to tasks that involved (social-)cognitive processing. In line with these

findings, the sociability, but not positive affect, facet of the extraversion trait had a significant

genetic relation to TPJ. At the same time, we observed negative genetic correlation of

avoidance and hyperactivity with TPJ structure. In line with extraversions facets, extraversion

does not only entail sociability but also 'action' components. Though only at trend level, also

action facets of extraversion related to TPJ structure. Indeed, functionally and anatomically

the TPJ is placed at a nexus. Its reconciling role combining attention, language, memory, and

social cognitive processing also relates to the simulation of action in self and others[60].

Abnormal structure might result in impaired functional processing which in turn results in

avoidance behavior or hyperactivity. Importantly, previous work has been able to identify

sub-components of in the TPJ [61,62], and further studies on the genetic differentiability of these

sub-regions will clarify its relation to social and non-social behavioral processes.

The functional specificity of neurogenetic correlates of personality traits suggests a direct

genetic link between variations in local brain morphometry, function of the respective brain

regions, and complex behavioral traits such as personality. The observed genetic correlation

possibly reflects a persistent effect of genetic factors on specific brain structures, which gives

rise to specific behaviors. On the other hand, however, it could also be that genetic

predispositions shape behavioral traits in other ways than through brain morphometry and

thus modulate brain structure of these regions only indirectly through use-dependent

plasticity, e.g., by parental influences[63] or other mechanisms that cannot be seen in MRI.

Interestingly, whole brain genetic correlations showed inverse effects of genetic and

environmental correlations, suggesting complex interactions between genes and

environmental are at play in the context of brain – behavior relationships. Environmental

factors can modulate genetic predisposion in various ways[64], including enhancement,

sensitivity, as well as (negative) interactions. Indeed, though the genetic overlap between

participants in our sample should already have maximized genetic influence on phenotypical

correlations between personality and brain, phenotypic correlations were not driven by

genetic processes alone, but rather a combination of both genetic and environmental effects.

Importantly, we found little consistency in the patterns of environmental influences on brain

morphometry and personality traits. This might be due to methodological limitations, since

we had insufficient data to test for household effects, and thus environmental correlation in

the current study results from unmeasured aspects of the environment or correlated

measurement errors[65]. Given the twin set-up as well as age range of the HCP sample, other

designs with broader age ranges and kinship relations might help to further understand

environmental contributions to brain-behavior relationships.

Nevertheless, previous studies have shown that environment plays an important role in

shaping brain-behavior relationships in the context of higher-cognitive and social skills. For

example, various studies have shown how social-environmental factors such as city-living

impact regulation of negative emotion and stress by mediation of genetic predispositions[66,67].

Also, improvement in socio-cognitive abilities following mental training have been shown to

result in structural changes of the TPJ[68].

In sum, we demonstrate a shared genetic basis of variance in brain structure and personality

traits. Functional decoding analysis indicated that regions genetically linked to personality

traits were functionally involved in to (socio-) cognitive and language processes, supporting

previous research on the biological basis of personality. Studying the genetic correlation

between brain and personality might help to better describe and understand the neural

endophenotype of personality traits, as well as the contribution of genetic and environmental

factors to inter-individual difference. Ultimately genetic neuroimaging endophenotypes of

personality traits can further help understand genetic predisposition and risk factors for mental

illness.

**Methods**

*Participants and study design*

For our analysis we used the publicly available data from the Human Connectome Project (HCP; http://www.humanconnectome.org/), which comprised data from 1206 individuals (656 females), 298 MZ twins, 188 DZ twins, and 720 singletons, with mean age 28.8 years (SD = 3.7, range = 22–37). We included for whom the scans and data had been released (humanconnectome.org) after passing the HCP quality control and assurance standards [69]. The full set of inclusion and exclusion criteria are described elsewhere[70,71].

After removing individuals with missing structural imaging or behavioral data our sample consisted of 1085 individuals (283 MZ-twins, 167 DZ-twins, and 579 non-twin siblings and 57 singles) with mean age of 28.8 years (SD =3.7, range =22-37).

*Structural imaging processing*

MRI protocols of the HCP are previously described in[70,71]. The pipeline used to obtain the Freesurfer-segmentation is described in detail in a previous article [70] and is recommended for the HCP-data. The pre-processing steps included co-registration of T1 and T2 scans, B1 (bias field) correction, and segmentation and surface reconstruction using FreeSurfer version 5.3-HCP to estimate cortical thickness.

We used a parcellation scheme[31] based on the combination of a local gradient approach and a global similarity approach using a gradient-weighted Markov Random models. The parcellation has been extensively evaluated with regards to stability and convergence with histological mapping and alternative parcellations. In the context of the current study, we focus on the granularity of 200 parcels, as averaging will improve signal-to-noise. 200 parcels are close to granularity used in Enigma (Desikan-atlas, 68 parcels) while at the same time providing more anatomical detail in functionally heterogeneous regions such as parietal and frontal cortex. In order to improve signal-to-noise and improve analysis speed, we opted to

average unsmoothed structural data within each parcel. Thus, cortical thickness of each ROI

was estimated as the trimmed mean (10 percent trim).

*Behavioral assessment*

The NEO-FFI personality traits were assessed using the NEO-Five-Factors-Inventory (NEO-

FFI)[72]. The NEO-FFI is composed of a subset of 60-items extracted from the full-length 240-

item NEO-PI-R. For each item, participants reported their level of agreement on a 5-point

Likert scale, from strongly disagree to strongly agree. The NEO instruments have been

previously validated in USA and several other countries[73]. As conventional scoring of the

NEO-FFI does not provide scores on more specific aspects of the broad-bandwidth factors,

we also computed facets based on Saucier et al[12]. Here, 13 factor-analytically derived scales

were found to replicate across halves of a sample of self-descriptions by adults (N = 732). The

scales demonstrated reliability and factor structure comparable to that of the 30 facet scales of

the NEO-PI-R.

For markers of adaptive behavior and problems, we used the Achenbach Adult Self-Report

(ASR) for Ages 18-59[36]. The ASR is a self-administered test examining diverse aspects of

adaptive functioning and problems. Scales are based on 2020 referred adults and normed on

1767 non-referred adults. The test-retest reliability of the ASR was supported by 1-week test-

retest that were all above 0.71. The ASR also has good internal consistency (0.83).

(https://aseba.org/wp-content/uploads/2019/01/ASEBA-Reliability-and-Validity-Adult.pdf).

As a proxy for IQ we used the NIH Toolbox Cognition[74], 'total composite score'. The

Cognitive Function Composite score is derived by averaging the normalized scores of each of

the Fluid and Crystallized cognition measures, then deriving scale scores based on this new

distribution. Higher scores indicate higher levels of cognitive functioning. Participant score is

normed to those in the entire NIH Toolbox Normative Sample (18 and older), regardless of

age or any other variable, where a score of 100 indicates performance that was at the national

average and a score of 115 or 85, indicates performance 1 SD above or below the national

average.

*Statistical analysis*
425
To investigate the heritability and genetic correlation of brain structure and personality traits,

we analyzed 200 parcels of cortical thickness, as well as personality trait score of each subject

in a twin-based heritability analysis. As in previous studies[65], the quantitative genetic analyses

were conducted using Sequential Oligogenic Linkage Analysis Routines (SOLAR)[34]. SOLAR

uses maximum likelihood variance-decomposition methods to determine the relative

importance of familial and environmental influences on a phenotype by modeling the

covariance among family members as a function of genetic proximity. This approach can

handle pedigrees of arbitrary size and complexity and thus, is optimally efficient with regard

to extracting maximal genetic information. To ensure that our traits, behavioral as well as of

cortical thickness parcels, conform to the assumptions of normality, an inverse normal

transformation was applied for all behavioral and neuroimaging traits[65].

Heritability ($h^2$) represents the portion of the phenotypic variance ($\sigma^2_p$) accounted for by the

total additive genetic variance ($\sigma^2_g$), i.e., $h^2 = \sigma^2_g/\sigma^2_p$. Phenotypes exhibiting stronger

covariances between genetically more similar individuals than between genetically less

similar individuals have higher heritability. Within SOLAR, this is assessed by contrasting the

observed covariance matrices for a neuroimaging measure with the structure of the covariance

matrix predicted by kinship. Heritability analyses were conducted with simultaneous

estimation for the effects of potential covariates. For this study, we included covariates

including age, sex, age × sex interaction, age$^2$, age$^2$ × sex interaction. Post-hoc we also

controlled for a proxy for intelligence, total cognitive score[74]. When investigating cortical

thickness, we additionally controlled for global thickness effects (mean cortical thickness).

Heritability estimates were corrected for multiple comparisons using Benjamin-Hochberg

FDR[75] at whole-brain analysis. Bonferroni correction was applied when investigating

behavior or in post-hoc brain analysis, and corrections across number of analysis were

applied.

To determine if variations in personality and brain structure were influenced by the same

genetic factors, genetic correlation analyses were conducted. More formally, bivariate

polygenic analyses were performed to estimate genetic ($\rho_g$) and environmental ($\rho_e$)

correlations, based on the phenotypical correlation ($\rho_p$), between brain structure and

personality with the following formula: $\rho_p = \rho_g\sqrt{(h^2_1 h^2_2)} + \rho_e\sqrt{[(1 - h^2_1)(1 - h^2_2)]}$,

where $h^2_1$ and $h^2_2$ are the heritability's of the parcel-based cortical thickness and the various

behavioral traits. The significance of these correlations was tested by comparing the log

likelihood for two restricted models (with either $\rho_g$ or $\rho_e$ constrained to be equal to 0) against

the log likelihood for the model in which these parameters were estimated. For our

phenotypical analysis we performed Spearman correlation analysis, controlling for the same

variables as in the genetic analysis, namely age, sex, age $\times$ sex interaction, $age^2$, $age^2 \times$ sex

interaction, as well as global thickness effects when investigating brain structure. A

significant genetic correlation (corrected for multiple comparisons using Benjamin-Hochberg

FDR[75] in case of whole-brain analysis and Bonferroni correction when investigating behavior

or in post-hoc brain analysis) is evidence suggesting that (a proportion of) both phenotypes

are influenced by a gene or set of genes[76].

467
*Functional decoding*

All significant parcels were in a last step functionally characterized using the Behavioral

Domain    meta-data    from    the    BrainMap    database    using    forward    inference

(http://www.brainmap.org[77,78]). To do so, volumetric counterparts of the surface-based parcels

were identified.  In particular, we identified those meta-data labels (describing the computed

contrast [behavioral domain]) that were significantly more likely than chance to result in

activation of a given parcel[22,33,35]. That is, functions were attributed to the identified

neurogenetic effects by quantitatively determining which types of experiments are associated

with activation in the respective parcellation region. Significance was established using a

binomial test ($p < 0.05$, corrected for multiple comparisons using false discovery rate (FDR)).

**References**

1. Mischel, W. Toward an integrative science of the person. *Annu Rev Psychol* **55**, 1-22, doi:10.1146/annurev.psych.55.042902.130709 (2004).
2. McCrae, R. R. & Costa, P. T., Jr. Updating Norman's "Adequate Taxonomy": intelligence and personality dimensions in natural language and in questionnaires. *J Pers Soc Psychol* **49**, 710-721 (1985).
3. McCrae, R. R. & Costa, P. T., Jr. Validation of the five-factor model of personality across instruments and observers. *J Pers Soc Psychol* **52**, 81-90 (1987).
4. Asendorpf, J. B. & Wilpers, S. Personality effects on social relationships. *Journal of Personality and Social Psychology* **74**, 1531-1544 (1998).
5. Rothmann, S. & Coetzer, E. P. The big five personality dimensions and job performance. *SA Journal of Industrial Psychology* **29** (2003).
6. Miller, J. D., Lynam, D. R., Widiger, T. A. & Leukefeld, C. Personality disorders as extreme variants of common personality dimensions: can the Five-Factor Model adequately represent psychopathy? *J Pers* **69**, 253-276 (2001).
7. Trull, T. J. Dimensional models of personality: the five-factor model and the DSM-5. *Dialogues Clin Neurosci* **15**, 135-146 (2013).
8. Roelofs, J., Huibers, M., Peeters, F., Arntz, A. & van Os, J. Rumination and worrying as possible mediators in the relation between neuroticism and symptoms of depression and anxiety in clinically depressed individuals. *Behav Res Ther* **46**, 1283-1289, doi:10.1016/j.brat.2008.10.002 (2008).
9. Brainstorm, C. *et al.* Analysis of shared heritability in common disorders of the brain. *Science* **360**, doi:10.1126/science.aap8757 (2018).
10. Gunthert, K. C., Cohen, L. H. & Armeli, S. The role of neuroticism in daily stress and coping. *J Pers Soc Psychol* **77**, 1087-1100 (1999).
11. Lo, M. T. *et al.* Genome-wide analyses for personality traits identify six genomic loci and show correlations with psychiatric disorders. *Nat Genet* **49**, 152-156, doi:10.1038/ng.3736 (2017).
12. Saucier, G. Replicable Item-Cluster Subcomponents in the NEO Five-Factor Inventory. *Journal of Personality Assessment* **70**, 263-276 (1998).
13. Bouchard, T. J., Jr. Genes, environment, and personality. *Science* **264**, 1700-1701 (1994).
14. Bouchard, T. J., Jr. & McGue, M. Genetic and environmental influences on human psychological differences. *J Neurobiol* **54**, 4-45, doi:10.1002/neu.10160 (2003).
15. Jang, K. L., Livesley, W. J. & Vernon, P. A. Heritability of the big five personality dimensions and their facets: a twin study. *J Pers* **64**, 577-591 (1996).
16. Loehlin, J. C., McCrae, R. R., Costa, P. T. & John, O. P. Heritabilities of Common and Measure-Specific Components of the Big Five Personality Factors. *Journal of Research in Personality* **32**, 431-453 (1998).
17. Vukasovic, T. & Bratko, D. Heritability of personality: A meta-analysis of behavior genetic studies. *Psychol Bull* **141**, 769-785, doi:10.1037/bul0000017 (2015).
18. Bouchard, T. J., Jr. & Loehlin, J. C. Genes, evolution, and personality. *Behav Genet* **31**, 243-273 (2001).
19. de Moor, M. H. *et al.* Meta-analysis of genome-wide association studies for personality. *Molecular psychiatry* **17**, 337-349, doi:10.1038/mp.2010.128 (2012).
20. van den Berg, S. M. *et al.* Meta-analysis of Genome-Wide Association Studies for Extraversion: Findings from the Genetics of Personality Consortium. *Behav Genet* **46**, 170-182, doi:10.1007/s10519-015-9735-5 (2016).

528  21    DeYoung, C. G. *et al.* Testing predictions from personality neuroscience. Brain
529            structure and the big five. *Psychol Sci* **21**, 820-828, doi:10.1177/0956797610370159
530            (2010).
531  22    Nostro, A. D., Muller, V. I., Reid, A. T. & Eickhoff, S. B. Correlations Between
532            Personality and Brain Structure: A Crucial Role of Gender. *Cereb Cortex* **27**, 3698-
533            3712, doi:10.1093/cercor/bhw191 (2017).
534  23    Riccelli, R., Toschi, N., Nigro, S., Terracciano, A. & Passamonti, L. Surface-based
535            morphometry reveals the neuroanatomical basis of the five-factor model of
536            personality. *Soc Cogn Affect Neurosci* **12**, 671-684, doi:10.1093/scan/nsw175 (2017).
537  24    Bjornebekk, A. *et al.* Neuronal correlates of the five factor model (FFM) of human
538            personality: Multimodal imaging in a large healthy sample. *Neuroimage* **65**, 194-208,
539            doi:10.1016/j.neuroimage.2012.10.009 (2013).
540  25    Sneve, M. H. *et al.* High-Expanding Regions in Primate Cortical Brain Evolution
541            Support Supramodal Cognitive Flexibility. *Cereb Cortex*, doi:10.1093/cercor/bhy268
542            (2018).
543  26    Baare, W. F. *et al.* Quantitative genetic modeling of variation in human brain
544            morphology. *Cereb Cortex* **11**, 816-824 (2001).
545  27    Glahn, D. C., Thompson, P. M. & Blangero, J. Neuroimaging endophenotypes:
546            strategies for finding genes influencing brain structure and function. *Hum Brain Mapp*
547            **28**, 488-501, doi:10.1002/hbm.20401 (2007).
548  28    Strike, L. T. *et al.* Genetic Complexity of Cortical Structure: Differences in Genetic
549            and Environmental Factors Influencing Cortical Surface Area and Thickness. *Cereb*
550            *Cortex*, doi:10.1093/cercor/bhy002 (2018).
551  29    Thompson, P. M. *et al.* Genetic influences on brain structure. *Nat Neurosci* **4**, 1253-
552            1258, doi:10.1038/nn758 (2001).
553  30    Teeuw, J. *et al.* Genetic Influences on the Development of Cerebral Cortical
554            Thickness During Childhood and Adolescence in a Dutch Longitudinal Twin Sample:
555            The Brainscale Study. *Cereb Cortex*, doi:10.1093/cercor/bhy005 (2018).
556  31    Schaefer, A. *et al.* Local-Global Parcellation of the Human Cerebral Cortex from
557            Intrinsic Functional Connectivity MRI. *Cereb Cortex* **28**, 3095-3114,
558            doi:10.1093/cercor/bhx179 (2018).
559  32    Eickhoff, S. B., Constable, R. T. & Yeo, B. T. T. Topographic organization of the
560            cerebral cortex and brain cartography. *Neuroimage* **170**, 332-347,
561            doi:10.1016/j.neuroimage.2017.02.018 (2018).
562  33    Genon, S., Reid, A., Langner, R., Amunts, K. & Eickhoff, S. B. How to Characterize
563            the Function of a Brain Region. *Trends Cogn Sci* **22**, 350-364,
564            doi:10.1016/j.tics.2018.01.010 (2018).
565  34    Almasy, L. & Blangero, J. Multipoint quantitative-trait linkage analysis in general
566            pedigrees. *Am J Hum Genet* **62**, 1198-1211, doi:10.1086/301844 (1998).
567  35    Fox, P. T., Lancaster, J. L., Laird, A. R. & Eickhoff, S. B. Meta-analysis in human
568            neuroimaging: computational modeling of large-scale databases. *Annu Rev Neurosci*
569            **37**, 409-434, doi:10.1146/annurev-neuro-062012-170320 (2014).
570  36    Achenbach, T. M. & Rescorla, L. A.   (ed Youth and Families Research center for
571            Children) (University of Vermont, Burlington, VT, USA, 2003).
572  37    Varikuti, D. P. *et al.* Evaluation of non-negative matrix factorization of grey matter in
573            age prediction. *Neuroimage* **173**, 394-410, doi:10.1016/j.neuroimage.2018.03.007
574            (2018).
575  38    Van Essen, D. C. & Glasser, M. F. Parcelling Cerebral Cortex: How Invasive
576            Animal Studies Inform Noninvasive Mapmaking in Humans. *Neuron* **99**, 640-663,
577            doi:10.1016/j.neuron.2018.07.002 (2018).

| 578 | 39 | Seidlitz, J. *et al.* Morphometric Similarity Networks Detect Microscale Cortical |
| 579 | | Organization and Predict Inter-Individual Cognitive Variation. *Neuron* **97**, 231-247 |
| 580 | | e237, doi:10.1016/j.neuron.2017.11.039 (2018). |
| 581 | 40 | Kremen, W. S. *et al.* Genetic and environmental influences on the size of specific |
| 582 | | brain regions in midlife: the VETSA MRI study. *Neuroimage* **49**, 1213-1223, |
| 583 | | doi:10.1016/j.neuroimage.2009.09.043 (2010). |
| 584 | 41 | Eyler, L. T. *et al.* A comparison of heritability maps of cortical surface area and |
| 585 | | thickness and the influence of adjustment for whole brain measures: a magnetic |
| 586 | | resonance imaging twin study. *Twin Res Hum Genet* **15**, 304-314, |
| 587 | | doi:10.1017/thg.2012.3 (2012). |
| 588 | 42 | Panizzon, M. S. *et al.* Distinct genetic influences on cortical surface area and cortical |
| 589 | | thickness. *Cereb Cortex* **19**, 2728-2735, doi:10.1093/cercor/bhp026 (2009). |
| 590 | 43 | Winkler, A. M. *et al.* Cortical thickness or grey matter volume? The importance of |
| 591 | | selecting the phenotype for imaging genetics studies. *Neuroimage* **53**, 1135-1146, |
| 592 | | doi:10.1016/j.neuroimage.2009.12.028 (2010). |
| 593 | 44 | Hofer, E. & al., e. Genetic Determinants of Cortical Structure (Thickness, Surface |
| 594 | | Area and Volumes) among Disease Free Adults in the CHARGE Consortium. *bioRxiv* |
| 595 | | *preprint first posted online Sep. 9, 2018; doi: http://dx.doi.org/10.1101/409649.* |
| 596 | | (2018). |
| 597 | 45 | Fjell, A. M. *et al.* Development and aging of cortical thickness correspond to genetic |
| 598 | | organization patterns. *Proc Natl Acad Sci U S A* **112**, 15462-15467, |
| 599 | | doi:10.1073/pnas.1508831112 (2015). |
| 600 | 46 | Martin-Raugh, M. P., Kell, H. J. & Motowidlo, S. J. Prosocial knowledge mediates |
| 601 | | effects of agreeableness and emotional intelligence on prosocial behavior. *Pers Individ* |
| 602 | | *Dif.* **90**, 41-49 (2016). |
| 603 | 47 | Hilbig, B. E., Glockner, A. & Zettler, I. Personality and prosocial behavior: linking |
| 604 | | basic traits and social value orientations. *J Pers Soc Psychol* **107**, 529-539, |
| 605 | | doi:10.1037/a0036074 (2014). |
| 606 | 48 | Binder, J. R. The Wernicke area: Modern evidence and a reinterpretation. *Neurology* |
| 607 | | **85**, 2170-2175, doi:10.1212/WNL.0000000000002219 (2015). |
| 608 | 49 | Yeo, B. T. *et al.* The organization of the human cerebral cortex estimated by intrinsic |
| 609 | | functional connectivity. *J Neurophysiol* **106**, 1125-1165, doi:10.1152/jn.00338.2011 |
| 610 | | (2011). |
| 611 | 50 | Nachev, P., Wydell, H., O'Neill, K., Husain, M. & Kennard, C. The role of the pre- |
| 612 | | supplementary motor area in the control of action. *Neuroimage* **36 Suppl 2**, T155-163, |
| 613 | | doi:10.1016/j.neuroimage.2007.03.034 (2007). |
| 614 | 51 | Mars, R. B. & Grol, M. J. Dorsolateral prefrontal cortex, working memory, and |
| 615 | | prospective coding for action. *J Neurosci* **27**, 1801-1802 (2007). |
| 616 | 52 | Frith, C. D. The role of metacognition in human social interactions. *Philos Trans R* |
| 617 | | *Soc Lond B Biol Sci* **367**, 2213-2223, doi:10.1098/rstb.2012.0123 (2012). |
| 618 | 53 | Cole, M. W., Yarkoni, T., Repovs, G., Anticevic, A. & Braver, T. S. Global |
| 619 | | connectivity of prefrontal cortex predicts cognitive control and intelligence. *J* |
| 620 | | *Neurosci* **32**, 8988-8999 (2012). |
| 621 | 54 | MacDonald, A. W., 3rd, Cohen, J. D., Stenger, V. A. & Carter, C. S. Dissociating the |
| 622 | | role of the dorsolateral prefrontal and anterior cingulate cortex in cognitive control. |
| 623 | | *Science* **288**, 1835-1838 (2000). |
| 624 | 55 | Hart, H., Radua, J., Nakao, T., Mataix-Cols, D. & Rubia, K. Meta-analysis of |
| 625 | | functional magnetic resonance imaging studies of inhibition and attention in attention- |
| 626 | | deficit/hyperactivity disorder: exploring task-specific, stimulant medication, and age |
| 627 | | effects. *JAMA Psychiatry* **70**, 185-198, doi:10.1001/jamapsychiatry.2013.277 (2013). |

56  Bzdok, D. *et al.* Left inferior parietal lobe engagement in social cognition and
language. *Neurosci Biobehav Rev* **68**, 319-334, doi:10.1016/j.neubiorev.2016.02.024
(2016).
57  Mar, R. A. The neural bases of social cognition and story comprehension. *Annu Rev*
*Psychol* **62**, 103-134, doi:10.1146/annurev-psych-120709-145406 (2011).
58  Kanske, P., Bockler, A., Trautwein, F. M. & Singer, T. Dissecting the social brain:
Introducing the EmpaToM to reveal distinct neural networks and brain-behavior
relations for empathy and Theory of Mind. *Neuroimage* **122**, 6-19,
doi:10.1016/j.neuroimage.2015.07.082 (2015).
59  Valk, S. L. *et al.* Socio-Cognitive Phenotypes Differentially Modulate Large-Scale
Structural Covariance Networks. *Cereb Cortex*, doi:10.1093/cercor/bhv319 (2016).
60  Carter, R. M. & Huettel, S. A. A nexus model of the temporal-parietal junction.
*Trends Cogn Sci* **17**, 328-336, doi:10.1016/j.tics.2013.05.007 (2013).
61  Krall, S. C. *et al.* The role of the right temporoparietal junction in attention and social
interaction as revealed by ALE meta-analysis. *Brain Struct Funct* **220**, 587-604,
doi:10.1007/s00429-014-0803-z (2015).
62  Mars, R. B. *et al.* Connectivity-based subdivisions of the human right
"temporoparietal junction area": evidence for different areas participating in different
cortical networks. *Cereb Cortex* **22**, 1894-1903, doi:10.1093/cercor/bhr268 (2012).
63  Kong, A. *et al.* The nature of nurture: Effects of parental genotypes. *Science* **359**, 424-
428, doi:10.1126/science.aan6877 (2018).
64  Reiss, D., Leve, L. D. & Neiderhiser, J. M. How genes and the social environment
moderate each other. *Am J Public Health* **103 Suppl 1**, S111-121,
doi:10.2105/AJPH.2013.301408 (2013).
65  Glahn, D. C. *et al.* Genetic control over the resting brain. *Proc Natl Acad Sci U S A*
**107**, 1223-1228, doi:10.1073/pnas.0909969107 (2010).
66  Meyer-Lindenberg, A. & Tost, H. Neural mechanisms of social risk for psychiatric
disorders. *Nat Neurosci* **15**, 663-668, doi:10.1038/nn.3083 (2012).
67  Tost, H., Champagne, F. A. & Meyer-Lindenberg, A. Environmental influence in the
brain, human welfare and mental health. *Nat Neurosci* **18**, 1421-1431,
doi:10.1038/nn.4108 (2015).
68  Valk, S. L. *et al.* Structural plasticity of the social brain: Differential change after
socio-affective and cognitive mental training. *Sci Adv* **3**, e1700489,
doi:10.1126/sciadv.1700489 (2017).
69  Marcus, D. S. *et al.* Human Connectome Project informatics: quality control, database
services, and data visualization. *Neuroimage* **80**, 202-219,
doi:10.1016/j.neuroimage.2013.05.077 (2013).
70  Glasser, M. F. *et al.* The minimal preprocessing pipelines for the Human Connectome
Project. *Neuroimage* **80**, 105-124, doi:10.1016/j.neuroimage.2013.04.127 (2013).
71  Van Essen, D. C. *et al.* The WU-Minn Human Connectome Project: an overview.
*Neuroimage* **80**, 62-79, doi:10.1016/j.neuroimage.2013.05.041 (2013).
72  McCrae, R. R. & Costa, P. T. A contemplated revision of the NEO Five-Factor
Inventory. *Pers Individ Dif.* **36**, 587-596 (2004).
73  McCrae, R. R. & Terracciano, A. Personality profiles of cultures: aggregate
personality traits. *J Pers Soc Psychol* **89**, 407-425, doi:10.1037/0022-3514.89.3.407
(2005).
74  Weintraub, S. *et al.* Cognition assessment using the NIH Toolbox. *Neurology* **80**, S54-
64, doi:10.1212/WNL.0b013e3182872ded (2013).
75  Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and
Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society.*
*Series B (Methodological)* **57**, 289-300 (1995).

76   Almasy, L., Dyer, T. D. & Blangero, J. Bivariate quantitative trait linkage analysis:
pleiotropy versus co-incident linkages. *Genet Epidemiol* **14**, 953-958,
doi:10.1002/(SICI)1098-2272(1997)14:6<953::AID-GEPI65>3.0.CO;2-K (1997).
77   Laird, A. R. *et al.* The BrainMap strategy for standardization, sharing, and meta-
analysis of neuroimaging data. *BMC Res Notes* **4**, 349, doi:10.1186/1756-0500-4-349
(2011).
78   Laird, A. R. *et al.* ALE Meta-Analysis Workflows Via the Brainmap Database:
Progress Towards A Probabilistic Functional Brain Atlas. *Front Neuroinform* **3**, 23,
doi:10.3389/neuro.11.023.2009 (2009).
688
689
690

691 **Contributions**

692 S.L.V. and S.B.E. conceived this study. S.L.V. developed and implements the analysis, with

693 input from F.H., B.T.T.Y. and P.K. J.C. implemented the Quantitative functional decoding

694 analysis. S.L.V. wrote the paper, and S.L.V., F.E., J.C., B.T.T.Y, P.K., and S.B.E. were

695 involved in interpreting results and editing the manuscript.