PERSONALITY AND LOCAL BRAIN STRUCTURE: THEIR SHARED GENETIC BASIS AND REPRODUCIBILITY

Sofie L. Valk[1,2*]; Felix Hoffstaedter[1,2]; Julia A. Camilleri[1,2]; Peter Kochunov[3]; B.T. Thomas Yeo[4,5,6]; Simon B. Eickhoff[1,2]

[1)] *Institute of Neuroscience and Medicine (INM-7: Brain and Behaviour), Research Centre Jülich, 52425 Jülich, Germany*
[2)] *Institute of Systems Neuroscience, Heinrich Heine University Düsseldorf, 40225 Düsseldorf, Germany*
[3)] *Maryland Psychiatric Research Center, University of Maryland School of Medicine, Baltimore, Maryland, USA.*
[4)] *Department of Electrical and Computer Engineering, Centre for Sleep & Cognition, Clinical Imaging Research Centre, and N.1 Institute for Health, National University of Singapore, Singapore, Singapore.*
[5)] *Athinoula A. Martinos Center for Biomedical Imaging, Massachusetts General Hospital, Charlestown, MA, USA.*
[6)] *NUS Graduate School for Integrative Sciences and Engineering, National University of Singapore, Singapore, Singapore.*

Abstract

Local variation in cortical architecture is highly heritable and distinct genes are

associated with specific cortical regions. Total surface area has been shown to be

genetically correlated with complex cognitive capacities, suggesting cortical brain

structure is a viable endophenotype linking genes to behavior. However, to what extend

local brain structure has a genetic association with cognitive and emotional functioning is

incompletely understood. Here, we study the genetic correlation between personality

traits and local cortical structure in a large-scale twin sample (Human Connectome

Project, n=1106, 22-37y). We found a genetic overlap between personality traits and

local cortical structure in 10 of 17 observed phenotypic associations in predominantly

frontal cortices. To evaluate the robustness of observed personality-brain associations we

studied two independent age-matched samples (GSP: n=926, age=19-35y, eNKI: n=210,

age: 19-39y). We observed anecdotal to moderate evidence for a successful replication of

the negative association between surface area in medial prefrontal cortex and

Neuroticism in both samples. Quantitative functional decoding indicated this region is

implicated in emotional and socio-cognitive functional processes. In sum, our

observations suggest that associations between local brain structure and personality are,

in part, under genetic control. However, associations are weak and only the relation

between frontal surface area and Neuroticism was consistently observed across three

independent samples of young adults.

Personality and local brain structure

Introduction

The local macro-anatomical structure of the cerebral cortex is largely heritable, and has a

highly polygenetic architecture (Grasby et al., 2020; Panizzon et al., 2009; Strike et al.,

2019; Winkler et al., 2010). Recently, it has been shown that common genetic variants

that influence surface area also affect various behavioral traits, suggesting that brain

structure is an essential endophenotype linking genes and behavior (Grasby et al., 2020).

However, to what extend the correlation between local cortical structure on the one hand

and cognitive and emotional functioning on the other is driven by shared genetic factors

is incompletely understood.

One of the most broadly used summaries of an individual's characteristic patterns of

behavior, thought, and emotions is personality (Funder, 2001). Behavioral science

establishes personality structure by parcellating the individual variability in goals,

cognition, and emotion into independent components (Mischel, 2004). A widely used

personality taxonomy is the Big Five Personality inventory (John and Srivastava, 1999;

McCrae and Costa, 1997; Saucier and Srivastava, 2015). The Five-factor personality

structure derives five orthogonal dimensions or traits of Agreeableness,

Conscientiousness, Extraversion, Neuroticism, and Openness (John et al., 2008; Saucier

and Srivastava, 2015). Personality traits have been related to the quality of social

relationships (Asendorpf and Wilpers, 1998), job performance (Rothmann and Coetzer,

2003), risk for mental disorders (Miller et al., 2001; Trull, 2013), general health and

wellbeing, and reproductive success (Alvergne et al., 2010; Strickhouser et al., 2017).

Personality has both stable and malleable features (Damian et al., 2019; Harris et al.,

2016; Penke and Jokela, 2016) and has been found heritable with approximately 40% of

the variance attributable to additive genetic factors (Bouchard, 1994; Bouchard and

Loehlin, 2001; Bouchard and McGue, 2003; Jang et al., 1996; Loehlin et al., 1998;

Vukasovic and Bratko, 2015). Evolutionary causes for variability in personality traits

have been suggested to be due to balancing selection, where selection pressures in

different directions affect the same traits enabling adaptation to changing environmental

demands (Penke and Jokela, 2016). Indeed, genome-wide association studies (GWAS)

have reported a large number of genetic variants associated with personality traits with

each contributing to the heritability of personality (Consortium, 2015; de Moor et al.,

2012; Lo et al., 2017; van den Berg et al., 2016; Verweij et al., 2012).

The biological basis of personality in humans has also been studied in relation to

macroscale brain structure and function using magnetic resonance imaging (MRI)

(Bjornebekk et al., 2013; DeYoung et al., 2010; Dubois et al., 2018; Ferschmann et al.,

2018; Kong et al., 2019; Nostro et al., 2017; Riccelli et al., 2017; Wu et al., 2019).

Various studies have reported a phenotypic relationship between local brain structure and

personality traits (Bjornebekk et al., 2013; DeYoung et al., 2010; Gray et al., 2019;

Nostro et al., 2017; Riccelli et al., 2017). Using the Human Connectome Project, young

adult (HCP) sample, including monozygotic and dizygotic twins, Owens and colleagues

(Owens et al., 2019) report significant phenotypic relationships between personality traits

and various markers of cortical structure. For example, Owens and colleagues observed

associations between Agreeableness, Conscientiousness, Neuroticism, and Openness with

morphometry in prefrontal areas. However findings on the relationship between

personality traits and local brain structure have been inconsistent, for instance, Avinun

and colleagues failed to observe significant relations between personality and various

markers of brain structure using the largest sample to date (Avinun et al., bioarXiv). In

line with this report, Kharabian et al. have recently shown that, in general, relationships

between local brain structure and psychometric variables are not robust and highly

dependent on sample and effect size (Kharabian Masouleh et al., 2020; Kharabian

Masouleh et al., 2019). At the same time, it has recently been shown that traits such as

neuroticism, general cognitive function, educational attainment, and depressive

symptoms show a genetic correlation with total surface area, suggesting brain structure is

a key phenotype reflecting individual differences in behavior (Grasby et al., 2020).

Taken together contemporary theory suggests that (a) individual variation in both local

brain structure and personality can be, in part, attributed to genetic effects (b) brain

structure is a viable endophenotype linking genes and behavior (c) personality relates to

macro-scale brain structure and function, but local relationships are weak and vary as a

function of sample and effect size. However, whether regional brain structure and

personality have a shared genetic basis remains unclear. To answer our research question,

we studied the relationship between the Big Five personality traits and local cortical

thickness and surface area. We captured variations in brain morphometry using an atlas-

based approach, dividing the cortex in 200 functionally-defined parcels (Eickhoff et al.,

2018; Schaefer et al., 2018). We studied three independent samples of young adults, the

Human Connectome Project Young Adult (HCP, n=1106), Brain Genomics

superstructure project (GSP, n=926) and enhanced Nathan Kline Institute dataset (eNKI,

n=210). The HCP sample is unique in that it provided us with high quality neuroimaging

and personality trait (NEO-FFI) data in a large number of twins, siblings, and unrelated

individuals, enabling us to compute genetic correlation between personality and local

brain structure. Analysis of heritability and genetic correlation was performed using

maximum likelihood variance-decomposition methods using Sequential Oligogenic

Linkage Analysis Routines (www.solar-eclipse-genetics.org; Solar Eclipse 8.4.0.).

Second, to assess whether observed associations between personality and local brain

structure in the HCP sample reflect generalizable relationships between personality and

local brain structure, we selected two samples (Brain Genomics superstructure project –

GSP (n=926) and enhanced Nathan Kline Institute dataset – eNKI (n=210)) of unrelated

individuals between 20 and 40 years of age in which we studied whether personality-

brain relationships observed in the HCP sample would replicate in two independent

samples. Last, we performed functional decoding to further evaluate the functional

mechanisms underlying brain regions robustly associated with personality.

Personality and local brain structure

Materials and methods

<u>HCP sample:</u>

Participants and study design

For our analysis we used the publicly available data from the Human Connectome Project

S1200 release (HCP; http://www.humanconnectome.org/), which comprised data from

1206 individuals (656 females), 298 MZ twins, 188 DZ twins, and 720 singletons, with

mean age 28.8 years (SD = 3.7, min-max = 22–37). We included individuals for whom

the scans and data had been released (humanconnectome.org) after passing the HCP

quality control and assurance standards (Marcus et al., 2013). The full set of inclusion

and exclusion criteria are described elsewhere (Glasser et al., 2013; Van Essen et al.,

2013). In short, the primary participant pool comes from healthy individuals born in

Missouri to families that include twins, based on data from the Missouri Department of

Health and Senior Services Bureau of Vital Records. Additional recruiting efforts were

used to ensure participants broadly reflect ethnic and racial composition of the U.S.

population. Healthy is broadly defined, in order to gain a sample generally representative

of the population at large. Sibships with individuals having severe neurodevelopmental

disorders (e.g., autism), documented neuropsychiatric disorders (e.g. schizophrenia or

depression) or neurologic disorders (e.g. Parkinson's disease) are excluded, as well as

individuals with diabetes or high blood pressure. Twins born prior 34 weeks of gestation

and non-twins born prior 37 weeks of gestation are excluded as well. After removing

individuals with missing structural imaging or behavioral data our sample consisted of

1106 individuals (including 286 MZ-twins and 170 DZ-twins) with mean age of 28.8

years (SD =3.7, min-max =22-37). See further Table 1.

Personality and local brain structure

Structural imaging processing

MRI protocols of the HCP are previously described (Glasser et al., 2013; Van Essen et

al., 2013). The pipeline used to obtain the Freesurfer-segmentation is described in detail

in a previous article (Glasser et al., 2013) and is recommended for the HCP-data. In short,

the pre-processing steps included co-registration of T1 and T2 scans, B1 (bias field)

correction, and segmentation and surface reconstruction to estimate cortical thickness.

The HCP structural pipelines use Freesurfer 5.1 software

(http://surfer.nmr.mgh.harvard.edu/) (Dale et al., 1999; Fischl, 2013; Fischl and Dale,

2000; Fischl et al., 1999) plus a series of customized steps that combine information from

T_1w as well as T_2w scans for more accurate white and pial surfaces (Glasser et al., 2013).

Next, the individual cortical thickness and surface area maps were standardized to

fsaverage5 for further analysis. Segmentation maps were visually inspected (S.L.V.) for

inaccuracies.

Five Factor Model of Personality

The Big Five personality traits were assessed using the NEO-Five-Factors-Inventory

(NEO-FFI)(McCrae and Costa, 2004). The NEO-FFI is composed of a subset of 60-items

extracted from the full-length 240-item NEO-PI-R. For each item, participants reported

their level of agreement on a 5-point Likert scale, from strongly disagree to strongly

agree. The NEO instruments have been previously validated in USA and several other

countries (McCrae and Terracciano, 2005). See further **Table 1**.

As a proxy for IQ we used the NIH Toolbox Cognition (Weintraub et al., 2013), 'total

composite score'. The Cognitive Function Composite score is derived by averaging the

normalized scores of each of the Fluid and Crystallized cognition measures, then deriving

scale scores based on this new distribution. Higher scores indicate higher levels of

cognitive functioning. Participant score is normed to those in the entire NIH Toolbox

Normative Sample (18 and older), regardless of age or any other variable, where a score

of 100 indicates performance that was at the national average and a score of 115 or 85,

indicates performance 1 SD above or below the national average. See further **Table 1**.

Measure	n	mean±SD (min-max)
Males/Females	506/600	-
Age	1106	28.8±3.7 (22-36)
Intelligence (Composite score)	1089	121.9±14.6 (84.6-153.4)
Agreeableness	1106	33.5±5.8 (10-48)
Conscientiousness	1106	34.5±5.9 (11-48)
Extraversion	1106	30.7±6 (10-47)
Neuroticism	1106	16.6±7.4 (0-43)
Openness	1106	28.3±6.2 (10-47)

180   **Table 1. Behavioral characteristics of the HCP sample.** Behavioral characteristics for
181   gender, age, intelligence as well as the NEO-FFI scores in the HCP sample.
182

183   GSP sample

184
Participants and study design

To evaluate the cross-sample reproducibility of observations we additionally investigated

the association between personality and local cortical brain structure in the Brain

Genomics Superstruct Project (GSP) (Holmes et al., 2015). In short, between 2008 and

2012 young adults (ages 18 to 35) with normal or corrected-to-normal vision were

recruited from the Boston community to participate in the GSP. The 1,570 individuals

included in the data release (Holmes et al., 2015) were selected from a larger databased

of individuals who participated in the ongoing GSP data collection initiative. Participants

included well-educated individuals with relatively high IQs (many of the college age

students are from local colleges). Participants provided written informed consent in

accordance with guidelines established by the Partners Health Care Institutional Review

Board and the Harvard University Committee on the Use of Human Subjects in Research

(See Supplementary Appendix A in (Holmes et al., 2015)).

Structural imaging processing

All imaging data were collected on matched 3T Tim Trio scanners (Siemens Healthcare,

Erlangen, Germany) at Harvard University and Massachusetts General Hospital using the

vendor-supplied 12-channel phased-array head coil. Structural data included a high-

resolution (1.2mm isotropic) multi-echo T1-weighted magnetization-prepared gradient-

echo image (multi-echo MPRAGE, see further: (Holmes et al., 2015)). The low

participant burden resulting from the use of multi-echo MPRAGE anatomical scans

makes this sequence well suited for high-throughput studies. The morphometric features

derived through conventional 6-min 1mm MPRAGE and the 2-min 1.2mm multi-echo

MPRAGE are highly consistent (r2>0.9 for most structures) suggesting that rapid

acquisition multi-echo MPRAGE can be used for many purposes in place of longer

anatomical scans without degradation of the quantitative morphometric estimates. All T1

scans   pre-processed   using   the   Freesurfer   software   library

(http://surfer.nmr.mgh.harvard.edu/) version 6.0.0 (Dale et al., 1999; Fischl, 2013; Fischl

and Dale, 2000; Fischl et al., 1999). Next, the individual cortical thickness and surface

area maps were standardized to fsaverage5 for further analysis. Segmentation maps were

visually inspected (S.L.V.) for inaccuracies and 1 individual was excluded due to

regional abnormalities.

Five Factor Model of Personality

The Big Five personality traits were assessed using the full-length 240-item Revised

NEO Personality Inventory NEO-Five-Factors-Inventory (NEO-PI-R)(Costa and

McCrae, 1992), the full-length 240-item NEO-PI-R. For each item, participants reported

their level of agreement on a 5-point Likert scale, from strongly disagree to strongly

agree. The NEO instruments have been previously validated in USA and several other

countries (McCrae and Terracciano, 2005). As a proxy for IQ we used the estimated IQ

derived through the Oklahoma Premorbid Intelligence Estimate--3 (OPIE3) formula

(Schoenberg et al., 2002). Reported values are in integers and binned. It is of note that

distribution of IQ values is positively skewed relative to the general population and that

many personality traits, including negative affect and Neuroticism were observed to have

distribution that would be expected of a clinically-screened population-based sample

(Holmes et al., 2015). See further **Table 2**.

Measure	n	mean±SD (min-max)
Males/Females	536/390	-
Age	926	21.6±3.9 (19-35)
Estimated IQ	891	108.7±8.1 (77-129)
Agreeableness	926	32.0±6.6 (9-47)
Conscientiousness	926	31.7±7.2 (8-48)
Extraversion	926	30.7±6.5 (9-48)
Neuroticism	926	20.3±8.8 (0-48)
Openness	926	31.6±6.1 (14-47)

232   **Table 2. Behavioral characteristics of the GSP sample.** Behavioral characteristics for
233   gender, age, intelligence as well as the NEO-FFI scores in the GSP sample.
234

eNKI sample:

Participants and study design

To evaluate the cross-sample reproducibility of observations we additionally investigated

correspondence between personality and cortical brain structure in the eNKI where we

selected adults between 20 and 40 years of age to match the age-range of the HCP and

GSP samples. The sample was made available by the Nathan-Kline Institute (NKY, NY,

USA), as part of the '*enhanced NKI-Rockland sample*'

(https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3472598/). In short, eNKI was designed

to yield a community-ascertained, lifespan sample in which age, ethnicity, and

socioeconomic status are representative of Rockland County, New York, U.S.A. ZIP-

code based recruitment and enrollments efforts were being used to avoid over-

representation of any portion of the community. Participants below 6 years were

excluded to balance data losses with scientific yield, as well as participants above the age

of 85, as chronic illness was observed to dramatically increase after this age. All

approvals regarding human subjects' studies were sought following NKI procedures.

Scans were acquired from the International Neuroimaging Data Sharing Initiative (INDI)

online database http://fcon_1000.projects.nitrc.org/indi/enhanced/studies.html For our

phenotypic analyses, we selected individuals with complete personality and imaging data

within the age-range of 18-40 years to match the age-range of the HCP and GSP samples.

Our sample for phenotypic correlations consisted of 210 (121 females) individuals with

mean age of 26.0 years (SD =6.1, min-max =18-39). Please see **Table 3** for demographic

characteristics.

Structural imaging processing

3D magnetization-prepared rapid gradient-echo imaging (3D MP-RAGE) structural

scans(Mugler and Brookeman, 1990) were acquired using a 3.0☐T Siemens Trio scanner

with TR=2500☐ms, TE=3.5☐ms, Bandwidth=190☐Hz/Px, field of view=256 ×

256☐mm, flip angle=8°, voxel size=1.0 × 1.0 × 1.0☐mm. More details on image

acquisition are available at http://fcon_1000.projects.nitrc.org/indi/enhanced/studies.html.

All T1 scans were pre-processed using the Freesurfer software library

(http://surfer.nmr.mgh.harvard.edu/) version 6.0.0 (Dale et al., 1999; Fischl, 2013; Fischl

and Dale, 2000; Fischl et al., 1999) to compute cortical thickness and surface area. Next,

the individual cortical thickness and surface area maps were standardized to fsaverage5

for further analysis. Segmentations were visually inspected for anatomical errors

(S.L.V.).

Five Factor Model of Personality

The Big Five personality traits were assessed using the NEO-FFI3(McCrae and Costa,

2004; McCrae and Terracciano, 2005).

For an assessment of intelligence we used the Wechsler Abbreviated Scale of Intelligence

(WASI-II)(Wechsler, 1999), full scale IQ. The WASI is a general intelligence, or IQ test

designed to assess specific and overall cognitive capabilities and is individually

administered to children, adolescents and adults (ages 6-89). It is a battery of four

subtests: Vocabulary (31-item), Block Design (13-item), Similarities (24-item) and

Matrix Reasoning (30-item). In addition to assessing general, or Full Scale, intelligence,

the WASI is also designed to provide estimates of Verbal and Performance intelligence

consistent with other Wechsler tests. Specifically, the four subtests comprise the full scale

and yield the Full Scale IQ (FSIQ-4), see further **Table 3**.

Measure	n	mean±SD (min-max)
Males/Females	121/210	-
Age	210	26.0±6.1 (18-39)
Intelligence (WASI)	210	100.3±12.3 (69-135)
Agreeableness	210	33.6±6.1 (18-48)
Conscientiousness	210	33.9±7.2 (13-48)
Extraversion	210	30.5±6.3 (7-44)
Neuroticism	210	19.7±8.1(2-42)
Openness	210	33.0±6.3 (12-48)

284   **Table 3. Behavioral characteristics of the eNKI sample**. Behavioral characteristics for
285   gender, age, intelligence as well as the NEO-FFI scores in the eNKI sample.
286

Parcellation approach

In all three samples, we used a parcellation scheme (Schaefer et al., 2018) based on the

combination of a local gradient approach and a global similarity approach using gradient-

weighted Markov Random models. The parcellation has been extensively evaluated with

regards to stability and convergence with histological mapping and alternative

parcellations. In the context of the current study, we focus on the granularity of 200

parcels, as averaging will improve signal-to-noise ratio. In order to improve signal-to-

noise ratio and to accelerate analysis speed, we opted to average unsmoothed structural

data within each parcel. Thus, cortical thickness of each ROI was estimated as the

trimmed mean (10 percent trim) and surface area as the sum of area within an ROI.

Phenotypic correlation analysis.

As in previous structural MRI analyses (Bernhardt et al., 2014; Valk et al., 2016a; Valk et

al., 2016b), we used SurfStat for Matlab [R2017a, The Mathworks, Natick, MA](Worsley

et al., 2009). Phenotypic correlation analyses between personality traits and local brain

structure were carried out per parcel, using a 200 parcel-parcellation scheme (Schaefer et

al., 2018) on surface area and cortical thickness. We controlled for the same variables as

in the genetic analysis, namely age, sex, age $\times$ sex interaction, age^2, age$^2 \times$ sex

interaction, as well as global thickness effects when investigating cortical thickness and

intracranial volume when assessing surface area. Results were corrected for multiple

comparisons using Benjamini-Hochberg FDR (Benjamini and Hochberg, 1995) at whole-

brain analysis, where we corrected for number of analysis within the current step and

report FDR thresholds. When investigating personality or in post-hoc brain analysis, we

corrected for number of analysis x ROIs. Post-hoc we also controlled for a proxy for

intelligence, total cognitive score (Weintraub et al., 2013). We displayed significant

(FDRq<0.05) findings on the brain surface.

Heritability and genetic correlation analysis

To investigate the heritability and genetic correlation of brain structure and personality

traits, we analyzed 200 parcels of cortical thickness and surface area, as well as

personality trait score of each subject in a twin-based heritability analysis. As in previous

studies (Glahn et al., 2010), the quantitative genetic analyses were conducted using

Sequential Oligogenic Linkage Analysis Routines (SOLAR) (Almasy and Blangero,

1998). SOLAR uses maximum likelihood variance-decomposition methods to determine

the relative importance of familial and environmental influences on a phenotype by

modeling the covariance among family members as a function of genetic proximity. This

approach can handle pedigrees of arbitrary size and complexity and thus, is optimally

efficient with regard to extracting maximal genetic information. To ensure that our traits,

behavioral as well as of brain structure, were conform to the assumptions of normality, an

inverse normal transformation was applied for all behavioral and neuroimaging traits

(Glahn et al., 2010).

Heritability (h^2) represents the portion of the phenotypic variance (σ^2_p) accounted for by

the total additive genetic variance (σ^2_g), i.e., $h^2 = \sigma^2_g/\sigma^2_p$. Phenotypes exhibiting stronger

covariances between genetically more similar individuals than between genetically less

similar individuals have higher heritability. Heritability analyses were conducted with

simultaneous estimation for the effects of potential covariates. For this study, we included

covariates including age, sex, age $\times$ sex interaction, age^2, age^2 $\times$ sex interaction. Post-hoc

we also controlled for a proxy for intelligence, total cognitive score(Weintraub et al.,

2013). When investigating cortical thickness, we additionally controlled for global

thickness effects (mean cortical thickness) and in case of surface area we controlled for

intracranial volume.

To determine if variations in personality and brain structure were influenced by the same

genetic factors, genetic correlation analyses were conducted. More formally, bivariate

polygenic analyses were performed to estimate genetic (ρ_g) and environmental (ρ_e)

correlations, based on the phenotypic correlation (ρ_p), between brain structure and

personality with the following formula: $\rho_p = \rho_g\sqrt{(h^2_1 h^2_2)} + \rho_e\sqrt{[(1 - h^2_1)(1 - h^2_2)]}$,

where h^2_1 and h^2_2 are the heritability's of the parcel-based cortical thickness and the

various behavioral traits. The significance of these correlations was tested by comparing

the log likelihood for two restricted models (with either ρ_g or ρ_e constrained to be equal to

0) against the log likelihood for the model in which these parameters were estimated. A

significant genetic correlation (corrected for multiple comparisons using Benjamin-

Hochberg FDR (Benjamini and Hochberg, 1995)) is evidence suggesting that (a

proportion of) both phenotypes are influenced by a gene or set of genes (Almasy et al.,

1997).

To compute the contribution of genetic effects relative to the phenotypic correlation, we

computed the contribution of the genetic path to the phenotypic correlation ($\sqrt{h^2_1} \times \rho_g \times$

$\sqrt{h^2_2}$) ($\rho_{ph}g$) divided by the phenotypic correlation. For the relative contribution of

environmental correlation to the phenotypic correlation we computed ($\sqrt{1-h^2_1} \times \rho_e \times \sqrt{1-}$

h^2_2) ($\rho_{ph}e$) divided by the phenotypic correlation (Zheng et al., 2019).

Bayes Factors of replication

To compare the evidence that the personality-local brain structure could be replicated in

two independent samples (H1, replication, and H0, no replication), we additionally

quantified personality-brain associations within each ROI, using Bayes factors (Verhagen

and Wagenmakers, 2014). In line with previous work of our group (Kharabian Masouleh

et al., 2020; Kharabian Masouleh et al., 2019) Bayes factors (BF) were summarized into

four categories. These categories are used to simplify the interpretation and comparison

of replication rates. For example, a BF_{01} lower than 1/3 shows that the data is three times

or more likely to have happened under H1 than H0. "Successful" replication is defined as

a replication lower than 1 in both replication samples.

Personality and local brain structure

Functional decoding

Parcel that were significantly replicated in at least one sample were functionally

characterized using the Behavioral Domain meta-data from the BrainMap database

(http://www.brainmap.org (Laird et al., 2011; Laird et al., 2009)). To do so, volumetric

counterparts, delineating the surface-based parcels in volume space, as provided by

Schaefer, (Schaefer et al., 2018)

(https://github.com/ThomasYeoLab/CBIG/tree/master/stable_projects/brain_parcellation/

Schaefer2018_LocalGlobal/Parcellations) were used. In particular, we identified those

meta-data labels (describing the computed contrast [behavioral domain as well as

paradigm]) that were significantly more likely than chance to result in activation of a

given parcel (Fox et al., 2014; Genon et al., 2018; Nostro et al., 2017). That is, functions

were attributed to the parcels by quantitatively determining which types of experiments

are associated with activation in the respective parcellation region. Significance was

established using a binomial test ($q < 0.05$, corrected for multiple comparisons using false

discovery rate, FDR).

Personality and local brain structure

Results

Association between personality traits and cortical brain structure.

To assess the association between personality and macroscale cortical brain structure we

first evaluated distribution of behavioral measures. Using the Kolmogorov-Smirnov test

we found that all personality traits in the HCP sample (n=1106 including 286 MZ-twins

and 170 DZ-twins) were conform to normal distributions (KS-score between 0.97 and 1)

(**Figure 1**). We observed significant phenotypic correlation between all personality traits,

with the exception of Openness and Neuroticism (r =0.01) (**Figure 1**, Supplementary

Table 1).

398
399   **Figure 1. Distribution of personality traits in the full HCP sample.** Distribution of
400   NEO-FFI personality traits in the HCP dataset, score on x-axis, number of occurrences on
401   the y-axis, as well as the correlation between NEO-FFI traits in the HCP sample.
402

403   Next, we assessed phenotypic correlation between personality traits and cortical structure,

404   specifically cortical thickness and surface area. Distribution of cortical thickness values

Personality and local brain structure

summarized in 200 functionally informed parcels (Schaefer et al., 2018) showed highest

thickness in anterior insula, and relatively low values in occipital regions (**Figure 2A**).

At the regional level, we observed correlations between Agreeableness, Neuroticism, and

Openness and local cortical thickness (**Figure 2B**). Specifically, Agreeableness related

negatively to variations in thickness in left lateral and bilateral medial prefrontal cortex

(FDRq<0.01). Neuroticism related positively to thickness in dorsolateral frontal areas and

left posterior operculum, and negatively to thickness in left posterior occipital regions

(FDRq<0.015). Openness related negatively to thickness in left ventrolateral cortex, and

positively to right temporal pole (FDRq<0.01). We did not observe significant

associations between mean cortical thickness and personality scores (**Table 4**).

Total surface area had a negative relation with conscientiousness (t=-2.59, p<0.005) and a

positive association with openness (t=2.94, p<0.002) (**Table 4**). Regionally, we found a

negative relation between Neuroticism and local surface area in bilateral medial frontal

cortex, left inferior frontal gyrus, and left posterior insula (FDRq<0.02).

	Average cortical thickness	**Total surface area**
Agreeableness	-0.24	0.66
Conscientiousness	0.55	-2.59**
Extraversion	0.66	1.14
Neuroticism	1.90*	-2.03*
Openness	-1.28	2.94**

419   **Table 4. Association between personality traits and whole brain summaries of**
420   **surface area and cortical thickness in the full HCP sample.** T-values of the association
421   between average cortical thickness and total surface area and personality traits. **
422   indicates FDRq<0.05, * indicates p<0.05.
423

Personality and local brain structure

424
**Figure 2. Relation between personality traits and local brain structure in the full
HCP sample.** A) Mean cortical thickness of each parcel and the distribution of average
cortical thickness across participants; B) Regional associations between personality traits
and cortical thickness; C) Average surface area sum per parcel and the distribution of
total surface area across participants; D) Regional associations between surface area and
personality traits. Positive associations between local brain structure and each personality
trait are displayed in red and negative associations displayed in blue. Multiple
comparisons were accounted for by using FDR corrections at q<0.05 correcting for the
number of parcels (200) and only significant associations are displayed.
434

Personality and local brain structure

To test stability of our findings we additionally evaluated the robustness of phenotypic

associations between personality and global and local brain structure while controlling

for total cognitive score and the other personality traits (Supplementary Materials,

Supplementary Table 2 and 3). While all local associations remained significant at

p<0.01, strength of associations was generally reduced and few regions reached

FDRq<0.05 significance levels.

Genetic relationship between personality traits and cortical brain structure

Subsequently, we sought to evaluate whether the phenotypic correlations observed in the

twin-sample were due to shared genetic or environmental effects on grey matter brain

structure and personality traits. All personality traits were significantly heritable in our

current sample (**Figure 3C**, Supplementary Table 4), as were mean cortical thickness

(h^2=0.85) and total surface area (h^2=0.93), and we confirmed also local cortical thickness

(h^2: mean±std: 0.33±0.11) and surface area (h^2: mean±std: 0.41±0.13) to be heritable in

our parcel-based approach (**Figure 3A-B**, Supplementary Table 5 and 11).

450
451   **Figure 3. Heritability of local cortical structure and personality traits.** A) Heritability
452   of local cortical thickness; B) Heritability of surface area; C) Heritability of NEO-FFI:
453   A= agreeableness, C=conscientiousness, E=extraversion, N=Neuroticism, O=Openness
454
455

Personality and local brain structure

Following, we assessed genetic correlation between personality traits and cortical

structure. We did not observe genetic or environmental associations between personality

and global thickness (**Table 5**), however, the phenotypic association between total

surface area and conscientiousness was observed to be driven by shared genetic effects

(ρ_g =-0.12, p<0.03) whereas the association between openness and total surface area was

driven by environmental effects (ρ_e=0.18, p<0.025).

	Global thickness		Total surface area	
Agreeableness	ρ_e= 0.03, p=ns;	ρ_g= -0.03, p=ns	ρ_e=-0.08, p=ns;	ρ_g =0.07, p=ns
Conscientiousness	ρ_e=-0.02, p=ns;	ρ_g = 0.05, p=ns	**ρ_e=0.04, p=ns;**	**ρ_g =-0.12, p=0.03**
Extraversion	ρ_e=-0.03, p=ns;	ρ_g = 0.06, p=ns	ρ_e=0.03, p=ns;	ρ_g =0.04, p=ns
Neuroticism	**ρ_e=0.10, p=ns;**	**ρ_g = 0.02, p=ns**	ρ_e=-0.09, p=ns;	ρ_g =-0.06, p=ns
Openness	ρ_e=-0.03, p=ns;	ρ_g =-0.03, p=ns	**ρ_e=0.18, p=0.025;**	**ρ_g =0.08, p=ns**

462 **Table 5. Genetic and environmental correlation between personality traits and**
463 **whole brain summaries of surface area and cortical thickness.** Genetic and
464 environmental correlations are computed in the HCP sample, and exact p-values are
465 reported, associations that did not show phenotypic correlation at p<0.05 threshold are in
466 bold.
467

Last, we evaluated the genetic correlation of regions that showed phenotypic correlations

between personality and local brain structure. We observed that 10 out of 17 phenotypic

correlations showed a genetic correlation (p≤0.05), and 2 out of 17 phenotypic correlates

related to an environmental correlation (p≤0.05) (**Table 6**). More specifically, we found a

negative genetic correlation between Agreeableness and bilateral superior frontal

thickness (p<0.04), a positive genetic correlation between Neuroticism and right superior

and lateral frontal cortex thickness (p<0.05) and a positive genetic correlation between

right temporal pole thickness and Openness (p<0.005). Neuroticism had a negative

genetic correlation between local surface area in left posterior insula, and bilateral

superior frontal cortex, and right medial frontal regions (p<0.05). See Supplementary

Personality and local brain structure

478   Tables (6-10 and 12-16) for genetic and environmental correlations between personality

479   traits and all parcels.

Cortical thickness	ROI	Environmental correlation	Genetic correlation	Genetic contribution to phenotypic correlation
Agreeableness	LH Cont_PFCl_4	ρ_e -0.12, p=0.04*	ρ_g -0.08, p=0.61	20 %
	LH Default_PFC_9	ρ_e -0.05, p=0.45	ρ_g -0.21, p=0.08	72%
	LH Default_PFC_11	ρ_e 0.01, p=0.90	ρ_g -0.36, p=0.004**	100%
	LH Default_PFC_13	ρ_e -0.03, p=0.72	ρ_g -0.26, p=0.04*	87%
	LH Default_PFCm_5	ρ_e 0.03, p= 0.64	ρ_g -0.33, p=0.005**	100%
Neuroticism	LH Vis_14	ρ_e -0.08, p=0.22	ρ_g -0.18, p=0.12	61%
	LH Default_PFC_9	ρ_e 0.08, p=0.23	ρ_g 0.17, p=0.13	61%
	RH Cont_PFCl_6	ρ_e 0.03, p=0.63	ρ_g 0.26, p=0.05*	82%
	RH Default_PFCm_5	ρ_e 0.04, p=0.54	ρ_g 0.21, p=0.05*	79%
Openness	LH Cont_PFCl_4	ρ_e -0.13, p= 0.05*	ρ_g -0.15, p= 0.17	46%
	RH Limbic_TempPole_1	ρ_e -0.01, p= 0.94	ρ_g 0.37, p<. 0.005**	100%
Surface area				
Neuroticism	LH SomMot_3	ρ_e 0.02, p=0.78	ρ_g -0.28, p=0.01*	100%
	LH Default_PFC_3	ρ_e -0.04, p=0.52	ρ_g -0.18, p=0.12	74%
	LH Default_PFC_9	ρ_e 0.13, p=0.07	ρ_g -0.45, p=0.0002**	100%
	LH Default_PFC_13	ρ_e -0.03, p=0.60	ρ_g -0.21, p=0.14	76%
	RH Default_PFCm_4	ρ_e 0.02, p=0.80	ρ_g -0.25, p=0.02*	100%
	RH Default_PFCm_5	ρ_e -0.04, p=0.55	ρ_g -0.30, p=0.02*	82%

Table 6. Genetic and environmental correlation of personality brain associations in
the full HCP sample. Genetic and environmental correlations are computed in the HCP
sample, and exact p-values are reported. ** denotes a significant genetic correlation at
FDRq<0.05, corrected for the number of ROIs associated with the respective personality
trait within the structural marker. * indicated an association of p<0.05. The genetic
contribution of phenotypic correlation was computed using the respective heritability of
the personality trait and the local parcel as well as their genetic and phenotypic
correlation.

Cross-sample reproducibility of the association between personality trait and local brain

structure.

In the previous analysis steps, we could show that a) there is a significant relationship

between local cortical structure and personality traits in a large-scale twin sample (HCP)

and that b) this relationship can be, in part, attributed to shared genetic factors.

Following, to study whether associations between local brain structure and personality

traits are generalizable, we evaluate the phenotypic correlation between personality traits

and cortical phenotypes observed in the HPC sample are reproducible in two age-

matched samples of young adults (GSP and eNKI). To formalize the level of

reproducibility, we computed Bayer Factors (BF) summarizing the evidence of a

successful reproduction across samples (Verhagen and Wagenmakers, 2014).

We found moderate to anecdotal evidence of replication for only one personality-brain

association in both samples; the relationship between local surface area in right medial

frontal cortex and Neuroticism (GSP: t=-1.54, p<0.07; BF=0.68; and eNKI: t=-1.99,

p<0.025; BF=0.16). Various associations between local cortical thickness and personality

traits could be reproduced in one of both replication samples (**Table 7**). Specifically, in

GSP, the association between thickness in right superior frontal cortex and Agreeableness

(t=-1.71, p<0.05; BF=0.52), and between thickness of right dorsal lateral PFC and

Neuroticism (t=2.08, p<0.02; BF=0.18). In the eNKI sample we observed some evidence

of successful replication of the association between left visual cortex and Neuroticism

(t=-1.73, p<0.05; BF=0.24), left dorsolateral prefrontal thickness and Openness (t=-0.96,

p>0.1, BF=0.82), surface area of left sensory-motor cortex (t=-1.09, p>0.1, BF=0.58) and

Personality and local brain structure

513   left prefrontal cortex (t=-1.09, p>0.1, BF=0.58) and Neuroticism. Global measures of

514   cortical thickness and surface area did not replicate out of sample, only in case of the

515   positive association between total surface area and Openness we observed anecdotal

516   evidence of successful replication in the eNKI sample (t=1.55, p<0.1, BF=0.33)

517   (Supplementary Table 17).

Cortical thickness	ROI	GSP t-value (BF)	eNKI t-value (BF)
Agreeableness	LH Cont_PFCl_4	1.36 (>3)	-0.01 (>3)
	LH Default_PFC_9	-0.33 (>3)	0.36 (>3)
	LH Default_PFC_11	-0.22 (>3)	-0.31 (>2.5)
	LH Default_PFC_13	-0.44 (>3)	-0.56 (>1)
	LH Default_PFCm_5	-1.71*(0.52)	1.38 (>3)
Neuroticism	LH Vis_14	1.97* (>3)	-1.73* (0.24)
	LH Default_PFC_9	0.92 (>3)	0.72 (>1)
	RH Cont_PFCl_6	2.08*(0.18)	-1.81* (>3)
	RH Default_PFCm_5	-0.15 (>3)	-0.54 (>3)
Openness	LH Cont_PFCl_4	0.70 (>3)	-0.96 (0.82)
	RH Limbic_TempPole_1	0.62 (>3)	0.55 (>1)
Surface area			
Neuroticism	LH SomMot_3	-1.06 (>2)	-1.09 (0.58)
	LH Default_PFC_3	2.32* (>3)	0.39 (>3)
	LH Default_PFC_9	-0.22 (>3)	-0.42 (>1)
	LH Default_PFC_13	-0.13 (>3)	-1.09 (0.58)
	RH Default_PFCm_4	-1.54 (0.68)	-1.99* (0.16)
	RH Default_PFCm_5	0.03 (>3)	-0.44 (>2)

Table 7. Replication of personality brain associations. Replication in the GSP and
eNKI sample of significant associations between personality and local brain structure
observed in the HCP sample, t-values as well as Bayes Factors (BF) are reported. If a
BF_{01} is between 0 and 1/3 there is a moderate/strong evidence for H1 (replication),
between 1/3 and 1 anecdotal evidence for H1, between 1 and 3 anecdotal evidence for H0
(no replication) and >3 moderate to strong evidence of H0. We underlined replications
with a correct sign, and made replications bold if there is evidence
(anecdotal/moderate/strong) of H1. ** indicates a significant correlation at FDRq<0.05, *
is p<0.05.
527

Personality and local brain structure

Quantitative functional decoding.

Last, we performed quantitative functional mapping of the personality – brain

relationships for which we observed a) phenotypic and genetic correlation in the HCP

sample b) an association (p<0.05) in combination with a BF of <1 in at least one

additional sample.

The right medial frontal cortex, where we observed a robust association between surface

area and Neuroticism, was functionally involved in various emotional domains, social

cognition, and memory, and active in paradigms involving self-reflection, Theory of

Mind, and emotion induction (FDRq<0.05) (**Figure 4**).

Figure 4. Quantitative functional decoding of consistent associations between
personality and local brain structure. Both forward inference and reverse inference of
activation-domain and paradigm-domain contrasts are reported for the right medial
frontal cortex which showed evidence of successful replication in two samples.
Discussion

Both local brain structure and personality are heritable. Moreover, a large body of

evidence has suggested a relationship between personality and local brain structure.

However, effects are weak and vary as a function of sample and effect size. In the current

study, we used the large scale and openly available HCP dataset which included

monozygotic and dizygotic twins to study whether there is a genetic correlation between

local brain structure and personality traits. Second, we evaluated the robustness of

personality-brain relationships in two additional age-matched samples.

First, we identified phenotypic associations between personality traits and local cortical

structure. Associations between personality on the one hand and cortical thickness and

surface area on the other were predominantly observed in frontal cortices. Performing

genetic correlation analysis, we found that 10 of 17 phenotypic associations could be

explained by shared genetic effects. To evaluate whether observed relationship between

personality traits and local brain structure were generalizable to other samples, we

additionally studied phenotypic correlations between personality traits and brain structure

in two independent age-matched samples of unrelated individuals (GSP and eNKI). Here,

we found that surface area in right medial prefrontal cortex was robustly associated with

Neuroticism across all three samples. In sum, our findings suggest that part of phenotypic

associations between personality and local brain structure can be attributed to shared

genetic effects in a large-scale twin sample. However, associations were weak and the

association between surface area in right medial prefrontal cortex and Neuroticism

replicated in two independent samples.

We assessed the genetic basis of the association between personality and cortical

thickness using compressed surface-based MRI data based on the parcellation scheme of

Schaefer et al. (2018). Using compressed features of structural MRI has been suggested

to both improve signal-to-noise ratio of brain measures (cf. (Eickhoff et al., 2018) and

(Genon et al., 2018)), and optimize analysis scalability. The Schaefer parcellation is

derived using functional MRI data from ~1500 subjects and integrates local approaches

detecting abrupt transitions in functional connectivity patterns and global approaches that

cluster similar functional connectivity patterns (Schaefer et al., 2018). Indeed, a

combination of within-area micro circuitry, proxied by brain morphometry, and between-

area connectivity enables each area to perform a unique set of computations (Van Essen

and Glasser, 2018). Therefore, a parcellation approach that considers both local and

global connectivity might benefit structural image analysis, as it reduces signal-to-noise

both within and across individuals and makes control for multiple comparisons more

straightforward (Genon et al., 2018). Based on the findings in our study, we suggest our

approach might be a fruitful first exploratory step to investigate the genetic relation

between brain structure and behavior, and locate mechanisms of interest. Future studies

can subsequently verify these results by exploring more specific genetic mechanisms, as

well as neuroanatomical features.

Though we could establish phenotypic correlations between personality traits and local

cortical thickness, associations were weak and phenotypic associations ranged between t-

values of 3.5 and -3.5. In the HCP dataset phenotypic correlations between

predominantly frontal regions and personality traits of Agreeableness, Neuroticism, and

Openness have been previously reported using a non-parcel-based method by Owens and

colleagues (Owens et al., 2019). Frontal cortices are functionally involved in a number of

tasks involving higher cognitive functioning, such as executive functioning, memory,

metacognition and social cognition (Amodio and Frith, 2006; Baird et al., 2013; Bludau

et al., 2014; Buckner et al., 2008; Fleming and Dolan, 2012; Valk et al., 2016a). We

additionally observed various relationships between global measures of surface area and

cortical thickness on the one hand and personality traits on the other. Indeed, we could

replicate a recently reported association between total surface area and Neuroticism in

phenotypic correlation analysis (Grasby et al., 2020). However, associations between

global measures of cortical structure and personality were not consistent across samples.

We extend previously reported phenotypic observations by showing that these phenotypic

relationships between personality and local cortical structure are driven, in part, by shared

additive genetic effects rather than environmental factors alone. The contribution of

genetic effects on phenotypic correlations is dependent on the heritability of each of the

correlated markers. In our sample, between 30 % and 60 % (on average 42 %) of variance

in personality traits was explained by additive genetic factors. This is in line with

previous studies using twin and family samples (Jang et al., 1996) as well as genome-

wide approaches (Lo et al., 2017). A recent meta-analysis (Vukasovic and Bratko, 2015)

confirmed that on average 40 % of the variance in personality traits is of genetic origin.

Also, conform with previous studies (Eyler et al., 2012; Kremen et al., 2010; Panizzon et

al., 2009; Strike et al., 2018; Winkler et al., 2010), we observe heritability of local

cortical thickness, with highest values in primary sensory areas. Heritability patterns

followed previously described patterns with relatively strong genetic influence on cortical

thickness in unimodal cortices, whereas variance in association cortices is on average less

influenced by genetic factors (Eyler et al., 2012; Grasby et al., 2020; Hofer and al., 2018;

Kremen et al., 2010; Panizzon et al., 2009; Strike et al., 2018; Winkler et al., 2010). Also,

local surface area was heritable, with lowest heritability values in dorsolateral PFC and

temporal-parietal regions (Eyler et al., 2012; Grasby et al., 2020; Hofer and al., 2018;

Kremen et al., 2010; Panizzon et al., 2009; Strike et al., 2018; Winkler et al., 2010).

Performing genetic correlation analysis, we observed that the phenotypic correlation

between personality and local brain structure in 10 out of 17 regions was driven by

genetic factors. These regions were predominantly located in frontal areas, suggesting a

genetic link between local structure in frontal cortices and personality. Indeed, various

studies have suggested a relationship between personality and the frontal lobe in humans

(DeYoung et al., 2010; Owens et al., 2019; Riccelli et al., 2017) and in chimpanzees

(Latzman et al., 2015). There are various ways in which a genetic process would affect

the relationship between personality and cortical macrostructure and it is likely the

observed genetic correlations between local cortical structure and personality traits in the

HCP sample are be due to mediated pleiotropy (a gene affects A which affects B). On

the one hand, it could be a genetic factor affects grey matter macrostructure and

associated function and, as a consequence, personality. On the other hand, it could be that

genetic variation affects brain function which in turn modulates both macroscale structure

Personality and local brain structure

as well as personality, or a genetic mechanism affects an unknown biological factor

which in turn affects personality and brain structure. Recent work using GWAS and

genetic correlation in a large sample of individuals could found a genetic association

between cortical brain structure and various markers of behavior, providing first evidence

of a direct link of genes and behavior via cortical brain structure (Grasby et al., 2020).

Here, Grasby et al found evidence that genetic regulatory elements influencing local

surface area and local cortical thickness stem from different devepemental mechanisms.

Whereas surface area is associated with genetic variants active during fetal development,

cortical thickness may reflect genetic processes underlying myelination, branching and

pruning. Such differential mechanistic timing effects on cortical structure might

contribute to the understanding of which biological mechanisms underlie personality, and

further dissociate factors that shape personality across the life-span.

As various studies have indicated relationships between local brain structure and

psychometric variables are not robust (Avinun et al., bioarXiv; Kharabian Masouleh et

al., 2020; Kharabian Masouleh et al., 2019), we further evaluated the robustness of

phenotypic associations between personality and local brain structure in two age-matched

samples of unrelated individuals. Indeed, though most associations did not replicate

across all three samples, the association between medial prefrontal surface area and

Neuroticism was observed in all three samples. Functional decoding indicated that this

region is functionally involved in (social)-cognitive and emotional processing.

Additionally, we found anecdotal to moderate evidence for successful replication of

various associations cortical thickness and personality in either GSP or eNKI sample.

However, given the inconsistency across samples, these replications are challenging to

interpret.

Limitations and outlook

Moving forward, there are various limitations and challenges in operationalizing

personality that might have resulted in a lack of consistent findings across samples.

Though our samples all were from WEIRD (Western, educated, industrialized, rich, and

democratic) populations (Laajaj et al., 2019), it might be that personality traits probed are

not comparable across samples due to challenges to reliably operationalize personality,

and that confounding environmental and noise effects vary across samples. For example,

it is possible inconsistent or lack of findings with regard to macroscale neuroanatomical

associations of personality may be a function of the assessment of personality used (in

this case, the NEO-FFI/NEO-PI-R) rather than a true null or unreliable finding (Avinun et

al., bioarXiv). The five-factor personality model and the subsequent operationalizations

in instruments such as the NEO are based on a lexical approach. Though such an

approach might be able to dissociate various personality traits, it is debated whether

lexical taxonomy has a direct relation to neurobiology (Perkins et al., 2020; Yarkoni,

2015). Future studies might benefit from using personality instruments developed in

concordance with brain structure and function such as Hierarchical Taxonomy of

Psychopathology (HiTOP) (Perkins et al., 2020).

Second, a recent review on the neurobiology of personality suggested that rather than

focusing on a one-to-one mapping between personality and neurobiology, as done in the

current study, studies that seek to identify mechanisms contributing to particular clusters

of behaviors might be a more fruitful approach to capture the neurobiological

mechanisms underlying personality traits (Yarkoni, 2015). For example, though brain

structure is a viable endophenotype of personality, correlation between personality and

macro-scale cortical structure is weak. Thus, further study of the relationship between

personality and functional activity and functional dynamics might further contribute to

understanding the biological basis of personality and other complex traits (Dubois et al.,

2018; Kebets et al., 2019; Kong et al., 2019; Wu et al., 2019).

Third, only 40% of personality variance in the current sample could attributed to genetic

effects. Environment, such as family environment, peer-groups, and stress have been

reported to influence personality (Hopwood et al., 2011; Nakao et al., 2000), and also

local cortical structure and associated behavior has been reported to change as a

consequence of changing environments in adulthood (Valk et al., 2017). Though genetic

and gene by environment effects are not to be excluded in this context, is likely such

environmental mechanisms further shape the relation between personality traits and brain

structure, above and beyond direct additive genetic effects. Longitudinal designs might

help to further understand the environmental relationship between personality and brain

structure and function.

Taken together, in the current study we report evidence of a shared genetic basis of

personality traits and local brain structure within the HCP sample, and a robust

association of local surface area in medial prefrontal regions and Neuroticism across

three independent samples. It is of note that our study on the shared genetic basis of

personality and brain structure was made possible by the open HCP, GSP, and eNKI

Personality and local brain structure

neuroimaging repositories. These initiatives offer cognitive neuroimaging communities

an unparalleled access to large datasets for the investigation of the brain basis of

individual difference. They have also enabled us to highlight variability across samples

and validation experiments to verify stability of our observations. Notably, the use of

multiple datasets enabled us to test robustness of our findings. Given that reproducibility

is increasingly important, our study illustrates the advantages of open data to increase

understanding of complex traits.

Acknowledgements

We would like to thank the various contributors to the open access databases that our data

was downloaded from. Specifically; HCP data were provided by the Human Connectome

Project, Washington University, the University of Minnesota, and Oxford University

Consortium (Principal Investigators: David Van Essen and Kamil

Ugurbil;1U54MH091657) funded by the 16 NIH Institutes and Centers that support the

NIH Blueprint for Neuroscience Research; and by the McDonnell Center for Systems

Neuroscience at Washington University. GSP data were provided by the Brain Genomics

Superstruct Project of Harvard University and the Massachusetts General Hospital,

(Principal Investigators: Randy Buckner, Joshua Roffman, and Jordan Smoller), with

support from the Center for Brain Science Neuroinformatics Research Group, the

Athinoula A. Martinos Center for Biomedical Imaging, and the Center for Human

Genetic Research. 20 individual investigators at Harvard and MGH generously

contributed data to the overall project.

For enhanced NKI, we would like to thank the principal support for the enhanced NKI-

RS project is provided by the NIMH BRAINS R01MH094639-01 (PI Milham). Funding

for key personnel was also provided in part by the New York State Office of Mental

Health and Research Foundation for Mental Hygiene. Funding for the decompression and

augmentation of administrative and phenotypic protocols provided by a grant from the

Child Mind Institute (1FDN2012-1). Additional personnel support provided by the

Center for the Developing Brain at the Child Mind Institute, as well as NIMH

R01MH081218, R01MH083246, and R21MH084126. Project support also provided by

the NKI Center for Advanced Brain Imaging (CABI), the Brain Research Foundation

Personality and local brain structure

731   (Chicago, IL), and the Stavros Niarchos Foundation. Last, we want to thank L.C.Valk for

732   proofreading the manuscript.

Personality and local brain structure

734 **References**
735
Almasy, L., Blangero, J., 1998. Multipoint quantitative-trait linkage analysis in general
pedigrees. Am J Hum Genet 62, 1198-1211.
Almasy, L., Dyer, T.D., Blangero, J., 1997. Bivariate quantitative trait linkage analysis:
pleiotropy versus co-incident linkages. Genet Epidemiol 14, 953-958.
Alvergne, A., Jokela, M., Lummaa, V., 2010. Personality and reproductive success in a
high-fertility human population. Proc Natl Acad Sci U S A 107, 11745-11750.
Amodio, D.M., Frith, C.D., 2006. Meeting of minds: the medial frontal cortex and social
cognition. Nat Rev Neurosci 7, 268-277.
Asendorpf, J.B., Wilpers, S., 1998. Personality effects on social relationships. Journal of
Personality and Social Psychology 74, 1531-1544.
Avinun, R., Isreal, S., Knodt, A.R., Hariri, A.R., bioarXiv. No evidence for associations
between big five personality traits and variability in brain gray or white matter.
Baird, B., Smallwood, J., Gorgolewski, K.J., Margulies, D.S., 2013. Medial and lateral
networks in anterior prefrontal cortex support metacognitive ability for memory and
perception. J Neurosci 33, 16657-16665.
Benjamini, Y., Hochberg, Y., 1995. Controlling the False Discovery Rate: A Practical
and Powerful Approach to Multiple Testing. Journal of the Royal Statistical Society.
Series B (Methodological) 57, 289-300.
Bernhardt, B.C., Valk, S.L., Silani, G., Bird, G., Frith, U., Singer, T., 2014. Selective
disruption of sociocognitive structural brain networks in autism and alexithymia. Cereb
Cortex 24, 3258-3267.
Bjornebekk, A., Fjell, A.M., Walhovd, K.B., Grydeland, H., Torgersen, S., Westlye, L.T.,
2013. Neuronal correlates of the five factor model (FFM) of human personality:
Multimodal imaging in a large healthy sample. Neuroimage 65, 194-208.
Bludau, S., Eickhoff, S.B., Mohlberg, H., Caspers, S., Laird, A.R., Fox, P.T., Schleicher,
A., Zilles, K., Amunts, K., 2014. Cytoarchitecture, probability maps and functions of the
human frontal pole. Neuroimage 93 Pt 2, 260-275.
Bouchard, T.J., Jr., 1994. Genes, environment, and personality. Science 264, 1700-1701.
Bouchard, T.J., Jr., Loehlin, J.C., 2001. Genes, evolution, and personality. Behav Genet
31, 243-273.
Bouchard, T.J., Jr., McGue, M., 2003. Genetic and environmental influences on human
psychological differences. J Neurobiol 54, 4-45.
Buckner, R.L., Andrews-Hanna, J.R., Schacter, D.L., 2008. The brain's default network:
anatomy, function, and relevance to disease. Ann N Y Acad Sci 1124, 1-38.
Consortium, G.o.P., 2015. Meta-analysis of genome-wide association studies for
neuroticism, and the polygenic association with major depressive disorder. JAMA
Psychiatry.
Costa, P.T., McCrae, R.R., 1992. NEO PI-R Professional Manual. Psychological
Assessment Resources, Inc.
Dale, A.M., Fischl, B., Sereno, M.I., 1999. Cortical surface-based analysis. I.
Segmentation and surface reconstruction. Neuroimage 9, 179-194.
Damian, R.I., Spengler, M., Sutu, A., Roberts, B.W., 2019. Sixteen going on sixty-six: A
longitudinal study of personality stability and change across 50 years. J Pers Soc Psychol
117, 674-695.

de Moor, M.H., Costa, P.T., Terracciano, A., Krueger, R.F., de Geus, E.J., Toshiko, T.,
Penninx, B.W., Esko, T., Madden, P.A., Derringer, J., Amin, N., Willemsen, G.,
Hottenga, J.J., Distel, M.A., Uda, M., Sanna, S., Spinhoven, P., Hartman, C.A., Sullivan,
P., Realo, A., Allik, J., Heath, A.C., Pergadia, M.L., Agrawal, A., Lin, P., Grucza, R.,
Nutile, T., Ciullo, M., Rujescu, D., Giegling, I., Konte, B., Widen, E., Cousminer, D.L.,
Eriksson, J.G., Palotie, A., Peltonen, L., Luciano, M., Tenesa, A., Davies, G., Lopez,
L.M., Hansell, N.K., Medland, S.E., Ferrucci, L., Schlessinger, D., Montgomery, G.W.,
Wright, M.J., Aulchenko, Y.S., Janssens, A.C., Oostra, B.A., Metspalu, A., Abecasis,
G.R., Deary, I.J., Raikkonen, K., Bierut, L.J., Martin, N.G., van Duijn, C.M., Boomsma,
D.I., 2012. Meta-analysis of genome-wide association studies for personality. Mol
Psychiatry 17, 337-349.
DeYoung, C.G., Hirsh, J.B., Shane, M.S., Papademetris, X., Rajeevan, N., Gray, J.R.,
2010. Testing predictions from personality neuroscience. Brain structure and the big five.
Psychol Sci 21, 820-828.
Dubois, J., Galdi, P., Han, Y., Paul, L.K., Adolphs, R., 2018. Resting-state functional
brain connectivity best predicts the personality dimension of openness to experience.
Personal Neurosci 1.
Eickhoff, S.B., Constable, R.T., Yeo, B.T.T., 2018. Topographic organization of the
cerebral cortex and brain cartography. Neuroimage 170, 332-347.
Eyler, L.T., Chen, C.H., Panizzon, M.S., Fennema-Notestine, C., Neale, M.C., Jak, A.,
Jernigan, T.L., Fischl, B., Franz, C.E., Lyons, M.J., Grant, M., Prom-Wormley, E.,
Seidman, L.J., Tsuang, M.T., Fiecas, M.J., Dale, A.M., Kremen, W.S., 2012. A
comparison of heritability maps of cortical surface area and thickness and the influence
of adjustment for whole brain measures: a magnetic resonance imaging twin study. Twin
Res Hum Genet 15, 304-314.
Ferschmann, L., Fjell, A.M., Vollrath, M.E., Grydeland, H., Walhovd, K.B., Tamnes,
C.K., 2018. Personality Traits Are Associated With Cortical Development Across
Adolescence: A Longitudinal Structural MRI Study. Child Dev 89, 811-822.
Fischl, B., 2013. FreeSurfer. Neuroimage 62, 774-781.
Fischl, B., Dale, A.M., 2000. Measuring the thickness of the human cerebral cortex from
magnetic resonance images. Proc Natl Acad Sci U S A 97, 11050-11055.
Fischl, B., Sereno, M.I., Dale, A.M., 1999. Cortical surface-based analysis. II: Inflation,
flattening, and a surface-based coordinate system. Neuroimage 9, 195-207.
Fleming, S.M., Dolan, R.J., 2012. The neural basis of metacognitive ability. Philos Trans
R Soc Lond B Biol Sci 367, 1338-1349.
Fox, P.T., Lancaster, J.L., Laird, A.R., Eickhoff, S.B., 2014. Meta-analysis in human
neuroimaging: computational modeling of large-scale databases. Annu Rev Neurosci 37,
409-434.
Funder, D.C., 2001. Personality. Annu Rev Psychol 52, 197-221.
Genon, S., Reid, A., Langner, R., Amunts, K., Eickhoff, S.B., 2018. How to Characterize
the Function of a Brain Region. Trends Cogn Sci 22, 350-364.
Glahn, D.C., Winkler, A.M., Kochunov, P., Almasy, L., Duggirala, R., Carless, M.A.,
Curran, J.C., Olvera, R.L., Laird, A.R., Smith, S.M., Beckmann, C.F., Fox, P.T.,
Blangero, J., 2010. Genetic control over the resting brain. Proc Natl Acad Sci U S A 107,
1223-1228.

Personality and local brain structure

Glasser, M.F., Sotiropoulos, S.N., Wilson, J.A., Coalson, T.S., Fischl, B., Andersson,
J.L., Xu, J., Jbabdi, S., Webster, M., Polimeni, J.R., Van Essen, D.C., Jenkinson, M.,
Consortium, W.U.-M.H., 2013. The minimal preprocessing pipelines for the Human
Connectome Project. Neuroimage 80, 105-124.
Grasby, K.L., Jahanshad, N., Painter, J.N., Colodro-Conde, L., Bralten, J., Hibar, D.P.,
Lind, P.A., Pizzagalli, F., Ching, C.R.K., McMahon, M.A.B., Shatokhina, N., Zsembik,
L.C.P., Thomopoulos, S.I., Zhu, A.H., Strike, L.T., Agartz, I., Alhusaini, S., Almeida,
M.A.A., Alnaes, D., Amlien, I.K., Andersson, M., Ard, T., Armstrong, N.J., Ashley-
Koch, A., Atkins, J.R., Bernard, M., Brouwer, R.M., Buimer, E.E.L., Bulow, R., Burger,
C., Cannon, D.M., Chakravarty, M., Chen, Q., Cheung, J.W., Couvy-Duchesne, B., Dale,
A.M., Dalvie, S., de Araujo, T.K., de Zubicaray, G.I., de Zwarte, S.M.C., den Braber, A.,
Doan, N.T., Dohm, K., Ehrlich, S., Engelbrecht, H.R., Erk, S., Fan, C.C., Fedko, I.O.,
Foley, S.F., Ford, J.M., Fukunaga, M., Garrett, M.E., Ge, T., Giddaluru, S., Goldman,
A.L., Green, M.J., Groenewold, N.A., Grotegerd, D., Gurholt, T.P., Gutman, B.A.,
Hansell, N.K., Harris, M.A., Harrison, M.B., Haswell, C.C., Hauser, M., Herms, S.,
Heslenfeld, D.J., Ho, N.F., Hoehn, D., Hoffmann, P., Holleran, L., Hoogman, M.,
Hottenga, J.J., Ikeda, M., Janowitz, D., Jansen, I.E., Jia, T., Jockwitz, C., Kanai, R.,
Karama, S., Kasperaviciute, D., Kaufmann, T., Kelly, S., Kikuchi, M., Klein, M., Knapp,
M., Knodt, A.R., Kramer, B., Lam, M., Lancaster, T.M., Lee, P.H., Lett, T.A., Lewis,
L.B., Lopes-Cendes, I., Luciano, M., Macciardi, F., Marquand, A.F., Mathias, S.R.,
Melzer, T.R., Milaneschi, Y., Mirza-Schreiber, N., Moreira, J.C.V., Muhleisen, T.W.,
Muller-Myhsok, B., Najt, P., Nakahara, S., Nho, K., Olde Loohuis, L.M., Orfanos, D.P.,
Pearson, J.F., Pitcher, T.L., Putz, B., Quide, Y., Ragothaman, A., Rashid, F.M., Reay,
W.R., Redlich, R., Reinbold, C.S., Repple, J., Richard, G., Riedel, B.C., Risacher, S.L.,
Rocha, C.S., Mota, N.R., Salminen, L., Saremi, A., Saykin, A.J., Schlag, F., Schmaal, L.,
Schofield, P.R., Secolin, R., Shapland, C.Y., Shen, L., Shin, J., Shumskaya, E., Sonderby,
I.E., Sprooten, E., Tansey, K.E., Teumer, A., Thalamuthu, A., Tordesillas-Gutierrez, D.,
Turner, J.A., Uhlmann, A., Vallerga, C.L., van der Meer, D., van Donkelaar, M.M.J., van
Eijk, L., van Erp, T.G.M., van Haren, N.E.M., van Rooij, D., van Tol, M.J., Veldink,
J.H., Verhoef, E., Walton, E., Wang, M., Wang, Y., Wardlaw, J.M., Wen, W., Westlye,
L.T., Whelan, C.D., Witt, S.H., Wittfeld, K., Wolf, C., Wolfers, T., Wu, J.Q., Yasuda,
C.L., Zaremba, D., Zhang, Z., Zwiers, M.P., Artiges, E., Assareh, A.A., Ayesa-Arriola,
R., Belger, A., Brandt, C.L., Brown, G.G., Cichon, S., Curran, J.E., Davies, G.E.,
Degenhardt, F., Dennis, M.F., Dietsche, B., Djurovic, S., Doherty, C.P., Espiritu, R.,
Garijo, D., Gil, Y., Gowland, P.A., Green, R.C., Hausler, A.N., Heindel, W., Ho, B.C.,
Hoffmann, W.U., Holsboer, F., Homuth, G., Hosten, N., Jack, C.R., Jr., Jang, M., Jansen,
A., Kimbrel, N.A., Kolskar, K., Koops, S., Krug, A., Lim, K.O., Luykx, J.J., Mathalon,
D.H., Mather, K.A., Mattay, V.S., Matthews, S., Mayoral Van Son, J., McEwen, S.C.,
Melle, I., Morris, D.W., Mueller, B.A., Nauck, M., Nordvik, J.E., Nothen, M.M.,
O'Leary, D.S., Opel, N., Martinot, M.P., Pike, G.B., Preda, A., Quinlan, E.B., Rasser,
P.E., Ratnakar, V., Reppermund, S., Steen, V.M., Tooney, P.A., Torres, F.R., Veltman,
D.J., Voyvodic, J.T., Whelan, R., White, T., Yamamori, H., Adams, H.H.H., Bis, J.C.,
Debette, S., Decarli, C., Fornage, M., Gudnason, V., Hofer, E., Ikram, M.A., Launer, L.,
Longstreth, W.T., Lopez, O.L., Mazoyer, B., Mosley, T.H., Roshchupkin, G.V.,
Satizabal, C.L., Schmidt, R., Seshadri, S., Yang, Q., Alzheimer's Disease Neuroimaging,
I., Consortium, C., Consortium, E., Consortium, I., Consortium, S.Y.S., Parkinson's

Progression Markers, I., Alvim, M.K.M., Ames, D., Anderson, T.J., Andreassen, O.A.,
Arias-Vasquez, A., Bastin, M.E., Baune, B.T., Beckham, J.C., Blangero, J., Boomsma,
D.I., Brodaty, H., Brunner, H.G., Buckner, R.L., Buitelaar, J.K., Bustillo, J.R., Cahn, W.,
Cairns, M.J., Calhoun, V., Carr, V.J., Caseras, X., Caspers, S., Cavalleri, G.L., Cendes,
F., Corvin, A., Crespo-Facorro, B., Dalrymple-Alford, J.C., Dannlowski, U., de Geus,
E.J.C., Deary, I.J., Delanty, N., Depondt, C., Desrivieres, S., Donohoe, G., Espeseth, T.,
Fernandez, G., Fisher, S.E., Flor, H., Forstner, A.J., Francks, C., Franke, B., Glahn, D.C.,
Gollub, R.L., Grabe, H.J., Gruber, O., Haberg, A.K., Hariri, A.R., Hartman, C.A.,
Hashimoto, R., Heinz, A., Henskens, F.A., Hillegers, M.H.J., Hoekstra, P.J., Holmes,
A.J., Hong, L.E., Hopkins, W.D., Hulshoff Pol, H.E., Jernigan, T.L., Jonsson, E.G.,
Kahn, R.S., Kennedy, M.A., Kircher, T.T.J., Kochunov, P., Kwok, J.B.J., Le Hellard, S.,
Loughland, C.M., Martin, N.G., Martinot, J.L., McDonald, C., McMahon, K.L., Meyer-
Lindenberg, A., Michie, P.T., Morey, R.A., Mowry, B., Nyberg, L., Oosterlaan, J.,
Ophoff, R.A., Pantelis, C., Paus, T., Pausova, Z., Penninx, B., Polderman, T.J.C.,
Posthuma, D., Rietschel, M., Roffman, J.L., Rowland, L.M., Sachdev, P.S., Samann,
P.G., Schall, U., Schumann, G., Scott, R.J., Sim, K., Sisodiya, S.M., Smoller, J.W.,
Sommer, I.E., St Pourcain, B., Stein, D.J., Toga, A.W., Trollor, J.N., Van der Wee,
N.J.A., van 't Ent, D., Volzke, H., Walter, H., Weber, B., Weinberger, D.R., Wright, M.J.,
Zhou, J., Stein, J.L., Thompson, P.M., Medland, S.E., Enhancing NeuroImaging Genetics
through Meta-Analysis Consortium -Genetics working, g., 2020. The genetic architecture
of the human cerebral cortex. Science 367.
Gray, J.C., Owens, M.M., Hyatt, C.S., Miller, J.D., 2019. No evidence for morphometric
associations of the amygdala and hippocampus with the five-factor model personality
traits. psyarXiv.
Harris, M.A., Brett, C.E., Johnson, W., Deary, I.J., 2016. Personality stability from age
14 to age 77 years. Psychol Aging 31, 862-874.
Hofer, E., al., e., 2018. Genetic Determinants of Cortical Structure (Thickness, Surface
Area and Volumes) among Disease Free Adults in the CHARGE Consortium. bioRxiv
preprint first posted online Sep. 9, 2018; doi: http://dx.doi.org/10.1101/409649.
Holmes, A.J., Hollinshead, M.O., O'Keefe, T.M., Petrov, V.I., Fariello, G.R., Wald, L.L.,
Fischl, B., Rosen, B.R., Mair, R.W., Roffman, J.L., Smoller, J.W., Buckner, R.L., 2015.
Brain Genomics Superstruct Project initial data release with structural, functional, and
behavioral measures. Sci Data 2, 150031.
Hopwood, C.J., Donnellan, M.B., Blonigen, D.M., Krueger, R.F., McGue, M., Iacono,
W.G., Burt, S.A., 2011. Genetic and environmental influences on personality trait
stability and growth during the transition to adulthood: a three-wave longitudinal study. J
Pers Soc Psychol 100, 545-556.
Jang, K.L., Livesley, W.J., Vernon, P.A., 1996. Heritability of the big five personality
dimensions and their facets: a twin study. J Pers 64, 577-591.
John, O.P., Naumann, L.P., Soto, C.J., 2008. Paradigm Shift to the Itegrative Big-Five
Trait Taxonomy: History, Measurement, and Conceptual Issues. Guilford Press, New
York, NY.
John, O.P., Srivastava, S., 1999. The Big Five Trait taxonomy: History, measurement,
and theoretical perspectives. In: Pervin, L.A., John, O.P. (Eds.), Handbook of
Personality: Theory and Research. Guilford Press, pp. 102-138.

Kebets, V., Holmes, A.J., Orban, C., Tang, S., Li, J., Sun, N., Kong, R., Poldrack, R.A.,
Yeo, B.T.T., 2019. Somatosensory-Motor Dysconnectivity Spans Multiple
Transdiagnostic Dimensions of Psychopathology. Biol Psychiatry 86, 779-791.
Kharabian Masouleh, S., Eickhoff, S.B., Genon, S., 2020. Searching for replicable
associations between cortical thickness and psychometric variables in healthy adults:
empirical facts. Biorxiv.
Kharabian Masouleh, S., Eickhoff, S.B., Hoffstaedter, F., Genon, S., Alzheimer's Disease
Neuroimaging, I., 2019. Empirical examination of the replicability of associations
between brain structure and psychological variables. elife.
Kong, R., Li, J., Orban, C., Sabuncu, M.R., Liu, H., Schaefer, A., Sun, N., Zuo, X.N.,
Holmes, A.J., Eickhoff, S.B., Yeo, B.T.T., 2019. Spatial Topography of Individual-
Specific Cortical Networks Predicts Human Cognition, Personality, and Emotion. Cereb
Cortex 29, 2533-2551.
Kremen, W.S., Prom-Wormley, E., Panizzon, M.S., Eyler, L.T., Fischl, B., Neale, M.C.,
Franz, C.E., Lyons, M.J., Pacheco, J., Perry, M.E., Stevens, A., Schmitt, J.E., Grant,
M.D., Seidman, L.J., Thermenos, H.W., Tsuang, M.T., Eisen, S.A., Dale, A.M.,
Fennema-Notestine, C., 2010. Genetic and environmental influences on the size of
specific brain regions in midlife: the VETSA MRI study. Neuroimage 49, 1213-1223.
Laajaj, R., Macours, K., Pinzon Hernandez, D.A., Arias, O., Gosling, S.D., Potter, J.,
Rubio-Codina, M., Vakis, R., 2019. Challenges to capture the big five personality traits in
non-WEIRD populations. Science Advances 5.
Laird, A.R., Eickhoff, S.B., Fox, P.M., Uecker, A.M., Ray, K.L., Saenz, J.J., Jr., McKay,
D.R., Bzdok, D., Laird, R.W., Robinson, J.L., Turner, J.A., Turkeltaub, P.E., Lancaster,
J.L., Fox, P.T., 2011. The BrainMap strategy for standardization, sharing, and meta-
analysis of neuroimaging data. BMC Res Notes 4, 349.
Laird, A.R., Eickhoff, S.B., Kurth, F., Fox, P.M., Uecker, A.M., Turner, J.A., Robinson,
J.L., Lancaster, J.L., Fox, P.T., 2009. ALE Meta-Analysis Workflows Via the Brainmap
Database: Progress Towards A Probabilistic Functional Brain Atlas. Front Neuroinform
3, 23.
Latzman, R.D., Hecht, L.K., Freeman, H.D., Schapiro, S.J., Hopkins, W.D., 2015.
Neuroanatomical correlates of personality in chimpanzees (Pan troglodytes):
Associations between personality and frontal cortex. Neuroimage 123, 63-71.
Lo, M.T., Hinds, D.A., Tung, J.Y., Franz, C., Fan, C.C., Wang, Y., Smeland, O.B.,
Schork, A., Holland, D., Kauppi, K., Sanyal, N., Escott-Price, V., Smith, D.J.,
O'Donovan, M., Stefansson, H., Bjornsdottir, G., Thorgeirsson, T.E., Stefansson, K.,
McEvoy, L.K., Dale, A.M., Andreassen, O.A., Chen, C.H., 2017. Genome-wide analyses
for personality traits identify six genomic loci and show correlations with psychiatric
disorders. Nat Genet 49, 152-156.
Loehlin, J.C., McCrae, R.R., Costa, P.T., John, O.P., 1998. Heritabilities of Common and
Measure-Specific Components of the Big Five Personality Factors. Journal of Research
in Personality 32, 431-453.
Marcus, D.S., Harms, M.P., Snyder, A.Z., Jenkinson, M., Wilson, J.A., Glasser, M.F.,
Barch, D.M., Archie, K.A., Burgess, G.C., Ramaratnam, M., Hodge, M., Horton, W.,
Herrick, R., Olsen, T., McKay, M., House, M., Hileman, M., Reid, E., Harwell, J.,
Coalson, T., Schindler, J., Elam, J.S., Curtiss, S.W., Van Essen, D.C., Consortium, W.U.-

M.H., 2013. Human Connectome Project informatics: quality control, database services,
and data visualization. Neuroimage 80, 202-219.
McCrae, R.R., Costa, P.T., 2004. A contemplated revision of the NEO Five-Factor
Inventory. Pers Individ Dif. 36, 587-596.
McCrae, R.R., Costa, P.T., Jr., 1997. Personality trait structure as a human universal. Am
Psychol 52, 509-516.
McCrae, R.R., Terracciano, A., 2005. Personality profiles of cultures: aggregate
personality traits. J Pers Soc Psychol 89, 407-425.
Miller, J.D., Lynam, D.R., Widiger, T.A., Leukefeld, C., 2001. Personality disorders as
extreme variants of common personality dimensions: can the Five-Factor Model
adequately represent psychopathy? J Pers 69, 253-276.
Mischel, W., 2004. Toward an integrative science of the person. Annu Rev Psychol 55, 1-
22.
Mugler, J.P., 3rd, Brookeman, J.R., 1990. Three-dimensional magnetization-prepared
rapid gradient-echo imaging (3D MP RAGE). Magn Reson Med 15, 152-157.
Nakao, K., Takaishi, J., Tatsuta, K., Katayama, H., Iwase, M., Yorifuji, K., Takeda, M.,
2000. The influences of family environment on personality traits. Psychiatry Clin
Neurosci 54, 91-95.
Nostro, A.D., Muller, V.I., Reid, A.T., Eickhoff, S.B., 2017. Correlations Between
Personality and Brain Structure: A Crucial Role of Gender. Cereb Cortex 27, 3698-3712.
Owens, M.M., Hyatt, C.S., Gray, J.C., Carter, N.T., MacKillop, J., Miller, J.D., Sweet,
L.H., 2019. Cortical morphometry of the five-factor model of personality: findings from
the Human Connectome Project full sample. Soc Cogn Affect Neurosci 14, 381-395.
Panizzon, M.S., Fennema-Notestine, C., Eyler, L.T., Jernigan, T.L., Prom-Wormley, E.,
Neale, M., Jacobson, K., Lyons, M.J., Grant, M.D., Franz, C.E., Xian, H., Tsuang, M.,
Fischl, B., Seidman, L., Dale, A., Kremen, W.S., 2009. Distinct genetic influences on
cortical surface area and cortical thickness. Cereb Cortex 19, 2728-2735.
Penke, L., Jokela, M., 2016. The evolutionary genetics of personality revisited. Current
Opinion in Psychology 7, 104-109.
Perkins, E.R., Latzman, R.D., Patrick, C.J., 2020. Interfacing neural constructs with the
Hierarchical Taxonomy of Psychopathology: 'Why' and 'how'. Personal Ment Health 14,
106-122.
Riccelli, R., Toschi, N., Nigro, S., Terracciano, A., Passamonti, L., 2017. Surface-based
morphometry reveals the neuroanatomical basis of the five-factor model of personality.
Soc Cogn Affect Neurosci 12, 671-684.
Rothmann, S., Coetzer, E.P., 2003. The big five personality dimensions and job
performance. SA Journal of Industrial Psychology 29.
Saucier, G., Srivastava, S., 2015. What makes a good structural model of personality?
Evaluating the big five and alternatives. In: Mikulincer, M., Shaver, P.R. (Eds.), APA
Handbook of Personality and Social Psychology: Vol 4. Personality Processes and
Individual Differences. American Psychological Association.
Schaefer, A., Kong, R., Gordon, E.M., Laumann, T.O., Zuo, X.N., Holmes, A.J.,
Eickhoff, S.B., Yeo, B.T.T., 2018. Local-Global Parcellation of the Human Cerebral
Cortex from Intrinsic Functional Connectivity MRI. Cereb Cortex 28, 3095-3114.

Schoenberg, M.R., Scott, J.G., Duff, K., Adams, R.L., 2002. Estimation of WAIS-III
intelligence from combined performance and demographic variables: development of the
OPIE-3. Clin Neuropsychol 16, 426-437.
Strickhouser, J.E., Zell, E., Krizan, Z., 2017. Does personality predict health and well-
being? A metasynthesis. Health Psychol 36, 797-810.
Strike, L.T., Hansell, N.K., Couvy-Duchesne, B., Thompson, P.M., de Zubicaray, G.I.,
McMahon, K.L., Wright, M.J., 2018. Genetic Complexity of Cortical Structure:
Differences in Genetic and Environmental Factors Influencing Cortical Surface Area and
Thickness. Cereb Cortex.
Strike, L.T., Hansell, N.K., Couvy-Duchesne, B., Thompson, P.M., de Zubicaray, G.I.,
McMahon, K.L., Wright, M.J., 2019. Genetic Complexity of Cortical Structure:
Differences in Genetic and Environmental Factors Influencing Cortical Surface Area and
Thickness. Cereb Cortex 29, 952-962.
Trull, T.J., 2013. Dimensional models of personality: the five-factor model and the DSM-
5. Dialogues Clin Neurosci 15, 135-146.
Valk, S.L., Bernhardt, B.C., Bockler, A., Kanske, P., Singer, T., 2016a. Substrates of
metacognition on perception and metacognition on higher-order cognition relate to
different subsystems of the mentalizing network. Hum Brain Mapp 37, 3388-3399.
Valk, S.L., Bernhardt, B.C., Bockler, A., Trautwein, F.M., Kanske, P., Singer, T., 2016b.
Socio-Cognitive Phenotypes Differentially Modulate Large-Scale Structural Covariance
Networks. Cereb Cortex.
Valk, S.L., Bernhardt, B.C., Trautwein, F.M., Bockler, A., Kanske, P., Guizard, N.,
Collins, D.L., Singer, T., 2017. Structural plasticity of the social brain: Differential
change after socio-affective and cognitive mental training. Sci Adv 3, e1700489.
van den Berg, S.M., de Moor, M.H., Verweij, K.J., Krueger, R.F., Luciano, M., Arias
Vasquez, A., Matteson, L.K., Derringer, J., Esko, T., Amin, N., Gordon, S.D., Hansell,
N.K., Hart, A.B., Seppala, I., Huffman, J.E., Konte, B., Lahti, J., Lee, M., Miller, M.,
Nutile, T., Tanaka, T., Teumer, A., Viktorin, A., Wedenoja, J., Abdellaoui, A., Abecasis,
G.R., Adkins, D.E., Agrawal, A., Allik, J., Appel, K., Bigdeli, T.B., Busonero, F.,
Campbell, H., Costa, P.T., Smith, G.D., Davies, G., de Wit, H., Ding, J., Engelhardt,
B.E., Eriksson, J.G., Fedko, I.O., Ferrucci, L., Franke, B., Giegling, I., Grucza, R.,
Hartmann, A.M., Heath, A.C., Heinonen, K., Henders, A.K., Homuth, G., Hottenga, J.J.,
Iacono, W.G., Janzing, J., Jokela, M., Karlsson, R., Kemp, J.P., Kirkpatrick, M.G.,
Latvala, A., Lehtimaki, T., Liewald, D.C., Madden, P.A., Magri, C., Magnusson, P.K.,
Marten, J., Maschio, A., Mbarek, H., Medland, S.E., Mihailov, E., Milaneschi, Y.,
Montgomery, G.W., Nauck, M., Nivard, M.G., Ouwens, K.G., Palotie, A., Pettersson, E.,
Polasek, O., Qian, Y., Pulkki-Raback, L., Raitakari, O.T., Realo, A., Rose, R.J.,
Ruggiero, D., Schmidt, C.O., Slutske, W.S., Sorice, R., Starr, J.M., St Pourcain, B., Sutin,
A.R., Timpson, N.J., Trochet, H., Vermeulen, S., Vuoksimaa, E., Widen, E., Wouda, J.,
Wright, M.J., Zgaga, L., Generation, S., Porteous, D., Minelli, A., Palmer, A.A., Rujescu,
D., Ciullo, M., Hayward, C., Rudan, I., Metspalu, A., Kaprio, J., Deary, I.J., Raikkonen,
K., Wilson, J.F., Keltikangas-Jarvinen, L., Bierut, L.J., Hettema, J.M., Grabe, H.J.,
Penninx, B.W., van Duijn, C.M., Evans, D.M., Schlessinger, D., Pedersen, N.L.,
Terracciano, A., McGue, M., Martin, N.G., Boomsma, D.I., 2016. Meta-analysis of
Genome-Wide Association Studies for Extraversion: Findings from the Genetics of
Personality Consortium. Behav Genet 46, 170-182.

Personality and local brain structure

Van Essen, D.C., Glasser, M.F., 2018. Parcellating Cerebral Cortex: How Invasive
Animal Studies Inform Noninvasive Mapmaking in Humans. Neuron 99, 640-663.
Van Essen, D.C., Smith, S.M., Barch, D.M., Behrens, T.E., Yacoub, E., Ugurbil, K.,
Consortium, W.U.-M.H., 2013. The WU-Minn Human Connectome Project: an
overview. Neuroimage 80, 62-79.
Verhagen, J., Wagenmakers, E.J., 2014. Bayesian tests to quantify the result of a
replication attempt. J Exp Psychol Gen 143, 1457-1475.
Verweij, K.J., Yang, J., Lahti, J., Veijola, J., Hintsanen, M., Pulkki-Raback, L.,
Heinonen, K., Pouta, A., Pesonen, A.K., Widen, E., al., e., 2012. Maintenance of genetic
variation in human personality: testing evolutionary models by estimating heritability due
to common causal variants and investigating the effect of distant inbreeding. Evolution
66, 3288-3251.
Vukasovic, T., Bratko, D., 2015. Heritability of personality: A meta-analysis of behavior
genetic studies. Psychol Bull 141, 769-785.
Wechsler, D., 1999. Wechsler Abbreviated Scale of Intelligence. The Psychological
Corporation., San Antonio, TX.
Weintraub, S., Dikmen, S.S., Heaton, R.K., Tulsky, D.S., Zelazo, P.D., Bauer, P.J.,
Carlozzi, N.E., Slotkin, J., Blitz, D., Wallner-Allen, K., Fox, N.A., Beaumont, J.L.,
Mungas, D., Nowinski, C.J., Richler, J., Deocampo, J.A., Anderson, J.E., Manly, J.J.,
Borosh, B., Havlik, R., Conway, K., Edwards, E., Freund, L., King, J.W., Moy, C., Witt,
E., Gershon, R.C., 2013. Cognition assessment using the NIH Toolbox. Neurology 80,
S54-64.
Winkler, A.M., Kochunov, P., Blangero, J., Almasy, L., Zilles, K., Fox, P.T., Duggirala,
R., Glahn, D.C., 2010. Cortical thickness or grey matter volume? The importance of
selecting the phenotype for imaging genetics studies. Neuroimage 53, 1135-1146.
Worsley, K., Taylor, J.E., Carbonell, F., Chung, M.K., Duerden, E., Bernhardt, B.C.,
Lyttelton, O.C., Boucher, M., Evans, A.C., 2009. SurfStat: A Matlab toolbox for the
statistical analysis of univariate and multivariate surface and volumetric data using linear
mixed effects models and random field theory. . Neuroimage 47.
Wu, X., He, H., Shi, L., Xia, Y., Zuang, K., Feng, Q., Zhang, Y., Ren, Z., Wei, D., Qiu,
J., 2019. Personality traits are related with dynamic functional connectivity in major
depression disorder: A resting-state analysis. J Affect Disord 245, 1032-1042.
Yarkoni, T., 2015. Neurobiological substrates of personality: A critical overview. In: M.,
M., R., S.P., L., C.M., J., L.R. (Eds.), APA handbooks in psychology®. APA handbook
of personality and social psychology. American Psychological Association, pp. 61–83.
Zheng, D., Chen, J., Wang, X., Zhou, Y., 2019. Genetic contribution to the phenotypic
correlation between trait impulsivity and resting-state functional connectivity of the
amygdala and its subregions. Neuroimage 201, 115997.
1089

Personality and local brain structure