**Evolutionary dynamics of the trypsin superfamily and the origins of novel secretory cell**

**function in the internalized ectoderm of the sea anemone *Nematostella vectensis***

Leslie S. Babonis[1,3], Joseph F. Ryan[1,2], Camille Enjolras[1], and Mark Q. Martindale[1,2]

**Affiliations:**

[1]Whitney Laboratory for Marine Bioscience, University of Florida, St. Augustine, FL 32080

[2]Department of Biology, University of Florida, Gainesville, FL 32611, USA.

[3]Corresponding author: babonis@whitney.ufl.edu

**Abstract**

Understanding the drivers of morphological diversity is a persistent challenge in evolutionary

biology. Here, we investigate functional diversification of secretory cells in the sea anemone

*Nematostella vectensis* to understand the mechanisms promoting cellular specialization across

animals. We demonstrate regionalized expression of gland cell subtypes in the internal ectoderm

of *N. vectensis* and show that adult gland cell identity is acquired very early in development. A

phylogenetic survey of trypsins across animals suggests this gene family has undergone

numerous expansions. We reveal unexpected diversity in trypsin protein structure and show that

trypsin diversity arose through independent acquisitions of non-trypsin domains. Finally, we

show that trypsin diversification in *N. vectensis* was effected through a combination of tandem

duplication, exon shuffling, and retrotransposition. Together we reveal that numerous

evolutionary mechanisms drove trypsin duplication and divergence during the morphological

specialization of cell types, and suggest the secretory cell phenotype is highly adaptable as a

vehicle for novel secretory products.

**Introduction**

The development of new tissue layers provides the opportunity to spatially segregate cell types

enabling the compartmentalization of different functions. Cnidarians are diploblasts, comprised

of an internal endodermal epithelium separated from an external ectodermal epithelium by a

largely acellular matrix called mesoglea. Anthozoans (corals, sea anemones, and their kin) are

unusual among cnidarians in their possession of internal tissues (pharynx and mesenteries) that

arise by secondary epithelial fold morphogenesis following completion of gastrulation (Magie et

al., 2007). Additional growth and differentiation of both internalized layers results in the

morphogenesis of the pharynx and mesenteries and in an adult form quite different from that of

medusozoans. In anthozoans, both layers (endoderm and ectoderm) are in contact with the

gastric cavity, whereas in medusozoans (and, indeed, most other animals), the gastrovascular

cavity is lined only by endoderm.

*Nematostella vectensis,* the starlet sea anemone, has become a valuable model for studies of

animal body plan evolution (Finnerty et al., 2004; Fritzenwanker et al., 2007; Kusserow et al.,

2005; Leclere and Rentzsch, 2014; Wijesena et al., 2017); yet, little is known about the extent of

cell diversity in the tissues that comprise the pharynx and mesenteries. The endodermal

component of the mesenteries houses the germ cell precursors and two types of muscle cells and

the few recent studies of the mesenteries in *Nematostella* have focused largely on these

endodermal functions (Jahnel et al., 2014; Moiseeva et al., 2017; Renfer et al., 2010). The

ectodermal component of the mesenteries is known to be populated by cnidocytes and gland cells

(Frank and Bleakney, 1976) and two recent studies demonstrated the expression of multiple

proteases in the mesenteries of *N. vectensis* (Babonis et al., 2016; Steinmetz et al., 2017).

Trypsins are the largest family of proteases, and although they have diverse functions, most

trypsins are secreted to the extracellular environment and are therefore expressed in zymogen-

type gland cells (Page and Di Cera, 2008b). A previous study cataloging trypsin diversity from

prokaryotes and eukaryotes identified 75 trypsins in the genome of *N. vectensis* (Page and Di

Cera, 2008a),  suggesting that the few cell types identified anatomically as zymogen gland cells

(Frank and Bleakney, 1976) may belie the digestive capacity of the mesenteries.

We sought to understand the evolutionary mechanisms promoting functional diversification at

the cell and tissue levels in the mesenteries of *N. vectensis,* and to characterize the evolutionary

history of a large (super)family of proteases expressed abundantly in the mesenteries.  Building

on a previous study using RNA-seq to characterize the expression profile of the mesenteries in *N.*

*vectensis* (Babonis et al., 2016), we show that the continuous epithelium comprising the internal

ectoderm in *N. vectensis* is functionally partitioned into different regions associated with distinct

morphologies and functions. Additionally, we show numerous lineage-specific expansions of

trypsins and that trypsin diversification arises through novel domain acquisition. Finally, we

propose a model by which the expansion of trypsins may have promoted specialization of gland

cell subtype in cnidarians.

**Results**

*Morphology and function of the internal ectoderm*

We examined the fine structure of the internal and external ectoderm in the region of the mouth

of *N. vectensis* during feeding for evidence of morphological and functional variation (Figure 1).

Cells in the external ectoderm around the mouth are organized into a low cuboidal type

epithelium that covers the closed mouth between feeding events (Figure 1A-D). In the presence

of prey, the pharynx is partially everted, exposing the tall columnar epithelium of the pharyngeal

ectoderm (Figure 1E-G). After passing through the pharynx (Figure 1H), ingested prey remains

in contact with the ectodermal portion of the mesenteries, which is populated by cnidocytes and

gland cells (Figure 1I-K).

The pharyngeal ectoderm is populated by numerous distinct electron dense (zymogen-secreting)

and electron lucent (mucus-secreting) gland cells (Figure 2A-F). The adjacent non-secretory cells

in this epithelium have distinctive apical electron-dense vesicles (Figure 2F). The proximal

region of the mesentery (adjacent to the body wall) is comprised of endoderm while the distal

portion (the free edge) is comprised of ectoderm (Figure 2G). The ectodermal region gives rise to

both the cnidoglandular tract at the most distal extent (Figure 2H-J, M-O) and the ciliated tract

more proximally (Figure 2K,L). Thin sections of the ectodermal mesentery in the oral region

(near the pharynx) show abundant zymogen gland cells (Figure 2H-J), some of which contain

secretory vesicles with heterogeneous contents (Figure 2J). Ciliated tracts are short and are

present only in the oral end of each mesentery. Cells of the ciliated tract are highly proliferative

and have apical motile cilia but do not have other distinguishing features (Figure 2K,L). The

aboral mesentery lacks a ciliated tract but the cnidoglandular tract still contains numerous

distinct zymogen gland cells, some with motile apical cilia (Figure 2M-O). Mucus-secreting cells

were found in the pharyngeal ectoderm (Figure 2D) and in the external ectoderm of the body

wall and tentacles (Supplemental file 1new), but never in the endoderm.

*Proteolytic enzymes are expressed in the developing mesenteries*

We previously identified numerous genes encoding different classes of proteases to be

upregulated in the adult mesentery of *N. vectensis* (Babonis et al., 2016) . Using *in situ*

hybridization we examined the spatial and temporal expression of 10 proteases of various classes

during early development of the pharynx/mesenteries to understand the ontogeny of digestive

function and the onset of terminal gut cell differentiation. All genes examined were expressed in

individual ectodermal cells of the mesenteries at the primary polyp stage, just after

metamorphosis (Figure 3A,B); two protease genes (NVJ_82725 and NVJ_83864) were also

expressed in the pharyngeal ectoderm of the primary polyp. There was surprisingly little

variation in the onset of protease expression, although serine proteases (trypsins) consistently

exhibited expression in the early planula stage before differentiation of the presumptive pharynx

and mesenteries (Figure 3B). Double fluorescent *in situ* hybridization for two metalloprotease

genes (NVJ_88668 and NVJ_2109) indicates both co-expression of these two enzymes in few

cells at the aboral end of the pharynx and independent expression of the two genes in distinct

cells of the ectodermal mesenteries in the late tentacle bud stage (Figure 3C).

The surprising lack of any obvious spatial segregation in protease expression led us to

hypothesize that many proteases may be co-expressed together in the few anatomically

distinguishable gland cells identified above (Figure 2). Using the raw data from a single-cell

RNA-Seq study published previously (Sebe-Pedros et al., 2018), we show co-expression of six of

the ten proteases we studied by *in situ* hybridization in a single putative gland cell (Figure 3D).

Using the raw data from the same study and a very low cutoff for gene expression ($N \geq 1$ read),

we examined more fully the co-expression of the large superfamily of trypsin proteases and

found 6,727 cells expressing at least one trypsin gene. Nearly 50% of the trypsin-expressing cells

(3,282/6,727) appear to express only a single trypsin, while the remaining cells exhibited co-

expression of up to 24 trypsins (Figure 3E). For each trypsin, we then examined the relationship

between the ubiquity of expression (the total number of cells in which that trypsin is expressed)

and the number of cells in which it is co-expressed with other trypsins and found a strong

positive correlation (Figure 3F), confirming that the trypsins with the broadest expression

profiles were most likely to be co-expressed with other trypsins.

*The tryptome is lineage-specific*

To characterize the tryptome (all proteins with a trypsin domain) of *N. vectensis*, we searched the

JGI gene models (https://genome.jgi.doe.gov/Nemve1/Nemve1.home.html) for all sequences

containing a significant Trypsin or Trypsin_2 domain using hmmsearch (HMMER 3.1b2;

http://hmmer.org) and constructed domain architecture diagrams for each protein (Figure 4). Of

the 72 trypsin gene models that remained after curation (see Methods), 28 encode a trypsin

domain but lack any other conserved domains and the other 44 encode a trypsin domain and at

least one additional conserved domain. In total, trypsin domains were found in association with

24 other domains in *N. vectensis*. To determine if any of these associated domains were

overrepresented in the tryptome, we compared the abundance of trypsin-associated domains in

the tryptome and in the proteins predicted from the JGI gene models (N = 27,273 protein

predictions). Six domains were found to be represented in high abundance ($\geq$10%) in the

tryptome: DIM, ShK, Lustrin_cystein, Sushi, MAM and SRCR (Figure 4A). The DIM and

Lustrin_cysteine domains are present in low abundance throughout the predicted proteome (1

and 4 total domains, respectively), artificially inflating their perceived abundance in the

tryptome. For ShK, Sushi, MAM, and SRCR, $\geq$15% of the domains found in the proteome were

associated with trypsins.

To determine whether the makeup of the tryptome was unique to *N. vectensis*, we searched for

proteins with these same domain architectures in representatives from all domains of life (other

cnidarians, bilaterians, non-metazoan eukaryotes, and a selection of prokaryotes). Two domain

architectures were found to be present across taxa: those with only a trypsin domain, and those

with a trypsin and a PDZ domain (Figure 4B). Trypsins appear to have expanded considerably

after the origin of animals, as both choanoflagellate lineages had fewer than five trypsins but the

ctenophore *Mnemiopsis leidyi* and the placozoan *Trichoplax adhaerens* (representing two of the

earliest diverging animal lineages) both have at least twenty. Among the cnidarians, other

actinarians (sea anemones) shared more trypsin proteins in common with *N. vectensis* than any

other lineage; however, we identified three trypsins (NVJ_105548, NVJ_105271, and

NVJ_199428) specific to *N. vectensis* that were absent even from *Edwardsiella lineata*

(representing the genus sister to *Nematostella*).

*Trypsins diversified independently in cnidarians and bilaterians*

To characterize the diversification of animal trypsins, we built a phylogeny of trypsin domains

from taxa representing each of the five major animal lineages: bilaterians, cnidarians,

placozoans, sponges, and ctenophores. Using this tree, we identify six clades of trypsins and

classify them by their function in human: a non-catalytic group, the intracellular trypsins,

tryptases and transmembrane trypsins, trypsins involved in coagulation and immune response,

chymotrypsins, and the clade including granzymes, pancreatic trypsins, kallikreins, hepatocyte

growth factors, and elastases (Figure 5A). Each of these includes representatives from

bilaterians, cnidarians, and at least one placozoan, sponge, or ctenophore and likely represent the

suite of trypsin clades present in the last common ancestor of animals. The *N. vectensis* tryptome

includes representatives of 5 of 6 clades likely present in the common ancestor of animals. *N.*

*vectensis* may have lost representatives of the tryptase/transmembrane clade as this these trypsins

appear to be present in *M. leidyi*, *A. digitifera*, and bilaterians (Figure 5A). Proteins with only a

trypsin domain were distributed throughout the tree, rather than being clustered in a single clade.

Two proteins from *N. vectensis* (NVJ_203589 and NVJ_23745) had divergent trypsin domains

and were detected only by the Trypsin_2 HMM, both of which appear to be part of the clade that

includes trypsin-PDZ proteins (Supplemental file 2).

We compared the distribution of conserved domains from different clades of trypsins in *N.*

*vectensis* and *H. sapiens* (Figure 5B). In *N. vectensi*s, domain diversity is greatest among the

trypsins that group with human chymotrypsins, as trypsin domains from this clade co-occur with

14 associated domains. Chymotrypsins from *N. vectensis* share four domains in common with

the immune/coagulation group, which is represented by 10 domains. The "pancreatic" group

(including granzymes, kallikreins, HGF, and elastase) is characterized by only 5 domains, 3 of

which are shared with chymotrypsins. Trypsins from the non-catalytic clade lacks associated

domains and the intracellular clade uses unique domains (Death and PDZ). Four trypsin-

associated domains (Sushi, EGF_CA, CUB, and FXa_inhibition) were found in the

immune/coagulation clade of trypsins in both *N. vectensis* and *H. sapiens* and the PDZ domain is

restricted to the intracellular clade of trypsins in both taxa. Surprisingly, there were no other

domains found in common between *N. vectensis* and *H. sapiens* trypsins from the same clade.

(See supplemental file 3 for distribution of human trypsin domain architectures.)

These analyses of trypsin distribution across animals revealed a surprisingly large tryptome in

sea anemones (*N. vectensis* and *E. lineata*) relative to most other taxa (including another

cnidarian, *Hydra magnipapillata*). To determine whether the tryptome of *N. vectensis* is

reflective of other cnidarians, we built a phylogeny using representatives of each class within

Cnidaria (Figure 6). We identify 16 clades of trypsins that include representatives of at least two

lineages of anthozoans and two lineages of medusozoans, suggesting these clades may have been

present in the stem cnidarian. Two clades (the trypsin-MAM and trypsin-ShK clades) seem to

have undergone further expansion in anthozoans after their divergence from medusozoans.

*The Nematostella tryptome diversified through numerous mechanisms*

To understand the mechanisms generating trypsin diversity in *N. vectensis*, we examined the

evolutionary relationships of the 72 trypsin proteins in the tryptome (Figure 7A). Among the 72

predicted proteins, 85% (61/72) had all three conserved residues constituting the catalytic triad

and are likely to function as proteases, 79% (57/72) were predicted to have a signal peptide and

are presumably secreted, and 7% (5/72) were predicted to have a transmembrane domain (see

Supplemental File 4). Four of the five clades of trypsins from *N. vectensis* (excluding the

intracellular clade) include secreted trypsins, membrane-bound trypsins, and trypsins with

divergent sequence that have likely lost their catalytic function. While most (70/72) of the

trypsin domains were encoded across multiple exons (Supplemental file 5), two genes

(NVJ_128003 and NVJ_216003) lack introns completely.

Numerous trypsins from the "pancreatic" and chymotrypsin clades were associated with ShK

domains. Likewise, over 30% (26/82) of the ShK domains in *N. vectensis* are associated with

trypsins (Figure 4A). To determine if the combination of the trypsin and ShK domains may have

duplicated together, we built a phylogeny of all 108 ShK domains from the *N. vectensis* proteins

predicted from gene models (Figure 7B). Despite the abundance of trypsin-ShK associations, the

distribution of trypsin-ShK proteins suggests that this domain combination may have duplicated

together only once, giving rise to NVJ_218669 and NVJ_218670; Figure 7A).

*Trypsin diversity increases through new associations with old domains*

Gene age can be estimated using a phylostratigraphic approach; in such analyses, the minimum

age of a gene is inferred by identifying the last common ancestor in which the gene is present

(Domazet-Loso et al., 2007; Long et al., 2003). We examined the age of the trypsins found in *N.*

*vectensis* and the age of each associated domain across all domains of life to understand the

evolution of trypsin diversity. Trypsin-PDZ and a subset of the trypsin-only proteins likely arose

before bacteria/archaea split from eukaryotes, over 2 billion years ago (Figure 8). While trypsin-

only proteins are present in every lineage examined, trypsin-PDZ proteins appear to have been

lost in several taxa including *C. owczarzaki*, *M. leidyi*, *A. vanhoeffeni*, and *C. cruxmelitensis*

(Figure 4). All other associations between trypsin and other conserved domains appear to have

originated after the stem metazoan diverged from the rest of life (~800 million years ago). Many

of the trypsin-associated domains originated long before they became associated with trypsin; for

example, the astacin domain was present in the ancestor of all life but the trypsin-astacin

association likely did not arise until the origin of Cnidaria (Figure 8A). By contrast, the SRCR

domain and its association with trypsin likely arose in the stem metazoan as trypsin-SRCR

proteins were found in *M. leidyi* (Supplemental file 6).

There does not appear to be a relationship between the age of the domain and the origin of its

association with trypsin (Figure 8B). Two trypsin associations were found only in *N. vectensis*:

trypsin-DIM (NVJ_199428) and trypsin-WSC (NVJ_105271), and one association was found

only in *Edwarsiidae* (*Nematostella + Edwardsiella)*: trypsin-Lustrin_cystein (NVJ_164017). The

WSC domain is present throughout eukaryotes (Figure 8A) but was associated with trypsin in

only in *N. vectensis*. The Lustrin_cystein domain seems to have arisen in the last common

ancestor of parahoxozoa (Placozoa + Cnidaria + Bilateria). These two associations represent

extreme cases whereby trypsin diversity in *N. vectensis* arose through acquisition of both young

(Lustrin_cystein) and old (WSC) domains.

**Discussion**

*The not-so-simple cnidarian ectoderm*

Although cnidarian body plans develop from only two tissue layers, morphological diversity

varies widely across taxa. Similarly, only a dozen or so morphologically unique cell types have

been described (Frank and Bleakney, 1976; Marlow et al., 2009), but cnidarian genomic and

functional diversity rival that of any bilaterian lineage (Kusserow et al., 2005; Putnam et al.,

2007). While the ectodermal layer comprising the external and pharyngeal epithelia may be

contiguous, these regions are morphologically and functionally distinct in *N. vectensis* (Figure 1)

(Frank and Bleakney, 1976; Steinmetz et al., 2017). In this study, we further demonstrate that the

continuous layer of internal ectoderm from the pharynx through the mesenteries is equally

heterogeneous. The pharyngeal ectoderm houses numerous zymogen and mucous cells while the

ectoderm of the mesenteries houses only the former (Figure 2). This anatomical heterogeneity is

supported by variable gene expression: some proteases are expressed throughout the pharyngeal

and mesentery ectoderm, while others are restricted only to the mesentery ectoderm (Figure

3A,B) (also see (Moran et al., 2013; Steinmetz et al., 2017). Furthermore, the combinatorial

expression of only two proteases can result in the development of at least three distinct cell types

(Figure 3C). Together, the combination of a diverse tryptome and extensive trypsin co-

expression (Figure 3E) suggests cell functional diversity in cnidarians may well exceed historical

expectations.

We found no evidence of endodermal gland cells (zymogen type or mucous type) in our TEM or

*in situ* hybridization results (Figure 2,3, Supplemental file 1). Indeed, all non-neuronal secretory

cells (including mucous cells, zymogen cells, and cnidocytes), are restricted to the ectoderm in

*N. vectensis* but their distribution is heterogeneous. Zymogen cell diversity, for example, is much

higher in the internal than the external ectoderm (Figure 2, Supplemental file 1). This is

consistent with the histological analyses of Frank and Bleakney (1976) but seems to be in

contrast with the distribution of gland cells in medusozoans. In *Hydra*, for example, zymogen

gland cells are found exclusively in the endoderm (Haynes and Davis, 1969). These observations

suggest that the internalization of the ectoderm in anthozoans was a pivotal event in the

diversification of specialized zymogen cells. Cell products secreted from the tentacle ectoderm

may quickly become diluted in the water column, whereas the closed environment of the

gastrovascular cavity limits the space over which secreted products can diffuse; thus,

internalization created distinct selective pressures in different regions of the ectoderm. The

selective pressure to secrete digestive enzymes into the enclosed gastrovascular cavity may have

driven the development of gland cells in the internal ectoderm of anthozoans and the endoderm

of medusozoans (and many bilaterians). As such, we see no reason to homologize the ectoderm

of anthozoan mesenteries and the endodermal lining of the vertebrate midgut/pancreas

(Steinmetz et al., 2017). We consider it more likely that these tissues have converged on similar

morphologies and gene expression profiles in response to similar selection pressures associated

with extracellular digestion.

*N. vectensis trypsins have many putative functions*

The trypsin domain catalyzes the cleavage of polypeptides at internal amino acid residues and is

therefore essential for processing large proteins into smaller peptide chains. Digestive trypsins

are synthesized in secretory cells with zymogen type secretory granules where they are packaged

into vesicles for release into the gut. We show that there are at least 10 morphologically distinct

zymogen gland cell types in the pharyngeal and mesentery ectoderm of *N. vectensis* (Figure 2).

Further, we demonstrate that numerous proteases are expressed in the same tissues (Figure 3)

and that the vast majority of trypsins in *N. vectensis* encode a signal peptide (Figure 7A). Using

published single-cell expression data (Sebe-Pedros et al., 2018), we identified ten putative gland

cells that express trypsins, at least two of which also express synaptotagmin (Supplemental file

4), which facilitates fusion of the vesicle with the cell membrane during regulated secretion.

These combined features point to the expression of trypsins in secretory cells of the internal

ectoderm and strongly support a role for trypsins in extracellular protein degradation in *N.*

*vectensis*.

Numerous trypsins were expressed outside of the putative gland cells identified by Sebe-Pedros

et al (2018). At least 20 cells categorized by these authors as neural cells exhibited trypsin

expression but unlike gland cells, the maximum number of trypsins expressed by any putative

neuron is three (Supplemental file 4). We show trypsin-expressing cells differentiating very early

in development, in the invaginating pharynx/mesenteries (Figure 3). Neurons expressing

RFamide and Elav are also undergoing terminal differentiation in this tissue at this

developmental stage (Marlow et al., 2009; Nakanishi et al., 2012). Indeed, the trypsin protease

NVJ_99932 (Figure 3) is co-expressed with two other trypsins (NVJ_230861 and NVJ_130234)

in a putative neuron expressing GABA and dopamine receptors (Supplementary file 7). In

vertebrates, secretion of neurotrypsin from the pre-synaptic membrane facilitates degradation of

the extracellular matrix during synaptic plasticity and axon guidance (Almonte and Sweatt,

2011). Although 17 different trypsins were expressed in putative neurons, none of the trypsins

from *N. vectensis* clustered with human neurotrypsin (Figure 5), suggesting this function may

have been acquired independently from different ancestral trypsins.

Trypsins are important regulators of tissue remodeling; as such, upregulation of trypsins and

other proteases occurs coincident with wound healing and tissue regeneration (Daley et al.,

2008).  Recent studies of regeneration in *N. vectensis* demonstrated that a new pharynx will

regenerate from the oral ends of the mesenteries after amputation (Amiel et al., 2015) and that

many proteases are expressed abundantly during this process (Schaffer et al., 2016). Thus, the

mesenteries appear to play an important role in directing the tissue remodeling process in *N.*

*vectensis*. In support of this, a study of wound healing in response to a body wall injury

demonstrated that the mesenteries come into direct contact with damaged tissue during the

healing process (Dubuc et al., 2014b). This study also showed that two trypsins (NVJ_107554

and NVJ_112683) are among the top genes undergoing upregulation during wound healing in *N.*

*vectensis*. While NVJ_112683 was not reported in the single-cell dataset, NVJ_107554 is

expressed in two putative gland cells (metacells C12 and C19, Supplemental file 4). One of these

(metacell C12) is also the site of expression of three proteases examined by *in situ* hybridization

in this study (Figure 3). Thus, mesentery-expressed trypsins play important roles in the cell and

tissue biology of *N. vectensis* during wound healing and regeneration and these roles may vary

through ontogeny.

Beyond their roles in digestion and tissue remodeling, trypsins are an important component of

the innate immune system. In vertebrates, immune trypsins play a role in blood coagulation and

are part of the complement system which recognizes foreign particles (Heutinck et al., 2010). In

symbiotic cnidarians, immune trypsins play a role in the beneficial interaction between the host

and the alga (Poole et al., 2016). While *N. vectensis* does not host symbiotic algae, a previous

study aimed an understanding the origin of the innate immune system reported the expression of

three immune system trypsins in *N. vectensis*: MASP (NVJ_138799) and two paralogs of Factor

B (NVJ_41116, NVJ_204186), each of which were expressed in the endoderm (gastrodermis) of

juvenile polyps (Kimura et al., 2009). We found that the two factor B orthologs were also co-

expressed in a single putative gastrodermal cell (Supplemental file 4) further supporting a role

for the endoderm in the immune response of *N. vectensis*. One trypsin (NVJ_127465) was not

reported in the single-cell dataset (Sebe-Pedros et al., 2018) but was among the genes found to be

significantly upregulated in the tissue-specific transcriptome of nematosomes, which may also

play a role in the immune system of *N. vectensis* (Babonis et al., 2016). This gene clustered with

human chymotrypsin genes, not the immune system trypsins (Figure 5); as such it may have

acquired a role in the immune system secondarily.

*Trypsin functional diversity has undergone numerous expansions*

Two groups of trypsins (intracellular and non-catalytic) are found in all domains of life (Figure

4), both of which form monophyletic groups in animals (Figure 5). Trypsin diversity expanded

rapidly with the origin of animals, as representatives of two of the earliest diverging animal

lineages (ctenophores and placozoans) have at least 20 trypsins (Figure 4). The phylogeny of

animal trypsins suggests the last common ancestor of animals may have had at least six major

groups of trypsins (Figure 5). Sponges are unusual among animals in that they have only three

trypsins – two trypsin-PDZ paralogs and a trypsin-Sushi protein (Supplemental file 6). This

suggests either extensive loss of trypsins in Porifera or independent diversification of trypsins in

ctenophores and in the stem of parahoxozoa. The evolutionary history of trypsin-Sushi, trypsin-

SRCR, and trypsin-ShK proteins sheds little light on this topic; while all three associations are

found in ctenophores, trypsin-Sushi proteins are missing from placozoans and trypsin-SRCR and

trypsin-ShK proteins are missing from both sponges and placozoans (Figure 4,8, Supplemental

file 6). Presently, it is difficult to determine if these associations were either lost in sponges and

placozoans or arose independently in ctenophores and the stem of planulozoa. Considering the

association between the ShK and trypsin domains also seems to have been lost in the bilaterian

lineage (Figure 4, Supplemental file 6), we think independent gain of ShK domains in

ctenophores and cnidarians is likely.

The ancestral cnidarian may have had a far more diverse suite of trypsins than the ancestral

animal. Indeed, our data suggest there were at least 17 lineages of trypsins present in the last

common cnidarian ancestor (Figure 6) and 12 of the associations between trypsin and another

conserved domain in *N. vectensis* are specific to cnidarian lineages (Figure 8). Furthermore,

while the diversity of trypsins in *N. vectensis* rivals that of *H. sapiens* (Figure 4), there is little

conservation in the associated domains in these taxa (Figure 5B). Furthermore, within each clade

of trypsins, sequences from cnidarians and bilaterians form distinct groups, suggesting that

secretory cell function expanded independently in the cnidarian and bilaterian stem lineages.

There was extensive divergence in the trypsin gene superfamily during the diversification of

cnidarians but anthozoans seem to have undergone additional radiations in at least two trypsin

clades. Anthozoans are the most speciose group of cnidarians and are largely sessile; thus,

selection for trophic specialization and sympatric niche diversification may be stronger among

anthozoans than medusozoans.

*Numerous evolutionary mechanisms contributed to the rise of trypsin diversity*

Multidomain proteins are more common than proteins with only a single domain as domain

recombination increases versatility in protein function (Vogel et al., 2005). Selection to maintain

the catalytic activity of the trypsin domain while allowing the context in which this domain is

expressed to vary was a primary driver of diversification of this gene family. Surprisingly, there

was little conservation in trypsin-associated domains across animals, even among cnidarians

(Figure 4, Supplemental file 6), suggesting the trypsin domain underwent significant duplication

before the diversification of cnidarians and that the associated domains have been continuously

gained and lost in each cnidarian lineage. Furthermore, nearly 40% (28/72) of the proteins

comprising the *N. vectensis* tryptome have only a trypsin domain (Figure 7) yet these trypsin-

only proteins did not form a monophyletic group (Figure 5,6). These results suggest that trypsin

domains themselves may be rapidly gained and lost from evolutionarily unrelated proteins,

further underscoring the selective advantage of a trypsin domain.

Trypsins were associated with 24 other conserved protein domains in the *N. vectensis* tryptome,

only few of which were over-represented in the tryptome (Figure 4A). The ShK domain is a

short peptide found in a K-channel inhibitor originally isolated from the sea anemone

*Stichodactyla helianthus* (Castaneda and Harvey, 2009). The ShK phylogeny (Figure 7B) further

suggests this domain is gained and lost easily, as ShK domains from sister trypsins were almost

never monophyletic. Consistent with this, every ShK domain in the tryptome of *N. vectensis* was

encoded by only a single exon (Supplemental file 5), supporting the possibility of rapid evolution

through exon shuffling. Two trypsin-ShK proteins (NVJ_218669 and NVJ_218670) were found

to be sister in both phylogenies, suggesting they arose by duplication of the combined domains.

These two genes are encoded on the same scaffold and are separated by approximately 1000bp

of genomic DNA; thus, they are likely the result of a recent tandem duplication event. What role

the ShK domain plays when it is paired with the trypsin domain is not known but the

overabundance of these two combined domains in cnidarian tryptomes (Supplemental file 6)

combined with the multiple independent origins of this domain combination in *N. vectensis*

(Figure 7B) suggest the pairing provides a strong selective advantage in the biology of

cnidarians.

We also found evidence of trypsin diversification independent of the acquisition of associated

domains. One gene (NVJ_127465) encodes three trypsin domains, all of which form a

monophyletic group suggesting this gene structure arose through tandem duplication of the

trypsin domain. Likewise, we identify four cases where sister trypsins are found on the same

scaffold, suggesting tandem gene duplication. Additionally, two trypsins were found to lack

introns (NVJ_128003 and NVJ_216003), suggesting these two arose through recent

retrotransposition. These two genes are also on the same scaffold, suggesting retrotransposition

may have been followed by tandem gene duplication. The tryptome from *H. sapiens* also

includes two proteins with three trypsin domains each (Supplemental file 6). All six of these

trypsin domains from *H. sapiens* are found in the tryptase/transmembrane clade (Supplemental

file 3), whereas the three domains in NVJ_127465 group with chymotrypsins. Despite their

similar domain architecture, triple-trypsin domain proteins appear to have evolved multiple

times.

Diversification of the trypsin superfamily through gene duplication and divergence has been

continuous, suggesting an important role for trypsins in the evolutionary success of animal

lineages. Early expansions, before the diversification of animals may have promoted the origins

of the innate immune system and extracellular digestion, facilitating the evolution of large body

size and increased longevity in this lineage. Lineage-specific expansions (between genera, for

example) in the digestive trypsins enable the development of specialized of taxon-specific

tryptomes, which can support niche specialization in otherwise similar taxa (Figure 4,6,8). The

expansion of the tryptome of *N. vectensis* was facilitated by gene duplication followed by the

acquisition of additional domains (Figure 8). In some cases, the acquisition of an associated

domain by a trypsin protein occurred soon after the origin of the acquired domain (e.g., trypsin-

SRCR), while in other cases this acquisition occurred long after the associated domain first

appeared (e.g., trypsin-astacin). Indeed, we found no relationship between the age of the domain

and the age of the association with trypsin (Figure 8B), further supporting the idea that trypsin

domain architectures diversify continuously and are not dependent on the origin of novel

domains.

New genes are thought to arise rapidly *via* duplication and divergence (Tautz and Domazet-Loso,

2011) but the proportion of novel genes that become functionally integrated into signaling

networks may be very small. The factors that promote retention of new genes following

duplication are not well understood but may involve complimentary degenerative mutations

promoting subfunctionalization in the duplicates (Force et al., 1999). Under this framework,

sister trypsins that result from a recent duplication event should be expressed in differing

contexts, for example in different developmental stages or in different cell types. Consistent with

this, analysis of the single-cell expression of NVJ_218669 and NVJ_218670, which appear to

have undergone recent duplication, suggests that NVJ_218669 is expressed in a subset of the

cells that express NVJ_218670 (Supplemental file 4). Only 51 of the 72 trypsins in *N. vectensis*

were reported in the single-cell study (Sebe-Pedros et al., 2018), making a detailed analysis of

shared expression patterns in sister trypsins premature; however, a targeted study of this protein

superfamily in the future may reveal novel cis-regulatory regions, further enlightening the

processes involved in the diversification of this protein family.

*Secretory cells and the evolution of cnidarians*

Resolving the embryological origin of cnidarian gland cells will be important for understanding

the evolution of life history in Cnidaria. If the anthozoan polyp body plan is ancestral to all

cnidarians (Kayal et al., 2018), then the origin of strobilation (medusa formation) and its

associated tissue remodeling in the stem medusozoan may have necessitated the sacrifice of the

internalized tissue layers of the ancestral pharynx and mesenteries. In this case, the stem

medusozoan may have overcome this loss by shifting the development of their gland cell

population to the endoderm without sacrificing the selective advantage of secreting their

products into the gastrovascular cavity. In support of this hypothesis, gland cells in *Hydra* are

known to undergo differentiation in a location-specific manner, suggesting the identity of this

cell lineage is highly sensitive to positional cues from other cells in their environment (Siebert et

al., 2008). Furthermore, a recent study of single-cell dynamics in *Hydra* demonstrated that gland

cells acquire their identity in the endoderm only after their precursor migrates out of the

ectoderm and across the mesoglea (Siebert et al., 2018). Both of these studies point to the highly

plastic nature of gland cell identity in *Hydra* but similar analyses in more medusozoans are

needed to understand the relationship between gland cell development and cnidarian life history

evolution.

**Conclusions**

The transition from unicellular to multicellular life was marked by many transitions that enabled

functional specialization. Unicellular taxa used trypsins for intracellular protein regulation but

the origin of the regulated secretion system created new opportunities for protease activity in

multiple tissue compartments. Secretion of molecules to the extracellular space enabled the

development of the nervous, endocrine, immune, and digestive systems, and permitted spatial

and temporal separation of multiple functions performed by a single cell. The diversification of

animals was associated with a large expansion of trypsins. Trypsins with transmembrane

domains first appear in the choanoflagellates but trypsins with signal peptides did not appear

until the origin of animals. Subsequent duplication and divergence (e.g., through exon shuffling

and retrotransposition) of genes encoding secreted proteases enabled nuanced variation in the

function of these secretory cells before the increase in anatomical diversity (Figure 9).

**Methods**

*Electron microscopy, cell proliferation assay, and* in situ *hybridization*

Adult polyps were immobilized for 10 mins in 7.5% $MgCl_2$ and processed for transmission

electron microscopy as described previously (Dubuc et al., 2014a). Samples were imaged on a

Hitachi HT7700 at the University of Hawaii's Biological Electron Microscopy facility. To

identify proliferating nuclei, live polyps were incubated in 100uM EdU (in 1/3X seawater) for 30

mins at room temperature. Animals were then immobilized and fixed briefly (1.5 min) at 25C in

4% paraformaldehyde with 0.2% glutaraldehyde in phosphate buffered saline with 0.1% Tween-

20  (PTw) followed by a long fixation (60 min) in 4% paraformaldehyde in PTw at 4C. Fixed

tissues were analyzed using the Click-IT EdU kit (#C10340, Invitrogen, USA) following the

manufacturer's protocol. Nuclei were counter stained in a 30 min incubation in DAPI at room

temperature and samples were imaged on a Zeiss 710 confocal microscope at the Whitney Lab

for Marine Bioscience. To characterize the localization of target genes, we performed *in situ*

hybridization following a standard protocol for *N. vectensis* (Wolenski et al., 2013).

*Protein domain analysis*

To identify trypsin-domain proteins from *N. vectensis* we first searched the JGI protein models

using the default settings with hmmsearch (HMMER 3.1b2; http://hmmer.org/) and two target

HMMs: Trypsin (PF00089) and Trypsin_2 (PF13365). This approach yielded 99 putative

trypsin-domain containing proteins (hereafter referred to as trypsins) with an E-value $\leq$ 1e-05

(Koch et al., 2012). Where multiple partial non-overlapping trypsin domains were identified

from the same protein, we assumed these represented one single contiguous domain (Triant and

Pearson, 2015). Based on a reciprocal BLAST comparison with transcriptome data available

publicly (Babonis et al., 2016), we found 68/99 of the JGI gene models coding for trypsin

proteins were incomplete. We manually corrected these sequences using the transcriptome data

and used these corrected sequences for downstream analyses. We then used the transcriptome

data to search protein models for evidence of pseudogenes (with premature stop codons) using

the translation and alignment features in Geneious v 7.1.8 (https://www.geneious.com) and

manually examined models for duplicate predictions using the JGI genome viewer. Based on

these analyses, we removed 27 sequences, resulting in a final set of 72 curated trypsin protein

models (FASTA file available at: https://github.com/josephryan/2019-Babonis_et_al_trypsins).

We examined the domain architecture of trypsin proteins from *N. vectensis* by searching for non-

Trypsin domains in the amino acid sequences using hmmscan (HMMER 3.1b2) and the complete

Pfam-A database (downloaded Oct 27, 2017). Hmmscan identifies regions of similarity between

protein queries and domain models (protein profiles) derived from numerous proteins within the

family from a range of animals (Bateman et al 2004). Following the protocol of Koch et al

(2012), we ran hmmscan using the default parameters and report only those domains with an

independent (domain-specific) E-value ≤ 0.05 that were found in a protein containing a

significant Trypsin (or Trypsin_2) domain. Domains that overlapped by ≤ 20% were both

retained; when the overlap was >20% the domain with the lower E-value was retained. In

addition to domain analysis, we manually searched an alignment of the corrected set of trypsin

protein models from *N. vectensis* for the conserved residues that comprise the trypsin catalytic

triad (necessary for inferring protease activity): H-57, D-102, or S-195.  Finally, we searched the

corrected amino acid sequences for signal peptides and transmembrane domains using SignalP

v4.1 (Petersen et al., 2011) and TMHMM v2.0 (Krogh et al., 2001), respectively.

To characterize the origin of trypsin domain architecture, we hmmscan with the same approach

described above to identify and characterize trypsins from representatives across all domains of

life. We sampled three bilaterians (*Capitella teleta*, *Branchiostoma floridae*, *Homo sapiens*), ten

cnidarians (*N. vectensis*, *Edwardsiella lineata*, *Aiptasia pallida*, *Anthopleura elegantissima*,

*Acropora digitifera*, *Renilla renilla*, *Hydra magnipapillata*, *Calvadosia cruxmelitensis*, *Atolla*

*vanhoeffeni, Alatina alata*), three non-planulozoan animals (*Mnemiopsis leidyi*, *Amphimedon*

*queenslandica*, *Trichoplax adhaerens),* five non-metazoan eukaryotes (*Dictyostelium discoidum*,

*Schizosaccharomyces pombe*, *Capsaspora owczarzaki*, *Monosiga brevicolis*, *Salpingoeca*

*rosetta*) and a combined database of representative archeaea and bacteria (*Candidatus aquiluna*,

*Candidatus nitrosopumilus*, *Candidatus pelagibacter*, *Glaciecola pallidula*, *Marinobacter*

*adhaerens,* a marine gamma proteobacterium, and a marine group I thaumarchaeote). Protein

models were predicted from transcriptome data previously for *N. vectensis, E. lineata, A. pallida,*

*A. elegantissima, A. alatina, A. vanhoeffeni*, and *P. carnea (Babonis et al., 2016)*. Proteomes for

*R. renilla* and *C. cruxmelitensis* were predicted from the transcriptome data reported by Kayal et

al (2018) using the same methods. For all other taxa, protein models were downloaded directly

(commands available at: https://github.com/josephryan/2019-Babonis_et_al_trypsins).

*Phylotocol (phylogenetic transparency)*

All phylogenetic investigations were planned prior to running any analyses and all are reported

in this manuscript. In most cases, these analyses were outlined beforehand in a phylotocol (Ryan

and Debiasse, 2018) that is posted on our GitHub site: https://github.com/josephryan/2019-

Babonis_et_al_trypsins. Any analyses performed prior to being added to our phylotocol were

later added to the document and justified.

*Phylogenetics*

To understand the diversification of animal trypsins, we built a phylogeny using predicted

proteins from *M. leidyi, A. queenslandica, T. adhaerens, N. vectensis, E. lineata, H.*

*magnipapillata, C. teleta, B. floridae,* and *H. sapiens*. First, we used a custom script to generate

alignments from these protein files using the Trypsin HMM (commands available at:

https://github.com/josephryan/2019-Babonis_et_al_trypsins). All trees were constructed using a

maximum likelihood framework with RAxML and IQ-TREE (Kalyaanamoorthy et al., 2017;

Nguyen et al., 2015; Stamatakis, 2014). We used the model finder function with IQ-TREE (-m

MF) to determine the best substitution model for the alignment and then ran three approaches in

parallel: RAxML with 25 parsimony starting trees, RAxML with 25 random starting trees, a

single run with IQ-TREE (which, by default, uses a broad sampling of initial starting trees). We

selected the best tree by comparing the maximum likelihood scores of all three approaches.

Using the Trypsin HMM we recovered 97% (70/72) of the curated trypsin proteins from *N.*

*vectensis*. The two remaining trypsin proteins (NVJ_23745 and NVJ_203589) were recovered

using the Trypsin_2 HMM. (Note: the Trypsin_2 HMM recovered only 89% (64/72) of the

curated trypsins.) To understand the evolutionary relationships of these two proteins to the rest of

the trypsin family, we generated another phylogeny using the same procedure as above and an

alignment built using the Trypsin_2 HMM. This best tree recovered using the Trypsin_2 HMM

is provided in Supplemental file 2. After inspecting both trees, we removed sequences from *B.*

*floridae* for ease of viewing and re-ran the full analyses. All tree files and alignment files are

available on our Github site (https://github.com/josephryan/2019-Babonis_et_al_trypsins).

To evaluate whether *N. vectensis* has undergone lineage-specific expansion of trypsins or if the

common ancestor of all cnidarians had an equally diverse trypsin protein repertoire, we built a

phylogeny of trypsin proteins from cnidarians only using a subset of the proteomes listed above.

Specifically, we used four species of anthozoans (*N. vectensis, E. lineata, R. renilla, A.*

*digitifera*) and four medusozoans (*H. magnipapillata, C. cruxmelitensis, A. vanhoeffeni, A.*

*alata*). We then pruned all non-*Nematostella* taxa from this tree using Phyutility v.2.2.6 (Smith

and Dunn, 2008) to generate a tree for *N. vectensis* trypsins only. To examine the evolutionary

history of ShK domains from *N. vectensis*, we used hmmsearch with the ShK HMM and a

custom script (as above) to identify and align all ShK domains from the predicted proteome. We

then used the approach described above to produce a phylogeny of ShK domains.

**Acknowledgements**

We are grateful to Tina Carvahlo and Dr. Marilyn Dunlap of the Biological Electron Microscopy

Facility at the University of Hawai'i Manoa for their assistance with electron microscopy. This

work was supported by the National Aeronautics and Space Administration (grant

NNX14AG70G to MQM) and the National Science Foundation (grant 1542597 to JFR).

**Authors' contributions**

Study design/concept: LSB, MQM, JFR; animal/tissue methods: LSB, CE; phylogenetics: LSB,

JFR; other analyses: LSB; writing: LSB; editing: MQM, JFR, CE. All authors read and approved

the final manuscript.

**Competing interests**

The authors have no competing interests.

**Figures**

**Figure 1.** Morphology of the pharynx and mesenteries. (A) Adult polyp; the pharynx/mesentery

transition is denoted by the dotted line. (B-D) Polyp at rest; the pharyngeal ectoderm (green) is

retracted inside the oral ectoderm (yellow). (E-G) Partial eversion of pharynx occurs during

capture/handling of prey (*Artemia* sp., indicated by *). (H-J) Ingested prey passes through the

mouth and pharynx and remains in contact with the ectoderm (white arrows) of the mesenteries

during digestion; colored arrow indicates endoderm of mesenteries (pigmentation from

consumption of *Artemia*). (K) Cnidocytes (black arrow) and gland cells (green arrow) are

restricted to the ectoderm of the mesenteries. C,D,F,G,K are DIC micrographs. D,G are false

colored. E,H are oral views, the remaining images are lateral views. Dotted lines in D,G,K

denote position of mesoglea. White arrowheads point to the mouth, black arrowheads denote

transition from external to internal ectoderm.

**Figure 2.** Fine structure of the pharyngeal ectoderm and ectodermal mesenteries. (A)

Diagrammatic representation of regions where thin sections were examined through the pharynx

(I) and at two positions in the mesenteries (II, III). Top – sagittal view of a primary polyp,

middle – cross-section at position I, bottom – partial cross sections at positions II (left) and III

(right). (B-F) TEMs of a region of the pharyngeal ectoderm (I) indicated by the box in A. (B)

Two zymogen gland cells are false colored for emphasis. (C) Cross sections of cilia emerging

from the pharyngeal ectoderm into the pharyngeal canal. (D) Mucus-secreting cell, false colored.

(E) Ten zymogen-type gland cells and two cnidocytes. (F) Electron-dense vesicles (black

arrowheads in E,F,N,O) in the apex of non-glandular ectodermal cells. (G) SEM of a mesentery

from position II in panel A. The ectodermal (EC) portion has two parts: the ciliated tract (black

line) and cnidoglandular tract (white line); the endodermal (EN) portion is false colored. (H-J)

TEMs of the cnidoglandular tract of a mesentery from position II in A. Sections correspond to

the position of the dotted line in G. (I) One zymogen type gland cell is false colored. (J) Some

zymogen vesicles have heterogeneous contents (black arrow). (K) 3D reconstruction of a

confocal z-stack through the ciliated tract of a mesentery from position II in A; pink – nuclei of

proliferating cells (EdU), blue – quiescent nuclei (DAPI). (L) TEM section through the ciliated

tract at the position indicated by the dotted line in K and by the box in H. Nuclei corresponding

to panel K are false colored pink. (M-O) TEMs of a mesentery lacking a ciliated tract, position

III in panel A. (M) Endoderm is false colored. (N) Zymogen gland cells are false colored. (O)

Ciliary rootlet (black arrow) in the apex of a zymogen gland cell. Scale bars: white line – 10 um,

black line – 500 nm, grey line (panels G, K) – 50 um. Panels B and H are composites of multiple

micrographs. White arrowheads indicate cnidocytes throughout.

**Figure 3.** Expression of proteases in *N. vectensis*. (A-C) *In situ* hybridization of proteases in the

(A) M family – metallopeptidases, astacins, and carboxypeptidases and (B) the S1 family –

trypsins during early development. The oral pole is to the left in all images; black arrows indicate

expression in the ectoderm of the pharynx. (C) Double fluorescent *in situ* hybridization showing

the co-expression (yellow arrow) of two metalloprotease genes (Nv_88668 and Nv_2109) in the

aboral ectoderm of the pharynx and independent expression of both genes in individual cells of

the ectodermal mesentery (red and green arrows). Nuclei are labeled with DAPI. (D) Single-cell

expression of genes from panels A and B in metacells identified by Sebe-Pedros et al (2018).

Metacells C9-C12 are part of the "gland" cell cluster and C26 is part of the "neuron" cell cluster.

Two genes (NVJ_200868 and NVJ_2109) are clearly expressed in the ectodermal mesenteries

but were not reported by Sebe-Pedros et al 2018 (indicated by dotted lines). (E) Histogram of the

number of expressed trypsins per cell (from Sebe-Pedros et al 2018); red font highlights a small

second mode in the distribution. (F) Scatterplot of the number of cells exhibiting co-expression

of multiple trypsins as a function of the number of total cells in which a particular trypsin is

expressed.

**Figure 4.** Characterization of the 72 proteins comprising the *N. vectensis* tryptome. (A)

Abundance of the 24 trypsin-associated domains in the tryptome (black) and the remaining

proteins in the predicted proteome (white). Numbers inside the bars reflect domain counts in

both datasets; grey numbers to the right of the bars indicate total proteins containing the

indicated domain. Pfam domains are listed on the left; see Supplemental file 4 for pfam domains

IDs. (B) Phylogenetic distribution of trypsin domain architectures. Domain architecture models

from the *N. vectensis* tryptome are shown on the right; proteins that differ only in the number of

copies of associated domains were collapsed into a single row and these protein IDs are

underlined. The number of proteins with identical domain composition are indicated for each

taxon in the table. Actinarians (sea anemones) are highlighted in blue. [1]Maximum number of

trypsins from any one taxon in the bacteria/archaea database. [2]El,Ae,Ap,Av have trypsin-TSP_1-

ShK proteins which were included in this count. Bac – bacteria/archaea, Dd – *Dictyostelium*

*discoidum*, Sp – *Schizosaccharomyces pombe*, Co – *Capsaspora owczarzaki*, Sr – *Salpingoeca*

*rosetta*, Mb – *Monosiga brevicollis*, Ml – *Mnemiopsis leidyi*, Aq – *Amphimedon queenslandica*,

Ta – *Trichoplax adhaerens*, El – *Edwardsiella lineata*, Ap – *Aiptasia pallida*, Ae – *Anthopleura*

*elegantissima*, Ad – *Acropora digitifera*, Rr – *Renilla renilla*, Hm – *Hydra magnipapillata*, Pc –

*Podocoryna carnea*, Av – *Atolla vanhoeffeni*, Aa – *Alatina alata*, Cc – *Calvadosia*

*cruxmelitensis*, Ct – *Capitella teleta*, Hs – *Homo sapiens*

**Figure 5.** Evolutionary history of animal trypsins. (A) Phylogeny of animal trypsins; clades are

named after the orthologs from *H. sapiens*. Domain architectures for *N. vectensis* proteins are

shown. Proteins with multiple trypsin domains are polyphyletic; in such cases, the domain

architecture diagram points to a single trypsin domain and the position of the other trypsin

domains is indicated by open/closed circles. Open stars indicate clades that descended from

single genes present in the last common ancestor of animals. N indicates the position of human

neurotrypsin (NP_003610.2). The trypsin-death protein groups with trypsin-PDZ proteins but is

not found on this tree. *N. vectensis* proteins characterized by *in situ* hybridization are indicated:

arrows (this study), † (Kimura et al 2009), or * (Steinmetz et al 2017). (B) Comparison of the

domain content of trypsins from *N. vectensis* and *H. sapiens*. The non-catalytic trypsins do not

have associated domains and are not included. The tryptase/transmembrane trypsins are not

represented in *N. vectensis* and are also not included. The "pancreatic" group includes granzyme,

kallikrein, HGF, and elastase.

**Figure 6.** Gene phylogeny of cnidarian trypsin domains. Domain architectures for *N. vectensis*

proteins are shown; domains are colored as shown in Figure 4A. Proteins with multiple trypsin

domains are polyphyletic; in such cases, the domain architecture diagram points to a single

trypsin domain and the position of the other trypsin domains is indicated by open/closed circles.

Open stars indicate clades that descended from single genes present in the last common ancestor

of cnidarians; closed stars represent possible anthozoan expansions. Clades are labeled as in

Figure 5A; unshaded clades do not include any taxa from Figure 5 and cannot be resolved. The

"pancreatic" group includes proteins that are sister to human granzyme, kallikrein, HGF, and

elastase. Chymotrypsins are not monophyletic on this tree but are shaded together to facilitate

comparison with Figure 5A.

**Figure 7**. Relationships among trypsin and ShK proteins in *N. vectensis*. (A) Phylogeny of *N.*

*vectensis* trypsin domains; domain architecture diagrams are shown to the right. Proteins with

multiple trypsin domains appear multiple times in the tree; in these cases, the domain represented

at each position in the tree is bolded. SP – signal peptide, TM - transmembrane domain. Trypsin

domains lacking H-57, D-102, or S-195 are designated non-catalytic (see Supplemental file 4).

Hypothesized ancestral states (models with dotted lines) are indicated in two positions for

trypsin-ShK proteins; a revised hypothesis is indicated by the trypsin-ShK model with solid

lines. (B) Phylogeny of *N. vectensis* ShK domains from the predicted proteome. Proteins with

multiple ShK domains appear multiple times in the tree; in these cases, the domain represented at

each position is indicated with an arrowhead. Proteins that have both trypsin and ShK domains

are colored green. Colored symbols to the right of the domain architectures are provided to

facilitate comparisons of identical proteins in panels A and B. Domains are colored as shown in

Figure 4A.

**Figure 8.** Evolutionary history of the tryptome. (A) Domain architectures at ancestral nodes

indicate the origin of each domain combination; domains are reported at nodes where they

appear for the first time with an independent (domain-specific) E-value ≤ 0.05. The origin of the

SRCR domain and the origin of its association with trypsin are indicated by white arrows; black

arrows indicate the origin of the astacin domain and the association between astacin and trypsin.

E – Edwardsiidae only (*Edwardsiella* + *Nematostella*), N – *Nematostella* only. WAP domains (*)

either evolved twice (in *D. discoideum* and the ancestor of animals) or this domain was lost in

the intervening lineages. The DIM domain (†) has a complex distribution in bacteria, fungi,

cnidarians, and insects. (B) There is no relationship between the phlyostratigraphic age of the

domain and the age of its association with trypsin.

**Figure 9.** Secretory vesicles permit functional expansion without anatomical variation. Blue and

white cells reflect the intracellular expression of blue and white gene products. The origin of the

signal peptide directing proteins to the regulated secretion system/secretory vesicle (white arrow)

permitted segregation of gene products into two distinct compartments: intracellular and

intravesicular. The subsequent duplication of only few genes (black arrow) could result in the

acquisition of numerous new cell types through unique and combinatorial expression. Green –

co-expression of blue and yellow, purple – co-expression of blue and red, brown – co-expression

of blue, yellow, and red.

**Supplemental Material**

**Supplemental file 1:** Gland cells of the external ectoderm. (A-C) Mucus cells (false colored

yellow) in the tentacle ectoderm. (D) Zymogen cell (false colored green) in the body wall

ectoderm. White arrows point to cnidocytes, black arrowheads point to electron dense apical

vesicles in cells adjacent to gland cells. A, B – SEM, C,D – TEM. Scale bars: black – 5um, white

– 10um.

**Supplemental file 2**: Phylogeny of Trypsin_2 domains across animals. Protein models for *N.*

*vectensis* are shown. NVJ_203589 and NVJ_23745 were not detected by the trypsin HMM and

do not appear in Figure 5. Colors as in Figure 4A.

**Supplemental file 3:** Human trypsin domain architecture mapped on the animal trypsin

phylogeny. Proteins with multiple trypsin domains are polyphyletic; in such cases, the diagram

points to a single trypsin domain and the position of the other trypsin domains is indicated by

symbols. *N. vectensis* proteins characterized by *in situ* hybridization are indicated by arrows (this

study), † (Kimura et al., 2009), or * (Steinmetz et al., 2017).

**Supplemental file 4**: Excel file tabulating: presence/absence of signal peptides and

transmembrane domains in *N. vectensis* trypsins, amino acid sequences for trypsin catalytic

domains for all taxa, pfam IDs for all domains, and a summary of single-cell expression of

trypsins from Sebe-Pedros et al. (2018).

**Supplemental file 5:** All trypsin domains are encoded by multiple exons in *N. vectensis*

(excluding the NVJ_128003 and NVJ_216003) but many of the associated domains are encoded

by a single exon (indicated by triangle). Domains that span intron/exon boundaries by 10 or

fewer nucleotides were considered to be encoded by a single exon.

**Supplemental file 6:** Trypsin protein domain architectures from all taxa examined in this study.

**Supplemental file 7:** Excel file of trypsin protein IDs from all taxa examined in this study.

**References**

Almonte, A.G., Sweatt, J.D., 2011. Serine proteases, serine protease inhibitors, and protease-
activated receptors: Roles in synaptic function and behavior. Brain Res 1407, 107-122.

Amiel, A.R., Johnston, H.T., Nedoncelle, K., Warner, J.F., Ferreira, S., Rottinger, E., 2015.
Characterization of Morphological and Cellular Events Underlying Oral Regeneration in
the Sea Anemone, Nematostella vectensis. Int J Mol Sci 16, 28449-28471.

Babonis, L.S., Martindale, M.Q., Ryan, J.F., 2016. Do novel genes drive morphological novelty?
An investigation of the nematosomes in the sea anemone Nematostella vectensis. Bmc
Evol Biol 16.

Castaneda, O., Harvey, A.L., 2009. Discovery and characterization of cnidarian peptide toxins
that affect neuronal potassium ion channels. Toxicon 54, 1119-1124.

Daley, W.P., Peters, S.B., Larsen, M., 2008. Extracellular matrix dynamics in development and
regenerative medicine. J Cell Sci 121, 255-264.

Domazet-Loso, T., Brajkovic, J., Tautz, D., 2007. A phylostratigraphy approach to uncover the
genomic history of major adaptations in metazoan lineages. Trends Genet 23, 533-539.

Dubuc, T.Q., Dattoli, A.A., Babonis, L.S., Salinas-Saavedra, M., Rottinger, E., Martindale,
M.Q., Postma, M., 2014a. In vivo imaging of Nematostella vectensis embryogenesis and
late development using fluorescent probes. Bmc Cell Biol 15.

Dubuc, T.Q., Traylor-Knowles, N., Martindale, M.Q., 2014b. Initiating a regenerative response;

cellular and molecular features of wound healing in the cnidarian Nematostella vectensis.

Bmc Biol 12.

Finnerty, J.R., Pang, K., Burton, P., Paulson, D., Martindale, M.Q., 2004. Origins of bilateral

symmetry: Hox and dpp expression in a sea anemone. Science 304, 1335-1337.

Force, A., Lynch, M., Pickett, F.B., Amores, A., Yan, Y.L., Postlethwait, J., 1999. Preservation

of duplicate genes by complementary, degenerative mutations. Genetics 151, 1531-1545.

Frank, P., Bleakney, J., 1976. Histology and sexual reproduction of the anemone Nematostella

vectensis Stephenson 1935. Journal of Natural History 10, 441-449.

Fritzenwanker, J.H., Genikhovich, G., Kraus, Y., Technau, U., 2007. Early development and axis

specification in the sea anemone Nematostella vectensis. Dev Biol 310, 264-279.

Haynes, J.F., Davis, L.E., 1969. Ultrastructure of Zymogen Cells in Hydra Viridis. Zeitschrift

Fur Zellforschung Und Mikroskopische Anatomie 100, 316-&.

Heutinck, K.M., ten Berge, I.J.M., Hack, C.E., Hamann, J., Rowshani, A.T., 2010. Serine

proteases of the human immune system in health and disease. Mol Immunol 47, 1943-

1955.

Jahnel, S.M., Walzl, M., Technau, U., 2014. Development and epithelial organisation of muscle

cells in the sea anemone Nematostella vectensis. Front Zool 11.

Kalyaanamoorthy, S., Minh, B.Q., Wong, T.K.F., von Haeseler, A., Jermiin, L.S., 2017.

ModelFinder: fast model selection for accurate phylogenetic estimates. Nat Methods 14,

587-+.

Kayal, E., Bentlage, B., Pankey, M.S., Ohdera, A.H., Medina, M., Plachetzki, D.C., Collins,

A.G., Ryan, J.F., 2018. Phylogenomics provides a robust topology of the major cnidarian

lineages and insights on the origins of key organismal traits. Bmc Evol Biol 18.

Kimura, A., Sakaguchi, E., Nonaka, M., 2009. Multi-component complement system of

Cnidaria: C3, Bf, and MASP genes expressed in the endodermal tissues of a sea

anemone, Nematostella vectensis. Immunobiology 214, 165-178.

Koch, B.J., Ryan, J.F., Baxevanis, A.D., 2012. The Diversification of the LIM Superclass at the

Base of the Metazoa Increased Subcellular Complexity and Promoted Multicellular

Specialization. Plos One 7.

Krogh, A., Larsson, B., von Heijne, G., Sonnhammer, E.L.L., 2001. Predicting transmembrane

protein topology with a hidden Markov model: Application to complete genomes. J Mol

Biol 305, 567-580.

Kusserow, A., Pang, K., Sturm, C., Hrouda, M., Lentfer, J., Schmidt, H.A., Technau, U., von

Haeseler, A., Hobmayer, B., Martindale, M.Q., Holstein, T.W., 2005. Unexpected

complexity of the Wnt gene family in a sea anemone. Nature 433, 156-160.

Leclere, L., Rentzsch, F., 2014. RGM Regulates BMP-Mediated Secondary Axis Formation in

the Sea Anemone Nematostella vectensis. Cell Rep 9, 1921-1930.

Long, M., Betran, E., Thornton, K., Wang, W., 2003. The origin of new genes: Glimpses from

the young and old. Nat Rev Genet 4, 865-875.

Magie, C.R., Daly, M., Martindale, M.Q., 2007. Gastrulation in the cnidarian Nematostella

vectensis occurs via invagination not ingression. Dev Biol 305, 483-497.

Marlow, H.Q., Srivastava, M., Matus, D.Q., Rokhsar, D., Martindale, M.Q., 2009. Anatomy and

Development of the Nervous System of Nematostella vectensis, an Anthozoan Cnidarian.

Dev Neurobiol 69, 235-254.

Moiseeva, E., Rabinowitz, C., Paz, G., Rinkevich, B., 2017. Histological study on maturation,

fertilization and the state of gonadal region following spawning in the model sea

anemone, Nematostella vectensis. Plos One 12.

Moran, Y., Praher, D., Schlesinger, A., Ayalon, A., Tal, Y., Technau, U., 2013. Analysis of

Soluble Protein Contents from the Nematocysts of a Model Sea Anemone Sheds Light on

Venom Evolution. Mar Biotechnol 15, 329-339.

Nakanishi, N., Renfer, E., Technau, U., Rentzsch, F., 2012. Nervous systems of the sea anemone

Nematostella vectensis are generated by ectoderm and endoderm and shaped by distinct

mechanisms. Development 139, 347-357.

Nguyen, L.T., Schmidt, H.A., von Haeseler, A., Minh, B.Q., 2015. IQ-TREE: A Fast and

Effective Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. Mol

Biol Evol 32, 268-274.

Page, M.J., Di Cera, E., 2008a. Evolution of Peptidase Diversity. J Biol Chem 283, 30010-

30014.

Page, M.J., Di Cera, E., 2008b. Serine peptidases: Classification, structure and function. Cell

Mol Life Sci 65, 1220-1236.

Petersen, T.N., Brunak, S., von Heijne, G., Nielsen, H., 2011. SignalP 4.0: discriminating signal

peptides from transmembrane regions. Nat Methods 8, 785-786.

Poole, A.Z., Kitchen, S.A., Weis, V.M., 2016. The Role of Complement in Cnidarian-
Dinoflagellate Symbiosis and Immune Challenge in the Sea Anemone Aiptasia pallida.
Front Microbiol 7.

Putnam, N.H., Srivastava, M., Hellsten, U., Dirks, B., Chapman, J., Salamov, A., Terry, A.,
Shapiro, H., Lindquist, E., Kapitonov, V.V., Jurka, J., Genikhovich, G., Grigoriev, I.V.,
Lucas, S.M., Steele, R.E., Finnerty, J.R., Technau, U., Martindale, M.Q., Rokhsar, D.S.,
2007. Sea anemone genome reveals ancestral eumetazoan gene repertoire and genomic
organization. Science 317, 86-94.

Renfer, E., Amon-Hassenzahl, A., Steinmetz, P.R.H., Technau, U., 2010. A muscle-specific
transgenic reporter line of the sea anemone, Nematostella vectensis. P Natl Acad Sci
USA 107, 104-108.

Ryan, J.F., Debiasse, M.B., 2018. Phylotocol: Promoting transparency and overcoming bias
through publicly posted, a priori methodological protocols in phylogenetics. Integr Comp
Biol 58, E410-E410.

Schaffer, A.A., Bazarsky, M., Levy, K., Chalifa-Caspi, V., Gat, U., 2016. A transcriptional time-
course analysis of oral vs. aboral whole-body regeneration in the Sea anemone
Nematostella vectensis. Bmc Genomics 17.

Sebe-Pedros, A., Saudemont, B., Chomsky, E., Plessier, F., Mailhe, M.P., Renno, J., Loe-Mie,
Y., Lifshitz, A., Mukamel, Z., Schmutz, S., Novault, S., Steinmetz, P.R.H., Spitz, F.,
Tanay, A., Marlow, H., 2018. Cnidarian Cell Type Diversity and Regulation Revealed by
Whole-Organism Single-Cell RNA-Seq. Cell 173, 1520-+.

Siebert, S., Anton-Erxleben, F., Bosch, T.C.G., 2008. Cell type complexity in the basal metazoan

Hydra is maintained by both fly stem cell based mechanisms and transdifferentiation.

Dev Biol 313, 13-24.

Siebert, S., Farrell, J.A., Cazet, J.F., Abeykoon, Y.L., Primack, A.S., Schnitzler, C.E., Juliano,

C.E., 2018. Stem cell differentiation trajectories in Hydra resolved at single-cell

resolution. bioRxiv, 460154.

Smith, S.A., Dunn, C.W., 2008. Phyutility: a phyloinformatics tool for trees, alignments and

molecular data. Bioinformatics 24, 715-716.

Stamatakis, A., 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of

large phylogenies. Bioinformatics 30, 1312-1313.

Steinmetz, P.R.H., Aman, A., Kraus, J.E.M., Technau, U., 2017. Gut-like ectodermal tissue in a

sea anemone challenges germ layer homology. Nat Ecol Evol 1, 1535-1542.

Tautz, D., Domazet-Loso, T., 2011. The evolutionary origin of orphan genes. Nat Rev Genet 12,

692-702.

Triant, D.A., Pearson, W.R., 2015. Most partial domains in proteins are alignment and

annotation artifacts. Genome Biol 16.

Vogel, C., Teichmann, S.A., Pereira-Leal, J., 2005. The relationship between domain duplication

and recombination. J Mol Biol 346, 355-365.

Wijesena, N., Simmons, D.K., Martindale, M.Q., 2017. Antagonistic BMP-cWNT signaling in

the cnidarian Nematostella vectensis reveals insight into the evolution of mesoderm. P

Natl Acad Sci USA 114, E5608-E5615.

Wolenski, F.S., Layden, M.J., Martindale, M.Q., Gilmore, T.D., Finnerty, J.R., 2013.

Characterizing the spatiotemporal expression of RNAs and proteins in the starlet sea

anemone, Nematostella vectensis. Nat Protoc 8, 900-915.

## Figure 1

Figure 2

Figure 3

Figure 4

Figure 6

**Anthozoa**
*N. vectensis*
*E. lineata*
*R. renilla*
*A. digitifera*

**Medusozoa**
*H. magnipapillata*
*C. cruxmelitensis*
*A. vanhoeffeni*
*A. alatina*

chymotrypsinogen

non-catalytic

intracellular

immune/coagulation

"pancreatic"

# Figure 7

Figure 8

Trypsin | TSP_1 | Ldl_a | VWA | ShK | Ig_2 | Sushi | cEGF | EGF_CA | CUB | WSC | SRCR | PLAT | Lustrin | Death | WAP | PDZ | Astacin | FXa | I-set | Ldl_b | MAM | Thy | DIM | F5F8

# Figure 9