Disease Resistance Genetics and Genomics in Octoploid Strawberry

Christopher Barbey[*,‡], Seonghee Lee[†], Sujeet Verma[†], Kevin A. Bird[§,**], Alan E. Yocca[††], Patrick P. Edger[§,**], Steven J. Knapp[‡‡], Vance M. Whitaker[†] and Kevin M. Folta[*,‡]

[*] Horticultural Sciences Department, University of Florida, Gainesville FL USA

[†] Gulf Coast Research and Education Center, University of Florida, Wimauma, FL USA

[‡] Graduate Program in Plant Molecular and Cellular Biology, University of Florida, Gainesville, FL USA

[§] Department of Horticulture, Michigan State University, East Lansing, MI, USA

[**] Ecology, Evolutionary Biology and Behavior, Michigan State University, East Lansing, MI, USA

[††] Department of Plant Biology, Michigan State University, East Lansing, MI, USA

[‡‡] Department of Plant Sciences, University of California, Davis, Davis, CA, USA

Octoploid Strawberry R-genes

Strawberry, Disease Resistance, R-gene, eQTL, Subgenome Dominance, RenSeq

Dr. Christopher Barbey

Horticultural Sciences Department

University of Florida

Gainesville FL 32611

208-585-7967

cbarbey@ufl.edu

ABSTRACT

Octoploid strawberry (*Fragaria ×ananassa*) is a valuable specialty crop, but profitable production and availability are threatened by many pathogens. Efforts to identify and introgress useful disease resistance genes (R-genes) in breeding programs are complicated by strawberry's complex octoploid genome. Recently-developed resources in strawberry, including a complete octoploid reference genome and high-resolution octoploid genotyping, enable new analyses in strawberry disease resistance genetics. This study characterizes the complete R-gene collection in the genomes of commercial octoploid strawberry and two diploid ancestral relatives, and introduces several new technological and data resources for strawberry disease resistance research. These include octoploid R-gene transcription profiling, dN/dS analysis, eQTL analysis and RenSeq analysis in cultivars. Octoploid fruit transcript expression quantitative trait loci (eQTL) were identified for 77 putative R-genes. R-genes from the ancestral diploids *Fragaria vesca* and *Fragaria iinumae* were compared, revealing differential inheritance and retention of various octoploid R-gene subtypes. The mode and magnitude of natural selection of individual *F. ×ananassa* R-genes was also determined via dN/dS analysis. R-gene sequencing using enriched libraries (RenSeq) has been used recently for R-gene discovery in many crops, however this technique somewhat relies upon *a priori* knowledge of desired sequences. An octoploid strawberry capture-probe panel, derived from the results of this study, is validated in a RenSeq experiment and is presented for community use. These results give unprecedented insight into crop disease resistance genetics, and represent an advance towards exploiting variation for strawberry cultivar improvement.

INTRODUCTION

Cultivated strawberry (*Fragaria ×ananassa*) is an important specialty crop that is cultivated world-wide for its sweet and flavorful fruit. However, marketable yields and post-harvest quality are significantly affected by disease. The strawberry fruit presents a vulnerable target for microbial pathogens (Farzaneh *et al.*, 2015), as it is soft, moist, carbohydrate rich, and subject to damage from forces as seemingly innocuous as rain (Herrington *et al.*, 2011). Genetic disease resistance has been a long-standing breeding priority. While breeders have made progress in producing varieties with tolerance to some pathogens, growers remain dependent on exogenous crop protection strategies to reduce pathogen loads (Cordova *et al.*, 2017).

Plant R-genes are mediators of resistance to specific pathogens via effector triggered immunity, which results in the hypersensitive response and cell death (Amil-Ruiz *et al.*, 2018). R-genes require a high degree of regulation to maintain homeostatic transcript levels to mitigate off-target protein interactions (Hammond-Kosack and Jones, 1997). For this reason, many classes of functional R-genes are expressed at low levels unless elicited by pathogens (Lai and Eulgem, 2017), contributing to the challenges of R-gene genomic and functional annotation. About 60% of characterized plant R-genes contain nucleotide-binding (NB-ARC) and leucine-rich-repeat (LRR) domains, and are referred to NLR genes (Funk *et al.*, 2018). Plant R-genes are frequent targets for genetic improvement via breeding and genetic engineering (Djian-Caporalino *et al.*, 2014, Baumgartner *et al.*, 2015), and gene editing methods may accelerate their introduction into already-elite varieties. However, progress has been hindered because relatively few R-genes conferring novel resistance have been characterized (Amil-Ruiz *et al.*, 2018). This problem is appreciable in strawberry, where the genetic complexity of octoploid cultivars presents unique challenges for functional identification and cloning of causal variants. An analysis of diploid R-genes across the Rosaceae genus was previously conducted (Arya *et al.*, 2014). New genetic resources for high-resolution genotyping in octoploid strawberry have resulted in the recent identification of several disease resistance loci (Roach *et al.*, 2016, Mangandi *et al.*, 2017, Anciro *et al.*, 2018, Pincot *et al.*, 2018, Salinas *et al.*, 2018, Verma *et al.*, 2018). However, the specific genes mediating resistance in these QTL intervals typically remain unresolved, as genomic resources for octoploid strawberry have not kept pace with genetic mapping.

Cultivated strawberry shares common ancestors with the extant diploid species *F. vesca, F. iinumae, F. nipponica,* and *F. viridis* (Edger *et al.*, 2019) . A high-quality octoploid strawberry

genome has been recently developed (Edger *et al.*, 2019), enabling new kinds analyses and

improved resolution compared with previous studies involving *Fragaria* NLRs (Jia *et al.*, 2015,

Zhong *et al.*, 2018). Analysis of this *F. ×ananassa* 'Camarosa' genome identified the repertoire

of octoploid R-gene sequences and further demonstrated a general genomic retention bias towards

F. vesca-like sequences (Edger *et al.*, 2019).

This research compares R-genes from octoploid strawberry with its diploid ancestors and provides

additional analysis into the genetic control of R-gene expression and retention patterns. Additional

bias towards retention of *F. vesca*-like R-genes was detected in octoploid strawberry, beyond the

bias observed in non-R-gene coding sequences. This finding provides insight into potential

practical drivers of biased gene retention. Conserved domains were compared to describe specific

R-gene phylogenic relationships. The octoploid genome was used to assemble 61 fruit

transcriptomes, and used to discover subgenomic expression quantitative trait loci (eQTL) for R-

genes expressed in octoploid fruit. Data from the octoploid 'Camarosa' strawberry gene expression

atlas (Sánchez-Sevilla et al., 2017) was also used to determine R-gene transcript accumulation

throughout the strawberry plant.

Resistance gene enrichment and sequencing (RenSeq) is an advantageous method for sequencing

R-genes (Andolfo *et al.*, 2014), and is likely to be very useful for *de novo* resolution of causal

mutations (Witek *et al.*, 2016). This method can be used to identify casual mutations within

existing disease resistance QTL. For this purpose, a novel octoploid strawberry RenSeq capture

probe library was designed using the R-genes identified in this analysis. This panel was

experimentally validated using the University of Florida breeding germplasm. The results

demonstrate robust capture and resequencing of octoploid and diploid R-genes using only short

second-generation sequence reads and with relatively deep genomic multiplexing.

This report characterizes the complete R-gene collection in the genomes of commercial octoploid

strawberry and two diploid ancestral relatives, providing the genome-level resolution necessary

for fully exploiting genetic disease resistance in strawberry. This research introduces several new

technology and data resources that now may be applied in study of strawberry disease resistance.

RESULTS

Octoploid and Diploid R-genes

The genomes of octoploid 'Camarosa', diploid *F. vesca,* and diploid *F. iinumae* were analyzed

for R-gene signatures. The *F.iinumae* genome was selected to represent the closely-related 'old

world' diploid ancestors *F.iinumae, F. nipponica* and *F. viridis*, which each have highly similar

but fragmented genomic assemblies.

Putative R-genes were identified based on protein domain and motif analysis, which identified

gene models with traditional NLR-type domains, including coiled coil (CC), Toll Interleukin

Receptor-like (TIR), Leucine Rich Repeat (LRR), and Nucleotide Binding - APAF-1 (apoptotic

protease-activating factor-1), R proteins and CED-4 (*Caenorhabditis elegans* death-4 protein)

(NB-ARC). Gene models with NLR-type domains that are not highly specific to NLR sequences

(e.g. LRR domains) were included if there was also supporting evidence of an additional NLR-

associated motif. BLAST2GO annotated disease resistance associated genes not meeting these

criteria were analyzed manually, leading to the intentional inclusion of many putative Receptor-

like Kinase (RLK-type) R-genes in this analysis.

Octoploid *F. ×ananassa* 'Camarosa' carries 1,962 putative resistance genes (1.82% of all genes),

including 975 complete or truncated NLR genes (Table 1). NLR gene content is similar in genic

proportion to the 367 complete or truncated NLR genes in *F. vesca* (1.09% of all genes) and 387

in *F. iinumae* (0.5% of all genes) (Table 1). Traditional NLR domains including Coiled Coil (CC),

Toll Interleukin Receptor-like (TIR), Leucine Rich Repeat (LRR), and Nucleotide Binding -

APAF-1 (apoptotic protease-activating factor-1), R proteins and CED-4 (*Caenorhabditis*

elegans death-4 protein) (NB-ARC) domains comprise the majority of domain classes in all

predicted resistance gene models in diploid and octoploid strawberry accessions (Figure 1). In

many categories, the three genomes show somewhat dissimilar ratios of relative NLR-subtype

content (Table 1). These include biases towards TIR-only proteins in *F. vesca* and CNL-type R-

genes in *F. innumae*. Octoploid 'Camarosa' is proportionally intermediate for many NLR

categories, relative to *F. vesca and F. innumae*. A high proportion of TIR-NB and TIR-NB-LRR-

type NLR-genes is observed in 'Camarosa'. However, the overall proportion of TIR-containing

genes is similar. The Resistance to Powdery Mildew 8 (RPW8) domain, a disease resistance

domain associated with broad-spectrum mildew resistance in Arabidopsis, appears frequently in

strawberry and is present in 136 (13.9%) of octoploid NLRs (Table 1). Basic trends in NLR-

subtype genomic content in 'Camarosa' does not more strongly resemble either *F. vesca* or *F.*
innumae.

In 'Camarosa', 750 genes contain at least one NB-ARC domain, which is the most
characteristic domain of NLR-type R-genes (Table 1). The ratio of 'Camarosa' NB-ARC-
containing genes to total predicted gene content (1:144) is higher than in *F. vesca* (1:171) and *F.*
iinumae (1:262), possibly indicating diversifying selection of NLR genes in octoploid *F.*
×*ananassa*. A substantial number of atypical domains are present on strawberry R-genes, including
malectin-like carbohydrate-binding domains, RNA-binding domains, transcription factor-like
WRKY and F-box domains, and several types of protein kinase domains (Figure 1).

Tandem clusters of R-genes were observed in all three of the analyzed strawberry genomes.
The phenomena of R-gene expansion through tandem duplication is exemplified in the RPW8-
containing R-gene class. Of the seventy-one RPW-containing R-genes in *F. vesca*, all but seven
reside in one of a few genomic clusters (Figure S1). The major RPW cluster observed in *F. vesca*
chromosome 1 is strongly retained in 'Camarosa' (Figure S2). Similar R-gene hotspots are
observed throughout the diploid and octoploid strawberry reference genomes. Genome annotations
for all NLR domains in 'Camarosa' and *F. vesca* genotype Hawaii 4 v2.0 (Tennessen *et al.*, 2014)
are provided in File S1. Annotations are also available on the JBrowse web-based genome browser
at the Genome Database for the Rosaceae (www.rosaceae.org).

Phylogenetic Analysis of Strawberry R-genes
The conserved NB-ARC domains from 'Camarosa', *F. vesca*, and *F. iinumae* were compared via
maximum likelihood analysis to examine evolutionary trends among NLR genes. NB-ARCs from
all three genomes phylogenetically organized mostly according to their extended R-gene domain
structures, with TNLs, CNLs, and RPW-associated NB-ARCs forming clades based on this criteria
(Figure 2). Minor NLR subtypes, such as WRKY-associated NLR genes, also sorted into a unique
subclade based only on NB-ARC sequence. Multiple distinct clades with identical domain
architectures were detected, and in a few cases these subclades are relatively distant from one
another.

R-gene Transcript Accumulation
Raw RNAseq expression data from different tissues of 'Camarosa', derived from the octoploid
strawberry gene expression atlas (Sánchez-Sevilla *et al.*, 2017), were reassembled based on the

'Camarosa' genome. A majority of 'Camarosa' NLR genes are predominantly expressed in the

roots and leaves (Figure 3A-B). Comparatively few NLRs are predominantly or specifically

expressed in the mature receptacle. Most NLR type R-genes are broadly specific to only one or

two tissues. Expressed NLR genes from root, leaf, green and white receptacles show remarkably

poor overlap. Overall NLR transcript accumulation is correlated across all stages of receptacle

development, with strongest expression in the earlier stages and decreasing with maturity in both

the receptacle and achene. Complete R-gene expression values for each tissue are provided in

Table S1.

Mature receptacle transcriptomes from 61 field-grown individuals of three octoploid populations

reveal broadly stable R-gene expression levels (SD 0.09). R-genes comprise 1.8% of the predicted

gene models in the 'Camarosa' genome, but represent an average of only 0.48% of total transcripts

in the mature receptacle (Figure 3C). Minute but statistically significant absolute differences were

observed between each of the three populations [$F(2,59) = 19.06$, $p < .00001$]. To explore possible

biases in the gene expression analysis caused by confounding environmental factors, principal

component analysis (PCA) was performed on all RNAseq assemblies including two replicates of

'Mara des Bois' fruit harvested in different seasons (Figure S2). Total transcript-accumulation

variation clusters most strongly according to familial relationship, with the 'Mara des Bois'

replicates showing similar expression patterns. A measured amount of variation due to

environmental influence can also be seen, as the two 'Mara des Bois' RNAseq replicates cluster

somewhat more closely with their co-harvested progeny.

R-gene eQTL in 61 Strawberry Fruit Transcriptomes

eQTL analysis was performed to evaluate heritable genotypic effects on R-gene transcription using

61 octoploid IStraw35 genotypes and mature fruit transcriptomes. This analysis identified 77 R-

gene-like sequences with at least one highly-significant locus explaining differential expression

(3.9% of octoploid genome-predicted R-genes). These R-genes include 53 NB-ARC containing

genes, comprised of 25 TNL's, 14 CNL's, 3 NBS-RPW proteins, and 11 NB-ARC-only proteins

(File S1). The majority of remaining eQTL genes are TIR, LRR, and RPW-only genes. As the

'Camarosa' genomic locus of each transcript is known, *cis* vs *trans* eQTL status was determined.

Of 77 significant R-gene eQTL transcripts, all 77 R-genes are regulated via a *cis*-genetic locus

(Table 2), of which 24 R-genes are also under regulation of an additional second *trans*-eQTL

(Table 3). No solely *trans*-eQTL were discovered among this set of R-genes. The most significant

IStraw35 SNP marker name and position for each R-gene transcript is provided with the eQTL

phase, minor allele frequency, p-value (FDR-adjusted), heritability estimate, expression in parental

lines, and BLAST2GO description.

A representative eQTL R-gene (maker-Fvb5-1-snap-gene-191.37) is detailed in Figure 6. Analysis

by BLAST2GO indicates an "RPM1-like disease resistance gene" whose *Arabidopsis thaliana*

homolog confers resistance to *Pseudomonas syringae*. This gene is hereafter referred to as

"*FaRPM1.1*". An unusual double NB-ARC structure is predicted for *FaRPM1.1* (Figure 6A). An

eQTL was detected for this gene relative to chromosome 5 on the *F. vesca* genome position (Figure

6B). This eQTL is superficially analogous to the physical position of octoploid *FaRPM1.1* on

chromosome 5, homoeolog 1 (Figure 6C). The significant eQTL markers are not included in the

'Holiday' × 'Korona' octoploid genetic map, impeding a recombination-based subgenomic genetic

association (van Dijk *et al.*, 2014). However, the associated marker physical sequences match

uniquely to the 'Camarosa' chromosome 5 homoeolog 1 locus, within several kilobases of the

FaRPM1.1, confirming a *cis*-eQTL designation (Figure 6D).

Evolutionary Pressure on *F. ×ananassa* R-genes

Elevated median dN/dS ratios were observed across all 1,962 predicted *F. × ananassa* R-genes

(0.4232) compared to non R-genes (0.3526) (Figure 4). Fewer R-genes exhibited extremely low

dN/dS ratios, indicating that high degrees of R-gene conservation are less common. However, a

similar rate of hypervariable genes (dN>>dS) was observed between R-genes and non R-genes. A

complete list of dN/dS ratios for each *F. ×ananassa* R-gene is provided in Supplementary Table

1.

R-gene dN/dS values were compared against transcript accumulation across various strawberry

tissues and receptacle stages. R-genes with low transcript accumulation across all tissues were

correlated with higher dN/dS ratios (Pearson's r = -0.69, p < .0001) (Figure S5). In other words,

R-genes with poor evidence of expression also have higher ratios of non-synonymous mutation

capable of altering amino acid sequences and affecting protein function.

Subgenome Dominance in Octoploid Strawberry

Polyploidization is associated with rapid genome remodeling events to establish a new

homeostasis, including selective gene loss and methylation. While R-gene expansiveness is often

considered evolutionarily favorable, genes that are stoichiometrically or dosage sensitive are more

commonly retained in duplicate after polyploidization (Edger and Pires, 2009, Birchler and Veitia,

2012, Edger *et al.*, 2017a). The 'Camarosa' octoploid genome, in comparison with the genomes

from its diploid *F. vesca*-like and *F. iinumae*-like ancestors, has provided an ideal platform to

study the general biological phenomena of post-hybridization genome remodeling and subgenome

dominance (Edger *et al.*, 2019). To gauge R-gene post-hybridization retention specifically, a gene-

focused baseline assessment of subgenome dominance in the 'Camarosa' octoploid genome was

necessary. Putative gene ancestry was predicted based on gene-by-gene sequence comparisons to

determine the closest 'Camarosa' gene homologs in *F. vesca* (Fragaria_vesca_v2.0.a2.cds) and *F.*

iinumae (FII_r1.1cds), which is representative of the highly similar 'old world' subgenomes. This

gene-by-gene putative orthology analysis was selected over a total comparison of homeologous

chromosomes, as extensive genetic transfer from the *F. vesca*-like subgenome has strongly

converted all subgenomes to contain *F. vesca*-like genes over time (Tennessen *et al.*, 2014), and

because the *F. iinumae* FII_r1.1 genome is incompletely assembled and is not amenable to whole-

genome alignment. By this facile coding-sequence comparison method, a significant bias towards

the retention and/or expansion of *F. vesca*-like genes is observed in the 'Camarosa' genome

(Figure 5A), with an even stronger bias towards *F. vesca*-like fruit gene expression (Figure 5B)

consistent with previous analyses (Edger *et al.*, 2019). Of 108,087 *F. ×ananassa* 'Camarosa'

predicted gene models, 68,664 genes (63.5%) were most similar to an *F. vesca* gene model, with

35,377 (32.7%) most similar to an *F. iinumae* gene model, with a minority of genes not closely

matching either. A single homoeologous chromosome with significantly more *F. vesca*-like genes

(~80% *F. vesca*-like) was seen in every chromosomal group. In a majority of cases, this putative

F. vesca-derived chromosome possesses the greatest total gene content of the chromosome group.

In 61 fruit transcriptomes, 73.7% of total transcripts derived from a gene sequence most similar to

F. vesca, corresponding to a 10.2% expression increase relative to the baseline genomic retention

bias. This bias towards the expression of *F. vesca*-like sequences was seen on every subgenome

(Figure 5B, yellow highlight).

NLR-gene Subgenome Dominance

Significant gene retention bias towards R-genes that are more *F. vesca*-like is observed in

'Camarosa' gene models (Figure 6). Of the 750 predicted NLR-gene models, 71.7% more closely

resemble a *F. vesca* gene rather than an *F. iinumae* gene (Figure 6A). This is somewhat higher

than the baseline retention bias towards *F. vesca*-like genes in octoploid (63.8%) from this

analysis. In every chromosome group, the *F. vesca*-like homoeologous chromosomes (yellow
highlight) retained the greatest number of NLRs. Of expressed R-genes, 1,125 demonstrate the
highest sequence identity with an *F. vesca* gene, 444 show highest sequence identity with an *F.*
iinumae gene, and 2 (an RPW-only gene, and an LRR_8-only gene) are without significant
matches to either diploid genome. While *F. vesca*-like genes contribute the most to total mature
fruit NLR expression (70.5% of transcripts), this is proportional to *F. vesca*-like NLR genome
content (71.3%) and is similar in magnitude to general *F. vesca* expression bias (73.7%) (Figure
6B). In other words, *F. vesca*-like NLR genes are retained in the octoploid genome somewhat
above the baseline bias, but do not experience the additional expression bias that is a generic
feature of *F. vesca*-like transcripts.

RenSeq for Strawberry Resistance Genes
A panel of sequence capture probes was designed based on putative R-gene sequences discovered
in the genomes of *F. ×ananassa* 'Camarosa', *F. vesca* genotype Hawaii 4, *F. iinumae*, and de novo
fruit transcriptomes from *F. ×ananassa* cultivars 'Mara des Bois' and 'Florida Elyana'. Benchtop
RenSeq capture on genomic DNA was performed on a collection of sixteen strawberry genotypes,
including twelve *F. ×ananassa* advanced breeding selections, three *F. ×ananassa* disease-resistant
cultivars, and a diploid *F. vesca*. As a preliminary validation of capture efficiency with this novel
RenSeq probe panel, multiplexed Illumina sequencing was performed on captured R-gene
genomic libraries. An average of 2.60 million reads (2×100bp) was obtained for each of sixteen
libraries from a single lane. Reads from octoploid and diploid lines were mapped to their respective
annotated genomic references. An average R-gene resequencing depth of 26x was achieved in the
'Camarosa' RenSeq line and 30x in the *F. vesca*, with similar coverage ranges in the other diverse
octoploid accessions (Figure 8). In the 'Camarosa' RenSeq line, 68% of reads mapped to an
annotated resistance gene, while an additional 20% of reads mapped to a non-R-gene gene model.
In *F. vesca* this efficiency was lower, where 36% of reads mapped to an annotated R-gene. A
FASTA of RenSeq probes is provided for use in File S2. Example probe coverage is detailed in
Figure S6.

DISCUSSION

These results provide a characterization of the R-gene complement of cultivated octoploid strawberry and the relationship to the extant diploid relatives, *F. vesca* and *F. iinumae*. Commercial strawberry is hypothesized to contain a single *F. vesca*-like subgenome, and three highly-similar 'old world' subgenomes which are likely derived from *F. iinumae*, *F. viridis,* and *F. nipponica* (Edger *et al.*, 2019). Polyploidization is associated with massive genome remodeling events including gene loss (Edger *et al.*, 2017a, Edger *et al.*, 2018). Linkage-map comparisons in octoploid and diploid strawberry have uncovered extensive unidirectional homoeologous exchanges which have broadly converted the three *'old world' F. innumae*-like subgenomes to be more *F. vesca*-like (Tennessen *et al.*, 2014). This finding explains the difficulties of clear ancestral delineation of strawberry homoeologs (Vining *et al.*, 2017). Recent analysis of the octoploid genome reveals that biased homologous exchanges have converted other subgenomes to be more like the dominant *F. vesca*-like subgenome (Edger *et al.*, 2019). The present gene-level homology and expression analysis shows the majority of *F. vesca*-like dominance is derived from *F. vesca*-like genes residing on alternate subgenomes. For NLR genes in particular, the bias towards *F. vesca*-like genomic retention was more pronounced. Unlike general octoploid genes, expression of *F. vesca*-like and *F. iinumae*-like NLRs is proportional to their genomic representation. This finding provides potential insight into the practical drivers of subgenome conversion. Consolidation of redundant genes and maintenance of stoichiometrically sensitive genes has been hypothesized as a driver for gene retention bias. NLRs are involved in consequential and sensitive protein-level interactions, including signaling functions requiring homo- and hetero-dimerizations. Avoidance of dysfunctional NLR molecular interactions may have contributed to the observed biases in NLR retention and expression, post-polyploidization.

Multiple distinct NLR clades with identical domain architectures were detected, likely distinguishing intra-subgenome homologs from different subgenomes. These likely reflect broad ancestral sequence divergences prior to hybridization. Comparison of R-genes in octoploid and diploid strawberry reveals enrichment of different subtypes. The 'Camarosa' genome shows a large increase in complete TNL-type R-genes and a concomitant decrease in truncated TIR-only genes, relative to its diploid ancestral relatives. The *F. iinumae* genome shows a considerably larger amount of CNL-types. TNLs have been nearly eliminated from most monocot genomes in bias towards CNL-types (Nepal *et al.*, 2017). The reasons for emerging divisions in TNL/CNL

content in plant genomes remains unclear. In hybrid *F. ×ananassa,* it is possible that relatively high number of complete TNL genes is a result of higher rate of retention post-polyploidization in this category. Many of the non-classical domains discovered in 'Camarosa' R-genes have also been found and characterized in the R-genes of other species. These include an LRR/Malectin-like RLK protein, which mediates powdery mildew resistance in barley and wheat (Rajaraman *et al.*, 2016). Atypical R-gene domains physically associated with NB-ARCs have been implicated in a variety of active disease resistance functions, including signal transduction and defense gene activation, and serving as decoy endogenous sequences to bait pathogen effectors into direct interaction and detection (Khan *et al.*, 2016).

A large proportion of strawberry NLR genes from octoploid and diploid genomes are associated with RPW8 domains. It has been suggested that the RPW8 domain emerged with the earliest land plants and subsequently merged with NLR genes, however their prevalence across plant genomes varies widely (Zhong and Cheng, 2016). The RPW domain appears to have been completely lost in monocots, and is rare in many other species. R-gene genomic studies frequently neglect to assess the presence of NLR-associated RPW domains. Two NBS-RPW8 proteins conferring mildew-resistance have been described in the *Arabidopsis thaliana* genome (Xiao *et al.*, 2001) that retained their function when expressed in grape (Hu *et al.*, 2018). The *At*RPW8.2 gene was recently shown to induce the expression of defense-related genes when expressed in strawberry leaves (Cui *et al.*, 2017). This R-gene subtype has apparently expanded in strawberry, possibly due to unusually high mildew disease pressure exerted on strawberry species and intense selection for resistance. However, R-gene domain content is not reliably predictive of resistance specificity, and close R-gene paralogs are known to confer resistance to pathogens in entirely different kingdoms (Wen *et al.*, 2015). Interestingly, the strawberry RPW8 domain is frequently found in association with NB-ARC-containing genes but never with TIRs. The purpose of RPW8 gene expansion in strawberry remains an interesting open question.

Octoploid NLR transcript accumulation is low throughout the strawberry plant, but is particularly low in the mature receptacle. This is a somewhat unexpected result due to the many pathogens targeting this susceptible organ. It is possible that only certain R genes are highly upregulated in the response to pathogen attack. Another possibility is that resistance based on the hypersensitive response may be less effective at mature stages, where cell wall disruption has already initiated with ripening and the intercellular environment is conducive to pathogen growth. Transcriptional

response to *Botrytis cinerea* infection in the mature octoploid receptacle led to differential
expression of over 1,500 genes, including secondary metabolism and pathogenesis-related (PR)
genes, but only 15 NLR genes (Xiong *et al.*, 2018). In the present study, elevated NLR
transcription in the green, white, and turning stages suggest NLR-based resistance may be more
prevalent at these earlier developmental stages. The highest levels of NLR expression were seen
in the roots and leaves, indicating this mode of resistance may be more common in these tissues.
NLR expression overlaps poorly between tissues, supporting the concept that NLRs are optimized
for each tissue. It would be interesting to examine the patterns of tissue specific expression of R-
genes against different strawberry pathogens.

The genetics of differential fruit expression of R-genes in strawberry cultivars was examined via
eQTL analysis. In many cases, the identified genetic markers described presence/absence of R-
gene expression. The identified eQTLs were often due to a *cis* variant at a single detectable locus,
very close to the physical position of the gene itself. This is suggestive of a mutation in a *cis*-
regulatory element, such as the gene promoter or 5'-UTR, or a genic presence/absence structural
variation. Such presence/absence variation affects nearly 20% of genes in the *Brassica oleracea*
pangenome and is a major contributor of agronomic trait diversity (Golicz *et al.*, 2016). As these
strawberry R-gene eQTL are derived from crosses of cultivars with differing ranges of pathogen
susceptibility, these eQTL genes represent strong candidates for functional disease resistance and
potential genetic improvement. These disclosed R-gene eQTL marker sequences may be cross-
referenced with existing disease-resistance QTL to potentially identify causal R-genes. As
categories of R-genes are expressed at very low levels unless induced by pathogens (Lai and
Eulgem, 2017), the genotype × pathogen interaction may have lowered confidence values or
introduced possible type II errors in eQTL detection. However, the reproducibility of *cis*-eQTL
tends to be particularly high in related populations (Peirce *et al.*, 2006). Additional replicates and
infected/non-infected challenge conditions will likely reveal additional eQTL associations and
greatly improve the confidence of heritability estimates, and may be used to validate pathogen-
induced R-gene candidates.

F. ×*ananassa* predicted R-genes (NLRs and other R-gene types) have elevated average dN/dS
ratios compared to non R-genes, indicating greater overall tendency towards divergent selection.
R-genes with very low dN/dS ratios are likely to be conserved disease resistance genes. This active
evolutionary selection is highly indicative of function. Of particular interest are strawberry R-

genes demonstrating both low dN/dS values and low transcript levels across all tissues (Table S1).
Many functional R-genes are expressed at low levels, either constitutively or until elicited by the
proper pathogen (Lai and Eulgem, 2017). Such R-genes may be difficult to distinguish from
pseudogenes on a purely transcriptional bases. Low dN/dS values demonstrate selective pressure
to maintain these sequences, offering evidence of maintained function despite low expression. The
results of this combinatorial analysis can be used help identify novel sources of R-gene-based
resistance which may be otherwise difficult to detect. It should be noted that this analysis is
performed in the context of a single cultivar, which has undergone several centuries of artificial
selection. It is possible that wild octoploid species may reveal different and more natural patterns
of disease-resistance selection. More sequenced accessions from geographically diverse wild and
cultivated germplasm are needed. Further analysis on the octoploid pangenome will reveal more
detailed selection patterns, and more importantly, reveal recent selection sweep events which may
have occurred in certain R-gene groups.

Many R-genes were discovered clustered in the genomes of both octoploid and diploid strawberry,
highlighting the challenges of resolving individual R-genes via association mapping and positional
cloning. The difficulty of isolating functional R-genes from strawberry disease resistance QTL
was the principle motivator of this analysis. A thorough identification of R-genes in the octoploid
genome is necessary for future genomics and genetics analysis in strawberry disease-resistance
breeding programs. Additionally, this information is prerequisite for creating a RenSeq probe
panel, to facilitate targeted R-gene sequencing in breeding programs.

A novel strawberry RenSeq capture-probe library was developed based on the R-gene sequences
identified from genomic and transcriptomic resources.   This 39,501-probe panel was
experimentally validated using octoploid and diploid genomes and resulted in an average ~20× R-
gene resequencing depth per genomic library, using only multiplexed short reads. RenSeq
assembly in 'Camarosa' and the *F. vesca* genotype Hawaii 4 resulted in significant coverage of R-
genes. Interestingly, the capture efficiency (R-gene reads over total reads) was somewhat lower in
F. vesca, likely representing the saturation of capture probes in the smaller *F. vesca* genome.
Similar rates of perfect sequence matching along the entire read in 'Camarosa' and *F. vesca*
(66.24% and 69.68%, respectively) indicates that theoretical octoploid reference sequence errors
are not likely promoting RenSeq assembly error in 'Camarosa'. However, 14.4% of mapped
'Camarosa' R-gene reads have an equally valid alternative R-gene mapping locus, compared with

just 4.83% in *F. vesca*. This difference indicates that homoeologous sequence redundancy is an

appreciable issue for mapping short-reads in polyploids, even with an isogenic (but not haplotype-

specific) mapping reference. Longer sequencing read-lengths, spanning less well-conserved non-

coding sequences, will assist in de novo resolution of similar loci in octoploid strawberry.

Combining RenSeq with longer-read third generation sequencing technologies will allow for

improved de novo assembly of R-gene loci, and will greatly facilitate causal mutation detection

within disease resistance QTL in octoploid strawberry.

MATERIALS AND METHODS

Plant Populations and Genetic Materials

Three pedigree-connected and segregating strawberry populations were created from crosses 'Florida Elyana' × 'Mara de Bois', 'Florida Radiance' × 'Mara des Bois', and 'Strawberry Festival' × 'Winter Dawn' (Figure S7). These cultivars and 54 progeny were selected for RNAseq and Istraw35 SNP genotyping analysis (Verma *et al.*, 2017), and were used to identify expressed genes and R-gene eQTL. De novo assemblies of 'Mara des Bois' and 'Florida Elyana' were also used to help design RenSeq capture probes.

For RenSeq, 14 disease resistant octoploid cultivars and elite breeding lines were selected from the University of Florida breeding program, and supplemented with 'Camarosa' and with the ancestral diploid *F. vesca*. The RenSeq lines are *F. vesca* genotype Hawaii 4, 'Camarosa', Sweet Sensation® 'Florida127', 'Florida Elyana', 11.28-34, 11.77-96, 11.98-41, 12.115-10, 12.121-5, 13.26-134, 13.42-5, 13.55-195, 14.100-58, 14.100-59, 14.101-154, and 14.101-225.

Identification of R-genes in Strawberry spp.

R-genes were predicted from the strawberry octoploid 'Camarosa' draft genome "F_ana_Camarosa_6-28-17.rm" (Edger *et al.*, 2019), the diploid *F. vesca* reassembly "Fragaria_vesca_v2.0.a2" (Tennessen *et al.*, 2014), and the diploid *F. innumae* assembly "FII_r1.1" (Hirakawa *et al.*, 2014). Domain-level analysis was performed using the CLC Genomics Workbench 11 HMM implementation to search for Pfam- v29 domains on translated gene models from all genomic and transcriptomic strawberry resources. Motif search was performed on all translated gene models, using 56 R-gene-associated motifs collected from (Lukasik and Takken, 2009, Jupe *et al.*, 2012, Van Ghelder and Esmenjaud, 2016). The CLC Genomics Workbench 11 (CLC Bio, Denmark) pattern discovery tool was trained on a preliminary list of strawberry R-genes, and novel motifs were reiterated back to all protein models. The Ncoils sequence analysis algorithm (Lupas *et al.*, 1991) was used to detect coiled-coil domains, and the output was parsed into GFF3 format for protein list reannotation. BLAST2GO annotation (Conesa *et al.*, 2005) was performed to assign putative functions to all genes and confirm sequence association with disease resistance in a cross-referenced database.

Protein models containing canonical R-gene domains (eg. NB-ARC domain) were selected for inclusion as R-genes, as were gene models with more common domains (eg. LRR) with supporting

450 evidence of an R-gene-associated motif. BLAST2GO annotated disease resistance associated
451 genes not meeting the domain and motif-level criteria were analyzed for potential inclusion,
452 leading to the inclusion of many LRR-containing RLK putative R-genes.

NB-ARC Phylogenetic Analysis

NB-ARC domains were extracted from *F. iinumae*, *F. vesca*, and *F. ×ananassa* 'Camarosa'. The
CIPRES Science Gateway (Miller *et al.*, 2010) was utilized for full-length protein sequence
alignment using MUSCLE v3.7 (Edgar, 2004) and Maximum likelihood analysis using RAxML
v8.2.10 (Stamatakis, 2014). Tree construction was performed using the PROTGAMMA rate
distribution model with 100 bootstrap replicates, and rooted with human APAF-1. This process
was replicated five times using different random number seeds. Trees were visualized in CLC
Genomic Workbench 11 with a 50% threshold bootstrap value. Word clouds were generated per
clade based on the relative domain content of the full proteins.

Subgenome Dominance in Octoploid Strawberry R-genes

The closest homolog for each *F. ×ananassa* 'Camarosa' gene in either Fragaria_vesca_v2.0.a2.cds
or FII_r1.1cds was determined via BLAST analysis (e-value threshold < 0.1, word size=25,
match=1, mismatch=l, existence=0, extension=2). *F. vesca*-like and *F. iinumae*-like gene counts
and TPMs were independently calculated for each octoploid chromosome. This process was
performed first on all genes in the 'Camarosa' genome to establish the baseline gene retention and
expression bias, and then repeated using only predicted NLR genes.

***d*N/*d*S Analysis**

*d*N   and   *d*S   values   were   computed   using   a   set   of   custom   scripts
(https://github.com/Aeyocca/ka_ks_pipe/). Orthologous genes between the *F. ×ananassa and F.*
vesca v4 (Edger *et al.*, 2017b) genomes were identified using the compara module in JCVI utilities
library (Tang et al., 2015). Filtering of the JCVI utilities output was performed using a custom perl
script to identify the best syntenic ortholog and best blast hit below e-value 1e-4. Alignment of
each orthologous gene pair was performed using MUSCLE v3.8.31 (Edgar, 2004), followed by
PAL2NAL (v14) (Suyama *et al.*, 2006) to convert the peptide alignment to a nucleotide alignment.
Finally, *d*N and *d*S values were computed between those gene pairs using codeml from PAML
Version 4.9h (Yang, 2007) with parameters specified in the control file found in the GitHub
repository listed above.

Tissue-specific Transcriptome Analysis

Raw short read RNAseq libraries from various 'Camarosa' tissue (Sánchez-Sevilla *et al.*, 2017)

with the study reference PRJEB12420 were download from the European Nucleotide Archive

(https://www.ebi.ac.uk/ena). The complete 54 library RNAseq experiment consisted of six

independent green receptacle libraries, six white receptacle libraries, six turning receptacle

libraries, six red receptacle libraries, three root libraries, three leaf libraries, and six achene libraries

each for all corresponding fruit stages. Raw RNAseq reads were assembled to the 'Camarosa'

reference using the same pipeline as previously described for fruit transcriptome population

analysis. Expression values from biologically-replicated libraries were averaged. Clustvis

(Metsalu and Vilo, 2015) was used for tissue-based RNAseq clustering and heatmap visualization

using correlation distance and average linkage with scaling applied using default parameters.

Fruit Transcriptome Analysis

61 fruit transcriptomes were sequenced via Illumina paired-end RNAseq (Avg. 65million reads,

2x100bp), and consisted of parents and progeny from crosses of 'Florida Elyana' × 'Mara de Bois',

'Florida Radiance' × 'Mara des Bois', and 'Strawberry Festival' × 'Winter Dawn'. Reads were

trimmed and mapped to the *F. ×ananassa* octoploid 'Camarosa' annotated genome using CLC

Genomic Workbench 11 (mismatch cost of 2, insertion cost of 3, deletion cost of 3, length fraction

of 0.8, similarity fraction of 0.8, 1 maximum hit per read). Reads that mapped equally well to more

than one locus were discarded from the analysis. RNAseq counts were calculated in Transcripts

Per Million (TPM). Three-dimensional principle component analysis (PCA) was performed on all

RNAseq assemblies, including two replicates of 'Mara des Bois' fruit harvested three years apart

and sequenced independently (Figure S2). Transcript abundances were normalized via the Box-

Cox transformation algorithm performed in R (R. Development Core Team, 2014) prior to eQTL

analysis. The BLAST2GO pipeline was used to annotate the full 'Camarosa' predicted gene

complement.

Genotyping and Genetic Association of Octoploid Fruit R-genes

505 The Affymetrix IStraw35 Axiom® SNP array (Verma *et al.*, 2017) was used to genotype 60

506 individuals, including six parental lines from three independent biparental RNAseq populations

507 (Figure S7). Sequence variants belonging to the Poly High Resolution (PHR) and No Minor

508 Homozygote (NMH) marker classes were included for association mapping. Mono High

509 Resolution (MHR), Off-Target Variant (OTV), Call Rate Below Threshold (CRBT), and Other

marker quality classes, were discarded and not used for mapping. Individual marker calls inconsistent with Mendelian inheritance from parental lines were removed. The *F. vesca* physical map was used to orient marker positions as current octoploid maps do not include a majority of the available IStraw35 markers. A genome-wide analysis study (GWAS) was performed using GAPIT v2 (Tang *et al.*, 2016) performed in R. R-gene eQTL were evaluated for significance based on the presence of multiple co-locating markers of p-value < 0.05 after false discovery rate correction for multiple comparisons. *Cis* vs *trans* eQTL determinations were made by corroborating known 'Camarosa' physical gene position with the eQTL position the *F. vesca* map. In the example case of *FaRPM1.1*, subgenomic localization was confirmed via BLAST of the associated markers to the correct 'Camarosa' homoeologous chromosome.

RenSeq Probe Design and Validation

A panel of 39,501 of 120mer-length capture probes were designed based on the set of discovered strawberry R-genes from *F. ×ananassa* 'Camarosa', *F. vesca* genotype Hawaii 4, *F. iinumae* genomes, and de novo fruit transcriptomes from *F. ×ananassa* 'Mara des Bois' and 'Florida Elyana'. A proprietary algorithm was used to select for capture probes of ideal hybridization thermodynamics and screened for potential off-target capture in the intergenic regions of 'Camarosa' and *F. vesca* (Rapid Genomics LLC, Gainesville FL). Probes were designed to not span exon-exon junction, to facilitate cross-utility for both genomic and cDNA libraries (Figure S6). A minimum baseline of 1x probe coverage was provided across the length of every predicted R-gene coding sequence, and additional probes were designed against conserved R-gene domains in order to promote capture of unknown and divergent R-genes across diverse octoploid accessions. RenSeq capture was performed on genomic libraries from fifteen octoploid disease-resistant cultivars and advanced breeding selections, and *F. vesca*, based on conditions set by (Jupe *et al.*, 2014), with optimizations provided by Rapid Genomics LLC. Captured libraries were sequenced via 16x multiplexed Illumina Hiseq (2 × 100bp) and mapped to their respective annotated genomic references using CLC Genomic Workbench 11 (CLCBio, Aarhus, Denmark) (Similarity fraction = 0.9, Length fraction = 0.9, Match score = 1, Mismatch cost = 2, Insertion cost = 3, Deletion cost = 3).

Data Availability

Supplementary figures, tables, files, and raw data are available at FigShare. Custom scripts used for performing *d*N/*d*S analysis are available at Github: https://github.com/Aeyocca/ka_ks_pipe/.

Raw short read RNAseq data from fruit transcriptomes are available from the NCBI Short Read Archive under project SRP039356 (http://www.ncbi.nlm.nih.gov/sra/?term=SRP039356). Raw short read RNAseq data from the 'Camarosa' gene expression atlas (Sánchez-Sevilla et al., 2017) are available at the European Nucleotide Archive (https://www.ebi.ac.uk/ena) with the study reference PRJEB12420. Results derived from these data are compiled in Table S1. File S1 contains BED files for annotating the octoploid genome (Edger *et al.*, 2019) with R-genes and R-gene domains. Renseq probe sequences are provided in File S2. Istraw35 markers, map positions, and sample genotypes used in eQTL analysis are available in File S3.

ACKNOWLEDGEMENTS

We acknowledge Aristotole Koukoulidis for ncoils prediction and annotation, Matthew Robinson for assistance with data normalization, Rapid Genomics LLC for technical assistance in probe generation and sequence capture, Anne Schwartz, Max Hogshead, Kiran Sharma and Nadia Mourad for assistance in data compilation, Ben Harrison and Max Hogshead for assistance in gDNA isolation of eQTL lines, and Dr. Alan Chambers and Dr. Jeremy Pillet for RNA isolation and RNAseq line selection. This research is supported by grants to SJK, VMW, and SL from the United Stated Department of Agriculture (http://dx.doi.org/10.13039/100000199) National Institute of Food and Agriculture (NIFA) Specialty Crops Research Initiative (#2017-51181-26833) and to SJK from the California Strawberry Commission (http://dx.doi.org/10.13039/100006760).

CONFLICT OF INTEREST

The authors declare no conflicts of interest.

LITERATURE CITED

Amil-Ruiz, F., Departamento de Bioquímica y Biología Molecular e Instituto Andaluz de Biotecnología, C.d.E.I.A.C.C.d.R.E.S.O.U.d.C.C.S., Blanco-Portales, R., Muñoz-Blanco, J. and Caballero, J.L. (2018) The Strawberry Plant Defense Mechanism: A Molecular Review. *Plant and Cell Physiology*, **52**, 1873-190310.1093/pcp/pcr136.

Anciro, A., Mangandi, J., Verma, S., Peres, N., Whitaker, V.M. and Lee, S. (2018) FaRCg1: a quantitative trait locus conferring resistance to Colletotrichum crown rot caused by Colletotrichum gloeosporioides in octoploid strawberry. *Theoretical and Applied Genetics*10.1007/s00122-018-3145-z.

Andolfo, G., Jupe, F., Witek, K., Etherington, G.J., Ercolano, M.R. and Jones, J.D.G. (2014) Defining the full tomato NB-LRR resistance gene repertoire using genomic and cDNA RenSeq. *BMC Plant Biology*, **14**,

12010.1186/1471-2229-14-120.

Arya, P., Kumar, G., Acharya, V. and Singh, A.K. (2014) Genome-Wide Identification and Expression Analysis
of NBS-Encoding Genes in Malus x domestica and Expansion of NBS Genes Family in Rosaceae. *PLOS*
ONE, **9**, e10798710.1371/journal.pone.0107987.

Baumgartner, I.O., Patocchi, A., Frey, J.E., Peil, A. and Kellerhals, M. (2015) Breeding Elite Lines of Apple
Carrying Pyramided Homozygous Resistance Genes Against Apple Scab and Resistance Against Powdery
Mildew and Fire Blight. *Plant Molecular Biology Reporter*, **33**, 1573-158310.1007/s11105-015-0858-x.

Birchler, J.A. and Veitia, R.A. (2012) Gene balance hypothesis: Connecting issues of dosage sensitivity across
biological disciplines. *Proceedings of the National Academy of Sciences*, **109**,
1474610.1073/pnas.1207726109.

Conesa, A., Götz, S., García-Gómez, J.M., Terol, J., Talón, M. and Robles, M. (2005) Blast2GO: a universal tool
for annotation, visualization and analysis in functional genomics research. *Bioinformatics*, **21**, 3674-
367610.1093/bioinformatics/bti610.

Cordova, L.G., Amiri, A. and Peres, N.A. (2017) Effectiveness of fungicide treatments following the Strawberry
Advisory System for control of Botrytis fruit rot in Florida. *Crop Protection*, **100**, 163-
167https://doi.org/10.1016/j.cropro.2017.07.002.

Cui, M.-Y., Wei, W., Gao, K., Xie, Y.-G., Guo, Y. and Feng, J.-Y. (2017) A rapid and efficient Agrobacterium-
mediated transient gene expression system for strawberry leaves and the study of disease resistance proteins.
Plant Cell, Tissue and Organ Culture (PCTOC), **131**, 233-24610.1007/s11240-017-1279-3.

Djian-Caporalino, C., Palloix, A., Fazari, A., Marteu, N., Barbary, A., Abad, P., Sage-Palloix, A.-M., Mateille,
T., Risso, S., Lanza, R., Taussig, C. and Castagnone-Sereno, P. (2014) Pyramiding, alternating or mixing:
comparative performances of deployment strategies of nematode resistance genes to promote plant resistance
efficiency and durability. *BMC Plant Biology*, **14**, 5310.1186/1471-2229-14-53.

Edgar, R.C. (2004) MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic acids*
research, **32**, 1792-1797

Edger, P.P., McKain, M.R., Bird, K.A. and VanBuren, R. (2018) Subgenome assignment in allopolyploids:
challenges and future directions. *Current Opinion in Plant Biology*, **42**, 76-
80https://doi.org/10.1016/j.pbi.2018.03.006.

Edger, P.P. and Pires, J.C. (2009) Gene and genome duplications: the impact of dosage-sensitivity on the fate of
nuclear genes. *Chromosome Research*, **17**, 69910.1007/s10577-009-9055-9.

Edger, P.P., Poorten, T.J., VanBuren, R., Hardigan, M.A., Colle, M., McKain, M.R., Smith, R.D., Teresi, S.J.,
Nelson, A.D.L., Wai, C.M., Alger, E.I., Bird, K.A., Yocca, A.E., Pumplin, N., Ou, S., Ben-Zvi, G., Brodt,
A., Baruch, K., Swale, T., Shiue, L., Acharya, C.B., Cole, G.S., Mower, J.P., Childs, K.L., Jiang, N.,
Lyons, E., Freeling, M., Puzey, J.R. and Knapp, S.J. (2019) Origin and evolution of the octoploid
strawberry genome. *Nature Genetics*, **51**, 541-54710.1038/s41588-019-0356-4.

Edger, P.P., Smith, R., McKain, M.R., Cooley, A.M., Vallejo-Marin, M., Yuan, Y., Bewick, A.J., Ji, L., Platts,
A.E., Bowman, M.J., Childs, K.L., Washburn, J.D., Schmitz, R.J., Smith, G.D., Pires, J.C. and Puzey,
J.R. (2017a) Subgenome Dominance in an Interspecific Hybrid, Synthetic Allopolyploid, and a 140-Year-
Old Naturally Established Neo-Allopolyploid Monkeyflower. *The Plant Cell*, **29**, 2150

Edger, P.P., VanBuren, R., Colle, M., Poorten, T.J., Wai, C.M., Niederhuth, C.E., Alger, E.I., Ou, S., Acharya,
C.B., Wang, J., Callow, P., McKain, M.R., Shi, J., Collier, C., Xiong, Z., Mower, J.P., Slovin, J.P.,
Hytönen, T., Jiang, N., Childs, K.L. and Knapp, S.J. (2017b) Single-molecule sequencing and optical
mapping yields an improved genome of woodland strawberry (Fragaria vesca) with chromosome-scale
contiguity. *GigaScience*, **7**10.1093/gigascience/gix124.

Farzaneh, M., Kiani, H., Sharifi, R., Reisi, M. and Hadian, J. (2015) Chemical composition and antifungal effects
of three species of Satureja (S. hortensis, S. spicigera, and S. khuzistanica) essential oils on the main
pathogens of strawberry fruit. *Postharvest Biology and Technology*, **109**, 145-
151https://doi.org/10.1016/j.postharvbio.2015.06.014.

Funk, A., Galewski, P. and McGrath, J.M. (2018) Nucleotide-binding resistance gene signatures in sugar beet,
insights from a new reference genome. *The Plant Journal*, **95**, 659-67110.1111/tpj.13977.

Golicz, A.A., Bayer, P.E., Barker, G.C., Edger, P.P., Kim, H., Martinez, P.A., Chan, C.K.K., Severn-Ellis, A.,
McCombie, W.R., Parkin, I.A.P., Paterson, A.H., Pires, J.C., Sharpe, A.G., Tang, H., Teakle, G.R.,
Town, C.D., Batley, J. and Edwards, D. (2016) The pangenome of an agronomically important crop plant
Brassica oleracea. *Nature Communications*, **7**, 1339010.1038/ncomms13390

https://www.nature.com/articles/ncomms13390#supplementary-information.

Hammond-Kosack, K.E. and Jones, J.D.G. (1997) Plant Disease Resistance Genes. *Annual Review of Plant*
Physiology and Plant Molecular Biology, **48**, 575-60710.1146/annurev.arplant.48.1.575.

Herrington, M.E., Hardner, C., Wegener, M., Woolcock, L.L. and Dieters, M.J. (2011) Rain Damage to
Strawberries Grown in Southeast Queensland: Evaluation and Genetic Control. *HortScience*, **46**, 832-837

Hirakawa, H., Shirasawa, K., Kosugi, S., Tashiro, K., Nakayama, S., Yamada, M., Kohara, M., Watanabe, A.,
Kishida, Y., Fujishiro, T., Tsuruoka, H., Minami, C., Sasamoto, S., Kato, M., Nanri, K., Komaki, A.,
Yanagi, T., Guoxin, Q., Maeda, F., Ishikawa, M., Kuhara, S., Sato, S., Tabata, S. and Isobe, S.N. (2014)
Dissection of the Octoploid Strawberry Genome by Deep Sequencing of the Genomes of Fragaria Species.
DNA Research, **21**, 169-18110.1093/dnares/dst049.

Hu, Y., Li, Y., Hou, F., Wan, D., Cheng, Y., Han, Y., Gao, Y., Liu, J., Guo, Y., Xiao, S., Wang, Y. and Wen, Y.-
Q. (2018) Ectopic expression of Arabidopsis broad-spectrum resistance gene RPW8.2 improves the
resistance to powdery mildew in grapevine (Vitis vinifera). *Plant Science*, **267**, 20-
31https://doi.org/10.1016/j.plantsci.2017.11.005.

Jia, Y., Yuan, Y., Zhang, Y., Yang, S. and Zhang, X. (2015) Extreme expansion of NBS-encoding genes in
Rosaceae. *BMC Genetics*, **16**, 4810.1186/s12863-015-0208-x.

Jupe, F., Chen, X., Verweij, W., Witek, K., Jones, J.D.G. and Hein, I. (2014) Genomic DNA Library Preparation
for Resistance Gene Enrichment and Sequencing (RenSeq) in Plants. In *Plant-Pathogen Interactions:*
Methods and Protocols (Birch, P., Jones, J.T. and Bos, J.I.B. eds). Totowa, NJ: Humana Press, pp. 291-303.

Jupe, F., Pritchard, L., Etherington, G.J., MacKenzie, K., Cock, P.J.A., Wright, F., Sharma, S.K., Bolser, D.,
Bryan, G.J., Jones, J.D.G. and Hein, I. (2012) Identification and localisation of the NB-LRR gene family
within the potato genome. *BMC Genomics*, **13**, 7510.1186/1471-2164-13-75.

Khan, M., Subramaniam, R. and Desveaux, D. (2016) Of guards, decoys, baits and traps: pathogen perception in
plants by type III effector sensors - ScienceDirect. *Current Opinion in Microbiology*, **29**, 49-
5510.1016/j.mib.2015.10.006.

Lai, Y. and Eulgem, T. (2017) Transcript-level expression control of plant NLR genes. *Molecular Plant Pathology*,
19, 1267-128110.1111/mpp.12607.

Lukasik, E. and Takken, F.L.W. (2009) STANDing strong, resistance proteins instigators of plant defence. *Current*
opinion in plant biology, **12**, 427-436

Lupas, A., Van Dyke, M. and Stock, J. (1991) Predicting Coiled Coils from Protein Sequences. *Science*, **252**, 1162-
1164

Mangandi, J., Verma, S., Osorio, L., Peres, N.A., van de Weg, E. and Whitaker, V.M. (2017) Pedigree-Based
Analysis in a Multiparental Population of Octoploid Strawberry Reveals QTL Alleles Conferring Resistance
to Phytophthora cactorum. *G3: Genes|Genomes|Genetics*, **7**, 1707

Metsalu, T. and Vilo, J. (2015) ClustVis: a web tool for visualizing clustering of multivariate data using Principal
Component Analysis and heatmap. *Nucleic Acids Research*, **43**, W566-W57010.1093/nar/gkv468.

Miller, M.A., Pfeiffer, W. and Schwartz, T. (2010) Creating the CIPRES Science Gateway for inference of large
phylogenetic trees. In *2010 Gateway Computing Environments Workshop (GCE)*, pp. 1-8.

Nepal, M.P.A., E. J., Neupane, S. and Benson, B.V. (2017) Comparative Genomics of Non-TNL Disease Resistance
Genes from Six Plant Species. In *Genes (Basel)*.

Peirce, J.L., Li, H., Wang, J., Manly, K.F., Hitzemann, R.J., Belknap, J.K., Rosen, G.D., Goodwin, S., Sutter,
T.R., Williams, R.W. and Lu, L. (2006) How replicable are mRNA expression QTL? *Mammalian Genome*,
17, 643-65610.1007/s00335-005-0187-8.

Pincot, D.D.A., Poorten, T.J., Hardigan, M.A., Harshman, J.M., Acharya, C.B., Cole, G.S., Gordon, T.R.,
Stueven, M., Edger, P.P. and Knapp, S.J. (2018) Genome-Wide Association Mapping Uncovers
Fw1, a Dominant Gene Conferring Resistance to Fusarium Wilt in Strawberry. *G3:*
Genes|Genomes|Genetics, **8**, 181710.1534/g3.118.200129.

R. Development Core Team (2014) R: A language and environment for statistical computing.

Rajaraman, J., Douchkov, D., Hensel, G., Stefanato, F.L., Gordon, A., Ereful, N., Caldararu, O.F., Petrescu,
A.J., Kumlehn, J., Boyd, L.A. and Schweizer, P. (2016) An LRR/Malectin Receptor-Like Kinase Mediates
Resistance to Non-adapted and Adapted Powdery Mildew Fungi in Barley and Wheat. *Front Plant Sci*,
710.3389/fpls.2016.01836.

Roach, J.A., Verma, S., Peres, N.A., Jamieson, A.R., van de Weg, W.E., Bink, M.C.A.M., Bassil, N.V., Lee, S.
and Whitaker, V.M. (2016) FaRXf1: a locus conferring resistance to angular leaf spot caused by

Xanthomonas fragariae in octoploid strawberry. *Theoretical and Applied Genetics*, **129**, 1191-
120110.1007/s00122-016-2695-1.

Salinas, N., Verma, S., Peres, N. and Whitaker, V.M. (2018) FaRCa1: a major subgenome-specific locus conferring
resistance to Colletotrichum acutatum in strawberry. *Theoretical and Applied Genetics*10.1007/s00122-018-
3263-7.

Stamatakis, A. (2014) RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies.
Bioinformatics, **30**, 1312-131310.1093/bioinformatics/btu033.

Suyama, M., Torrents, D. and Bork, P. (2006) PAL2NAL: robust conversion of protein sequence alignments into
the corresponding codon alignments. *Nucleic acids research*, **34**, W609-W612

Sánchez-Sevilla, J.F., Vallarino, J.G., Osorio, S., Bombarely, A., Posé, D., Merchante, C., Botella, M.A., Amaya,
I. and Valpuesta, V. (2017) Gene expression atlas of fruit ripening and transcriptome assembly from RNA-
seq data in octoploid strawberry (Fragaria × ananassa). *Scientific Reports*, **7**, 1373710.1038/s41598-017-
14239-6.

Tang, Y., Liu, X., Wang, J., Li, M., Wang, Q., Tian, F., Su, Z., Pan, Y., Liu, D., Lipka, A.E., Buckler, E.S. and
Zhang, Z. (2016) GAPIT Version 2: An Enhanced Integrated Tool for Genomic Association and Prediction.
The Plant Genome, **9**10.3835/plantgenome2015.11.0120.

Tennessen, J.A., Govindarajulu, R., Ashman, T.-L. and Liston, A. (2014) Evolutionary Origins and Dynamics of
Octoploid Strawberry Subgenomes Revealed by Dense Targeted Capture Linkage Maps. *Genome Biology*
and Evolution, **6**, 3295-331310.1093/gbe/evu261.

van Dijk, T., Pagliarani, G., Pikunova, A., Noordijk, Y., Yilmaz-Temel, H., Meulenbroek, B., Visser, R.G.F.
and van de Weg, E. (2014) Genomic rearrangements and signatures of breeding in the allo-octoploid
strawberry as revealed through an allele dose based SSR linkage map. *BMC Plant Biology*, **14**,
5510.1186/1471-2229-14-55.

Van Ghelder, C. and Esmenjaud, D. (2016) TNL genes in peach: insights into the post-LRR domain. *BMC*
Genomics, **17**, 31710.1186/s12864-016-2635-0.

Verma, S., Bassil, N.V., van de Weg, E., Harrison, R.J., Monfort, A., Hidalgo, J.M., Amaya, I., Denoyes, B.,
Mahoney, L., Davis, T.M., Fan, Z., Knapp, S. and Whitaker, V.M. (2017) Development and evaluation
of the Axiom® IStraw35 384HT array for the allo-octoploid cultivated strawberry Fragaria ×ananassa:
International Society for Horticultural Science (ISHS), Leuven, Belgium, pp. 75-82.

Verma, S., Osorio, L.F., Lee, S., Bassil, N.V. and Whitaker, V.M. (2018) Genome-Assisted Breeding in the
Octoploid Strawberry. In *The Genomes of Rosaceous Berries and Their Wild Relatives* (Hytönen, T., Graham,
J. and Harrison, R. eds). Cham: Springer International Publishing, pp. 161-184.

Vining, K.J., Salinas, N., Tennessen, J.A., Zurn, J.D., Sargent, D.J., Hancock, J. and Bassil, N.V. (2017)
Genotyping-by-sequencing enables linkage mapping in three octoploid cultivated strawberry families. *PeerJ*,
5, e373110.7717/peerj.3731.

Wen, Z., Yao, L., Wan, R., Li, Z., Liu, C. and Wang, X. (2015) Ectopic Expression in Arabidopsis thaliana of an
NB-ARC Encoding Putative Disease Resistance Gene from Wild Chinese Vitis pseudoreticulata Enhances
Resistance to Phytopathogenic Fungi and Bacteria. *Frontiers in Plant Science*, **6**, 1087

Witek, K., Jupe, F., Witek, A.I., Baker, D., Clark, M.D. and Jones, J.D.G. (2016) Accelerated cloning of a potato
late blight–resistance gene using RenSeq and SMRT sequencing. *Nature Biotechnology*, **34**,
65610.1038/nbt.3540
https://www.nature.com/articles/nbt.3540#supplementary-information.

Xiao, S., Ellwood, S., Calis, O., Patrick, E., Li, T., Coleman, M. and Turner, J.G. (2001) Broad-Spectrum Mildew
Resistance in Arabidopsis thaliana Mediated by RPW8. 10.1126/science.291.5501.118.

Xiong, J.-S., Zhu, H.-Y., Bai, Y.-B., Liu, H. and Cheng, Z.-M. (2018) RNA sequencing-based transcriptome
analysis of mature strawberry fruit infected by necrotrophic fungal pathogen Botrytis cinerea. *Physiological*
and Molecular Plant Pathology, **104**, 77-85https://doi.org/10.1016/j.pmpp.2018.08.005.

Yang, Z. (2007) PAML 4: phylogenetic analysis by maximum likelihood. *Molecular biology and evolution*, **24**, 1586-
1591

Zhong, Y. and Cheng, Z.-M.M. (2016) A unique RPW8-encoding class of genes that originated in early land plants
and evolved through domain fission, fusion, and duplication. *Scientific Reports*, **6**,
32923doi:10.1038/srep32923.

Zhong, Y., Zhang, X. and Cheng, Z.-M. (2018) Lineage-specific duplications of NBS-LRR genes occurring before
the divergence of six Fragaria species. *BMC Genomics*, **19**, 12810.1186/s12864-018-4521-4.

Table 1 NLR-gene subtype distribution across three strawberry species

	cv. Camarosa	F. vesca	F. iinumae
NB	19.8%	17.7%	11.6%
NB-LRR	2.3%	1.9%	5.2%
NB-only type	**22.1%**	**19.6%**	**16.8%**
CC-NB	16.7%	12.3%	32.6%
CC-NB-LRR	3.0%	2.5%	2.1%
CNL-type	**19.7%**	**14.7%**	**34.6%**
RPW	9.3%	10.9%	5.9%
RPW-NB	4.6%	8.4%	3.4%
RPW-type	**13.9%**	**19.3%**	**9.3%**
TIR	13.7%	37.1%	22.2%
TIR-NB	20.0%	4.9%	13.2%
TIR-NB-LRR	10.6%	4.4%	3.9%
TNL-type	**44.3%**	**46.3%**	**39.3%**

Relative content of NLR-gene subtypes and truncated subtypes are shown.

738
739

Table 2 Cis Expression-QTL for Strawberry Fruit

R-gene Name	eQTL phase	IStraw35 AX-	MAF	p-value (FDR)	h² estimate	Mara	Elya-na	Radi-ance	Festi-val	Winter dawn	Description
augustus_masked-fvb1-2-processed-gene-27.2	cis	89817565	0.43	0.0418	30.4%	36.6	113.3	32.2	21.3	34.3	uncharacterized protein LOC101293711
augustus_masked-fvb1-4-processed-gene-7.11	cis	166502627	0.16	0.0150	43.4%	2.3	0.6	4.0	3.8	1.6	PLT5_ARATH Sugar-proton symporter PLT5
augustus_masked-fvb2-4-processed-gene-105.5	cis	166503861	0.29	0.0114	80.9%	0.0	0.0	2.8	1.8	1.3	TMVRN_NICGU TMV resistance N
augustus_masked-fvb3-1-processed-gene-107.3	cis	166511589	0.50	0.0294	41.8%	0.0	0.5	0.7	0.8	0.3	DRL30_ARATH Probable disease resistance At5g04720
augustus_masked-fvb3-3-processed-gene-283.7	cis	89786873	0.38	0.0009	30.4%	0.8	0.4	0.0	2.6	0.4	DRL1_ARATH Probable disease resistance At1g12280
augustus_masked-fvb3-3-processed-gene-38.8	cis	166513199	0.23	0.0032	81.7%	6.0	3.0	5.5	0.8	4.9	Y4294_ARATH LRR receptor-like serine threonine- kinase
augustus_masked-fvb3-4-processed-gene-19.4	cis	166504873	0.28	0.0014	100%	0.1	4.8	5.6	8.0	0.1	DR100_ARATH DNA damage-repair toleration DRT100
augustus_masked-fvb4-1-processed-gene-166.2	cis	166505902	0.43	0.0083	91.5%	0.6	0.4	2.5	1.4	1.8	MKKA_DICDI Mitogen-activated kinase kinase
augustus_masked-fvb4-2-processed-gene-107.10	cis	166527457	0.18	0.0202	100%	0.0	0.0	1.2	1.0	0.1	RGA1_SOLBU disease resistance RGA1 RGA3-blb
augustus_masked-fvb4-2-processed-gene-258.8	cis	166505336	0.35	0.0015	25.1%	4.6	0.0	1.4	1.3	0.1	LRX2_ARATH Leucine-rich repeat extensin 2
augustus_masked-fvb5-1-processed-gene-238.7	cis	123365994	0.09	0.0012	74.1%	0.8	0.8	1.4	2.8	0.5	HSL1_ARATH Receptor kinase HSL1 HAESA-LIKE1
augustus_masked-fvb5-1-processed-gene-71.8	cis	166524323	0.48	0.0080	35.5%	1.7	0.0	10.3	4.4	3.0	TIR_ARATH Toll interleukin-1 receptor
augustus_masked-fvb5-3-processed-gene-135.4	cis	166523635	0.26	0.0124	63.7%	0.5	0.2	0.1	0.0	0.0	MKKA_DICDI Mitogen-activated kinase kinase
augustus_masked-fvb5-4-processed-gene-18.1	cis	166518037	0.35	0.0110	26.6%	0.8	1.4	0.7	2.5	0.8	RGA1_SOLBU disease resistance RGA1 RGA3-blb
augustus_masked-fvb5-4-processed-gene-241.6	cis	123525092	0.44	0.0007	64.9%	1.4	0.1	2.5	2.1	1.8	LRX2_ARATH Leucine-rich repeat extensin 2 2 LRR
augustus_masked-fvb6-1-processed-gene-345.10	cis	166524541	0.21	0.0050	31.7%	1.8	0.7	4.1	4.1	2.2	HSL1_ARATH Receptor kinase HSL1 HAESA-LIKE1
augustus_masked-fvb7-1-processed-gene-57.2	cis	166509572	0.12	0.0003	73.3%	0.5	0.2	0.6	0.2	13.2	TIR_ARATH Toll interleukin-1 receptor
augustus_masked-fvb7-2-processed-gene-302.13	cis	166517211	0.29	0.0083	69.2%	2.3	0.2	2.1	0.1	0.0	MKKA_DICDI Mitogen-activated kinase kinase
augustus_masked-fvb7-2-processed-gene-53.6	cis	166509530	0.49	0.0118	35.7%	1.2	0.2	0.5	0.4	2.2	RGA1_SOLBU disease resistance RGA1 RGA3-blb
augustus_masked-fvb7-2-processed-gene-54.1	cis	166509530	0.49	0.0121	100%	9.9	0.1	0.4	0.3	11.5	LRX2_ARATH Leucine-rich repeat extensin 2 2 LRR
maker-fvb1-4-augustus-gene-30.48	cis	123365909	0.33	0.0175	63.8%	2.5	0.4	3.1	4.2	2.5	HSL1_ARATH Receptor kinase HSL1 HAESA-LIKE1
maker-fvb2-1-augustus-gene-182.42	cis	89877559	0.46	0.0225	80.9%	2.1	3.8	2.2	3.2	0.8	MKKA_DICDI Mitogen-activated kinase
maker-fvb2-1-snap-gene-111.27	cis	166503168	0.29	0.0002	100%	0.5	0.0	0.0	2.0	1.2	RGA1_SOLBU disease resistance RGA1 RGA3-blb
maker-fvb7-1-snap-gene-223.45	cis	123540423	0.45	0.0077	70.3%	2.4	0.1	0.1	0.0	1.7	LRX2_ARATH Leucine-rich repeat extensin 2 2 LRR
maker-fvb7-4-snap-gene-48.49	cis	166508582	0.34	0.0357	75.1%	0.3	0.4	0.1	0.0	0.2	HSL1_ARATH Receptor kinase HSL1 HAESA-LIKE1
maker-fvb7-4-snap-gene-59.59	cis	166518351	0.50	0.0064	4.7%	0.9	0.4	1.8	0.1	1.6	TIR_ARATH Toll interleukin-1 receptor
maker-fvb7-4-snap-gene-59.63	cis	166518351	0.50	0.0092	44.8%	0.1	0.0	0.6	0.0	0.4	MKKA_DICDI Mitogen-activated kinase
maker-fvb7-4-snap-gene-69.51	cis	123359450	0.17	0.0002	82.6%	2.3	1.8	2.1	3.6	2.2	RGA1_SOLBU disease resistance RGA1 RGA3-blb
snap_masked-fvb1-2-processed-gene-79.33	cis	123359751	0.08	0.0068	98.3%	0.0	0.0	0.0	0.0	2.0	LRX2_ARATH Leucine-rich repeat extensin 2 2 LRR
snap_masked-fvb2-1-processed-gene-107.14	cis	89780995	0.22	0.0011	75.1%	2.1	2.6	8.5	4.0	3.4	HSL1_ARATH Receptor kinase HSL1 HAESA-LIKE1
snap_masked-fvb3-2-processed-gene-11.25	cis	166509770	0.37	0.0084	11.4%	0.2	0.2	1.1	1.5	0.7	TIR_ARATH Toll interleukin-1 receptor
snap_masked-fvb3-3-processed-gene-288.15	cis	123361033	0.45	0.0095	98.9%	0.9	0.5	2.0	1.4	1.3	MKKA_DICDI Mitogen-activated kinase kinase kinase A
snap_masked-fvb6-1-processed-gene-37.31	cis	166519417	0.34	0.0411	24.2%	13.3	19.2	0.9	0.5	0.0	RGA1_SOLBU disease resistance RGA1
snap_masked-fvb7-2-processed-gene-254.35	cis	123357141	0.33	0.0011	36.5%	0.0	0.0	0.0	0.0	0.0	LRX2_ARATH Leucine-rich repeat extensin 2 2
maker-fvb7-1-snap-gene-273.51	cis	166508667	0.43	0.0319	35.7%	0.0	0.0	0.0	0.4	0.0	Y3475_ARATH LRR receptor-like serine threonine- kinase
maker-fvb3-4-augustus-gene-265.40	cis	166513103	0.13	0.0007	54.7%	4.4	5.3	2.6	18.8	5.2	GLO5_ARATH Peroxisomal(S)-2-hydroxy-acid oxidase GLO5
maker-fvb5-2-augustus-gene-191.37	cis	166523649	0.13	0.0000	36.5%	0.0	0.0	0.0	0.0	0.1	RPM1_ARATH Disease resistance RPM1
maker-fvb5-2-augustus-gene-59.20	cis	123364094	0.46	0.0036	46.0%	0.0	0.3	4.0	12.0	0.8	probable LRR receptor-like serine threonine- kinase At5g48740
maker-fvb5-2-augustus-gene-61.13	cis	123364094	0.46	0.0014	71.7%	0.0	0.3	2.2	1.6	0.8	P2B10_ARATH F-box PP2-B10 PHLOEM PROTEIN 2-LIKE
maker-fvb5-2-augustus-gene-63.17	cis	123364094	0.46	0.0072	16.5%	0.0	0.0	3.5	3.7	1.1	P2B11_ARATH F-box PP2-B11 PHLOEM PROTEIN 2-LIKE
maker-fvb5-2-snap-gene-4.75	cis	166506813	0.32	0.0000	29.9%	2.1	9.3	37.2	43.0	1.4	DRL28_ARATH Probable disease resistance At4g27220
maker-fvb5-2-snap-gene-61.17	cis	123364094	0.46	0.0291	78.4%	0.0	0.2	1.3	1.4	0.3	P2B11_ARATH F-box PP2-B11 PHLOEM PROTEIN 2-LIKE
maker-fvb5-3-augustus-gene-135.25	cis	89832439	0.37	0.0040	11.5%	0.0	0.0	1.6	0.3	1.3	RPM1_ARATH Disease resistance RPM1
maker-fvb4-3-snap-gene-155.68	cis	123524810	0.21	0.0016	100%	0.1	0.2	0.6	0.1	0.5	TMVRN_NICGU TMV resistance N
maker-fvb5-3-snap-gene-221.67	cis	89893608	0.11	0.0045	94.4%	0.5	0.2	0.1	1.6	0.1	TMVRN_NICGU TMV resistance N
maker-fvb5-3-snap-gene-254.50	cis	166523796	0.21	0.0155	51.8%	0.1	0.1	0.4	0.5	0.1	TMVRN_NICGU TMV resistance N
maker-fvb5-4-snap-gene-125.42	cis	166506186	0.18	0.0078	93.2%	0.6	0.0	2.0	2.3	1.4	RGA3_SOLBU RGA3 Blight resistance B149
maker-fvb6-1-augustus-gene-153.32	cis	166507404	0.17	0.0000	76.3%	0.0	0.0	23.5	6.8	17.5	RGA3_SOLBU RGA3 Blight resistance B149
maker-fvb5-2-augustus-gene-61.14	cis	123358673	0.47	0.0167	44.9%	0.0	0.1	0.7	1.2	0.3	P2B10_ARATH F-box PP2-B10 PHLOEM PROTEIN 2-LIKE
augustus_masked-fvb7-1-processed-gene-284.2	cis	123359573	0.4	0.0139	95.1%	0.3	0	0.1	0.3	0.2	EMS1_ARATH Leucine-rich repeat receptor kinase EMS1
snap_masked-fvb6-2-processed-gene-263.31	cis	89781514	0.24	0.0084	54.9%	0	0	4.3	4.3	2.2	TMVRN_NICGU TMV resistance N
maker-fvb7-2-snap-gene-161.50	cis	123364494	0.39	0.0052	38.9%	2.4	1	4	2	2.6	MAP1A_ARATH Methionine aminopeptidase 1A MAP 1A 1A
maker-fvb6-1-augustus-gene-160.45	cis	166515747	0.21	0.0340	38.4%	12.6	8.8	11.9	4.2	10.3	DGK5_ARATH Diacylglycerol kinase 5

Genetic association results for 61 transcriptomes are shown, detailing cis genetic factors controlling differentially expressed R-genes. The most significant marker name, minor allele frequency, FDR-adjusted p-value, narrow sense heritability, transcript accumulation in cultivars, and BLAST2GO description are shown.

Table 3 Trans and Cis Expression-QTL for Strawberry Fruit

R-gene Name	eQTL phase	IStraw35 AX-	MAF	p-value (FDR)	h² estimate	Mara	Elya-na	Radi-ance	Festi-val	Winter dawn	Description
maker-fvb4-2-snap-gene-5.61	cis	166505436	0.38	0.0214	34.3%	1.5	0.9	1.6	3.0	2.3	PSKR1_ARATH LRR receptor kinase 1 serine-threonine
	trans	123361503	0.38	0.0214							
maker-fvb5-1-augustus-gene-139.47	cis	123524951	0.28	0.0482	77.5%	0.3	0.4	5.8	1.5	1.0	PIRL5_ORYSJ Plant intracellular Ras-group-related LRR 5
	trans	123361742	0.14	0.0482							
maker-fvb5-1-augustus-gene-263.34	cis	123357041	0.18	0.0415	14.6%	8.6	4.0	3.3	3.7	3.2	R13L1_ARATH disease resistance RPP13 1
	trans	166506561	0.18	0.0415							
maker-fvb5-2-snap-gene-213.46	cis	89817904	0.10	0.0006	79.8%	1.7	1.6	3.3	0.0	1.4	DRL21_ARATH disease resistance At3g14460
	trans	123539826	0.10	0.0006							
maker-fvb5-2-snap-gene-46.64	cis	166506808	0.48	0.0192	67.1%	3.4	0.0	1.3	1.0	1.4	GSO1_ARATH LRR receptor-like serine threonine- kinase
	trans	166522785	0.48	0.0192							
maker-fvb5-3-snap-gene-244.48	cis	123367068	0.48	0.0046	39.5%	0.4	0.5	4.7	5.6	2.8	Y4265_ARATH Probable LRR receptor-like serine threonine-
	trans	166524268	0.48	0.0046							
maker-fvb5-4-snap-gene-110.38	cis	166523511	0.21	0.0001	100%	1.0	1.3	0.4	1.6	0.3	PIRL5_ORYSJ Plant intracellular Ras-group-related LRR 5
	trans	166524147	0.29	0.0014							
maker-fvb6-1-augustus-gene-27.58	cis	123363787	0.38	0.0028	91.2%	0.3	0.0	0.2	0.5	0.0	DRL42_ARATH Probable disease resistance At5g66900
	trans	123358884	0.38	0.0028							
maker-fvb6-3-snap-gene-412.71	cis	123614270	0.27	0.0457	74.4%	1.8	1.1	1.4	2.1	0.9	TIR_ARATH Toll interleukin-1 receptor
	trans	166525307	0.27	0.0457							
augustus_masked-fvb6-2-processed-gene-268.10	cis	166525890	0.48	0.0021	91.2%	0.8	0.8	0.3	1.7	1.2	HSL1_ARATH Receptor kinase HSL1 HAESA-LIKE1
	trans	123362183	0.48	0.0021							
augustus_masked-fvb6-3-processed-gene-176.9	cis	166515622	0.33	0.0004	65.6%	0.4	0.0	7.9	20.0	0.1	ADT3_ARATH ADP,ATP carrier mitochondrial ADP ATP
	trans	123525691	0.33	0.0005							
augustus_masked-fvb7-2-processed-gene-63.0	cis	166508452	0.30	0.0047	27.7%	0.5	0.1	1.3	0.1	0.0	TMVRN_NICGU TMV resistance N
	trans	89823698	0.28	0.0047							
augustus_masked-fvb7-3-processed-gene-243.6	cis	166517344	0.38	0.0204	66.5%	47.5	14.1	0.1	8.1	0.7	DR100_ARATH DNA damage-repair toleration DRT100
	trans	89894427	0.38	0.0204							
maker-fvb1-2-snap-gene-63.40	cis	166510935	0.17	0.0015	86.2%	0.5	0.1	1.3	0.1	0.0	TMVRN_NICGU TMV resistance N
	trans	166525497	0.16	0.0015							
maker-fvb1-2-snap-gene-96.40	cis	166517617	0.18	0.0020	67.6%	0.0	0.0	1.0	0.0	0.4	DRL42_ARATH Probable disease resistance
	trans	166525528	0.18	0.0020							
maker-fvb1-4-snap-gene-66.72	cis	123363545	0.28	0.0280	68.4%	3.9	0.2	12.0	12.0	7.4	U496I_ARATH UPF0496 At2g18630
	trans	166516240	0.18	0.0093							
maker-fvb1-4-snap-gene-76.45	cis	123357162	0.18	0.0001	100%	0.7	0.2	0.0	0.0	0.0	DRL43_ARATH Probable disease resistance
	trans	166516240	0.18	0.0001							
maker-fvb7-2-augustus-gene-136.47	cis	166526312	0.43	0.0116	79.9%	0.3	0.0	0.4	0.4	0.2	TMVRN_NICGU TMV resistance N
	trans	123359434	0.46	0.0371							
maker-fvb7-2-augustus-gene-147.47	cis	123359385	0.45	0.0003	45.2%	1.1	0.2	5.5	3.1	2.1	TMVRN_NICGU TMV resistance N
	trans	123365359	0.42	0.0011							
maker-fvb7-2-augustus-gene-163.44	cis	123364494	0.39	0.0011	52.3%	0.7	0.3	2.0	0.9	0.8	RGA3_SOLBU disease resistance RGA3 Blight resistance
	trans	123365359	0.42	0.0055							
maker-fvb7-2-augustus-gene-65.21	cis	166512110	0.21	0.0031	69.2%	5.3	0.3	0.9	1.2	0.5	RP8L2_ARATH Probable disease resistance RPP8 2
	trans	166509598	0.21	0.0031							
maker-fvb7-2-snap-gene-161.50	cis	123364494	0.39	0.0052	38.9%	2.4	1.0	4.0	2.0	2.6	MAP1A_ARATH Methionine aminopeptidase 1A MAP 1A 1A
	trans	123365359	0.42	0.0154							
snap_masked-fvb3-4-processed-gene-8.19	cis	166521734	0.16	0.0023	69.6%	2.2	4.3	6.2	5.2	4.0	TMVRN_NICGU TMV resistance N
	trans	89826525	0.17	0.0033							
snap_masked-fvb6-1-processed-gene-352.19	cis	123366334	0.27	0.0306	76.8%	0.1	0.0	0.8	1.1	0.8	TMVRN_NICGU TMV resistance N
	trans	123357007	0.42	0.0465							

Genetic association results for 61 transcriptomes are shown, detailing cis genetic factors controlling differentially expressed R-genes. The most significant marker name, minor allele frequency, FDR-adjusted p-value, narrow sense heritability, transcript accumulation in cultivars, and BLAST2GO description are shown.

Figure 1 Canonical and non-Canonical Domains in Strawberry R-Genes. Classic TNL/CNL-type R-gene domains (TIR, NB-ARC, LRR, etc.) comprise the majority of domain classes in predicted R-genes, however a number of atypical domains are observed in high frequency. Domains below a count of five are not shown.

Figure 2 Phylogenetic Relationship of NB-ARC domains in *F. vesca*, *F. iinumae*, & *F. × ananassa* 'Camarosa'. A. Full-length NB-ARC domains from strawberry *spp*. organize into clades based on NLR-gene subtype (CC, TIR, NB-ARC, LRR, and RPW-containing combinations). Maximum likelihood bootstrap values (100 replicates) above a threshold of 50% are shown with the NB-ARC domain from human *Apaf1* as the outgroup. Word sizes correspond to relative domain content within each clade.

A.

B.

	Green Receptacle	White receptacle	Turning receptacle	Red receptacle	Root	Leaf	Green achene	White achene	Turning achene	Red achene
Average TPM	1.73	1.55	1.05	0.88	3.04	1.88	1.35	0.31	0.27	0.37
Expressed NLRs	213 (28.1%)	198 (26.2%)	108 (14.3%)	116 (15.3%)	321 (42.2%)	275 (36.4%)	194 (25.7%)	48 (6.3%)	40 (5.3%)	56 (7.4%)

C.

Figure 3 RNAseq Expression of Octoploid NLR genes. A. Tissue-based heatmap of scaled
NLR transcript accumulation. B. Averaged NLR transcript abundance (TPM) and total number
of putative expressed NLRs (TPM >1) are reported. The percent of all 756 NLR genes putatively
expressed in each tissue are shown in parentheses. C. Generalized R-gene fruit expression across
three segregating populations (n=61).

Figure 4 Evolutionary Pressures on *F. × ananassa* R-genes. The median *d*N/*d*S ratio for R-
genes (0.4142) is higher than for non R-genes (0.3547). Density curves for *F. ×ananassa* R-genes
(blue) and non R-genes (red) are calculated based on comparison to the closest ancestral diploid
homolog from *F. vesca*.

A.   **Gene Content**

	*F. iinumae-*like	*F. vesca-*like	niether	% *F. vesca-*like	Total genes
Fvb1-1	1,188	1,736	108	57.3%	3,032
Fvb1-2	1,443	1,930	112	55.4%	3,485
Fvb1-3	1,243	1,798	106	57.1%	3,147
Fvb1-4	504	2,593	132	80.3%	3,229
Fvb2-1	1,293	2,124	109	60.2%	3,526
Fvb2-2	588	2,836	163	79.1%	3,587
Fvb2-3	1,217	1,895	110	58.8%	3,222
Fvb2-4	1,631	2,017	118	53.6%	3,766
Fvb3-1	1,487	2,270	114	58.6%	3,871
Fvb3-2	1,830	2,102	139	51.6%	4,071
Fvb3-3	1,463	2,156	132	57.5%	3,751
Fvb3-4	697	3,206	138	79.3%	4,041
Fvb4-1	961	1,386	82	57.1%	2,429
Fvb4-2	1,245	1,743	148	55.6%	3,136
Fvb4-3	634	3,534	191	81.1%	4,359
Fvb4-4	1,319	1,537	130	51.5%	2,986
Fvb5-1	728	3,671	198	79.9%	4,597
Fvb5-2	1,199	2,018	118	60.5%	3,335
Fvb5-3	1,492	2,066	139	55.9%	3,697
Fvb5-4	1,232	1,902	137	58.1%	3,271
Fvb6-1	927	4,820	233	80.6%	5,980
Fvb6-2	1,782	2,919	203	59.5%	4,904
Fvb6-3	2,555	3,091	188	53.0%	5,834
Fvb6-4	1,810	2,670	145	57.7%	4,625
Fvb7-1	1,483	2,730	161	62.4%	4,374
Fvb7-2	1,046	3,853	231	75.1%	5,130
Fvb7-3	1,414	1,990	118	56.5%	3,522
Fvb7-4	966	2,071	143	65.1%	3,180
Total Genes	35,377	68,664	4,046	63.5%	108,087
Percent	32.7%	63.5%	3.7%		

B.   **Fruit Expression**

	F. iinumae-like TPM (12,762 expressed genes)	*F. vesca*-like TPM (32,663 expressed genes)	niether TPM (559 expressed genes)	Total TPM
Fvb1-1	8,360	24,301	277	32,939
Fvb1-2	9,064	15,374	87	24,525
Fvb1-3	7,666	15,364	243	23,272
Fvb1-4	2,296	33,504	133	35,933
Fvb2-1	8,291	24,535	138	32,963
Fvb2-2	1,864	33,012	2,456	37,333
Fvb2-3	7,893	25,382	210	33,485
Fvb2-4	10,471	26,313	527	37,312
Fvb3-1	16,794	19,921	388	37,103
Fvb3-2	23,304	19,340	158	42,802
Fvb3-3	17,609	21,526	231	39,365
Fvb3-4	3,948	30,368	166	34,482
Fvb4-1	6,457	37,194	31	43,682
Fvb4-2	7,317	21,620	322	29,259
Fvb4-3	4,118	37,246	231	41,595
Fvb4-4	11,480	20,677	214	32,371
Fvb5-1	3,275	33,971	284	37,531
Fvb5-2	7,435	16,610	347	24,392
Fvb5-3	11,851	17,926	220	29,997
Fvb5-4	5,311	15,460	49	20,820
Fvb6-1	3,522	65,588	348	69,458
Fvb6-2	9,648	28,090	552	38,290
Fvb6-3	26,622	32,002	555	59,178
Fvb6-4	13,270	33,673	195	47,138
Fvb7-1	7,579	20,190	612	28,381
Fvb7-2	1,548	26,160	285	27,993
Fvb7-3	11,301	20,874	74	32,248
Fvb7-4	4,835	20,881	437	26,152
Total Genes	253,128	737,102	9,770	1,000,000
Percent	25.31%	73.71%	0.98%	

Figure 5 General Retention and Expression Bias in Octoploid Strawberry. 'Camarosa' gene models are categorized as either more *F. vesca*-like, more *F. iinumae*-like, or neither. Red-green color scale indicates low-to-high gene content, respectively. Yellow highlight indicates the most *F. vesca*-like homoeolog. **A.** Gene content per homeologous chromosome, by putative ancestral gene similarity. **B.** Relative transcript accumulation of all genes in the fruit, by putative ancestral similarity.

A. NLR Gene Content

	F. iinumae-like	F. vesca-like	niether	% F. vesca-like	Total NLRs
Fvb1-1	6	7	0	53.8%	13
Fvb1-2	4	17	0	81.0%	21
Fvb1-3	6	13	0	68.4%	19
Fvb1-4	4	21	0	84.0%	25
Fvb2-1	8	9	0	52.9%	17
Fvb2-2	10	13	0	56.5%	23
Fvb2-3	5	4	0	44.4%	9
Fvb2-4	11	10	0	47.6%	21
Fvb3-1	6	20	0	76.9%	26
Fvb3-2	7	16	0	69.6%	23
Fvb3-3	8	19	0	70.4%	27
Fvb3-4	14	32	0	69.6%	46
Fvb4-1	2	3	0	60.0%	5
Fvb4-2	6	10	0	62.5%	16
Fvb4-3	3	15	0	83.3%	18
Fvb4-4	5	4	1	40.0%	10
Fvb5-1	9	27	0	75.0%	36
Fvb5-2	4	16	0	80.0%	20
Fvb5-3	11	19	0	63.3%	30
Fvb5-4	12	12	0	50.0%	24
Fvb6-1	3	50	0	94.3%	53
Fvb6-2	3	27	0	90.0%	30
Fvb6-3	8	26	1	74.3%	35
Fvb6-4	4	16	0	80.0%	20
Fvb7-1	15	31	0	67.4%	46
Fvb7-2	18	48	0	72.7%	66
Fvb7-3	11	30	0	73.2%	41
Fvb7-4	7	23	0	76.7%	30
Total NLRs	210	538	2	71.7%	750
Percent	28.0%	71.7%	0.3%		

B. NLR Gene Fruit Expression

	F. iinumae-like TPM (12,762 expressed genes)	F. vesca-like TPM (32,663 expressed genes)	niether TPM (559 expressed genes)	Total TPM
Fvb1-1	1	11	0	12
Fvb1-2	6	19	0	25
Fvb1-3	3	16	0	19
Fvb1-4	2	39	0	41
Fvb2-1	62	51	0	113
Fvb2-2	72	92	0	164
Fvb2-3	5	2	0	7
Fvb2-4	54	30	0	85
Fvb3-1	2	34	0	36
Fvb3-2	3	11	0	14
Fvb3-3	2	14	0	16
Fvb3-4	17	33	0	50
Fvb4-1	0	3	0	3
Fvb4-2	4	13	0	17
Fvb4-3	0	40	0	40
Fvb4-4	13	3	0	16
Fvb5-1	1	20	0	21
Fvb5-2	5	25	0	29
Fvb5-3	9	8	0	17
Fvb5-4	5	15	0	20
Fvb6-1	4	68	0	72
Fvb6-2	1	13	0	13
Fvb6-3	5	25	0	30
Fvb6-4	1	15	0	16
Fvb7-1	15	19	0	34
Fvb7-2	3	28	0	31
Fvb7-3	1	43	0	45
Fvb7-4	3	21	0	24
Total TPM	297	712	0	1,009
Percent	29.5%	70.5%	0.0%	

Figure 6 NLR-gene Retention and Expression Bias in Octoploid Strawberry. 'Camarosa' gene models are categorized as either more *F. vesca*-like, more *F. iinumae*-like, or neither. Red-green color scale indicates low-to-high gene content, respectively. Yellow highlight indicates the most *F. vesca*-like homoeolog. **A**. NLR-gene content per homeologous chromosome, by putative ancestral gene similarity. **B.** Relative transcript accumulation of NLR-genes in the fruit, by putative ancestral similarity.

Figure 7 Example *cis*-eQTL of a Fruit-Expressed Strawberry R-gene. A. Domain analysis of the 'Camarosa' putative resistance gene *FaRPM1.1* indicates two NB-ARC domains. Grey lines delineate exon-exon borders in the mature predicted transcript. **B.** Octoploid fruit expression of *FaRPM1.1* associates with a single locus on chromosome 5. **C.** The *Fvb5-1* subgenomic location of *FaRPM1.1* in the octoploid 'Camarosa' genome is indicated (purple vertical line). **D**. Three equally highly-significant GWAS markers (p-value 8.09E-06, post-FDR adjustment) show close subgenome co-localization with *FaRPM1.1*.

803
804
Figure 8 RenSeq Increases Sequencing Depth for R-gene Loci in Multiplexed Octoploid
Genomes. Sixteen disease-resistant strawberry genomic libraries (fifteen octoploid accessions and
diploid F. vesca) were enriched for R-genes and sequenced via Illumina Hiseq yielding an average
of 2.60 million reads per genomic library. Roughly half of all sequencing reads mapped to a
previously-identified R-gene locus, representing a substantial sequence enrichment relative to R-
gene genomic representation.
811