# BOLD and EEG Signal Variability at Rest
# Differently Relate to Aging in the Human Brain

D. Kumral[1,2], F. Şansal[3,1], E. Cesnaite[1], K. Mahjoory[1,4], E. Al[2,1], M. Gaebler[1,2], V. V. Nikulin[1,5,6], A. Villringer[1,2,7,8]

[1]Department of Neurology, Max Planck Institute for Human Cognitive and Brain Sciences, Leipzig, Germany

[2]MindBrainBody Institute at the Berlin School of Mind and Brain, Humboldt-Universität zu Berlin, Berlin, Germany

[3]International Graduate Program Medical Neurosciences, Charité-Universitätsmedizin, Berlin, Germany

[4]Institute for Biomagnetism and Biosignalanalysis, University of Muenster, Muenster, Germany

[5]Neurophysics Group, Department of Neurology, Campus Benjamin Franklin, Charité Universitätsmedizin Berlin, Berlin, Germany

[6]Centre for Cognition and Decision Making, National Research University Higher School of Economics, Moscow, Russian Federation

[7]Center for Stroke Research Berlin, Charité – Universitätsmedizin Berlin, Berlin, Germany

[8]Department of Cognitive Neurology, University Hospital Leipzig

**Corresponding author:** Deniz Kumral, Department of Neurology, Max Planck Institute for Human Cognitive and Brain Sciences, Stephanstrasse 1A, 04103, Leipzig, Germany.
Email: dkumral@cbs.mpg.de

**Abbreviations:** BOLD – Blood Oxygenation Level Dependent; CBF – cerebral blood flow; CBV– cerebral blood volume; CCA – canonical correlation analysis; $CMRO_2$ – cerebral metabolic rate of oxygen; CVR – cerebrovascular reactivity; DMN – Default Mode Network; EEG – electroencephalography; EC – eyes closed; EO – eyes open; framewise displacement – FD; FDR - false discovery rate; FEM – finite element method; fMRI – functional Magnetic Resonance Imaging; fNIRS – functional Near-Infrared Spectroscopy; FWHM – full-width half-maximum; ICBM – International Consortium for Brain Mapping; MEG – magnetoencephalography; MNI – Montreal Neurological Institute; rho – Spearman's rank correlation coefficient; PET –Positron-emission tomography; ROI – regions of interests; rs – resting state; SD – standard deviation; SVD – Singular Value Decomposition; TIV–Total Intracranial Volume

**Abstract**

Variability of neural activity is regarded as a crucial feature of healthy brain function, and several neuroimaging approaches have been employed to assess it noninvasively. Studies on the variability of both evoked brain response and spontaneous brain signals have shown remarkable changes with aging but it is unclear if the different measures of brain signal variability – identified with either hemodynamic or electrophysiological methods – reflect the same underlying physiology. In this study, we aimed to explore age differences of spontaneous brain signal variability with two different imaging modalities (EEG, fMRI) in healthy younger (25±3 years, N=135) and older (67±4 years, N=54) adults. Consistent with the previous studies, we found lower blood oxygenation level dependent (BOLD) variability in the older subjects as well as less signal variability in the amplitude of low-frequency oscillations (1–12 Hz), measured in source space. These age-related reductions were mostly observed in the areas that overlap with the default mode network. Moreover, age-related increases of variability in the amplitude of beta-band frequency EEG oscillations (15–25 Hz) were seen predominantly in temporal brain regions. There were significant sex differences in EEG signal variability in various brain regions while no significant sex differences were observed in BOLD signal variability. Bivariate and multivariate correlation analyses revealed no significant associations between EEG- and fMRI-based variability measures. In summary, we show that both BOLD and EEG signal variability reflect aging-related processes but are likely to be dominated by different physiological origins, which relate differentially to age and sex.

**Keywords:** brain signal variability, resting state, BOLD, fMRI, EEG, aging, sex, default mode network

# 1. Introduction

Functional neuroimaging methods such as fMRI, PET, fNIRS, EEG, or MEG have allowed the non-invasive assessment of functional changes in the aging human brain (Cabeza, 2001; Cabeza et al., 2018). Most previous functional neuroimaging studies on aging have employed a task-based design (Grady, 2012) and in their data analysis the central tendency has typically been assumed to be the most representative value in a distribution (e.g., mean) (Speelman and McGann, 2013) or the "signal" within distributional "noise". In recent years, also the variability of brain activation in task-dependent and task-independent measurements (as spontaneous variations of background activity) has been shown to provide relevant information about the brain's functional state (Garrett et al., 2013b; Grady and Garrett, 2018; Nomi et al., 2017). These studies primarily measured the blood oxygen level dependent (BOLD) signal using fMRI. For example, it has been demonstrated that the variance of the task-evoked BOLD response was differentially related to aging as well as cognitive performance (Armbruster-Genc et al., 2016; Garrett et al., 2013a). Similarly, spontaneous signal variability in resting state fMRI (rsfMRI) has been found to decrease with age (Grady and Garrett, 2018; Nomi et al., 2017), in individuals with stroke (Kielar et al., 2016), and 22q11.2 deletion syndrome (Zöller et al., 2017). An increase of fMRI variability has been shown to occur in inflammation induced state-anxiety (Labrenz et al., 2018) and to parallel symptom severity in Attention Deficit Hyperactivity Disorder (Nomi et al., 2018). From these studies, it was concluded that changes in BOLD signal variability might serve as an index for alterations in neural processing and cognitive flexibility (Grady and Garrett, 2014).

The conclusions of aforementioned studies imply that BOLD signal variability is mainly determined by *neuronal* variability. To a large extent, this is based on the premise that BOLD is related to neuronal activity: The evoked BOLD signal in task-based fMRI reflects the decrease of the deoxyhemoglobin concentration to changes in local brain activity, which is determined by vascular (blood velocity and volume: "neurovascular coupling") and metabolic (oxygen consumption: "neurometabolic coupling") factors (Logothetis and Wandell, 2004; Villringer and Dirnagl, 1995). The BOLD signal is therefore only an indirect measure of neural activity (Logothetis, 2008). For the variability of task-evoked BOLD signal and for spontaneous variations of the BOLD signal, in principle, the same considerations apply regarding their relationship to underlying neural processes (Murayama et al., 2010). However, since in rsfMRI there is no explicit external trigger for evoked brain activity to which time-locked averaging could be applied, the time course of rsfMRI signals is potentially more susceptible to contributions of "physiological noise", such as cardiac and respiratory signals

(Birn et al., 2008; Chang et al., 2009), but also spontaneous fluctuations of vascular tone, which is found even in isolated arterial vessels (Failla et al., 1999; Hudetz et al., 1998; Wang et al., 2006). In the same vein, the variability of task-evoked fMRI is not necessarily reflecting only the variability of evoked neuronal activity, as it may also – at least partly – reflect the variability of the spontaneous background signal on which a constant evoked response is superimposed.

In aging, non-neuronal signal fluctuations may also introduce spurious common variance across the rsfMRI time series (Caballero-Gaudes and Reynolds, 2017), thus confounding estimates of "neural" brain signal variability. Previous evidence suggests that the relationship between neuronal activity and the vascular response is attenuated with age – and so is, as a consequence, the BOLD signal (for review see D'Esposito et al., 2003). For instance, aging has been associated with altered cerebrovascular ultrastructure, reduced elasticity of vessels, and atherosclerosis (Farkas and Luiten, 2001) but also with a decrease in resting cerebral blood flow (CBF), cerebral metabolic rate of oxygen ($CMRO_2$), and cerebrovascular reactivity (CVR) (Liu et al., 2013). Taken together, age-related changes in BOLD signal or BOLD signal variability are related to a mixture of alterations in non-neural spontaneous fluctuations of vascular signals, neural activity, neurovascular coupling, and/or neurometabolic coupling (D'Esposito et al., 2003; Geerligs et al., 2017; Tsvetanov et al., 2015).

While BOLD fMRI signal and specifically variance measures based on fMRI are only partially and indirectly related to neural activity (Liu, 2013; Logothetis, 2008), electrophysiological methods such as EEG can provide a more direct assessment of neural activity with a higher temporal but poorer spatial resolution (Cohen, 2017). EEG measures neuronal currents resulting from the synchronization of dendritic postsynaptic potentials across the neural population; the cerebral EEG rhythms thereby reflect the underlying brain neural network activity (Steriade, 2006). Resting state (rs)EEG is characterized by spontaneous oscillations ("brain rhythms") at different frequencies. Previously, the mean amplitude of low-frequency bands (e.g., delta and/or theta, 1-7 Hz) has been shown to correlate negatively with age (Vlahou et al., 2015), while higher-frequency bands (e.g., beta, 15-25 Hz) show the reverse pattern (Rossiter et al., 2014). However, less is known about the within-subject variability of EEG measures and their association with aging. Several studies have addressed the variability in the spectral amplitudes of different frequency bands using variance (Hawkes and Prescott, 1973; Oken and Chiappa, 1988), coefficient of variation (Burgess and Gruzelier, 1993; Maltez et al., 2004), and complexity (Fernández et al., 2012;

Sleimen-Malkoun et al., 2015). For instance, reductions of the complexity in rsEEG signal

have been found not only in healthy aging (Yang et al., 2013; Zappasodi et al., 2015) but also

in age-related pathologies such as mild cognitive impairment (McBride et al., 2014) and

Alzheimer's disease (Smits et al., 2016). Accordingly, it has been suggested that irregular

(e.g., variable) systems indicate a normal and healthy state (more integrated information)

while highly regular systems often mark dysfunction or disease (Lipsitz and Goldberger,

1992; Vaillancourt and Newell, 2002).

The different methodological approaches, fMRI based "vascular" approaches on the

one hand and electrophysiological methods such as EEG and MEG, on the other hand,

indicate alterations of brain signal variability with aging. However, it remains unclear whether

these different measures of brain variability at rest reflect the same underlying physiological

changes. Evidently, there are some correlations between the two signal sources (for a review

see, Jorge et al., 2014; Ritter and Villringer, 2006). For instance, in task-based EEG-fMRI

simultaneous recordings, a relationship between BOLD responses and amplitude of evoked

potentials has been demonstrated (e.g., Ritter et al., 2009; Seaquist et al., 2007), while in

resting state EEG-fMRI studies, a negative association between spontaneous modulations of

alpha rhythm and BOLD signal has also been established (e.g., Chang et al., 2013; Goldman

et al., 2002; Gonçalves et al., 2006; Moosmann et al., 2003). Further, differential correlation

patterns have been noted for the various rhythms of different frequencies in EEG/MEG and

the fMRI signal, such that low-frequency oscillations show a negative (Deligianni et al., 2014;

Mantini et al., 2007; Meyer et al., 2013), while higher frequencies oscillations demonstrate a

positive correlation with the BOLD signal (Niessing et al., 2005; Scheeringa et al., 2011).

Regarding the known age-related changes in BOLD and EEG signal variability,

respectively, the question arises whether these alterations are dominated by joint signal

sources of fMRI and EEG or by – potentially different – signal contributions that relate to

each of these two methods. Given the – potentially large – non-neuronal signal contribution,

this issue is particularly relevant for rsfMRI studies. Here, we addressed this question by

analyzing rsfMRI and EEG measures of variability in healthy younger and older subjects. To

our knowledge, the only study that compared variability in a "vascular" imaging method

(rsfMRI) and an electrophysiological method (rsMEG at the sensor space) concluded that the

effects of aging on BOLD signal variability were mainly driven by vascular factors (e.g.,

heart rate variability) and not well-explained by the changes in neural variability (Tsvetanov

et al., 2015). The main aims of the present study were to explore i) age differences of brain

signal variability measures, as well as to investigate ii) how neural variability derived from

rsEEG related to the analogous parameters of BOLD signal variability derived from rsfMRI.

We used rsfMRI and rsEEG from the "Leipzig Study for Mind-Body-Emotion Interactions"

(Babayan et al., 2019). As an explanatory analysis, we further investigated sex-related

differences of brain signal variability measures. To measure brain signal variability, we

calculated the standard deviation (SD) of both the BOLD signal and of the amplitude

envelope of the filtered rsEEG time series for a number of standard frequency bands at the

source space. We hypothesized that brain signal variability would generally decrease with

aging. In addition, based on the premise that BOLD fMRI signal variability reflects *neural*

variability as measured by rsEEG, we expected that the corresponding changes in both signal

modalities would demonstrate moderate to strong similarity in their spatial distribution. Given

the confounding effects of vascular factors during aging on the fMRI signal (D'Esposito et al.,

2003; Liu, 2013; Thompson, 2018), we further expected to find the relationship between

BOLD and EEG signal variability to be stronger in younger than older adults.

# 2. Method

## 2.1. Participants

The data of the "Leipzig Study for Mind-Body-Emotion Interactions" (LEMON;

Babayan et al., 2019) comprised 227 subjects in two age groups (younger: 20-35, older: 59-

77). Only participants who did not report any neurological disorders, head injury, alcohol or

other substance abuse, hypertension, pregnancy, claustrophobia, chemotherapy and malignant

diseases, current and/or previous psychiatric disease or any medication affecting the

cardiovascular and/or central nervous system in a telephone pre-screening were invited to the

laboratory. The study protocol conformed to the Declaration of Helsinki and was approved by

the ethics committee at the medical faculty of the University of Leipzig (reference number

154/13-ff).

RsEEG recordings were available for 216 subjects who completed the full study

protocol. We excluded data from subjects that had missing event information (N=1), different

sampling rate (N=3), mismatching header files or insufficient data quality (N=9). Based on

the rsfMRI quality assessment, we further excluded data from subjects with faulty

preprocessing (N=7), ghost artefacts (N=2), incomplete data (N=1), or excessive head motion

(N=3) (criterion: mean framewise displacement (FD) $\leq 0.5$ mm; Power et al., 2012)

(Supplementary Figure 1). The final sample included 135 younger ($M = 25.10 \pm 3.70$ years,

42 females) and 54 older subjects ($M = 67.15 \pm 4.52$ years, 27 females).

## 2.1. fMRI Acquisition

Brain imaging was performed on a 3T Siemens Magnetom Verio MR scanner

(Siemens Medical Systems, Erlangen, Germany) with a standard 32-channel head coil. The

participants were instructed to keep their eyes open and not fall asleep while looking at a low-

contrast (light grey on dark grey background) fixation cross.

The structural image was recorded using an MP2RAGE sequence (Marques et al., 2010) with

the following parameters: TI 1 = 700 ms, TI 2 = 2500 ms, TR = 5000 ms, TE = 2.92 ms, FA 1

= 4°, FA 2 = 5°, band width = 240 Hz/pixel, FOV = 256 × 240 × 176 mm$^3$, voxel size = 1 x 1

x 1 mm$^3$. The functional images were acquired using a T2*-weighted multiband EPI sequence

with the following parameters: TR = 1400 ms, TE = 30 ms, FA= 69°, FOV = 202 mm,

imaging matrix=88 × 88, 64 slices with voxel size = 2.3 x 2.3 x 2.3 mm$^3$, slice thickness = 2.3

mm, echo spacing = 0.67 ms, bandwidth=1776 Hz/Px, partial fourier 7/8, no pre-scan

normalization, multiband acceleration factor = 4, 657 volumes, duration = 15 min 30 s. A

gradient echo field map with the sample geometry was used for distortion correction (TR =

680 ms, TE 1 = 5.19 ms, TE 2 = 7.65 ms).

## 2.2. fMRI Preprocessing

Preprocessing was implemented in Nipype (Gorgolewski et al., 2011), incorporating tools from FreeSurfer (Fischl, 2012), FSL (Jenkinson et al., 2012), AFNI (Cox, 1996), ANTs (Avants et al., 2011), CBS Tools (Bazin et al., 2014), and Nitime (Rokem et al., 2009). The pipeline comprised the following steps: (I) discarding the first five EPI volumes to allow for signal equilibration and steady state, (II) 3D motion correction (FSL mcflirt), (III) distortion correction (FSL fugue), (IV) rigid body coregistration of functional scans to the individual T1-weighted image (Freesurfer bbregister), (V) denoising including removal of 24 motion parameters (CPAC, Friston et al., 1996), motion, signal intensity spikes (Nipype rapidart), physiological noise in white matter and cerebrospinal fluid (CSF) (CompCor; Behzadi et al., 2007), together with linear and quadratic signal trends, (VI) band-pass filtering between 0.01-0.1 Hz (FSL fslmaths), (VII) spatial normalization to MNI152 (Montreal Neurological Institute) standard space (2 mm isotropic) via transformation parameters derived during structural preprocessing (ANTS). (VIII) The data were then spatially smoothed with a 6-mm full-width half-maximum (FWHM) Gaussian kernel (FSL fslmaths). Additionally, we calculated total intracranial volume (TIV) of each subject using the Computational Anatomy Toolbox (CAT12: http:// dbm.neuro.uni-jena.de/cat/) running on Matlab 9.3 (Mathworks, Natick, MA, USA) and used it as a covariate for further statistical analyses (Malone et al., 2015).

*BOLD Signal Variability ($SD_{BOLD}$).* Standard deviation (SD) quantifies the amount of variation or dispersion in a set of values (Garrett et al., 2015; Grady and Garrett, 2018). Higher SD in rsfMRI signal indicates greater intensity of signal fluctuation or an increased level of activation in a given area (Garrett et al., 2011). We first calculated $SD_{BOLD}$ across the whole time series for each voxel and then within 96 boundaries of preselected atlas-based regions of interests (ROIs) based on the Harvard-Oxford cortical atlas (Desikan et al., 2006). The main steps of deriving brain signal variability ($SD_{BOLD}$) from the preprocessed fMRI signal are shown in Figure 1.

The reproducible workflows containing fMRI preprocessing details can be found here: https://github.com/NeuroanatomyAndConnectivity/pipelines/releases/tag/v2.0.

## 2.3. EEG Recordings

Sixteen minutes of rsEEG were acquired on a separate day with BrainAmp MR-plus amplifiers using 61 ActiCAP electrodes (both Brain Products, Germany) attached according to the international standard 10-20 localization system (Jurcak et al., 2007) with FCz (fronto-central or cephalic electrode) as the reference. The ground electrode was located at the

sternum. Electrode impedance was kept below 5 kΩ. Continuous EEG activity was digitized

at a sampling rate of 2500 Hz and band–pass filtered online between 0.015 Hz and 1 kHz.

The experimental session was divided into 16 blocks, each lasting 60 s, with two

conditions interleaved, eyes closed (EC) and eyes open (EO), starting with the EC condition.

Changes between blocks were announced with the software Presentation (v16.5,

Neurobehavioral Systems Inc., USA). Participants were asked to sit comfortably in a chair in

a dimly illuminated, sound-shielded Faraday recording room. During the EO periods,

participants were instructed to stay awake while fixating on a black cross presented on a white

background. To maximize comparability, only EEG data from the EO condition were

analyzed, since rsfMRI data were collected only in the EO condition.

**2.4.EEG Data Analysis**

EEG processing and analyses were performed with custom Matlab (The MathWorks,

Inc, Natick, Massachusetts, USA) scripts using functions from the EEGLAB environment

(version 14.1.1b; Delorme and Makeig, 2004). The continuous EEG data were down-sampled

to 250 Hz, band-pass filtered within 1–45 Hz ($4^{th}$ order back and forth Butterworth filter) and

split into EO and EC conditions. Segments contaminated by large artefacts due to facial

muscle tensions and gross movements were removed following visual inspection, resulting in

a rejection of on average 6.6% of the recorded data. Rare occasions of artifactual channels

were excluded from the analysis. The dimensionality of the data was reduced using principal

component analysis (PCA) by selecting at least 30 principal components explaining 95% of

the total variance. Next, using independent component analysis (Infomax; Bell and

Sejnowski, 1995), the confounding sources e.g. eye-movements, eye-blinks, muscle activity,

and residual ballistocardiographic artefacts were rejected from the data.

**2.5.EEG Source Reconstruction**

Before conducting source reconstruction, preprocessed EEG signals were re-

referenced to a common average. We incorporated a standard highly detailed finite element

method (FEM) volume conduction model as described by Huang et al. (2016).

The geometry of the FEM model was based on an extended MNI/ICBM152 (International

Consortium for Brain Mapping) standard anatomy, where the source space constrained to

cortical surface and parceled to 96 ROIs based on the Harvard-Oxford atlas (Desikan et al.,

2006). We used eLORETA (exact low resolution brain electromagnetic tomography) as

implemented in as implemented in as implemented in the M/EEG Toolbox of Hamburg

(METH; Haufe and Ewald, 2016; Pascual-Marqui, 2007) to compute the cortical electrical

distribution from the scalp EEG recordings. The leadfield matrix was calculated between

1804 points located on the cortical surface to the 61 scalp electrodes. We filtered into several frequency bands, associated with brain oscillations: delta (1–3 Hz), theta (4–8 Hz), alpha (8–12 Hz), and beta (15–25 Hz). Following the singular value decomposition (SVD) of each voxel's three-dimensional time course, the dominant orientation of the source signal was identified by preserving the first SVD component. The amplitude envelope of filtered oscillations was extracted using the Hilbert transform (Rosenblum et al., 2001). Next, we applied temporal coarse graining by averaging data points in non-overlapping windows of length 0.5 s (Figure 1).

*EEG Variability ($SD_{EEG}$).* We calculated the SD of amplitude envelope of band-pass filtered oscillations on the coarse-grained signal. RsEEG signal variability ($SD_{EEG}$) was obtained for different frequency bands ($SD_{DELTA}$, $SD_{THETA}$, $SD_{ALPHA}$, $SD_{BETA}$) in each of 96 ROIs. Further, in our study we investigated variability in the amplitude of oscillatory signals from one time segment to the other. If amplitude (or power) of each signal stays the same, the variability (SD) in the amplitude of such segments will be zero. Therefore, the average amplitude of a signal is not indicative of its variability. Although amplitude and its standard deviation mathematically are different, they can show some correlation due to size effects (Immer, 1937).

Main steps toward deriving brain signal variability from the preprocessed EEG signal are shown in Figure 1. The raw and preprocessed fMRI and EEG data samples can be found at https://ftp.gwdg.de/pub/misc/MPI-Leipzig_Mind-Brain-Body-LEMON/

**Figure 1.** Main steps of deriving brain signal variability from the preprocessed resting state

fMRI and EEG signal. We calculated the standard deviation of the blood oxygen level

dependent (BOLD) signal and of the coarse-grained amplitude envelope of the rsEEG time

series for a number of standard frequency bands at the source space. Each sample of coarse-

grained amplitude envelope of the rsEEG (represented in numbers) is generated by averaging

the samples in non-overlapping windows of length 0.5 s.

**2.6. Statistical Analyses**

*Mean $SD_{BOLD}$ and $SD_{EEG}$.* For the topographic information (based on ROIs), the mean BOLD

and EEG variability were calculated by I) log-transforming the SD values, II) averaging

separately for younger and older subjects, and III) then back-transforming the values

(McDonald, 2014).

*Age and Sex Effects.* A series of non-parametric analyses of covariance (ANCOVAs, type III)

were applied to brain signal variability values in each 96 ROIs for $SD_{BOLD}$ and $SD_{EEG}$ using

age group and sex as variables of interest, adjusting for TIV and mean FD. The significance

level was controlled for using false discovery rate (FDR) correction according to Benjamini

and Hochberg (1995). Significant group differences were further examined by Tukey HSD

post-hoc comparisons. The signal variability values were log-transformed to normalize

$SD_{BOLD}$ and $SD_{EEG}$ before further analyses (assessed by Lilliefors tests at a significance

threshold of 0.05). All analyses were performed using the *aovp* function in the *lmperm*

package (Wheeler, 2016) as implemented in R (R core team, 2018).

*$SD_{BOLD}$ – $SD_{EEG}$ Correlation.* To investigate the association between each ROI of $SD_{BOLD}$ and

$SD_{EEG}$, we used pairwise Spearman's rank correlation separately for younger and older

subjects, corrected for FDR (96 ROIs). We further applied sparse canonical correlation

analysis (CCA) to show that the relationship between $SD_{BOLD}$ and $SD_{EEG}$ is not missed when

only mass bivariate correlations are used. CCA is a multivariate method to find the

independent linear combinations of variables such that the correlation between variables is

maximized (Witten et al., 2009). The sparse CCA criterion is obtained by adding a Lasso

Penalty function ($l_1$), which performs continuous shrinkage and automatic variable selection

and can solve statistical problems such as multicollinearity and overfitting (Tibshirani, 2011).

We used $l_1$ penalty as the regularization function to obtain sparse coefficients, that is, the

canonical vectors (i.e., translating from full variables to a data matrix's low-rank components

of variation) will contain exactly zero elements. Sparse CCA was performed using the R

package PMA (Penalized Multivariate Analysis; Witten et al., 2009; http://cran.r-

project.org/web/packages/PMA/). In our analyses, the significance of the correlation was

estimated using the permutation approach (N=1000) as implemented in the CCA.permute

function in R ($p_{perm} < 0.05$).

*Cognition.* The Trail Making Test (TMT) is a cognitive test measuring executive function,

including processing speed and mental flexibility (Reitan, 1955; Reitan and Wolfson, 1995).

In the first part of the test (TMT-A) the targets are all numbers, while in the second part

(TMT-B), participants need to alternate between numbers and letters. In both TMT-A and B,

the time to complete the task quantifies the performance, and lower scores indicate better

performance. Based on the previous literature, we focused on $SD_{BOLD}$, $SD_{DELTA}$, and $SD_{THETA}$

(Vlahou et al., 2015) and selected different ROIs from two research papers about the neural

correlates of the TMT: Zakzanis et al., (2005) and Jacobson et al., (2011) (Table 1). To reduce

the number of multiple comparisons (Nguyen and Holmes, 2019), these ROIs were

decomposed into singular values using the *prcomp* function belonging to *factoextra* package

(R core team, 2018), which performs SVD on the centered values. As a criterion, the

minimum total variance explained over 70% was selected (Jollife and Cadima, 2016). This

resulted in three principle components (PC) in $SD_{BOLD}$ (52.82%, 10.34%, and 7%), two PCs

in $SD_{DELTA}$ (67.37%, 10.95%), and one PC in $SD_{THETA}$ (75.63%). We also ran multiple linear

regression using task completion time in TMT-A and TMT-B as the dependent variables with

the PC scores (for $SD_{BOLD}$, $SD_{DELTA}$, and $SD_{THETA}$) and their interaction with continuous age

as independent variables. Since the residuals from the regression models fitted to the data

were not normally distributed, the TMT values were log-transformed prior to the final

analyses. These tests were conducted using the *lmp* function in *lmperm* package implemented

in R (R core team, 2018).

**Table 1.** Selected region of interests (ROIs) derived from the previous fMRI literature, and

their corresponding ROIs in Harvard-Oxford atlas to investigate the age-dependent

relationship between TMT and $SD_{BOLD}$ or $SD_{EEG}$.

| Literature | Region | Hemisphere | Harvard-Oxford Atlas |
|---|---|---|---|
| Zakzanis et al., 2005 | Middle Frontal Gyrus | Left | Middle Frontal Gyrus |
| | Precentral Gyrus | Left | Precentral Gyrus |
| | Cingulate Gyrus | Left/Right | Cingulate Gyrus, anterior/posterior |
| | Superior Frontal Gyrus | Left | Superior Frontal Gyrus |
| | Medial Frontal Gyrus | Left | Frontal Medial Cortex |
| | Insula | Left/Right | Insular Cortex |
| | Middle Temporal Gyrus | Left | Middle Temporal Gyrus, anterior/posterior/temporooccipital |
| | Superior Temporal Gyrus | Left | Superior Temporal Gyrus, anterior/posterior |
| Jacobson et al., 2011 | Fusiform Gyrus | Right | Occipital Fusiform Gyrus |
| | Inferior Middle Frontal Gyrus | Right | Middle Frontal Gyrus |
| | Precentral Gyrus | Right | Precentral Gyrus |

## 3. Results

*Mean $SD_{BOLD}$ and $SD_{EEG}$.* The topographic distribution of $SD_{BOLD}$ in younger adults revealed the largest brain signal variability values in fronto-temporal regions while in older adults it was in the frontal and occipital areas. Further, we found strongest variability across younger subjects in occipito-temporal regions for $SD_{DELTA}$, $SD_{THETA}$, $SD_{ALPHA,}$ and in medial frontal brain regions for $SD_{BETA}$, while older adults showed strongest brain signal variability in the fronto-central brain regions for $SD_{DELTA}$, in parietal-central brain regions for $SD_{THETA}$, $SD_{ALPHA}$, and in medial frontal brain regions for $SD_{BETA}$. The details of the mean values of $SD_{BOLD}$ and $SD_{EEG}$ across age groups and their topographic distributions are given in Supplementary Table 1, Supplementary Figure 2 and 3, and are also available at Neurovault (https://neurovault.org/collections/WWOKVUDV/).

*Age and Sex Effects.* The nonparametric ANCOVAs with $SD_{BOLD}$ as dependent variable demonstrated that there was a significant main effect of age group in 72 ROIs in frontal, temporal, and occipital brain regions (F-values: 13.32–61.14; Figure 2). However, there was no significant main effect of sex on $SD_{BOLD}$ and no significant interaction between age group and sex (all $p_{FDR}>0.05$). Tukey HSD post-hoc analyses showed that older subjects had decreased $SD_{BOLD}$ compared to younger adults which were presented in both sexes ($n_{ROI}=35$). The nonparametric ANCOVAs with $SD_{EEG}$ as dependent variable showed significant main effects of age group in all frequency bands: $SD_{DELTA}$ in 14 ROIs in occipital (F-values: 12.57–20.94), $SD_{THETA}$ in 16 ROIs in frontal and parietal (F-values: 13.16–40.30), $SD_{ALPHA}$ in 20 ROIs in occipital (F-values: 12.69–20.12), and $SD_{BETA}$ in 19 ROIs in central and temporal brain regions (F-values: 12.50–21.61), as shown in Figure 2. There were also significant main effects of sex in all frequency bands: $SD_{DELTA}$ in 21 ROIs in temporal and occipital (F-values: 13.24–26.63), $SD_{THETA}$ in 74 ROIs in frontal, occipital, and temporal (F-values: 12.68–30.06), $SD_{ALPHA}$ in 4 ROIs in frontal (F-values: 12.88–16.51), and $SD_{BETA}$ in 69 ROIs in temporal, occipital, and central brain regions (F-values: 12.54–35.72), as shown in Figure 3. No significant interaction effects between age group and sex on $SD_{EEG}$ were observed in any frequency band ($p_{FDR}>0.05$). Tukey HSD post-hoc analyses on $SD_{EEG}$ showed that older subjects had less brain signal variability, which was present in both sexes for $SD_{DELTA}$ ($n_{ROI}=12$), $SD_{THETA}$ ($n_{ROI}=10$), and $SD_{ALPHA}$ ($n_{ROI}=11$). Additionally, older adults showed higher $SD_{BETA}$, driven by female subjects ($n_{ROI}=15$). With regard to sex differences, post-hoc analyses showed that females had higher $SD_{DELTA}$, $SD_{THETA}$, $SD_{ALPHA}$, and $SD_{BETA}$ than males. Sex differences in $SD_{DELTA}$ ($n_{ROI}=13$) and $SD_{THETA}$ ($n_{ROI}=54$) were mostly pronounced

in younger adults, while the effect of sex in $SD_{BETA}$ ($n_{ROI}$=21) were mainly presented in older

adults ($p<0.05$). The graphical distribution of the F-values for the significant effects of age

group or sex for each ROIs are shown in Supplementary Figure 4. Additional information of

$SD_{BOLD}$ and $SD_{EEG}$ for each frequency band and for each of the 96 ROIs, split up by age group

and sex, are presented in the Supplementary Tables 2-6.

**Figure 2**. Spatial maps of significant age group differences in $SD_{BOLD}$ and $SD_{EEG}$.

We calculated the standard deviation (SD) of the blood oxygen level dependent (BOLD)

signal and of the coarse-grained amplitude envelope of the rsEEG time series for the delta (1–

3 Hz), theta (4–8 Hz), alpha (8–12 Hz), and beta (15–25 Hz) frequency bands at the source

space. Statistical significance was determined using nonparametric ANCOVAs corrected for

multiple comparisons by false discovery rates (FDR; Benjamini and Hochberg, 1995). Blue

color indicates areas where brain signal variability was lower in older than in younger adults,

while red color indicates the opposite.

**Figure 3**. Spatial maps of significant sex differences in $SD_{BOLD}$ and $SD_{EEG}$.

We calculated the standard deviation (SD) of the blood oxygen level dependent (BOLD)

signal and of the coarse-grained amplitude envelope of the rsEEG time series for the delta (1–

3 Hz), theta (4–8 Hz), alpha (8–12 Hz), and beta (15–25 Hz) frequency bands at the source

space. Statistical significance was determined using nonparametric ANCOVAs corrected for

multiple comparisons by false discovery rates (FDR; Benjamini and Hochberg, 1995). Yellow

color indicates areas where brain signal variability was higher in female subjects as compared

to male subjects in EEG.

*$SD_{BOLD} - SD_{EEG}$ Correlation*. The correlation coefficient of pairwise associations for 96 ROIs

of $SD_{BOLD}$ with $SD_{DELTA}$, $SD_{THETA}$, $SD_{ALPHA}$, and $SD_{BETA}$ ranged in younger adults from rho=-

0.200 to rho=0.223 (Supplementary Table 7) and in older adults from rho=0.386 to

rho=0.349 (Supplementary Table 8). None of the pairwise associations between $SD_{BOLD}$ and

$SD_{EEG}$ remained significant after the correction for multiple comparison corrections.

Confirmatory multivariate sparse CCA further showed that correlations between $SD_{BOLD}$ and

$SD_{EEG}$ across all subjects were rather low, highly sparse, and non-significant ($SD_{DELTA}$;

r=0.145, $p_{perm}$=0.750, $l_1$=0.367; $SD_{THETA}$; r=0.143, $p_{perm}$=0.713 $l_1$=0.7; $SD_{ALPHA}$; r=0.153,

$p_{perm}$=0.528, $l_1$=0.1; $SD_{BETA}$; r=0. 232, $p_{perm}$=0.096, $l_1$=0.633).

**Figure 4.** Distribution of correlation coefficients (rho) for the association between $SD_{BOLD}$

and $SD_{EEG}$ for A) younger (N=135) and B) older (N=54) adults for different frequency bands

across each pair of 96 regions of interests. The correlations between $SD_{BOLD}$ and $SD_{EEG}$ were

tested using pairwise Spearman's rank correlation corrected for multiple comparison by false

discovery rates (FDR; Benjamini and Hochberg, 1995).

*Cognition.* There was a significant interaction between age and $SD_{BOLD}$ in PC2 on the TMT-B

performance (adjusted $R^2 = 0.395$, $F(7,181) = 18.60$, p <.001, interaction: $\beta$= -0.002, p =

0.027). For older, but not younger participants, stronger $SD_{BOLD}$ was associated with faster

completion time in PC2, driven mainly by the left temporal gyrus as well as the left anterior

and posterior cingulate cortex (Figure 5). The regression analyses in $SD_{DELTA}$ and $SD_{THETA}$

did not show a significant association between cognition and brain signal variability

measures. The contributions of selected ROIs to the PCs resulted from SVD analyses can be

found in Supplementary Table 9. The complete multiple linear regression results can be found

in Supplementary Table 10.

**Figure 5.** Age-dependent relationship between cognitive performance and BOLD signal

variability.

The scatterplot shows the significant association between task completion time in TMT-B (x-

axis) and $SD_{BOLD}$ (adjusted $R^2 = 0.395$, $F(7,181) = 18.60$, p <.001, interaction: $\beta$= -0.002, p =

0.027) in PC2, driven mainly by the left anterior and posterior temporal gyrus, bilateral

anterior and posterior cingulate cortex.

## 4. Discussion

Comparing healthy younger and older adults, we found widespread variability reductions in BOLD signal as well as in the amplitude envelope of delta, theta, and alpha frequency of rsEEG, whereas increased variability with aging was observed in the beta-band frequency. As a complementary analysis, we also explored sex differences and found that female subjects exhibited higher EEG signal variability than male subjects; no significant sex differences were found in BOLD signal variability. There were no significant correlations between hemodynamic ($SD_{BOLD}$) and electrophysiological ($SD_{EEG}$) measures of brain signal variability, neither in the younger nor in the older adults. Our results suggest that variability measures of rsfMRI and rsEEG – while both related to aging – are dominated by different physiological origins and relate differently to age and sex.

### 4.1. BOLD Signal Variability

The first aim of our study was to investigate the effect of age on BOLD signal variability, as measured by SD of spontaneous fluctuations during rsfMRI. Consistent with recent rsfMRI studies demonstrating that BOLD signal variability decreases with age in large-scale networks (Grady and Garrett, 2018; Nomi et al., 2017), we found that older subjects had reduced $SD_{BOLD}$ in temporal and occipital brain regions but also in cortical midline structures like the precuneus, anterior and posterior cingulate cortices, as well as orbitofrontal cortex compared to younger adults. These age-related reductions in BOLD signal variability were thus especially apparent in regions of the Default Mode (DMN) and the Fronto-Parietal Network (FPN). The DMN is an intrinsically correlated network of brain regions, that is particularly active during rest or fixation blocks (Biswal et al., 2010). It reflects the systematic integration of information across the cortex (Margulies et al., 2016) and has been frequently associated with psychological functions like self-referential thought or mind-wandering, and also memory retrieval (Andrews-Hanna et al., 2014; Raichle, 2015). The FPN is involved in cognitive control processes (Spreng et al., 2013), and closely interacts with the DMN, for example during mind-wandering state (Golchert et al., 2017). Previous studies in healthy aging noted that older subjects showed lower functional connectivity in DMN and FPN regions (Damoiseaux, 2017; Damoiseaux et al., 2008; Meunier et al., 2009; Petersen et al., 2014). Similarly, an altered functional connectivity in the DMN has been found in different pathologies, for example, in Alzheimer's disease (Greicius et al., 2004) or mild cognitive impairment (Das et al., 2015). Further, we found a significant interaction between age and $SD_{BOLD}$ in temporal and cingulate cortices for performance on the cognitive task (TMT-B),

suggesting that the relationship between brain signal variability and cognitive performance
depends on the participants' age. We speculate that – in the elderly – reduced BOLD signal
variability in the DMN and the FPN, particularly in the overlapping frontal brain regions,
could be related to locally impaired function that is reflected in impaired cognitive
performance (Campbell et al., 2012). Such findings support the notion that local BOLD signal
variability may be a valuable biomarker of neurocognitive health (and disease) in aging.

Sex-specific differences in brain structure and function have been previously shown
(for a review see, Gong et al., 2011; Ruigrok et al., 2014; Sacher et al., 2013). For example,
larger total brain volume has been reported in male as compared to female subjects (Gong et
al., 2011), whereas higher cerebral blood flow (Gur et al., 1982; Rodriguez et al., 1988) and
stronger functional connectivity in the DMN (Tomasi and Volkow, 2012) were found in
females. In our exploratory analysis, we did not find significant sex differences in BOLD
signal variability when controlling for total intracranial volume as an approximation of overall
brain size.

**4.2.Electrophysiological Signal Variability**

Measures of neural variability were derived from rsEEG for several main frequency
bands (delta, theta, alpha, beta) as the standard deviation of their amplitude of envelope time
series data, analogously to the BOLD signal variability. Multimodal imaging studies have
shown that the amplitude envelope of neural oscillatory activity across frequency bands
relates to different rsfMRI networks (Brookes et al., 2011; Deligianni et al., 2014), confirming
the neurophysiological origin of the resting state networks measured with BOLD fMRI.
Additionally, these studies also concluded that different frequency bands can be related to the
same functional network, but also differentially to distinct networks (Brookes et al., 2011;
Laufs et al., 2006; Mantini et al., 2007; Meyer et al., 2013). For instance, Mantini et. al.
(2007) reported that the visual network is associated with all frequency bands except gamma
rhythm, while the sensorimotor network is primarily associated with beta-band oscillations.

In our analysis, we found age-dependent EEG signal variability changes within
networks which were associated with more than one frequency band, thus confirming that
neurons generating oscillations at different frequencies may contribute to the same network.
More precisely, we found age-related reductions in $SD_{DELTA}$ and $SD_{ALPHA}$ mainly in a visual
network (including calcarine regions, cuneal cortex, and occipital pole), $SD_{THETA}$ in posterior
DMN (e.g., posterior cingulate cortex), while an enhancement of $SD_{BETA}$ was mainly seen in
the temporal (e.g., superior/middle temporal gyrus), and central/sensorimotor (e.g.,

supramarginal gyrus) regions. These results align with previous reports of age-dependent

changes of electrophysiological activity using spectral power (Dustman et al., 1993; Vlahou et

al., 2015), and signal variability (Dustman et al., 1999; Tsvetanov et al., 2015).

Age-related decreases of alpha amplitude and alpha band variability (measured by SD

of the oscillatory signal) were previously found in posterior and occipital brain regions

(Babiloni et al., 2006; Tsvetanov et al., 2015). Alpha rhythm is a classical EEG hallmark of

resting wakefulness (Laufs et al., 2003) that is modulated by thalamo-cortical and cortico-

cortical interactions (Bazanova and Vernon, 2014; Goldman et al., 2002; Lopes Da Silva et

al., 1997; Moosmann et al., 2003). It has been suggested that the posterior alpha-frequency

plays an important role in the top-down control of cortical activation and excitability

(Klimesch, 1999). Accordingly, decreased alpha variability in occipital regions might be

associated with altered functioning of the cholinergic basal forebrain, affecting thalamo-

cortical and cortico-cortical processing. Our finding of higher temporal and sensorimotor

$SD_{BETA}$ in the elderly is in line with previous findings (Rossiter et al., 2014; Tsvetanov et al.,

2015). Aging has previously been associated with an increase in movement-related beta-band

attenuation, suggesting an enhanced motor cortex GABAergic inhibitory activity in older

individuals (Rossiter et al., 2014). Similarly, beta-band activity is thought to play a key role in

signaling maintenance of the status quo of the motor system, despite the absence of

movement (Engel and Fries, 2010). Therefore, greater $SD_{BETA}$ in sensorimotor brain regions

could be interpreted as a compensatory mechanism to account for a decline of motor

performance during aging (Quandt et al., 2016).

It should be noted that the present findings of age-related alterations of brain signal

variability at different frequencies might be influenced by several anatomical factors which

might influence EEG-generators such as reduced cortical gray matter (Babiloni et al., 2013;

Moretti et al., 2012), white-matter (Nunez et al., 2015; Valdés-Hernández et al., 2010), and

increased amount of cerebrospinal fluid (CSF; Hartikainen et al., 1992; Stomrud et al., 2010),

but also alterations of cerebral glucose metabolism (Dierks et al., 2000). Localized or global

disturbances of brain anatomy and function might lead to deviations in the EEG sources,

resulting in EEG amplitude changes. A methodological improvement for future studies will

therefore be the application of individual head models (Ziegler et al., 2014).

In addition to the effect of age on rsEEG signal variability, an exploratory analysis

showed sex differences in distinct brain regions and EEG frequencies. More precisely, we

found higher $SD_{DELTA}$ and $SD_{THETA}$ in occipito-temporal, $SD_{ALPHA}$ in frontal, and $SD_{BETA}$ in

frontal as well as occipito-temporal brain regions in female compared to male subjects. While

some studies demonstrated higher alpha (Aurlien et al., 2003), delta (Armitage, 1995), theta

(Carrier et al., 2001; Duffy et al., 1993), and beta power (Jaušovec and Jaušovec, 2010;

Matsuura et al., 1985; Veldhuizen et al., 1993) in female relative to male subjects, other

studies reported the opposite pattern (Brenner et al., 1995; Latta et al., 2005; Zappasodi et al.,

2006). These differences in EEG signal variability could be a result of different mechanisms

(biological/hormonal, cultural or developmental) involved in shaping sex differences.

Unfortunately, based on our dataset we cannot differentiate which of these potential

mechanisms might be most relevant for the observed changes.

**4.3. Association between BOLD and EEG Variability**

We further assessed how neural variability in source-reconstructed rsEEG related to

the analogous parameters of BOLD signal variability in rsfMRI using univariate and

multivariate correlation analyses. Previously, simultaneous EEG-fMRI studies have shown

meaningful relationships between fluctuations in EEG power, frequency, phase, and local

BOLD changes (for a review see, Jorge et al., 2014; Ritter and Villringer, 2006). Due to age-

related physiological (particularly cardiovascular) alterations in the brain, we expected the

relationship between BOLD and EEG signal variability to be stronger in younger than older

adults. However, in the present study, both univariate and multivariate analyses showed no

significant correlations between $SD_{BOLD}$ and $SD_{EEG}$ neither in the younger nor in the older

adults. This finding was supported by the distinct anatomical distributions of age-related

changes in BOLD and EEG signal variability, that barely showed a spatial overlap, suggesting

different underlying physiological processes. What could they be? Clearly, neuronal activity

is the main signal source for EEG- and MEG recordings as well as for EEG/MEG-based

variability measures. BOLD signal variability, however, can reflect both vascular and neural

processes (Garrett et al., 2017). While neuronal activity clearly contributes to the BOLD

signal at rest (Ma et al., 2016; Mateo et al., 2017), our results indicate, however, that neuronal

activity which is captured by EEG (or more specifically by our EEG-based measures), is not

the major determinant of BOLD variability in the resting state. Other factors that could

contribute to BOLD variability are (i) neuronal activity which is not captured by EEG and (ii)

non-neural factors such as vasomotion, or cardiac and respiratory signals (Murphy et al.,

2013). In the elderly, additional factors related to the known morphological and functional

changes of blood vessels as well as age-related metabolic changes are known to affect CBF

(Ances et al., 2009; Martin et al., 1991), $CMRO_2$ (Aanerud et al., 2012), and CVR (Liu et al.,

2013) and therefore are likely to also influence BOLD variability. Thus, given different

underlying physiology, joint EEG and fMRI variability studies might provide complementary information for a comprehensive assessment of neuronal as well as vascular factors related to aging.

## 5. Limitations

There are several limitations of our study: EEG and MRI scans were not recorded simultaneously. Therefore, we could not directly relate the two signals in a cross-correlation analysis. Furthermore, EEG and MRI were performed with different body postures (fMRI; supine, EEG; seated) known to affect brain function, for example, changes in the amplitude of the EEG signal have been related to different body postures presumably due to the shifts in cerebrospinal fluid layer thickness (Rice et al., 2013). Similarly, other experimental (e.g., visual display; Nir et al., 2006), environmental (e.g., acoustic noise in MRI; Andoh et al., 2017; Cho et al., 1998) and subject-related factors (e.g., changes of vigilance; Tagliazucchi and Laufs, 2014; Wong et al., 2013) could have introduced unintended variations in our results (Yan et al., 2013) and the influence of these factors is probably not the same for the different methods, e.g., noise in MRI or poor "control" of vigilance in MRI. For instance, given the well-known relationship between vigilance or arousal and fMRI signal fluctuations (Bijsterbosch et al., 2017; Chang et al., 2016; Haimovici et al., 2017), it is likely that the observed age-related differences in BOLD signal variability might be confounded by such within-subject (state) variability. Therefore, future rsfMRI studies may benefit from obtaining arousal-related (e.g., self-report) measures and an explicit measurement of eye movements and eye opening/closure to account for the influence of arousal on the BOLD amplitude changes. Another option would be to combine EEG and fMRI simultaneously. Yet, resting state measures of EEG (Näpflin et al., 2007) and fMRI (Shehzad et al., 2009; Zuo et al., 2010) have been shown to be reliable within-individuals across time.

In our study, the computation of the source reconstructed rsEEG required the parcellation of the brain into relatively large anatomical ROIs. It could well be that the analysis with a higher spatial resolution (e.g., at the voxel-level) with individual head models may provide additional insights about brain signal variability.

Finally, while our study aimed at comparing analogous variability measures in EEG and fMRI, future research using rsEEG and rsfMRI in the same subjects would benefit from the addition of connectivity-based measures including graph theory-based (Yu et al., 2016) or sliding-window methods (Chang et al., 2013; Qin et al., 2019).

## 6. Conclusion

In this study, we report age and sex differences of brain signal variability obtained with rsfMRI and rsEEG from the same subjects. We demonstrate extensive age-related reduction of $SD_{BOLD}$, $SD_{DELTA}$, $SD_{THETA}$, and $SD_{ALPHA}$ mainly in the DMN and the visual network, while a significant increase of $SD_{BETA}$ was mainly seen in temporal brain regions. We could not demonstrate significant associations between $SD_{BOLD}$ and $SD_{EEG}$. Our findings indicate that measurements of BOLD and EEG signal variability, respectively, are likely to stem from different physiological origins and relate differentially to age and sex. While the two types of measurements are thus not interchangeable, it seems, however, plausible that both markers of brain variability may provide complementary information about the aging process.

## 7. Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

## 8. Acknowledgements

We gratefully acknowledge the Mind-Body- Emotion group at the Max Planck Institute for Human Cognitive and Brain Sciences.

## 9.  References

Aanerud, J., Borghammer, P., Mallar Chakravarty, M., Vang, K., Rodell, A.B., Jónsdottir, K.Y.,
Møller, A., Ashkanian, M., Vafaee, M.S., Iversen, P., Johannsen, P., Gjedde, A., 2012. Brain
energy metabolism and blood flow differences in healthy aging. J. Cereb. Blood Flow Metab.
https://doi.org/10.1038/jcbfm.2012.18

Ances, B.M., Liang, C.L., Leontiev, O., Perthen, J.E., Fleisher, A.S., Lansing, A.E., Buxton, R.B.,
2009. Effects of aging on cerebral blood flow, oxygen metabolism, and blood oxygenation level
dependent responses to visual stimulation. Hum. Brain Mapp. 30, 1120–1132.
https://doi.org/10.1002/hbm.20574

Andoh, J., Ferreira, M., Leppert, I.R., Matsushita, R., Pike, B., Zatorre, R.J., 2017. How restful is it
with all that noise? Comparison of Interleaved silent steady state (ISSS) and conventional
imaging in resting-state fMRI. Neuroimage 147, 726–735.
https://doi.org/10.1016/j.neuroimage.2016.11.065

Andrews-Hanna, J.R., Smallwood, J., Spreng, R.N., 2014. The default network and self-generated
thought: Component processes, dynamic control, and clinical relevance. Ann. N. Y. Acad. Sci.
1316, 29–52. https://doi.org/10.1111/nyas.12360

Armbruster-Genc, D.J.N., Ueltzhoffer, K., Fiebach, C.J., 2016. Brain Signal Variability Differentially
Affects Cognitive Flexibility and Cognitive Stability. J. Neurosci. 36, 3978–3987.
https://doi.org/10.1523/JNEUROSCI.2517-14.2016

Armitage, R., 1995. The distribution of EEG frequencies in REM and NREM sleep stages in healthy
young adults. Sleep 18, 334–341. https://doi.org/10.1093/sleep/18.5.334

Aurlien, H., Gjerde, I.., Aarseth, J.., Eldøen, G., Karlsen, B., Skeidsvoll, H., Gilhus, N.., 2003. EEG
background activity described by a large computerized database. Clin. Neurophysiol. 115, 665–
673. https://doi.org/10.1016/j.clinph.2003.10.019

Avants, B.B., Tustison, N.J., Song, G., Cook, P.A., Klein, A., Gee, J.C., 2011. A reproducible
evaluation of ANTs similarity metric performance in brain image registration. Neuroimage 54,
2033–2044. https://doi.org/10.1016/j.neuroimage.2010.09.025

Babayan, A., Erbey, M., Kumral, D., Reinelt, J.D., Reiter, A.M.F., Röbbig, J., Lina Schaare, H.,
Uhlig, M., Anwander, A., Bazin, P.L., Horstmann, A., Lampe, L., Nikulin, V. V., Okon-Singer,
H., Preusser, S., Pampel, A., Rohr, C.S., Sacher, J., Thöne-Otto, A., Trapp, S., Nierhaus, T.,
Altmann, D., Arelin, K., Blöchl, M., Bongartz, E., Breig, P., Cesnaite, E., Chen, S., Cozatl, R.,
Czerwonatis, S., Dambrauskaite, G., Dreyer, M., Enders, J., Engelhardt, M., Fischer, M.M.,
Forschack, N., Golchert, J., Golz, L., Guran, C.A., Hedrich, S., Hentschel, N., Hoffmann, D.I.,
Huntenburg, J.M., Jost, R., Kosatschek, A., Kunzendorf, S., Lammers, H., Lauckner, M.E.,
Mahjoory, K., Kanaan, A.S., Mendes, N., Menger, R., Morino, E., Näthe, K., Neubauer, J.,
Noyan, H., Oligschläger, S., Panczyszyn-Trzewik, P., Poehlchen, D., Putzke, N., Roski, S.,
Schaller, M.C., Schieferbein, A., Schlaak, B., Schmidt, R., Gorgolewski, K.J., Schmidt, H.M.,

Schrimpf, A., Stasch, S., Voss, M., Wiedemann, A., Margulies, D.S., Gaebler, M., Villringer, A.,

2019. Data descriptor: A mind-brain-body dataset of MRI, EEG, cognition, emotion, and

peripheral physiology in young and old adults. Sci. Data 6, 180308.

https://doi.org/10.1038/sdata.2018.308

Babiloni, C., Binetti, G., Cassarino, A., Dal Forno, G., Del Percio, C., Ferreri, F., Ferri, R., Frisoni, G.,

Galderisi, S., Hirata, K., Lanuzza, B., Miniussi, C., Mucci, A., Nobili, F., Rodriguez, G.,

Romani, G.L., Rossini, P.M., Forno, G.D., Percio, C. Del, Ferreri, F., Ferri, R., Frisoni, G.,

Galderisi, S., Hirata, K., Lanuzza, B., Miniussi, C., Mucci, A., Nobili, F., Rodriguez, G.,

Romani, G.L., Rossini, P.M., 2006. Sources of cortical rhythms in adults during physiological

aging: A multicentric EEG study. Hum. Brain Mapp. 27, 162–172.

https://doi.org/10.1002/hbm.20175

Babiloni, C., Carducci, F., Lizio, R., Vecchio, F., Baglieri, A., Bernardini, S., Cavedo, E., Bozzao, A.,

Buttinelli, C., Esposito, F., Giubilei, F., Guizzaro, A., Marino, S., Montella, P., Quattrocchi,

C.C., Redolfi, A., Soricelli, A., Tedeschi, G., Ferri, R., Rossi-Fedele, G., Ursini, F., Scrascia, F.,

Vernieri, F., Pedersen, T.J., Hardemark, H.G., Rossini, P.M., Frisoni, G.B., 2013. Resting state

cortical electroencephalographic rhythms are related to gray matter volume in subjects with mild

cognitive impairment and Alzheimer's disease. Hum. Brain Mapp. 34, 1427–1446.

https://doi.org/10.1002/hbm.22005

Bazanova, O.M., Vernon, D., 2014. Interpreting EEG alpha activity. Neurosci. Biobehav. Rev. 44, 94–

110. https://doi.org/10.1016/j.neubiorev.2013.05.007

Bazin, P.L., Weiss, M., Dinse, J., Schäfer, A., Trampel, R., Turner, R., 2014. A computational

framework for ultra-high resolution cortical segmentation at 7 Tesla. Neuroimage 93, 201–209.

https://doi.org/10.1016/j.neuroimage.2013.03.077

Behzadi, Y., Restom, K., Liau, J., Liu, T.T., 2007. A component based noise correction method

(CompCor) for BOLD and perfusion based fMRI. Neuroimage 37, 90–101.

https://doi.org/10.1016/j.neuroimage.2007.04.042

Bell, A.J., Sejnowski, T.J., 1995. An Information-Maximization Approach to Blind Separation and

Blind Deconvolution. Neural Comput. 7, 1129–1159. https://doi.org/10.1162/neco.1995.7.6.1129

Benjamini, Y., Hochberg, Y., 1995. Controlling the False Discovery Rate : A Practical and Powerful

Approach to Multiple Testing Author ( s ): Yoav Benjamini and Yosef Hochberg Source :

Journal of the Royal Statistical Society . Series B ( Methodological ), Vol . 57 , No . 1 Published

by : J. R. Stat. Soc. 57, 289–300.

Bijsterbosch, J., Harrison, S., Duff, E., Alfaro-Almagro, F., Woolrich, M., Smith, S., 2017.

Investigations into within- and between-subject resting-state amplitude variations. Neuroimage

159, 57–69. https://doi.org/10.1016/j.neuroimage.2017.07.014

Birn, R.M., Murphy, K., Bandettini, P.A., 2008. The effect of respiration variations on independent

component analysis results of resting state functional connectivity. Hum. Brain Mapp. 29, 740–

750. https://doi.org/10.1002/hbm.20577

Biswal, B.B., Mennes, M., Zuo, X.-N., 2010. Toward discovery science of human brain function.

Proc. Natl. Acad. Sci. U. S. A. 107, 4734–4739. https://doi.org/10.1073/pnas.0911855107

Brenner, R.P., Ulrich, R.F., Reynolds, C.F., 1995. EEG spectral findings in healthy, elderly men and

women - sex differences. Electroencephalogr. Clin. Neurophysiol. 94, 1–5.

https://doi.org/10.1016/0013-4694(94)00234-C

Brookes, M.J., Woolrich, M., Luckhoo, H., Price, D., Hale, J.R., Stephenson, M.C., Barnes, G.R.,

Smith, S.M., Morris, P.G., 2011. Investigating the electrophysiological basis of resting state

networks using magnetoencephalography. Proc. Natl. Acad. Sci. 108, 16783–16788.

https://doi.org/10.1073/pnas.1112685108

Burgess, A., Gruzelier, J., 1993. Individual reliability of amplitude distribution in topographical

mapping of EEG. Electroencephalogr. Clin. Neurophysiol. 86, 219–223.

https://doi.org/10.1016/0013-4694(93)90101-Z

Caballero-Gaudes, C., Reynolds, R.C., 2017. Methods for cleaning the BOLD fMRI signal.

Neuroimage 154, 128–149. https://doi.org/10.1016/j.neuroimage.2016.12.018

Cabeza, R., 2001. Cognitive neuroscience of aging: Contributions of functional neuroimaging. Scand.

J. Psychol. 42, 277–286. https://doi.org/10.1111/1467-9450.00237

Cabeza, R., Albert, M., Belleville, S., Craik, F.I.M., Duarte, A., Grady, C.L., Lindenberger, U.,

Nyberg, L., Park, D.C., Reuter-Lorenz, P.A., Rugg, M.D., Steffener, J., Rajah, M.N., 2018.

Maintenance, reserve and compensation: the cognitive neuroscience of healthy ageing. Nat. Rev.

Neurosci. 19, 701–710. https://doi.org/10.1038/s41583-018-0068-2

Campbell, K.L., Grady, C.L., Ng, C., Hasher, L., 2012. Age differences in the frontoparietal cognitive

control network: Implications for distractibility. Neuropsychologia 50, 2212–2223.

https://doi.org/10.1016/j.neuropsychologia.2012.05.025

Carrier, J., Land, S., Buysse, D.J., Kupfer, D.J., Monk, T.H., 2001. The effects of age and gender on

sleep EEG power spectral density in the middle years of life (ages 20-60 years old).

Psychophysiology 38, 232–242. https://doi.org/10.1017/S0048577201991838

Chang, C., Cunningham, J.P., Glover, G.H., 2009. Influence of heart rate on the BOLD signal: The

cardiac response function. Neuroimage 44, 857–869.

https://doi.org/10.1016/j.neuroimage.2008.09.029

Chang, C., Leopold, D.A., Schölvinck, M.L., Mandelkow, H., Picchioni, D., Liu, X., Ye, F.Q., Turchi,

J.N., Duyn, J.H., 2016. Tracking brain arousal fluctuations with fMRI. Proc. Natl. Acad. Sci.

113, 4518–4523. https://doi.org/10.1073/pnas.1520613113

Chang, C., Liu, Z., Chen, M.C., Liu, X., Duyn, J.H., 2013. EEG correlates of time-varying BOLD

functional connectivity. Neuroimage 72, 227–236.

https://doi.org/10.1016/j.neuroimage.2013.01.049

Cho, Z., Chung, S., Lim, D., Wong, E.K., 1998. Effects of the Acoustic Noise of the Gradient Systems

on fMRI: Magn. Reson. Med. 39 331–335.

Cohen, M.X., 2017. Where Does EEG Come From and What Does It Mean? Trends Neurosci. 40,

208–218. https://doi.org/10.1016/j.tins.2017.02.004

Cox, R.W., 1996. AFNI: Software for Analysis and Visualization of Functional Magnetic Resonance

Neuroimages. Comput. Biomed. Res. 29, 162–173. https://doi.org/10.1006/cbmr.1996.0014

D'Esposito, M., Deouell, L.Y., Gazzaley, A., 2003. Alterations in the BOLD fMRI signal with ageing

and disease: a challenge for neuroimaging. Nat. Rev. Neurosci. 4, 863–872.

https://doi.org/10.1038/nrn1246

Damoiseaux, J.S., 2017. Effects of aging on functional and structural brain connectivity. Neuroimage

160, 32–40. https://doi.org/10.1016/j.neuroimage.2017.01.077

Damoiseaux, J.S., Beckmann, C.F., Arigita, E.J.S., Barkhof, F., Scheltens, P., Stam, C.J., Smith, S.M.,

Rombouts, S.A.R.B., 2008. Reduced resting-state brain activity in the "default network" in

normal aging. Cereb. Cortex 18, 1856–1864. https://doi.org/10.1093/cercor/bhm207

Das, S.R., Pluta, J., Mancuso, L., Kliot, D., Yushkevich, P.A., Wolk, D.A., 2015. Anterior and

posterior MTL networks in aging and MCI. Neurobiol. Aging 36, S141–S150.

https://doi.org/10.1016/j.neurobiolaging.2014.03.041

Deligianni, F., Centeno, M., Carmichael, D.W., Clayden, J.D., 2014. Relating resting-state fMRI and

EEG whole-brain connectomes across frequency bands. Front. Neurosci. 8, 1–16.

https://doi.org/10.3389/fnins.2014.00258

Delorme, A., Makeig, S., 2004. EEGLAB: an open sorce toolbox for analysis of single-trail EEG

dynamics including independent component anlaysis. J. Neurosci. Methods 134, 9–21.

https://doi.org/10.1016/j.jneumeth.2003.10.009

Desikan, R.S., Ségonne, F., Fischl, B., Quinn, B.T., Dickerson, B.C., Blacker, D., Buckner, R.L.,

Dale, A.M., Maguire, R.P., Hyman, B.T., Albert, M.S., Killiany, R.J., 2006. An automated

labeling system for subdividing the human cerebral cortex on MRI scans into gyral based regions

of interest. Neuroimage 31, 968–980. https://doi.org/10.1016/j.neuroimage.2006.01.021

Dierks, T., Jelic, V., Pascual-Marqui, R.D., Wahlund, L.O., Julin, P., Linden, D.E.J., Maurer, K.,

Winblad, B., Nordberg, A., 2000. Spatial pattern of cerebral glucose metabolism (PET)

correlates with localization of intracerebral EEG-generators in Alzheimer's disease. Clin.

Neurophysiol. 111, 1817–1824. https://doi.org/10.1016/S1388-2457(00)00427-2

Duffy, F.H., McAnulty, G.B., Albert, M.S., 1993. The pattern of age-related differences in

electrophysiological activity of healthy males and females. Neurobiol. Aging 14, 73–84.

https://doi.org/10.1016/0197-4580(93)90025-7

Dustman, R.E., Shearer, D.E., Emmerson, R.Y., 1999. Life-span changes in EEG spectral amplitude,

amplitude variability and mean frequency. Clin. Neurophysiol. 110, 1399–1409.

https://doi.org/10.1016/S1388-2457(99)00102-9

Dustman, R.E., Shearer, D.E., Emmerson, R.Y., 1993. EEG and event-related potentials in normal

aging. Prog. Neurobiol. 41, 369–401. https://doi.org/10.1016/0301-0082(93)90005-D

Engel, A.K., Fries, P., 2010. Beta-band oscillations-signalling the status quo? Curr. Opin. Neurobiol.

20, 156–165. https://doi.org/10.1016/j.conb.2010.02.015

Failla, M., Grappiolo, A., Emanuelli, G., Vitale, G., Fraschini, N., Bigoni, M., Grieco, N., Denti, M.,

Giannattasio, C., Mancia, G., 1999. Sympathetic tone restrains arterial distensibility of healthy

and atherosclerotic subjects. J. Hypertens. 17, 1117–1123. https://doi.org/10.1097/00004872-

199917080-00011

Farkas, E., Luiten, P.G.M., 2001. Cerebral microvascular pathology in aging and Alzheimer's disease,

Progress in Neurobiology. https://doi.org/10.1016/S0301-0082(00)00068-X

Fernández, A., Zuluaga, P., Abásolo, D., Gómez, C., Serra, A., Méndez, M.A., Hornero, R., 2012.

Brain oscillatory complexity across the life span. Clin. Neurophysiol. 123, 2154–2162.

https://doi.org/10.1016/j.clinph.2012.04.025

Fischl, B., 2012. FreeSurfer. Neuroimage 62, 774–781.

https://doi.org/10.1016/j.neuroimage.2012.01.021

Friston, K.J., Williams, S., Howard, R., Frackowiak, R.S.J., Turner, R., 1996. Movement-related

effects in fMRI time-series. Magn. Reson. Med. 35, 346–355.

https://doi.org/10.1002/mrm.1910350312

Garrett, D.D., Kovacevic, N., McIntosh, A.R., Grady, C.L., 2013a. The modulation of BOLD

variability between cognitive states varies by age and processing speed. Cereb. Cortex 23, 684–

693. https://doi.org/10.1093/cercor/bhs055

Garrett, D.D., Lindenberger, U., Hoge, R.D., Gauthier, C.J., 2017. Age differences in brain signal

variability are robust to multiple vascular controls. Sci. Rep. 7, 10149.

https://doi.org/10.1038/s41598-017-09752-7

Garrett, D.D., Nagel, I.E., Preuschhof, C., Burzynska, A.Z., Marchner, J., Wiegert, S., Jungehülsing,

G.J., Nyberg, L., Villringer, A., Li, S.-C., Heekeren, H.R., Bäckman, L., Lindenberger, U., 2015.

Amphetamine modulates brain signal variability and working memory in younger and older

adults. Proc. Natl. Acad. Sci. 112, 7593–7598. https://doi.org/10.1073/pnas.1504090112

Garrett, D.D., Samanez-Larkin, G.R., MacDonald, S.W.S., Lindenberger, U., McIntosh, A.R., Grady,

C.L., 2013b. Moment-to-moment brain signal variability: A next frontier in human brain

mapping? Neurosci. Biobehav. Rev. 37, 610–624.

https://doi.org/10.1016/j.neubiorev.2013.02.015

Geerligs, L., Tsvetanov, K.A., Cam-CAN, Henson, R.N., 2017. Challenges in measuring individual

differences in functional connectivity using fMRI: The case of healthy aging. Hum. Brain Mapp.

38, 4125–4156. https://doi.org/10.1002/hbm.23653

Golchert, J., Smallwood, J., Jefferies, E., Seli, P., Huntenburg, J.M., Liem, F., Lauckner, M.E.,

Oligschläger, S., Bernhardt, B.C., Villringer, A., Margulies, D.S., 2017. Individual variation in

intentionality in the mind-wandering state is reflected in the integration of the default-mode,

fronto-parietal, and limbic networks. Neuroimage 146, 226–235.

https://doi.org/10.1016/j.neuroimage.2016.11.025

Goldman, R.I., Stern, J.M., Engel, J., Cohen, M.S., 2002. Simultaneous EEG and fMRI of the alpha

rhythm. Neuroreport 13, 2487–2492. https://doi.org/10.1097/00001756-200212200-00022

Gonçalves, S.I., De Munck, J.C., Pouwels, P.J.W., Schoonhoven, R., Kuijer, J.P.A., Maurits, N.M.,

Hoogduin, J.M., Van Someren, E.J.W., Heethaar, R.M., Lopes Da Silva, F.H., 2006. Correlating

the alpha rhythm to BOLD using simultaneous EEG/fMRI: Inter-subject variability. Neuroimage

30, 203–213. https://doi.org/10.1016/j.neuroimage.2005.09.062

Gong, G., He, Y., Evans, A.C., 2011. Brain connectivity: Gender makes a difference. Neuroscientist

17, 575–591. https://doi.org/10.1177/1073858410386492

Gorgolewski, K., Burns, C.D., Madison, C., Clark, D., Halchenko, Y.O., Waskom, M.L., Ghosh, S.S.,

2011. Nipype: A Flexible, Lightweight and Extensible Neuroimaging Data Processing

Framework in Python. Front. Neuroinform. 5, 13. https://doi.org/10.3389/fninf.2011.00013

Grady, C., 2012. The cognitive neuroscience of ageing. Nat. Rev. Neurosci. 13, 491–505.

https://doi.org/10.1038/nrn3256

Grady, C.L., Garrett, D.D., 2018. Brain signal variability is modulated as a function of internal and

external demand in younger and older adults. Neuroimage 169, 510–523.

https://doi.org/10.1016/j.neuroimage.2017.12.031

Grady, C.L., Garrett, D.D., 2014. Understanding variability in the BOLD signal and why it matters for

aging. Brain Imaging Behav. https://doi.org/10.1007/s11682-013-9253-0

Greicius, M.D., Srivastava, G., Reiss, A.L., Menon, V., 2004. Default-mode network activity

distinguishes Alzheimer's disease from healthy aging: evidence from functional MRI. Proc. Natl.

Acad. Sci. U. S. A. 101, 4637–42. https://doi.org/10.1073/pnas.0308627101

Gur, R.C., Gur, R.E., Obrist, W.D., Hungerbuhler, J.P., Younkin, D., Rosen, A.D., Skolnick, B.E.,

Reivich, M., 1982. Sex and handedness differences in cerebral blood flow during rest and

cognitive activity. Science (80-. ). 217, 659–661. https://doi.org/10.1126/science.7089587

Haimovici, A., Tagliazucchi, E., Balenzuela, P., Laufs, H., 2017. On wakefulness fluctuations as a

source of BOLD functional connectivity dynamics. Sci. Rep. 7. https://doi.org/10.1038/s41598-

017-06389-4

Hartikainen, P., Soininen, H., Partanen, J., Helkala, E.L., Riekkinen, P., 1992. Aging and spectral

analysis of EEG in normal subjects: a link to memory and CSF AChE. Acta Neurol. Scand. 86,

148–155. https://doi.org/10.1111/j.1600-0404.1992.tb05057.x

Haufe, S., Ewald, A., 2016. A Simulation Framework for Benchmarking EEG-Based Brain

Connectivity Estimation Methodologies. Brain Topogr. 1–18. https://doi.org/10.1007/s10548-

016-0498-y

Hawkes, C.H., Prescott, R.J., 1973. EEG variation in healthy subjects. Electroencephalogr. Clin.

Neurophysiol. 34, 197–199. https://doi.org/10.1016/0013-4694(73)90048-5

Huang, Y., Parra, L.C., Haufe, S., 2016. The New York Head—A precise standardized volume
conductor model for EEG source localization and tES targeting. Neuroimage 140, 150–162.
https://doi.org/10.1016/j.neuroimage.2015.12.019

Hudetz, A.G., Biswal, B.B., Shen, H., Lauer, K.K., Kampine, J.P., 1998. Spontaneous Fluctuations in
Cerebral Oxygen Supply. Springer, Boston, MA, pp. 551–559. https://doi.org/10.1007/978-1-
4615-4863-8_66

Immer, F.R., 1937. Correlation between Means and Standard Deviations in Field Experiments. J. Am.
Stat. Assoc. 32, 525–531. https://doi.org/10.1080/01621459.1937.10502321

Jacobson, S.C., Blanchard, M., Connolly, C.C., Cannon, M., Garavan, H., 2011. An fMRI
investigation of a novel analogue to the Trail-Making Test. Brain Cogn. 77, 60–70.
https://doi.org/10.1016/j.bandc.2011.06.001

Jaušovec, N., Jaušovec, K., 2010. Resting brain activity: Differences between genders.
Neuropsychologia 48, 3918–3925. https://doi.org/10.1016/j.neuropsychologia.2010.09.020

Jenkinson, M., Beckmann, C.F., Behrens, T.E.J., Woolrich, M.W., Smith, S.M., 2012. Fsl.
Neuroimage 62, 782–790. https://doi.org/10.1016/j.neuroimage.2011.09.015

Jollife, I.T., Cadima, J., 2016. Principal component analysis: A review and recent developments.
Philos. Trans. R. Soc. A Math. Phys. Eng. Sci. 374, 20150202.
https://doi.org/10.1098/rsta.2015.0202

Jorge, J., Van der Zwaag, W., Figueiredo, P., 2014. EEG-fMRI integration for the study of human
brain function. Neuroimage 102, 24–34. https://doi.org/10.1016/j.neuroimage.2013.05.114

Jurcak, V., Tsuzuki, D., Dan, I., 2007. 10/20, 10/10, and 10/5 systems revisited: Their validity as
relative head-surface-based positioning systems. Neuroimage 34, 1600–1611.
https://doi.org/10.1016/j.neuroimage.2006.09.024

Kielar, A., Deschamps, T., Chu, R.K.O., Jokel, R., Khatamian, Y.B., Chen, J.J., Meltzer, J.A., 2016.
Identifying dysfunctional cortex: Dissociable effects of stroke and aging on resting state
dynamics in MEG and fmri. Front. Aging Neurosci. 8. https://doi.org/10.3389/fnagi.2016.00040

Klimesch, W., 1999. EEG alpha and theta oscillations reflect cognitive and memory performance: A
review and analysis. Brain Res. Rev. 29, 169–195. https://doi.org/10.1016/S0165-
0173(98)00056-3

Labrenz, F., Ferri, F., Wrede, K., Forsting, M., Schedlowski, M., Engler, H., Elsenbruch, S., Benson,
S., Costantini, M., 2018. Altered temporal variance and functional connectivity of BOLD signal
is associated with state anxiety during acute systemic inflammation. Neuroimage 184, 916–924.
https://doi.org/10.1016/j.neuroimage.2018.09.056

Latta, F., Leproult, R., Tasali, E., Hofmann, E., Van Cauter, E., 2005. Sex differences in delta and
alpha EEG activities in healthy older adults. Sleep 28, 1525–1534.
https://doi.org/10.1093/sleep/28.12.1525

Laufs, H., Kleinschmidt, A., Holt, J.L., Elfont, R., Krams, M., Paul, J.S., Krakow, K., 2006. Where the

BOLD signal goes when alpha EEG leaves. Neuroimage 31, 1408–1418.

https://doi.org/10.1016/j.neuroimage.2006.02.002

Laufs, H., Krakow, K., Sterzer, P., Eger, E., Beyerle, A., Kleinschmidt, A., 2003.

Electroencephalographic signatures of attentional and cognitive default modes in spontaneous

brain activity fluctuations at rest 100.

Lipsitz, L.A., Goldberger, A.L., 1992. Loss of 'Complexity' and Aging: Potential Applications of

Fractals and Chaos Theory to Senescence. JAMA J. Am. Med. Assoc. 267, 1806–1809.

https://doi.org/10.1001/jama.1992.03480130122036

Liu, P., Hebrank, A.C., Rodrigue, K.M., Kennedy, K.M., Section, J., Park, D.C., Lu, H., 2013. Age-

related differences in memory-encoding fMRI responses after accounting for decline in vascular

reactivity. Neuroimage 78, 415–425. https://doi.org/10.1016/j.neuroimage.2013.04.053

Liu, T.T., 2013. Neurovascular factors in resting-state functional MRI. Neuroimage 80, 339–348.

https://doi.org/10.1016/j.neuroimage.2013.04.071

Logothetis, N.K., 2008. What we can do and what we cannot do with fMRI. Nature 453, 869–878.

https://doi.org/10.1038/nature06976

Logothetis, N.K., Wandell, B.A., 2004. Interpreting the BOLD Signal. Annu. Rev. Physiol. 66, 735–

769. https://doi.org/10.1146/annurev.physiol.66.082602.092845

Lopes Da Silva, F.H., Pijn, J.P., Velis, D., Nijssen, P.C.G., 1997. Alpha rhythms: Noise, dynamics and

models. Int. J. Psychophysiol. 26, 237–249. https://doi.org/10.1016/S0167-8760(97)00767-8

Lopez-Larson, M.P., Anderson, J.S., Ferguson, M.A., Yurgelun-Todd, D., 2011. Local brain

connectivity and associations with gender and age. Dev. Cogn. Neurosci. 1, 187–197.

https://doi.org/10.1016/j.dcn.2010.10.001

Ma, Y., Shaik, M.A., Kozberg, M.G., Kim, S.H., Portes, J.P., Timerman, D., Hillman, E.M.C., 2016.

Resting-state hemodynamics are spatiotemporally coupled to synchronized and symmetric neural

activity in excitatory neurons. Proc. Natl. Acad. Sci. 113, E8463–E8471.

https://doi.org/10.1073/pnas.1525369113

Malone, I.B., Leung, K.K., Clegg, S., Barnes, J., Whitwell, J.L., Ashburner, J., Fox, N.C., Ridgway,

G.R., 2015. Accurate automatic estimation of total intracranial volume: A nuisance variable with

less nuisance. Neuroimage 104, 366–372. https://doi.org/10.1016/j.neuroimage.2014.09.034

Maltez, J., Hyllienmark, L., Nikulin, V. V., Brismar, T., 2004. Time course and variability of power in

different frequency bands of EEG during resting conditions. Neurophysiol. Clin. 34, 195–202.

https://doi.org/10.1016/j.neucli.2004.09.003

Mantini, D., Perrucci, M.G., Del Gratta, C., Romani, G.L., Corbetta, M., 2007. Electrophysiological

signatures of resting state networks in the human brain. Proc. Natl. Acad. Sci. 104, 13170–

13175. https://doi.org/10.1073/pnas.0700668104

Margulies, D.S., Ghosh, S.S., Goulas, A., Falkiewicz, M., Huntenburg, J.M., Langs, G., Bezgin, G.,

Eickhoff, S.B., Castellanos, F.X., Petrides, M., Jefferies, E., Smallwood, J., 2016. Situating the

900 default-mode network along a principal gradient of macroscale cortical organization. Proc. Natl.
901 Acad. Sci. 113, 12574–12579. https://doi.org/10.1073/pnas.1608282113

902 Martin, J., Friston, K.J., Colebatch, J.G., Frackowiak, R.S.J., Unit, M.R.C.C., Hospital, H., 1991.
903 Decreases in Regional Cerebral Blood Flow with Normal Aging 684–689.

904 Mateo, C., Knutsen, P.M., Tsai, P.S., Shih, A.Y., Kleinfeld, D., 2017. Entrainment of Arteriole
905 Vasomotor Fluctuations by Neural Activity Is a Basis of Blood-Oxygenation-Level-Dependent
906 "Resting-State" Connectivity. Neuron 96, 936-948.e3.
907 https://doi.org/10.1016/j.neuron.2017.10.012

908 Matsuura, M., Yamamoto, K., Fukuzawa, H., Okubo, Y., Uesugi, H., Moriiwa, M., Kojima, T.,
909 Shimazono, Y., 1985. Age development and sex differences of various EEG elements in healthy
910 children and adults - Quantification by a computerized waveform recognition method.
911 Electroencephalogr. Clin. Neurophysiol. 60, 394–406. https://doi.org/10.1016/0013-
912 4694(85)91013-2

913 McBride, J.C., Zhao, X., Munro, N.B., Smith, C.D., Jicha, G.A., Hively, L., Broster, L.S., Schmitt,
914 F.A., Kryscio, R.J., Jiang, Y., 2014. Spectral and complexity analysis of scalp EEG
915 characteristics for mild cognitive impairment and early Alzheimer's disease. Comput. Methods
916 Programs Biomed. 114, 153–163. https://doi.org/10.1016/j.cmpb.2014.01.019

917 McDonald, J.H., 2014. Handbook of Biological Statistics. Sparky House Publ. 180–185.

918 Meunier, D., Achard, S., Morcom, A., Bullmore, E., 2009. Age-related changes in modular
919 organization of human brain functional networks. Neuroimage 44, 715–723.
920 https://doi.org/10.1016/j.neuroimage.2008.09.062

921 Meyer, M.C., Janssen, R.J., Van Oort, E.S.B., Beckmann, C.F., Barth, M., 2013. The Quest for EEG
922 Power Band Correlation with ICA Derived fMRI Resting State Networks. Front. Hum. Neurosci.
923 7, 315. https://doi.org/10.3389/fnhum.2013.00315

924 Moosmann, M., Ritter, P., Krastel, I., Brink, A., Thees, S., Blankenburg, F., Taskin, B., Obrig, H.,
925 Villringer, A., 2003. Correlates of alpha rhythm in functional magnetic resonance imaging and
926 near infrared spectroscopy. Neuroimage 20, 145–158. https://doi.org/10.1016/S1053-
927 8119(03)00344-6

928 Moretti, D. V., Paternicò, D., Binetti, G., Zanetti, O., Frisoni, G.B., 2012. EEG markers are associated
929 to gray matter changes in thalamus and basal ganglia in subjects with mild cognitive impairment.
930 Neuroimage 60, 489–496. https://doi.org/10.1016/j.neuroimage.2011.11.086

931 Murayama, Y., Bießmann, F., Logothetis, N.K., Oeltermann, A., Müller, K.-R., Meinecke, F.C.,
932 Augath, M., 2010. Relationship between neural and hemodynamic signals during spontaneous
933 activity studied with temporal kernel CCA. Magn. Reson. Imaging 28, 1095–1103.
934 https://doi.org/10.1016/j.mri.2009.12.016

935 Murphy, K., Birn, R.M., Bandettini, P.A., 2013. Resting-state fMRI confounds and cleanup.
936 Neuroimage 80, 349–359. https://doi.org/10.1016/j.neuroimage.2013.04.001

Näpflin, M., Wildi, M., Sarnthein, J., 2007. Test-retest reliability of resting EEG spectra validates a
statistical signature of persons. Clin. Neurophysiol. 118, 2519–2524.
https://doi.org/10.1016/j.clinph.2007.07.022

Nguyen, L.H., Holmes, S., 2019. Ten quick tips for effective dimensionality reduction. PLOS Comput.
Biol. 15, e1006907. https://doi.org/10.1371/journal.pcbi.1006907

Niessing, J., Ebisch, B., Schmidt, K.E., Niessing, M., Singer, W., Galuske, R.A.W., 2005.
Neuroscience: Hemodynamic signals correlate tightly with synchronized gamma oscillations.
Science (80-. ). 309, 948–951. https://doi.org/10.1126/science.1110948

Nir, Y., Hasson, U., Levy, I., Yeshurun, Y., Malach, R., 2006. Widespread functional connectivity and
fMRI fluctuations in human visual cortex in the absence of visual stimulation. Neuroimage 30,
1313–1324. https://doi.org/10.1016/j.neuroimage.2005.11.018

Nomi, J.S., Bolt, T.S., Ezie, C.E.C., Uddin, L.Q., Heller, A.S., 2017. Moment-to-Moment BOLD
Signal Variability Reflects Regional Changes in Neural Flexibility across the Lifespan. J.
Neurosci. 37, 5539–5548. https://doi.org/10.1523/JNEUROSCI.3408-16.2017

Nomi, J.S., Schettini, E., Voorhies, W., Bolt, T.S., Heller, A.S., Uddin, L.Q., 2018. Resting-State
Brain Signal Variability in Prefrontal Cortex Is Associated With ADHD Symptom Severity in
Children. Front. Hum. Neurosci. 12, 90. https://doi.org/10.3389/fnhum.2018.00090

Nunez, P.L., Srinivasan, R., Fields, R.D., 2015. EEG functional connectivity, axon delays and white
matter disease. Clin. Neurophysiol. 126, 110–120. https://doi.org/10.1016/j.clinph.2014.04.003

Oken, B.S., Chiappa, K.H., 1988. Short-term variability in EEG frequency analysis.
Electroencephalogr. Clin. Neurophysiol. 69, 191–198. https://doi.org/10.1016/0013-
4694(88)90128-9

Petersen, S.E., Savalia, N.K., Chan, M.Y., Park, D.C., Wig, G.S., 2014. Decreased segregation of
brain systems across the healthy adult lifespan. Proc. Natl. Acad. Sci. 111, E4997–E5006.
https://doi.org/10.1073/pnas.1415122111

Qin, Y., Jiang, S., Zhang, Q., Dong, L., Jia, X., He, H., Yao, Y., Yang, H., Zhang, T., Luo, C., Yao,
D., 2019. BOLD-fMRI activity informed by network variation of scalp EEG in juvenile
myoclonic epilepsy. NeuroImage Clin. 22, 101759. https://doi.org/10.1016/j.nicl.2019.101759

Quandt, F., Bönstrup, M., Schulz, R., Timmermann, J.E., Zimerman, M., Nolte, G., Hummel, F.C.,
2016. Spectral variability in the aged brain during fine motor control. Front. Aging Neurosci. 8.
https://doi.org/10.3389/fnagi.2016.00305

Raichle, M.E., 2015. The Brain's Default Mode Network. Annu. Rev. Neurosci. 38, 433–447.
https://doi.org/10.1146/annurev-neuro-071013-014030

Reitan, R.M., 1955. Certain differential effects of left and right cerebral lesions in human adults. J
Comp Physiol Psychol 48, 474–477.

Reitan, R.M., Wolfson, D., 1995. Category Test and Trail Making Test as Measures of Frontal Lobe
Functions. Clin. Neuropsychol. 9, 50–56. https://doi.org/10.1080/13854049508402057

Rice, J.K., Rorden, C., Little, J.S., Parra, L.C., 2013. Subject position affects EEG magnitudes.
Neuroimage 64, 476–484. https://doi.org/10.1016/j.neuroimage.2012.09.041

Ritter, P., Moosmann, M., Villringer, A., 2009. Rolandic alpha and beta EEG rhythms' strengths are
inversely related to fMRI-BOLD signal in primary somatosensory and motor cortex. Hum. Brain
Mapp. 30, 1168–1187. https://doi.org/10.1002/hbm.20585

Ritter, P., Villringer, A., 2006. Simultaneous EEG-fMRI. Neurosci. Biobehav. Rev. 30, 823–838.
https://doi.org/10.1016/j.neubiorev.2006.06.008

Rodriguez, G., Warkentin, S., Risberg, J., Rosadini, G., 1988. Sex differences in regional cerebral
blood flow. J. Cereb. Blood Flow Metab. 8, 783–789. https://doi.org/10.1038/jcbfm.1988.133

Rokem, a, Trumpis, M., Perez, F., 2009. Nitime: time-series analysis for neuroimaging data. Proc. 8th
Python Sci. Conf. (SciPy 2009) 1–8.

Rosenblum, M., Pikovsky, A., Kurths, J., Schäfer, C., Tass, P.A., 2001. Phase synchronization: From
theory to data analysis. Handb. Biol. Phys. 4, 279–321. https://doi.org/10.1016/S1383-
8121(01)80012-9

Rossiter, H.E., Davis, E.M., Clark, E. V., Boudrias, M.H., Ward, N.S., 2014. Beta oscillations reflect
changes in motor cortex inhibition in healthy ageing. Neuroimage 91, 360–365.
https://doi.org/10.1016/j.neuroimage.2014.01.012

Ruigrok, A.N.V., Salimi-Khorshidi, G., Lai, M.C., Baron-Cohen, S., Lombardo, M. V., Tait, R.J.,
Suckling, J., 2014. A meta-analysis of sex differences in human brain structure. Neurosci.
Biobehav. Rev. 39, 34–50. https://doi.org/10.1016/j.neubiorev.2013.12.004

Sacher, J., Neumann, J., Okon-Singer, H., Gotowiec, S., Villringer, A., 2013. Sexual dimorphism in
the human brain: Evidence from neuroimaging. Magn. Reson. Imaging 31, 366–375.
https://doi.org/10.1016/j.mri.2012.06.007

Scheeringa, R., Fries, P., Petersson, K.M., Oostenveld, R., Grothe, I., Norris, D.G., Hagoort, P.,
Bastiaansen, M.C.M., 2011. Neuronal Dynamics Underlying High- and Low-Frequency EEG
Oscillations Contribute Independently to the Human BOLD Signal. Neuron 69, 572–583.
https://doi.org/10.1016/j.neuron.2010.11.044

Seaquist, E.R., Chen, W., Benedict, L.E., Ugurbil, K., Kwag, J.H., Zhu, X.H., Nelson, C.A., 2007.
Insulin reduces the BOLD response but is without effect on the VEP during presentation of a
visual task in humans. J. Cereb. Blood Flow Metab. 27, 154–160.
https://doi.org/10.1038/sj.jcbfm.9600316

Shehzad, Z., Kelly, A.M.C., Reiss, P.T., Gee, D.G., Gotimer, K., Uddin, L.Q., Lee, S.H., Margulies,
D.S., Roy, A.K., Biswal, B.B., Petkova, E., Castellanos, F.X., Milham, M.P., 2009. The Resting
Brain: Unconstrained yet Reliable. Cereb. Cortex 19, 2209–2229.
https://doi.org/10.1093/cercor/bhn256

Sleimen-Malkoun, R., Perdikis, D., Muller, V., Blanc, J.-L., Huys, R., Temprado, J.-J., Jirsa, V.K.,
2015. Brain Dynamics of Aging: Multiscale Variability of EEG Signals at Rest and during an

Auditory Oddball Task. eNeuro 2. https://doi.org/10.1523/ENEURO.0067-14.2015

Smits, F.M., Porcaro, C., Cottone, C., Cancelli, A., Rossini, P.M., Tecchio, F., 2016.

Electroencephalographic fractal dimension in healthy ageing and Alzheimer's disease. PLoS One

11, 1–16. https://doi.org/10.1371/journal.pone.0149587

Speelman, C.P., McGann, M., 2013. How mean is the mean? Front. Psychol. 4, 1–12.

https://doi.org/10.3389/fpsyg.2013.00451

Spreng, R.N., Sepulcre, J., Turner, G.R., Stevens, W.D., Schacter, D.L., 2013. Intrinsic Architecture

Underlying the Relations among the Default, Dorsal Attention, and Frontoparietal Control

Networks of the Human Brain. J. Cogn. Neurosci. 25, 74–86.

https://doi.org/10.1162/jocn_a_00281

Steriade, M., 2006. Grouping of brain rhythms in corticothalamic systems. Neuroscience 137, 1087–

1106. https://doi.org/10.1016/j.neuroscience.2005.10.029

Stomrud, E., Hansson, O., Minthon, L., Blennow, K., Rosén, I., Londos, E., 2010. Slowing of EEG

correlates with CSF biomarkers and reduced cognitive speed in elderly with normal cognition

over 4 years. Neurobiol. Aging 31, 215–223.

https://doi.org/10.1016/j.neurobiolaging.2008.03.025

Tagliazucchi, E., Laufs, H., 2014. Decoding Wakefulness Levels from Typical fMRI Resting-State

Data Reveals Reliable Drifts between Wakefulness and Sleep. Neuron 82, 695–708.

https://doi.org/10.1016/j.neuron.2014.03.020

Thompson, G.J., 2018. Neural and metabolic basis of dynamic resting state fMRI. Neuroimage 180,

448–462. https://doi.org/10.1016/j.neuroimage.2017.09.010

Tibshirani, R., 2011. Regression shrinkage and selection via the lasso: a retrospective. J. R. Stat. Soc.

Ser. B (Statistical Methodol. 73, 273–282. https://doi.org/10.1111/j.1467-9868.2011.00771.x

Tomasi, D., Volkow, N.D., 2012. Aging and functional brain networks. Mol. Psychiatry 17, 549–558.

https://doi.org/10.1038/mp.2011.81

Tsvetanov, K.A., Henson, R.N.A., Tyler, L.K., Davis, S.W., Shafto, M.A., Taylor, J.R., Williams, N.,

Cam-Can, Rowe, J.B., 2015. The effect of ageing on fMRI: Correction for the confounding

effects of vascular reactivity evaluated by joint fMRI and MEG in 335 adults. Hum. Brain Mapp.

36, 2248–2269. https://doi.org/10.1002/hbm.22768

Vaillancourt, D.E., Newell, K.M., 2002. Changing complexity in human behavior and physiology

through aging and disease. Neurobiol Aging 23, 1–11. https://doi.org/10.1016/S0197-

4580(02)00052-0

Valdés-Hernández, P.A., Ojeda-González, A., Martínez-Montes, E., Lage-Castellanos, A., Virués-

Alba, T., Valdés-Urrutia, L., Valdes-Sosa, P.A., 2010. White matter architecture rather than

cortical surface area correlates with the EEG alpha rhythm. Neuroimage 49, 2328–2339.

https://doi.org/10.1016/j.neuroimage.2009.10.030

Veldhuizen, R.J., Jonkman, E.J., Poortvliet, D.C.J., 1993. Sex differences in age regression parameters

of healthy adults-normative data and practical implications. Electroencephalogr. Clin.

Neurophysiol. 86, 377–384. https://doi.org/10.1016/0013-4694(93)90133-G

Villringer, A., Dirnagl, U., 1995. Coupling of brain activity and cerebral blood flow: basis of

functional neuroimaging. Cerebrovasc. Brain Metab. Rev. 7, 240—276.

Vlahou, E.L., Thurm, F., Kolassa, I.-T., Schlee, W., 2015. Resting-state slow wave power, healthy

aging and cognitive performance. Sci. Rep. 4, 5101. https://doi.org/10.1038/srep05101

Wang, R., Foniok, T., Wamsteeker, J.I., Qiao, M., Tomanek, B., Vivanco, R.A., Tuor, U.I., 2006.

Transient blood pressure changes affect the functional magnetic resonance imaging detection of

cerebral activation. Neuroimage 31, 1–11. https://doi.org/10.1016/j.neuroimage.2005.12.004

Wheeler, R.E., 2016. lmPerm: Permutation tests for linear models. http://CRAN.R-

project.org/package=lmPerm. 24 p.

Witten, D.M., Tibshirani, R., Hastie, T., 2009. A penalized matrix decomposition, with applications to

sparse principal components and canonical correlation analysis. Biostatistics 10, 515–534.

https://doi.org/10.1093/biostatistics/kxp008

Wong, C.W., Olafsson, V., Tal, O., Liu, T.T., 2013. The amplitude of the resting-state fMRI global

signal is related to EEG vigilance measures. Neuroimage 83, 983–990.

https://doi.org/10.1016/j.neuroimage.2013.07.057

Yan, C.G., Craddock, R.C., Zuo, X.N., Zang, Y.F., Milham, M.P., 2013. Standardizing the intrinsic

brain: Towards robust measurement of inter-individual variation in 1000 functional

connectomes. Neuroimage 80, 246–262. https://doi.org/10.1016/j.neuroimage.2013.04.081

Yang, A.C., Huang, C.C., Yeh, H.L., Liu, M.E., Hong, C.J., Tu, P.C., Chen, J.F., Huang, N.E., Peng,

C.K., Lin, C.P., Tsai, S.J., 2013. Complexity of spontaneous BOLD activity in default mode

network is correlated with cognitive function in normal male elderly: A multiscale entropy

analysis. Neurobiol. Aging 34, 428–438. https://doi.org/10.1016/j.neurobiolaging.2012.05.004

Yu, Q., Wu, L., Bridwell, D.A., Erhardt, E.B., Du, Y., He, H., Chen, J., Liu, P., Sui, J., Pearlson, G.,

Calhoun, V.D., 2016. Building an EEG-fMRI Multi-Modal Brain Graph: A Concurrent EEG-

fMRI Study. Front. Hum. Neurosci. 10, 1–17. https://doi.org/10.3389/fnhum.2016.00476

Zakzanis, K., Mraz, R., Graham, S., 2005. An fMRI study of the trail making test. Neuropsychologia.

Zappasodi, F., Marzetti, L., Olejarczyk, E., Tecchio, F., Pizzella, V., 2015. Age-related changes in

electroencephalographic signal complexity. PLoS One 10, 1–13.

https://doi.org/10.1371/journal.pone.0141995

Zappasodi, F., Pasqualetti, P., Tombini, M., Ercolani, M., Pizzella, V., Rossini, P.M., Tecchio, F.,

2006. Hand cortical representation at rest and during activation: Gender and age effects in the

two hemispheres. Clin. Neurophysiol. 117, 1518–1528.

https://doi.org/10.1016/j.clinph.2006.03.016

Ziegler, E., Chellappa, S.L., Gaggioni, G., Ly, J.Q.M., Vandewalle, G., André, E., Geuzaine, C.,

Phillips, C., 2014. A finite-element reciprocity solution for EEG forward modeling with realistic

individual head models. Neuroimage 103, 542–551.

https://doi.org/10.1016/j.neuroimage.2014.08.056

Zöller, D., Schaer, M., Scariati, E., Padula, M.C., Eliez, S., Van De Ville, D., 2017. Disentangling

resting-state BOLD variability and PCC functional connectivity in 22q11.2 deletion syndrome.

Neuroimage 149, 85–97. https://doi.org/10.1016/j.neuroimage.2017.01.064

Zuo, X.N., Di Martino, A., Kelly, C., Shehzad, Z.E., Gee, D.G., Klein, D.F., Castellanos, F.X.,

Biswal, B.B., Milham, M.P., 2010. The oscillating brain: Complex and reliable. Neuroimage 49,

1432–1445. https://doi.org/10.1016/j.neuroimage.2009.09.037

1093