APEC: An accesson-based method for single-cell chromatin accessibility analysis

Bin Li[1,3], Young Li[1,3], Kun Li[1], Lianbang Zhu[1], Qiaoni Yu[1], Jingwen Fang[1], Pengfei Cai[1], Chen Jiang[1], Kun Qu[1,2,*]

Affiliation

[1]Division of Molecular Medicine, Hefei National Laboratory for Physical Sciences at Microscale, The CAS Key Laboratory of Innate Immunity and Chronic Disease, School of Life Sciences, University of Science and Technology of China

[2]CAS Center for Excellence in Molecular Cell Sciences, University of Science and Technology of China

[3]Cofirst authors

*Correspondence: Kun Qu (qukun@ustc.edu.cn)

Contact Information

Kun Qu, Ph.D.

Division of Molecular Medicine, Hefei National Laboratory for Physical Sciences at Microscale, The CAS Key Laboratory of Innate Immunity and Chronic Disease, School of Life Sciences, University of Science and Technology of China

Hefei, Anhui, China, 230027

Email: qukun@ustc.edu.cn

Phone: +86-551-63606257

Abstract:

The development of sequencing technologies has promoted the survey of genome-wide

chromatin accessibility at single-cell resolution; however, comprehensive analysis of single-cell

epigenomic profiles remains a challenge. Here, we introduce an accessibility pattern-based

epigenomic clustering (APEC) method, which classifies each individual cell by groups of

accessible regions with synergistic signal patterns termed "accessons". By integrating with other

analytical tools, this python-based APEC package greatly improves the accuracy of unsupervised

single-cell clustering for many different public data sets. APEC also identifies significant

differentially accessible sites, predicts enriched motifs, and projects pseudotime trajectories.

Furthermore, we developed a fluorescent tagmentation- and FACS-sorting-based single-cell

ATAC-seq technique named ftATAC-seq and investigated the per cell regulome dynamics of

mouse thymocytes. Associated with ftATAC-seq, APEC revealed a detailed epigenomic

heterogeneity of thymocytes, characterized the developmental trajectory and predicted the

regulators that control the stages of maturation process. Overall, this work illustrates a powerful

approach to study single-cell epigenomic heterogeneity and regulome dynamics.

INTRODUCTION

As a technique for probing genome-wide chromatin accessibility in a small number of cells *in vivo*, the assay for transposase-accessible chromatin with high-throughput sequencing (ATAC-seq) has been widely applied to investigate the cellular regulomes of many important biological processes[1], such as hematopoietic stem cell (HSC) differentiation[2-4], embryonic development[5, 6], neuronal activity and regeneration[7-10], tumor cell metastasis[11], and patient responses to anticancer drug treatment[12]. Recently, several experimental schemes have been developed to capture chromatin accessibility at single-cell/nucleus resolution, i.e., single-cell ATAC-seq (scATAC-seq)[13], single-nucleus ATAC-seq (snATAC-seq)[14, 15], and single-cell combinatorial indexing ATAC-seq (sci-ATAC-seq)[16, 17], which significantly extended researchers' ability to uncover cell-to-cell epigenetic variation and other fundamental mechanisms that generate heterogeneity from identical DNA sequences. By contrast, the in-depth analysis of single-cell chromatin accessibility profiles for this purpose remains a challenge. Numerous efficient algorithms have been developed to accurately normalize, cluster and visualize cells from single-cell transcriptome sequencing profiles, including but not limited to SCnorm[18], Seurat[19], SC3[20], SIMLR[21], bigSCale[22], and SCANPY[23]. However, most of these algorithms are not directly compatible with a single-cell ATAC-seq dataset, for which the signal matrix is much sparser. To characterize scATAC-seq data, the Greenleaf lab developed an algorithm named chromVAR[24], which aggregates mapped reads at accessible sites based on annotated motifs of known transcription factors (TFs) and thus projects the sparse per accessible peak per cell matrix to a bias-corrected deviation per motif per cell matrix and significantly stabilizes the data matrix for downstream clustering analysis. Other mathematical tools, such as the latent semantic indexing (LSI) and density-based clustering methods, have also been applied to process single-cell/nucleus ATAC-seq data[15, 17]. However, none of these methods can precisely distinguish cells from low sequencing depth without prior knowledge of essential principle components or TF motifs. Therefore, a refined algorithm is urgently needed to better categorize cell subgroups with minor differences under low coverage, thereby providing a deeper mechanistic understanding of single-cell epigenetic heterogeneity and regulation.

RESULTS

Accesson-based algorithm improves single-cell clustering

Here, we introduce a new single-cell chromatin accessibility analysis toolkit named APEC (accessibility pattern-based epigenomic clustering), which combines peaks with the same signal fluctuation among all single cells into peak groups, termed "accessons", and converts the original sparse cell-peak matrix to a much denser cell-accesson matrix for cell type categorization (Figure 1a). In contrast to previous motif-based methods (e.g., chromVAR), this accesson-based reduction scheme naturally groups synergistic accessible regions together without a priori knowledge of genetic information (such as TF motifs) and provides more efficient, accurate and rapid cell clustering from single-cell ATAC-seq profiles. More conveniently, APEC integrates all necessary procedures, from raw sequence trimming, alignment, and quality control (Supplementary Figure 1a-1c) to cell clustering, motif enrichment, and pseudotime trajectory prediction into a head-to-toe program package that has been made available on GitHub (https://github.com/QuKunLab/APEC).

To test the performance of APEC, we first obtained data from previous publications, which performed scATAC-seq on lymphoid-primed multipotent progenitors (LMPPs), monocytes, HL-60 lymphoblastoid cells (HL60), and blast cells and leukemic stem cells (LSCs) from two acute myeloid leukemia (AML) patients[24]. Compared to the motif-based method chromVAR[24, 25], this new accesson-based algorithm more precisely and clearly clustered cells into their corresponding identities (Figure 1b-1e). For instance, distinct cell types, such as LMPPs, monocytes and HL60 cells, were more vividly separated from each other (Adjusted Rand Index (ARI)=0.95, compared to ARI=0.59 for chromVAR); similar cells, such as the blast cells and LSCs from two AML patients, were ambiguous in chromVAR (ARI=0.36) but were more clearly categorized in both the hierarchical clustering heatmap and the tSNE scattering plot in APEC (ARI=0.69). The contribution of the minor differences between similar cells is aggregated in accessons but diluted in motifs. For example, APEC identified prominent superenhancers around the E3 ligase inhibitor gene *N4BP1*[26] and the MLL fusion gene *GPHN*[27] in the LSC cells from AML patient 1 (P1-LSC) but not in the other cell types (Figure 1f, Supplementary Figure 1d). We noticed that all peaks in these superenhancers were classified into one accesson that was critical for distinguishing P1-LSCs from P2-LSCs, P1-blast cells and P2-blast cells. However, these peaks were distributed in multiple TF motifs, which significantly diluted the contributions of the minor differences (Supplementary Figure 1e-f). To test the robustness of APEC at low sequencing depth, we

randomly selected reads from the raw data and calculated the ARI values for each sampled data.

Compared with chromVAR, the APEC algorithm exhibits better robustness at sequencing depth

as low as 10% of the original data (Supplementary Figure 1g).

APEC is applicable to multiple single-cell chromatin detection techniques

To evaluate the compatibility and performance of APEC with other single-cell chromatin

accessibility detection techniques, such as snATAC-seq[15], transcript-indexed scATAC-seq[28] and

sciATAC-seq[16], APEC was also tested with the data sets generated by those experiments. For

example, APEC discovered 10 cell subpopulations in adult mouse forebrain snATAC-seq data[15],

including three clusters of excitatory neurons (EX1-3), five groups of inhibitory neurons (IN1-4),

astroglia cells (AC), oligodendrocyte cells (OC), and microglial cells (MG; Figure 2a & 2b), as

defined by the chromatin accessibilities at the loci of cell type-specific genes (Figure 2c).

Compared to published results[15], APEC identified 4 rather than 2 distinct inhibitory subpopulations,

among which IN1 and IN4 were more similar and IN2 and IN3 were more distinct (Figure 2d). The

motif enrichment analysis module in APEC identified cell type-specific regulators that are

consistent with previous publications[15]. For example, the NEUROD1 and OLIG2 motifs were

generally enriched on excitatory clusters (EX1~3); the MEF2C motif was more enriched on EX3

than on EX1/2 neurons; the motifs of MEIS2 and DLX2 were differentially enriched on two

subtypes of inhibitory neurons (IN2 and IN3, respectively); and the NOTO, SOX2, and ETS1

motifs were enriched on the AC, OC, and MG clusters, respectively (Figure 2e). These results

suggest that APEC is capable of identifying cell subtype-specific regulators.

Since the divergence of the gene expression levels in a single cell is much greater than

that of the chromatin accessibilities, single-cell transcriptome analysis usually identifies more cell

subpopulations. Therefore, it is critical to anchor the cell types identified from scATAC-seq to

those from scRNA-seq. Lake et al. identified dozens of excitatory and inhibitory neuronal subtypes

in the adult human brain using snDrop-seq and scTHS-seq experiments[14] and provided tens of

signature genes that distinguished these cell types. Interestingly, the accessons that represent

these signature genes were also distinctly enriched at corresponding clusters of neurons. For

example, the upright part of the EX1 cell cluster in snATAC-seq enriched accessons represents

the genes *Cbln2* and *Col5a2*, which are specific genes in clusters Ex1/2/3a that were defined in

the scDrop-seq data (Figure 2f). The left part of the EX1 cell cluster in snATAC-seq matched the

Ex3b/3c/3e clusters in scDrop-seq (marked by *Nefm*), EX2 matched Ex4/5/6 (marked by *Foxp2*

and *Pcp4*), and EX3 matched Ex3d (marked by *Phactr2*). The same method also works to anchor

inhibitory neurons, as the IN2 cells in the snATAC-seq data corresponded to the In1/2 clusters in

the scDrop-seq data (marked by *Cck* and *Cnr1*), the IN3 cells corresponded to the In6b/8 clusters

(marked by *Stxbp6* and *Tac1*), the IN4 cells corresponded to the In1c/3 clusters (marked by *Vip*

and *Tshz2*), and the low right branch of IN1 corresponded to the In7 cluster (marked by *Npy*)

(Figure 2g). These results highlight the potential advantages of the accesson-based method for

integrative analysis of scRNA-seq and scATAC-seq data.

In addition, due to the sparser per-cell-per-peak fragment count matrix, more than 29.7%

(946 out of 3034) of high-quality cells were previously unable to be correctly assigned into any

subpopulation of interest[15], but APEC successfully categorized all cells into their corresponding

subtypes, confirming its high sensitivity. In contrast, chromVAR misclustered AC and EX4 with

inhibitory neurons, although the same parameters were applied (Supplementary Figure 2a-2c).

These results confirm that this accesson-based APEC method can better distinguish and

categorize single cells with great sensitivity and reliability.

APEC constructs a pseudotime trajectory that predicts cell differentiation lineage

Cells are not static but dynamic entities, and they have a history, particularly a

developmental history. Although single-cell experiments often profile a momentary snapshot, a

number of remarkable computational algorithms have been developed to pseudo-order cells

based on the different points they were assumed to occupy in a trajectory, thereby leveraging

biological asynchrony[29, 30]. For instance, Monocle[30, 31] constructs the minimum spanning tree, and

Wishbone[32] and Spring[33] construct the nearest neighbor graph from single-cell transcriptome

profiles. These tools have been widely used to depict neurogenesis[34], hematopoiesis[35, 36] and

reprogramming[37]. APEC integrates the Monocle algorithm into the accesson-based method and

enables pseudotime prediction from scATAC-seq data[38] and was applied to investigate HSC

differentiation linages (Figure 3a). Principal component analysis (PCA) of the accesson matrix

revealed multiple stages of the lineage during HSC differentiation (Figure 3b) and was consistent

with previous publications[3, 38]. After utilizing the Monocle package, APEC provided more precise

pathways from HSCs to the differentiated cell types (Figure 3c). In addition to the differentiation

pathways to MEP cells through the CMP state and to CLP cells through the LMPP state, MPP

cells may differentiate into GMP cells through two distinct trajectories: Path A through the CMP

state and Path B through the LMPP state, which is consistent with the composite model of HSC

and blood lineage commitment[39]. Notably, APEC suggested that CD34$^+$ plasmacytoid dendritic

cells (pDCs) from the bone marrow (Supplementary Figure 3) were derived from CLP cells on the

psuedotime trajectory (Figure 3c), which also agrees with a previous report[40]. Furthermore, APEC

is capable of evaluating the deviation of each TF along the single-cell trajectory to determine the

regulatory mechanisms during HSC differentiation. As expected, the HOX motif is highly enriched

in the accessible sites of HSCs/MPP cells, as are the GATA1, CEBPB and TCF4 motifs, which

exhibit gradients that increase along the erythroid, myeloid and lymphoid differentiation pathways,

respectively[38] (Figure 3d). In addition, we can see that the TF regulatory strategies of the two

paths from MMPs towards GMP cells were very different. Finally, we generated a hematopoiesis

tree based on the APEC analysis (Figure 3e).

APEC reveals the single-cell regulatory heterogeneity of thymocytes

T cells generated in the thymus play a critical role in the adaptive immune system, and

the development of thymocytes can be divided into 3 main stages based on the expression of the

surface markers CD4 and CD8, namely, CD4 CD8 double-negative (DN), CD4 CD8 double-

positive (DP) and CD4 or CD8 single-positive (CD4SP or CD8SP, respectively) stages[41]. However,

due to technical limitations, our genome-wide understanding of thymocyte development at single-

cell resolution remains unclear. Typically, more than 80% of thymocytes stay in the DP stage in

the thymus, whereas DN cells account for only approximately 3% of the thymocyte population. To

eliminate the impacts of great differences in proportion, we developed a fluorescent tagmentation-

and FACS-sorting-based scATAC-seq strategy (ftATAC-seq), which combined the advantages of

ATAC-see[42] and Pi-ATAC-seq[43] to manipulate the desired number of target cells by indexed

sorting (Figure 4a). Tn5 transposomes were fluorescently labeled in each cell to evaluate the

tagmentation efficiency so that cells with low ATAC signals could be gated out easily

(Supplementary Figure 4a, Figure 4b). With ftATAC-seq, we acquired high-quality chromatin

accessibility data for 352 index-sorted DN, DP, CD4SP, and CD8SP single cells and 352 mixed

thymocytes (Figure 4b). We applied APEC to mouse thymocyte ftATAC-seq data to investigate

the chromatin accessibility divergence during the developmental process and to reveal refined

regulome heterogeneity at single-cell resolution. Taking into account all 130685 peaks called from

the raw sequencing data, APEC aggregated 600 accessons and successfully assigned over 92%

of index-sorted DN, DP, CD4SP and CD8SP cells into the correct subpopulations (Figure 4c, 4d),

providing a much better classification than chromVAR (Supplementary Figure 4b, 4c), for which

this rate was only 56%. As expected, the majority of randomly sorted and mixed thymocytes were

classified into DP subtypes based on similarity hierarchical clustering, which was consistent with

the cellular subtype proportions in the thymus. APEC further classified all thymocytes into 14

subpopulations, including 2 DN, 7 DP, 1 CD4SP, 2 CD8SP, 1 coherence (Coh.A) and 1 transition

(Tran.A) state, suggesting that extensive epigenetic heterogeneity exists among cells with the

same CD4 and CD8 surface markers (Figure 4e). For instance, there are four main subtypes of

DN cells, according to the expression of the surface markers CD44 and CD25[44], while two clusters

were identified in ftATAC-seq. The accessibility signals around the *Il2ra* (Cd25) and *Cd44* gene

loci demonstrated that DN.A1 comprised CD44$^+$CD25$^-$ and CD44$^+$CD25$^+$ DN subtypes (DN1 and

DN2), and DN.A2 cells comprised CD44$^-$CD25$^+$ and CD44$^-$CD25$^-$ subtypes (DN3 and DN4),

suggesting significant chromatin changes between DN2 and DN3 cell development (Figure 4f).

Many TFs have been reported to be essential in regulating thymocyte development, and

we found that their motifs were remarkably enriched at different stages during the process (Figure

4g). For instance, Runx3 is well known for regulating CD8SP cells[45], and we observed significant

enrichment of the RUNX motif on DN cells and a group of CD8SP cells. Similarly, the TCF[46, 47],

RORC[48] and NFkB[49] family in regulating the corresponding stages during this process. More

enriched TF motifs in each cell subpopulation were also observed, suggesting significant

regulatory divergence in thymocytes (Supplementary Figure 4d). Interestingly, two clusters of

CD8SP cells appear to be differentially regulated based on motif analysis, in which CD8.A1 cells

are closer to DP cells, while CD8.A2 cells are more distant at the chromatin level, suggesting that

CD8.A2 cells are more mature CD8SP cells. In addition to the well-defined subtypes, APEC also

found a mixed cell population without specific features that was termed the coherence state

(Coh.A) and a transitional population between DP and SP cells (Tran.A).

APEC is capable of integrating single-cell transcriptional and epigenetic information by

scoring gene sets of interest based on their nearby peaks from scATAC-seq, thereby converting

the chromatin accessibility signals to values that are comparable to gene expression profiles

(online **Methods**). To test the performance of this integrative analysis approach and to evaluate

the accuracy of thymocyte classification by APEC, we assayed the transcriptomes of single

thymocytes and obtained 357 high-quality scRNA-seq profiles using the SMART-seq2 protocol[50].

Unsupervised analysis of gene expression profiles clustered these thymocytes into 13 groups in

Seurat[19] (Supplementary Figure 5a, 5b), and each subpopulation was identified based on known

feature genes (Supplementary Figure 5c, 5d). We then compared the adjusted scores obtained

from APEC with the single-cell RNA expression profile and observed a strong correlation between

the subtypes identified from the transcriptome and the subtypes identified from chromatin

accessibility (Figure 4h), confirming the reliability and stability of cellular classification using APEC.

APEC reconstructs the thymocyte developmental trajectory from ftATAC-seq profiles

APEC is capable of constructing a pseudotime trajectory and then predicting the cell differentiation lineage from a "snapshot" of single-cell epigenomes (Figure 3). We applied APEC to recapitulate the developmental trajectory and thereby reveal the single-cell regulatory dynamics during the maturation of thymocytes. Psuedotime analysis based on single-cell ftATAC-seq data shaped thymocytes into 5 developing stages (Figure 5a, Supplementary Figure 6a-b), where most of the cells in stages 1, 2, 4, and 5 were DN, DP, CD8SP and CD4SP cells, respectively. APEC also identified a transitional stage 3, which consisted of DP, coherence and transitional cells. Interestingly, the psuedotime trajectory suggests three developmental pathways for this process, one of which started with stage 1 (DN) and ended in stage 2 (DP), and the other two of which started with stage 1 (DN), went through a transitional stage 3 (a mixture of DP, Coh and Tran) and then bifurcated into stage 4 (CD8SP) and 5 (CD4SP). The predicted developmental trajectory could also be confirmed by the gene expression of surface markers, such as Cd4, Cd8, Runx3 and Ccr7 (Figure 5b). To evaluate the gene ontology (GO) enrichments over the entire process, we implemented an accesson-based GO module in APEC, which highlights the significance of the association between cells and biological function (Figure 5c). For instance, T cells selections, including β-selection, positive selection and negative selection, start from the DN3 stage. Consistent with this process, we observed a strong "T cell selection" GO term on the trajectory path after DN.A1. Since TCR signals are essential for T cell selection, we also observed the "T cell activation" GO term accompanied by "T cell selection". Meanwhile, the regulation of protein binding signal was also decreased at SP stages, indicating the necessity of weak TCR signal for the survive of SP T cells during negative selection.

To further uncover the regulatory mechanism underlying this developmental process, APEC was implemented to identify stage-specific enriched TFs along the trajectory and pinpoint the "pseudotime" at which the regulation occurs. In addition to the well-studied TFs mentioned above (Figure 4g, Supplemental Figure 4c), APEC also identified Zeb1[51], Ctcf[52] and Id4 as potential stage-specific regulators (Figure 5d). Interestingly, the Id4 motif enriched on DP cells was also reported to regulate apoptosis in other cell types[53, 54]. Associated with the fact that the vast majority of DP thymocytes die because of a failure of positive selection[55], we hypothesize that stage 2 may be the path towards DP cell apoptosis. We then checked the distribution of DP cells along the stage 2 trajectory and found that most DP.A1 cells were scattered in "early" stage 2, and they were enriched with GO terms such as "T cell selection", "cell activation" and "differentiation" (Figure 5e, Supplementary Figure 6c). However, most DP.A5-A6 cells were

distributed at the end of stage 2, and their principle accessons were enriched with GO terms such
as "apoptosis" and "chromatin modification". These results suggest that a majority of DP
thymocytes undergo T cell selection and enter an apoptosis state. Although it is believed that
more than 95% of DP thymocytes are subjected to death in positive selection, only a small
proportion of apoptotic cells could be detected in a snapshot of the thymus. By comparing the
number of cells near stage 3 with all the cells in stage 2, we estimated that ~3-5% of cells would
survive positive selection, which is consistent with previous publications[56, 57]. Our data suggest
that before entering an apoptotic stage, DP thymocytes that fail selection could have already
committed to apoptosis at the chromatin level.
285
286   **DISCUSSION**
287
Here, we introduced an accesson-based algorithm for single-cell chromatin accessibility
analysis. Without any prior information (such as motifs), this approach generated more refined
cell groups with reliable biological functions and properties. Integrating the new algorithm with all
necessary chromatin sequencing data processing tools, APEC provides a comprehensive
solution for transforming raw experimental single-cell data into final visualized results. In addition
to better clustering of subtle cell subtypes, APEC is also capable of locating potential specific
superenhancers, searching enriched motifs, estimating gene opening scores, and building time-
dependent cell developmental trajectories, and it is compatible with many existing single-cell
accessibility datasets. Despite these advantages, the biological implications of accessons are still
obscure, especially for those that involve only a small number of peaks; therefore, further
investigations may require uncovering the biology that underlies accessons.

To evaluate the performance of this approach in the context of the immune system, we
also adopted APEC with scATAC-seq technology to investigate the regulome dynamics of the
thymic development process. We developed a novel method of ftATAC-seq that captures Tn5-
tagged single cells of interest and outlines the chromatin accessibility heterogeneity and dynamics
during this process. Coordinated with essential cell surface markers, APEC provided a much more
in-depth classification of thymocytes than the conventional DN, DP, CD4SP and CD8SP stages
based on single-cell chromatin status. By reconstructing the developmental pseudotime trajectory,
APEC discovered a transitional stage before thymocytes bifurcate into CD4SP and CD8SP cells
and inferred that one of the stages leads to cell apoptosis. APEC analysis suggested that DP cells

were gradually programmed to undergo apoptosis at the chromatin level; however, further studies
are needed to fully understand the regulatory mechanism of this process.

Acknowledgments

This work was supported by the National Key R&D Program of China (2017YFA0102903 to K.Q.)
and by National Natural Science Foundation of China grants 91640113 (to K.Q.) and
31771428 (to K.Q.). It was also supported by Anhui Provincial Natural Science Foundation grant
BJ2070000097 (to B.L.). We thank the Howard Chang lab at Stanford University for helpful
discussion. We thank the USTC supercomputing center and the School of Life Science
Bioinformatics Center for providing supercomputing resources for this project.

Authors' contributions

KQ, BL and YL conceived the project, BL developed the APEC software and performed all data
analysis with helps from QY, JF, PC, and CJ. YL developed ftATAC-seq technique and performed
all scATAC-seq and scRNA-seq experiments with helps from LZ. KL analyzed scRNA-seq data.
BL, YL and KQ wrote the manuscript with inputs from all other authors.

Data and code availability

Mouse thymocytes ftATAC-seq data can be obtained from the Genome Sequence Archive at BIG
Data Center with the accession number CRA001267 and is available via
http://bigd.big.ac.cn/gsa/s/yp1164Et. Other published data sets used in this study are available
from NIH GEO with accession numbers GSE74310[3], GSE65360[13], GSE96772[38], and
GSE100033[15]. APEC pipeline can be downloaded from the GitHub website
(https://github.com/QuKunLab/APEC).

REFERENCES

1. Buenrostro, J.D., Giresi, P.G., Zaba, L.C., Chang, H.Y. & Greenleaf, W.J. Transposition of native chromatin for fast and sensitive epigenomic profiling of open chromatin, DNA-binding proteins and nucleosome position. *Nat Methods* **10**, 1213-1218 (2013).

2. Ye, M. et al. Hematopoietic Differentiation Is Required for Initiation of Acute Myeloid Leukemia. *Cell Stem Cell* **17**, 611-623 (2015).

3. Corces, M.R. et al. Lineage-specific and single-cell chromatin accessibility charts human hematopoiesis and leukemia evolution. *Nat Genet* **48**, 1193-1203 (2016).

4. Yu, V.W.C. et al. Epigenetic Memory Underlies Cell-Autonomous Heterogeneous Behavior of Hematopoietic Stem Cells. *Cell* **167**, 1310-1322 e1317 (2016).

5. Wu, J. et al. The landscape of accessible chromatin in mammalian preimplantation embryos. *Nature* **534**, 652-657 (2016).

6. Wu, J. et al. Chromatin analysis in human early development reveals epigenetic transition during ZGA. *Nature* **557**, 256-260 (2018).

7. Jorstad, N.L. et al. Stimulation of functional neuronal regeneration from Muller glia in adult mice. *Nature* **548**, 103-107 (2017).

8. Su, Y. et al. Neuronal activity modifies the chromatin accessibility landscape in the adult brain. *Nat Neurosci* **20**, 476-483 (2017).

9. Xiang, Y. et al. Fusion of Regionally Specified hPSC-Derived Organoids Models Human Brain Development and Interneuron Migration. *Cell Stem Cell* **21**, 383-398 e387 (2017).

10. de la Torre-Ubieta, L. et al. The Dynamic Landscape of Open Chromatin during Human Cortical Neurogenesis. *Cell* **172**, 289-304 e218 (2018).

11. Denny, S.K. et al. Nfib Promotes Metastasis through a Widespread Increase in Chromatin Accessibility. *Cell* **166**, 328-342 (2016).

12. Qu, K. et al. Chromatin Accessibility Landscape of Cutaneous T Cell Lymphoma and Dynamic Response to HDAC Inhibitors. *Cancer Cell* **32**, 27-41 e24 (2017).

13. Buenrostro, J.D. et al. Single-cell chromatin accessibility reveals principles of regulatory variation. *Nature* **523**, 486-490 (2015).

14. Lake, B.B. et al. Integrative single-cell analysis of transcriptional and epigenetic states in the human adult brain. *Nat Biotechnol* **36**, 70-80 (2018).

15. Preissl, S. et al. Single-nucleus analysis of accessible chromatin in developing mouse forebrain reveals cell-type-specific transcriptional regulation. *Nat Neurosci* **21**, 432-439 (2018).

16. Cusanovich, D.A. et al. Multiplex single cell profiling of chromatin accessibility by combinatorial cellular indexing. *Science* **348**, 910-914 (2015).

17. Cusanovich, D.A. et al. The cis-regulatory dynamics of embryonic development at single-cell resolution. *Nature* **555**, 538-542 (2018).

18. Bacher, R. et al. SCnorm: robust normalization of single-cell RNA-seq data. *Nat Methods* **14**, 584-586 (2017).

19. Butler, A., Hoffman, P., Smibert, P., Papalexi, E. & Satija, R. Integrating single-cell transcriptomic data across different conditions, technologies, and species. *Nat Biotechnol* **36**, 411-420 (2018).

20. Kiselev, V.Y. et al. SC3: consensus clustering of single-cell RNA-seq data. *Nat Methods* **14**, 483-486 (2017).

21. Wang, B., Zhu, J., Pierson, E., Ramazzotti, D. & Batzoglou, S. Visualization and analysis of single-cell RNA-seq data by kernel-based similarity learning. *Nat Methods* **14**, 414-416 (2017).

22. Iacono, G. et al. bigSCale: an analytical framework for big-scale single-cell data. *Genome Res* **28**, 878-890 (2018).

23.   Wolf, F.A., Angerer, P. & Theis, F.J. SCANPY: large-scale single-cell gene expression data analysis.
Genome Biol **19**, 15 (2018).

24.   Schep, A.N., Wu, B., Buenrostro, J.D. & Greenleaf, W.J. chromVAR: inferring transcription-factor-
associated accessibility from single-cell epigenomic data. *Nat Methods* **14**, 975-978 (2017).

25.   Zamanighomi, M. et al. Unsupervised clustering and epigenetic classification of single cells. *Nat*
Commun **9**, 2410 (2018).

26.   Oberst, A. et al. The Nedd4-binding partner 1 (N4BP1) protein is an inhibitor of the E3 ligase Itch.
Proc Natl Acad Sci U S A **104**, 11280-11285 (2007).

27.   Eguchi, M. et al. GPHN, a novel partner gene fused to MLL in a leukemia with t(11;14)(q23;q24).
Genes Chromosomes Cancer **32**, 212-221 (2001).

28.   Satpathy, A.T. et al. Transcript-indexed ATAC-seq for precision immune profiling. *Nat Med* **24**,
580-590 (2018).

29.   Bendall, S.C. et al. Single-cell trajectory detection uncovers progression and regulatory
coordination in human B cell development. *Cell* **157**, 714-725 (2014).

30.   Trapnell, C. et al. The dynamics and regulators of cell fate decisions are revealed by
pseudotemporal ordering of single cells. *Nat Biotechnol* **32**, 381-386 (2014).

31.   Qiu, X. et al. Reversed graph embedding resolves complex single-cell trajectories. *Nat Methods*
14, 979-982 (2017).

32.   Setty, M. et al. Wishbone identifies bifurcating developmental trajectories from single-cell data.
Nat Biotechnol **34**, 637-645 (2016).

33.   Weinreb, C., Wolock, S. & Klein, A.M. SPRING: a kinetic interface for visualizing high dimensional
single-cell expression data. *Bioinformatics* **34**, 1246-1248 (2018).

34.   Habib, N. et al. Div-Seq: Single-nucleus RNA-Seq reveals dynamics of rare adult newborn neurons.
Science **353**, 925-928 (2016).

35.   Olsson, A. et al. Single-cell analysis of mixed-lineage states leading to a binary cell fate choice.
Nature **537**, 698-+ (2016).

36.   Zhou, F. et al. Tracing haematopoietic stem cell formation at single-cell resolution. *Nature* **533**,
487-+ (2016).

37.   Treutlein, B. et al. Dissecting direct reprogramming from fibroblast to neuron using single-cell
RNA-seq. *Nature* **534**, 391-+ (2016).

38.   Buenrostro, J.D. et al. Integrated Single-Cell Analysis Maps the Continuous Regulatory Landscape
of Human Hematopoietic Differentiation. *Cell* **173**, 1535-1548 e1516 (2018).

39.   Adolfsson, J. et al. Identification of Flt3+ lympho-myeloid stem cells lacking erythro-
megakaryocytic potential a revised road map for adult blood lineage commitment. *Cell* **121**, 295-
306 (2005).

40.   Chistiakov, D.A., Orekhov, A.N., Sobenin, I.A. & Bobryshev, Y.V. Plasmacytoid dendritic cells:
development, functions, and role in atherosclerotic inflammation. *Front Physiol* **5**, 279 (2014).

41.   Germain, R.N. T-cell development and the CD4-CD8 lineage decision. *Nat Rev Immunol* **2**, 309-322
(2002).

42.   Chen, X. et al. ATAC-see reveals the accessible genome by transposase-mediated imaging and
sequencing. *Nat Methods* **13**, 1013-1020 (2016).

43.   Chen, X. et al. Joint single-cell DNA accessibility and protein epitope profiling reveals
environmental regulation of epigenomic heterogeneity. *Nat Commun* **9**, 4590 (2018).

44.   Godfrey, D.I., Kennedy, J., Suda, T. & Zlotnik, A. A developmental pathway involving four
phenotypically and functionally distinct subsets of CD3-CD4-CD8- triple-negative adult mouse
thymocytes defined by CD44 and CD25 expression. *J Immunol* **150**, 4244-4252 (1993).

45.   Taniuchi, I. et al. Differential requirements for Runx proteins in CD4 repression and epigenetic
silencing during T lymphocyte development. *Cell* **111**, 621-633 (2002).

46. Ioannidis, V., Beermann, F., Clevers, H. & Held, W. The beta-catenin--TCF-1 pathway ensures
CD4(+)CD8(+) thymocyte survival. *Nat Immunol* **2**, 691-697 (2001).
47. Yu, S. et al. The TCF-1 and LEF-1 transcription factors have cooperative and opposing roles in T cell
development and malignancy. *Immunity* **37**, 813-826 (2012).
48. Sun, Z. et al. Requirement for RORgamma in thymocyte survival and lymphoid organ development.
Science **288**, 2369-2373 (2000).
49. Gerondakis, S., Fulford, T.S., Messina, N.L. & Grumont, R.J. NF-kappaB control of T cell
development. *Nat Immunol* **15**, 15-25 (2014).
50. Picelli, S. et al. Smart-seq2 for sensitive full-length transcriptome profiling in single cells. *Nat
Methods* **10**, 1096-1098 (2013).
51. Higashi, Y. et al. Impairment of T cell development in deltaEF1 mutant mice. *J Exp Med* **185**, 1467-
1479 (1997).
52. Heath, H. et al. CTCF regulates cell cycle progression of alphabeta T cells in the thymus. *EMBO J*
27, 2839-2850 (2008).
53. Andres-Barquin, P.J., Hernandez, M.C. & Israel, M.A. Id4 expression induces apoptosis in astrocytic
cultures and is down-regulated by activation of the cAMP-dependent signal transduction pathway.
Exp Cell Res **247**, 347-355 (1999).
54. Carey, J.P., Knowell, A.E., Chinaranagari, S. & Chaudhary, J. Id4 promotes senescence and
sensitivity to doxorubicin-induced apoptosis in DU145 prostate cancer cells. *Anticancer Res* **33**,
4271-4278 (2013).
55. Surh, C.D. & Sprent, J. T-cell apoptosis detected in situ during positive and negative selection in
the thymus. *Nature* **372**, 100-103 (1994).
56. Huesmann, M., Scott, B., Kisielow, P. & von Boehmer, H. Kinetics and efficacy of positive selection
in the thymus of normal and T cell receptor transgenic mice. *Cell* **66**, 533-540 (1991).
57. Shortman, K., Vremec, D. & Egerton, M. The kinetics of T cell antigen receptor expression by
subgroups of CD4+8+ thymocytes: delineation of CD4+8+3(2+) thymocytes as post-selection
intermediates leading to mature T cells. *J Exp Med* **173**, 323-332 (1991).
455

FIGURES

Figure 1. The accesson matrix constructed from the sparse fragment count matrix improved the
clustering of scATAC-seq data. (**a**) Step-by-step workflow of APEC. Peaks were grouped into
accessons by their accessibility pattern among cells with the K nearest neighbors (KNN) method.
(**b, c**) Hierarchical clustering of cell-cell correlations based on the accesson matrix (from APEC)
and the motif matrix (from chromVAR). The scATAC-seq data include leukemic stem cells (LSCs),
leukemia blast cells, lymphoid-primed multipotential progenitors (LMPPs), HL60 cells, and
monocytes. P1, acute myeloid leukemia (AML) patient 1 (SU070); P2, AML patient 2 (SU353).
The cells are labeled by their fluorescence indices. (**d, e**) t-Distributed Stochastic Neighbor
Embedding (tSNE) diagrams based on the accesson matrix and the motif matrix. (**f**) Fragment
counts were specifically enriched in the superenhancer region upstream of N4BP1 in P1-LSCs.

Figure 2. APEC improved the cell type classification of adult mouse forebrain snATAC-seq data. (**a**) A tSNE diagram demonstrates the KNN clustering of forebrain cells. (**b**) Hierarchical clustering of the cell-cell correlation matrix. The side bar denotes cell clusters from the KNN method. (**c**) Average of the marker gene scores for each cell cluster, normalized by the standard score (z-score). The top row lists the cell numbers for each cluster. (**d**) Hierarchical clustering of the cluster-cluster correlation matrix. (**e**) Differential enrichments of cell type-specific motifs in different clusters. (**f, g**) Intensity of representative accessons associated with signature genes of excitatory (Ex) and inhibitory (In) neuron subtypes. The subtypes listed in parentheses were defined by the signature genes in the results from scRNA-seq data[14].

Figure 3. APEC constructed a differentiation pathway from scATAC-seq data from human hematopoietic cells. (**a**) The pseudotime trajectory construction scheme based on the accesson matrix and Monocle. (**b**) Principal component analysis (PCA) of the accesson matrix for human hematopoietic cells. The first principal component is not shown here because it was highly correlated with sequencing depth[38]. HSC, hematopoietic stem cell; MPP, multipotent progenitor; LMPP, lymphoid-primed multipotential progenitor; CMP, common myeloid progenitor; CLP, common lymphoid progenitor; pDC, plasmacytoid dendritic cell; GMP, granulocyte-macrophage

487   progenitor; MEP, megakaryocyte-erythroid progenitor; unknown, unlabeled cells. (**c**) Pseudotime

488   trajectory for the same data constructed by calling Monocle on the accesson matrix. Paths A and

489   B represent different pathways for GMP cell differentiation. (**d**) The deviations of significant

490   differential motifs (HOXA9, GATA1, CEBPB, and TCF4) plotted on the pseudotime trajectory. (**e**)

491   Modified schematic of human hematopoietic differentiation.

accesson-based KNN method clustered thymocytes into 14 subtypes. DN.A1&A2, double-negative clusters; DP.A1~A7, double-positive clusters; Coh.A, coherent state; Tran.A, transition state; CD8.A1&A2, CD8$^+$ single-positive clusters; CD4.A, CD4$^+$ single-positive cluster. (**f**) Average fragment counts of two DN clusters around the marker genes Cd44 and Il2ra. (**g**) Differential enrichment of the motifs Runx, Tcf, Rorc, and Nfkb in the cell clusters. (**h**) Correlation of the cell clusters identified by data from single-cell transcriptome (SMART-seq) and chromatin accessibility (ftATAC-seq) analysis.

Figure 5. APEC depicted the developmental pathways of *Mus musculus* thymocytes by pseudotime analysis. (**a**) Pseudotime trajectory based on the accesson matrix of thymocyte ftATAC-seq data. Cell colors were defined by the developmental stages along pseudotime. Pie charts show the proportion of cell clusters at each stage. (**b**) Normalized scores of important marker genes (Cd8a, Cd4, Runx3, and Ccr7) along each branch of the pseudotime trajectory. (**c**) Accesson weight scores of important functional GO terms along each branch of the pseudotime trajectory. (**d**) Enrichment of specific motifs searched from the differential accessons of each cell subtype. (**e**) On the stage 2 branch, the cell number distribution of clusters DP.A1~A7 along pseudotime (upper panel) and the intensity of marker accessons of DP.A1 and DP.A5/6 (lower panel).

METHODS

Mice. C57BL/6 mice were purchased from Beijing Vital River Laboratory Animal Technology and maintained under specific pathogen-free conditions until the time of experiments. All mouse experiments in this study were reviewed and approved by the Institutional Animal Care and Use Committee of the University of Science and Technology of China.

ftATAC-seq on mouse thymocytes. Alexa fluor 488-labeled adaptor oligonucleotides were synthesized at Sangon Biotech as follows: Tn5ME, 5'-[phos]CTGTCTCTTATACACATCT-3'; AF488-R1, 5'-AF488- TCGTCGGCAGCGTCAGATGTGTATAAGAGACAG-3'; and AF488-R2, 5'-AF488-GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAG-3'. Then, 50 µM of AF488-R1/Tn5ME and AF488-R2/Tn5ME were denatured separately in TE buffer (Qiagen) at 95 °C for 5 min and cooled down to 22 °C at 0.1 °C/s. AF488-labeled adaptors were assembled onto Robust Tn5 transposase (Robustnique) according to the user manual to form fluorescent transposomes.

Thymus tissues isolated from 6- to 8-week-old male mice were gently ground in 1 mL of RPMI-1640. Thymocytes in a single-cell suspension were counted after passing through a 40 µm nylon mesh. A total of 1×10^6 thymocytes were stained with PerCP-Cy5.5-anti-CD45, PE-anti-CD8a and APC-Cy7-anti-CD4 antibodies (Biolegend) and then fixed in 1× PBS containing 1% methanol at room temperature for 5 min. After washing twice with 1× PBS, the cells were counted again. A total of 1×10^5 fixed cells were resuspended in 40 µL of 1× TD buffer (5 mM Tris-HCl , pH 8.0, 5 mM $MgCl_2$, and 10% DMF) containing 0.1% NP-40. Then, 10 µL of fluorescent transposomes were added and mixed gently. Fluorescent tagmentation was conducted at 55 °C for 30 min and stopped by adding 200 µL of 100 mM EDTA directly to the reaction mixture. The cells were loaded on a Sony SH800S sorter, and single cells of the CD45$^+$/AF488-Tn5^{hi} population were index-sorted into each well of 384-well plates. The 384-well plates used to acquire sorted cells were loaded with 2 µL of release buffer (50 mM EDTA, 0.02% SDS) before use. After sorting, the cells in the wells were incubated for 1 min. Plates that were not processed immediately were preserved at -80 °C.

To prepare a single-cell ATAC-seq library, plates containing fluorescently tagmented cells were incubated at 55 °C for 30 min. Then, 4.2 µL of PCR round 1 buffer (1 µL of 100 µM $MgCl_2$, 3 µL of 2× I-5 PCR mix [MCLAB], and 0.1 µL each of 10 µM R1 and R2 primers) were added to each well, followed by PCR: 72 °C for 10 min; 98 °C for 3 min; 10 cycles of 98 °C for 10 s, 63 °C for 30 s and 72 °C for 1 min; 72 °C for 3 min; and holding at 4 °C. Thereafter, each well received 4 µL of PCR round 2 buffer (2 µL of I-5 PCR Mix, 0.5 µL each of Ad1 and barcoded Ad2 primers, and 1 µL of ddH$_2$O), and final PCR amplification was carried out: 98 °C for 3 min; 12 cycles of 98 °C

for 10 s, 63 °C for 30 s and 72 °C for 1 min; 72 °C for 3 min; and holding at 4 °C. Wells containing

different Ad2 barcodes were collected together and purified with a QIAquick PCR purification kit

(Qiagen). Libraries were sequenced on an Illumina HiSeq X Ten system.

SMART-seq on thymocytes. Thymocytes were stained and sorted directly into 384-well plates

without fixation. SMART-seq was performed as described with some modifications.[58] Reverse

transcription and the template-switch reaction were performed at 50 °C for 1 hr with Maxima H

Minus Reverse Transcriptase (Thermo Fisher); for library construction, 0.5-1 ng of cDNA was

fragmented with 0.05 μL of Robust Tn5 transposome in 20 μL of TD buffer at 55 °C for 10 min,

then purified with 0.8× VAHTS DNA Clean Beads (Vazyme Biotech), followed by PCR

amplification with Ad1 and barcoded Ad2 primers and purification with 0.6× VAHTS DNA Clean

Beads. Libraries were sequenced on an Illumina HiSeq X Ten system.

Data source. All experimental raw data used in this paper are available online. The single-cell

data for mouse thymocytes captured by the ftATAC-seq experiment can be obtained from the

Genome Sequence Archive at BIG Data Center with the accession number CRA001267 and is

available via http://bigd.big.ac.cn/gsa/s/yp1164Et. Other published data sets used in this study

are available from NIH GEO: (1) scATAC-seq data for LSCs and leukemic blast cells from patients

SU070 and SU353, LMPP cells, and monocytes from GSE74310[3]; (2) scATAC-seq data for HL-

60 cells from GSE65360[13]; and (3) scATAC-seq data for hematopoietic development (HSCs,

MPPs, CMPs, LMPPs, GMPs, EMPs, CLPs and pDCs) from GSE96772[38]. APEC is also

compatible with a preprocessed fragment count matrix from the snATAC-seq data for the

forebrain of adult mice (p56) from GSE100033[15].

Preparing the fragment count matrix from the raw data. APEC adopted the general mapping,

alignment, peak calling and motif searching procedures to process the scATAC-seq data. To trim

the adapters in the raw data (in paired-end fastq format files for each single-cell sample), we

implanted the python version trimming code from our previous published pipeline (ATAC-pipe)[59].

Then, APEC used BOWTIE2 to map the trimmed sequencing data to the corresponding genome

index and used PICARD for the sorting, duplicate removal, and fragment length counting of the

aligned data. The pipeline called peaks from the merged file of all cells by MACS2, ranked and

filtered out the low quality peaks based on the false discovery rate (Q-value). Genomic locations

of the peaks were annotated by HOMER, and motifs searched by FIMO. APEC calculates the

number of fragments and the percent of reads mapped to the TSS region (±2000 BP) for each

cell, and filters out high quality cells for downstream analysis. All required files for the hg19 and

mm10 assembly have been integrated into the pipeline. If users want to process data from other

species, they can also download corresponding reference files from the UCSC website. By
combining existing tools, APEC made it possible to finish all of the above data processing steps
by one command line, and generate a fragment count matrix for subsequent cell clustering and
differential analysis.

Accesson-based clustering algorithm. We define accesson as a set of peaks with similar
accessibility patterns across all single cells, similar to the definition of gene modules for RNA-seq
data. The peaks of a same accesson can be distant from each other on the genome, and
sometimes on multiple chromosomes. After preprocessing, a filtered fragment count matrix **M** is
obtained, and APEC groups peaks to construct accessons and then performs cell clustering
analysis as follows:

(1) Normalization of the fragment count matrix. Each matrix element M_{ij} represents the
number of raw reads in cell i and peak j, and element M_{ij} was then normalized by the
total number of reads in each cell i, as if there are 10,000 reads in each cell.

$$M'_{ij} = log_2\left(\frac{M_{ij} \times 10000}{\sum_{j'} M_{ij'}} + 1\right)$$

(2) Constructing accessons. The top 40 principal components of the normalized matrix **M'**
were used to construct the connectivity matrix (**C$_{peak}$**) of peaks by the K-nearest-neighbor
(KNN) method. Based on the matrix **C$_{peak}$**, all peaks were grouped by agglomerative
clustering with the Ward's method, and the sum of one peak group was an accesson. In
processing of all datasets in this study, the default number of accessons was set to 600.
We recommend using a flexible number of accessons so that you can accumulate enough
peaks in one accesson while avoiding incorrect grouping of differential peaks. However,
not all accessons were used for cell clustering in the next step. Sparse accessons with 4
or fewer peaks were discarded since they will interfere with the clustering accuracy. Only
accessons containing 5 or more peaks were retained in the accesson count matrix **M$_a$**.
Each row of **M$_a$** is an accesson, each column is a cell and the elements of **M$_a$** represents
the cumulative read counts of each accesson in each cell. If less than 30% of the
accessons contain enough number of peaks, the users may consider to reduce the default
accesson number to avoid sparse accessons.
(3) Cell clustering. From the accesson matrix **M$_a$**, APEC calculated the Pearson correlation
between each pair of cells, and then performed both hierarchical and KNN clustering on
the correlation matrix to categorize cells into different clusters. The number of cell clusters
can be predicted by the Louvain method, or inputted by the users. By default, cell

clustering was performed in the high-dimensional PCA transformed space, but also
supports clustering in the tSNE space.

(4) Comparison with other clustering methods. To investigate the accuracy of clusters
generated by different algorithms, APEC provides two ways to compare cell clusters: a)
The contingency matrix, in which each element represents the number of common cells
between two clusters from different methods (e.g., hierarchy and KNN clustering, or
accesson-based and motif-based clustering); b) The ARI value, which evaluates the
similarity of clustering results from two different algorithms[20]. Moreover, one clustering
method can be compared with known cell types in the original single-cell data (such as
the FACS index) to confirm the accuracy of the cell type classification algorithm.

Gene score evaluated by peaks around the TSS. To evaluate the accessibility score of one
gene, we calculated the average count of all peaks around its TSS (±20000 BP) as its raw score
(S_{ij} for cell i and gene j). Then, we obtained the gene accessibility score by normalizing the raw
score ($S'_{ij} = S_{ij} * 10000/\sum_i S_{ij}$), which is in a range comparable to the gene expression from
scRNA-seq data. The average score of all cells in one cluster represents the accessibility of a cell
type ($\bar{S}_{kj}$ for cell cluster k and gene j). We normalized the gene score matrix $\bar{S}$ by caculating z-
score for each row and column, and the final matrix $\bar{S}_z$ represents the relative strength of gene
accessibility for each cell type.

Significant differential peaks, genes and motifs. APEC used the Student's t-test to estimate
the significance of the fragment count differences between cell clusters, with P-value and fold
changes, and one can determine the thresholds to identify significant differential peaks for each
cluster. The significant differential genes of each cell cluster can also be acquired from the
accessibility score ($\bar{S}_{kj}$) by the same method. To accurately quantify the enrichment of motifs on
each cell, APEC applied the bias-corrected deviation algorithm from chromVAR[24]; thus, the
chromVAR algorithm has been embedded into the pipeline to facilitate the calculation of the
corrected deviation of the motifs. In this python version of chromVAR, permuted sampling and
background deviation calculation can be run in parallel on multiple processors to reduce the
computer time. The differentially enriched motifs were defined by a fold change >1 in the average
motif deviation between one cluster and another.

Potential super-enhancers. Here, we defined a super-enhancer as a long continuous genomic
area containing many accessible regions and have the same accessibility pattern in different cells.
Many different motifs appear in one super-enhancer, therefore, the motif-based clustering method

cannot reflect the critical contributions from super-enhancers for cell clustering. However, the
accesson-based algorithm can group most peaks in one super-enhancer to one accesson since
they always present the same accessibility pattern between cells. APEC identified super-
enhancers by counting the number of peaks in a 1 million BP genomic area that belong to a same
accesson. It also requires that more than 3/4 of the putative peaks in one super-enhancer be
adjacent on the initial peak list. The pipeline can also aggregate bam files by cell types/clusters
and convert them to BigWig format for users to upload to the UCSC genome browser for
visulization.

Pseudotime trajectory. As a tool to simulate the time-dependent variation of gene expression
and the cell development pathway, Monocle has been widely used for the analysis of single-cell
RNA-seq experiments[30, 60]. APEC reduced the dimension of the accesson count matrix $\mathbf{M_a}$ by
PCA, and then performed pseudotime analysis using the Monocle program. For complex datasets,
it is necessary to limit the number of principal components, since too many features will cause
too many branches on the pseudo-time trajectory, and makes it difficult for a user to identify the
biological significance of each branch. For the hematopoietic single cell data and thymocyte data,
we used the top 5 principal components of the accesson matrix to construct the developmental
and differentiation trajectories.

Parameter settings for each analysis. In the quality control (QC) step, cells are filtered by two
constraints: the percentage of the fragments in peaks (P_f) and the total number of valid fragments
(N_f). However, there is no fixed cutoff for these two parameters since the quality of different cell
types and/or experiment batches are completely different. The total number of peaks is usually
limited to approximately 50000 to reduce computer time, but we recommend using all peaks if the
users want to obtain better cell clusters. (1) For the scATAC-seq data from leukemic cells (P1/P2
LSCs and blast cells, LMPPs, HL60 cells, and monocytes), the threshold of -log(Q-value) was set
to 8 to retain 42139 high-quality peaks for subsequent processing. In the QC step, we set the P_f
cutoff to 0.05 and the N_f cutoff to 800. (2) For the snATAC-seq data from the adult mouse forebrain,
all peaks and the raw count matrix obtained from the original data source were adopted in the
analysis. (3) For the data set from hematopoietic cells, the -log(Q-value) threshold of high-quality
peaks was set to 35 to retain 54212 peaks, and the cutoff values of P_f and N_f were 0.1 and 1000,
respectively. (4) For the ftATAC-seq data from thymocytes, all 130685 peaks called by MACS2
were reserved for the fragment count matrix (Q-value<0.05), and we retained cells with P_f >0.2
and N_f >2000.

SMART-seq data analysis with Seurat. For the analysis of SMART-seq data from mouse
thymocytes, we employed STAR (version 2.5.2a) with the ratio of mismatches to mapped length
(outFilterMismatchNoverLmax) less than or equal to 0.05, translated output alignments into
transcript coordinates (i.e., quantMode TranscriptomeSAM) for mapping[61] (Dobin et al., 2013) and
used RSEM[62] (Bo et al., 2011) to calculate the TPM of genes. For QC, we excluded cells in which
fewer than 2000 genes were detected and genes that were expressed in only 3 or fewer cells.
Seurat filtered cells with several specific parameters to limit the number of genes detected in each
cell to 2000~6000 and the proportion of mitochondrial genes in each cell was set to less than 0.4
(i.e., low.thresholds=c(2000,-Inf), high.thresholds=c(6000,0.4)). Additionally, the top 12 principal
components were used for dimension reduction with a resolution of 3.2 (dims.use =1:12,
resolution=3.2), followed by cell clustering and differential expressed gene analysis[63].

Association of cell clusters from scATAC-seq and scRNA-seq data. To determine the
association between cell clusters from epigenomics and transcriptome sequencing, we calculated
the P-values of Fisher's exact test of marker/nonmarker genes between each pair of cell clusters
from scATAC-seq and scRNA-seq data. For example, for cell cluster i from ftATAC-seq and cell
cluster j from SMART-seq, if the number of consensus marker genes in both cluster i and j is G_{11},
the number of genes that are not markers in either cluster i or j is G_{22}, and the number of markers
in either cluster i (or cluster j) is G_{12} (or G_{21}), then the 2 by 2 matrix **G** can be directly used for
Fisher's exact test to evaluate the P-valuie A_{ij} between cluster i and j. We calculated the logarithm
of matrix **A** to obtain matrix **A'**, then calculated the z-score for each row and column of **A'** to
determine the correlation of cell clusters from different experiments.

Biological function of accesson. We defined the functional characteristics of each accesson by
the GO terms and motifs enriched on its peaks. The GO terms of an accesson were obtained by
submitting all of its peaks to the GREAT website[64]. The logarithm of the P-value of each GO term
in each accesson was filled into a (GO terms) × (accessons) matrix **L**. The significance of each
GO term on each cell was evaluated by the product of the matrix **L** and the accesson reads count
matrix $\mathbf{M_a}$. Then we calculated the z-score for each row of this product matrix, and plotted the z-
score as the GO-term score on the trajectory diagram. To assess the motif enrichment of the
accessons, we used the Centrimo tool of MEME suite[65] to search for the enriched motifs for the
peaks of each accesson, and applied the same algorithm as the GO term score to obtain the motif
score.

REFERENCES

58. Picelli, S. et al. Full-length RNA-seq from single cells using Smart-seq2. *Nat Protoc* **9**, 171-181 (2014).

59. Zuo, Z. et al. ATAC-pipe: general analysis of genome-wide chromatin accessibility. *Briefings in Bioinformatics*, bby056-bby056 (2018).

60. Qiu, X. et al. Single-cell mRNA quantification and differential analysis with Census. *Nat Methods* **14**, 309-315 (2017).

61. Dobin, A. et al. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21 (2013).

62. Li, B. & Dewey, C.N. RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. *Bmc Bioinformatics* **12**, 323 (2011).

63. Macosko, E.Z. et al. Highly Parallel Genome-wide Expression Profiling of Individual Cells Using Nanoliter Droplets. *Cell* **161**, 1202-1214 (2015).

64. McLean, C.Y. et al. GREAT improves functional interpretation of cis-regulatory regions. *Nat Biotechnol* **28**, 495-501 (2010).

65. Bailey, T.L. et al. MEME SUITE: tools for motif discovery and searching. *Nucleic Acids Res* **37**, W202-208 (2009).