APEC: An accesson-based method for single-cell chromatin
accessibility analysis

Bin Li[1,3], Young Li[1,3], Kun Li[1], Lianbang Zhu[1], Qiaoni Yu[1], Pengfei Cai[1], Jingwen Fang[1], Wen Zhang[1],
Pengcheng Du[1], Chen Jiang[1], Kun Qu[1,2,]*

Affiliation

[1]Division of Molecular Medicine, Hefei National Laboratory for Physical Sciences at Microscale, The CAS
Key Laboratory of Innate Immunity and Chronic Disease, School of Life Sciences, University of Science
and Technology of China

[2]CAS Center for Excellence in Molecular Cell Sciences, University of Science and Technology of China

[3]Cofirst authors

*Correspondence: Kun Qu (qukun@ustc.edu.cn)

Contact Information

Kun Qu, Ph.D.

Division of Molecular Medicine, Hefei National Laboratory for Physical Sciences at Microscale, The CAS
Key Laboratory of Innate Immunity and Chronic Disease, School of Life Sciences, University of Science
and Technology of China

Hefei, Anhui, China, 230027

Email: qukun@ustc.edu.cn

Phone: +86-551-63606257

Abstract:

The development of sequencing technologies has promoted the survey of genome-wide chromatin accessibility at single-cell resolution; however, comprehensive analysis of single-cell epigenomic profiles remains a challenge. Here, we introduce an accessibility pattern-based epigenomic clustering (APEC) method, which classifies each individual cell by groups of accessible regions with synergistic signal patterns termed "accessons". By integrating with other analytical tools, this python-based APEC package greatly improves the accuracy of unsupervised single-cell clustering for many different public data sets. APEC also identifies significant differentially accessible sites, predicts enriched motifs, and projects pseudotime trajectories. Furthermore, we developed a fluorescent tagmentation- and FACS-sorting-based single-cell ATAC-seq technique named ftATAC-seq and investigated the per cell regulome dynamics of mouse thymocytes. Associated with ftATAC-seq, APEC revealed a detailed epigenomic heterogeneity of thymocytes, characterized the developmental trajectory and predicted the regulators that control the stages of maturation process. Overall, this work illustrates a powerful approach to study single-cell epigenomic heterogeneity and regulome dynamics.

INTRODUCTION

As a technique for probing genome-wide chromatin accessibility in a small number of cells *in vivo*, the assay for transposase-accessible chromatin with high-throughput sequencing (ATAC-seq) has been widely applied to investigate the cellular regulomes of many important biological processes[1], such as hematopoietic stem cell (HSC) differentiation[2], embryonic development[3], neuronal activity and regeneration[4, 5], tumor cell metastasis[6], and patient responses to anticancer drug treatment[7]. Recently, several experimental schemes have been developed to capture chromatin accessibility at single-cell/nucleus resolution, i.e., single-cell ATAC-seq (scATAC-seq)[8], single-nucleus ATAC-seq (snATAC-seq)[9, 10], and single-cell combinatorial indexing ATAC-seq (sci-ATAC-seq)[11, 12], which significantly extended researchers' ability to uncover cell-to-cell epigenetic variation and other fundamental mechanisms that generate heterogeneity from identical DNA sequences. By contrast, the in-depth analysis of single-cell chromatin accessibility profiles for this purpose remains a challenge. Numerous efficient algorithms have been developed to accurately normalize, cluster and visualize cells from single-cell transcriptome sequencing profiles, including but not limited to Seurat[13], SC3[14], SIMLR[15], and SCANPY[16]. However, most of these algorithms are not directly compatible with a single-cell ATAC-seq dataset, for which the signal matrix is much sparser. To characterize scATAC-seq data, the Greenleaf lab developed an algorithm named chromVAR[17], which aggregates mapped reads at accessible sites based on annotated motifs of known transcription factors (TFs) and thus projects the sparse per accessible peak per cell matrix to a bias-corrected deviation per motif per cell matrix and significantly stabilizes the data matrix for downstream clustering analysis. Other mathematical tools, such as the latent semantic indexing (LSI)[11, 12], Cicero[18], and cisTopic[19] have also been applied to process single-cell/nucleus ATAC-seq data[10, 12]. However, none of these methods can precisely distinguish cells without prior knowledge of essential principle components or TF motifs using limited computational resources. Therefore, a refined algorithm is urgently needed to better categorize cell subgroups with minor differences, thereby providing a deeper mechanistic understanding of single-cell epigenetic heterogeneity and regulation.

RESULTS

Accesson-based algorithm improves single-cell clustering

Here, we introduce a new single-cell chromatin accessibility analysis toolkit named APEC (accessibility pattern-based epigenomic clustering), which combines peaks with the same signal fluctuation among all single cells into peak groups, termed "accessons", and converts the original sparse cell-peak matrix to a much denser cell-accesson matrix for cell type categorization (Fig. 1a). In contrast to previous methods (e.g., chromVAR[17], LSI[11, 12], Cicero[18], and cisTopic[19]), this accesson-based reduction scheme naturally groups synergistic accessible regions together without a priori knowledge of genetic information (such as TF motifs or genomic distance) and provides more efficient, accurate, robust and rapid cell clustering from single-cell ATAC-seq profiles. More conveniently, APEC integrates all necessary procedures, from raw sequence trimming, alignment, and quality control (Supplementary Fig. 1a-i) to cell clustering, motif enrichment, and pseudotime trajectory prediction into a head-to-toe program package that has been made available on GitHub (https://github.com/QuKunLab/APEC).

To test the performance of APEC, we first obtained data from previous publications that performed scATAC-seq on a variety of cell types with known identity during hematopoietic stem cell (HSC) differentiation[20] as a gold standard. Compared to other state-of-the-art single cell chromatin accessibility analysis methods such as chromVAR[17, 21], LSI[11, 12], Cicero[18] and cisTopic[19], this new accesson-based algorithm more precisely and clearly clustered cells into their corresponding identities according to the Adjusted Rand Index (ARI) (Fig. 1b-c). CisTopic is the second most accurate method for predicting cell identities; however, the cisTopic algorithm is 15-50 times slower than the other methods, given the same computational resources. To further confirm the superiority of APEC, we performed the same comparison analysis with another scATAC-seq dataset on three distinct cell types, namely, lymphoid-primed multipotent progenitors (LMPPs), monocytes, and HL-60 lymphoblastoid cells (HL60), and two similar cell types, namely, blast cells and leukemic stem cells (LSCs) from two acute myeloid leukemia (AML) patients[17]. Again, we see that APEC outperformed all the other tools in cell clustering with an overall ARI=0.770/0.763 (Supplementary Fig. 2a). Interestingly, APEC, cisTopic and LSI were all capable of almost perfectly separating the three distinct cell types (LMPPs, monocytes, and HL60), with ARI = 1.000/0.994, 0.987/0.987 and 0.969, respectively, while the other two Cicero (ARI=0.917) and chromVAR (ARI=0.705) were not as good. However, in terms of clustering the two similar cell types from two AML patients, APEC (ARI=0.590/0.550) clearly outperformed cisTopic (ARI=0.477/0.497) and LSI (ARI=0.382) (Fig. 1d, Supplementary Fig. 2b), suggesting that APEC was the most sensitive among all the tools. Since each method can generate varying numbers of clusters depending on the parameters used, we benchmarked

the performance of all the methods using ARI across a wide range of parameters to ensure the

reliability of their predictions (Supplementary Fig. 2c-d). To further test the robustness of APEC

at low sequencing depth, we randomly selected reads from the raw data and calculated the ARI

values for each down-sampled data. APEC exhibits better performance at sequencing depths

as low as 20% of the original data (Supplementary Fig. 2e), confirming the sensitivity of the

algorithm.

Compare with chromVAR, the contribution of the minor differences between similar cells

is aggregated in accessons but diluted in motifs. For example, APEC identified prominent super-

enhancers around the E3 ligase inhibitor gene *N4BP1*[22] and the MLL fusion gene *GPHN*[23] in the

LSC cells from AML patient 1 (P1-LSC) but not in the other cell types (Supplementary Fig. 3a-b).

We noticed that all peaks in these super-enhancers were classified into one accesson that was

critical for distinguishing P1-LSCs from P2-LSCs, P1-blast cells and P2-blast cells. However,

these peaks were distributed in multiple TF motifs, which significantly diluted the contributions of

the minor differences (Supplementary Fig. 3c-d). In contrast to Cicero, which aggregates peaks

based on their co-accessibilities within a certain range of genomic distance[18], APEC aggregates

synergistic peaks genome-wide. Buenrostro et al. showed that the covariation of the accessible

sites across all the cells may reflect the spatial distance between the corresponding peaks[8]. By

integrating the chromatin conformation profiles from Hi-C experiments with the scATAC-seq

profile for the same cells, we found that peaks in the same accesson are spatially much closer

to each other than randomly selected peaks (Supplementary Fig. 4a, P-value<10^{-7}), suggesting

that they may belong to the same topologically associated domains (TADs) (Supplementary Fig.

4b).

Speed and scalability have become extremely important for single-cell analytical tools

due to the rapid growth in the number of cells sequenced in each experiment. We benchmarked

APEC and all the other tools based on a random sampling of the single-cell atlas of *in vivo*

mammalian chromatin accessibility dataset[24], which contains over 80,000 cells. With 1 CPU

thread and 32 GB RAM, APEC took 210 min to finish the cell clustering, ~4 times faster than

chromVAR and ~6-13 times slower than Cicero and LSI (Supplementary Fig. 4c). CisTopic, on

the other hand, took 1500 min to finish clustering 30,000 cells using 5 CPU threads, and we

estimate it would take 30 days to finish clustering 80,000 cells if no parallel computing is applied.

In addition, APEC is very stable for a wide range of parameter values used in the algorithm,

such as the number of accessons, nearest neighbors and principle components (Supplementary

Fig. 4d-f).

APEC is applicable to multiple single-cell chromatin detection techniques

To evaluate the compatibility and performance of APEC with other single-cell chromatin accessibility detection techniques, such as snATAC-seq[10], transcript-indexed scATAC-seq[25] and sciATAC-seq[11], APEC was also tested with the data sets generated by those experiments. For example, APEC discovered 13 cell subpopulations in adult mouse forebrain snATAC-seq data[10], including four clusters of excitatory neurons (EX1-4), five groups of inhibitory neurons (IN1-5), astroglia cells (AC1&2), oligodendrocyte cells (OC), and microglial cells (MG; Fig. 2a-b), defined by the chromatin accessibilities at the loci of cell type-specific genes (Fig. 2c). Compared to published results[10], APEC identified 4 rather than 3 excitatory subpopulations and 5 rather than 2 distinct inhibitory subpopulations, and all the cell groups were clearly distinguished from each other by hierarchical clustering (Fig. 2d). The motif enrichment analysis module in APEC identified cell type-specific regulators that are also consistent with previous publications[10]. For example, the NEUROD1 and OLIG2 motifs were generally enriched on excitatory clusters (EX1\3\4); the MEF2C motif was more enriched on EX1/2 and the left part of EX3 than other excitatory neurons; the motifs of MEIS2 and DLX2 were differentially enriched on two subtypes of inhibitory neurons (IN2 and IN4, respectively); and the NOTO, SOX2, and ETS1 motifs were enriched on the AC1, OC, and MG clusters, respectively (Fig. 2e). These results suggest that APEC is capable of identifying cell subtype-specific regulators.

Since single-cell transcriptome analysis is also capable to identify novel cell subpopulations, it is critical to anchor the cell types identified from scATAC-seq to those from scRNA-seq. Hrvatin et al. identified dozens of excitatory and inhibitory neuronal subtypes in the mouse visual cortex using single cell inDrops sequencing[26] and provided tens of signature genes that distinguished these cell types. Interestingly, the accessons that represent these signature genes were also distinctly enriched at corresponding clusters of neurons. For example, cells in the EX1 cluster in snATAC-seq-enriched accessons represent the gene *Enpp2*, which is the marker gene for cluster Excl23 that was defined in the inDrops-seq data (Fig. 2f, Supplementary Fig. 5a). Cluster EX1 in scATAC-seq also matched with Excl5_3 in inDrops-seq (marked by *Deptor* and *Fam3c*). Similarly, cluster EX2 matched Excl4 (marked by *Rorb* and *Tshz1*), EX3 matched Excl5_1 (marked by *Bmp3*), and EX4 matched Excl6 (marked by *Col5a1*, *Foxp2* and *Pcp4*). The same method also works to anchor inhibitory neurons, as the IN1 cells in the snATAC-seq data corresponded to the Int_Vip and Int_Npy clusters in the inDrops-seq data (marked by *Vip* and *Npy*). Cluster IN2 matched Int_Cck (marked by *Cck*), IN3 matched Int_Pv (marked by *Pvalb*), IN4 matched Int_Sst_2 (marked by *Crhbp*), and IN5 matched Int_Sst_1

(marked by *Lypd6b*) (Fig. 2g, Supplementary Fig. 5b). These results highlight the potential

advantages of the accesson-based approach for the integrative analysis of scATAC-seq and

scRNA-seq data.

In addition, due to the sparser per-cell-per-peak fragment count matrix, more than 29.7%

(946 out of 3034) of high-quality cells were previously unable to be correctly assigned into any

subpopulation of interest[10], but APEC successfully categorized all cells into their corresponding

subtypes, confirming its high sensitivity. In contrast, although chromVAR also identified 11 cell

clusters (Supplementary Fig. 6a-b), it mis-clustered AC and EX4 with inhibitory neurons

(Supplementary Fig. 6c). Cicero was also applied to distinguish these cells; however, since it

failed to predict the activities of several critical genes (e.g., *Neurod6*, *C1qb* and *Ctss*), the

algorithm was not able to clearly identify 20% of cells or to cluster them correctly

(Supplementary Fig. 6e-i). These results confirm that APEC can better distinguish and

categorize single cells with great sensitivity and reliability.

APEC constructs a pseudotime trajectory that predicts cell differentiation lineage

Cells are not static but dynamic entities, and they have a history, particularly a

developmental history. Although single-cell experiments often profile a momentary snapshot, a

number of remarkable computational algorithms have been developed to pseudo-order cells

based on the different points they were assumed to occupy in a trajectory, thereby leveraging

biological asynchrony[27, 28]. For instance, Monocle[28, 29] constructs the minimum spanning tree,

and Wishbone[30] and SPRING[31] construct the nearest neighbor graph from single-cell

transcriptome profiles. These tools have been widely used to depict neurogenesis[32],

hematopoiesis[33, 34] and reprogramming[35]. APEC integrates the Monocle algorithm into the

accesson-based method and enables pseudotime prediction from scATAC-seq data[20] and was

applied to investigate HSC differentiation linages (Fig. 3a). Principal component analysis (PCA)

of the accesson matrix revealed multiple stages of the lineage during HSC differentiation (Fig.

3b) and was consistent with previous publications[2, 20]. After utilizing the Monocle package,

APEC provided more precise pathways from HSCs to the differentiated cell types (Fig. 3c). In

addition to the differentiation pathways to MEP cells through the CMP state and to CLP cells

through the LMPP state, MPP cells may differentiate into GMP cells through two distinct

trajectories: Path A through the CMP state and Path B through the LMPP state, which is

consistent with the composite model of HSC and blood lineage commitment[36]. Notably, pDCs

from bone marrow are CD34[+] (Supplementary Fig. 7), indicative of precursors of pDCs. APEC
suggested that pDC precursors were derived from CLP cells on the pseudotime trajectory (Fig.
3c), which also agrees with previous reports[37]. Furthermore, APEC incorporated the chromVAR
algorithm to determine the regulatory mechanisms during HSC differentiation by evaluating the
deviation of each TF along the single-cell trajectory. As expected, the HOX motif is highly
enriched in the accessible sites of HSCs/MPP cells, as are the GATA1, CEBPB and TCF4
motifs, which exhibit gradients that increase along the erythroid, myeloid and lymphoid
differentiation pathways, respectively[20] (Fig. 3d). In addition, we can see that the TF regulatory
strategies of the two paths from MMPs towards GMP cells were very different. Finally, we
generated a hematopoiesis tree based on the APEC analysis (Fig. 3e).

Furthermore, we benchmarked the performance of APEC and of all the other tools in
constructing a pseudotime trajectory from the scATAC-seq profile on the same dataset. We
found that (1) when the raw peak count matrix was invoked into Monocle, almost none
developmental pathways were constructed (Supplementary Fig. 8a), suggesting that the peak
aggregation step in APEC greatly improves the pseudotime estimation; (2) APEC + Monocle
provides the most precise pathways from HSCs to differentiated cells, compared to those of
other peak aggregation methods, such as chromVAR, Cicero, LSI and cisTopic (Supplementary
Fig. 8b-e); and (3) when we applied other pseudotime trajectory construction methods, such as
SPRING[31], after APEC, a similar though less clear cell differentiation diagram was also obtained,
suggesting the reliability of our prediction (Supplementary Fig. 8f).

APEC reveals the single-cell regulatory heterogeneity of thymocytes

T cells generated in the thymus play a critical role in the adaptive immune system, and
the development of thymocytes can be divided into 3 main stages based on the expression of
the surface markers CD4 and CD8, namely, CD4 CD8 double-negative (DN), CD4 CD8 double-
positive (DP) and CD4 or CD8 single-positive (CD4SP or CD8SP, respectively) stages[38].
However, due to technical limitations, our genome-wide understanding of thymocyte
development at single-cell resolution remains unclear. Typically, more than 80% of thymocytes
stay in the DP stage in the thymus, whereas DN cells account for only approximately 3% of the
thymocyte population. To eliminate the impacts of great differences in proportion, we developed
a fluorescent tagmentation- and FACS-sorting-based scATAC-seq strategy (ftATAC-seq), which
combined the advantages of ATAC-see[39] and Pi-ATAC-seq[40] to manipulate the desired number
of target cells by indexed sorting (Fig. 4a). Tn5 transposomes were fluorescently labeled in each

cell to evaluate the tagmentation efficiency so that cells with low ATAC signals could be gated

out easily (Fig. 4b, Supplementary Fig. 9a). With ftATAC-seq, we acquired high-quality

chromatin accessibility data for 352 index-sorted DN, DP, CD4SP, and CD8SP single cells and

352 mixed thymocytes (Supplementary Fig. 9b-d). Correlation analysis with the published bulk

ATACs-eq data of thymocytes [41] indicates that the cells we sorted in ftATAC-seq were correctly

labeled (Supplementary Fig. 9e). We then applied APEC on this dataset to investigate the

chromatin accessibility divergence during developmental process and to reveal refined

regulome heterogeneity of mouse thymocytes at single-cell resolution. Taking into account all

the 130685 peaks called from the raw sequencing data, APEC aggregated 600 accessons and

successfully assigned over 82% of index-sorted DN, DP, CD4SP and CD8SP cells into the

correct subpopulations (Fig. 4c-d). As expected, the majority of randomly sorted and mixed

thymocytes were classified into DP subtypes based on hierarchical clustering of cell-cell

correlation matrix, which was consistent with the cellular subtype proportions in the thymus.

APEC further classified all thymocytes into 11 subpopulations, including 2 DN, 6 DP, 1 CD4SP,

2 CD8SP, suggesting that extensive epigenetic heterogeneity exists among cells with the same

CD4 and CD8 surface markers (Fig. 4e). For instance, there are four main subtypes of DN cells,

according to the expression of the surface markers CD44 and CD25[42], while two clusters were

identified in ftATAC-seq. The accessibility signals around the *Il2ra* (Cd25) and *Cd44* gene loci

demonstrated that DN.A1 comprised $CD44^+CD25^-$ and $CD44^+CD25^+$ DN subtypes (DN1 and

DN2), and DN.A2 cells comprised $CD44^-CD25^+$ and $CD44^-CD25^-$ subtypes (DN3 and DN4),

suggesting significant chromatin changes between DN2 and DN3 cell development (Fig. 4f).

Many TFs have been reported to be essential in regulating thymocyte development, and

we found that their motifs were remarkably enriched at different stages during the process (Fig.

4g). For instance, Runx3 is well known for regulating CD8SP cells[43], and we observed

significant enrichment of the RUNX motif on DN cells and a group of CD8SP cells. Similarly, the

TCF[44, 45], RORC[46] and NFkB[47] family in regulating the corresponding stages during this process.

More enriched TF motifs in each cell subpopulation were also observed, suggesting significant

regulatory divergence in thymocytes (Supplementary Fig. 10a). Interestingly, two clusters of

CD8SP cells appear to be differentially regulated based on motif analysis, in which CD8.A1 cells

are closer to DP cells, while CD8.A2 cells are more distant at the chromatin level, suggesting

that CD8.A2 cells are more mature CD8SP cells, and CD8.A1 cells are in a transitional state

between DP and SP cells.

APEC is capable of integrating single-cell transcriptional and epigenetic information by

scoring gene sets of interest based on their nearby peaks from scATAC-seq, thereby converting

the chromatin accessibility signals to values that are comparable to gene expression profiles

(Online **Methods**). To test the performance of this integrative analysis approach and to evaluate

the accuracy of thymocyte classification by APEC, we assayed the transcriptomes of single

thymocytes and obtained 357 high-quality scRNA-seq profiles using the SMART-seq2 protocol[48].

Unsupervised analysis of gene expression profiles clustered these thymocytes into 13 groups in

Seurat[13] (Supplementary Fig. 11a-b), and each subpopulation was identified based on known

feature genes (Supplementary Fig. 11c-d). We then compared the adjusted gene scores

obtained from APEC with the single-cell RNA expression profile and observed a strong

correlation between the subtypes identified from the transcriptome and the subtypes identified

from chromatin accessibility (Fig. 4h), confirming the reliability and stability of cellular

classification using APEC. We also applied Cicero to predict gene activities; however, the

correspondence between transcriptional and epigenomic cell clusters was not revealed by the

Cicero gene scores (Supplementary Fig. 10b-c).

APEC reconstructs the thymocyte developmental trajectory from ftATAC-seq profiles

APEC is capable of constructing a pseudotime trajectory and then predicting the cell

differentiation lineage from a "snapshot" of single-cell epigenomes (Fig. 3). We applied APEC to

recapitulate the developmental trajectory and thereby reveal the single-cell regulatory dynamics

during the maturation of thymocytes. Pseudotime analysis based on single-cell ftATAC-seq data

shaped thymocytes into 5 developing stages (Fig. 5a, Supplementary Fig. 12a-b), where most

of the cells in stages 1, 2, 4, and 5 were DN, DP, CD8SP and CD4SP cells, respectively. APEC

also identified a transitional stage 3, which was mainly consisted of last stages of DP cells.

Besides Monocle, a similar developmental pathway can also be constructed by SPRING[31]

based on the accesson matrix (Supplementary Fig 12c). Interestingly, the pseudotime trajectory

suggests three developmental pathways for this process, one of which started with stage 1 (DN)

and ended in stage 2 (DP), and the other two of which started with stage 1 (DN), went through a

transitional stage 3 and then bifurcated into stage 4 (CD8SP) and 5 (CD4SP). The predicted

developmental trajectory could also be confirmed by the gene expression of surface markers,

such as Cd4, Cd8, Runx3 and Ccr7 (Fig. 5b). To evaluate the gene ontology (GO) enrichments

over the entire process, we implemented an accesson-based GO module in APEC, which

highlights the significance of the association between cells and biological function (Fig. 5c). For

instance, T cells selections, including β-selection, positive selection and negative selection, start

from the DN3 stage. Consistent with this process, we observed a strong "T cell selection" GO

term on the trajectory path after DN.A1 (Supplemental Fig. 12d). Since TCR signals are
essential for T cell selection, we also observed the "T cell activation" GO term accompanied by
"T cell selection". Meanwhile, the regulation of protein binding signal was also decreased at SP
stages, indicating the necessity of weak TCR signal for the survival of SP T cells during
negative selection.

To further uncover the regulatory mechanism underlying this developmental process,
APEC was implemented to identify stage-specific enriched TFs along the trajectory and pinpoint
the "pseudotime" at which the regulation occurs. In addition to the well-studied TFs mentioned
above (Fig. 4g, Supplemental Fig. 10a), APEC also identified Zeb1[49], Ctcf[50] and Id4 as potential
stage-specific regulators (Fig. 5d). Interestingly, the Id4 motif enriched on DP cells was also
reported to regulate apoptosis in other cell types[51, 52]. Associated with the fact that the vast
majority of DP thymocytes die because of a failure of positive selection[53], we hypothesize that
stage 2 may be the path towards DP cell apoptosis. We then checked the distribution of DP
cells along the stage 2 trajectory and found that most DP.A1 cells were scattered in "early"
stage 2, and they were enriched with GO terms such as "T cell selection", "cell activation" and
"differentiation" (Fig. 5e, Supplementary Fig. 12e). However, most DP.A4/A5 cells were
distributed at the end of stage 2, and their principle accessons were enriched with GO terms
such as "apoptosis" and "chromatin modification". Although it is believed that more than 95% of
DP thymocytes die during positive selection, only a small proportion of apoptotic cells could be
detected in a snapshot of the thymus, which in our data are the cells at the end of stage 2. By
comparing the number of cells near stage 3 with all the cells in stage 2, we estimated that ~3-5%
of cells would survive positive selection, which is consistent with the findings reported in
previous publications[54, 55]. Our data suggest that before entering the final apoptotic stage, DP
thymocytes under selection could have already been under apoptotic stress at the chromatin
level, which explains why DP cells are more susceptible to apoptosis than other thymocyte
subtypes[56].

DISCUSSION

Here, we introduced an accesson-based algorithm for single-cell chromatin accessibility
analysis. Without any prior information (such as motifs), this approach generated more refined
cell groups with reliable biological functions and properties. Integrating the new algorithm with
all necessary chromatin sequencing data processing tools, APEC provides a comprehensive
solution for transforming raw experimental single-cell data into final visualized results. In

addition to improving the clustering of subtle cell subtypes, APEC is also capable of locating

potential specific super-enhancers, searching enriched motifs, estimating gene activities, and

constructing time-dependent cell developmental trajectories, and it is compatible with many

existing single-cell accessibility datasets. Compared with all the other state-of-the-art single cell

chromatin accessibility analysis methods, APEC clearly shows superiority in correctly predicting

cell identities and precisely constructing developmental trajectories. APEC is also very robust

and stable and is scalable to clustering a large number of cells using limited computational

resources. Despite these advantages, the biological implications of accessons are still obscure,

especially for those that involve only a small number of peaks. Although we noticed peaks in the

same accesson may belong to the same TADs, further investigations are still required to fully

uncover the biology that underlies accessons.

To evaluate the performance of this approach in the context of the immune system, we

adopted APEC with scATAC-seq technology to investigate the regulome dynamics of the thymic

development process. We developed a novel method of ftATAC-seq that captures Tn5-tagged

single cells of interest and outlines the chromatin accessibility heterogeneity and dynamics

during this process. Coordinated with essential cell surface markers, APEC provided a much

more in-depth classification of thymocytes than the conventional DN, DP, CD4SP and CD8SP

stages based on single-cell chromatin status. By reconstructing the developmental pseudotime

trajectory, APEC discovered a transitional stage before thymocytes bifurcate into CD4SP and

CD8SP cells and inferred that one of the stages leads to cell apoptosis. Considering that more

than 95% of DP cells undergo apoptosis, which is a programmed cell death process, we believe

that before DP cells enter the final apoptotic state, there would already be some intracellular

changes towards apoptosis, which might not be detected by traditional biochemistry methods,

such as annexin V and TUNEL staining. Our data suggested that DP cells were gradually

programmed to undergo apoptosis at the chromatin level; however, further studies are needed

to fully understand the regulatory mechanism of this process.

Acknowledgments

This work was supported by the National Key R&D Program of China (2017YFA0102900 to

K.Q.) and by National Natural Science Foundation of China grants (81788101, 91640113,

31771428 to K.Q.). It was also supported by Anhui Provincial Natural Science Foundation grant

BJ2070000097 (to B.L.) and 1908085QH326 (to Y.L.). We thank the Howard Chang lab at

Stanford University for helpful discussion. We thank the USTC supercomputing center and the
School of Life Science Bioinformatics Center for providing supercomputing resources for this
project.

Authors' contributions

385 KQ, BL and YL conceived the project, BL developed the APEC software and performed all data
386 analysis with helps from KL, QY, PC, JF, WZ, PD and CJ. YL developed ftATAC-seq technique
387 and performed all scATAC-seq and scRNA-seq experiments with helps from LZ. KL analyzed
388 scRNA-seq data. BL, YL and KQ wrote the manuscript with inputs from all other authors.

Data and code availability

Mouse thymocytes ftATAC-seq data can be obtained from the Genome Sequence Archive at
BIG Data Center with the accession number CRA001267 and is available via
http://bigd.big.ac.cn/gsa/s/yp1164Et. Other published data sets used in this study are available
from NIH GEO with accession numbers GSE74310[2], GSE65360[8], GSE96772[20], GSE100033[10],
GSE111586[24], and GSE63525[57]. APEC pipeline can be downloaded from the GitHub website
(https://github.com/QuKunLab/APEC).

REFERENCES

1.   Buenrostro, J.D., Giresi, P.G., Zaba, L.C., Chang, H.Y. & Greenleaf, W.J. Transposition of native
chromatin for fast and sensitive epigenomic profiling of open chromatin, DNA-binding proteins
and nucleosome position. *Nat Methods* **10**, 1213-1218 (2013).
2.   Corces, M.R. et al. Lineage-specific and single-cell chromatin accessibility charts human
hematopoiesis and leukemia evolution. *Nat Genet* **48**, 1193-1203 (2016).
3.   Wu, J. et al. The landscape of accessible chromatin in mammalian preimplantation embryos.
Nature **534**, 652-657 (2016).
4.   Jorstad, N.L. et al. Stimulation of functional neuronal regeneration from Muller glia in adult mice.
Nature **548**, 103-107 (2017).
5.   Su, Y. et al. Neuronal activity modifies the chromatin accessibility landscape in the adult brain.
Nat Neurosci **20**, 476-483 (2017).
6.   Denny, S.K. et al. Nfib Promotes Metastasis through a Widespread Increase in Chromatin
Accessibility. *Cell* **166**, 328-342 (2016).
7.   Qu, K. et al. Chromatin Accessibility Landscape of Cutaneous T Cell Lymphoma and Dynamic
Response to HDAC Inhibitors. *Cancer Cell* **32**, 27-41 e24 (2017).

8.   Buenrostro, J.D. et al. Single-cell chromatin accessibility reveals principles of regulatory variation.
Nature **523**, 486-490 (2015).

9.   Lake, B.B. et al. Integrative single-cell analysis of transcriptional and epigenetic states in the
human adult brain. *Nat Biotechnol* **36**, 70-80 (2018).

10.   Preissl, S. et al. Single-nucleus analysis of accessible chromatin in developing mouse forebrain
reveals cell-type-specific transcriptional regulation. *Nat Neurosci* **21**, 432-439 (2018).

11.   Cusanovich, D.A. et al. Multiplex single cell profiling of chromatin accessibility by combinatorial
cellular indexing. *Science* **348**, 910-914 (2015).

12.   Cusanovich, D.A. et al. The cis-regulatory dynamics of embryonic development at single-cell
resolution. *Nature* **555**, 538-542 (2018).

13.   Butler, A., Hoffman, P., Smibert, P., Papalexi, E. & Satija, R. Integrating single-cell transcriptomic
data across different conditions, technologies, and species. *Nat Biotechnol* **36**, 411-420 (2018).

14.   Kiselev, V.Y. et al. SC3: consensus clustering of single-cell RNA-seq data. *Nat Methods* **14**, 483-
486 (2017).

15.   Wang, B., Zhu, J., Pierson, E., Ramazzotti, D. & Batzoglou, S. Visualization and analysis of single-
cell RNA-seq data by kernel-based similarity learning. *Nat Methods* **14**, 414-416 (2017).

16.   Wolf, F.A., Angerer, P. & Theis, F.J. SCANPY: large-scale single-cell gene expression data analysis.
Genome Biol **19**, 15 (2018).

17.   Schep, A.N., Wu, B., Buenrostro, J.D. & Greenleaf, W.J. chromVAR: inferring transcription-factor-
associated accessibility from single-cell epigenomic data. *Nat Methods* **14**, 975-978 (2017).

18.   Pliner, H.A. et al. Cicero Predicts cis-Regulatory DNA Interactions from Single-Cell Chromatin
Accessibility Data. *Mol Cell* **71**, 858-871 e858 (2018).

19.   Bravo González-Blas, C. et al. Cis-topic modelling of single cell epigenomes. *bioRxiv*
(2018).

20.   Buenrostro, J.D. et al. Integrated Single-Cell Analysis Maps the Continuous Regulatory Landscape
of Human Hematopoietic Differentiation. *Cell* **173**, 1535-1548 e1516 (2018).

21.   Zamanighomi, M. et al. Unsupervised clustering and epigenetic classification of single cells. *Nat
Commun* **9**, 2410 (2018).

22.   Oberst, A. et al. The Nedd4-binding partner 1 (N4BP1) protein is an inhibitor of the E3 ligase Itch.
Proc Natl Acad Sci U S A **104**, 11280-11285 (2007).

23.   Eguchi, M. et al. GPHN, a novel partner gene fused to MLL in a leukemia with t(11;14)(q23;q24).
Genes Chromosomes Cancer **32**, 212-221 (2001).

24.   Cusanovich, D.A. et al. A Single-Cell Atlas of In Vivo Mammalian Chromatin Accessibility. *Cell* **174**,
1309-+ (2018).

25.   Satpathy, A.T. et al. Transcript-indexed ATAC-seq for precision immune profiling. *Nat Med* **24**,
580-590 (2018).

26.   Hrvatin, S. et al. Single-cell analysis of experience-dependent transcriptomic states in the mouse
visual cortex. *Nat Neurosci* **21**, 120-129 (2018).

27.   Bendall, S.C. et al. Single-cell trajectory detection uncovers progression and regulatory
coordination in human B cell development. *Cell* **157**, 714-725 (2014).

28.   Trapnell, C. et al. The dynamics and regulators of cell fate decisions are revealed by
pseudotemporal ordering of single cells. *Nat Biotechnol* **32**, 381-386 (2014).

29.   Qiu, X. et al. Reversed graph embedding resolves complex single-cell trajectories. *Nat Methods*
14, 979-982 (2017).

30.   Setty, M. et al. Wishbone identifies bifurcating developmental trajectories from single-cell data.
Nat Biotechnol **34**, 637-645 (2016).

31.   Weinreb, C., Wolock, S. & Klein, A.M. SPRING: a kinetic interface for visualizing high dimensional
single-cell expression data. *Bioinformatics* **34**, 1246-1248 (2018).

32.   Habib, N. et al. Div-Seq: Single-nucleus RNA-Seq reveals dynamics of rare adult newborn
463   neurons. *Science* **353**, 925-928 (2016).

464   33.   Olsson, A. et al. Single-cell analysis of mixed-lineage states leading to a binary cell fate choice.
465   *Nature* **537**, 698-+ (2016).

466   34.   Zhou, F. et al. Tracing haematopoietic stem cell formation at single-cell resolution. *Nature* **533**,
467   487-+ (2016).

468   35.   Treutlein, B. et al. Dissecting direct reprogramming from fibroblast to neuron using single-cell
469   RNA-seq. *Nature* **534**, 391-+ (2016).

470   36.   Adolfsson, J. et al. Identification of Flt3+ lympho-myeloid stem cells lacking erythro-
471   megakaryocytic potential a revised road map for adult blood lineage commitment. *Cell* **121**, 295-
472   306 (2005).

473   37.   Chistiakov, D.A., Orekhov, A.N., Sobenin, I.A. & Bobryshev, Y.V. Plasmacytoid dendritic cells:
474   development, functions, and role in atherosclerotic inflammation. *Front Physiol* **5**, 279 (2014).

475   38.   Germain, R.N. T-cell development and the CD4-CD8 lineage decision. *Nat Rev Immunol* **2**, 309-
476   322 (2002).

477   39.   Chen, X. et al. ATAC-see reveals the accessible genome by transposase-mediated imaging and
478   sequencing. *Nat Methods* **13**, 1013-1020 (2016).

479   40.   Chen, X. et al. Joint single-cell DNA accessibility and protein epitope profiling reveals
480   environmental regulation of epigenomic heterogeneity. *Nat Commun* **9**, 4590 (2018).

481   41.   Yoshida, H. et al. The cis-Regulatory Atlas of the Mouse Immune System. *Cell* **176**, 897-912 e820
482   (2019).

483   42.   Godfrey, D.I., Kennedy, J., Suda, T. & Zlotnik, A. A developmental pathway involving four
484   phenotypically and functionally distinct subsets of CD3-CD4-CD8- triple-negative adult mouse
485   thymocytes defined by CD44 and CD25 expression. *J Immunol* **150**, 4244-4252 (1993).

486   43.   Taniuchi, I. et al. Differential requirements for Runx proteins in CD4 repression and epigenetic
487   silencing during T lymphocyte development. *Cell* **111**, 621-633 (2002).

488   44.   Ioannidis, V., Beermann, F., Clevers, H. & Held, W. The beta-catenin--TCF-1 pathway ensures
489   CD4(+)CD8(+) thymocyte survival. *Nat Immunol* **2**, 691-697 (2001).

490   45.   Yu, S. et al. The TCF-1 and LEF-1 transcription factors have cooperative and opposing roles in T
491   cell development and malignancy. *Immunity* **37**, 813-826 (2012).

492   46.   Sun, Z. et al. Requirement for RORgamma in thymocyte survival and lymphoid organ
493   development. *Science* **288**, 2369-2373 (2000).

494   47.   Gerondakis, S., Fulford, T.S., Messina, N.L. & Grumont, R.J. NF-kappaB control of T cell
495   development. *Nat Immunol* **15**, 15-25 (2014).

496   48.   Picelli, S. et al. Smart-seq2 for sensitive full-length transcriptome profiling in single cells. *Nat
497   Methods* **10**, 1096-1098 (2013).

498   49.   Higashi, Y. et al. Impairment of T cell development in deltaEF1 mutant mice. *J Exp Med* **185**,
499   1467-1479 (1997).

500   50.   Heath, H. et al. CTCF regulates cell cycle progression of alphabeta T cells in the thymus. *EMBO J*
501   **27**, 2839-2850 (2008).

502   51.   Andres-Barquin, P.J., Hernandez, M.C. & Israel, M.A. Id4 expression induces apoptosis in
503   astrocytic cultures and is down-regulated by activation of the cAMP-dependent signal
504   transduction pathway. *Exp Cell Res* **247**, 347-355 (1999).

505   52.   Carey, J.P., Knowell, A.E., Chinaranagari, S. & Chaudhary, J. Id4 promotes senescence and
506   sensitivity to doxorubicin-induced apoptosis in DU145 prostate cancer cells. *Anticancer Res* **33**,
507   4271-4278 (2013).

508   53.   Surh, C.D. & Sprent, J. T-cell apoptosis detected in situ during positive and negative selection in
509   the thymus. *Nature* **372**, 100-103 (1994).

54. Huesmann, M., Scott, B., Kisielow, P. & von Boehmer, H. Kinetics and efficacy of positive
selection in the thymus of normal and T cell receptor transgenic mice. *Cell* **66**, 533-540 (1991).
55. Shortman, K., Vremec, D. & Egerton, M. The kinetics of T cell antigen receptor expression by
subgroups of CD4+8+ thymocytes: delineation of CD4+8+3(2+) thymocytes as post-selection
intermediates leading to mature T cells. *J Exp Med* **173**, 323-332 (1991).
56. Chow, S.C., Snowden, R., Orrenius, S. & Cohen, G.M. Susceptibility of different subsets of
immature thymocytes to apoptosis. *Febs Lett* **408**, 141-146 (1997).
57. Rao, S.S. et al. A 3D map of the human genome at kilobase resolution reveals principles of
chromatin looping. *Cell* **159**, 1665-1680 (2014).
519

  FIGURES

Figure 1. The accesson matrix constructed from the sparse fragment count matrix improved the
clustering of scATAC-seq data. (**a**) Step-by-step workflow of APEC. Peaks were grouped into
accessons by their accessibility pattern among cells with the K nearest neighbors (KNN) method.
(**b**) t-Distributed Stochastic Neighbor Embedding (tSNE) diagrams based on the dimension-
transformed matrices from different algorithms, i.e., APEC: accesson matrix; cisTopic: topic
matrix; LSI: LSI matrix; chromVAR: bias-corrected deviation matrix; Cicero: aggregated model
matrix. The cells are FACS-indexed human hematopoietic cells, including HSCs (hematopoietic
stem cells), MPPs (multipotent progenitors), LMPPs (lymphoid-primed multipotential
progenitors), CMPs (common myeloid progenitors), CLPs (common lymphoid progenitors),
pDCs (plasmacytoid dendritic cells), GMPs (granulocyte-macrophage progenitors), MEPs
(megakaryocyte-erythroid progenitors), and unknown cells. (**c**) The average ARI (Adjusted Rand

533   Index) scores and computing time for the clustering of the human hematopoietic cells by

534   different algorithms. (**d**) The average ARI scores and computing time for the clustering of the

535   blast cells and LSCs (leukemic stem cells) from two AML (acute myeloid leukemia) patients (P1

536   and P2) by different algorithms. As in (c), cisTopic was performed using 5 CPU threads and all

537   other tools with 1 CPU thread.

Figure 2. APEC improved the cell type classification of adult mouse forebrain snATAC-seq data.
(**a**) A tSNE diagram demonstrates the APEC clustering of forebrain cells. (**b**) Hierarchical
clustering of the cell-cell correlation matrix. The side bar denotes cell clusters from APEC. (**c**)
Average of the marker gene scores for each cell cluster, normalized by the standard score (z-
score). The top row lists the cell numbers for each cluster. (**d**) Hierarchical clustering of the
cluster-cluster correlation matrix. (**e**) Differential enrichments of cell type-specific motifs in
different clusters. (**f, g**) Intensities of representative accessons of the excitatory (EX) and
inhibitory (IN) clusters from snATAC-seq associated with the activities of signature genes of the
excitatory (Excl) and inhibitory (Int) neuron subtypes defined in inDrops-seq[26].

Figure 3. APEC constructed a differentiation pathway from scATAC-seq data from human hematopoietic cells. (**a**) The pseudotime trajectory construction scheme based on the accesson matrix and Monocle. (**b**) Principal component analysis (PCA) of the accesson matrix for human hematopoietic cells. The first principal component is not shown here because it was highly correlated with sequencing depth[20]. HSC, hematopoietic stem cell; MPP, multipotent progenitor; LMPP, lymphoid-primed multipotential progenitor; CMP, common myeloid progenitor; CLP, common lymphoid progenitor; pDC, plasmacytoid dendritic cell; GMP, granulocyte-macrophage

progenitor; MEP, megakaryocyte-erythroid progenitor; unknown, unlabeled cells. (**c**) Pseudotime trajectory for the same data constructed by applying Monocle on the accesson matrix. Paths A and B represent different pathways for GMP cell differentiation. (**d**) The deviations of significant differential motifs (HOXA9, GATA1, CEBPB, and TCF4) plotted on the pseudotime trajectory. (**e**) Modified schematic of human hematopoietic differentiation.

Figure 4. APEC accurately identified cell subtypes based on scATAC-seq data from *Mus musculus* thymocytes. (**a**) Experimental workflow of the fluorescent tagmentation- and FACS-sorting-based scATAC-seq strategy (ftATAC-seq). (**b**) Indexed sorting of double-negative (DN), double-positive (DP), CD4+ single-positive (CD4SP), and CD8+ single-positive (CD8SP) cells with strong tagmentation signals. (**c**) The tSNE of thymocyte single-cell ftATAC-seq data based on the accesson matrix, in which the cells are labeled by the sorting index. (**d**) Hierarchical clustering of the cell-cell correlation matrix. On the sidebar, each cell was colored by the sorting

index. (**e**) The accesson-based Louvain method clustered thymocytes into 11 subtypes. DN.A1

(dark green) & A2 (light green), double-negative clusters; DP.A1~A5 and DPsel.A, double-

positive clusters; CD8.A1 (dark blue) & A2 (light blue), CD8$^+$ single-positive clusters; CD4.A

(purple), CD4$^+$ single-positive cluster. (**f**) Average fragment counts of two DN clusters around

the marker genes *Cd44* and *Il2ra*. (**g**) Differential enrichment of the motifs Runx, Tcf, Rorc, and

Nfkb in the cell clusters. (**h**) Z-score of the Fisher exact test -log(p-value) of the common

differential genes between the cell clusters from different experiments. The row and column

clusters were identified by data from single-cell transcriptome (SMART-seq) and chromatin

accessibility (ftATAC-seq) analysis respectively.

Figure 5. APEC depicted the developmental pathways of *Mus musculus* thymocytes by pseudotime analysis. (**a**) Pseudotime trajectory based on the accesson matrix of thymocyte ftATAC-seq data. Cell colors were defined by the developmental stages along pseudotime. Pie charts show the proportion of cell clusters at each stage. (**b**) APEC scores of important marker genes (*Cd8a*, *Cd4*, *Runx3*, and *Ccr7*) along each branch of the pseudotime trajectory. (**c**) Weighted scores of important functional GO terms along each branch of the pseudotime trajectory. (**d**) Enrichment of specific motifs searched from the differential accessons of each cell subtype. (**e**) On the stage 2 branch, the cell number distribution of clusters DP.A1~A5 along pseudotime (upper panel) and the intensity of marker accessons of DP.A1 and DP.A4/A5 (lower panel) with top enriched GO terms.

METHODS

Mice. C57BL/6 mice were purchased from Beijing Vital River Laboratory Animal Technology
and maintained under specific pathogen-free conditions until the time of experiments. All mouse
experiments in this study were reviewed and approved by the Institutional Animal Care and Use
Committee of the University of Science and Technology of China.

ftATAC-seq on mouse thymocytes. Alexa fluor 488-labeled adaptor oligonucleotides were
synthesized at Sangon Biotech as follows: Tn5ME, 5'-[phos]CTGTCTCTTATACACATCT-3';
AF488-R1, 5'-AF488- TCGTCGGCAGCGTCAGATGTGTATAAGAGACAG-3'; and AF488-R2,
5'-AF488-GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAG-3'. Then, 50 µM of AF488-
R1/Tn5ME and AF488-R2/Tn5ME were denatured separately in TE buffer (Qiagen) at 95 °C for
5 min and cooled down to 22 °C at 0.1 °C/s. AF488-labeled adaptors were assembled onto
Robust Tn5 transposase (Robustnique) according to the user manual to form fluorescent
transposomes.

Thymus tissues isolated from 6- to 8-week-old male mice were gently ground in 1 mL of RPMI-
1640. Thymocytes in a single-cell suspension were counted after passing through a 40 µm
nylon mesh. A total of 1×10^6 thymocytes were stained with PerCP-Cy5.5-anti-CD45, PE-anti-
CD8a and APC-Cy7-anti-CD4 antibodies (Biolegend) and then fixed in 1× PBS containing 1%
methanal at room temperature for 5 min. After washing twice with 1× PBS, the cells were
counted again. A total of 1×10^5 fixed cells were resuspended in 40 µL of 1× TD buffer (5 mM
Tris-HCl, pH 8.0, 5 mM $MgCl_2$, and 10% DMF) containing 0.1% NP-40. Then, 10 µL of
fluorescent transposomes were added and mixed gently. Fluorescent tagmentation was
conducted at 55 °C for 30 min and stopped by adding 200 µL of 100 mM EDTA directly to the
reaction mixture. The cells were loaded on a Sony SH800S sorter, and single cells of the
$CD45^+/AF488-Tn5^{hi}$ population were index-sorted into each well of 384-well plates. The 384-well
plates used to acquire sorted cells were loaded with 2 µL of release buffer (50 mM EDTA, 0.02%
SDS) before use. After sorting, the cells in the wells were incubated for 1 min. Plates that were
not processed immediately were preserved at -80 °C.

To prepare a single-cell ATAC-seq library, plates containing fluorescently tagmented cells were
incubated at 55 °C for 30 min. Then, 4.2 µL of PCR round 1 buffer (1 µL of 100 µM $MgCl_2$, 3 µL
of 2× I-5 PCR mix [MCLAB], and 0.1 µL each of 10 µM R1 and R2 primers) were added to each
well, followed by PCR: 72 °C for 10 min; 98 °C for 3 min; 10 cycles of 98 °C for 10 s, 63 °C for
30 s and 72 °C for 1 min; 72 °C for 3 min; and holding at 4 °C. Thereafter, each well received 4

µL of PCR round 2 buffer (2 µL of I-5 PCR Mix, 0.5 µL each of Ad1 and barcoded Ad2 primers,
and 1 µL of ddH$_2$O), and final PCR amplification was carried out: 98 °C for 3 min; 12 cycles of
98 °C for 10 s, 63 °C for 30 s and 72 °C for 1 min; 72 °C for 3 min; and holding at 4 °C. Wells
containing different Ad2 barcodes were collected together and purified with a QIAquick PCR
purification kit (Qiagen). Libraries were sequenced on an Illumina HiSeq X Ten system.

SMART-seq on thymocytes. Thymocytes were stained and sorted directly into 384-well plates
without fixation. SMART-seq was performed as described with some modifications[58]. Reverse
transcription and the template-switch reaction were performed at 50 °C for 1 hr with Maxima H
Minus Reverse Transcriptase (Thermo Fisher); for library construction, 0.5-1 ng of cDNA was
fragmented with 0.05 µL of Robust Tn5 transposome in 20 µL of TD buffer at 55 °C for 10 min,
then purified with 0.8× VAHTS DNA Clean Beads (Vazyme Biotech), followed by PCR
amplification with Ad1 and barcoded Ad2 primers and purification with 0.6× VAHTS DNA Clean
Beads. Libraries were sequenced on an Illumina HiSeq X Ten system.

Data source. All experimental raw data used in this paper are available online. The single-cell
data for mouse thymocytes captured by the ftATAC-seq experiment can be obtained from the
Genome Sequence Archive at BIG Data Center with the accession number CRA001267 and is
available via http://bigd.big.ac.cn/gsa/s/yp1164Et. Other published data sets used in this study
are available from NIH GEO: (1) scATAC-seq data for LSCs and leukemic blast cells from
patients SU070 and SU353, LMPP cells, and monocytes from GSE74310[2]; (2) scATAC-seq
data for HL-60 cells from GSE65360[8]; and (3) scATAC-seq data for hematopoietic development
(HSCs, MPPs, CMPs, LMPPs, GMPs, EMPs, CLPs and pDCs) from GSE96772[20]. (4) APEC is
also compatible with a preprocessed fragment count matrix from the snATAC-seq data for the
forebrain of adult mice (p56) from GSE100033[10]. (5) The computational efficiency of APEC and
other methods was tested using data from the single-cell atlas of mouse chromatin accessibility
(sciATAC-seq) from GSE111586[24]. (6) The scATAC-seq (GSE65360[8]) and Hi-C (GSE63525[57])
data of GM12878 cells were used to generate the spatial correlation of peaks in the same or in
different accessons.

Preparing the fragment count matrix from the raw data. APEC adopted the general mapping,
alignment, peak calling and motif searching procedures to process the scATAC-seq data from
ATAC-pipe[59]. We also implemented the python script in ATAC-pipe[59] to trim the adapters in the
raw data (in paired-end fastq format files for each single-cell sample). APEC used BOWTIE2 to
map the trimmed sequencing data to the corresponding genome index and used PICARD for
the sorting, duplicate removal, and fragment length counting of the aligned data. The pipeline

called peaks from the merged file of all cells by MACS2, ranked and filtered out the low quality

peaks based on the false discovery rate (Q-value). Genomic locations of the peaks were

annotated by HOMER, and motifs searched by FIMO. APEC calculates the number of

fragments and the percent of reads mapped to the TSS region (±2000 BP) for each cell, and

filters out high quality cells for downstream analysis. All required files for the hg19 and mm10

assembly have been integrated into the pipeline. If users want to process data from other

species, they can also download corresponding reference files from the UCSC website. By

combining existing tools, APEC made it possible to finish all of the above data processing steps

by one command line, and generate a fragment count matrix for subsequent cell clustering and

differential analysis. APEC has been made available on GitHub

(https://github.com/QuKunLab/APEC).

Accesson-based clustering algorithm. We define accesson as a set of peaks with similar

accessibility patterns across all single cells, similar to the definition of gene module for RNA-seq

data. After preprocessing, a filtered fragment count matrix **B** is obtained, and APEC groups

peaks to construct accessons and then performs cell clustering analysis as follows:

(1) *Normalization of the fragment count matrix.* Each matrix element B_{ij} represents the number

of raw reads in cell i and peak j, and element B_{ij} was then normalized by the total number

of reads in each cell i, as if there are 10,000 reads in each cell.

$$B'_{ij} = log_2\left(\frac{B_{ij} \times 10000}{\sum_{j'} B_{ij'}} + 1\right)$$

(2) *Constructing accessons.* The top 40 principal components of the normalized matrix were

used to construct the connectivity matrix (**C**_{peak}) of peaks by the K-nearest-neighbor (KNN)

method with K=10. The grouping of peaks is insensitive to the number of principal

components and the number of nearest neighbors, so it is usually not necessary to change

these two parameters for different data sets. Based on the matrix **C**_{peak}, all peaks were

grouped by agglomerative clustering with Euclidean distance and Ward linkage method,

and the sum of one peak group was an accesson. For most data sets, we recommend

setting the number of accessons to a value between 500 and 1500, and the default was set

to 600, however the cell clustering result is not sensitive to the choice of accesson number

within this range. We then built the accesson count matrix **M** by summing the fragment

count of all peaks in one accesson. Thus, each column of matrix **M** is an accesson, each

row is a cell, and each element represents the cumulative fragment count of each accesson

in each cell.

(3) *Accesson filtering and normalization.* Not all accessons were used for cell clustering, and
those with low dispersion were filtered out to improve cell clustering. The Gini coefficient[60,]
[61] was used to measure the dispersion/inequality of the fragment count numbers of each
accesson among cells, i.e.,

$$Gini_j = \frac{Mean\left(\left|\vec{M}_j \otimes \vec{M}_j\right|\right)}{Mean\left(\vec{M}_j\right)}, \quad j = 1 \sim N_{accesson}$$

694   where $\vec{M}_j$ is the jth column of the accesson count matrix **M**. Since the Gini coefficient
695   increases with the mean count of the low dispersion accessons, we fitted the Gini
696   coefficients and the mean counts of all accessons into a linear equation, i.e.,

$$Gini_j = a \cdot Mean\left(\vec{M}_j\right) + b, \quad j = 1 \sim N_{accesson}$$

and selected the accessons above the line, i.e., the accessons with high dispersion, from
the accesson count matrix **M**. The filtered accesson matrix was then normalized by
calculating the z-score or probability of the fragment count for each row (i.e., each cell) to
generate normalized matrix **M$_a$** for the next step of cell clustering.

(4) *Cell clustering.* From the filtered and normalized accesson matrix **M$_a$**, APEC established the
connectivity matrix by computing the k-neighbor graph of all cells. Since the Louvain
algorithm was proven to be a reliable single-cell clustering method in Seurat[13] and Scanpy[16],
we adopted it in APEC to automatically predict the number of clusters from the connectivity
matrix and defined each Louvain community as a cell cluster. APEC uses the Louvain
algorithm to predict cluster number and perform cell clustering as default. Meanwhile, if
users want to artificially define the number of cell clusters, APEC can also perform KNN
clustering on the connectivity matrix.

(5) *Compare the performance of APEC with that of other methods on cells with known identity.*
To investigate the accuracy of the cell clusters predicted by different algorithms, we used
the ARI value, which evaluates the similarity of clustering results with all known types of
cells[14]. The ARI value can be calculated as follows:

$$ARI = \frac{\sum_{ij}\binom{n_{ij}}{2} - \sum_i\binom{a_i}{2} \cdot \sum_j\binom{b_j}{2}\Big/\binom{n}{2}}{\left[\sum_i\binom{a_i}{2} + \sum_j\binom{b_j}{2}\right]\Big/2 - \sum_i\binom{a_i}{2} \cdot \sum_j\binom{b_j}{2}\Big/\binom{n}{2}}$$

where n_{ij} is the element from the contingency matrix, a_i and b_j are the sums of the ith row
and jth column, respectively, and $\binom{x}{y}$ denotes a binomial coefficient. The higher the ARI
value is, the more accurate the clustering method is.

(6) *Characteristics of accesson*. The peaks of a same accesson can be distant from each other

on the genome, and sometimes even on multiple chromosomes. The average number of

peaks per accesson depends on the total number of peaks in the dataset and the number

of accessons set in the program (default 600). Usually the total number of peaks can vary

between ~40,000-150,000 depending on the total number of cells and the sequencing

depth for each cell, thereby the average number of peaks per accesson is around ~60-250.

Beside, we chose the top 40 principle components (PCs) of the normalized matrix to

construct the connectivity matrix since the first 3~5 PCs are usually not sufficient to capture

the detailed features of a single cell dataset, as described in Seurat and many other single

cell analysis tools. As described on the scikit-learn website (https://scikit-

[learn.org/stable/modules/clustering.html)](learn.org/stable/modules/clustering.html), the use of the KNN and then 'Euclidean' distance

and 'Ward' linkage methods to build the connectivity matrix between cells and perform the

agglomerative clustering for the connectivity matrix usually provides better clustering

results for different types of datasets. Although default values were chosen to provide

better clustering results based on analysis of multiple datasets, users can adjust these

parameters as needed.

Sampling of accesson number. To test if the APEC clustering result is sensitive to the choice

of accesson numbers, we sampled 100 different accesson numbers from 500 to 1500 in steps

of 10 and clustered the cells of each dataset 100 times. APEC generated stable clustering

results in terms of the average ARI on these datasets, with a wide range of different accesson

numbers (Supplementary Fig. 4d). We used accesson number = 600 as the default in APEC.

Parameter settings for other algorithms. To quantify the cell clustering performance of APEC,

we compared APEC with other state-of-the-art single-cell epigenomic algorithms on the same

datasets with gold standards, including cisTopic[19], LSI[11, 12], chromVAR[17] and Cicero[18]. Since

most of them have no cell clustering algorithm within their original codes, we applied the

Louvain clustering algorithm on their transformed matrices to fairly compare their performance.

We adopted the default settings of these tools for most of the comparisons in this paper, except

for some parameters that were manually defined as necessary, such as the random seed in

cisTopic, the number of top components in LSI, and the peak aggregation distance in Cicero.

Therefore, we sampled these parameters multiple times to obtain the average ARI of the

clustering results for each tool, just as we sampled the accesson number for APEC. We set the

same parameters for all the datasets as follows:

(1) *cisTopic.* The scanning range of the topic number was set to [10, 40], the number of
parallel CPUs was set to 5, and the random seed was sampled 100 times from 100 to
600 in steps of 5. We kept the topic matrices normalized by z-score and probability and
provided the performances based on both normalization methods. We then applied the
Louvain algorithm as we did in APEC to cluster cells from the normalized topic matrix
generated by cisTopic. The average ARI of the 100 clustering results from 100 random
seeds was reported.

(2) *LSI.* We performed truncated SVD (singular value decomposition) analysis on the TF-
IDF (term frequency-inverse document frequency) matrix and chose N_{SVD} top
components to generate the LSI matrix. N_{SVD} was sampled 6 times from 6 to 11. The first
component was ignored since it is always related to read depth, and the LSI scores were
capped at ±1.5. Then, we used the Louvain algorithm to cluster the cells of the LSI-
processed matrix. Its average ARI was calculated based on the 6 clustering results with
different numbers of components.

(3) *chromVAR.* The number of background iterations was set to 50, and the number of
parallel CPUs was set to 1. We then used the Louvain algorithm to cluster cells based
on the bias corrected deviation matrix generated by chromVAR. Its ARI value was
calculated based on the clustering result with the above parameters.

(4) *Cicero.* The genome window was set to 500k BPs, the normalization method was set to
"log", the number of sample regions was set to 100, the number of dimensions was set
to 40, and the peak aggregation distance was sampled at 20 values from 1k to 20k BPs
in steps of 1k BPs. Then, we used the Louvain algorithm to cluster cells based on the
aggregated model matrix generated by Cicero. Its average ARI was calculated based on
the 4 clustering results with different aggregation distances.

To test the robustness of each algorithm, we randomly sampled 20%~90% of the raw sequence
reads from the dataset of AML cells and 3 cell lines (LMPP, HL60 and monocyte) and calculated
the ARI accordingly. This random sampling experiment was performed 50 times for each
method and average ARIs were reported. The manually defined parameters for each method
were set to: APEC, 600 accessons; cisTopic, random seed 100; LSI, top 2-6 principle
components; Cicero, 10k BPs aggregation distance.

Gene score evaluated by peaks around the TSS. To evaluate the accessibility score of one
gene, we calculated the average count of all peaks around its TSS (±20000 BP) as its raw score
(S_{ij} for cell i and gene j). Then, we obtained the gene accessibility score by normalizing the raw

score ($S'_{ij} = S_{ij} * 10000/\sum_i S_{ij}$), which is in a range comparable to the gene expression from scRNA-seq data. The average score of all cells in one cluster represents the accessibility of a cell type ($\bar{S}_{kj}$ for cell cluster k and gene j). We normalized the gene score matrix $\bar{S}$ by caculating the z-score for each row and column, and the final matrix $\bar{S}_z$ represents the relative strength of the gene accessibility in each cell type. To compare the gene scoring methods between APEC and Cicero, we ran Cicero on both the mouse forebrain and thymocyte datasets. The parameter settings in Cicero were the same as described in the previous section, and the genomic distance for gene scoring was set to the default value (250k BPs).

Significant differential peaks, genes and motifs. APEC used the Student's t-test to estimate the significance of the fragment count differences between cell clusters, with P-value and fold changes, and one can determine the thresholds to identify significant differential peaks for each cluster. The significant differential genes of each cell cluster can also be acquired from the gene score ($\bar{S}_{kj}$) by the same method. To accurately quantify the enrichment of motifs on each cell, APEC applied the bias-corrected deviation algorithm from chromVAR[17]; thus, the chromVAR algorithm has been embedded into the pipeline to facilitate the calculation of the corrected deviation of the motifs. In this python version of chromVAR, permuted sampling and background deviation calculation can be run in parallel on multiple processors to reduce the computer time. The differentially enriched motifs were defined by a fold change >1 in the average motif deviation between one cluster and another.

Potential super-enhancers. Here, we defined a super-enhancer as a long continuous genomic area containing many accessible regions and have the same accessibility pattern in different cells. Many different motifs appear in one super-enhancer, therefore, the motif-based clustering method cannot reflect the critical contributions from super-enhancers for cell clustering. However, the accesson-based algorithm can group most peaks in one super-enhancer to one accesson since they always present the same accessibility pattern between cells. APEC identified super-enhancers by counting the number of peaks in a 1 million BP genomic area that belong to a same accesson. It also requires that more than 3/4 of the putative peaks in one super-enhancer be adjacent on the initial peak list. The pipeline can also aggregate bam files by cell types/clusters and convert them to BigWig format for users to upload to the UCSC genome browser for visualization.

Spatial correlation of peaks in the same accesson. To test whether peaks in the same accesson are closer in space, we integrated the Hi-C[57] data (GSE63525) and scATAC-seq[8]

data (GSE65360) on GM12878 cells. The spatial correlation of different windows, both intra-
and inter-chromosomal, can be directly extracted by Juicer[62]. The Pearson's correlation matrix
of the intra-chromosomal or inter-chromosomal windows can be calculated from the
corresponding observed/expected matrix. We constructed 600 accessons by grouping peaks in
the GM12878 scATAC-seq data in APEC, and removed the accessons that contained more
than 1000 peaks or less than 5 peaks. The width of the window was set to 500k BPs, and peaks
were then assigned to each window. Next, we collected Hi-C correlations between windows that
contained peaks in the same accesson, termed "Accesson" correlations. For comparison, we
also shuffled all peaks in different accessons to make fake accessons and re-collected the Hi-C
correlations between windows that contained peaks in each fake accesson, termed "Shuffled"
correlations. Meanwhile, we also collected Hi-C correlations between windows that contained no
peaks, termed "Non-accesson" correlations. We made boxplots for these three types of
correlations for intra-chromosomal and inter-chromosomal peaks and found that co-accessible
peaks are spatially closer to each other than random peaks.

Pseudotime trajectory constructed by APEC. As a tool to simulate the time-dependent
variation of gene expression and the cell development pathway, Monocle has been widely used
for the analysis of single-cell RNA-seq experiments[28, 63]. APEC reduced the dimension of the
accesson count matrix **M** by PCA, and then performed pseudotime analysis using the Monocle
program. For complex datasets, it is necessary to limit the number of principal components,
since too many features will cause too many branches on the pseudotime trajectory, and makes
it difficult for a user to identify the biological significance of each branch. For the hematopoietic
single cell data and thymocyte data, we used the top 5 principal components of the accesson
matrix to construct the developmental and differentiation trajectories.

Pseudotime trajectory constructed by other algorithms. To check whether other algorithms
can provide solutions to construct cell developmental pathways, we combined their transformed
count matrix with Monocle to build the pseudotime trajectory from scATAC-seq data. A similar
preprocessing method was applied to ensure the fairness of the comparison:

(1) Raw fragment count matrix. We normalized the raw count matrix **B** exactly as in the first
step of APEC, i.e., $B'_{ij} = log_2\left(\frac{B_{ij} \times 10000}{\sum_{j'} B_{ij'}} + 1\right)$, and performed PCA analysis on the
normalized matrix B'. Only the top 5 PCs were subjected to Monocle to construct the
trajectory.

(2) cisTopic. The topic matrix generated by cisTopic was normalized by making the sum of
each row the same (i.e., the probability). Then, we performed PCA analysis on the
normalized topic matrix and subjected the top 5 PCs to Monocle to build the trajectory.

(3) LSI. We chose the 2nd~6th principle components of the SVD transformation of the LSI
matrix and subjected them to Monocle to construct the trajectory.

(4) ChromVAR. After the PCA analysis of the bias corrected deviation matrix generated by
chromVAR, the top 5 PCs were combined with Monocle to construct the trajectory.

(5) Cicero. We performed PCA analysis on the aggregated matrix generated by Cicero and
used the top 5 PCs in Monocle to build the trajectory.

In addition, to confirm the reliability of the APEC + Monocle prediction of the developmental
pathway, we applied another pseudotime trajectory constructing method, SPRING[31], to the
accesson count matrix **M** from APEC to reconstruct the pathways for the hematopoietic
differentiation dataset and thymocyte developmental dataset. We performed PCA analysis of
the accesson matrix **M** and subjected the top 5 PCs to SPRING to generate the trajectories. The
number of edges per node in SPRING was set to 5.

Parameter settings for each dataset. In the quality control (QC) step, cells are filtered by two
constraints: the percentage of the fragments in peaks (P_f) and the total number of valid
fragments (N_f). However, there is no fixed cutoff for these two parameters since the quality of
different cell types and/or experiment batches are completely different. The total number of
peaks is usually limited to approximately 50000 to reduce computer time, but we recommend
using all peaks if the users want to obtain better cell clusters. (1) For the data set from
hematopoietic cells, the -log(Q-value) threshold of high-quality peaks was set to 35 to retain
54212 peaks, and the cutoff values of P_f and N_f were 0.1 and 1000, respectively. (2) For the
scATAC-seq data on the two types of cells from 2 AML patients (P1-LSC, P1-Blast, P2-LSC,
P2-Blast), the threshold of -log(Q-value) was set to 5 to retain 38683 high-quality peaks for
subsequent processing. When LMPPs, HL60 and monocytes were added to this dataset with
the AML cells, the threshold of -log(Q-value) was set to 8 to retain 42139 peaks. In the QC step,
we set the P_f cutoff to 0.05 and the N_f cutoff to 800. (3) For the snATAC-seq data from the adult
mouse forebrain, all peaks and the raw count matrix obtained from the original data source were
adopted in the analysis. (4) For the ftATAC-seq data from thymocytes, all 130685 peaks called
by MACS2 were reserved for the fragment count matrix (Q-value<0.05), and we retained cells
with P_f >0.2 and N_f >2000.

SMART-seq data analysis with Seurat. For the analysis of SMART-seq data from mouse thymocytes, we employed STAR (version 2.5.2a) with the ratio of mismatches to mapped length (outFilterMismatchNoverLmax) less than or equal to 0.05, translated output alignments into transcript coordinates (i.e., quantMode TranscriptomeSAM) for mapping[64] (Dobin et al., 2013) and used RSEM[65] (Bo et al., 2011) to calculate the TPM of genes. For QC, we excluded cells in which fewer than 2000 genes were detected and genes that were expressed in only 3 or fewer cells. Seurat filtered cells with several specific parameters to limit the number of genes detected in each cell to 2000~6000 and the proportion of mitochondrial genes in each cell was set to less than 0.4 (i.e., low.thresholds=c(2000, Inf), high.thresholds=c(6000, 0.4)). Additionally, the top 12 principal components were used for dimension reduction with a resolution of 3.2 (dims.use =1:12, resolution=3.2), followed by cell clustering and differential expressed gene analysis[66].

Association of cell clusters from scATAC-seq and scRNA-seq data. To determine the association between cell clusters from the epigenomics and transcriptomic sequencing, we calculated the P-values of Fisher's exact test of marker/non-marker genes between each pair of cell clusters from scATAC-seq and scRNA-seq data. For example, for cell cluster a from ftATAC-seq and cell cluster b from SMART-seq, if the number of consensus marker genes in both cluster a and b is G_{11}, the number of genes that are not markers in either cluster a or b is G_{22}, and the number of markers in only cluster a (or cluster b) is G_{12} (or G_{21}), then the 2×2 matrix **G** can be directly used for Fisher's exact test to evaluate the P-valuie A_{ab} between cluster a and b. After constructing a matrix **A** filled with the negative logarithm of A_{ab} for ftATAC-seq cluster a and SMART-seq cluster b, we calculated the z-score for each row and column of **A** to determine the correlation between cell clusters from different sequencing experiments.

GO term analysis of cells along pseudotime trajectory. We defined the functional characteristics of each accesson by the GO terms and motifs enriched on its peaks. The GO terms of an accesson were obtained by submitting all of its peaks to the GREAT website[67]. The negative logarithm of the P-value of each GO term in each accesson was filled into a (GO terms) × (accessons) matrix **L**. The significance of each GO term on each cell was evaluated by the product of the matrix **L** and the accesson count matrix **M**, i.e.

$$GO_{ij} = \sum_k L_{ik} \cdot M_{kj}$$

where i is the ith GO term, j is the jth cell, and k is the kth accesson. Then we calculated the z-

score for each row of this product matrix, and plotted the z-score as the GO-term score on the

trajectory diagram.

Motif enrichment of cells along pseudotime trajectory. To assess the motif enrichment of

the accessons, we used the Centrimo tool of the MEME suite[68] to search for the enriched motifs

for the peaks of each accesson and applied the same algorithm as to the GO term score to

obtain the motif score. The negative logarithm of the E-value (product of adjusted P-value and

motif number)[68] of each motif in each accesson was used to construct a (motifs) × (accessons)

matrix **F**. The enrichment of each motif on each cell was evaluated by the product of the matrix

F and the accesson count matrix **M**, i.e.,

$$Motif_{ij} = \sum_k F_{ik} \cdot M_{kj}$$

915   where i is the ith motif, j is the jth cell, and k is the kth accesson. Then, we calculated the z-

916   score for each row of this product matrix and plotted the z-score as the motif score on the

917   trajectory diagram.

919   # REFERENCES

920   58.   Picelli, S. et al. Full-length RNA-seq from single cells using Smart-seq2. *Nat Protoc* **9**, 171-181
921   (2014).
922   59.   Zuo, Z. et al. ATAC-pipe: general analysis of genome-wide chromatin accessibility. *Briefings in*
923   *Bioinformatics*, bby056-bby056 (2018).
924   60.   Shaffer, S.M. et al. Rare cell variability and drug-induced reprogramming as a mode of cancer
925   drug resistance. *Nature* **546**, 431-435 (2017).
926   61.   Huang, M. et al. SAVER: gene expression recovery for single-cell RNA sequencing. *Nat Methods*
927   **15**, 539-542 (2018).
928   62.   Durand, N.C. et al. Juicer Provides a One-Click System for Analyzing Loop-Resolution Hi-C
929   Experiments. *Cell Syst* **3**, 95-98 (2016).
930   63.   Qiu, X. et al. Single-cell mRNA quantification and differential analysis with Census. *Nat Methods*
931   **14**, 309-315 (2017).
932   64.   Dobin, A. et al. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21 (2013).
933   65.   Li, B. & Dewey, C.N. RSEM: accurate transcript quantification from RNA-Seq data with or without
934   a reference genome. *Bmc Bioinformatics* **12**, 323 (2011).
935   66.   Macosko, E.Z. et al. Highly Parallel Genome-wide Expression Profiling of Individual Cells Using
936   Nanoliter Droplets. *Cell* **161**, 1202-1214 (2015).
937   67.   McLean, C.Y. et al. GREAT improves functional interpretation of cis-regulatory regions. *Nat*
938   *Biotechnol* **28**, 495-501 (2010).
939   68.   Bailey, T.L. et al. MEME SUITE: tools for motif discovery and searching. *Nucleic Acids Res* **37**,
940   W202-208 (2009).

Supplementary Figures

Supplementary Figure 1. Quality control diagrams generated by APEC for the scATAC-seq datasets. (**a**) Number of final mapped reads and percentage of fragments in peaks for each cell in the human hematopoietic dataset. (**b**) Average count of scATAC-seq insertions around TSS regions in the same dataset. (**c**) Statistical distribution of fragment lengths in the same dataset. (**d-f**) Quality control diagrams, as in (a-c), for the blast and LSC cells from two AML patients. (**g-i**) Quality control diagrams, as in (a-c), for the cells from AML patients and three cell lines (LMPP, HL60 and Monocyte).

Supplementary Figure 2. Clustering performance of the dimension-transformed matrices generated by different algorithms. (**a**) The tSNE diagrams of the cells from AML patients and 3 cell lines. Different algorithms provided different dimension-transformed matrices for tSNE analysis, i.e., APEC: accesson matrix; cisTopic: topic matrix; LSI: LSI matrix; chromVAR: bias corrected deviation matrix; Cicero: aggregated model matrix. The table below the diagrams contains the average ARI of the cell clustering results for each algorithm. (**b**) The tSNE diagrams and ARI table for the two cell types from AML patients only, as in (a). (**c-d**) Box-plots showing the ARI values for the clustering of human hematopoietic cells (c) and the blast and LSC cells from two AML patients (d). We sampled different parameters for different algorithms. APEC: 100 different accesson numbers from 500 to 1500 in steps of 10; cisTopic: 100 different random seeds from 100 to 600 in steps of 5; LSI: 6 different numbers of top SVD components from 6 to 11; Cicero: 20 different genomic distances from 1k BPs to 20k BPs in steps of 1k BPs; chromVAR: no sampling. Z-score and probability denote different methods of normalizing the dimension-transformed matrices. Center line, median; box limits, upper and lower quartiles; whiskers, 1.5x interquartile range; points, outliers. (**e**) The average ARI values calculated by down-sampling 50 times from the raw data of the AML cells and three cell lines for each method. The X-axis represents the percentage of down-sampled sequencing reads. Shaded error band: 95% confidence interval.

Supplementary Figure 3. Super-enhancers predicted by APEC for the scATAC-seq data of cells from AML patients. (**a-b**) The genome browser track shows the aggregated scATAC-seq signal of the super-enhancer of P1-LSC cells upstream of *N4BP1* (a) and *GPHN* (b). (**c-d**) The motifs associated with peaks in the super-enhancer upstream of *N4BP1* (c) and *GPHN* (d).

Supplementary Figure 4. Biological insights of the accesson and stability and scalability

analysis of APEC. (**a**) Box plot showing the average spatial distance between peaks in the same

accesson (from scATAC-seq) versus randomly shuffled peaks versus non-accessible genomic

regions. Spatial distance was estimated from chromosome conformation capture (Hi-C)

technology. Both Hi-C and scATAC-seq data were generated from the same cell line GM12878.

Left panel: intra-chromosomal correlation of windows in the Hi-C data; right panel: inter-

chromosomal correlation of windows in the Hi-C data. Accesson: The correlation between two

windows that contain peaks in the same accesson; Shuffled: The correlation calculated by

randomly shuffling peaks in each accesson; Non-accesson: The correlation between two

windows that contain no peaks. Center line, median; box limits, upper and lower quartiles;

whiskers, 1.5x interquartile range; points, outliers. (**b**) The Hi-C profile of windows between

chr1:500,000-21,500,000. The black bars below the Hi-C track denote peaks in the same

accesson from APEC. Dotted boxes indicate examples of peaks in the same accesson that are

distant in genomic position but close in space. (**c**) The computing time required for different

algorithms to cluster cell numbers from 10,000 to 80,000 to test the scalability of each method.

The data were sampled from the single-cell atlas of *in vivo* mammalian chromatin accessibility.

cisTopic was performed using 5 CPU threads and all the other tools with 1 CPU thread. (**d-f**)

The ARI values of the clustering results that used different numbers of accessons (d), nearest

neighbors (e), and principle components (f). The dataset includes the cells from two AML

patients and three cell lines. Default values are noted in red.

Supplementary Figure 5. (**a-b**) Intensity of the accessons associated with signature genes of

excitatory (Excl) and inhibitory (Int) subtypes. The subtypes listed in parentheses were defined

by the signature genes in the results from inDrops-seq data[26].

Supplementary Figure 6. The clustering and cell-type classification of the mouse forebrain

dataset by chromVAR and Cicero. (**a**) The mouse forebrain cells clustered by chromVAR,

illustrated in the tSNE diagram. (**b**) The z-scores of the average gene scores of cell clusters

generated by chromVAR. The gene scores were calculated by APEC. (**c**) The hierarchical

clustering of the correlation between cell clusters generated by chromVAR. The correlations

were calculated based on the gene scores generated by APEC. (**d**) The mouse forebrain cells

clustered by Cicero, illustrated in the tSNE diagram. (**e**) The z-scores of the average gene

scores of cell clusters generated by Cicero. The gene scores were calculated by Cicero. (**f**) The

hierarchical clustering of the correlation between cell clusters generated by Cicero. The

correlations were calculated based on the gene scores generated by Cicero. (**g**) The z-scores of

the average gene scores of cell clusters generated by APEC. The gene scores were calculated
by Cicero. (**h**) The hierarchical clustering of the correlation between cell clusters generated by
APEC. The correlations were calculated based on the gene scores generated by Cicero.

Supplementary Figure 7. UCSC genome browser track diagram of the normalized fragment
count around gene *CD34* for each hematopoietic cell type.

Supplementary Figure 8. Cell differentiation trajectories of the human hematopoietic dataset
constructed by different algorithms. (**a-e**) The pseudotime trajectories constructed by the
combination of Monocle and the raw peak count matrix, the topic matrix from cisTopic, the LSI
matrix, the aggregated model matrix from Cicero, and the bias corrected deviation matrix from
chromVAR, respectively. (**f**) The pseudotime trajectory constructed by the combination of
SPRING and the accesson matrix from APEC.

Supplementary Figure 9. (**a**) Gating strategy of the mouse thymocytes in ftATAC-seq. (**b-d**)
Quality control diagrams for the mouse thymocyte data, similar to Supplementary Fig. (1a-1c).
(**e**) The z-score of correlation between the cell types from ftATAC-seq and bulk ATAC-seq data.

Supplementary Figure 10. (**a**) Selected significant motifs enriched in different thymocyte
subtypes obtained by the APEC algorithm. (**b-c**) The correlation of cell clusters from scRNA-seq
and scATAC-seq data, by gene scores generated by Cicero with genomic distance = 20k BPs
(b) and with genomic distance = 250k BPs (c).

Supplementary Figure 11. Single-cell transcriptome analysis of *Mus musculus* thymocytes
from SMART-seq. (**a**) tSNE diagram of the single-cell expression matrix of *Mus musculus*
thymocytes, labeled by the FACS index of each cell. (**b**) Louvain clustering of the same single-
cell dataset obtained by Seurat. The cell types of these clusters were classified by the
expression of corresponding marker genes. (**c**) Important marker genes were differentially
expressed in different cell clusters. (**d**) Heatmap of the expressions of all genes significantly
differentially expressed between cell clusters. The top color bar used the same scheme
described in (b) to render cells of different clusters.

Supplementary Figure 12. Developmental characteristics of single-cell samples captured by
APEC. (**a, b**) Pseudotime trajectory of scATAC-seq data from *Mus musculus* thymocytes
labeled with the FACS index and APEC cluster index. (**c**) Pseudotime trajectory constructed by
applying SPRING to the accesson matrix. The colors of cells denote their stages in the APEC
trajectory results. (**d**) Z-scores of the -log(P-value) of the GO terms along the pseudotime

trajectory of stage 1 cells. (**e**) Logarithm of the P-value of GO terms searched from peaks in

accessons ac1~ac7, which are the marker accessons of cluster DP. A1 and DP. A4/A5 of

thymocytes.