Full title: Severity Index for Suspected Arbovirus (SISA): machine learning for accurate

prediction of hospitalization in subjects suspected of arboviral infection

Short title: Severity index for hospitalization of suspected arbovirus infections

Rachel Sippy[1, 2, *], Daniel F Farrell[3], Daniel Lichtenstein[3], Ryan Nightingale[3], Megan Harris[3],

Joseph Toth[3], Paris Hantztidiamantis[3], Nicholas Usher[4], Cinthya Cueva[1], Julio Barzallo Aguilar[5],

Anthony Puthumana[3], Timothy Endy[1, 3], Sadie J. Ryan[2], Anna M. Stewart Ibarra[1, 3, *]

1. Institute for Global Health and Translational Science, SUNY Upstate Medical University,

9   Syracuse, NY, USA

10   2. Department of Geography, University of Florida, Gainesville, FL, USA

11   3. Department of Medicine, SUNY Upstate Medical University, Syracuse, NY, USA

12   4. Department of Microbiology, Cornell University, Ithaca, NY, USA

13   5. Teofilo Davila Hospital, Ministry of Health, Machala, El Oro Province, Ecuador

15   [*]Corresponding authors: sippyr@upstate.edu, stewarta@upstate.edu

Abstract

Background: Dengue, chikungunya, and Zika are arboviruses of major global health concern. Decisions regarding the clinical management of suspected arboviral infection are challenging in resource-limited settings, particularly when deciding on patient hospitalization. The objective of this study was to determine if hospitalization of individuals with suspected arboviral infections could be predicted using subject intake data. Methodology/Principal Findings: Two prediction models were developed using data from a surveillance study in Machala, a city in southern coastal Ecuador with a high burden of arboviral infections. Data was obtained from subjects who presented at sentinel medical centers with suspected arboviral infection (November 2013 to September 2017). The first prediction model—called the Severity Index for Suspected Arbovirus (SISA)—used only demographic and symptom data. The second prediction model—called the Severity Index for Suspected Arbovirus with Laboratory (SISAL)—incorporated laboratory data. These models were selected by comparing the prediction ability of seven machine learning algorithms; the area under the receiver operating characteristic curve from the prediction of a test dataset was used to select the final algorithm for each model. After eliminating those with missing data, the SISA dataset had 534 subjects, and the SISAL dataset had 98 subjects. For SISA, the best prediction algorithm was the generalized boosting model, with an AUC of 0.91. For SISAL, the best prediction algorithm was the elastic net with an AUC of 0.94. A sensitivity analysis revealed that SISA and SISAL are not directly comparable to one another. Conclusions/Significance: Both SISA and SISAL were able to predict arbovirus hospitalization with a high degree of accuracy in our dataset. These algorithms will need to be tested and validated on new data from future patients. Machine learning is a powerful prediction tool and provides an excellent option for new management tools and clinical assessment of arboviral infection.

41   **Keywords:** machine learning, arboviral disease, dengue fever, undifferentiated febrile illness

Author Summary

Patient triage is a critical decision for clinicians; patients with suspected arbovirus infection are

difficult to diagnose as symptoms can be vague and molecular testing can be expensive or

unavailable. Determining whether these patients should be hospitalized or not can be challenging,

especially in resource-limited settings. Our study included data from 543 subjects with a diagnosis

of suspected dengue, chikungunya, or Zika infection. Using a machine learning approach, we

tested the ability of seven algorithms to predict hospitalization status based on the signs,

symptoms, and laboratory data that would be available to a clinician at patient intake. Using only

signs and symptoms, we were able to predict hospitalization with high accuracy (94%). Including

laboratory data also resulted in highly accurate prediction of hospitalization (92%). This tool

should be test in future studies with new subject data. Upon further development, we envision a

simple mobile application to aid in the decision-making process for clinicians in areas with limited

resources.

Introduction:

Undifferentiated febrile illness is a common clinical scenario in tropical medicine, with a long list of potential pathogens sharing similar symptoms. Arthropod-borne viruses (arboviruses), including dengue virus (DENV), chikungunya virus (CHIKV), and Zika virus (ZIKV), share common mosquito vectors (*Aedes aegypti* and *Ae. albopictus*) and often present with fever, rash, myalgias, and arthralgias. Dengue virus is endemic in the tropical Americas, and the emergence of CHIKV in 2013 in Saint Martin and ZIKV in 2015 in Brazil has brought these arboviruses to the forefront of international attention [1–3]. There were over 400,000 cases of dengue fever in Andean Latin America in 2013 [4], with transmission risk expected to increase sharply over the next 50 years [5]. Ecuador in particular has a high burden of arboviral illness, with 86,306 total cases of dengue from 2014—2018 [6–8]. There is also a high prevalence of asymptomatic DENV infections and co-infections in southern coastal Ecuador [1,9]. In 2014, CHIKV was introduced, with 35,555 cases from 2014—2018, followed by the introduction of ZIKV in 2016, with 5,304 cases from 2016—2018 [6–8].

Clinical decision-making in the context of arboviral infection is particularly challenging in resource-limited settings such as Ecuador. For instance, there may be limited healthcare professionals relative to the high disease burden, which may impact the ability to provide optimal subject care. Ecuador has 22 physicians for every 10,000 people, though this ranges by province, from 13 to 32 physicians per 10,000 people [10]. This is just above the 1 physician per 1,000 people benchmark of the World Health Organization [11], and physicians are likely concentrated in urban areas. Moreover, molecular diagnostics are often unavailable outside of large urban centers. Of Ecuador's 4,168 healthcare establishments, 1,045 (25.1%) have a clinical laboratory [10], leaving many healthcare providers in Ecuador without these crucial diagnostic tools (*i.e.* PCR

or ELISA). These infrastructural limitations create a challenging clinical environment, especially

as healthcare providers need to determine whether a patient with suspected arboviral illness should

be hospitalized or not. Efficient and effective triaging is essential for good clinical care in

resource-limited settings [12].

Current practice in Ecuador is to hospitalize subjects suspected of dengue infection when

they exhibit any of the World Health Organization (WHO) 2009 dengue warning signs, any signs

of shock, or severe thrombocytopenia [13]. While treatment for dengue is supportive, proper

inpatient management of severe dengue can reduce mortality dramatically [14,15]. Dengue has a

wide spectrum of clinical presentations, with the majority patients recovering following a self-

limited clinical course, and a small percentage progressing to severe disease characterized by

plasma leakage. In the latter cases, prompt intravenous rehydration can reduce the case fatality rate

to less than 1% [14]. Similarly, management of CHIKV and ZIKV infections is largely supportive,

but both infecitons may result in potentially serious complications, such as adverse neonatal effects

[16–19]. Deciding whether to hospitalize a subject with a suspected DENV, CHIKV, or ZIKV

infection is thus an important clinical decision. This decision has other non-clinical and indirect

consequences, including the utilization of hospital resources that could otherwise be used for other

patients, as well as increasing the financial cost of the case when compared to less costly outpatient

care. Globally, an estimated 18% of dengue cases are admitted to the hospital, with 48% managed

as outpatients and 34% not seeking medical attention [20]. The average cost to manage a case of

dengue is tripled if the patient is hospitalized [14,20].

Machine learning is a tool that combines statistics with computer science to make efficient

use of massive data sets [21]. It differs from traditional statistical modeling (e.g. regression

models) in that there are fewer assumptions about the underlying distribution of the data and the

relationships between variables. While model interpretability is often a goal of traditional

statistical models, this is not important in machine learning. The only goal is to create highly

accurate predictions of an outcome of interest, often using as many variables as possible [22,23].

In modeling relationships with a machine learning approach, the computer incorporates

connections not obvious to human beings to successfully predict an outcome of interest. Machine

learning is applicable in many fields and has been previously used in medical applications, to

estimate clinical risk, guide triage or diagnose disease [21,24–26]. Clinical applications of machine

learning for arboviral illnesses specifically have included analysis of patient genomes for dengue

prognosis [27], scanning of patient sera for DENV [28] or Zika diagnosis [29], thermal image

scanning for detection of hemodynamic shock [30], analysis of body temperature patterns for

diagnosis of undifferentiated fever etiology [31], and analysis of patient data for dengue fever

diagnosis [25]. No studies have yet attempted to use machine learning for prediction of

hospitalization among arboviral illness or undifferentiated fever patients, although it has been used

to predict critical care and hospitalization outcomes based on emergency department triage data in

children and adults [32,33].

The objective of this study was to determine if the hospitalization of individuals with

suspected arboviral infections could be predicted using subject intake data. In this study, we take

a retrospective view of arboviral infection management in a tropical city in southern coastal

Ecuador using data from an ongoing prospective surveillance study. Using actual clinical practice

as a guide, we assessed the ability of seven machine learning algorithms to determine

hospitalization using basic symptom and demographic data that was collected via standard intake

of subjects with suspected DENV, CHIKV, or ZIKV infections. The machine learning approach

and algorithms developed here could potentially support physicians faced with complex clinical

management decisions in areas where multiple arboviruses co-circulate, such as Ecuador.

Methods

Ethics statement

This study protocol was reviewed and approved by Institutional Review Boards at the State

University of New York (SUNY) Upstate Medical University, the Luis Vernaza Hospital in

Guayaquil, Ecuador, and the Ecuadorean Ministry of Health (MoH). Clinical and demographic

data from study subjects was obtained following informed consent (and/or assent, as applicable)

per the protocol of an ongoing arboviral surveillance study in Machala, Ecuador (described

previously) [1].

Study design and data source

We conducted a retrospective analysis of data from a prospective arbovirus surveillance

study, which included subjects (age >6 months) recruited from Ecuadorean MoH clinical sites

from November 2013 to September 2017 in Machala, Ecuador. Subjects were identified as a part

of an ongoing, multi-year arbovirus surveillance project, a description of which has been published

previously [1]. Briefly, subjects were invited to enroll in the study if they presented at the reference

hospital or one of four outpatient clinics and were diagnosed with arboviral infection by MoH

physicians. In 2014 and 2015, we recruited subjects who were clinically diagnosed with dengue

fever by MoH physicians. Following the emergence of CHIKV (2014) and ZIKV (2016), the

inclusion criterion in 2016 and 2017 was expanded to include subjects clinically diagnosed with

DENV, CHIKV or ZIKV infection. At the time of enrollment, subject demographic information,

clinical history, and symptoms present during current illness were collected using a questionnaire

administered by trained study personnel. Subjects were asked about symptoms in the past 7 days:

headache, anorexia or nausea, muscle or joint pain, rash, bleeding (defined as bleeding from

respiratory, digestive, or genitourinary mucosa), rhinorrhea, vomiting, lethargy or drowsiness,

cough, abdominal pain, diarrhea, and retro-orbital pain. Conjunctivitis was later added to the

enrollment survey after the emergence of ZIKV but was not included in this analysis. Laboratory

data (hematocrit, white blood cell count, neutrophils, lymphocytes, and platelet count) was

collected at the time of enrollment if the subject had copies of recent laboratory evaluation (for

outpatients), or the first labs on admission to the hospital (for hospitalized subjects) were used.

Additional laboratory data were available for hospitalized subjects, but analysis was limited to the

aforementioned three parameters as these were consistently available among a subset of the non-

hospitalized subjects. Laboratory arboviral diagnostic data were not included, as these data are

often not available at the time that a physician decides whether or not to hospitalize a patient, and

we utilized only rely the data that would realistically be available. Data from enrollment surveys

was used for the analysis of the non-hospitalized outpatients in the current study. Laboratory data

on the hospitalized subjects was verified by review of medical records and managed using

REDCap software [34] hosted at SUNY Upstate Medical University.

Exclusion criteria

Hospitalized subjects whose physical medical records could not be located and subjects

with incomplete enrollment survey data (*i.e.* missing hospitalization status, symptom survey

questions) were excluded. The subset of the non-hospitalized subjects who had available

laboratory data were included in a second analysis with the same hospitalized cohort, all of whom

had available laboratory data.

Statistical analysis

The outcome variable was hospitalization status. Variables of interest included

demographic data, presenting symptoms, past medical history, and laboratory data (hematocrit,

white blood cell count, neutrophils, lymphocytes, and platelet count). A prediction algorithm was

developed using demographic and symptom data only (28 total predictors), called the Severity

Index for Suspected Arbovirus (SISA, in Spanish the *Severidad de Infecciones Sospechosas por*

Arbovirus). A second prediction algorithm was developed using demographic, symptom, and

laboratory data (33 total predictors), called the Severity Index for Suspected Arbovirus with

Laboratory (SISAL, in Spanish the *Severidad de Infecciones Sospechosas por Arbovirus con datos*

del Laboratorio). Characteristics for hospitalized and non-hospitalized subjects among these

subject groups were compared using a two-sample t-test (continuous) or Fisher's exact test

(categorical).

In machine learning, 10-fold cross-validation with holdout data results in an unbiased

estimate of model validity and accuracy [22,35]. Thus, our datasets were divided by random

sampling into training and testing (holdout) data sets. For SISA, the training set was 85% of the

full dataset and the testing set was the remaining 15%. For SISAL, the training set was 70% of the

full dataset and the testing set was the remaining 30%. With the training dataset, we used repeated

10-fold cross validation to test the ability of six algorithms with diverse statistical approaches—

bagged trees (bags) [36], *k* nearest neighbors regression (knn) [37], random forest [38], elastic net

regression [39], generalized boosting models (gbm) [40], and neural networks [41]—to predict

hospitalization. Because we have no prior assumptions about the nature of the relationship between

the available predictors and the outcome, we use a variety statistical approaches to improve the

likelihood that we will find an algorithm that works well with these data. Following a published

criticism of machine learning prediction compared to logistic regression [42], we added logistic

regression to our list of algorithms to test (seven total algorithms). For models with tuning

parameters (knn, random forest, elastic net, and gbm), tuning was performed using another layer

of 10-fold cross-validation [43]. Each algorithm was then used to predict hospitalization outcomes

within the testing dataset. Measures of discrimination [44], including accuracy, Cohen's kappa,

and area under the curve (AUC) for the receiver operating characteristic (ROC) were calculated

for each of the ten cross-validations and summarized as mean values. The best algorithm for each

dataset was chosen based on the highest AUC as calculated from the test set. Model residual plots

were examined. Data analysis and visualization was performed using SAS version 9.2 (SAS

Institute, Cary, NC) and R version 3.2.2 (R Foundation for Statistical Computing, Vienna, Austria)

including packages haven [45], caret [37], MASS [41], ipred [36], randomForest [38], elasticnet

[39], gbm [40], nnet [41], mgcv [46,47], kernlab [48], glmnet [49], and pROC [50].

We compared the prediction abilities of SISA versus SISAL to assess whether laboratory

data could improve our ability to predict subject hospitalization status. Because there may be some

selection bias for subjects with available laboratory data (e.g. more severe symptoms, more similar

subject data, or different socioeconomic status compared to typical patients with clinical arboviral

diagnosis), the subject groups in SISA and SISAL may not be exchangeable [51]. We performed

a sensitivity analysis to determine if the selected algorithm and prediction ability of SISA is the

same when using all SISA subjects or SISAL subjects (without laboratory data) for the training

and testing steps.

Results

General Characteristics

Between November 20, 2013 to September 13, 2017, 592 subjects were recruited into the

arboviral surveillance study. After exclusions (Figure 1), 534 subjects were included in the dataset

for SISA, of which 59 were hospitalized and 475 were not hospitalized. The SISA training dataset

included 455 subjects and the test dataset included 79 subjects. The SISAL dataset included 98

subjects, of which 59 were hospitalized and 39 were outpatients. The SISAL training dataset

included 70 subjects and the test dataset included 28 subjects. Demographics and symptoms for

the two datasets are in Table 1. Presenting temperature was higher, and presence of mucosal

bleeding, vomiting, and abdominal pain were significantly more common in hospitalized subjects

in the SISA dataset.

Table 1. Overview of demographics, symptoms, and laboratory values for subjects enrolled in the

study (n=543). Numerical data are shown as means and were analyzed with Welch 2-Sample T-

test. Categorical data shown as percentages and analyzed with Fisher's Exact Test. Bold p-values

indicate $p<0.05$.

	SISA (N=543)			SISAL (N=98)		
	Hospitalized (n=59)	Outpatient (n=484)	p-value	Hospitalized (n=59)	Outpatient (n=39)	p-value
Age (years)	23.3	25	0.38	23.3	22.3	0.75
Height (cm)	154.2	147.9	0.01	154.2	154.2	0.99
Weight (kg)	58.2[A]	56.5	0.61	58.2[A]	57	0.78
MUA Circumference (cm)	25.4	25.9	0.5	25.4	25.9	0.64
Waist Circumference (cm)	83.9	78.4	0.05	83.9	78	0.1
Temperature (ºC)	37.7	37.4	0.01	37.7	37.2	0.02
Hematocrit (%)				37.6	39.8	0.02
WBC Count (cells/µL)				5435	6863	0.18
Neutrophils (%)				58.1	54.5	0.38
Lymphocytes (%)				30.6	33.5	0.38
Platelet Count				117627	203333	<0.01
Gender (% female)	63	54	0.27	63	51	0.30
Fever in past 7 days	97	94[C]	0.56	97	97	1.00
Head pain	64	79[A]	0.01	64	82	0.07
Nausea	54	53[A]	0.09	54	59	0.68
Muscle or joint pain	70	81[A]	0.04	70	82	0.24
Rash	20	29[A]	0.22	20	15	0.60
Bleeding	24	4[A]	<0.01	24	5	0.02
Rhinorrhea	19	26[A]	0.27	19	21	1.00
Vomiting	56	27[B]	<0.01	56	31	0.02
Drowsiness or lethargy	80	84[B]	0.46	80	92	0.15
Coughing	37	35[B]	0.77	37	26	0.27
Abdominal pain	70	50[B]	<0.01	70	56	0.20
Diarrhea	29	22[B]	0.33	29	44	0.19
Retroorbital pain	56	66[C]	0.15	56	77	0.05
Positive tourniquet test	17	3[D]	<0.01	17	5	0.01
History of allergies	20	20[A]	0.86	20	18	0.80
History of hypertension	5	6[A]	1.00	5	10	0.43
History of asthma	2	4[A]	0.71	2	3	1.00
History of cancer	0	2[A]	0.61	0	0	1.00
History of diabetes	2	3[A]	1.00	2	0	1.00
History of dengue in household	17	12[A]	0.30	17	5	0.12
History of dengue	19	22[A]	0.74	19	18	1.00
Pregnancy (self-reported)	19	2	<0.01	19	3	0.22

SISA= Severity Index for Suspected Arbovirus or Severidad de Infeccion Sospechosa de

Arbovirus, SISAL= Severity Index for Suspected Arbovirus with Laboratory or Severidad de

Infeccion Sospechosa de Arbovirus con Laboratorio, cm=centimeters, kg=kilograms, MUA=mid-

upper arm, °C=degrees Celsius, WBC=white blood cell, µL=microliters. [A] Missing n=1, [B] Missing

n=2, [C] Missing n=3, [D] Missing n=4, % Pregnant is taken from the total population (male and

female)

Prediction of Hospitalization Status

Accuracy, Cohen's kappa, and AUC for the training and test sets are shown in Figure 2. For SISA, using only symptoms and demographics, generalized boosting model, elastic net, neural networks, and logistic regression performed well with the test set (accuracy: 89.8—96.2%, Cohen's kappa: 0.00—0.77, AUC: 0.50—0.91). The generalized boosting model algorithm was found to have the best AUC (0.91) among the test dataset and was the second-best algorithm in the training set. The sensitivity for this algorithm was 95.8%, and the specificity was 87.5% when predicting hospitalization of subjects in the test dataset. The calibration plot for this prediction is in Figure 3. The variables with the greatest influence on the final SISA model were drowsiness, bleeding, and temperature.

Results for SISAL, where laboratory parameters were included as well as symptoms and demographics, are in Figure 4. All models except neural networks and k nearest neighbors performed well with the test set (accuracy: 64.3—92.6%, Cohen's kappa: 0.25—0.85, AUC: 0.62—0.94). The elastic net algorithm had the best AUC (0.94) among the test dataset and was the third-best algorithm in the training set. The sensitivity for SISAL was 100% and the specificity was 88.2% when predicting hospitalization of subjects in the test dataset. The calibration plot for this prediction is in Figure 5. The variables with the greatest influence on the final SISAL model were drowsiness, orbital pain, and platelet count.

The results for SISA when trained with the SISAL subjects (without laboratory data) are in the S1 Figure. All models except k nearest neighbors and logistic regression performed well (test set accuracy: 53.5—92.9%, Cohen's kappa: 0.04—0.86, AUC: 0.51—0.94). The bagged trees, random forest, generalized boosting models, and elastic net algorithm had identical AUC (0.94). The sensitivity was 100% and the specificity was 88.2% when predicting hospitalization

of subjects in the test dataset. If the SISA and SISAL subjects were exchangeable, we would expect

the SISAL subject group (without laboratory data) to produce the same results as the SISA

analysis. Because these results differ from those obtained in the SISA analysis, we conclude that

the SISAL subjects are not exchangeable with the SISA subjects.

Discussion

Suspected arboviral infections impose large health and financial burdens on populations in

which the diseases are endemic. In 2013, the estimated global cost of dengue illness was US $8—

9 billion [20]. In many arbovirus endemic regions, DENV, CHIK, and ZIKV infections are

diagnosed based on clinical presentation and basic laboratory results, which can be difficult due to

nonspecific symptoms and limited availability of definitive diagnostic tools [52]. In this study, we

demonstrate that our machine learning algorithms were able to predict hospitalization status among

our cohort of subjects with suspected arboviral illness with up to 96% accuracy using only

symptom and demographic data. We thus describe the early development of a new tool,

SISA/SISAL, which in the future may be utilized by clinicians in resource-limited settings when

triaging subjects with suspected arboviral illness.

In our cohort, we found that hospitalized cases had statistically significant - though

clinically insignificant - elevations in temperature at presentation in both SISA and SISAL. This

demonstrates an algorithm's ability to make use of data that might be overlooked by the clinician.

In the SISA dataset, mucosal bleeding, vomiting, and abdominal pain were more common in

hospitalized subjects than in outpatients. In the SISAL dataset, while hospitalized subjects

experienced more mucosal bleeding and vomiting than outpatients, presence of abdominal pain

did not differ between groups. This could suggest that the outpatient subjects who were sent for

laboratory testing represented cases of serious concern, as abdominal pain may qualify those with

suspected or confirmed dengue for hospital admission [13,14]. For SISAL, hospitalized subjects

had lower hematocrit and platelet counts when compared to non-hospitalized subjects, with the

lower platelet counts to be expected in hospitalized dengue cases. The WBC counts showed no

significant difference, which could be due to leukocytosis or leukopenia obscuring the mean

difference between hospitalized and non-hospitalized subjects.

Our sensitivity analysis revealed that SISA produced different results when its

training/testing dataset was restricted to those subjects with laboratory data available. This result

is unsurprising, as we expect that selection bias is contributing to the subjects available for the

SISA and SISAL datasets. All hospitalized subjects had laboratory data available, and we would

additionally expect that subjects with laboratory data had some signs or symptoms that would

prompt the attending physician to request laboratory diagnostics, setting them apart from subjects

without laboratory data. These signs and symptoms are also likely linked to whether subjects were

eventually hospitalized or not, meaning these groups of subjects are not directly comparable to one

another. When we used the SISA approach (symptoms and demographics only) for a dataset

comprised of the SISAL group of subjects (without laboratory data), we found that the AUC was

identical to the AUC from the SISAL approach. This would suggest that we are unable to improve

our prediction of hospitalization status by using subject laboratory data. The AUC was higher for

this group of subjects, but these improvements are likely due to fundamental differences between

the SISA/SISAL groups of subjects. These patient groups should continue to be analyzed with

separate algorithms.

This is the first use of machine learning to predict hospitalization status of subjects with

clinically diagnosed arboviral infections. Our models exhibit high accuracy, sensitivity, and

specificity in a region with a high burden of co-circulating of DENV, CHIKV, and ZIKV. These

algorithms, particularly SISA, use information that could easily be obtained in resource-limited

settings, suggesting the potential to develop a useful tool for clinicians. Our model's accuracy is

consistent with tools previously reported in the literature. Past predictive modeling of disease with

a machine learning approach had been efficacious in the diagnosis of pneumonia (95% sensitivity,

96.5% specificity), dengue (70% sensitivity, 80% specificity), hepatitis (96% accuracy), and

tuberculosis (95% accuracy) using clinical and laboratory parameters [25,53–55].

There has been criticism regarding the use of machine learning in prediction models: a

recent systematic review found that machine learning predictions had no advantage over logistic

regression predictions on average [42]. Christodoulou *et al*. do an excellent job of outlining some

common missteps in the use of machine learning for prediction and the somewhat alarming lack

of transparency in many published machine learning prediction models. We agree with many of

the assertions made by the authors and strive to improve reporting and validation in our own work

as a result. However, in this specific study, we did not find that logistic regression performed better

than other algorithms. Our overall approach differs from that of most machine learning papers in

that we did not assume that one particular algorithm would have superior prediction abilities for

our data, but rigorously compared multiple algorithms with the goal of finding an algorithm that

functions well with our predictors and outcome of interest, to be further validated with a new

dataset in future research. We have no illusions about the potential lack of generalizability of our

data and caution against any strong conclusions about the future utility SISA/SISAL in predicting

hospitalization status for future patients. We present the current study as preliminary yet promising

results in the development of a future tool that will need additional, vigorous validation in future

sets of subject data before use in the real world.

Numerous studies have looked at clinical and laboratory findings specific to certain

arbovirus diagnoses, yet few have proposed tools that can aid in management of unconfirmed

febrile illness [56–59]. A study in Puerto Rico of acute febrile illness emergency room cases found

the tourniquet test and leukopenia to be predictive of dengue diagnosis, yet dengue was confirmed

in only 11% of their total 284 cases [60]. In Thailand, fever, positive tourniquet test and leukopenia

differentiated confirmed dengue from other febrile illness initially suspected as dengue [61]. Also

in Thailand, among a sample of 172 children with acute fever without obvious cause, those with

dengue had several laboratory parameters that differentiated them from the other febrile illness

[52]. While these studies were able to distinguish dengue from other acute febrile illness, they

highlight the large proportion of cases that do not get a confirmed diagnosis and have not moved

beyond initial reports to demonstrate predictive abilities. With SISA/SISAL, the approach is more

empirical. Clinical diagnosis of DENV, CHIKV, or ZIKV was a starting point for the machine

learning used here. Given that timely laboratory diagnostics may not be available, grouping these

suspected subjects reflects the reality that physicians face in the clinic in arbovirus-endemic

regions. That such a model can accurately predict hospitalization outcome suggests that

SISA/SISAL could be expanded to undifferentiated fever. The ability of machine learning models

to predict hospital admission outcomes using only emergency department triage data lends support

to expanding our approach to undifferentiated fever [32,33]. Of the suspected arboviral cases

analyzed here, approximately 54% were confirmed as acute or recent DENV infection, 17% had

acute CHIKV infection, and 29% had no pathogen identified (based on analysis of subjects in 2014

and 2015) [1]. Results of the 2016 and 2017 subject samples are pending, but preliminary PCR

testing suggests predominance of CHIKV in 2016 and ZIKV in 2017.

Clinicians rely on tools to help make decisions about patient management, and simple tools

are can benefit physicians in limited-resource settings [62,63]. Smart phones are commonly used

in Ecuador and mobile health tools are a great option for physicians, with several popular apps that

include various triage rules and scores, such as MDCalc [64]. After further development and

validation of our algorithmic approach, and evaluation of its potential benefit in the clinic, we

conceive of its inclusion in a user-friendly mobile application to aid in the decision to hospitalize

patients with undifferentiated fever.

Limitations

The variables with the greatest influence on the final SISA model were drowsiness,

temperature, and nausea; for the SISAL model they were drowsiness, orbital pain, and platelet

count. An important caveat inherent to the nature of machine learning is that the exact weight of

each variable in the final prediction model is difficult to assess and interpret, thus we cannot

propose a causal relationship or correlation between these variables and our outcome of

hospitalization.

The SISA/SISAL models are presented here in the first iteration of their use. They have

not yet been validated beyond the current datasets, but the use of holdout data and 10-fold cross-

validation provides us with an unbiased estimate of model validity as well as prediction accuracy.

An external validation of these algorithms with a new dataset is ongoing, as well as the testing of

fewer prediction variables with the eventual goal of an easy-to-use online or mobile app for use in

the clinic.

In this study, we used the outcome of subject hospitalization for both prediction models.

The sensitivity and specificity of SISA/SISAL relies on the assumption that the subjects in this

dataset were correctly hospitalized. It is possible that some subjects were treated as outpatients

when they should have been hospitalized, or that some subjects were hospitalized unnecessarily.

For subjects that were incorrectly treated as outpatients, it is unlikely that the subject would not

have returned to a clinic to receive care, as their symptoms would likely drive them to do so.

Because our collection of medical records was retrospective, we were able to capture subject

hospitalization at any point, even if they were initially treated as outpatients. Hospital Teofilo

Davila is the reference MoH hospital in the province, and it is unlikely that these subjects would

have sought care at a hospital elsewhere. It is possible for some subjects to have been hospitalized

unnecessarily; we have no way of identifying these subjects or truly knowing if it was safe for

these subjects to have been treated as outpatients. As a result, our algorithms could thus

recommend hospitalization unnecessarily. Although hospitalization could place undue financial

burden on some patients and the health system, failure to hospitalize a serious case could results

in grave consequences and we would prefer to take a cautious approach in hospitalization decision-

making. Moreover, these algorithms are merely intended as a tool to inform clinical judgement,

not to replace important clinical triage decisions [65].

The time period during which our data were collected (2014—2017) included the

emergence of two important new arboviruse—CHIKV and ZIKV. The potential severity of these

infections and their novelty may have increased the number of patients willing to be hospitalized

or to seek healthcare in the first place. With ZIKV infection, physicians may have been more likely

to hospitalize pregnant women. This may limit the generalizability of SISA/SISAL in future

subject datasets, though as viral diseases continue to emerge globally, it is important to test the

ability of decision-making tools to function under these dynamic scenarios. For new diseases with

clear warning signs for potentially severe disease, we would expect SISA/SISAL to work well.

Conclusions

Clinicians in resource-limited settings commonly encounter subjects with a suspected

diagnosis of DENV, CHIKV, or ZIKV infection and often have limited tools at their disposal. A

subject may be unable or unwilling to provide a laboratory specimen, and diagnostic testing may

not always be available. The SISA/SISAL models are promising clinical tools, given the high

sensitivity and specificity for both models. Machine learning, if used thoughtfully, can be a

powerful method for building such prediction models, making the best use of real-world available

clinical data.

Acknowledgements

Many thanks to the Ministry of Health of Ecuador and SUNY Upstate's Institute for Global Health

and Translational Science, especially Tina Lupone, as well as the Upstate team in Machala.

Competing Interests

The authors have no competing interests to declare.

Supporting information captions

S1 Figure: SISA Analysis of SISAL Dataset Accuracy (blue), Cohen's kappa (red), and AUC

(green) were calculated for the 10-fold cross validation (left) and the holdout test dataset (right)

for prediction of hospitalization status in clinically diagnosed DENV, CHIKV or ZIKV infections.

bag=bagged trees, knn=k nearest neighbors, rf=random forest, gbm=generalized boosting models,

enet=elastic net, nnet=neural networks, log=logistic regression, DENV=dengue virus,

CHIKV=chikungunya virus, ZIKV=Zika virus

References

1. Stewart-Ibarra AM, Ryan SJ, Kenneson A, King CA, Abbott M, Barbachano-Guerrero A, et al. The Burden of Dengue Fever and Chikungunya in Southern Coastal Ecuador: Epidemiology, Clinical Presentation, and Phylogenetics from the First Two Years of a Prospective Study. Am J Trop Med Hyg. 2018;98: 1444–1459. doi:10.4269/ajtmh.17-0762

2. Staples JE, Fischer M. Chikungunya Virus in the Americas — What a Vectorborne Pathogen Can Do. N Engl J Med. 2014;371: 887–889. doi:10.1056/NEJMp1407698

3. Zanluca C, de Melo VCA, Mosimann ALP, dos Santos GIV, dos Santos CND, Luz K. First report of autochthonous transmission of Zika virus in Brazil. Mem Inst Oswaldo Cruz. 2015;110: 569–572. doi:10.1590/0074-02760150192

4. Stanaway JD, Shepard DS, Undurraga EA, Halasa YA, Coffeng LE, Brady OJ, et al. The global burden of dengue: an analysis from the Global Burden of Disease Study 2013. Lancet Infect Dis. 2016;16: 712–723. doi:10.1016/S1473-3099(16)00026-8

5. Ryan SJ, Carlson CJ, Mordecai EA, Johnson LR. Global expansion and redistribution of Aedes-borne virus transmission risk with climate change. PLoS Negl Trop Dis. 2019;13: e0007213. doi:https://doi.org/10.1371/journal.pntd.0007213

6. Ministerio de Salud Publica, Dirección Nacional de Vigilancia Epidemiológica. Enfermedades Trasmitidas por Vectores 1994-2017 [Internet]. Dirección Nacional de Vigilancia Epidemiológica, Ministerio de Salud Publica.; Available: https://public.tableau.com/profile/vvicentee80#!/vizhome/EnfermeddaesTropicales_vectoriales-2014/ANUARIO

7. Ministerio de Salud Publica, Dirección Nacional de Vigilancia Epidemiológica. Enfermedades Trasmitidas por Vectores SE 52/2018 [Internet]. Available: https://www.salud.gob.ec/wp-content/uploads/2019/01/GACETA-VECTORES-SE-52.pdf

8. Ministerio de Salud Publica, Dirección Nacional de Vigilancia Epidemiológica. Enfermedades Trasmitidas por Vectores SE 52/2017 [Internet]. Available: https://www.salud.gob.ec/wp-content/uploads/2017/07/Gaceta-Vectorial-SE52.pdf

9. Farrell DF, Lupone CD, Kenneson A, Cueva C, Heydari N, Barzallo Aguilera JH, et al. Case Report: An Acute Chikungunya Infection and a Recent Secondary Dengue Infection in a Peripartum Case in Ecuador. Am J Trop Med Hyg. 2018;98: 838–840. doi:10.4269/ajtmh.17-0781

10. Censos IN de E y. Actividades y Recursos de Salud. In: Instituto Nacional de Estadística y Censos [Internet]. [cited 28 Mar 2019]. Available: http://www.ecuadorencifras.gob.ec/actividades-y-recursos-de-salud/

11. WHO | Density of physicians (total number per 1000 population, latest available year)
[Internet]. [cited 11 Apr 2019]. Available:
https://www.who.int/gho/health_workforce/physicians_density/en/

12. Agyeman-Duah JNA, Theurer A, Munthali C, Alide N, Neuhann F. Understanding the
barriers to setting up a healthcare quality improvement process in resource-limited settings:
a situational analysis at the Medical Department of Kamuzu Central Hospital in Lilongwe,
Malawi. BMC Health Serv Res. 2014;14: 1. doi:10.1186/1472-6963-14-1

13. Dengue, Guías de atención para enfermos en la región de las Américas – Ministerio de
Salud Pública [Internet]. [cited 3 Apr 2019]. Available: https://www.salud.gob.ec/dengue-
guias-de-atencion-para-enfermos-en-la-region-de-las-americas/

14. WHO | Dengue guidelines for diagnosis, treatment, prevention and control: new edition. In:
WHO [Internet]. [cited 8 Jul 2017]. Available:
http://www.who.int/rpc/guidelines/9789241547871/en/

15. Dengue and severe dengue [Internet]. [cited 28 Mar 2019]. Available:
https://www.who.int/news-room/fact-sheets/detail/dengue-and-severe-dengue

16. Gérardin P, Barau G, Michault A, Bintner M, Randrianaivo H, Choker G, et al.
Multidisciplinary Prospective Study of Mother-to-Child Chikungunya Virus Infections on
the Island of La Réunion. PLOS Med. 2008;5: e60. doi:10.1371/journal.pmed.0050060

17. Hoz JM de la, Bayona B, Viloria S, Accini JL, Juan-Vergara HS, Viasus D. Fatal cases of
Chikungunya virus infection in Colombia: Diagnostic and treatment challenges. J Clin
Virol. 2015;69: 27–29. doi:10.1016/j.jcv.2015.05.021

18. Moore CA, Staples JE, Dobyns WB, Pessoa A, Ventura CV, Fonseca EB da, et al.
Characterizing the Pattern of Anomalies in Congenital Zika Syndrome for Pediatric
Clinicians. JAMA Pediatr. 2017;171: 288–295. doi:10.1001/jamapediatrics.2016.3982

19. Barbeito-Andrés J, Schuler-Faccini L, Garcez PP. Why is congenital Zika syndrome
asymmetrically distributed among human populations? PLOS Biol. 2018;16: e2006592.
doi:10.1371/journal.pbio.2006592

20. Shepard DS, Undurraga EA, Halasa YA, Stanaway JD. The global economic burden of
dengue: a systematic analysis. Lancet Infect Dis. 2016;16: 935–941. doi:10.1016/S1473-
3099(16)00146-8

21. Deo RC. Machine Learning in Medicine. Circulation. 2015;132: 1920–1930.
doi:10.1161/CIRCULATIONAHA.115.001593

22. Hastie T, Tibshirani R, Friedman J. The Elements of Statistical Learning: Data Mining,
Inference, and Prediction, Second Edition [Internet]. 2nd ed. New York: Springer-Verlag;
2009. Available: //www.springer.com/us/book/9780387848570

23. Breiman L. Statistical Modeling: The Two Cultures. Stat Sci. 2001;16: 199–215.

24.   Margolin AA, Bilal E, Huang E, Norman TC, Ottestad L, Mecham BH, et al. Systematic
analysis of challenge-driven improvements in molecular prognostic models for breast
cancer. Sci Transl Med. 2013;5: 181re1. doi:10.1126/scitranslmed.3006112

25.   Sa-ngamuang C, Haddawy P, Luvira V, Piyaphanee W, Iamsirithaworn S, Lawpoolsri S.
Accuracy of dengue clinical diagnosis with and without NS1 antigen rapid test: Comparison
between human and Bayesian network model decision. PLoS Negl Trop Dis. 2018;12.
doi:10.1371/journal.pntd.0006573

26.   Haddawy P, Kasantikul R, Hasan AHMI, Rattanabumrung C, Rungrun P, Suksopee N, et
al. Spatiotemporal Bayesian Networks for Malaria Prediction: Case Study of Northern
Thailand. Stud Health Technol Inform. 2016;228: 773–777.

27.   Davi CCM, Pastor A, Oliveira T, Lima Neto FB, Braga-Neto U, Bigham A, et al. Severe
Dengue Prognosis Using Human Genome Data and Machine Learning. IEEE Trans Biomed
Eng. 2019; doi:10.1109/TBME.2019.2897285

28.   Khan S, Ullah R, Khan A, Wahab N, Bilal M, Ahmed M. Analysis of dengue infection
based on Raman spectroscopy and support vector machine (SVM). Biomed Opt Express.
2016;7: 2249–2256. doi:10.1364/BOE.7.002249

29.   Melo CFOR, Navarro LC, de Oliveira DN, Guerreiro TM, Lima E de O, Delafiori J, et al. A
Machine Learning Application Based in Random Forest for Integrating Mass Spectrometry-
Based Metabolomic Data: A Simple Screening Method for Patients With Zika Virus. Front
Bioeng Biotechnol. 2018;6. doi:10.3389/fbioe.2018.00031

30.   Nagori A, Dhingra LS, Bhatnagar A, Lodha R, Sethi T. Predicting Hemodynamic Shock
from Thermal Images using Machine Learning. Sci Rep. 2019;9: 91. doi:10.1038/s41598-
018-36586-8

31.   Dakappa PH, Prasad K, Rao SB, Bolumbu G, Bhat GK, Mahabala C. A Predictive Model to
Classify Undifferentiated Fever Cases Based on Twenty-Four-Hour Continuous Tympanic
Temperature Recording. In: Journal of Healthcare Engineering [Internet]. 2017 [cited 9 Apr
2019]. doi:10.1155/2017/5707162

32.   Goto T, Camargo CA, Faridi MK, Freishtat RJ, Hasegawa K. Machine Learning-Based
Prediction of Clinical Outcomes for Children During Emergency Department Triage.
JAMA Netw Open. 2019;2: e186937. doi:10.1001/jamanetworkopen.2018.6937

33.   Raita Y, Goto T, Faridi MK, Brown DFM, Camargo CA, Hasegawa K. Emergency
department triage prediction of clinical outcomes using machine learning models. Crit Care.
2019;23. doi:10.1186/s13054-019-2351-7

34.   Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data
capture (REDCap) – A metadata-driven methodology and workflow process for providing
translational research informatics support. J Biomed Inf. 2009;42: 377–381.

35.   Boulesteix A-L, Schmid M. Machine learning versus statistical modeling. Biom J Biom Z.
2014;56: 588–593. doi:10.1002/bimj.201300226

36.   Peters A. ipred: Improved Predictors. 2017.

37.   Kuhn M. caret: Classification and Regression Training. 2017.

38.   Liaw A, Wiener M. Classification and Regression by RandomForest. Forest. 2001;23.

39.   Zou H, Hastie T. elasticnet: Elastic-Net for Sparse Estimation and Sparse PCA. 2012.

40.   Ridgeway G. gbm: Generalized Boosted Regression Models. 2017.

41.   Venables W, Ripley B. Modern Applied Statistics with S. 4th ed. New York: Springer-
Verlag; 2002.

42.   Christodoulou E, Ma J, Collins GS, Steyerberg EW, Verbakel JY, Van Calster B. A
systematic review shows no performance benefit of machine learning over logistic
regression for clinical prediction models. J Clin Epidemiol. 2019;110: 12–22.
doi:10.1016/j.jclinepi.2019.02.004

43.   Kuhn M. Building Predictive Models in R Using the caret Package. J Stat Softw. 2008;28:
1–26. doi:10.18637/jss.v028.i05

44.   Steyerberg EW, Vickers AJ, Cook NR, Gerds T, Gonen M, Obuchowski N, et al. Assessing
the Performance of Prediction Models: A Framework for Traditional and Novel Measures.
Epidemiology. 2010;21: 128–138. doi:10.1097/EDE.0b013e3181c30fb2

45.   Wickham H, Miller E. haven: Import and Export "SPSS", "Stata" and "SAS" Files. R
package version 1.1.0 ed2017.

46.   Wood SN. Fast stable restricted maximum likelihood and marginal likelihood estimation of
semiparametric generalized linear models. J R Stat Soc Ser B Stat Methodol. 2011;73: 3–
36. doi:10.1111/j.1467-9868.2010.00749.x

47.   N. Wood S. Thin Plate Regression Splines. J R Stat Soc Ser B. 2003;65: 95–114.
doi:10.1111/1467-9868.00374

48.   Karatzoglou A, Smola A, Hornik K, Zeileis A. kernlab - An S4 Package for Kernel
Methods in R. J Stat Softw. 2004;11: 1–20. doi:10.18637/jss.v011.i09

49.   Friedman J, Hastie T, Tibshirani R. Regularization Paths for Generalized Linear Models via
Coordinate Descent. J Stat Softw. 2010;33: 1–22.

50.   Robin X, Turck N, Hainard A, Tiberti N, Lisacek F, Sanchez J, et al. pROC: and open-
source package for R and S+ to analyze and compare ROC curves. BMC Bioinformatics.
2011;12: 77. doi:10.1186/1471-2105-12-77

51. Greenland S, Robins JM. Identifiability, exchangeability, and epidemiological confounding. Int J Epidemiol. 1986;15: 413–419. doi:10.1093/ije/15.3.413

52. Kalayanarooj S, Vaughn DW, Nimmannitya S, Green S, Suntayakorn S, Kunentrasai N, et al. Early clinical and laboratory indicators of acute dengue illness. J Infect Dis. 1997;176: 313–321.

53. Aronsky D, Haug PJ. Diagnosing community-acquired pneumonia with a Bayesian network. Proc AMIA Symp. 1998; 632–636.

54. Sartakhti JS, Zangooei MH, Mozafari K. Hepatitis disease diagnosis using a novel hybrid method based on support vector machine and simulated annealing (SVM-SA). Comput Methods Programs Biomed. 2012;108: 570–579. doi:10.1016/j.cmpb.2011.08.003

55. Er O, Temurtas F, Tanrikulu AÇ. Tuberculosis Disease Diagnosis Using Artificial Neural Networks. J Med Syst. 2008;34: 299–302. doi:10.1007/s10916-008-9241-x

56. Lee VJ, Chow A, Zheng X, Carrasco LR, Cook AR, Lye DC, et al. Simple Clinical and Laboratory Predictors of Chikungunya versus Dengue Infections in Adults. PLoS Negl Trop Dis. 2012;6. doi:10.1371/journal.pntd.0001786

57. Lee I-K, Liu J-W, Yang KD. Clinical and laboratory characteristics and risk factors for fatality in elderly patients with dengue hemorrhagic fever. Am J Trop Med Hyg. 2008;79: 149–153.

58. Thein T-L, Leo Y-S, Fisher DA, Low JG, Oh HML, Gan VC, et al. Risk Factors for Fatality among Confirmed Adult Dengue Inpatients in Singapore: A Matched Case-Control Study. PLOS ONE. 2013;8: e81060. doi:10.1371/journal.pone.0081060

59. Lee I-K, Liu J-W, Yang KD. Clinical characteristics, risk factors, and outcomes in adults experiencing dengue hemorrhagic fever complicated with acute renal failure. Am J Trop Med Hyg. 2009;80: 651–655.

60. Gregory CJ, Lorenzi OD, Colón L, Sepúlveda García A, Santiago LM, Cruz Rivera R, et al. Utility of the Tourniquet Test and the White Blood Cell Count to Differentiate Dengue among Acute Febrile Illnesses in the Emergency Room. PLoS Negl Trop Dis. 2011;5. doi:10.1371/journal.pntd.0001400

61. Sawasdivorn S, Vibulvattanakit S, Sasavatpakdee M, Iamsirithavorn S. Efficacy of Clinical Diagnosis of Dengue Fever in Paediatric Age Groups as Determined by WHO Case Definition 1997 in Thailand. 2001; Available: http://apps.who.int/iris/handle/10665/163627

62. Boillat-Blanco N, Mbarack Z, Samaka J, Mlaganile T, Mamin A, Genton B, et al. Prognostic value of quickSOFA as a predictor of 28-day mortality among febrile adult patients presenting to emergency departments in Dar es Salaam, Tanzania. PLoS ONE. 2018;13. doi:10.1371/journal.pone.0197982

63.   Seymour CW, Liu VX, Iwashyna TJ, Brunkhorst FM, Rea TD, Scherag A, et al.
608   Assessment of Clinical Criteria for Sepsis. JAMA. 2016;315: 762–774.
609   doi:10.1001/jama.2016.0288

610   64.   MDCalc - Medical calculators, equations, algorithms, and scores [Internet]. [cited 11 Apr
611   2019]. Available: https://www.mdcalc.com/

612   65.   Cabitza F, Rasoini R, Gensini GF. Unintended Consequences of Machine Learning in
613   Medicine. JAMA. 2017;318: 517–518. doi:10.1001/jama.2017.7797

 Figures

631 **Figure 1: Flow Diagram of Subject Selection** Subjects clinically diagnosed with arboviral

632 (dengue, chikungunya, Zika) infections were recruited from sentinel clinics in Machala, Ecuador.

633 A subset of subjects were used to test the Severity Index for Suspected Arbovirus (SISA) and

634 Severity Index for Suspected Arbovirus with Laboratory (SISAL) machine learning algorithms.

Figure 2: Results for SISA Dataset Accuracy (blue), Cohen's kappa (red), and AUC (green) were

calculated for the 10-fold cross validation (left) and the holdout test dataset (right) for prediction

of hospitalization status in clinically diagnosed dengue, chikungunya or Zika virus infections.

bag=bagged trees, knn=*k* nearest neighbors, rf=random forest, gbm=generalized boosting models,

enet=elastic net, nnet=neural networks, log=logistic regression

Figure 3: Calibration Plot for SISA Prediction For the final SISA algorithm (generalized

boosting model), the model prediction probability is compared to the actual hospitalization

outcome among the test set. Incorrect predictions are in red: there were three subjects incorrectly

classified for hospitalization and one subject incorrectly classified for non-hospitalization.

Figure 4: Results for SISAL Dataset Accuracy (blue), Cohen's kappa (red), and AUC (green)

were calculated for the 10-fold cross validation (left) and the holdout test dataset (right) for

prediction of hospitalization status in clinically diagnosed dengue, chikungunya or Zika virus

infections. bag=bagged trees, knn=k nearest neighbors, rf=random forest, gbm=generalized

boosting models, enet=elastic net, nnet=neural networks, log=logistic regression

Figure 5: Calibration Plot for SISAL Prediction For the final SISAL algorithm (elastic net

regression), the model prediction probability is compared to the actual hospitalization outcome

among the test set. Incorrect predictions are in red: there were two subjects incorrectly classified

for hospitalization.

683