Lanthanide transport, storage, and beyond: gene products and processes contributing to

lanthanide and methanol metabolism in *Methylorubrum extorquens* AM1

Paula Roszczenko-Jasińska,[a,*,£] Huong N. Vu,[b,*,£] Gabriel A. Subuyuj,[b,*] Ralph Valentine

Crisostomo,[b,*] Elena M. Ayala,[b] James Cai,[b] Nicholas F. Lien,[b] Erik J. Clippard,[b] Richard T.

Ngo,[b] Fauna Yarza,[b,*] Justin P. Wingett,[b] Charumathi Raghuraman,[b] Caitlin A. Hoeber,[b] Norma

C. Martinez-Gomez,[a]# Elizabeth Skovran[b]#

[a]Department of Microbiology and Molecular Genetics, Michigan State University, East Lansing,

USA

[b]Department of Biological Sciences, San José State University, San José, California, USA

Running Head: Lanthanide transport and storage

[£] Both authors contributed equally and share first authorship. Author order was determined both

alphabetically and in order of increasing seniority.

Address correspondence to: N. Cecilia Martinez-Gomez, mart1754@msu.edu and Elizabeth

Skovran, elizabeth.skovran@sjsu.edu

* Present address: PR-J: Institute of Microbiology, University of Warsaw, Warsaw, Poland;

HNV: Department of Microbiology, University of Georgia, Athens, Georgia; USA; GAS:

Department of Microbiology and Molecular Genetics, University of California at Davis, Davis,

California, USA; RVC: Molecular Biology Institute, University of California at Los Angeles,

Los Angeles, California, USA; FY: Department of Biochemistry and Biophysics, University of

California at San Francisco, San Francisco, California, USA

ABSTRACT

Lanthanide elements have been recently recognized as "new life metals" for diverse

environmental microorganisms including Gram-negative methylotrophic bacteria and strains of

Pseudomonas and *Bradyrhizobium*. Yet much remains unknown regarding lanthanide acquisition

and homeostasis. In *Methylorubrum extorquens* AM1, the periplasmic lanthanide-dependent

methanol dehydrogenase XoxF1 produces formaldehyde, which is lethal if allowed to

accumulate. This property enabled a transposon mutagenesis study to expand knowledge of the

metabolic network required for methanol oxidation when lanthanides are available. Growth

studies were conducted to detail the involvement of novel gene products that impact the ability

of XoxF-type enzymes to oxidize methanol to formaldehyde. The identified genes encode an

MxaD homolog, an ABC-type transporter, an aminopeptidase, a putative homospermidine

synthase, and two genes of unknown function annotated as *orf6* and *orf7*. Lanthanide transport

and trafficking genes were also identified. Growth and lanthanide uptake were measured using

strains lacking individual lanthanide transport cluster genes and transmission electron

microscopy was used to visualize lanthanide localization. We corroborated previous reports that

a TonB-ABC transport system is required for lanthanide incorporation to the cytoplasm.

However, cells are able to acclimate overtime and bypass the requirement for the TonB outer

membrane transporter to allow expression of *xoxF1* and growth. Transcriptional reporter fusions

show that excess lanthanides repress the gene encoding the TonB-receptor. Using growth studies

along with energy dispersive X-ray spectroscopy and transmission electron microscopy, we

demonstrate that lanthanides are stored as cytoplasmic inclusions that resemble polyphosphate

granules.

IMPORTANCE

The increasing genetic and biochemical evidence that lanthanide-dependent enzymes are widespread among numerous environmental microbes leads to the parallel questions of how these insoluble metals are scavenged, transported, and used by bacteria. Results herein describe the contribution of the different gene products that constitute the lanthanide utilization and transport machinery in the methylotroph *M. extorquens* AM1 and highlight possible redundancies by periplasmic components. The discovery and characterization of intracellular lanthanide storage in mineral form by these microbes opens the possibility of using methylotrophic platforms for concentration and recovery of these critical energy metals from diverse sources. In addition, methylotrophs are effective biotechnological platforms for the production of biofuels and bioplastics from pollutants such as methane, and inexpensive carbon feedstocks like methanol. Defining the lanthanide acquisition, transport, and storage machinery is a step forward in designing a sustainable platform to recover lanthanides efficiently.

INTRODUCTION

Lanthanide (Ln) metals have long been recognized for their magnetic and superconductive properties and have facilitated the advancement of our communication, green energy, and medical technologies (1–4). However, it has been less than a decade since an inherent role for Ln in methylotrophic bacteria has been described (5–7). Since these initial reports, bacterial strains that are not considered methylotrophs such as *Pseudomonas putida* and *Bradyrhizobium sp.* have been shown to similarly utilize Ln as cofactors in alcohol dehydrogenase (ADH) enzymes, suggesting the impact of Ln on microbial metabolism may be more widespread than initially thought (8–11). The discovery of Ln-dependent ADH enzymes has led to better identification and characterization of methylotrophs in bacterial communities and the cultivation of new bacterial species in the laboratory (6, 12–14).

Methylotrophic bacteria partake in the global carbon cycle by oxidizing methane, methanol, and other single-carbon substrates, and they have been shown to degrade environmental pollutants like methylmercury (15). They are also industrially important for the conversion of single-carbon waste streams and inexpensive feedstocks into value-added chemicals (16–18). As biological Ln use is a nascent field of study, efforts to understand Ln metabolism will further our understanding of bacterial-mediated environmental processes and community interactions, and can facilitate biological engineering efforts to develop methylotrophic platforms to recover critical Ln metals from waste streams like mining leachate, coal fly ash, and postconsumer electronics (2).

To use methanol as a carbon and energy source, many Gram-negative methylotrophic bacteria first oxidize methanol in the periplasmic space using pyrroloquinoline quinone (PQQ)-dependent ADH enzymes. MxaFI is a two-subunit Ca^{2+}-dependent ADH that has been studied

for over 60 years and was long considered to be the predominant methanol dehydrogenase

(MeDH) in nature until Ln-dependent XoxF enzymes were first described in 2011 (5). When

incorporated into the active site, Ln act as potent Lewis acids to facilitate a hydride transfer from

the alcohol to the catalytic PQQ cofactor to prompt alcohol oxidation (19–21).

XoxF-type MeDHs were named XoxF before their function was known and based on

their homology to MoxF which was the original designation for MxaF (MoxF = MxaF) (6, 19).

Since their role in methanol oxidation became apparent, XoxF enzymes have been classified into

five phylogenetically distinct clades (Type 1 to 5) (22, 23) yet recent metagenomic sequencing

efforts suggest additional XoxF clades may exist (18, 24). In 2016, a Ln-dependent ethanol

dehydrogenase was described and named as ExaF based on homology with the Ca^{2+}-dependent

Exa ethanol dehydrogenases from *Pseudomonas* and *Rhodopseudomonas* strains, and homology

with XoxF- and MxaF-type MeDHs (25, 26, 27). While these various ADHs have been

designated as MxaFI and XoxF MeDHs and ExaF ethanol dehydrogenases, there is often overlap

in the substrate range for the ADHs; XoxF and MxaFI enzymes can oxidize ethanol but have a

higher affinity for methanol (19, 22, 23, 25, 28–30), and ExaF ethanol dehydrogenases can

oxidize methanol but with less efficiency (26, 27).

All methylotrophic PQQ-ADHs are periplasmic enzymes associated with a cytochrome

c_L (MxaG, XoxG, and ExaG, respectively) that transfers electrons from PQQ to additional

cytochromes in the electron transport chain. Operons or genomic clusters that encode *xoxF* and

exaF genes often contain homologs of *mxaJ* (*xoxJ* and *exaJ* genes respectively), which encode

periplasmic binding proteins that have been suggested to function in the activation of the ADHs

(31). *mxa* operons encode additional proteins that are suggested to function in Ca^{2+} insertion or

to facilitate interactions between the MxaFI MeDH and its cytochrome (MxaACKLD), are

required for activation of the *mxa* operon (MxaB), or have unknown function (MxaRSE) (32).

Methylorubrum extorquens AM1 (formerly *Methylobacterium extorquens* AM1)

produces a MxaFI-type MeDH (encoded by *MexAM1_META1p4538* and

MexAM1_META1p4535), two XoxF (type 5) MeDHs (encoded by *xoxF1*:

MexAM1_META1p1740 and *xoxF2*: *MexAM1_META1p2757*), and an ExaF-type ethanol

dehydrogenase (encoded by *exaF*: *MexAM1_META1p1139*) (26, 27). Before XoxF enzymes

were classified into five clades, the XoxF enzymes from *M. extorquens* AM1, which share 90%

amino acid similarity, were named as XoxF1 and XoxF2 to distinguish them from one another

(33). Phenotypic studies suggest *M. extorquens* AM1 XoxF1 and XoxF2 have redundant

function. However, in the laboratory, XoxF1 appears to be the dominant XoxF-MeDH as *xoxF2*

is expressed at low levels (25, 27). A fifth putative PQQ-ADH encoded by

MexAM1_META1p4973 does not appear to contribute to methanol growth under laboratory

conditions tested (26).

When Ln are absent, MxaFI is the only known contributor to methanol oxidation in *M.*

extorquens AM1. When Ln are available, the XoxF enzymes produce formaldehyde from

methanol while ExaF oxidizes methanol further to formate (25). Good et al. showed that ExaF

can facilitate Ln-dependent methanol growth when *xoxF1* and *xoxF2* are deleted, but at a slower

rate (26).

The role of Ln in metabolism is not exclusively catalytic (34–36). When Ln are

transported into the cell, a transcriptional response occurs; this effect is referred to as "the Ln-

switch" or "rare earth-switch" (27, 37–39). During the Ln-switch, the *mxa* operon

(*mxaFJGIRSACKLDEHB*) is downregulated and transcript levels of the *xox1* operon genes

(*xoxF1*, *xoxG*, *xoxJ*) are upregulated (25, 27, 37–39).

In *M. extorquens* AM1 and closely related PA1 strain, the MxbDM two-component

system along with the *xoxF* genes themselves have been shown to be required for operation of

the Ln-switch (33, 40, 41). However, suppressor mutations in the *mxbD* sensor kinase encoding

gene can arise, which bypass the need for XoxF1 and XoxF2, presumably by constitutively

activating the MxbM response regulator (33, 36, 41). Loss of *xoxF2* alone does not impact

growth or operation of the Ln-switch. However, in the absence of *xoxF1*, there is an additive

effect when *xoxF2* is deleted from the genome suggesting *xoxF2* may be upregulated to

compensate for the absence of *xoxF1* (27, 33). Though much progress has been made regarding

the catalytic and regulatory roles of Ln in methylotrophic bacteria, relatively little is known

about how these Ln are acquired and incorporated into the enzymes that use them, and if they are

stored intracellularly.

The machinery necessary for Ln transport is in the early stages of characterization and is

predicted to be analogous to siderophore-mediated iron transport (4, 41) with the siderophore-

like molecule referred to as a lanthanophore (42). In *M. extorquens* AM1, ten genes predicted to

encode proteins involved in Ln transport and utilization are clustered together in the genome

(*MexAM1_META1p1778 – MexAM1_META1p1787*) and encode an ABC-type transporter, four

hypothetical periplasmic proteins of unknown function, a periplasmic protein that binds Ln

(encoded by *MexAM1_META1p1781*), a TonB-dependent transporter, lanmodulin (encoded by

MexAM1_META1p1786), and an additional periplasmic-binding protein capable of Ln binding

(4, 43).

Ochsner et al. initially characterized the Ln transport cluster in *M. extorquens* strain PA1

by generating deletions spanning multiple genes in the predicted transport system (41). Their

results showed that a TonB-dependent transporter and a putative ABC transporter were necessary

for Ln transport. A detailed analysis of the contribution of each gene from the transport cluster is

still lacking and Ln transport has not been quantified. Accordingly, Mattocks et al. identified

homologs of this system in *M. extorquens* AM1 and showed that a hypothetical periplasmic

protein encoded by *MexAM1_META1p1781* can efficiently bind Ln from lanthanum (La^{3+}) to

gadolinium (4). Additionally, they showed that the periplasmic binding component of the ABC

transporter encoded by *MexAM1_META1p1778* was unable to bind Ln, possibly because it

instead binds the predicted lanthanophore necessary to transport Ln into the cell.

In this study, we describe new pieces necessary to complete the Ln puzzle: the

identification of novel genes that contribute to Ln metabolism and methanol oxidation, and the

discovery and visualization of Ln storage in *M. extorquens* AM1. Detailed growth studies for

strains lacking each component of the first eight genes in the Ln transport cluster are described,

including the ability of these strains to mutate or acclimate (phenotypic change that is inducible

and reversible) to allow Ln transport through a predicted secondary mechanism. Ln uptake is also

quantified for several transport mutant strains. Finally, we show that *M. extorquens* AM1 stores

Ln with phosphate as crystalline cytoplasmic deposits.

RESULTS

A genetic study identifies gene products that contribute to methanol oxidation. A transposon

mutagenesis study was designed to take advantage of the *in vivo* formaldehyde production

capability of XoxF1 and XoxF2 to identify genes required for XoxF-dependent methanol

oxidation. The strain used to conduct the mutant hunt contained a mutation in *mxaF* to make

cells dependent on the exogenously provided La^{3+} for formaldehyde production, and a second

mutation in *fae*, which would result in formaldehyde accumulation and cell death when methanol

is oxidized to formaldehyde by XoxF1 and XoxF2 (Fig. 1). Transposon insertions that reduced or

eliminated formaldehyde production allowed survival and colony formation on media containing

both methanol and succinate, since methanol resistant strains could use succinate for growth. In

addition to genes required for XoxF1 and XoxF2 mediated methanol oxidation, this mutant hunt

had the potential to isolate insertions in an unknown formaldehyde import system, which would

theoretically reduce formaldehyde levels in the cytoplasm (Fig. 1). As ExaF oxidizes methanol

directly to formate (25), *exaF* and genes specific to ExaF function were not expected to be

identified though this genetic study.

Over six hundred transposon mutants were isolated, and their insertion locations mapped

to the *M. extorquens* AM1 genome. Since it is likely that a portion of these transposon mutants

became methanol-resistant due to spontaneous second-site suppressor mutations and not the

transposon insertion, only genes that were independently identified four or more times were

considered for further analysis and are listed in Table 1. From the twenty eight genes identified

in our transposon mutagenesis study (Table 1), mutations in twenty three genes were constructed

in *mxaF* and/or wild-type strain backgrounds and methanol growth in the presence of La^{3+} was

assessed (Table 2; Table 3).

Growth phenotypes of strains lacking genes identified in the transposon mutagenesis study.

Novel genes which impacted methanol growth when deleted from the *M. extorquens* AM1

genome were identified in the transposon mutagenesis study (Table 1). The identified genes

encode a LysR-type transcriptional regulator (*MexAM1_META1p0863*), two proteins of

unknown function annotated as *orf6* and *orf7* (*MexAM1_META1p1746* and

MexAM1_META1p1747 respectively), an MxaD homolog (*MexAM1_META1p1771*), a putative

homospermidine synthase (*MexAM1_META1p2024*), an ABC-type transporter

(*MexAM1_META1p2359*), and an aminopeptidase (*MexAM1_META1p3908*). Growth rates for

strains lacking these genes are shown in Table 2 and growth curves for mutant strains that had a

30% or greater reduction in growth rate are shown in Fig. 2. The identification of these genes

and initial growth studies lay the foundation to explore the specific roles and requirements for

these gene products in XoxF1 enzyme function.

Also included in the list of genes identified in the transposon mutagenesis study were

those that encode proteins with described roles in methanol oxidation (PQQ biosynthesis, *xoxF1,*

xoxG, xoxJ) and those with predicted functions based on sequence similarity (cytochrome

synthesis and Ln transport) (44). Deletion of genes encoding PQQ biosynthesis (Δ*pqqBCDE*;

Δ*pqqF*) and each of the three identified cytochrome *c* biogenesis and heme export clusters: *cycK*,

heme lyase; *ccmB*, heme exporter; and *ccmC*, heme exporter (reviewed in (44)) eliminated

methanol growth as expected (Table 2).

XoxF1 and XoxF2 catalytically contribute to methanol oxidation and growth when

methanol and Ln are provided (25, 27), and exert a regulatory role in the absence of Ln as

production of XoxF1 or XoxF2 is required for expression of the Ca^{2+}-dependent MxaFI-MeDH

(27, 33). Growth analysis and transcriptional reporter fusion studies were employed to see if

strains lacking *xoxG* or *xoxJ* had similar effects on growth and *mxa* operon expression. In

methanol La^{3+} medium, loss of either *xoxG* or *xoxJ* was equivalent to loss of both *xoxF1* and

xoxF2 (Fig. 3; Table 3), which is consistent with XoxG and XoxJ being essential for XoxF-

dependent methanol oxidation as suggested by recent biochemical studies (14, 31, 45).

Unlike the *xoxF1 xoxF2* double mutant strain, transcriptional reporter fusion studies that

assessed expression from the *mxa* promoter determined that the growth phenotypes observed for

the *xoxG* and *xoxJ* mutants grown in the absence of La^{3+} were not due to impaired *mxa*

expression (average relative fluorescence units (RFU)/OD_{600}: wild type, 322.9 ± 62.5; *xoxF1*

xoxF2, 3.1 ± 1.3; *xoxG,* 313.9 ± 24.0; *xoxJ,* 512.5 ± 54.9). Taken together, these phenotypes

suggest XoxG and XoxJ may play a broader role in methanol metabolism in *M. extorquens* AM1

independent from facilitating XoxF1 and XoxF2 catalytic and regulatory functions.

Detailed phenotypic characterization of Ln utilization and transport cluster mutant strains.

Consistent with previous reports for strain PA1, our transposon mutagenesis studies suggest that

genes encoding homologs of the TonB- and ABC-dependent Fe^{3+} scavenging systems play a role

in methanol metabolism when Ln are present (41). In addition to the transport system homologs,

two of six hypothetical periplasmic proteins encoded in the Ln transport cluster were identified

in the transposon mutagenesis study (Fig. 4A). We expanded Ochsner's study by dissecting the

contribution of individual genes (*MexAM1_META1p1778* through *MexAM1_META1p1785*).

Further, our results show that La^{3+} transport can occur in the absence of the TonB-ABC transport

system by acclimation and/or suppression. Based upon work detailed within, we propose to name

the Ln transport cluster genes as *lut*, for **L**n **u**tilization and **t**ransport.

As reported by Ochsner et al. (41), loss of *lutH* (*Mextp1853* in strain PA1) alone did not

result in a growth defect in medium containing methanol and La^{3+}, which is consistent with the

hypothesis that the TonB-dependent transporter is needed for transport of La^{3+} into the

periplasm. If La^{3+} does not enter the periplasm, the *mxaF* and *mxaI* genes are likely expressed

and used for methanol oxidation. Loss of both *mxaF* and *lutH* arrested growth, but after 90-120

h, growth of the *mxaF lutH* double mutant strain occurred in approximately 60% of the 24

cultures tested (Fig. 4C; Table 2). Consistent with an acclimation process and not a suppressor

mutation, the *mxaF lutH* double mutant strain lost the ability to grow in methanol medium with

La^{3+} after passage onto medium containing succinate. To determine if acclimated growth was

due to production of XoxF1, expression from the *mxa* and *xox1* promoters was measured before

and after acclimation. Before acclimation, *mxa* promoter expression occurred (RFU/OD$_{600}$: 363

$\pm$ 11) while *xox1* expression was repressed as the cells could not transport La^{3+} to promote the

Ln-switch. Once acclimation occurred and the strain began to grow, expression from the *mxa*

promoter was repressed and expression from the *xox1* promoter increased dramatically from

RFU/OD$_{600}$: 53 $\pm$ 7 to 1140 $\pm$ 73, suggesting the cells were able to uptake La^{3+} through the outer

membrane via an unknown mechanism.

LutA, a periplasmic binding protein, is predicted to traffic lanthanophore bound Ln

through the periplasm to the inner membrane transporter system (encoded by *lutE* and *lutF*) for

transport into the cytoplasm (Fig. 4B) (43). Strains lacking *lutE* or *lutF* in the *mxaF* mutant strain

background were unable to grow in medium containing methanol and La^{3+} (Fig. 4C). However,

after 200 h and 150 h respectively, second-site suppressor mutations arose which facilitated

growth albeit 88% slower than that of the wild-type strain (Fig. 4C; Table 2). In contrast, loss of

lutA in the absence of *mxaF* still allowed La^{3+}-dependent growth but at a reduced rate. Genes

lutB and *lutG*, encoding hypothetical periplasmic proteins, were also identified in the transposon

mutagenesis study. Growth of the *mxaF lutB* and *mxaF lutG* double mutant strains was similar to

that of the *mxaF lutA* double mutant strain, suggesting an equally important but non-essential

function. The growth observed for the *lutA*, *lutB*, and *lutG* mutant strains in the absence of *mxaF*

was not identified as acclimation or second-site suppression. To ensure that observed phenotypes

were not due to polarity, mutants lacking individual transport cluster genes (*lutABEFG*) were

complemented by expressing the respective gene in pCM62 (46) and growth similar to the wild-

type strain was restored in each case (Fig. S1).

Genes encoding LutC, LutD, and LanM were not identified in the transposon study but

LutD and LanM have been shown to bind Ln (4). Loss of *lutD* did not result in a significant

growth defect in the *mxaF* mutant and wild-type strain backgrounds while loss of *lutC* resulted in

a small growth defect when La^{3+} was provided (Fig. 4C; Table 2).

Quantification of La^{3+} uptake. To assess how loss of the Ln transport system affects La^{3+} uptake,

La^{3+} levels were quantified from the spent media at different phases of growth (Fig. 5). Strains

were grown in methanol medium containing 2 μM $LaCl_3$ and limiting succinate (3.75 mM) as *lutA,*

lutE, and *lutF* mutant strains are unable to grow or grow poorly with methanol as a sole carbon

source (Fig. 4C; Table 3). While significant decreases in La^{3+} levels were not observed for any

strain during early- to mid-exponential phase (Fig. 5B), as cultures continued to grow, and

succinate was likely depleted, significant differences in La^{3+} levels became apparent. In the wild-

type strain, La^{3+} uptake continued into stationary phase (Fig. 5B). Growth of the *lutE* and *lutF*

mutants (encoding ABC-transporter components) was arrested at an approximate OD_{600} of 0.7, yet

La^{3+} concentrations in the spent media continued to decrease to 0.7 μM eight hours into stationary

phase (Fig. 5A). Taken together with the growth phenotypes, these data suggest that in the absence

of the Lut-ABC-transporter system, Ln are likely transported through the outer membrane but

cannot enter the cytoplasm. Levels of La^{3+} in the supernatants from the *lutH* mutant (*lutH* encodes

the TonB-dependent transporter) were significantly higher than wild type in mid-exponential and

stationary phases (Fig. 5B). A decrease in La^{3+} content from the *lutH* mutant strain culture

supernatants was unexpected but may reflect adsorption of La^{3+} onto the surface of the cells or

interaction of La^{3+} with lipopolysaccharide, a phenomenon observed for other metals in different

bacterial species (47).

Expression from the *lutH* promoter is repressed by La^{3+}. Genes found within the *lut* cluster

are likely not expressed as a single transcript as suggested by the spacing between genes like

lutG and *lutH*, and by multiple promoters predicted by the bacterial promoter prediction tool,

BPROM (48). To test if expression of the of Ln uptake genes is regulated in a similar manner to

Fe^{3+} uptake genes, a fluorescent transcriptional reporter fusion was used to monitor expression

from the predicted *lutH* promoter region in methanol media. Addition of exogenous La^{3+}

repressed expression 4-fold (RFU/OD_{600}: 210 ± 7 without La^{3+} and 47 ± 5 with La^{3+}) showing

that when Ln are in excess, transport is down-regulated. This is consistent with the mechanism of

control for iron homeostasis (49, 50). Further, during the revision of this manuscript, similar

repression was reported for transcription of the TonB transporter of *Methylotuvimicrobium*

buryatense 5GB1C (51).

La^{3+} is stored as cytoplasmic crystalline deposits. The continued uptake of La^{3+} in stationary

phase suggested the possibility of Ln storage. To determine if La^{3+} storage could facilitate

methanol growth in the absence of exogenously supplied Ln, an *mxaF* mutant strain was grown

in methanol medium with 10-times the standard concentration of LaCl$_3$, resulting in a growth

rate of 0.15 ± 0.00 h^{-1}. Once in stationary phase, cells were washed and sub-cultured into

methanol medium lacking La^{3+}. As shown in Fig. 5C, a similar growth rate to that observed

when cells were grown with La^{3+} was obtained until the culture reached an approximate OD$_{600}$ of

1.0. A slower growth rate was noted as growth continued to an OD$_{600}$ of 1.5. After a second

subculture into media lacking La^{3+}, cells continued to grow but at a severely impaired rate (0.01

$\pm$ 0.0 h^{-1}) until they reached a peak OD$_{600}$ of 1.3. These data are consistent with the ability of Ln-

storage to supply Ln for methanol oxidation until the storage is depleted. As these storage growth

curves were carried out for >400 h, a control experiment was done to determine growth due to

either background levels of non-MxaF ADH activity or due to leaching of residual Ln from the

glassware. A growth rate of 0.004 ± 0.001 h^{-1} was observed when an *mxaF* mutant strain that had

not been previously grown with La^{3+} was tested.

Transmission electron microscopy (TEM) coupled with energy dispersive X-ray

spectroscopy (EDS) has been used to determine the elemental composition of cellular inclusions

(52, 53) while La^{3+} has been widely used as an intracellular and periplasmic stain for electron

microscopy (54–56). Here, we show that La^{3+} can be directly identified by TEM if accumulated

inside *M. extorquens* AM1 cells. Electron-dense deposits were observed in the cytoplasm from

M. extorquens AM1 cells grown with exogenous LaCl$_3$ (Fig. 6A-B). Samples were analyzed

using EDS and corroborated that the electron dense deposits contained La^{3+} (Fig. 6C). When

grown without La^{3+}, only a few cells showed smaller electron dense areas (Fig. 6E-F); however,

La^{3+} was not detected in EDS analysis in these cases (Fig. 6G). These data demonstrate that La^{3+}

can be stored in *M. extorquens* as metal deposits. Moreover, EDS analysis of electron dense

areas from the wild-type strain grown with La^{3+} determined a content of lanthanum (22.2 $\pm$

1.0%), phosphorous (15.1 $\pm$ 2.1%), and oxygen (51.1 $\pm$ 1.9%) suggesting La^{3+} is complexed with

phosphate (Fig. S2). Traces of chloride (3.0%), calcium (2.2%), and aluminum (3.4%) ions were

also detected. The copper, carbon, and silicon ion content from the support grids and embedding

medium were not considered for metal content calculations. High-resolution transmission

electron microscopy (HRTEM) images of the La^{3+} deposits showed an atomic lattice with a

Moiré fringes pattern, indicating a crystalline nature (57) (Fig. 6D). Together, these results

suggest that La^{3+} is embedded in inorganic phosphate crystals, which form the electron dense

deposits observed in the cytoplasm.

Visualization of La^{3+} accumulation in *lut* transporter mutants. To determine if La^{3+} could be

visualized in strains lacking the Lut-TonB-ABC transport system, TEM and EDS were employed

for the analysis of *lutA*, *lutE*, *lutF*, and *lutH* mutant strains. Strains were grown in methanol

medium containing La^{3+} and limiting succinate. Samples stained with OsO_4 and 2% uranyl

acetate allowed the outer and inner membrane of the bacterial cells to be distinguished (Fig. 7A-

E left subpanels), while visualization without staining enabled metal content analysis by

removing the interaction between Os^{8+} and phosphate, which interferes with La^{3+} measurements

(Fig. 7A-E right subpanels). Mutants lacking the TonB-dependent transporter (encoded by *lutH*)

did not display La^{3+} deposits (Fig. 7A). In contrast, localized La^{3+} deposits were visualized in the

periplasmic space in mutant strains lacking the ABC-transporter components (encoded by *lutA,*

lutE, and *lutF*) as shown in Fig. 7B-E. EDS microanalyses confirmed these electron dense

periplasmic deposition areas contained La^{3+} (Fig. 7F). Taken together, these findings directly

demonstrate a role for the TonB-dependent and ABC transporters in Ln transport.

DISCUSSION

The Ln-dependent XoxF1-MeDH has been shown to produce formaldehyde *in vivo* (25).

Here, we took advantage of this property which allowed lethal levels of formaldehyde to

accumulate when methanol was oxidized by the XoxF enzymes and *fae* was deleted from the

genome. These phenotypes enabled a genetic selection to identify gene products required for or

involved in methanol oxidation by the XoxF enzymes.

Novel genes were identified that when deleted from the genome, resulted in a 30% or

greater reduction in growth rate in medium containing methanol and La^{3+}. An ABC-transporter

of unknown function (encoded by *MexAM1_META1p2359*) was found to be essential for

methanol growth. Some possible functions for META1p2359 may include formaldehyde import,

or export of Ln or PQQ into the periplasm for incorporation into the ADHs. Further, a gene

encoding a putative homospermidine synthase (*hss*) was identified. Homospermidine synthases

function in polyamine biosynthesis and their products (e.g. putrescine, spermidine) have been

implicated in roles as diverse as host pathogen interactions, biofilm formation, siderophore

production, acid resistance, and free radical scavenging (58, 59).

Before the existence of Ln-dependent MeDHs was known, it was concluded that *orf6* and

orf7 gene products did not have a role in C_1-metabolism though *orf6* and *orf7* were proximal to

other methylotrophic genes (60). Results here clearly suggest that *orf6* and *orf7* are important for

Ln-dependent methylotrophy.

A homolog of MxaD (encoded by *MexAM1_META1p1771*) was also shown to be

important for methanol La^{3+} growth. MxaD is a 17-kDa periplasmic protein that directly or

indirectly stimulates the interaction between the MxaFI-MeDH and cytochrome c_L (61).

Interestingly, upstream of *MexAM1_META1p1771* is another *mxaD* homolog

(*MexAM1_META1p1772*) which was not identified in the transposon mutagenesis study. The

presence of multiple *mxaD* homologs may suggest redundancy or that the various MxaD

homologs may function with different ADH enzymes.

As expected, transposon insertions were found within genes with known or predicted

roles in methanol oxidation like PQQ biosynthesis, cytochrome biogenesis, and the *xoxF₁GJ*

genes. Recent biochemical and structural analyses of XoxG suggest that this cytochrome is tuned

specifically for light Ln (lanthanum to samarium), while XoxJ interacts with and may activate

XoxF (31). Here we show that loss of either *xoxG* or *xoxJ* resulted in growth that mirrored the

xoxF1 xoxF2 double mutant strain, consistent with XoxG and XoxJ as essential for the activity of

XoxF-enzymes.

Using growth and transcriptional reporter fusion studies, we show that unlike XoxF1,

XoxG and XoxJ are not required for expression of the *mxa* genes yet loss of the *xoxG* and *xoxJ*

genes impacts growth in the absence of La^{3+}. In the methanotroph *Methylomonas* sp. strain

LW13, loss of *xoxG* also results in a growth defect in methanol medium lacking Ln suggesting

an unknown role in metabolism in addition to functioning as a cytochrome for XoxF-mediated

methanol oxidation (45). Our results are in contrast to previous reports for *M. extorquens* PA1

and AM1, where loss of *xoxG* and *xoxJ* did not result in a growth phenotype in methanol

medium lacking Ln (41, 60). Notably, the previous AM1 studies were carried out on agar plates

where subtle growth defects may not be apparent, and the growth media used for the previous

PA1 and AM1 studies was not the same as used in these studies.

Growth phenotypes for *M. extorquens* AM1 PQQ biosynthesis mutants have been

reported previously in the absence of Ln (62) but *ccm* and *cyc* mutants have not been

characterized and the functions encoded by these genes inferred from sequence homology. Our

results using La^{3+} corroborate that the biosynthesis of PQQ, heme export, and cytochrome

biogenesis enable Ln-dependent ADH activities as predicted.

Many insertions identified in this study mapped to the previously identified Ln transport

gene cluster (4, 41). Characterization of strains lacking genes in the *lut* cluster is consistent with

the observations recently reported for *M. extorquens* PA1 (41) where a TonB-ABC transport

system contributes to Ln transport into the cytoplasm. It has been proposed that Ln may be

acquired via lanthanophores (42), which enter the periplasm through the TonB-dependent

transporter (4, 41). However, the acclimation of the *mxaF lutH* double mutant strain allowing

xoxF1 expression and growth suggests that an additional mechanism exists to take up Ln.

Additionally, we show that similar to the paradigm for regulation of siderophore mediated Fe^{3+}

uptake, expression from the TonB-dependent transporter promoter, *lutH*, is repressed by La^{3+}.

Since transposon mutants disrupted in the genes encoding lanmodulin or LutD were not

isolated, our observations are consistent with the findings in *M. extorquens* PA1 that suggest a

non-essential or redundant role for these proteins in Ln metabolism (41). Notably, the *mxaF fae*

transposon mutagenesis study did not identify obvious gene candidates for lanthanophore

biosynthesis though over 600 insertions were mapped to the genome. This may indicate that

more than one lanthanophore is produced by the cell or that the lanthanophore has an essential

role that is not yet understood. Additionally, mutations in the TonB-ExbB-ExbD energy

transducing system genes were not identified. Homologs are encoded in the *M. extorquens* AM1

genome in single copy (*MexAM1_META1p1510 – MexAM1_META1p1512*). Using a variety of

media conditions, repeated attempts were made to delete TonB without success, suggesting

TonB may be essential for *M. extorquens* AM1.

Our transport, TEM, and EDS analyses are consistent with LutH facilitating Ln transport

into the periplasm and the Ln-ABC transport system facilitating Ln transport into the cytoplasm.

La^{3+} concentrations found in the supernatant from strain variants lacking transport system

components suggest that once in the periplasm, significant concentrations of La^{3+} do not go back

outside of the cell as ABC transporter mutant strains showed uptake of La^{3+} from the medium

similar to the wild-type strain. Intriguingly, TEM and EDS studies with ABC transporter mutant

strains demonstrated localized accumulation of La^{3+} in the periplasmic space. It is not yet clear

how or why La^{3+} accumulate in specific areas rather than appear diffused throughout the

periplasm. Taken together, our phenotypic growth, transport, and visualization studies suggest

that Ln must enter the cytoplasm for the *xoxF* and *exaF* ADHs to be expressed and that Ln

uptake into the periplasmic space do not support methanol dehydrogenase activity but may be

enough to repress *mxa* expression. This work will facilitate new exploration of the Ln-switch.

It has been observed that bacteria such as *Bacillus licheniformis* (63), *Myxococcus*

xanthus (56), and *Pseudomonas aeruginosa* (64) can effectively adsorb Ln in mineral form onto

their cell surface when Ln are at high concentrations (mM). Unlike these microorganisms, our

TEM and EDS analyses demonstrate that *M. extorquens* AM1 stores Ln in the cytoplasm in

crystal form. This is consistent with recent studies for the non-methylotroph *Thermus*

scotoductus SA-01 which showed that europium (Eu^{3+}) can accumulate in the cytoplasm.

However, these studies were conducted using very high Eu^{3+} concentrations that are not typically

found in nature (53). Further, a role for Eu in the metabolism of *T. scotoductus* SA-01 remains to

be defined.

For many bacteria, biomineralization is a mechanism used to cope with toxicity of

different metals, manage waste products, sense and change orientations in accordance with

geomagnetic fields, and store important cations for growth (65–70). It has been reported that

some microorganisms store cations like Mg^{2+} and Ca^{2+} complexed to polyphosphate in the form

of volutin (also known as metachromatic granules) or acidocalcisomes (71–74). It is not yet

known if *M. extorquens* AM1 stores La^{3+} complexed to polyphosphate, however, the ratios of P,

O, and La^{3+} detected in our studies are consistent with La^{3+} phosphates. Detailed studies are

necessary to define the exact chemical structure of Ln storage deposits in *M. extorquens* AM1

and if these granules are membrane or lipid bound. Our current findings bring exciting

implications for bacterial metabolism and cell biology, and for the development of

bioremediation and biometallurgy strategies for Ln recovery.

MATERIALS AND METHODS

Bacterial strains and cultivation. Strains and plasmids used in this study are listed in Table 5.

Escherichia coli strains were cultivated in Lysogeny Broth (LB) medium (75) (BD, Franklin

Lakes, NJ) at 37°C. *M. extorquens* AM1 strains were grown in *Methylobacterium* PIPES

[piperazine-*N,N'*-bis(2-ethanesulfonic acid)] (MP) media (76) supplemented with succinate (15

mM) and/or methanol (125 mM) as described (26) unless otherwise stated. Conjugations took

place on Difco Nutrient Agar (Thermo Fisher Scientific, Waltham, MA). Cultures were grown in

round-bottom polypropylene or borosilicate glass culture tubes, or 250 mL polypropylene

Erlenmeyer flasks (Thermo Fisher Scientific, Waltham, MA). If glass tubes or flasks were used

to culture bacteria, they were pretreated to remove Ln as previously described (27). Liquid

cultures were grown at 29°C and shaken at 200 and 180 rpm in New Brunswick Innova 2300 and

Excella E25 shaking incubators (Eppendorf, Hauppauge, NY), respectively. LaCl$_3$ was

supplemented to a final concentration of 2 or 20 μM when indicated. When necessary, antibiotics

were added at the following concentrations: rifamycin (Rif, 50 μg/mL), tetracycline (Tc, 10

μg/mL for LB, 5 μg/mL for MP or 10 μg/mL when used together with Rif), kanamycin (Km, 50

μg/mL), ampicillin (Ap, 50 μg/mL).

Plasmid and strain construction. Primers used for plasmid and strain construction are listed in

Table 6. The allelic exchange plasmid pHV2 was constructed by cloning the *sacB* gene from

pCM433 (77) into the PscI site of pCM184 (78) in the same orientation as the Tc resistance gene.

Insertion and orientation of *sacB* was confirmed by colony PCR. The *lutH* transcriptional

reporter fusion was constructed by cloning the promoter region of *lutH* into the AclI and EcoRI

sites upstream of a promoter-less *venus* gene in pAP5 (33). To create overexpression constructs

for complementation studies, individual genes in the *lut* operon (*lutA*, *lutB*, *lutE*, *lutF*, and *lutG)*

were cloned into the KpnI and SacI sites downstream of a P$_{lac}$ promoter in pCM62 (46).

Diagnostic PCR was used to confirm successful integration of inserts. Plasmids were maintained

in *E. coli* TOP10 (Invitrogen, Carlsbad, CA).

Gene deletions were constructed using pCM184 or pHV2 as previously described (31)

except 5% sucrose was added for counter selection against single crossovers (79) when using

pHV2. Plasmids were conjugated into *M. extorquens* AM1 via biparental mating using *E. coli*

S17-1 (81) or triparental mating using *E. coli* TOP10 (Invitrogen, Carlsbad, CA) and *E. coli*

harboring the conjugative plasmid pRK2013 as described (31). When indicated, the Km

resistance cassette was resolved using pCM157 to achieve marker-less deletions (80).

Transposon mutagenesis. Suicide vector pCM639 carrying a mini transposon IS*phoA*/hah-Tc

(82) was conjugated into the *mxaF fae* strain background via triparental mating as described (31,

83). Dilutions of the mating mixtures were plated onto MP succinate (15 mM) plus methanol (50

mM) La^{3+} medium containing 10 μg/mL Tc to select for successful integration of the mini

transposon into the *M. extorquens* AM1 genome and 50 μg/mL Rif to counter select against *E.*

coli strains bearing pCM639 or pRK2013. Plates were incubated for 5-7 days at 29^{o}C.

Transposon mutant colonies were streaked onto MP succinate methanol La^{3+} Tc medium for

downstream studies.

Location of transposon insertions. To identify the transposon insertion sites, genomic DNA

was isolated using a Qiagen DNeasy UltraClean Microbial Kit (Qiagen, Germantown, MD).

Degenerate nested PCR was performed as described (81, 82) with the following exceptions: PCR

reactions contained 1 μM of each primer, 0.05 U/μL Dream Taq (Thermo Fisher Scientific,

Waltham, MA), and 5% dimethyl sulfoxide. Modifications to the PCR amplification parameters

included 2 minutes for the initial denaturation at 95°C, 6 cycles of annealing at 40°C followed by

25 cycles of annealing at 65°C for the first PCR reaction, and 30 cycles of annealing at 65°C for

the second PCR reaction. PCR products were purified using a Qiagen QIAquick 96 PCR

Purification Kit (Germantown, MD). Sequence analysis was performed using TransMapper, a

Python-based program developed in-house to identify transposon insertion locations and map

them to the *M. extorquens* AM1 genome for visualization using SnapGene Viewer (GSL Biotech

LLC, Chicago, IL).

Phenotypic analyses. Growth phenotypes were determined on solid or in liquid MP media using

a minimum of three biological replicates. On solid media, colony size was scored after four days.

Growth curve experiments were conducted at 29°C in an Excella E25 shaking incubator (New

Brunswick Scientific, Edison, NJ) using a custom-built angled tube rack holder as previously

described (26). Optical density (OD_{600}) was measured at 600 nm using a Spectronic 20D

spectrophotometer (Milton Roy Company, Warminster, PA). For strains with extended growth

lags, suppression and acclimation was assessed. Strains from the growth curves were streaked

onto methanol La^{3+} medium after they reached stationary phase. If the parent stock strain did not

grow on methanol La^{3+} medium and the strain post-growth curve grew, acclimation versus

suppression was tested. Strains were passaged from the methanol medium plate to a succinate

medium plate. After colonies grew on succinate medium, they were streaked back onto methanol

La^{3+} medium. If strains retained the ability to grow on methanol medium, it was concluded that

growth was due to a suppressor mutation. If strains lost the ability to grow on methanol medium

after succinate passage, it was concluded growth was due to acclimation and not a genetic

change.

For Ln storage growth experiments, *mxaF* deletion strains were grown with 20 μM LaCl$_3$

until six hours after transition into stationary phase. Cells were centrifuged and washed four

times with MP medium lacking La^{3+} and a carbon source, and sub-cultured into fresh MP

methanol medium without La^{3+}. Six hours post stationary phase, the cultures were streaked onto

MP methanol medium with and without La^{3+} to check for contamination. Cultures were again

washed and sub-cultured as described above. This process was repeated for two rounds of

growth without La^{3+}.

Transcriptional reporter fusion assays. *M. extorquens* AM1 strains carrying *mxa*, *xox1*, and

lutH transcriptional reporter fusions (Table 5) which use *venus* (83) as a fluorescent reporter

were grown in MP media supplemented with methanol only or methanol and succinate with and

without La^{3+} as indicated in the text. Expression was measured as relative fluorescent units

(RFU) using a SpectraMax M2 plate reader (Molecular Devices, Sunnyvale, CA) and normalized

to OD$_{600}$ as previously described (27).

La^{3+} depletion during *M. extorquens* AM1 growth. Overnight cultures of wild type, *lutA*, *lutE*,

lutF, and *lutH* mutant strains were inoculated 1:50 into 250 mL polycarbonate flasks (Corning

Inc., Corning, NY) containing 75 mL of MP medium (76). Succinate (3.75 mM) and methanol

(125 mM) were added as carbon sources with 2 μM LaCl$_3$. Flasks were incubated at 28°C at 200

rpm in Innova 2300 shaking incubators (Eppendorf, Hauppauge, NY) for 44 h. To monitor La^{3+}

depletion during *M. extorquens* AM1 cultivation, the Arsenazo III assay was used (84). 5 mL

samples were collected at four different time points (OD_{600} of 0.04, 0.4, 0.7, and 1.5) and the

concentration of La^{3+} remaining in the supernatant was calculated using the calibration curve

prepared as previously described (84). A control of three uninoculated flasks containing MP

medium with 2 µM $LaCl_3$ were considered to determine La^{3+} adsorption by the flasks which was

subtracted from the culture measurements. The initial concentration of La^{3+} in the media (before

growth) was measured using the Arsenazo III assay in the same way as described above.

Significant differences between depletion of La^{3+} by different strains were calculated using One-

way ANOVA followed by a T-test.

Cellular locations of Ln visualized using Transmission Electron Microscopy (TEM). Sample

preparation for TEM: wild type, *lutA*, *lutE*, *lutF*, and *lutH* mutant strains were grown in MP

medium containing 125 mM methanol and 3.75 mM succinate as carbon sources with or without

the addition of 20 µM $LaCl_3$ until they reached an OD_{600} of ~0.6. 3 mL of cells was harvested by

centrifugation for 3 min at $1500 \times g$ at room temperature and fixed for 30 min in 1 mL of 2.5%

(v/v) glutaraldehyde (Electron Microscopy Sciences, Hatfield, PA) in 0.1 M cacodylate buffer

(Electron Microscopy Sciences, Hatfield, PA). After fixation, cells were pelleted by centrifugation

for 3 min at $1500 \times g$ and washed with 1 mL of 0.1 M cacodylate buffer. Cell pellets were

embedded in 2% (w/v) agarose and washed three times with 0.1 M cacodylate buffer. When

indicated, pellets in agarose blocks were stained for 30 min in 1% osmium tetroxide in 0.1 M

cacodylate buffer. Samples were washed three times with 0.1 M cacodylate buffer, dehydrated in

acetone, and embedded in Spurr resin (Electron Microscopy Sciences, Hatfield, PA). Blocks were

polymerized at 60ºC for 48 h. 70 nm sections were obtained with a Power Tome XL

ultramicrotome (RMC Boeckeler Instruments, Tucson AZ), deposited on 200 mesh carbon coated

grids, and stained with 2% uranyl acetate (Electron Microscopy Sciences, Hatfield, PA). To assess

the presence of La^{3+} by EDS, sections were left unstained. To image the distribution of cellular

La^{3+}, a TEM JOEL 1400 Flash (Japan Electron Optics Laboratory, Tokyo, Japan) was used.

Detection of La^{3+} in the cells and high-resolution imaging were done with a JEOL 2200FS (Japan

Electron Optics Laboratory, Tokyo, Japan) operated at 200kV. X-ray energy dispersive

spectroscopy was performed using an Oxford Instruments INCA system (Abingdon, United

Kingdom).

ACKNOWLEDGMENTS

We would like to thank Dr. Lena Daumann and Dr. Nathan Good for critical review of this

manuscript. We would like to thank San José State University General Microbiology students who

isolated transposon mutants and purified genomic DNA for sequencing as part of a class research

project. The mutant hunt in this study was inspired by Dr. Elizabeth Skovran's undergraduate

research mentor, Dr. Marc Rott at the University of Wisconsin-LaCrosse who used transposon

mutagenesis in the teaching lab to identify genes involved in butanol oxidation in *Rhodobacter*

sphaeroides. We would like to thank Timothy Andriese for assistance with sequencing of the

transposon mutant DNA and all Skovran lab members for assistance with growth curves and

transcriptional reporter fusion assays. TEM work was done at the Center for Advanced

Microscopy, Michigan State University. We would like to thank Dr. Alicia Withrow for invaluable

assistance with TEM experiments and Dr. Xudong Fan for his expertise using TEM-EDS. E.S.,

N.C.M-G., P.R-J., and H.N.V. directed experiments. E.S., N.C.M-G., P.R-J., and H.N.V. wrote

the manuscript. P.R-J. conducted microscopy and Ln transport experiments. H.N.V, G.A.S, J.C.,

and E.C carried out transposon mutagenesis studies. H.N.V., G.A.S., R.C., J.C., C.R., E.M.A.,

E.C., N.F.L., and F.Y. constructed strains and conducted growth experiments. R.C., C.H., J.P.W.,

and G.A.S. conducted expression studies. R.T.N created TransMapper.

FUNDING SOURCES

This material is based upon work supported by the National Science Foundation under Grant No.

1750003 and by a California State University Program for Education and Research in

Biotechnology (CSUPERB) Joint Venture Grant. P.R-J. was supported by the National Science

Foundation under Grant No. 1750003. E.M.A and F.Y. were supported by the National Science

Foundation Research Initiative for Scientific Enhancement (RISE) award under Grant No.

R25GM071381. F.Y. was also supported by the National Institute of Health Maximizing Access

to Research Careers Undergraduate Student Training in Academic Research (MARC U-STAR)

award under Grant No. 4T34GM008253. Funding for transposon DNA isolation, PCR, and

sequencing was provided by San José State University through the Department of Biological

Sciences.

 TABLES

 Table 1. Genes identified four or more times via transposon mutagenesis.

Gene Designation	Gene Name	Predicted Function
MexAM1_META1p0863		LysR-type regulator
MexAM1_META1p1292	*cycL*	*c*-type cytochrome biogenesis
MexAM1_META1p1293	*cycK*	Heme lyase
MexAM1_META1p1294	*cycJ*	Periplasmic heme chaperone
MexAM1_META1p1740	*xoxF1*	Ln-dependent methanol dehydrogenase
MexAM1_META1p1741	*xoxG*	Cytochrome *c*
MexAM1_META1p1742	*xoxJ*	Periplasmic binding protein
MexAM1_META1p1746	*orf6*	Unknown
MexAM1_META1p1747	*orf7*	Unknown
MexAM1_META1p1748	*pqqE*	PQQ biosynthesis
MexAM1_META1p1749	*pqqCD*	PQQ biosynthesis
MexAM1_META1p1750	*pqqB*	PQQ biosynthesis
MexAM1_META1p1771		MxaD homolog
MexAM1_META1p1778	*lutA*	ABC transporter-periplasmic binding component
MexAM1_META1p1779	*lutB*	Exported protein
MexAM1_META1p1782	*lutE*	ABC transporter-ATP binding component
MexAM1_META1p1783	*lutF*	ABC transporter-membrane component
MexAM1_META1p1784	*lutG*	Exported protein
MexAM1_META1p1785	*lutH*	TonB-dependent receptor
MexAM1_META1p2024	*hss*	Homospermidine synthase
MexAM1_META1p2330	*pqqF*	Protease
MexAM1_META1p2331	*pqqG*	Protease
MexAM1_META1p2359		ABC transporter
MexAM1_META1p2732	*ccmC*	Heme export
MexAM1_META1p2734	*ccmG*	*c*-type cytochrome biogenesis
MexAM1_META1p2825	*ccmB*	Heme export
MexAM1_META1p2826	*ccmA*	Heme export
MexAM1_META1p3908		Leucyl aminopeptidase

607   Table 2. Growth parameters for strains grown in methanol medium with La^{3+}.

Strain	Growth Rate $(h^{-1})^£$
	MeOH + La^{3+}
Growth of control strains	
wild type	0.16 ± 0.01
mxaF	0.16 ± 0.01
xoxF1	6-9 h lag, 0.07 ± 0.00
xoxF1 xoxF2	6 h lag, 0.04 ± 0.01
mxaF xoxF1 xoxF2	6 h lag, 0.04 ± 0.00
Growth of strains lacking novel genes identified in genetic screen	
MexAM1_META1p0863	0.14 ± 0.00
MexAM1_META1p1771	0.11 ± 0.01
orf6	0.03 ± 0.00
orf7	9 h lag, 0.08 ± 0.01
MexAM1_META1p2359	NG*
hss	6 h lag, 0.10 ± 0.01
MexAM1_META1p3908	15 h lag, 0.07 ± 0.00
Growth of strains lacking genes with known function	
pqqBCDE	NG
pqqF	NG
cycK	NG
ccmB	NG
ccmC	NG
Growth of strains lacking lanthanide utilization and transport genes	
lutA	0.02 ± 0.01, 96 h; $0.08 \pm 0.01^{\#}$
mxaF lutA	0.02 ± 0.00
lutB	0.02 ± 0.00, 87 h; $0.05 \pm 0.01^{\#}$
mxaF lutB	0.02 ± 0.00
lutC	9 h lag, 0.11 ± 0.01
mxaF lutC	9 h lag, 0.11 ± 0.01
lutE	NG*
mxaF lutE	NG*
lutF	NG*
mxaF lutF	NG*
lutG	12 h lag, 0.03 ± 0.00
mxaF lutG	12 h lag, 0.03 ± 0.00
lutH	0.16 ± 0.00

Table 2. (continued)

	Growth Rate (h^{-1})$^{£}$
Strain	**MeOH + La^{3+}**
mxaF lutH	NG*

$^{£}$Data for a minimum of three biological replicates are reported.

* indicates if a suppressor mutation or acclimation event allowed eventual growth of the strain

(detailed in Table 4).

$^{#}$ indicates two growth rates were observed with an initial growth rate lasting for the time

indicated, followed by a second growth rate. Accurate suppression and acclimation

determinations could not be made for these strains as strains grew on methanol media prior to

inoculation. A small increase in colony size was observed after growth.

NG, no growth; MeOH, methanol.

± indicates standard deviations.

Table 3. Growth parameters for strains grown in methanol medium with and without La^{3+}.

Strain	Growth Rate (h^{-1})[£]	
	MeOH	MeOH + La^{3+}
Wild type	0.14 ± 0.01	0.16 ± 0.01
xoxF1 xoxF2	NG*	6 h lag, 0.04 ± 0.01
xoxG	3 h lag, 0.11 ± 0.01	0.04 ± 0.00
xoxJ	21 h lag, 0.13 ± 0.00	0.04 ± 0.01

620   [£]Data for a minimum of three biological replicates are reported.

621   * indicates a suppressor mutation allowed eventual growth of the strain.

622   NG, no growth; MeOH, methanol.

623   ± indicates standard deviations.

625   Table 4. Growth parameters of suppressor and acclimation events.

Strain	MeOH + La^{3+}	
	Time of S/A (h)	S/A Growth Rate (h^{-1}) [£]
MexAM1_META1p2359	S: 85	0.10 ± 0.01
lutE	S: 75-90	0.07 ± 0.01
mxaF lutE	S: 200-220	0.02 ± 0.00
lutF	S: 79-91	0.09 ± 0.02
mxaF lutF	S: 145-157	0.02 ± 0.01
mxaF lutH	A: 90-120	0.14 ± 0.02

626   [£]Data for a minimum of three biological replicates is reported.

627   Abbreviations are as follows: methanol, MeOH; suppression, S; acclimation, A.

628 Table 5. Strains and plasmids used in this study.

Strain or plasmid	Description	Reference
M. extorquens		
AM1	Rif^R derivative (wild type)	(85)
CM194.1	*ΔmxaF*	(86)
ES2813	*ΔmxaF Δfae*	This study
ES2432	*ΔxoxF1*	This study
ES2435	*ΔxoxF2 ΔxoxF1*	This study
ES2439	*ΔmxaF ΔxoxF2 ΔxoxF1*	This study
ES2386	*pqqBCDE::Km*	This study
ES2360	*pqqF::Km*	This study
ES3941	*cycK::Km*	This study
ES3968	*ccmB::Km*	This study
ES3965	*ccmC::Km*	This study
ES2817	*ΔxoxG*	This study
ES2821	*ΔxoxJ*	This study
ES2785	*ΔMETA1p0863*	This study
ES3947	*MexAM1_META1p1771::Km*	This study
ES3282	*Δorf6*	This study
ES2781	*Δorf7*	This study
ES3962	*MexAM1_META1p2359::Km*	This study
ES3959	*hss::Km*	This study
ES3257	*MexAM1_META1p3908::Km*	This study
ES2861	*MexAM1_META1_3909::Km*	This study
ES3950	*lutA::Km*	This study
ES4122	*lutB::Km*	This study
ES4444	*lutC::Km*	This study
ES4389	*lutD::Km*	This study
ES3953	*lutE::Km*	This study
ES4249	*lutF::Km*	This study
ES3956	*lutG::Km*	This study
ES2879	*lutH::Km*	This study
ES3983	*ΔmxaF lutA::Km*	This study
ES4124	*ΔmxaF lutB::Km*	This study
ES4447	*ΔmxaF lutC::Km*	This study
ES4392	*ΔmxaF lutD::Km*	This study
ES3986	*ΔmxaF lutE::Km*	This study

Table 5. (continued)

Strain or plasmid	Description	Reference
M. extorquens		
ES4246	Δ*mxaF lutF::Km*	This study
ES3989	Δ*mxaF lutG::Km*	This study
ES2962	Δ*mxaF lutH::Km*	This study
ES2231	AM1 / pCM62	This study
ES4293	*lutA::Km* / pCM62	This study
ES4295	*lutA::Km* / pES667	This study
ES5029	*lutB::Km* / pCM62	This study
ES5030	*lutB::Km* / pES810	This study
ES4301	*lutE::Km* / pCM62	This study
ES4305	*lutE::Km* / pES666	This study
ES5031	*lutF::Km* / pCM62	This study
ES5032	*lutF::Km* / pES815	This study
ES4309	*lutG::Km* / pCM62	This study
ES4315	*lutG::Km* / pES668	This study
ES1819	AM1 / pES502	(30)
ES4192	Δ*xoxF2* Δ*xoxF1* / pES502	This study
ES4194	Δ*xoxG* / pES502	This study
ES4196	Δ*xoxJ* / pES502	This study
ES3743	AM1 / pHV25	This study
E. coli		
TOP10	Cloning strain (Sm^R)	Invitrogen
S17-1	Helper strain (Tp^R, Sm^R)	(83)
Plasmid		
pRK2013	Conjugative helper plasmid (Km^R)	(33)
pCM184	Allelic exchange suicide vector (Km^R, Tc^R, Ap^R)	(78)
pHV2	Modified pCM184 expressing *sacB* (Km^R, Tc^R, Ap^R)	This study
pCM157	Modified pCM62 expressing *cre* (Tc^R)	(78)
pCM639	Modified pCM638 carrying IS*phoA*/hah-Tc (Tc^R)	(86)
pAP5	Modified pCM62 carrying promoter-less *venus* (Tc^R)	(33)
pCM62	Broad-host-range shuttle vector (Tc^R)	(88)
pES502	pAP5 carrying *mxa* promoter region upstream of *venus* (Tc^R)	(33)
pHV25	pAP5 carrying *lutH* promoter region upstream of *venus* (Tc^R)	This study
pES667	pCM62 carrying *lutA* downstream of P$_{lac}$ promoter (Tc^R)	This study
pES810	pCM62 carrying *lutB* downstream of P$_{lac}$ promoter (Tc^R)	This study

Table 5. (continued)

Strain or plasmid	Description	Reference
pES666	pCM62 carrying *lutE* downstream of P$_{lac}$ promoter (Tc^R)	This study
pES815	pCM62 carrying *lutF* downstream of P$_{lac}$ promoter (Tc^R)	This study
pES668	pCM62 carrying *lutG* downstream of P$_{lac}$ promoter (Tc^R)	This study

630   Table 6. Primers used in this study.

Primer name	Sequence (5' to 3')£	Plasmid	Gene
Strain construction (pCM184 backbone)			
xoxF_1740ULEcoRI	ta**gaattc**GCCCCGATGGCAGGAATTAAAGT	pES319-320	*xoxF1*
xoxF_1740URKpnI	at**ggtacc**CTGCTTCGGCTCGTACTTCCACA		
xoxF_1740DLHpaI	ta**gttaac**GCCTGTTCTACGTCCCCACCAAC		
xoxF_1740DRSacI	ta**gagctc**CTGCACCGACGAGACCAAGAAGA		
exaF_1139ULEcoRI	gt**gaattc**CCTCCGACAGCCTACTGATGAAC	pES3-4	*exaF*
exaF_1139URKpnI	ca**ggtac**CGTCTTGGCGTCGTTGAGGATGTC		
exaF_1139DLHpaI	ca**gttaac**GCCGTTCGTGTCCAACATCAACTG		
exaF_1139DRSacI	ta**gagctc**GTTGATGCCAGCCGTCGTCTTGG		
pqqBCDE_1748ULEcoRI	ta**gaattc**TCAACATCCGCCAAGTCAGCTTC	pES369-458	*pqqBCDE*
pqqBCDE_1748URKpnI	at**ggtacc**ATATCGGGAGAGGCGTTCAGCAG		
pqqBCDE_1748DLHpaI	tata**gttaac**GACGAATGTGCAAAACCCGTTGA		
pqqBCDE_1748DRSacI	tata**gagctc**ACGTCGGTCGGATAGTCGTTGTC		
pqqF_2330ULEcoRI	ta**gaattc**ACCGTCCCGCCATCACCTATCGC	pES789-555	*pqqF*
pqqF_2330URKpnI	at**ggtacc**GTGATTCAGGCCTCGCTGCGGTC		
pqqF_2330DLHpaI	ta**gttaac**GGTGACCGGCTACCTGACCAAGG		
pqqF_2330DRSacI	ta**gagctc**GTAGCGCAGGCTGGCGATCATCT		
lysR_0863ULEcoRI	at**gaattc**CAGGCGCTGTCGAAGCATTCAAG	pES431-432	*MexAM1_META1p0863*
lysR_0863URKpnI	at**ggtacc**GGGTGTTCTCCAGCCTGAGCTTC		
lysR_0863DLHpaI	ta**gttaac**AGCGAGATGCCGACGATGGAGAG		
lysR_0863DRSacI	ta**gagctc**ATGGACGTTGGAGAGGTGGATCT		
sacB_BegNcoI	tagca**ccatgg**CGATCCTTTTTAACCCATCACATA	pHV2	
sacB_EndNcoI	tata**ccatgg**GATATCGGCATTTTCTTTTGCG		
Strain construction (pHV2 backbone)			
fae1_1767UBMunI	tata**caattg**CGCGGTTCATCTCGCCTATCTCA	pES498-499	*fae1*
fae1_1767UENcoI	tata**ccatgg**CCGTGCTTGTTGTTGACCAGACC		
fae1_1767DBAgeI	ctct**accggt**TTGTCACGGAGCAGCGTAAGTCG		
fae1_1767DESacI	tata**gagctc**GTGACGCTGTGGGTCGATGC		
cycK_1293UBAatII	tata**gacgtc**CCAGCCTGAAGGAACGAGAAGAAGC	pES604-605	*cycK*
cycK_1293UENcoI	tata**ccatgg**CGGGCATCACAGCTTGCACCA		
cycK_1293DBBshTI	tata**accggt**CGGTCGGCGTCAGCCAGAT		
cycK_1293DESacI	tata**gagctc**GTGACGACGATGCGCCGGAC		
ccmB_2825UBAatII	tata**gacgtc**CCATGTGGCATCACCGAATCAATGC	pES622-623	*ccmB*
ccmB_2825UENcoI	taat**ccatgg**TCAGGAAGAAGACGAGCGAGAGC		
ccmB_2825DBBshTI	tata**accggt**GCTGATGGTCCCGACTCTGATCTT		
ccmB_2825DESacI	tata**gagctc**AACCGATGTGAGGCCGATCTCC		

Table 6. (continued)

Primer name	Sequence (5' to 3')£	Plasmid	Gene
ccmC_2732UBAatII	tata**gacgtc**AGAGACTGAGCGGGACCAAGC	pES620-621	*ccmC*
ccmC_2732UENcoI	tata**ccatgg**CCAGTAGGTCCCCCACATCGG		
ccmC_2732DBBshTI	tata**accggt**CTCGATGCTCTACCCGCTGCTC		
ccmC_2732DESacI	atat**gagctc**GGTCGCGTGCGGGATCTG		
ABC2_2359UBAatII	atat**gacgtc**TGCGTATGCGATGGTTCCAGACAA	pES618-619	*MexAM1_META1p2359*
ABC2_2359UENcoI	tata**ccatgg**AGCGGAAACACGATGAGTGGAATCG		
ABC2_2359DBBshTI	tata**accggt**CTGGACGGCATCTCCTTCACGG		
ABC2_2359DESacI	tata**gagctc**CGAGAGCATCCCGTTGATCCGAA		
25pep_3908UBBglII	ctag**agatct**GGTACACAGGTTCGCCGGTTTTC	pES467-468	*MexAM1_META1p3908*
25pep_3908UENcoI	tata**ccatgg**GCCTCGTCCTCGTCCTTCTTCTT		
25pep_3908DBAgeI	atat**accggt**CCTACAAGCCCAAGGCGATGATC		
25pep_3908DESacI	tata**gagctc**CGGTCAGGAAGACGACGAACAGG		
25mem_3909UBBglII	tata**agatct**GACCTACAAGCCCAAGGCGATGA	pES469-470	*MexAM1_META1p3909*
25mem_3909UENcoI	atat**ccatgg**CGGTCAGGAAGACGACGAACAGG		
25pep_3909DBAgeI	tata**accggt**CACGCCGATGGTGATCGCCTATG		
25pep_3909DESacI	tata**gagctc**CGCTGCCGTTCTCACTTCCCTTG		
hss_2024UBAatII	tata**gacgtc**CGTCATGGGATTAAGTGCGAAACAGG	pES616-617	*hss*
hss_2024UENcoI	atat**ccatgg**CGGCTCTTGTCGTACTCGAAATGGC		
hss_2024DBBshTI	tata**accggt**GACCTGCCACTACGCCTACCAC		
hss_2024DESacI	atat**gagctc**TGCGAAGGCCCGAACCGTAAC		
xoxG_1741UBbglII	tata**agatct**GCAACTTCATCGCCTGGGACAAC	pES459-460	*xoxG*
xoxG_1741UEKpnI	acgt**ggtacc**CGTCGTTGGTGATGTGGAAGGTC		
xoxG_1741DBAgeI	atat**accggt**GCGAGAAGGAAGACAAGCCCG		
xoxG_1741DESacI	tata**gagctc**AATCGAGCGGGTGGAATGAGAA		
xoxJ_1742UBBglII	atat**agatct**TCTTCTTGGTCTCGTCGGTGCAG	pES461-462	*xoxJ*
xoxJ_1742UEKpnI	tata**ggtacc**CGCAGCTTGACCTTCAGCTCGTC		
xoxJ_1742DBAgeI	tata**accggt**GCCACAACGAGAACGACTGGAAG		
xoxJ_1742DESacI	tata**gagctc**CTATCTCGGGCTCGGCAGCAATA		
xoxD_1771UBAatII	tata**gacgtc**GCCATGCCGAGTTCGAGGTG	pES608-609	*MexAM1_META1p1771*
xoxD_1771UENcoI	atta**ccatgg**ACCCTTGCCCTCGGTCTTCTC		
xoxD_1771DBBshTI	atat**accggt**GTGAAGGCCAAGGCCGAGAAG		
xoxD_1771DESacI	tata**gagctc**AGTCGGATTCGCGGTTGGCTAC		
orf6_1746UBMunI	atat**caattg**GAGCAGCCGCAGATGAAGAACAC	pES493-494	*MexAM1_META1p1746*
orf6_1746UENcoI	atat**ccatgg**GTCGGTCGGATAGTCGTTGTCGT		
orf6_1746DBAgeI	atac**accggt**CCTGCCCTATGACCGCTACGTT		

Table 6. (continued)

Primer name	Sequence (5' to 3')£	Plasmid	Gene
orf6_1746DESacI	atat**gagctc**CCGGCATTCCAACGAGATGTAGA		
orf7_1747UBMunI	atat**caattg**TTACGGCTGGGCCTATGTGAACC	pES495-496	*MexAM1_META1p1747*
orf7_1747UENcoI	atat**ccatgg**GTGTTCTTCATCTGCGGCTGCTC		
orf7_1747DBAgeI	atat**accggt**CCAACCCGCGCTACATCAAGAAC		
orf7_1747DESacI	atat**gagctc**TGACGTTGAGGATGCCCATTACC		
lutA_1778UBAatII	tata**gacgtc**GGACGGGCTTTAACTCGGTTTCC	pES610-611	*lutA*
lutA_1778UENcoI	tata**ccatgg**GGCGAAGGTCTGCTTGGTGAAG		
lutA_1778DBBshTI	taat**accggt**GACTGGGGCGATTATGTCCCGT		
lutA_1778DESacI	tata**gagctc**CCGTCGCTCAAACAAGAAACAGATGA		
lutB_1779UBAatII	atat**gacgtc**GCAAGTTCGGCCTCAAGATCACC	pES638-639	*lutB*
lutB_1779UENcoI	atat**ccatgg**GGCCTTCTCGTTGGTGACGTAGG		
lutB_1779DBAgeI	atat**accggt**CTCTACACGACCAACGGCACCTC		
lutB_1779DESacII	atat**gagctc**CCCCCTCACTTCACCGTAATCGT		
lutC_1780UBMunI	atat**caattg**GAGGTGATCGACAACGTGCTGGT	pES642-643	*lutC*
lutC_1780UENcoI	atat**ccatgg**GTTCTGCCGCTTCTTGGTGAGGT		
lutC_1780DBMluI	atat**acgcgt**CGGCACGATTACGGTGAAGTGAG		
lutC_1780DESacI	atat**gagctc**GCTGGCGTGGTAATGCAGGTTTT		
lutD_1781UBBglII	atat**agatct**TGTCCGAGAAGGCGATGCAAG	pES644-645	*lutD*
lutD_1781UEMunI	atat**caattg**CAAGCACCACGAGGGCAAGC		
lutD_1781DBAgeI	atat**accggt**GAGCAAGGCGAGGGCGAGTT		
lutD_1781DESacI	atgg**gagctc**ATCGAGGTGCCGAGCGTGTT		
lutE_1782UBAatII	tata**gacgtc**GTCCGAGAAGGCGATGCAAGAAC	pES612-613	*lutE*
lutE_1782UENcoI	tata**ccatgg**CGGCCCTCCTGATTGTTGTAGAGC		
lutE_1782DBBshTI	tata**accggt**CGACAAGAGAGCCGTGAAGTCCG		
lutE_1782DESacI	tata**gagctc**CAGAGCGGATATAGGGCAGAAGAGG		
lutF_1783UBAatII	tatc**gacgtc**ACCCGGCTCTACAACAATCAGGAG	pES640-641	*lutF*
lutF_1783UENcoI	atat**ccatgg**TTCTGGTGGAAGAAACGCAGCAG		
lutF_1783DBAgeI	atat**accggt**CGAATTGGTGCGTTTCGCTCTCT		
lutF_1783DESacI	atat**gagctc**TCGCAGACGTAGGAGACCGACTG		
lutG_1784UBAatII	tata**gacgtc**GCTGGTCTGGCTGTTCATCTTCG	pES614-615	*lutG*
lutG_1784UENcoI	atat**ccatgg**TAGCCCAGCGTCGTCACCTTG		
lutG_1784DBBshTI	tata**accggt**CGGACAAGGTGAGCGAGTTCAAGGA		
lutG_1784DESacI	tata**gagctc**GCAACATATTTCTGCCGTTCCGTCAA		
lutH_1785UBBglII	atat**agatct**TGCCGAATGATCCCGGTAAAGTC	pES465-466	*lutH*
lutH_1785UENcoI	tata**ccatgg**CGAACGAGGTGTTGAAACGAGCA		
lutH_1785DBAgeI	atat**accggt**CCCTGCTCGCTCACCAACTTCTT		

Table 6. (continued)

Primer name	Sequence (5' to 3')[£]	Plasmid	Gene
lutH_1785DESacI	atat**gagctc**TTCGGCCTCGCTTAGATTGCGTA		
Transcriptional reporter fusion (pAP5 backbone)			
lutH_1785BegAclI	tata**aacgtt**GCAGCCGTCGAAGCGATTTGC	pHV25	*lutH*
lutH_1785EndEcoRI	tata**gaattc**CCGAACGAGGTGTTGAAACGAGC		
Complementation (pCM62 backbone)			
lutA_1778BegKpnI	atat**ggtacc**CTTATGCTTCTCTCGTTGAGGCCG	pES667	*lutA*
lutA_1778EndSacI	atcg**gagctc**ATCGCCCGGATCGCTTCC		
lutB_1779BegKpnI	tattt**ggtacc**AAGCGATCCGGGCGATGC	pES810	*lutB*
lutB_1779EndSacI	tattt**gagctc**ACCTTTCGGGACGATGCCTTAGC		
lutE_1782BegKpnI	atat**ggtacc**GAGCAAGGCGAGGGCGAGTT	pES666	*lutE*
lutE_1782EndKpnI	atat**gagctc**GGCGGGATTCGAGGGAGACC		
lutF_1783BegKpnI	tattt**ggtacc**AAGGGTCGCATCGTCGCCCGTGG	pES815	*lutF*
lutF_1783EndSacI	tattt**gagctc**ACGAGCCGTAGCCCAGCGTCG		
lutG_1784BegKpnI	atat**ggtacc**GCCGCAATCGCCTACGATCC	pES668	*lutG*
lutG_1784EndSacI	atat**gagctc**CTTCGTCGCTTCCGACCGAAAG		

631   [£]Restriction endonuclease sites are in bolded font. Capitalized letters represent bases that anneal

632   to the target DNA sequence.

 FIGURE LEGENDS

Fig 1. Schematic representation of the metabolic processes relevant to the *mxaF fae*

transposon mutagenesis study. XoxF1 and XoxF2 oxidize methanol to formaldehyde, which

accumulates to lethal levels in the *fae* mutant strain. If a process required for XoxF-dependent

methanol oxidation is disrupted by a transposon insertion, formaldehyde is reduced or

eliminated, and cells use succinate for growth. A dashed line around the formaldehyde

transporter is to indicate that this function has not been demonstrated.

Fig 2. Growth of mutant strains lacking genes identified through transposon mutagenesis

in medium containing methanol and La^{3+}. Growth of *M. extorquens* AM1 wild-type is

represented by blue circles and depicted mutant strains (red triangles) have the following genes

deleted: A - *MexAM1_META1p1771*, *mxaD* homolog; B - *MexAM1_META1p1746, orf6*; C -

MexAM1_META1p1747, orf7; D - *MexAM1_META1p2024*, homospermidine synthase; E -

MexAM1_META1p2359, ABC transporter of unknown function; F - *MexAM1_META1p3908*,

aminopeptidase. Representative data from biological triplicates are shown. One-way analysis of

variance (ANOVA) determined that growth rate differences between mutant and parent strains

are significantly different ($p < 0.005$).

Fig 3. Growth of *xoxG* and *xoxJ* mutant strains in the presence (filled symbols) and

absence (open symbols) of La^{3+}. Representative data from biological triplicates are shown.

One-way analysis of variance (ANOVA) determined that growth rate differences between

mutant and parent strains are significantly different ($p < 0.005$).

Fig 4. Characterization of the *lut* cluster. A) Genomic map of the *lut* genes

(*MexAM1_META1p1778* to *MexAM1_META1p1787*). Black, genes encoding the ABC transport

system; green, gene encoding the TonB-dependent transporter; blue, genes encoding putative

periplasmic proteins identified by transposon mutagenesis; white, lanmodulin and additional

genes encoding periplasmic proteins not identified by transposon mutagenesis. B) Model for Ln

transport. The three-dimensional structures of monomers of LutH and LutA were predicted using

homology modeling (HHpredserver and MODELLER) (89) and homodimers of the ABC

transporter were predicted using GalaxyHomomer (90). C) Growth of *mxaF* and *mxaF lut* mutant

strains with 2 µM LaCl$_3$. Graphs depict representative data from three biological replicates.

Growth rate averages and standard deviations are shown in Table 2.

Fig 5. Growth studies and measurement of La^{3+} uptake suggest La^{3+} storage capability. A)

Growth of wild type (red dashed circles), *lutH* (white triangles), *lutA* (blue squares), *lutE* (gray

diamonds), and *lutF* (black circles) strains grown with limiting succinate (3.75 mM), methanol

(125 mM), and 2 µM LaCl$_3$. Data are the average of three biological replicates. Standard

deviations between biological replicates were less than ±0.02. B) La^{3+} concentrations (µM) in

culture supernatants of *lut* mutant strains depicted in same color code as in (A). Data depict the

average of three biological replicates with error bars showing the standard deviation. One-

way analysis of variance (ANOVA) followed by a T-test was used to represent statistical

significance. *The *p*-value is <0.005. C) Growth of the *mxaF* mutant strain with excess LaCl$_3$ (20

µM) (red filled circles). Stationary phase cells were washed and reinoculated into media lacking

Ln (subculture #1, black triangles) and the process was repeated (subculture #2, empty circles).

Representative data from three biological replicates is shown. Standard deviations for growth

rates between replicates were less than ± 0.02.

Fig 6. Visualization of La^{3+} storage. TEM of ultrathin sections of wild-type cells grown with

3.75 mM succinate and 125 mM methanol containing (A-B) and lacking (E-F) 20 µM La^{3+}. C)

and G) Elemental analysis of electron dense deposits from cells grown with (C) and without (G)

La^{3+}. D) High-resolution transmission electron microscopy analysis of the wild-type

strain shows an atomic lattice structure in electron-dense areas suggesting that La^{3+} is embedded

in cytoplasmic crystals.

Fig 7. Visualization of La^{3+} localization using TEM. Thin sections of A) *lutH*, B) *lutE*,

C) *lutF*, D) *lutA*, and G) wild-type strains grown with 3.75 mM succinate, 125 mM methanol,

and 20 µM LaCl$_3$. White arrows indicate deposits of electron scattering material in the

periplasm. Cells were fixed with 2.5% glutaraldehyde and stained with OsO$_4$ and uranyl acetate

to detect cell membranes (left subpanel), or left unstained for elemental analysis (right

subpanel). E) Magnification of the La^{3+}-deposits localized in the periplasmic space from the *lutA*

mutant strain; black arrows indicate the boundaries of the outer membrane and inner

membrane. F) Elemental analysis of the electron-dense deposits observed inside the periplasm.

REFERENCES

1. Zepf V, Reller A, Rennie C, Ashfield M, Simmons J. 2014. Materials Critical to the Energy Industry. An Introduction. 2nd ed. BP.
2. Martinez-Gomez NC, Vu HN, Skovran E. 2016. Lanthanide Chemistry: From Coordination in Chemical Complexes Shaping Our Technology to Coordination in Enzymes Shaping Bacterial Metabolism. Inorg Chem 55:10083–10089.
3. Teo RD, Termini J, Gray HB. 2016. Lanthanides: Applications in Cancer Diagnosis and Therapy. J Med Chem 59:6012–6024.
4. Mattocks JA, Ho J V, Cotruvo JA. 2019. A Selective, Protein-Based Fluorescent Sensor with Picomolar Affinity for Rare Earth Elements. J Am Chem Soc 141:2857–2861.
5. Hibi Y, Asai K, Arafuka H, Hamajima M, Iwama T, Kawai K. 2011. Molecular Structure of La3+-induced Methanol Dehydrogenase-like Protein in Methylobacterium radiotolerans. J Biosci Bioeng 111:547–9.
6. Pol A, Barends TRM, Dietl A, Khadem AF, Eygensteyn J, Jetten MSM, Op den Camp HJM. 2014. Rare Earth Metals are Essential for Methanotrophic Life in Volcanic Mudpots. Environ Microbiol 16:255–264.
7. Nakagawa T, Mitsui R, Tani A, Sasa K, Tashiro S, Iwama T, Hayakawa T, Kawai K. 2012. A Catalytic Role of XoxF1 as La3+-dependent Methanol Dehydrogenase in Methylobacterium extorquens strain AM1. PLoS One 7:e50480.
8. Fitriyanto NA, Nakamura M, Muto S, Kato K, Yabe T, Iwama T, Kawai K, Pertiwiningrum A. 2011. Ce 3+ -induced exopolysaccharide production by Bradyrhizobium sp. MAFF211645. J Biosci Bioeng 111:146–152.
9. Wehrmann M, Billard P, Martin-Meriadec A, Zegeye A, Klebensberger J. 2017. Functional Role of Lanthanides in Enzymatic Activity and Transcriptional Regulation of Pyrroloquinoline Quinone-Dependent Alcohol Dehydrogenases in Pseudomonas putida KT2440. MBio 8:e00570-17.
10. Wehrmann M, Berthelot C, Billard P, Klebensberger J. 2018. The PedS2/PedR2 Two-Component System Is Crucial for the Rare Earth Element Switch in Pseudomonas putida KT2440. mSphere 3:e00376-18.
11. Wang L, Suganuma S, Hibino A, Mitsui R, Tani A, Matsumoto T, Ebihara A, Fitriyanto NA, Pertiwiningrum A, Shimada M, Hayakawa T, Nakagawa T. 2019. Lanthanide-dependent Methanol Dehydrogenase from the Legume Symbiotic Nitrogen-fixing Bacterium Bradyrhizobium diazoefficiens strain USDA110. Enzyme Microb Technol 130:109371.
12. Skovran E, Martinez-Gomez NC. 2015. Just Add Lanthanides. Science 348:862–863.
13. van Teeseling MCF, Pol A, Harhangi HR, van der Zwart S, Jetten MSM, Op den Camp HJM, van Niftrik L. 2014. Expanding the Verrucomicrobial Methanotrophic World: Description of Three Novel Species of Methylacidimicrobium gen. nov. Appl Environ Microbiol 80:6782–6791.
14. Versantvoort W, Pol A, Daumann LJ, Larrabee JA, Strayer AH, Jetten MSM, van Niftrik L, Reimann J, Op den Camp HJM. 2019. Characterization of a Novel Cytochrome c as the Electron Acceptor of XoxF-MDH in the Thermoacidophilic Methanotroph Methylacidiphilum fumariolicum SolV. Biochim Biophys Acta - Proteins Proteomics 1867:595–603.
15. Lu X, Gu W, Zhao L, Farhan M, Haque U, Dispirito AA, Semrau JD, Gu B. 2017.

742   Methylmercury Uptake and Degradation by Methanotrophs. Sci Adv 3:e170004.

743   16. Whitaker WB, Sandoval NR, Bennett RK, Fast AG, Papoutsakis ET. 2015. Synthetic
744   Methylotrophy: Engineering the Production of Biofuels and Chemicals Based on the
745   Biology of Aerobic Methanol Utilization. Curr Opin Biotechnol 33:165–175.

746   17. Ochsner AM, Sonntag F, Buchhaupt M, Schrader J, Vorholt JA. 2015. Methylobacterium
747   extorquens: Methylotrophy and Biotechnological Applications. Appl Microbiol
748   Biotechnol 99:517–34.

749   18. Chistoserdova L, Kalyuzhnaya MG. 2018. Current Trends in Methylotrophy. Trends
750   Microbiol 26:703-714.

751   19. Lumpe H, Pol A, Op den Camp H, Daumann L. 2018. Impact of the Lanthanide
752   Contraction on the Activity of a Lanthanide-dependent Methanol Dehydrogenase – A
753   Kinetic and DFT Study and DFT Study. Dalt Trans 47:10463–10472.

754   20. Prejanò M, Iziana Marino T, Russo N, Marino T, Russo N. 2017. How Can Methanol
755   Dehydrogenase from Methylacidiphilum fumariolicum Work with the Alien Ce III Ion in
756   the Active Center? A Theoretical Study. Chem - A Eur J 23:8652–8657.

757   21. McSkimming A, Cheisson T, Carroll PJ, Schelter EJ, Roy P. 2018. Functional Synthetic
758   Model for the Lanthanide-Dependent Quinoid Alcohol Dehydrogenase Active Site. J Am
759   Chem Soc 140:1223–1226.

760   22. Huang J, Yu Z, Groom J, Cheng J-FF, Tarver A, Yoshikuni Y, Chistoserdova L. 2019.
761   Rare Earth Element Alcohol Dehydrogenases Widely Occur Among Globally Distributed,
762   Numerically Abundant and Environmentally Important Microbes. ISME J 13:2005–2017.

763   23. Keltjens JT, Pol A, Reimann J, Op Den Camp HJM. 2014. PQQ-dependent Methanol
764   Dehydrogenases: Rare-earth Elements Make a Difference. Appl Microbiol Biotechnol
765   98:6163–6183.

766   24. Chistoserdova L. 2016. Lanthanides: New Life Metals? World J Microbiol Biotechnol
767   32:138.

768   25. Good NM, Moore RS, Suriano CJ, Martinez-Gomez NC. 2019. Contrasting in vitro and in
769   vivo Methanol Oxidation Activities of Lanthanide-dependent Alcohol Dehydrogenases
770   XoxF1 and ExaF from Methylobacterium extorquens AM1. Sci Rep 9:4248.

771   26. Good NM, Vu HN, Suriano CJ, Subuyuj GA, Skovran E, Martinez-Gomez NC. 2016.
772   Pyrroloquinoline Quinone-Containing Ethanol Dehydrogenase in Methylobacterium
773   extorquens AM1 Extends Lanthanide-Dependent Metabolism to Multi-Carbon Substrates.
774   J Bacteriol 198:3109–3118.

775   27. Vu HN, Subuyuj GA, Vijayakumar S, Good NM, Martinez-Gomez NC, Skovran E. 2016.
776   Lanthanide-dependent Regulation of Methanol Oxidation Systems in Methylobacterium
777   extorquens AM1 and Their Contribution to Methanol Growth. J Bacteriol 198:1250–1259.

778   28. Deng YW, Ro SY, Rosenzweig AC. 2018. Structure and Function of the Lanthanide-
779   dependent Methanol Dehydrogenase XoxF from the Methanotroph Methylomicrobium
780   buryatense 5GB1C. J Biol Inorg Chem 23:1037–1047.

781   29. Taubert M, Grob C, Howat AM, Burns OJ, Dixon JL, Chen Y, Murrell JC. 2015. XoxF
782   Encoding an Alternative Methanol Dehydrogenase is Widespread in Coastal Marine
783   Environments. Environ Microbiol 17:3937–3948.

784   30. Huang J, Yu Z, Chistoserdova L. 2018. Lanthanide-dependent Methanol Dehydrogenases
785   of XoxF4 and XoxF5 Clades Are Differentially Distributed Among Methylotrophic
786   Bacteria and They Reveal Different Biochemical Properties. Front Microbiol 9:1366.

31. Featherston ER, Rose HR, McBride MJ, Taylor E, Boal AK, Cotruvo J. 2019.

Biochemical and Structural Characterization of XoxG and XoxJ and Their Roles in
Activity of the Lanthanide-dependent Methanol Dehydrogenase, XoxF. ChemBioChem
20:2360–2372.
32.   Chistoserdova L, Chen SW, Lapidus A, Lidstrom ME. 2003. Methylotrophy in
Methylobacterium extorquens AM1 From a Genomic Point of View. J Bacteriol
185:2980–2987.
33.   Skovran E, Palmer AD, Rountree AM, Good NM, Lidstrom ME. 2011. XoxF is Required
for Expression of Methanol Dehydrogenase in Methylobacterium extorquens AM1. J
Bacteriol 193:6032–6038.
34.   Chistoserdova L. 2019. New Pieces to the Lanthanide Puzzle. Mol Microbiol 111:1127–
1131.
35.   Picone N, Op den Camp HJ. 2019. Role of Rare Earth Elements in Methanol Oxidation.
Curr Opin Chem Biol 49:39–44.
36.   Skovran E, Raghuraman C, Martinez-Gomez NC. 2019. Lanthanides in Methylotrophy, p.
101–116. In Methylotrophs and Methylotroph Communities.
37.   Chu F, Beck DAC, Lidstrom ME. 2016. MxaY Regulates the Lanthanide-mediated
Methanol Dehydrogenase Switch in Methylomicrobium buryatense. PeerJ 4:e2435.
38.   Gu W, Haque MFU, DiSpirito AA, Semrau JD. 2016. Uptake and Effect of Rare Earth
Elements on Gene Expression in Methylosinus trichosporium OB3b. FEMS Microbiol
Lett 363:pii: fnw129.
39.   Masuda S, Suzuki Y, Fujitani Y, Mitsui R, Nakagawa T, Shintani M, Tani A. 2018.
Lanthanide-Dependent Regulation of Methylotrophy in Methylobacterium aquaticum
Strain 22A. mSphere 3:e00462-17.
40.   Springer AL, Morris CJ, Lidstrom ME. 1997. Molecular Analysis of mxbD and mxbM, a
Putative Sensor-regulator Pair Required for Oxidation of Methanol in Methylobacterium
extorquens AM1. Microbiology 143:1737–1744.
41.   Ochsner AM, Hemmerle L, Vonderach T, Nüssli R, Bortfeld-Miller M, Hattendorf B,
Vorholt JA. 2019. Use of Rare-earth Elements in the Phyllosphere Colonizer
Methylobacterium extorquens PA1. Mol Microbiol 111:1152–1166.
42.   Daumann LJ. 2019. Essential and Ubiquitous: The Emergence of Lanthanide
Metallobiochemistry. Angew Chem Int Ed Engl 58:12795–12802.
43.   Cotruvo, Jr. JA, Featherston ER, Mattocks JA, Ho J V., Laremore TN. 2018. Lanmodulin:
A Highly Selective Lanthanide-binding Protein from a Lanthanide-utilizing Bacterium. J
Am Chem Soc jacs.8b09842.
44.   Sawyer EB, Barker PD. 2012. Continued Surprises in the Cytochrome c Biogenesis Story.
Protein Cell 3:405–409.
45.   Zheng C, Huang Y, Zhao J, Chistoserdova F. 2018. Physiological Effect of XoxG(4) on
Lanthanide-Dependent Methanotrophy. MBio 9:e02430-17.
46.   Chou H-H, Marx CJ. 2012. Optimization of Gene Expression Through Divergent
Mutational Paths. Cell Rep 1:133–140.
47.   Langley S, Beveridge TJ. 1999. Effect of O-Side-Chain-Lipopolysaccharide Chemistry on
Metal Binding. Appl Environ Microbiol 65:489–498.
48.   Solovyev V SA. 2011. Automatic Annotation of Microbial Genomes and Metagenomic
Sequences, p. 458. In Li, RW (ed.), Metagenomics and its Applications in Agriculture,
Biomedicine and Environmental Studies. Nova Science Publisher's.
49.   Chen Z, Lewis KA, Shultzaberger RK, Lyakhov IG, Zheng M, Doan B, Storz G,

834   Schneider TD. 2007. Discovery of Fur Binding Site Clusters in Escherichia coli by
835   Information Theory Models. Nucleic Acids Res 35:6762–6777.

50. Young GM, Postle K. 1994. Repression of tonB Transcription During Anaerobic Growth Requires Fur Binding at the Promoter and a Second Factor Binding Upstream. Mol Microbiol 11:943–954.

51. Groom JD, Ford SM, Pesesky MW, Lidstrom ME. 2019. A Mutagenic Screen Identifies a TonB-dependent Receptor Required for the Lanthanide Metal Switch in the Type I methanotroph Methylotuvimicrobium buryatense 5GB1C. J Bacteriol 201:pii: e00120-19.

52. Klaus T, Joerger R, Olsson E, Granqvist C-G. 1999. Silver-based Crystalline Nanoparticles, Microbially Fabricated. Proc Natl Acad Sci 96:13611–13614.

53. Maleke M, Valverde A, Vermeulen JG, Cason E, Gomez-Arias A, Moloantoa K, Coetsee-Hugo L, Swart H, Van Heerden E, Castillo J. 2019. Biomineralization and Bioaccumulation of Europium by a Thermophilic Metal Resistant Bacterium. Front Microbiol 10:1–10.

54. Leeson TS, Higgs G. 1982. Lanthanum as an Intracellular Stain for Electron Microscopy. Histochem J 14:553–60.

55. Bayer ME, Bayer MH. 1991. Lanthanide Accumulation in the Periplasmic Space of Escherichia coli B. J Bacteriol 173:141–149.

56. Merroun ML, Ben Chekroun K, Arias JM, Gonz A Alez-Munoz MT. 2003. Lanthanum Fixation by Myxococcus xanthus: Cellular Location and Extracellular Polysaccharide Observation. Chemosphere 52:113–120.

57. Hashimoto H, Mannami M NT. 1961. Dynamical Theory of Electron Diffraction for the Electron Microscopic Image of Crystal Lattices II. Image of Superposed Crystals (Moiré Pattern). Philos Trans R Soc 253:490–516.

58. Krossa S, Faust A, Ober D, Scheidig AJ. 2015. Comprehensive Structural Characterization of the Bacterial Homospermidine Synthase-an Essential Enzyme of the Polyamine Metabolism. Sci Rep 6:19501.

59. Wortham BW, Patel, CN, Oliveira M. 2007. Polyamines in Bacteria: Pleiotropic Effects Yet Specific Mechanisms. Adv Exp Med Biol 603:106–15.

60. Chistoserdova L, Lidstrom ME. 1997. Molecular and Mutational Analysis of a DNA Region Separating Two Methylotrophy Gene Clusters in Methylobacterium extorquens AM1. Microbiology 143:1729–36.

61. Toyama H, Inagaki H, Matsushita K, Anthony C, Adachi O. 2003. The Role of the MxaD Protein in the Respiratory Chain of Methylobacterium extorquens During Growth on Methanol. Biochim Biophys Acta 1647:372–375.

62. Morris CJ, Biville F, Turlin E, Lee E, Ellermann K, Fan WH, Ramamoorthi R, Springer AL, Lidstrom ME. 1994. Isolation, Phenotypic Characterization, and Complementation Analysis of Mutants of Methylobacterium extorquens AM1 Unable to Synthesize Pyrroloquinoline Quinone and Sequences of pqqD, pqqG, and pqqC. J Bacteriol 176:1746–55.

63. Cheng Y, Zhang L, Bian X, Zuo H, Dong H. 2018. Adsorption and Mineralization of REE-lanthanum Onto Bacterial Cell Surface. Env Sci Pollut Res 25:22334–22339.

64. Mullen MD, Wolf DC, Ferris FG, Beveridge TJ, Flemming CA, Bailey3 GW. 1989. Bacterial Sorption of Heavy Metals. Appl Environ Microbiol 55:3143–3149.

65. Zhan G, Li D, Liang Zhang &. 2012. Aerobic Bioreduction of Nickel(II) to Elemental Nickel With Concomitant Biomineralization. Appl Microbiol Biotechnol 96:273– 281.

880   66.   Bai HJ, Zhang ZM, Guo Y, Yang GE. 2009. Biosynthesis of Cadmium Sulfide
881   Nanoparticles by Photosynthetic Bacteria Rhodopseudomonas palustris. Colloids Surfaces
882   B Biointerfaces 70:142–146.
883   67.   Sousa T, Chung A-P, Pereira A, Piedade AP, Morais P V. 2013. Aerobic Uranium
884   Immobilization by Rhodanobacter A2-61 Through Formation of Intracellular Uranium-
885   phosphate Complexes. Metallomics 5:390–397.
886   68.   Couradeau E, Benzerara K, Gérard E, Moreira D, Bernard S, Brown GE, López-García P.
887   2012. An Early-Branching Microbialite Cyanobacterium Forms Intracellular Carbonates.
888   Science 336:459–462.
889   69.   Debieux CM, Dridge EJ, Mueller CM, Splatt P, Paszkiewicz K, Knight I, Florance H,
890   Love J, Titball RW, Lewis RJ, Richardson DJ, Butler CS. 2011. A Bacterial Process for
891   Selenium Nanosphere Assembly. Proc Natl Acad Sci 108:13480–13485.
892   70.   Rahn-Lee L, Komeili A. 2013. The Magnetosome Model: Insights Into The Mechanisms
893   of Bacterial Biomineralization. Front Microbiol 4:352.
894   71.   Kulaev I. 1979. The Biochemistry of Inorganic Polyphosphates. John Wiley & Sons Ltd.
895   72.   Friedberg I, Avigad G. 1968. Structures Containing Polyphosphate in Micrococcus
896   lysodeikticus. J Bacteriol 96:544–553.
897   73.   Pallerla SR, Knebel S, Polen T, Klauth P, Hollender J, Wendisch VF, Schoberth SM.
898   2005. Formation of Volutin Granules in Corynebacterium glutamicum. FEMS Microbiol
899   Lett 243:133–140.
900   74.   Widra A. 1959. Metachromatic Granules of Microorganisms. J Bacteriol 78:664–670.
901   75.   Bertani G. 1951. Studies on Lysogenesis. I. The Mode of Phage Liberation by Lysogenic
902   Escherichia coli. J Bacteriol 62:293–300.
903   76.   Delaney NF, Kaczmarek ME, Ward LM, Swanson PK, Lee M-C, Marx CJ. 2013.
904   Development of an Optimized Medium, Strain and High-throughput Culturing Methods
905   for Methylobacterium extorquens. PLoS One 8:e62957.
906   77.   Marx CJ. 2008. Development of a Broad-host-range sacB-based Vector for Unmarked
907   Allelic Exchange. BMC Res Notes 1:1.
908   78.   Marx CJ, Lidstrom ME. 2002. Broad-host-range Cre-lox System for Antibiotic Marker
909   Recycling in Gram-negative Bacteria. Biotechniques 33:1062–1067.
910   79.   Simon R, Priefer U, Pühler A. 1983. A Broad Host Range Mobilization System for In
911   Vivo Genetic Engineering: Transposon Mutagenesis in Gram Negative Bacteria. Nat
912   Biotechnol 1:784–791.
913   80.   Jacobs MA, Alwood A, Thaipisuttikul I, Spencer D, Haugen E, Ernst S, Will O, Kaul R,
914   Raymond C, Levy R, Chun-Rong L, Guenthner D, Bovee D, Olson M V., Manoil C. 2003.
915   Comprehensive Transposon Mutant Library of Pseudomonas aeruginosa. Proc Natl Acad
916   Sci 100:14339–14344.
917   81.   Marx CJ, O'Brien BN, Breezee J, Lidstrom ME. 2003. Novel Methylotrophy Genes of
918   Methylobacterium extorquens AM1 Identified by Using Transposon Mutagenesis
919   Including a Putative Dihydromethanopterin Reductase. J Bacteriol 185:669–673.
920   82.   Manoil C, Traxler B. 2000. Insertion of In-frame Sequence Tags into Proteins Using
921   Transposons. Methods 20:55–61.
922   83.   Nagai T, Ibata K, Park ES, Kubota M, Mikoshiba K, Miyawaki A. 2002. A Variant of
923   Yellow Fluorescent Protein With Fast and Efficient Maturation for Cell-biological
924   Applications. Nat Biotechnol 20:87–90.
84.   Hogendoorn C, Roszczenko-Jasińska P, Martinez-Gomez NC, de Graaff J, Grassl P, Pol

A, Op den Camp HJM, Daumann LJ. 2018. Facile Arsenazo III-Based Assay for
Monitoring Rare Earth Element Depletion from Cultivation Media for Methanotrophic
and Methylotrophic Bacteria. Appl Environ Microbiol 84:e02887-17.
85.   Nunn DN, Lidstrom ME. 1986. Isolation and Complementation Analysis of 10 Methanol
Oxidation Mutant Classes and Identification of the Methanol Dehydrogenase Structural
Gene of Methylobacterium sp. Strain AM1. J Bacteriol 166:581–90.
86.   Marx CJ, Chistoserdova L, Lidstrom ME. 2003. Formaldehyde-detoxifying Role of the
Tetrahydromethanopterin-linked Pathway in Methylobacterium extorquens AM1. J
Bacteriol 185:7160–8.
87.   Figurski DH, Helinski DR. 1979. Replication of an Origin-containing Derivative of
Plasmid RK2 Dependent on a Plasmid Function Provided in Trans. Proc Natl Acad Sci
USA 76:1648–52.
88.   Marx CJ, Lidstrom ME. 2001. Development of Improved Versatile Broad-host-range
Vectors for Use in Methylotrophs and Other Gram-negative Bacteria. Microbiology
147:2065–75.
89.   Söding J, Biegert A, Lupas AN. 2005. The HHpred Interactive Server for Protein
Homology Detection and Structure Prediction. Nucleic Acids Res 33:W244–W248.
90.   Baek M, Park T, Heo L, Park C, Seok C. 2017. GalaxyHomomer: A Web Server for
Protein Homo-oligomer Structure Prediction From a Monomer Sequence or Structure.
Nucleic Acids Res 45:W320–W324.
946

+ La³⁺

No La³⁺